



DENMARK'S NATIONAL INVENTORY REPORT 2014

Emission Inventories 1990-2012 - Submitted under the United Nations Framework Convention on Climate Change and the Kyoto Protocol

Scientific Report from DCE - Danish Centre for Environment and Energy

No. 101

2014



AARHUS
UNIVERSITY

DCE - DANISH CENTRE FOR ENVIRONMENT AND ENERGY

[Blank page]

DENMARK'S NATIONAL INVENTORY REPORT 2014

Emission Inventories 1990-2012 - Submitted under the United Nations
Framework Convention on Climate Change and the Kyoto Protocol

Scientific Report from DCE – Danish Centre for Environment and Energy

No. 101

2014

Ole-Kenneth Nielsen¹
Marlene S. Plejdrup¹
Morten Winther¹
Malene Nielsen¹
Steen Gyldenkærne¹
Mette Hjorth Mikkelsen¹
Rikke Albrechtsen¹
Marianne Thomsen¹
Katja Hjelgaard¹
Leif Hoffmann¹
Patrik Fauser¹
Henrik G. Bruun¹
Vivian Kvist Johannsen²
Thomas Nord-Larsen²
Lars Vesterdal²
Inge Stupak Møller²
Ole Hjort Caspersen²
Erik Rasmussen³
Susanne Bødtker Petersen⁴
Lene Baunbæk⁵
Maria Gunnleivsdóttir Hansen⁶

¹ Aarhus University, Department of Environmental Science

² University of Copenhagen, Faculty of Sciences (IGN)

³ Danish Energy Agency

⁴ Danish Business Authority

⁵ Statistics Greenland

⁶ Umhvørvisstovan (The Environment Agency, The Faroe Islands)



Data sheet

Series title and no.:	Scientific Report from DCE – Danish Centre for Environment and Energy No. 101
Title:	Denmark's National Inventory Report 2014
Subtitle:	Emission Inventories 1990-2012 - Submitted under the United Nations Framework Convention on Climate Change and the Kyoto Protocol
Authors:	Ole-Kenneth Nielsen ¹ , Marlene S. Plejdrup ¹ , Morten Winther ¹ , Malene Nielsen ¹ , Steen Gyldenkaerne ¹ , Mette Hjorth Mikkelsen ¹ , Rikke Albrektsen ¹ , Marianne Thomsen ¹ , Katja Hjelgaard ¹ , Leif Hoffmann ¹ , Patrik Fayser ¹ , Henrik G. Bruun ¹ , Vivian Kvist Johannsen ² , Thomas Nord-Larsen ² , Lars Vesterdal ² , Inge Stupak Møller ² , Ole Hjort Caspersen ² , Erik Rasmussen ³ , Susanne Bødtker Petersen ⁴ , Lene Baunbæk ⁵ , Maria Gunnleivsdóttir Hansen ⁶
Institutions:	¹ Department of Environmental Science, Faculty of Science and Technology (ST), Aarhus University ² Department of Geosciences and Natural Resource Management, Faculty of Science (IGN), University of Copenhagen ³ Danish Energy Agency ⁴ Danish Business Authority ⁵ Statistics Greenland ⁶ Umhvørvisstovan (The Environment Agency, The Faroe Islands)
Publisher:	Aarhus University, DCE – Danish Centre for Environment and Energy ©
URL:	http://dce.au.dk/en/
Year of publication:	May 2014
Editing completed:	April 2014
Financial support:	No external financial support
Please cite as:	Nielsen, O.-K., Plejdrup, M.S., Winther, M., Nielsen, M., Gyldenkaerne, S., Mikkelsen, M.H., Albrektsen, R., Thomsen, M., Hjelgaard, K., Hoffmann, L., Fauser, P., Bruun, H.G., Johannsen, V.K., Nord-Larsen, T., Vesterdal, L., Møller, I.S., Caspersen, O.H., Rasmussen, E., Petersen, S.B., Baunbæk, L. & Hansen, M.G. 2014. Denmark's National Inventory Report 2014. Emission Inventories 1990-2012 - Submitted under the United Nations Framework Convention on Climate Change and the Kyoto Protocol. Aarhus University, DCE – Danish Centre for Environment and Energy, 1214pp. Scientific Report from DCE – Danish Centre for Environment and Energy. http://dce2.au.dk/pub/SR101.pdf
	Reproduction permitted provided the source is explicitly acknowledged
Abstract:	This report is Denmark's National Inventory Report 2014. The report contains information on Denmark's emission inventories for all years' from 1990 to 2012 for CO ₂ , CH ₄ , N ₂ O, HFCs, PFCs and SF ₆ , NO _x , CO, NMVOC, SO ₂
Keywords:	Emission Inventory; UNFCCC; IPCC; CO ₂ ; CH ₄ ; N ₂ O; HFCs; PFCs; SF ₆
Layout:	Ann-Katrine Holme Christoffersen
Front page photo:	Ann-Katrine Holme Christoffersen
ISBN:	978-87-7156-071-8
ISSN (electronic):	2245-0203
Number of pages:	1214
Internet version:	The report is available in electronic format (pdf) at http://dce2.au.dk/pub/SR101.pdf

Table of Contents

List of abbreviations	11
Acknowledgements	13
1 Introduction	37
1.1 Background information on greenhouse gas inventories and climate change	37
1.2 A description of the institutional arrangement for inventory preparation	40
1.3 Brief description of the process of inventory preparation. Data collection and processing, data storage and archiving	42
1.4 Brief general description of methodologies and data sources used	44
1.5 Brief description of key categories	54
1.6 Information on QA/QC plan including verification and treatment of confidential issues where relevant	57
1.7 General uncertainty evaluation, including data on the overall uncertainty for the inventory totals	72
1.8 General assessment of the completeness	86
1.9 References	86
2 Trends in greenhouse gas emissions	90
2.1 Description and interpretation of emission trends for aggregated greenhouse gas emissions	90
2.2 Description and interpretation of emission trends by gas	91
2.3 Description and interpretation of emission trends by source	93
2.4 Description and interpretation of emission trends for indirect greenhouse gases and SO ₂	95
2.5 Description and interpretation of emission trends for KP-LULUCF inventory in aggregate, by activity and by gas	97
3 Energy (CRF sector 1)	101
3.1 Overview of the sector	101
3.2 Stationary combustion (CRF sector 1A1, 1A2 and 1A4)	104
3.3 Transport and other mobile sources (CRF sector 1A2, 1A3, 1A4 and 1A5)	200
3.4 Additional information, CRF sector 1A Fuel combustion	283
3.5 Fugitive emissions (CRF sector 1B)	285
4 Industrial Processes (CRF sector 2)	319
4.1 Overview of the sector	319
4.2 Mineral products (2A)	321
4.3 Chemical industry (2B)	331
4.4 Metal production (2C)	333
4.5 Other production (2D)	334

4.6	Production of Halocarbons and SF ₆ (2E)	335
4.7	Metal production (2C) and consumption of Halocarbons and SF ₆ (2F)	335
4.8	Other (2G)	347
4.9	Uncertainty	349
4.10	Quality assurance/quality control (QA/QC)	351
4.11	References	355
5	Solvent and Other Product Use (CRF sector 3)	358
5.1	Introduction	358
5.2	Methodology	358
5.3	Emissions, activity data and emission factors	362
5.4	Uncertainties and time series consistency	366
5.5	Quality assurance/quality control (QA/QC) and verification	369
5.6	Recalculations	373
5.7	Planned improvements	373
5.8	References	373
6	Agriculture (CRF sector 4)	376
6.1	Overview of sector	376
6.2	Data references	378
6.3	CH ₄ emission from enteric fermentation (CRF sector 4A)	383
6.4	CH ₄ and N ₂ O emission from manure management (CRF sector 4B)	391
6.5	N ₂ O emission from agricultural soils (CRF sector 4D)	400
6.6	Field burning of agricultural residues (CRF sector 4F)	410
6.7	NMVOC emission	411
6.8	Uncertainties	411
6.9	Quality assurance and quality control (QA/QC)	414
6.10	Recalculation	424
6.11	Planned improvements	425
6.12	References	426
7	LULUCF (CRF sector 5)	431
7.1	Overview of the sector	431
7.2	Forest remaining forest (5.A.1)	435
7.3	Land converted to forests	449
7.4	Cropland (5B)	455
7.5	Grassland (5C)	473
7.6	Wetlands (5D)	475
7.7	Settlements (5E)	479
7.8	Other Land	481
7.9	Direct N ₂ O emissions from N fertilization of Forest Land and Other land use – 5(I)	481
7.10	Non-CO ₂ emissions from drainage of forest soils and wetlands – 5(II)	481
7.11	N ₂ O emissions from disturbance associated with land-use conversion to cropland – 5(III)	482
7.12	CO ₂ emissions from agricultural lime application – 5(IV)	483
7.13	Biomass burning – 5(V)	485
7.14	Uncertainty, Tier 2 analysis	485

7.15	References	486
8	Waste (CRF sector 6)	489
8.1	Overview of the sector	489
8.2	Solid waste disposal on land (CRF source category 6A)	492
8.3	Wastewater handling (CRF source category 6B)	502
8.4	Waste incineration (CRF source category 6C)	510
8.5	Waste Other (CRF Source Category 6D)	516
8.6	Uncertainties and time series consistency	528
8.7	QA/QC and verification	533
8.8	Source specific recalculations	539
8.9	Source specific planned improvements	541
8.10	References	542
9	Other (CRF sector 7)	553
10	Recalculations and improvements	554
10.1	Explanations and justifications for recalculations	554
10.2	Implications for emission levels	560
10.3	Implications for emission trends, including time series consistency	560
10.4	Recalculations, including those in response to the review process, and planned improvements to the inventory (e.g. institutional arrangements, inventory preparations)	564
10.5	Explanations, justifications and implications of recalculations for KP-LULUCF inventory	573
11	KP-LULUCF	575
11.1	General information	575
11.2	Land-related information	578
11.3	Afforestation, Reforestation & Deforestation (ARD)	580
11.4	Forest Management (FM)	582
11.5	Cropland Management (CM)	583
11.6	Grazing land management (GM)	585
11.7	Article 3.3	586
11.8	Article 3.4	587
11.9	Other information	588
11.10	Information relating to Article 6	589
12	Information on accounting of Kyoto units	590
12.1	Background information	590
12.2	Summary of information reported in the SEF tables	590
12.3	Discrepancies and notifications	590
12.4	Publicly accessible information	590
12.5	Calculation of the commitment period reserve	591
12.6	KP-LULUCF accounting	591
12.7	References	592
13	Information on changes in the national system	593
14	Information on changes in the national registry	594

15	Information on the minimization of adverse impacts in accordance with Article 3, paragraph 14	597
16	Methodology applied for the greenhouse gas inventory for Greenland	598
16.1	Introduction	598
16.2	Trends in Greenhouse Gas Emissions	608
16.3	Energy (CRF sector 1)	616
16.4	Industrial processes (CRF sector 2)	642
16.5	Solvent and other product use (CRF sector 3)	657
16.6	Agriculture (CRF sector 4)	663
16.7	LULUCF (CRF sector 5)	680
16.8	Waste (CRF sector 6)	692
16.9	Other	708
16.10	Recalculations and improvements	708
16.11	KP-LULUCF	708
16.12	Annex 1 Key categories	715
16.13	Annex 2 Detailed discussion of methodology and data for estimating CO ₂ emission from fossil fuel combustion	720
16.14	Annex 3 Other detailed methodological descriptions for individual source or sink categories	720
16.15	Annex 4 CO ₂ reference approach and comparison with sectoral approach, and relevant information on the national energy balance	721
16.16	Annex 5 Assessment of completeness and (potential) sources and sinks of greenhouse gas emissions and removals excluded	721
16.17	Annex 6 Additional information to be considered as part of the annual inventory submission and the supplementary information required under Article 7, paragraph 1, of the Kyoto Protocol or other useful reference information	721
16.18	Annex 7 Tables 6.1 and 6.2 of the IPCC good practice guidance	722
16.19	Annex 8 Results of a technical analysis conducted on the Greenlandic gas oil	724
17	Information regarding the aggregated submission for Denmark and Greenland	725
17.1	Trends in emissions	725
17.2	The reference approach	729
17.3	Uncertainties	729
17.4	Key category analysis	734
17.5	Recalculations	752
17.6	Technical description of the aggregation of the emission inventories of Denmark and Greenland	756
17.7	QA/QC of the aggregated submission for Denmark and Greenland	757
	Annex 1 - Key category analysis	760
	Description of the methodology used for identifying key categories	760

Annex 2 - Detailed discussion of methodology and data for estimation of CO₂ emission from fossil fuel combustion	811
Annex 3 - Other detailed methodological descriptions for individual source or sink categories (where relevant)	812
Annex 3A - Stationary combustion	813
Annex 3A-1 Correspondence list between SNAP and CRF source categories	814
Annex 3A-2 Fuel rate	817
Annex 3A-3 Default Lower Calorific Value (LCV) of fuels and fuel correspondence list	838
Annex 3A-4 Emission factors	842
Annex 3A-5 Large point sources	873
Annex 3A-6 Adjustment of CO ₂ emission	878
Annex 3A-7 Uncertainty estimates	879
Annex 3A-8 Emission inventory 2012 based on SNAP sectors	885
Annex 3A-9 Description of the Danish energy statistics	888
Annex 3A-10 EU ETS data for coal	896
Annex 3B-10 Actual vs. representative aircraft types, no. of LTO's from Danish airports, no. of flights between Denmark and Greenland/Faroe Islands, LTO and average cruise fuel consumption and emission factors	962
Annex 3B-16 Fuel consumption and emissions in CRF format	1078
Annex 3B-17 Uncertainty estimates	1098
Annex 3C - Industrial Processes	1101
Annex 3D - Solvents and Other Product Use	1102
Annex 3D-1 Tentative list of chemicals	1103
Annex 3D-2 NMVOC and CO ₂ equivalent emissions, 1990-2011	1110
Annex 3D-3 Activity data for solvent and product use, 1990-2011	1111
Annex 3D-4 Emissions from other product use, 1990-2012	1112
Annex 3D-5 Activity data for other product use, 1990-2012	1114
Annex 3E - Agriculture	1115
Table 3E-1 Changes in housing type 1990 – 2012	1116
Table 3E-2 Number of animals allocated on subcategories for 1990-2012, 1 000 head	1119
Table 3E-3 (a-d) NH ₃ emission factors for housing units, 2012	1120
Table 3E-4 NH ₃ emission factors for storage units, 2012	1121
Table 3E-5 Parameters for winter feeding plans	1122
Table 3E-6 Energy factors used for GE	1123
Table 3E-7 Feed intake 1990-2012, FU per animal per year	1123
Table 3E-8 Grazing animals 1990 – 2012, number of days on grass per year	1123
Table 3E-9a Average gross energy intake (GE) 1990 – 2012, MJ per head per day	1123
Table 3E-9b Average gross energy intake (GE) 1990 – 2012, MJ per head per day – Subcategories for cattle and swine	1124

Table 3E-10a	VS daily excretion (average) 1990 – 2012, kg dm per head per day – CRF categories	1124
Table 3E-10b	VS daily excretion (average) 1990 – 2012, kg dm per head per day – Subcategories	1124
Table 3E-11	Basic data from Sommer et al. (2001) used to estimation of lower emission of N ₂ O and CH ₄ from biogas treated slurry	1125
Table 3E-12a	Calculation of lower CH ₄ emission as a consequence of biogas treated slurry	1126
Table 3E-12b	Calculation of lower N ₂ O emission as a consequence of biogas treated slurry	1127
Table 3E-13	Background data for calculation of N content in nitrogen fixing crops	1128
Table 3E-14	Estimated share of nitrogen fixing plants in crops	1129
Table 3E-15	Area of N-fixing crops 1990-2012, ha	1129
Table 3E-16	Background data for estimation of N ₂ O emission from crop residue 2012	1130
Table 3E-17	Area of agricultural land, 1990 – 2012, ha	1131
Table 3E-18	Harvested straw, 1990 – 2012, 1 000 tonnes	1132
Table 3E-19	QA/QC procedure, stage I – III	1134
Annex 3F - LULUCF		1135
Table 3F.1	Estimation of forest percentage and forest area	1136
Table 3F.2	Estimation of forest area with a specific characteristic	1136
Table 3F.3	Estimation of diameter-height equations	1136
Table 3F.4	Estimation of quadratic mean diameter	1137
Table 3F.5	Estimation of biomass and carbon of trees	1137
Table 3F.6	Estimation of total biomass and carbon pools	1138
Table 3F.7	Estimation of biomass and carbon with a given characteristic	1138
Table 3F.8	Estimation of biomass and carbon content of dead wood	1139
Table 3F.9	Estimation of total biomass and carbon pools of dead wood	1139
Table 3F.10	Estimation of forest floor carbon	1140
Table 3F.11	Hectares grown in the different areas of Denmark	1140
Table 3F.12	Crop yield from Statistics Denmark in 2010 distributed regions, Hhg crop ha ⁻¹	1141
Table 3F.13	Area input format for Eastern Jutland to C-TOOL in 2010. FK represent the soil type (Color Code (Farve Kode))	1142
Table 3F.14	Average annual temperatures for Denmark, °C	1143
Annex 3G - Waste		1144
Annex 3G-1	Emissions from the waste sector, 1990-2012	1145
Annex 3G-2	Solid Waste Disposal on Land, 6A	1146
Annex 3G-3	Wastewater Handling, 6B	1155
Annex 3G-4	Waste Incineration, 6C	1158
Annex 3G-5	Waste Other, 6D	1160
Annex 3G-6	Recalculations to the waste sector	1168

Annex 4 - CO₂ reference approach and comparison with sectoral approach, and relevant information on the national energy balance	1171
Annex 5 - Assessment of completeness and (potential) sources and sinks of greenhouse gas emissions and removals excluded	1172
GHG inventory	1172
KP-LULUCF inventory	1172
Annex 6 - Additional information to be considered as part of the annual inventory submission and the supplementary information required under Article 7, paragraph 1, of the Kyoto Protocol or other useful reference information	1173
Annex 7 - Tables 6.1 and 6.2 of the IPCC good practice guidance	1180
Annex 8 - Methodology applied for the greenhouse gas inventory for the Faroe Islands	1187
Introduction	1187
Trends in Greenhouse Gas Emissions	1193
Energy (CRF sector 1)	1198
Industrial Processes (CRF Sector 2)	1199
Solvents and other product use (CRF Sector 3)	1201
Agriculture (CRF Sector 4)	1201
Land Use, Land Use Change and Forestry (CRF Sector 5)	1203
Waste Sector (CRF Sector 6)	1203
Other (CRF sector 7)	1204
Recalculations and improvements	1204
Annexes	1205

[Blank page]

List of abbreviations

BAT	Best Available Techniques
CH ₄	Methane
CHP	Combined Heat and Power
CHR	Central Husbandry Register
CLRTAP	Convention on Long-Range Transboundary Air Pollution
CO	Carbon monoxide
CO ₂	Carbon dioxide
COPERT	COmputer Programme to calculate Emissions from Road Transport
CORINAIR	CORe INventory on AIR emissions
CRF	Common Reporting Format
DAAS	Danish Agricultural Advisory Service
DAFA	Danish AgriFish Agency
DCA	Danish Centre for food and Agriculture
DCE	Danish Centre for Environment and energy
DEA	Danish Energy Agency
DEPA	Danish Environmental Protection Agency
DSt	Statistics Denmark
EEA	European Environment Agency
EF	Emission Factor
EIONET	European Environment Information and Observation Network
EMEP	European Monitoring and Evaluation Programme
ENVS	Department of ENVironmental Science, Aarhus University
EU ETS	European Union Emission Trading Scheme
FSE	Full Scale Equivalent
GE	Gross Energy
GHG	Greenhouse gas
GWP	Global Warming Potential
HCB	Hexachlorobenzene
HFCs	Hydrofluorocarbons
IDA	Integrated Database model for Agricultural emissions
IEF	Implied Emission Factor
IPCC	Intergovernmental Panel on Climate Change
KCA	Key Category Analysis
LPG	Liquefied Petroleum Gas
LRTAP	Long-Range Transboundary Air Pollution
LTO	Landing and Take Off
LULUCF	Land Use, Land-Use Change and Forestry
MCF	Methane Conversion Factor
MSW	Municipal Solid Waste
N ₂ O	Nitrous oxide
NFI	National Forest Inventory
NFR	Nomenclature For Reporting
NH ₃	Ammonia
NIR	National Inventory Report
NMVOC	Non-Methane Volatile Organic Compounds
NO _x	Nitrogen Oxides
PFCs	Perfluorocarbons
QA	Quality Assurance
QC	Quality Control
SCR	Selective Catalytic Reduction

SF ₆	Sulphur hexafluoride
SNAP	Selected Nomenclature for Air Pollution
SO ₂	Sulphur dioxide
SWDS	Solid Waste Disposal Sites
UNECE	United Nations Economic Commission for Europe
UNFCCC	United Nations Framework Convention on Climate Change
VS	Volatile Solids
WWTP	WasteWater Treatment Plant

Acknowledgements

The work of compiling the Danish greenhouse gas inventory requires the input of many individuals, companies and institutions. The authors of this report would in particular like to thank the following for their valuable input in the work process:

- The Danish Energy Agency, in particular Jane Rusbjerg, Ali A. Zarnaghi, Kaj Stærkind and Dorte Maimann for valuable discussions concerning energy balance data and EU ETS data.
- DTU Transport (Technical University of Denmark), in particular Thomas Jensen for valuable input and discussions on road transport fleet and mileage characterisation.
- Anette Holst, Statoil Refining Denmark A/S, for providing detailed data and information on calorific values and uncertainties related to processes at the refinery.
- Lis R. Rasmussen, A/S Danish Shell, Shell Refinery, for providing detailed data on emissions from the refinery.
- Andreas B. Jensen, Aalborg Supply - Gas, for providing information on town gas distribution.
- Torben Ahlmann-Laursen, Aalborg Portland, for providing information on production statistics.
- DCA - Danish Centre for Food and Agriculture, Aarhus University, in particular Hanne D. Poulsen for valuable input and discussions on N-excretions and ammonia emission factors for livestock production, Søren O. Petersen for discussion on emission factors from manure storages, organic soils and wetlands and Mogens H. Greve for valuable contributions to mapping of organic soils.
- Torkild Birkmose, AgroTech, Institute for Agri Technology and Food Innovation, for discussions on actual farming practice and ammonia emissions factors from manure management.
- The Danish AgriFish Agency for providing unrestricted access to all agricultural data.
- The European Environment Agency for granting permission for Denmark to use the CRF Aggregator tool for the submissions under the Kyoto Protocol and the UNFCCC.
- Carl Christian Hoffmann, Department of Bioscience, Aarhus University for valuable discussions on organic soils and wetlands.
- The IT section at Aarhus University for supporting the emission inventory data system.

Executive summary

ES.1 Background information on greenhouse gas inventories and climate change

ES.1.1 Reporting

This report is Denmark's National Inventory Report (NIR) 2014 for submission to the United Nations Framework Convention on Climate Change and the Kyoto Protocol, due April 15, 2014. The report contains detailed information about Denmark's inventories for all years from 1990 to 2012. The structure of the report is in accordance with the UNFCCC guidelines on reporting and review. The main difference between Denmark's NIR 2014 report to the European Commission, due March 15, 2014, and this report to UNFCCC is reporting of territories. The NIR 2014 to the EU Commission was for Denmark, while this NIR 2014 to the UNFCCC is for Denmark, Greenland and the Faroe Islands. The suggested outline provided by the UNFCCC secretariat has been followed to include the necessary information under the Kyoto Protocol. The report includes detailed and complete information on the inventories for all years from year 1990 to the year 2012, in order to ensure transparency.

The annual emission inventories for the years from 1990 to 2012 are reported in the Common Reporting Format (CRF). Within this submission separate CRF's are available for Denmark (EU), Greenland, the Faroe Islands, for Denmark and Greenland (KP) as well as for Denmark, Greenland and the Faroe Islands (UNFCCC). The CRF spreadsheets contain data on emissions, activity data and implied emission factors for each year. Emission trends are given for each greenhouse gas and for total greenhouse gas emissions in CO₂ equivalents.

The issues addressed in this report are: Trends in greenhouse gas emissions, description of each emission category of the CRF, uncertainty estimates, explanations on recalculations, planned improvements and procedure for quality assurance and control. The information presented in Chapters 2-9 and Chapter 11 refers to Denmark (EU) only. Specific information regarding the submission of Greenland and the Faroe Islands is included in Chapter 16 and Annex 8, respectively. Chapter 17 contains information (e.g. on trends, uncertainties and key category analysis) on the aggregated submission of Denmark and Greenland under the Kyoto Protocol.

This report itself does not contain the full set of CRF tables. The full set of CRF tables is available at the EIONET, Central Data Repository, kept by the European Environmental Agency:

http://cdr.eionet.europa.eu/dk/Air_Emission_Inventories

In the report English notation is used: "." (full stop) for decimal sign and mostly space for division of thousands. The English notation for division of thousand as "," (comma) is not used due to the risk of being misinterpreted by Danish readers.

ES.1.2 Institutions responsible

On behalf of the Ministry of the Environment and the Ministry of Climate, Energy and Building, the Danish Centre for Environment and Energy (DCE),

Aarhus University, is responsible for the calculation and reporting of the Danish national emission inventory to EU and the UNFCCC (United Nations Framework Convention on Climate Change) and UNECE CLRTAP (Convention on Long Range Transboundary Air Pollution) conventions. Hence, DCE prepares and publishes the annual submission for Denmark to the EU and UNFCCC of the National Inventory Report and the greenhouse gas (GHG) inventories in the Common Reporting Format, in accordance with the UNFCCC guidelines. Further, DCE is responsible for reporting the national inventory for the Kingdom of Denmark to the UNFCCC. DCE is also the body designated with overall responsibility for the national inventory under the Kyoto Protocol for Greenland and Denmark. Furthermore, DCE participates when reporting issues are discussed in the regime of UNFCCC and EU (Monitoring Mechanism).

The work concerning the annual greenhouse gas emission inventory is carried out in cooperation with Danish ministries, research institutes, organisations and companies. The Government of Greenland is responsible for finalising and transferring the inventory for Greenland to DCE. The Faroe Islands Environmental Agency is responsible for finalising and transferring the inventory for the Faroe Islands to DCE.

ES.1.3 Greenhouse gases

The greenhouse gases reported are those under the UN Climate Convention:

- Carbon dioxide CO_2
- Methane CH_4
- Nitrous oxide N_2O
- Hydrofluorocarbons HFCs
- Perfluorocarbons PFCs
- Sulphur hexafluoride SF_6

The global warming potential (GWP) for various greenhouse gases has been defined as the warming effect over a given time frame of a given weight of a specific substance relative to the same weight of CO_2 . The purpose of this measure is to be able to compare and integrate the effects of the individual greenhouse gases on the global climate. Typical lifetimes in the atmosphere of greenhouse gases are very different, e.g. approximately 12 and 120 years for CH_4 and N_2O , respectively. So the time perspective clearly plays a decisive role. The life frame chosen is typically 100 years. The effect of the various greenhouse gases can then be converted into the equivalent quantity of CO_2 , i.e. the quantity of CO_2 giving the same effect in absorbing solar radiation. According to the IPCC and their Second Assessment Report, which UNFCCC has decided to use as reference, the global warming potentials for a 100-year time horizon are:

- Carbon dioxide (CO_2): 1
- Methane (CH_4): 21
- Nitrous oxide (N_2O): 310

Based on weight and a 100-year period, CH_4 is thus 21 times more powerful a greenhouse gas than CO_2 and N_2O is 310 times more powerful than CO_2 . Some of the other greenhouse gases (hydrofluorocarbons, perfluorocarbons and sulphur hexafluoride) have considerably higher global warming potentials. For example, sulphur hexafluoride has a global warming potential of 23 900. The values for global warming potential used in this report are those

prescribed by UNFCCC. The indirect greenhouse gases reported are nitrogen oxides (NO_x), carbon monoxide (CO), non-methane volatile organic compounds (NMVOC) and sulphur dioxide (SO₂). Since no GWPs are assigned to these gases, they do not contribute to GHG emissions in CO₂ equivalents.

ES.2 Summary of national emission and removal trends

Summary ES.2-4 refers to the inventory for Denmark only. The inventories for Greenland, Denmark and Greenland and the Faroe islands are described in Chapter 16 and 17 and Annex 8, respectively.

ES.2.1 Greenhouse gas emissions inventory

The greenhouse gas emissions are estimated according to the IPCC guidelines and guidance and are aggregated into seven main sectors. According to decisions made under the UNFCCC and the Kyoto Protocol the greenhouse gas emissions are estimated according to the IPCC 1996 guidelines and the IPCC 2000 good practice guidance. The greenhouse gases include CO₂, CH₄, N₂O, HFCs, PFCs and SF₆. Figure ES.1 shows the estimated total greenhouse gas emissions in CO₂ equivalents from 1990 to 2012. The emissions are not corrected for electricity trade or temperature variations. CO₂ is the most important greenhouse gas contributing in 2012 to national total in CO₂ equivalents excluding LULUCF (Land Use and Land Use Change and Forestry) with 76.3 % followed by N₂O with 11.6 %, CH₄ 10.7 % and F-gases (HFCs, PFCs and SF₆) with 1.5 %. Seen over the time series from 1990 to 2012 these percentages have been increasing for CH₄ and F-gases and decreasing for N₂O. The percentages for CO₂ show larger fluctuations during the time series. Stationary combustion plants, Transport and Agriculture represent the largest contributing categories to emissions of greenhouse gases, followed by Industrial processes, Waste, Fugitive emissions and Solvents, see Figure ES.1. The net CO₂ uptake by LULUCF in 2012 is 0.2 % of the total emission in CO₂ equivalents excl. LULUCF. The national total greenhouse gas emission in CO₂ equivalents excluding LULUCF has decreased by 24.8 % from 1990 to 2012 and 30.5 % including LULUCF. Comments to the overall trends for the individual greenhouse gases etc. seen in Figure ES.1 are given in the sections below.




Figure ES.1 Greenhouse gas emissions in CO₂ equivalents distributed on main sectors (excl. LULUCF) for 2012 and time series for 1990 to 2012, Where data are given with or without LULUCF.

ES.2.2 KP-LULUCF activities

Net emissions from Afforestation, Reforestation and Deforestation (ARD) activities in 2012 were 148 Gg CO₂ equivalents, hereof 0.9 Gg CO₂ equivalents owe to N₂O emissions from disturbance of soils. Net removals from Forest

Matter (FM) were 4 479.6 Gg CO₂ equivalents (Table ES.1) hereof 12.1 Gg CO₂ equivalents owe to N₂O emissions from drainage of soils.

For Cropland Management (CM) the net emissions in 2012 were 2 958 Gg CO₂ equivalents compared to a net emission in 1990 of 4 845 Gg CO₂ equivalents.

For Grassland Management (GM) the net emissions in 2012 were 523 Gg CO₂ equivalents compared to a net emission in 1990 of 177 Gg CO₂ equivalents.

Table ES.1 Emissions and removals in 2012 for activities relating to Article 3.3 and Article 3.4.

	Net CO ₂ emissions/ removals	CH ₄	N ₂ O	Net CO ₂ equivalent emissions/ removals (Gg)
A. Article 3.3 activities				107,38
A.1. Afforestation and Reforestation	76,62	NO	IE,NA,NO	76,62
A.1.1. Units of land not harvested since the beginning of the commitment period	76,62	NO	IE,NA,NO	76,62
A.1.2. Units of land harvested since the beginning of the commitment period	IE,NO	NO	IE,NO	IE,NO
A.2. Deforestation	30,50	NO	0,00	30,76
B. Article 3.4 activities				5.088,29
B.1. Forest Management	49,60	0,03	0,05	66,24
B.2. Cropland Management	4.844,66	NO	NA,NO	4.844,66
B.3. Grazing Land Management	177,39	0,00	0,00	177,39
B.4. Re-vegetation	NA	NA	NA	NA

ES.3 Overview of source and sink category emission estimates and trends

ES.3.1 Greenhouse gas emissions inventory

Energy

The largest source of CO₂ emission is the energy sector, which includes the combustion of fossil fuels such as oil, coal and natural gas.

The emission of CO₂ from Energy Industries has decreased by 36.8 % from 1990 to 2012. The relatively large fluctuation in the emission is due to inter-country electricity trade. Thus, the high emissions in 1991, 1994, 1996, 2003 and 2006 reflect a large electricity export and the low emissions in 1990, 1992 and 2005, 2008, 2011 and 2012 are due to a large import of electricity. The main reason for this decrease owe to decreasing fuel consumption, mainly for coal and natural gas. This decrease is partly due to increasing import of electricity and partly to increasing production of wind power and other renewable energy sources.

The increasing emission of CH₄ during the nineties is due to the increasing use of gas engines in decentralised cogeneration plants. The CH₄ emissions from this sector have been decreasing from 2001 to 2012 due to the liberalisa-

tion of the electricity market. The CO₂ emission from the transport sector increased by 14.0 % from 1990 to 2012, mainly due to increasing road traffic.

Industrial processes

The GHG emissions from industrial processes, i.e. emissions from processes other than fuel combustion, amount in 2012 to 2.6 % of the total emission in CO₂ equivalents (excl. LULUCF). The main sources are cement production, refrigeration, foam blowing and calcination of limestone. The CO₂ emission from cement production – which is the largest source contributing in 2012 with 1.7 % of the national total – decreased by 1.3 % from 1990 to 2012. The second largest source has previously been N₂O from the production of nitric acid. However, the production of nitric acid/fertiliser ceased in 2004 and therefore the emission of N₂O also ceased.

The emission of HFCs, PFCs and SF₆ has increased by 140.7 % from 1995 until 2012, largely due to the increasing emission of HFCs. The use of HFCs, and especially HFC-134a, has increased several fold and thus HFCs have become the dominant F-gases, contributing 67 % to the F-gas total in 1995, rising to 83.9 % in 2012. HFC-134a is mainly used as a refrigerant. However, the use of HFC-134a is now stabilising. This is due to Danish legislation, which in 2007 banned new HFC-based refrigerant stationary systems. However, in contrast to this trend is the increasing use of air conditioning systems in mobile systems.

Solvent and other product use

The use of solvents in industries and households and other product use contribute 0.3 % of the total greenhouse gas emissions in CO₂ equivalents. There is a 34.6 % decrease in greenhouse gas emissions from solvent and other product use from 1990 to 2012. In 2012, N₂O comprises 11.5 % of the total CO₂ equivalent emissions for solvent and other product use.

Agriculture

The agricultural sector contributes in 2012 with 18.6 % of the total greenhouse gas emission in CO₂ equivalents (excl. LULUCF) and is the most important sector regarding the emissions of N₂O and CH₄. In 2012, the contribution of N₂O and CH₄ to the total emission of these gases was 90.5 % and 76.6 %, respectively. The N₂O emission from agriculture decreased by 34.9 % from 1990 to 2012. The main reason for the decrease is a legislative demand for an improved utilisation of nitrogen in manure. This result in less nitrogen excreted per livestock unit produced and a considerable reduction in the use of fertilisers. From 1990 to 2012, the emission of CH₄ from enteric fermentation has decreased due to decreasing numbers of cattle. However, the emission from manure management has increased due to changes in stable management systems towards an increase in slurry-based systems. Altogether, the emission of CH₄ for the agricultural sector has decreased by 0.7 % from 1990 to 2012.

Land Use and Land Use Change and Forestry (LULUCF)

The LULUCF sector alters between being a net sink and a net source of GHG. In 2012 LULUCF was a net sink with 1.6 % of the total GHG emission excluding LULUCF. In 2011 LULUCF was a net sink equivalent to 4.9 % of the total GHG emission (excluding LULUCF). The overall trend in the LULUCF sector without Forestry is a decrease of 30 % since 1990.

In 2012 Forest Land was a large sink of 4 441 CO₂ equivalents, while Cropland, Grassland, Wetlands and Settlements was net sources contributing with 2 956 Gg CO₂ equivalents, 554 Gg CO₂ equivalents, 2 Gg CO₂ equivalents and 91 Gg CO₂ equivalents, respectively.

Waste

The waste sector contributes in 2012 with 2.1 % to the national total of greenhouse gas emissions (excl. LULUCF), 15.7 % of the total CH₄ emission and 3.4 % of the total N₂O emission. The sector comprises solid waste disposal on land, wastewater handling, waste incineration without energy recovery (e.g. incineration of animal carcasses) and other waste (e.g. composting and accidental fires).

The GHG emission from the sector has decreased by 32.4 % from 1990 to 2012. This decrease is a result of (1) a decrease in the CH₄ emission from solid waste disposal sites (SWDS) by 48.9 % due to the increasing use of waste for power and heat production, and (2) a decrease in emission of N₂O from wastewater (WW) handling systems of 30.6 % due to upgrading of WW treatment plants. These decreases are counteracted by an increase in CH₄ from WW of 13.0 % due to increasing industrial load to WW systems. In 2012 the contribution of CH₄ from SWDS was 12.7 % of the total CH₄ emission. The CH₄ emission from WW amounts in 2012 to 1.3 % of the total CH₄ emissions. The emission of N₂O from WW in 2012 is 1.2 % of national total of N₂O. Since all incinerated waste is used for power and heat production, the emissions are included in the 1A CRF category.

ES.3.2 KP-LULUCF activities

In 2012 the activities under Article 3.3 was a net source of 148 Gg CO₂ equivalents and the activities under Article 3.4 was a net sink of 998 Gg CO₂ equivalents. A short overview of KP-LULUCF is given in Chapter ES.2.2 and a more detailed description is given in Chapter 11.

ES.4 Other information

ES.4.1 Quality assurance and quality control

A plan for Quality Assurance (QA) and Quality Control (QC) in greenhouse gas emission inventories is included in the report. The plan is in accordance with the guidelines provided by the UNFCCC (Good Practice Guidance and Uncertainty Management in National Greenhouse Gas Inventories and Guidelines for National Systems). ISO 9000 standards are also used as an important input for the plan.

The plan comprises a framework for documenting and reporting emissions in a way that emphasize transparency, consistency, comparability, completeness and accuracy. To fulfil these high criteria, the data structure describes the pathway, from the collection of raw data to data compilation and modelling and finally reporting.

As part of the Quality Assurance (QA) activities, emission inventory sector reports are being prepared and sent for review to national experts not involved in the inventory development. To date, the reviews have been completed for the stationary combustion plants sector, the fugitive emissions from fuels sector, the transport sector, the solvents and other product use sector and the agricultural sector. In order to evaluate the Danish emission

inventories, a project where emission levels and emission factors are compared with those in other countries has been conducted.

ES.4.2 Completeness

The Danish greenhouse gas emission inventories include all sources identified by the revised IPPC guidelines.

Please see Annex 5 for more information.

ES.4.3 Recalculations and improvements

The main improvements of the inventories are:

Energy

Stationary Combustion

For stationary combustion plants, the emission estimates for the years 1990-2011 have been updated according to the latest energy statistics published by the Danish Energy Agency. The update included both end use and transformation sectors as well as a source category update. The changes in the energy statistics are largest for the years 2009, 2010 and 2011.

For CO₂ the largest recalculation is in source category Manufacturing industries and constructions. The recalculation is related to liquid fuels and is a result of correction of an error. The consumption of residual oil was underestimated in the former inventories. The CO₂ emission from liquid fuels applied in manufacturing industries and construction for 2011 is 7% higher in the 2014 reporting than in the 2013 reporting.

The CH₄ emission from residential wood combustion has been recalculated based on improved emission factors for stoves. This has caused a 16 % increase of the CH₄ emission reported for biomass fuels in residential plants for 2011.

For N₂O the largest recalculation is in source category Manufacturing industries and constructions. This recalculation is also related to the former underestimate for residual oil. The N₂O emission from liquid fuels applied in manufacturing industries and constructions for 2011 is 13 % higher in the 2014 reporting than in the 2013 reporting.

Mobile sources

The following recalculations and improvements of the emission inventories have been made since the emission reporting in 2013.

Road transport

Based on the updated version of COPERT IV launched in 2013, new vehicle sub categories have been introduced in the emission inventories for mopeds and passenger cars. For mopeds a division is now made between 2-stroke and 4-stroke engine technologies and for passenger cars small engine sizes below 0.8 l. for gasoline and below 1.4 l. for diesel have been included. Also NO_x emission factors for euro 5 diesel passenger cars have been updated in the model based on the new COPERT IV version.

Small errors in input gasoline fuel consumption for the years 2009-2011 and for input diesel fuel consumption in the years 2010-2011 have been corrected.

The percentage emission change interval and year of largest percentage differences (low %; high %, year) for the different emission components are: CO₂ (-0.5 %; -0.05 %, 2008), CH₄ (-0.2 %; 2.4 %, 2011) and N₂O (-0.4 %; 0.3 %, 2008).

Navigation

Minor changes in ferry input data has been made for the years 2008-2011 causing minor emission changes for domestic navigation. The following largest percentage differences (in brackets) for domestic navigation are noted for: CO₂ (-1.0 %), CH₄ (-0.8 %) and N₂O (-1.1 %).

Agriculture/forestry/fisheries

The number and engine size of machine pool tractors has been updated for the years 2007-2011. The number of ATV's has been changed for the years 2009-2011. Errors in the fuel consumption for fisheries in 2000, 2010 and 2011 have been corrected.

In 2000 the following percentage differences (in brackets) for agriculture/forestry/fisheries are noted for: CO₂ (9.1 %), CH₄ (3.7 %) and N₂O (11.9 %) due to fuel consumption changes in fisheries.

For other years than 2000, the following largest percentage differences (in brackets) are noted for: CO₂ (-3.7 %), CH₄ (5 %) and N₂O (-4.9 %).

Industry

The number of mini loaders has been updated for the years 2004-2011.

The following largest percentage differences (in brackets) for industrial non road machinery are noted for: CO₂ (1.0 %), CH₄ (1.6 %) and N₂O (1.6 %).

Civil aviation

An error in the CH₄ emission factor has been corrected for the years 1985-2000. The emission factors are now in line with the factors proposed by the EMEP/EEA emission inventory guidebook. The CH₄ emission percentage differences are between -31 % and -42 %.

Military

Emission factors derived from the new road transport simulations have caused some emission changes from 1985-2010. The following largest percentage differences (in brackets) for military are noted for: CO₂ (0 %), CH₄ (0.6 %) and N₂O (0.2 %).

Fugitive emissions

In the emission inventory reported in 2014 for the years 1990-2012 the following recalculations regarding fugitive emissions from fuels have been applied:

Natural gas transmission and distribution

Activity data and IEF for the time series 1990-2011 has been updated for transmission and distribution according to annual environmental reports and the latest national energy statistics, respectively. Further, the CH₄ EFs are updated for one town gas distribution company following an update of the estimated fugitive losses per distribution. The recalculation has changed the CO₂ emission by 0.01 ktonnes and CH₄ emission by (-0.09) - 0.07 ktonnes,

corresponding to < 0.003% and (-2) % - 2% of the total fugitive CO₂ and CH₄ emission.

Venting

EFs for CH₄ have been added for the years 1990-1993 for one gas storage plant. In these years the plant is treated as an area source in the national system, while it is treated separately as a point source in the following years. EFs are based on data from annual reports for 1995-1999, as no data are available for the years 1990-1994. Further, a minor error has been applied for venting in 2011, according to the annual report from one of the natural gas storage facilities. The recalculation has changed the CH₄ emission by 0.06 ktonnes, corresponding to 2% of the total fugitive CH₄ emission.

Flaring in gas storage and treatment plants

EFs for CO₂ and CH₄ have been added for the years 1990-1993 for the gas treatment plant. In these years the plant is treated as an area source in the national system, while it is treated separately as a LPS in the following years. EFs are based on data from annual reports for 1995-1999, as no data are available for the years 1990-1994. The recalculation has changed the CO₂ emission by 2.2 ktonnes and the CH₄ emission by 0.01 ktonnes, corresponding to 0.4 % and 0.4 % of the total fugitive CO₂ and CH₄ emission.

Flaring in refineries

The CO₂ EF for one refinery has been updated for the years 1994-2006 and now reflects the average EF from the first five EU-ETS reports (2007-2011). The NO_x EF for two refineries has been changed to the standard EF in the EMEP/EEA Guidebook, as references to the previously applied EFs are outdated or not existing.

The CO₂ EF for a refinery that was shut down in 1996 has been changed for the years 1990-1996 to match the existing refineries, as no better data is available. CO₂ EF for 2010-2011 has been updated and now corresponds to the EU-ETS reporting. The recalculation has changed the CO₂ emission by (-0.05) - 0.49 ktonnes and CH₄ emission by (-0.06) ktonnes, corresponding to (-0.01) - 0.1% and (-2) % of the total fugitive CO₂ and CH₄ emission.

Industrial Processes

Lime production

EU-ETS data have been implemented for one lime production plant for 2011 leading to a minor decrease in the overall emission of CO₂ emission by 5.51 ktonnes. This change in methodology will be implemented for the previous years in the next submission.

Limestone and Dolomite Use

Activity data for flue gas cleaning have been changed for three power and waste incineration plants in 2011. Consumption of CaCO₃ has increased with 236 tonnes and CaCO₃ containing residues have increased with 669 tonnes resulting in an increased CO₂ emission of 162 tonnes.

An error in transferring data from the emission database to CRF Reporter has been corrected for 1995-2011 for two individual plants.

Chemical industry

The process emission has been adjusted to reflect the total emission reported in environmental reports minus the energy related emissions reported to EU-ETS.

Consumption of Halocarbons and SF₆ – SF₆ – Other

A minor correction has been made to emission of SF₆ from double glazed windows for 2010.

Potential emissions revised for 2005-7 to reflect the fact that potential emission of SF₆ is the same as yearly consumption. Consumption of SF₆ for double glaze windows stopped in 2001.

Solvents and Other Product Use

Recalculations have been made for Other Product Use, where changes were made for the activity data of all four emission sources; candles (2009-2011), fireworks (2009-2011), tobacco (1980-1999, 2011) and charcoal used for barbeques (1980-1987, 2009-2011). These changes have caused recalculations for NMVOC, N₂O and CO₂. NMVOC emissions for the years 1980-1999 have increased between 6.0 % (1997) and 9.1 % (1982) and decreased for 2009-2011 with 2.1 % (2009) to 5.7 % (2011). CO₂ emissions have decreased for the years 2009 (13.4 %) and 2010 (6.1 %) and increased for 2011 (0.7 %), no or minor recalculations were performed for 1980-2008. N₂O emissions have increased for 1980-1999 with between 0.5 % (1999) and 2.8 % (1982) and decreased for the years 2009-2011 with between 0.3 % (2011) and 1.0 % (2009).

Agriculture

Some changes of emissions from the agricultural sector have taken place. These changes reflect decreased emissions in the years 1990-2011 up to 0.2 % compared to the total CO₂ equivalent emission from the agricultural sector. The decrease in 1990-2011 is due to a decrease in the emissions of both N₂O and CH₄.

The CH₄ emission decreases both for emission from enteric fermentation and manure management. The number of geese has been changed for all years and the number of weaners and fattening pigs has been changed for 2011, this affects both emissions from enteric fermentation and manure management. The amount of straw used for bedding for heifers has been changed for 1990-2002 and the amount of biogas treated manure in 2010 has been changed. This affects the emission of CH₄ from manure management.

For the N₂O emission a range of changes have been made, which have both increasing and decreasing effect. EF for NH₃ from synthetic fertiliser has been changed for all years and this affects the emission of N₂O. The emission of N₂O from atmospheric deposition increases due to the change for NH₃ from synthetic fertiliser while the emission of N₂O from synthetic fertiliser decreases. Change in the number of geese decrease the emission of N₂O from grazing for all years. The change of amount of biogas treated manure decreases the N₂O emission in 2010. The emission from manure, manure on soil and leaching is decreased in 2011 due to updated numbers of weaners and fattening pigs.

LULUCF

An update of the LULUCF matrix has taken place for all years. The update was necessary because there were errors in the maps received last year from

the Danish Geodata Agency. The problems were especially allocated to a misclassification of recreational areas and a misclassification of parks inside some cities as being forests.

The forest area has been decreased slightly in 2011 (app. 0.3%) compared to the submission last year. This has only had a very limited effect on the total carbon stock in the Danish forest as the carbon stock is estimated in the National Forest Inventory, which is independent of the LULUCF matrix. Furthermore has some unclassified areas inside and around the cities which previous was classified as grassland now been removed to Settlement.

Cropland, grassland, wetlands and settlements

As the land use matrix is slightly changed the emissions from land use conversion for all sectors are changed slightly for the whole time series. These changes have no effect on the emissions from agricultural soils as these are based information from the EU Land Parcel Information System, i.e. the actual land use. Two minor technical errors have been found in the accounting estimate and corrected: living biomass in Settlements and the area accounted for in Cropland Management and Grazing Land Management under article 3.4. These errors have only a small impact on the inventory.

Waste

For the category 6A SWDS, an in depth disaggregation of deposited waste for the years 2010, 2011 and 2012 have been performed based on the new waste reporting system in Denmark. 18 categories have been identified of which eleven have been evaluated as inert waste. The overall results of this detailed characterisation and reallocation of the deposited waste results in a decrease in the CH₄ emission in 1990 of 7.6% and an increase in the CH₄ emission in 2011 of 4.6%.

For the category 6B wastewater handling, no recalculations were made for N₂O. For the methane emissions, the methane correction factor was decreased from 1 to 0.8, which is in accordance to the IPCC guidelines 2006, and which have been further justified by plant specific data. Besides the correction of MCF, smaller corrections in plant inlet TOW data have occurred corresponding to a change below 1% throughout the time series.

There are no recalculations in the waste incineration category, however; a correction in the rounding of decimals has caused an increase of 0.07 % of the N₂O emissions from animal cremation for all years 1990-2011.

For the category waste other; changes were made in the vehicle fires and composting source categories. For vehicle fires; the time series for vehicle population has been updated along with the estimated average weights of some vehicle types, the result is a decrease of CH₄ and CO₂ emissions from 1990-2006 and an increase for 2007-2011. Emission factors have been updated for composting of sludge and organic municipal waste which have resulted in increased emissions of CH₄ and N₂O. The joint effect of these recalculations is a decrease in CH₄ emissions from 1980 (2.4 %) to 1984 (2.0 %) and an increase from 1985 (16.7 %) to 2011 (275 %), for CO₂ the emissions have decreased from 1980 (1.1 %) to 2006 (0.004 %) and increased from 2007 (0.1 %) to 2011 (0.2 %), finally the N₂O emissions have increased for all years between 1985 (1.1 %) and 2011 (238 %).

KP-LULUCF

A recalculation for KP-LULUCF has been performed for all areas as a consequence of the new land area matrix.

Sammenfatning

S.1 Baggrund for opgørelse af drivhusgasemissioner og klimacændringer

S.1.1 Rapporteringen

Denne rapport er Danmarks årlige rapport – den såkaldte Nationale Inventory Report (NIR) for 2014. Rapporten beskriver drivhusgasopgørelsen som blev fremsendt til FN's konvention om klimacændringer (UNFCCC) og Kyoto-protokollen den 15. april 2014. Rapporten indeholder detaljerede informationer om Danmarks drivhusgasudslip for alle år fra 1990 til 2012. Rapportens struktur er i overensstemmelse med UNFCCC's retningslinjer for rapportering og review. Hovedforskellen mellem Danmarks NIR 2014 som blev fremsendt til EU-Kommissionen til den 15. marts 2014, og denne rapport til UNFCCC vedrører det territorium rapporteringen omfatter. NIR 2014 til EU-Kommissionen var for Danmark, mens NIR 2014 til UNFCCC er for Danmark, Grønland og Færøerne. For at sikre at opgørelserne er sammenhængende og gennemskelige indeholder rapporten detaljerede oplysninger om opgørelsesmetoder og baggrundsdata for alle årene fra 1990 og til 2012.

Denne emissionsopgørelse for årene 1990 til 2012, er som tidligere årlige opgørelser, rapporteret i formatet Common Reporting Format (CRF) som Klimakonventionen foreskriver anvendt. Emissionsopgørelsen i CRF foreligger med denne rapportering således, at der er separate CRF for Danmark (EU), Grønland, Færøerne, for Danmark og Grønland (KP) samt for Danmark, Grønland og Færøerne (Klimakonventionen). CRF-tabellerne indeholder oplysninger om emissioner, aktivitetsdata og emissionsfaktorer for hvert år, emissionsudvikling for de enkelte drivhusgasser samt den totale drivhusgasemission i CO₂-ækvivalenter.

Følgende emner er beskrevet i rapporten: Udviklingen i drivhusgasemissionerne, metoder mv. som anvendes til opgørelserne i de emissionskategorier som findes i CRF-formatet, usikkerheder, genberegninger, planlagte forbedringer og procedure for kvalitetssikring og -kontrol. Teksten i kapitel 2-9 og kapitel 11 omhandler kun Danmark som omfattet af EU. Oplysninger om emissionsopgørelsen for Grønland og Færøerne er inkluderet i henholdsvis kapitel 16 og annex 8. Kapitel 17 indeholder informationer (f.eks. om udviklingen i emissioner over tid, usikkerheder og identifikation af nøglekategorier) for den samlede aflevering for Danmark og Grønland under Kyoto-protokollen.

Denne rapport indeholder ikke det fulde sæt af CRF-tabeller. Det fulde sæt af CRF-tabeller er tilgængelige på EIONET, som er det Europæiske Miljøagenturs rapporterings-internetsite:

http://cdr.eionet.europa.eu/dk/Air_Emission_Inventories

Med hensyn til gengivelsen af tal i CRF-formatet, gøres opmærksom på at det er med dansk notation: “,” (komma) for decimaladskillelse og “.” (punktum) til adskillelse af tusinder. I rapporten er den engelske notation brugt: “.” (punktum) for decimaltegn og for det meste mellemrum for adskillelse af tusinder. Den engelske notation for adskillelse af tusinder med “,” (komma) er for det meste ikke brugt på grund af risikoen for fejltolkninger for danske læsere.

S.1.2 Ansvarlige institutioner

DCE - Nationalt Center for Miljø og Energi ved Aarhus Universitet er på vegne af Miljøministeriet samt Klima-, Energi- og Bygningsministeriet ansvarlig for udregning og afrapportering af den nationale emissionsopgørelse til EU og til UNFCCC (FN's konvention om klimaændringer) såvel som til UNECE-konventionen om langtransporteret grænseoverskridende luftforurening. Som følge heraf er DCE ansvarlig for udførelse og publicering af opgørelserne af drivhusgasemissioner og den årlige rapportering til EU og UNFCCC for Danmark. DCE er den centrale institution for Danmarks nationale system til drivhusgasopgørelser under Kyotoprotokollen. Ydermere er DCE ansvarlig for rapportering af drivhusgasemissionsopgørelser til Klimakonventionen for Kongeriget Danmark (Færøerne, Grønland og Danmark), samt Danmarks og Grønlands samlede rapportering til Kyotoprotokollen. DCE deltager desuden i arbejdet i regi af Klimakonventionen og Kyotoprotokollen, hvor retningslinjer for rapportering diskuteres og vedtages og i EU's monitoringsmekanisme for opgørelse af drivhusgasser, hvor retningslinjer for rapportering til EU reguleres.

Arbejdet med de årlige opgørelser udføres i samarbejde med andre danske ministerier, forskningsinstitutioner, organisationer og private virksomheder. Grønlands Klima- og Infrastrukturstyrelse er ansvarlig for levering af opgørelser for Grønland til DCE. Færøernes miljømyndighed (Umhvørvisstovan) er ansvarlig for de færøske opgørelser.

S.1.3 Drivhusgasser

Til Klimakonventionen rapporteres følgende drivhusgasser:

- | | |
|----------------------|------------------|
| • Kuldioxid | CO ₂ |
| • Metan | CH ₄ |
| • Lattergas | N ₂ O |
| • Hydrofluorcarboner | HFC'er |
| • Perfluorcarboner | PFC'er |
| • Svovlhexafluorid | SF ₆ |

Det globale opvarmningspotentiale, på engelsk Global Warming Potential (GWP), udtrykker klimapåvirkningen over en nærmere angivet tid af en vægtenhed af en given drivhusgas relativt til samme vægtenhed af CO₂. Drivhusgasser har forskellige karakteristiske levetider i atmosfæren, således for CH₄ ca. 12 år og for N₂O ca. 120 år. Derfor spiller tidshorisonten en afgørende rolle for størrelsen af GWP. Typisk vælges 100 år. Herefter kan effekten af de forskellige drivhusgasser omregnes til en ækvivalent mængde CO₂, dvs. til den mængde CO₂ der vil give samme klimapåvirkning. Til rapporteringen til Klimakonventionen er vedtaget at anvende GWP-værdier for en 100-årig tidshorisont, som ifølge IPCC's anden vurderingsrapport er:

- | | |
|--------------------------------|-----|
| • Kuldioxid, CO ₂ : | 1 |
| • Metan, CH ₄ : | 21 |
| • Lattergas, N ₂ O: | 310 |

Regnet efter vægt og over en 100-årig periode er metan således ca. 21 og lattergas ca. 310 gange så effektive drivhusgasser som kuldioxid. For andre drivhusgasser der indgår i rapporteringen, de såkaldte F-gasser (HFC, PFC, SF₆) findes væsentlig højere GWP-værdier. Under Klimakonventionen er der ligeledes vedtaget GWP-værdier for disse baseret på IPCC's anbefalinger.

Således har f.eks. SF₆ en GWP-værdi på 23 900. I denne rapport anvendes de GWP-værdier, som UNFCCC har vedtaget.


Endvidere rapporteres de indirekte drivhusgasser Kvælstofilte (NO_x), Kulilte (CO), Ikke-metan flygtige organiske forbindelser (NMVOC) og Svovldioxid (SO₂). Da der ikke tilskrives disse gasser GWP-værdier, medregnes disse ikke i drivhusgasemissioner i CO₂-ækvivalenter.

S.2 Udviklingen i drivhusgasemissioner og optag

Sammenfatning S.2.-4. omhandler alene opgørelsen for Danmark. Opgørelsen for Grønland, Danmark og Grønland samt for Færøerne beskrives i kapitel 16 og 17 samt i Annex 8.

S.2.1 Drivhusgasemissionsopgørelse

De danske opgørelser af drivhusgasemissioner følger metoderne som beskrevet i IPCC's retningslinjer. I den forbindelse skal nævnes at det under Klimakonventionen og Kyotoprotokollen er vedtaget at IPCC's 1996 retningslinjer og IPCC's 2000 anvisninger skal anvendes. Opgørelserne er opdelt i seks overordnede sektorer, 1. energi, 2. industrielle processer, 3. opløsningsmidler, 4. landbrug, 5. arealanvendelse for skove og jorder (Land Use Land Use Change and Forestry: LULUCF) og 6. affald. Drivhusgasserne omfatter CO₂, CH₄, N₂O og F-gasserne: HFC'er, PFC'er og SF₆. I Figur S.1 ses de estimerede drivhusgasemissioner for Danmark i CO₂-ækvivalenter for perioden 1990 til 2012. Figuren viser Danmarks totale udledning med og uden LULUCF-sektoren (Land Use and Land Use Change and Forestry). Til venstre i figur S.1 ses det relative bidrag til Danmarks totale udledning (uden LULUCF) i 2012 for sektorerne 1. – 4. og 6. For sektor 1. energi er vejtrafik vist særskilt. Sektor 5. LULUCF indgår ikke i denne figur da sektoren omfatter kilder der bidrager med både optag og udledninger.



Figur S.1 Danske drivhusgasemissioner. Bidrag til total emission fra hovedsektorer for 2012 og tidsrækker i CO₂-ækvivalenter for 1990-2012, hvor data er angivet med og uden LULUCF.

I overensstemmelse med retningslinjerne for opgørelserne er emissionerne ikke korrigerede for handel med elektricitet med andre lande og temperatursvingninger fra år til år. CO₂ er den vigtigste drivhusgas og bidrager i 2012 med 76,3 % af den nationale totale udledning uden LULUCF-sektoren, efterfulgt af N₂O med 11,6 % og CH₄ med 10,7 %, mens HFC'er, PFC'er og SF₆ kun udgør 1,5 % af de totale emissioner uden LULUCF-sektoren. Set over perioden 1990-2012 så har disse procenter været stigende for CH₄ og F-gasser og faldende for N₂O. For CO₂ har procenterne fluktueret mere gennem perioden. Netto CO₂-optaget fra LULUCF er i 2012 0,2 % af den nationale totale emission eksklusiv LULUCF. Med hensyn til sektorerne (figur S.1) så bidrager energi ekskl. vejtransport (hovedsageligt stationære for-

brændingsanlæg), transport og landbrug mest i 2012, efterfulgt af industrielle processer, affald, flygtige emissioner, og opløsningsmidler (Figur S.1). De nationale totale drivhusgasemissioner i CO₂-ækvivalenter er faldet med 24,8 % fra 1990 til 2012, hvis nettobidraget fra skovenes og jordernes udledninger og optag af CO₂ (LULUCF) ikke indregnes, og faldet med 30,5 % hvis LULUCF indregnes.

S.2.2 KP-LULUCF-aktiviteter

Den samlede udledning af drivhusgasser i skov omfattet af Kyotoprotokollens artikel 3.3 udgør 148 Gg CO₂-ækvivalenter i 2012, heraf stammer 0,9 Gg CO₂-ækvivalenter fra N₂O-udledning i forbindelse med skovrydning. Nettooptaget fra skov plantet før 1990 under Kyotoprotokollens artikel 3.4 udgør 4 479 Gg CO₂-ækvivalenter i 2012, heraf 12,1 Gg CO₂-ækvivalenter i form af N₂O fra dræning af jorde (tabel S.1).

Nettoemissionen fra landbrugsarealer under artikel 3.4 udgør 2 958 Gg CO₂-ækvivalenter i 2012. Til sammenligning var nettoemissionen fra samme kilde 4 845 Gg CO₂-ækvivalenter i 1990.

Det samlede emission fra permanente græsarealer under artikel 3.4 udgør 523 Gg CO₂-ækvivalenter i 2012. I 1990 var den tilsvarende emission på 177 Gg CO₂-ækvivalenter.

Tabel S.1 Emissioner og optag i 2012 for aktiviteter under Kyotoprotokollens artikel 3.3 og 3.4.

	Netto CO ₂ emission/ optag (Gg)	CH ₄	N ₂ O	Netto CO ₂ - ækvivalent emission/ optag
A. Aktiviteter under artikel 3.3				107,38
A.1. Skovrejsning	76,62	NO	IE,NA,NO	76,62
A.1.1. Arealer der ikke er afskovet siden starten af 2008	76,62	NO	IE,NA,NO	76,62
A.1.2. Arealer der er afskovet siden starten af 2008	IE,NO	NO	IE,NO	IE,NO
A.2. Skovrydning	30,50	NO	0,00	30,76
B. Aktiviteter under artikel 3.4				5.088,29
B.1. Forvaltning af skov plantet før 1990	49,60	0,03	0,05	66,24
B.2. Forvaltning af landbrugsarealer	4.844,66	NO	NA,NO	4.844,66
B.3. Forvaltning af permanente græsarealer	177,39	0,00	0,00	177,39
B.4. Gentilplantning	NA	NA	NA	NA

S.3 Oversigt over drivhusgasemissioner og optag fra sektorer

S.3.1 Drivhusgasemissionsopgørelse

Energi

Udledningen af CO₂ stammer altovervejende fra forbrænding af kul, olie, benzin og naturgas på kraftværker, i beboelsesejendomme, industri og vejtransport. CO₂-emissionen fra energisektorerne faldt med omkring 36,8 % fra 1990 til 2012. De relative store udsving i emissionerne fra år til år skyldes handel med elektricitet med andre lande, herunder særligt de nordiske. De høje emissioner i 1991, 1994, 1996, 2003 og 2006 er et resultat af stor eksport af elektricitet, mens de lave emissioner i 1990, 1992, 2005, 2008, 2011 og 2012 skyldes import af elektricitet. Den væsentligste årsag til dette fald skyldes faldende brændselsforbrug, hovedsageligt for kul og naturgas. Faldet skyldes delvist stigende import af elektricitet og stigende produktion af vindkraft.

Udledningen af CH₄ fra energiproduktion har været stigende på grund af øget anvendelse af gasmotorer, som har en stor CH₄-emission i forhold til andre forbrændingsteknologier. Anvendelsen af gasmotorer er dog blevet mindre siden liberaliseringen af elmarkedet, hvilket har ført til lavere CH₄-emissioner fra energisektoren. Transportsektorens CO₂-emissioner er steget med 14,0 % siden 1990 hovedsagelig på grund af voksende vejtrafik.

Industrielle processer

Emissionen fra industrielle processer – hvilket vil sige andre processer end forbrændingsprocesser – udgør i 2012 2,6 % af de totale danske drivhusgas-emissioner. De vigtigste kilder er cementproduktion, kølesystemer, opskumning af plast og kalcinering af kalksten. CO₂-emissionen fra cementproduktion – som er den største kilde – bidrager med 1,7 % af den totale emission i 2012. Emissionen fra cementproduktion er dog faldet med 1,3 % fra 1990 til 2012. Den anden største kilde har tidligere været N₂O fra produktion af salpetersyre. Produktionen af salpetersyre stoppede i midten af 2004, hvilket betyder, at N₂O-emissionen er nul for denne kilde fra 2005.

Emissionen af HFC'er, PFC'er og SF₆ er i perioden fra 1995 og til 2012 steget med 140,7 %, hovedsageligt på grund af stigende emissioner af HFC'er. Anvendelsen af HFC'er, og specielt HFC-134a, er steget kraftigt, hvilket har betydet, at andelen af HFC'er af den samlede F-gas-emission steg fra 67 % i 1995 og til 84 % i 2012. HFC'er anvendes primært inden for køleindustrien. Anvendelsen er dog nu stagnerende, som et resultat af dansk lovgivning, der forbyder anvendelsen af nye HFC-baserede stationære kølesystemer fra 2007. I modsætning til denne udvikling ses et stigende brug af airconditionssystemer i køretøjer. Den samlede effekt er, at emissionen forventes at falde fremover.

Opløsningsmidler og relaterede produkter

Forbrug af opløsningsmidler i industrier og husholdninger bidrager i 2012 med 0,3 % af totalmængden af emitterede drivhusgasser i CO₂-ækvivalenter. Der er en reduktion på 34,6 % i drivhusgasemissionen i perioden 1990 til 2012. Bidraget fra N₂O til den totale emission i CO₂-ækvivalenter for solventer og anden produktanvendelse er 11,5 %.

Landbrug

Landbrugssektoren bidrager i 2012 med 18,6 % til den totale drivhusgas-emission i CO₂-ækvivalenter og er den vigtigste sektor hvad angår emissioner af N₂O og CH₄. I 2012 var landbrugets bidrag til de totale emissioner af N₂O og CH₄ henholdsvis 90,5 % og 76,6 %. Fra 1990 til 2012 ses et fald på 34,9 % i N₂O-emissionen fra landbrug. Dette skyldes mindre brug af kvælstofhandelsgødning og bedre udnyttelse af kvælstof i husdyrgødningen, hvilket resulterer i mindre emissioner pr. produceret dyreenhed. Emissioner af CH₄ fra husdyrenes fordøjelsessystem er faldet fra 1990 til 2012 grundet et faldende antal kvæg. På den anden side har en stigende andel af gyllebase-rede staldsystemer bevirket at emissionerne fra husdyrgødning er steget. I alt er CH₄-emissionerne fra landbrugssektoren faldet med 0,7 % fra 1990 til 2012.

Arealanvendelse - skove og jorder (LULUCF)

LULUCF-sektoren skifter mellem at udgøre et nettooptag og en nettoudledning. I 2012 udgør LULUCF et nettooptag svarende til 1,6 % af den samlede drivhusgasudledning, eksklusiv LULUCF. I 2011 udgjorde LULUCF et net-

tooptag svarende til 4,9 % af den samlede drivhusgasudledning eksklusiv LULUCF. Siden 1990 er LULUCF sektoren eksklusiv skov faldet med 30 %.

I 2012 bidrager arealer med skov med et optag på 4 441 Gg CO₂-ækvivalenter, mens dyrkede jorder, græsning, vådområder og bebyggelse bidrager med emissioner på henholdsvis 2 956 Gg CO₂-ækvivalenter, 554 Gg CO₂-ækvivalenter, 2 Gg CO₂-ækvivalenter og 91 Gg CO₂-ækvivalenter.

Affald

Affaldssektoren udgør i 2012 2,1 % af den danske totalemission, 15,7 % af den totale CH₄-emission og 3,4 % af den totale N₂O-emission. Sektoren omfatter lossepladser, spildevandshåndtering, affaldsforbrænding uden energidnyttelse (f.eks. kremeringer af dyr), og andet affald (f.eks. kompostering og ildebrænde). Da al traditionel affaldsforbrænding bruges til produktion af elektricitet og varme, er emissionerne herfra inkluderet i CRF-kategorien 1A.

Drivhusgasemissionen fra sektoren er faldet med 32,4 % fra 1990 til 2012. Reduktionen skyldes især (1) et fald i CH₄-emissionen fra lossepladser på 48,9 % pga. reducerede mængder affald, der går til deponi, og (2) et fald i N₂O-emissionen fra spildevandshåndtering på 30,6 % pga. fornyelse af spildvandsanlæggene. Disse fald er delvist modvirket af en stigning i CH₄-emissionen fra spildevandshåndtering på 13,0 % pga. en stigning i det industrielle spildevand. I 2012 bidrog lossepladser med 12,7 % af den totale nationale CH₄-emission. CH₄-emissionen fra spildevandshåndtering udgør i 2012 1,3 % af den totale nationale CH₄-emission. Emissionen af N₂O fra spildevandshåndtering udgør i 2012 1,2 % af den totale nationale N₂O-emission. Da al affaldsforbrænding udnyttes til el- og varmeproduktion, indgår emissionerne i CRF kategorien 1A.

S.3.2 KP-LULUCF-aktiviteter

I 2012 udgjorde aktiviteterne under Kyotoprotokollens artikel 3.3 en netto-udledning på 148 Gg CO₂-ækvivalenter mens aktiviteterne under artikel 3.4 udgjorde et nettooptag på 998 Gg CO₂-ækvivalenter. En kort oversigt over KP-LULUCF findes i kapitel S.2.2 mens en mere detaljeret redegørelse findes i kapitel 11.

S.4 Andre informationer

S.4.1 Kvalitetssikring og -kontrol

Rapporten indeholder en plan for kvalitetssikring og -kontrol af emissions-opgørelserne. Kvalitetsplanen bygger på IPCC's retningslinjer og ISO 9000 standarderne. Planen skaber rammer for dokumentation og rapportering af emissionerne, så opgørelserne er gennemskuelige, konsistente, sammenlignelige, komplette og nøjagtige. For at opfylde disse kriterier, understøtter datastrukturen arbejdsgangen fra indsamling af data til sammenstilling, modellering og til sidst rapportering af data.

Som en del af kvalitetssikringen, udarbejdes der for emissionskilderne rapporter, der detaljeret beskriver og dokumenterer anvendte data og beregningsmetoder. Disse rapporter evalueres af personer uden for Aarhus Universitet, der har høj faglig ekspertise indenfor det pågældende område, men som ikke direkte er involveret i arbejdet med opgørelserne. Indtil nu er rapporter for stationære forbrændingsanlæg, transport og landbrug blevet evalueret. Desuden er der gennemført et projekt, hvor de danske opgørelsesme-

toder, emissionsfaktorer og usikkerheder sammenlignes med andre landes, for yderligere at verificere rigtigheden af opgørelserne.

S.4.2 Fuldstændighed i forhold til IPCC's retningslinjer for kilder og gasser

De danske opgørelser af drivhusgasemissioner indeholder alle de kilder, der er beskrevet i IPCC's retningslinjer.

I Annex 5 er der flere informationer om fuldstændigheden af den danske drivhusgasopgørelse.

S. 4.3 Rekalkulationer og forbedringer

De vigtigste forbedringer af opgørelserne er:

Energi

Stationær forbrænding

Den seneste officielle energistatistik er implementeret i opgørelsen for årene 1990-2011. Opdateringen omfatter både slutforbrug og konverteringssektoren samt opdatering af kilde kategorier. Ændringerne i energistatistikken er størst for årene 2009-2011.

For CO₂ er den største genberegning lavet i kategorien fremstillingsvirksomhed. Genberegningen er relateret til flydende brændsler og er resultatet af en fejlretning. Mængden af fuel olie har været underestimeret i de tidligere opgørelser. CO₂ emissionen fra fuel olie anvendt i fremstillingsvirksomhed er 7 % højere i denne rapportering sammenlignet med 2013 rapporteringen.

For CH₄ er emissionen fra træfyring i husholdninger blevet opdateret pga. opdaterede emissionsfaktorer for brændeovne. De opdaterede emissionsfaktorer har medført en stigning på 16 % i CH₄ emissionen fra forbrænding af biomasse i husholdninger.

For N₂O er den største genberegning lavet i kategorien fremstillingsvirksomhed og skyldes den samme fejlretning som beskrevet for CO₂. Konsekvensen er en stigning på 13 % for N₂O fra flydende brændsler i denne rapportering sammenlignet med 2013 rapporteringen.

Mobile kilder

Vejtransport

Baseret på den opdaterede version af COPERT IV, der blev lanceret i 2013, nye køretøjsunderkategorier er blevet introduceret i emissionsopgørelsen for passagerbiler og knallerter. For knallerter skelnes der nu mellem 2-taks og 4-taks motorer, mens der for passagerbiler er oprettet nye kategorier for benzinerbiler med motorer < 0,8 l og dieslbiler med motorer < 1,4 l. NO_x emissionsfaktorerne for euro 5 passagerbiler er blevet opdateret i modellen baseret på værdierne i den nye COPERT IV version.

Små fejl i data for benzinforbruget for årene 2009-2011 og for dieselforbruget for årene 2010-2011 er blevet rettet.

Minimum og maksimum procentvis difference og år for numerisk maksimum difference (min. %, maks. %, år med maks. %) for emissionskomponenterne er: CO₂ (-0,5 %, -0,05 %, 2008), CH₄ (-0,2 %, 2,4 %, 2011) og N₂O (-0,4 %, 0,3 %, 2008).

Søfart

Ændrede forudsætninger for færgeoverfart for årene 2008-2011 har medført mindre ændringer af emissionerne for national søfart. De følgende maksimale procentvise ændringer for national søfart (i parenteser) som følge af genberegningen er: CO₂ (-1,0 %), CH₄ (-0,8 %) og N₂O (-1,1 %).

Landbrug/skovbrug/fiskeri

Antallet og motorstørrelsen af traktorer på maskinstationer er opdateret for årene 2007-2011. Antallet af terrængående maskiner (ATV'er) er blevet opdateret for årene 2009-2011.

Der er rettet fejl angående brændselsforbruget for fiskeri for årene 2000, 2010 og 2011.

For 2000 er der følgende ændringer for landbrug/skovbrug/fiskeri: CO₂ (9,1 %), CH₄ (3,7 %) og N₂O (11,9 %), pga. ændringer i brændselsforbruget for fiskeri.

For de øvrige år er der følgende ændringer for landbrug/skovbrug/fiskeri, udtrykt ved maksimum procentvis difference for emissionskomponenterne er: CO₂ (-3,7 %), CH₄ (5 %) og N₂O (-4,9 %).

Industri

Antallet af minilæssere er opdateret for årene 2004-2011.

De følgende maksimale procentvise ændringer for national søfart (i parenteser) som følge af genberegningen er: CO₂ (1,0 %), CH₄ (1,6 %) og N₂O (1,6 %).

Luftfart

En mindre fejl i CH₄ emissionsfaktoren er blevet rettet for årene 1990-2000. Emissionsfaktorerne er nu i overensstemmelse med emissionsfaktorerne i EMEP/EEA Guidebook. De procentvise ændringer i CH₄ emissionen er mellem -31 % og -42 %.

Militær

Emissionsfaktorer afledt fra de nye modelsimulationer for vejtransport har medført små ændringer i emissionerne i perioden 1990-2011. Maksimum emissionsdifference er: CO₂ (0 %), CH₄ (0,6 %) og N₂O (0,2 %).

Flygtige emissioner

I forbindelse med rapporteringen i 2014 er der foretaget en række genberegninger som specificeret nedenfor.

Transmission og distribution af naturgas

Aktivitetsdata og afledte emissionsfaktorer for tidsserien 1990-2011 er blevet opdateret for transmission og distribution af naturgas baseret på oplysninger i grønne regnskaber og den seneste energistatistik. Derudover er CH₄ emissionsfaktorerne opdateret for et bygasselskab. Genberegningen har ændret CO₂ emissionen med 0,01 kton og CH₄ emissionen med (-0,09)-0,07 kton svarende til <0,003 % og (-2) % - 2 % af henholdsvis den totale CO₂ og CH₄ emission fra flygtige kilder.

Venting

Emissionsfaktorer for CH₄ er blevet tilføjet emissionsdatabasen for årene 1990-1993 for et gaslager. I disse år er anlægget inkluderet som en arealkilde i den nationale emissionsdatabase, mens det er behandlet som en punktkilde i de følgende år. Emissionsfaktorerne er baseret på data fra grønne regnskaber for 1995-1999, da der ikke er tilgængelige data for årene 1990-1994. Derudover er en mindre fejl blevet rettet i 2011 baseret på opdaterede oplysninger i det grønne regnskab fra et af de danske naturgaslagre.

Genberegningen har betydet en stigning i CH₄ emissionen på 0,06 kton svarende til 2 % af den totale CH₄ emission fra flygtige kilder.

Flaring i forbindelse med gaslagre og naturgasbehandlingsanlæg

Emissionsfaktorer for CO₂ og CH₄ er blevet tilføjet for årene 1990-1993 for gasbehandlingsanlægget. I disse år er anlægget inkluderet som en arealkilde i den nationale emissionsdatabase, mens det er behandlet som en punktkilde i de følgende år. Emissionsfaktorerne er baseret på data fra grønne regnskaber for 1995-1999, da der ikke er tilgængelige data for årene 1990-1994.

Genberegningen har betydet en stigning i CO₂ emissionen på 2,2 kton og en stigning i CH₄ emissionen på 0,01 kton svarende til 0,4 % af både den totale CO₂ og CH₄ emission fra flygtige kilder.

Flaring i forbindelse med raffinaderier

CO₂ emissionsfaktoren for et raffinaderi er opdateret for årene 1994-2006 og afspejler nu gennemsnittet af de første 5 års rapportering (2007-2011) under EU ETS. NO_x emissionsfaktorerne for de to raffinaderier er blevet opdateret til default emissionsfaktoren i EMEP/EEA Guidebook, da referencerne til de hidtidigt anvendte emissionsfaktorer er forældede eller ikke-eksisterende.

CO₂ emissionsfaktoren for et raffinaderi der blev lukket i 1996 er blevet ændret for årene 1990-1996, så den svarer til de øvrige to raffinaderier, da der ikke er bedre data tilgængelige.

CO₂ emissionsfaktorerne for 2010 og 2011 er opdateret i henhold til EU ETS indberetningerne.

Genberegningerne har medført ændringer i CO₂ emissionen på (-0,05) - 0,49 kton og i CH₄ emissionen på (-0,06) kton svarende til (-0,01) - 0,1 % og (-2) % af henholdsvis den samlede CO₂ og CH₄ emission fra flygtige kilder.

Industrielle processer

Produktion af brændt kalk

EU ETS data er blevet indarbejdet for en producent af brændt kalk for 2011, hvilket har medført et lille fald i den samlede CO₂ emission på 5,51 kton. Ændringen vil blive tilbageført for resten af tidsserien i 2015 afleveringen.

Anvendelse af kalksten og dolomit

Aktivitetsdata for anvendelse af kalksten til røggasrensning på tre kraftværker og affaldsforbrændingsanlæg er opdateret for 2011. Den anvendte kalkmængde er steget med 236 ton og mængden af CaCO₃ holdige restprodukter er steget med 669 ton resulterende i en stigning i CO₂ emissionen på 162 ton.

Kemisk industri

Procesemissionen er blevet korrigeret, så den afspejler den totale CO₂ emission rapporteret i det grønne regnskab fratrukket de energirelaterede emissioner rapporteret under EU ETS.

Anvendelse af HFC'er, PFC'er og SF₆

En mindre rettelse er foretaget for emissionen af SF₆ fra termoruder for 2010. Potentielle emissioner for 2005-2007 er ændret til at reflektere at den potentielle emission er identisk med den anvendte årlige mængde. Anvendelse af SF₆ i termoruder stoppede i 2001.

Opløsningsmidler og anden produktanvendelse

Der er foretaget genberegninger for øvrig produktanvendelse på baggrund af ændrede aktivitetsdata for alle fire produktanvendelser; lys (2009-2011), fyrværkeri (2009-2011), tobak (1980-1999, 2011) og trækul anvendt til grilning (1980-1987, 2009-2011). Disse ændringer har medført ændringer for NMVOC, N₂O og CO₂. NMVOC emissionen er steget for årene 1980-1999 med mellem 6 % (1997) og 9,1 % (1982) og faldet for årene 2009-2011 med mellem 2,1 % (2009) og 5,7 % (2011). CO₂ emissionen er faldet for 2009 (13,4 %) og 2010 (6,1 %) og steget for 2011 (0,7 %). N₂O emissionen er steget for årene 1980-1999 med mellem 0,5 % (1999) og 2,8 % (1982) og faldet for årene 2009-2011 med mellem 0,3 % (2011) og 1,0 % (2009).

Landbrug

Genberegninger for landbrugssektoren har medført et fald i emissionen for årene 1990-2011 på op til 0,2 % sammenlignet med den totale emission i CO₂-ækvivalenter fra landbrugssektoren.

CH₄ emissionen falder både for fordøjelse og gødningshåndtering. Antallet af gæs er opdateret for alle år og antallet af smågrise og slagtesvin er opdateret for 2011. Disse ændringer påvirker både emissionerne fra fordøjelsen og fra gødningshåndtering. Mængden af halm anvendt som strøelse for kvier er opdateret for årene 1990-2002 og mængden af biogas behandlet gylle i 2010 er opdateret. Disse ændringer påvirker emissionen fra gødningshåndtering.

For N₂O emissionen er der foretaget en række genberegninger, som påvirker emissionen i begge retninger. Emissionsfaktorerne for NH₃ fra handelsgødning er opdateret for alle år, og dette påvirker N₂O emissionen. Den stigende NH₃ emission fra handelsgødning medfører et fald i N₂O emissionen, men til gengæld en stigning i N₂O emissionen fra atmosfærisk deposition. Ændringen i antallet af gæs for alle år har medført et fald i N₂O emissionen fra græsning. Ændringen i mængden af biogas behandlet gylle i 2010 reducerer N₂O emissionen. Emissionerne fra gødningshåndtering, udbringning og udvaskning falder i 2011 på grund af opdaterede data for antallet af smågrise og slagtesvin.

Arealanvendelse (LULUCF)

Der er foretaget en opdatering af arealmatricen for alle år. Opdateringen var nødvendig, da der var fejl i det kortmateriale, der var leveret af Geodatastyrelsen. Fejlene var især angående klassificering af rekreative områder og fejlagtige klassificeringer af parker som skov.

Landbrugsarealer, græsningsarealer, vådområder og bebyggelse

Udover effekterne af den opdaterede arealmatrice, er der rettet to mindre fejl i forbindelse med levende biomasse i bebyggede områder samt arealerne in-

denfor Cropland Management og Grazing Land Management under Kyoto-protokollens artikel 3.4.

Affald

Deponier

Baseret på det nye affaldsrapporteringssystem er der foretaget en opdateret disaggregering af affaldskategorierne. Der opereres nu med 18 affaldskategorier, hvoraf 11 er inerte. Resultatet af den ændrede karakterisering og reallokering af affald er et fald i CH₄ emissionen i 1990 på 7,6 % og en stigning i CH₄ emissionen i 2011 på 4,6 %.

Spildevandshåndtering

For spildevandshåndtering er der ikke foretaget en genberegning af N₂O-emissionen. For CH₄ er MCF (Methane correction factor) værdien reduceret fra 1 til 0,8 i overensstemmelse med IPCC 2006 Guidelines og som er bekræftet af anlægsspecifikke data. Udover ændringer i MCF værdi, så er der foretaget mindre ændringer i TOW (Total organic waste) data svarende til en ændring under 1 % gennem tidsserien.

Affaldsforbrænding

Antallet af betydende cifre er blevet rettet for kremering af dyr. Genberegningen har resulteret i en stigning i N₂O emissionen for perioden 1990-2011 på 0,07 %.

Anden affaldsbehandling

For bilbrande er der foretaget en opdatering i bestandsdata, som har medført et lille fald i aktivitetsdata for brande i traktorer og mejetærskere. Derudover er gennemsnitsvægtene, der er antaget for campingvogne, campere, mejetærskere, motorcykler og knallerter blevet opdateret på baggrund af mere velfunderede ekspertvurderinger. Resultatet er et fald i emissionen af CH₄ og CO₂ fra 1990-2006 og en stigning for 2007-2011. Emissionsfaktorerne for kompostering af slam og organisk affald er opdateret, hvilket har medført stigende emissioner af CH₄ og N₂O. Den samlede effekt af genberegningerne er et fald i CH₄ emissionen fra 1980 (2,4 %) til 1984 (2,0 %) og en stigning fra 1985 (16,7 %) til 2011 (275 %). For CO₂ er emissionerne faldet mellem 1980 (1,1 %) og 2006 (0,004 %) og steget fra 2007 (0,1 %) til 2011 (0,2 %). For N₂O er emissionen steget for alle år mellem 1985 (1,1 %) og 2011 (238 %).

KP-LULUCF

Der er genberegnet for alle år som følge af den opdaterede arealmatrice.

1 Introduction

1.1 Background information on greenhouse gas inventories and climate change

1.1.1 Annual report

This report is Denmark's National Inventory Report (NIR) 2014 for submission to the United Nations Framework Convention on Climate change and the Kyoto Protocol, due April 15, 2014. The report contains detailed information about Denmark's inventories for all years from 1990 to 2012. The structure of the report is in accordance with the UNFCCC guidelines on reporting and review. The main difference between Denmark's NIR 2014 report to the European Commission, due March 15, 2014, and this report to UNFCCC is reporting of territories. The NIR 2014 to the EU Commission was for Denmark, while this NIR 2014 to the UNFCCC is for Denmark, Greenland and the Faroe Islands. The suggested outline provided by the UNFCCC secretariat has been followed to include the necessary information under the Kyoto Protocol. The report includes detailed and complete information on the inventories for all years from year 1990 to the year 2012, in order to ensure transparency.

The issues addressed in this report are trends in greenhouse gas emissions, a description of each IPCC category, uncertainty estimates, recalculations, planned improvements and procedures for quality assurance and control.

The annual emission inventories for the years from 1990 to 2012 are reported in the Common Reporting Format (CRF) as requested in the reporting guidelines. The CRF-spreadsheets contain data on emissions, activity data and implied emission factors for each year. Emission trends are given for each greenhouse gas and for the total greenhouse gas emissions in CO₂ equivalents.

According to the instrument of ratification, the Danish government has ratified the UNFCCC on behalf of Denmark, Greenland and the Faroe Islands. The Danish government has ratified the Kyoto Protocol on behalf of Denmark and Greenland. The information in the sectoral chapters in this report relates to Denmark only, while information for Greenland is included in Chapter 16 and for the Faroe Islands in Annex 8. Chapter 17 contains information (e.g. on trends, uncertainties and key category analysis) on the aggregated submission of Denmark and Greenland under the Kyoto Protocol.

This report itself does not contain the full set of CRF Tables. The full set of CRF tables is available at the EIONET, Central Data Repository, kept by the European Environmental Agency:

http://cdr.eionet.europa.eu/dk/Air_Emission_Inventories/Submission_UNFCCC

1.1.2 Greenhouse gases

The greenhouse gases reported under the Climate Convention are:

- Carbon dioxide CO₂
- Methane CH₄
- Nitrous Oxide N₂O

- Hydrofluorocarbons HFCs
- Perfluorocarbons PFCs
- Sulphur hexafluoride SF₆

The main greenhouse gas responsible for the anthropogenic influence on the heat balance is CO₂. The atmospheric concentration of CO₂ has increased from a pre-industrial value of about 280 ppm to 379 ppm in 2005 (an increase of about 35 %), and exceeds now the natural range of 180-300 ppm over the last 650 000 years as determined by ice cores (IPCC, Fourth Assessment Report, 2007). The main cause for the increase in CO₂ is the use of fossil fuels, but changing land use, including forest clearance, has also been a significant factor. The greenhouse gases CH₄ and N₂O are very much linked to agricultural production; CH₄ has increased from a pre-industrial atmospheric concentration of about 715 ppb to 1774 ppb in 2005 (an increase of about 140 %) and N₂O has increased from a pre-industrial atmospheric concentration of about 270 ppb to 319 ppb in 2005 (an increase of about 18 %) (IPCC, Fourth Assessment Report, 2007). Changes in the concentrations of greenhouse gases are not related in simple terms to the effect on the heat balance, however. The various gases absorb radiation at different wavelengths and with different efficiency. This must be considered in assessing the effects of changes in the concentrations of various gases. Furthermore, the lifetime of the gases in the atmosphere needs to be taken into account – the longer they remain in the atmosphere, the greater the overall effect. The global warming potential (GWP) for various gases has been defined as the warming effect over a given time of a given weight of a specific substance relative to the same weight of CO₂. The purpose of this measure is to be able to compare and integrate the effects of individual substances on the global climate. Typical lifetimes in the atmosphere of substances are very different, e.g. 12 and 120 years approximately for CH₄ and N₂O, respectively. So the time perspective clearly plays a decisive role. The time frame chosen is typically 100 years. The effect of the various greenhouse gases can, then, be converted into the equivalent quantity of CO₂, i.e. the quantity of CO₂ giving the same effect in absorbing solar radiation. According to the IPCC and their Second Assessment Report, which UNFCCC has decided to use as reference for reporting for inventory years throughout the commitment period 2008-2012, the global warming potentials for a 100-year time horizon are:

- Carbon dioxide (CO₂): 1
- Methane (CH₄): 21
- Nitrous oxide (N₂O): 310

Based on weight and a 100-year period, methane is thus 21 times more powerful a greenhouse gas than CO₂, and N₂O is 310 times more powerful. Some of the other greenhouse gases (hydrofluorocarbons, perfluorocarbons and sulphur hexafluoride) have considerably higher global warming potential values. For example, sulphur hexafluoride has a global warming potential of 23 900.

The indirect greenhouse gases reported are nitrogen oxides (NO_x), carbon monoxide (CO), non-methane volatile organic compounds (NMVOC) and sulphur dioxide (SO₂). Since no GWP is assigned these gases they do not contribute to GHG emissions in CO₂ equivalents.

1.1.3 The Climate Convention and the Kyoto Protocol

At the United Nations Conference on Environment and Development in Rio de Janeiro in June 1992, more than 150 countries signed the UNFCCC (the Climate Convention). On the 21st of December 1993, the Climate Convention was ratified by a sufficient number of countries, including Denmark, for it to enter into force on the 21st of March 1994. One of the provisions of the treaty was to stabilise the greenhouse gas emissions from the industrialised nations by the end of 2000. At the first conference under the UN Climate Convention in March 1995, it was decided that the stabilisation goal was inadequate. At the third conference in December 1997 in Kyoto in Japan, a legally binding agreement was reached committing the industrialised countries to reduce the six greenhouse gases by 5.2 % by 2008-2012 compared with the base year. For F-gases, the countries can choose freely between 1990 and 1995 as the base year. On May 16, 2002, the Danish parliament voted for the Danish ratification of the Kyoto Protocol. Denmark (including Greenland and excluding the Faroe Islands) is, thus, under a legal commitment to meet the requirements of the Kyoto Protocol, when it came into force on the 16th of February 2005. Hence, Denmark (including Greenland) is committed to reduce greenhouse gases with 8 %. The European Union is under the KP committed to reduce emissions of greenhouse gases by 8 %. However, within the EU member states have made a political agreement – the Burden Sharing Agreement – on the contributions to be made by each member state to the overall EU reduction level of 8 %.

Under the Burden Sharing Agreement, Denmark (excluding Greenland and the Faroe Islands) must reduce emissions by an average of 21 % in the period 2008-2012 compared with the base year emission level.

In accordance with the Kyoto Protocol, Denmark's base year emissions include the emissions of CO₂, CH₄ and N₂O in 1990 in CO₂ equivalents and Denmark has chosen 1995 as the base year for the emissions of HFCs, PFCs and SF₆.

1.1.4 The role of the European Union

The European Union (EU) is a party to the UNFCCC and the Kyoto Protocol. Therefore, the EU has to submit similar datasets and reports for the collective 15 EU Member States under the burden sharing. The EU imposes some additional guidelines and obligations to these EU Member States through Decision No. 280/2004/EC concerning a mechanism for monitoring community greenhouse gas emissions and for implementing the Kyoto Protocol (EU monitoring mechanism). In 2013 a new regulation was agreed regarding the reporting of information related to greenhouse gases in the EU, the regulation is Regulation (EU) No 525/2013.

1.1.5 Background information on supplementary information required under KP article 7.1

For the LULUCF activities under Article 3, paragraphs 3 and 4, of the Kyoto Protocol Denmark has chosen annual accounting. Article 3.3 covers direct, human induced afforestation (A), reforestation (R) and deforestation (D) activities, and accounting of these activities is mandatory. Under Article 3.4 Denmark has elected the activities Forest Management (FM), Cropland Management (CM) and Grazing Land Management (GM) for accounting in the first Commitment Period (CP). Net removals from FM can be used to compensate net emissions from activities under Article 3.3 (Article 3.3 off-

set), and through the issuance of removal units (RMUs) up to a cap value. Denmark's cap value for the CP is 916 667 tonnes CO₂ equivalents.

1.2 A description of the institutional arrangement for inventory preparation

On behalf of the Ministry of the Environment and the Ministry of Climate, Energy and Building the Danish Centre for Environment and Energy (DCE) is responsible for the calculation and reporting of the Danish national emission inventory to the EU, the UNFCCC (United Nations Framework Convention on Climate Change) and UNECE CLRTAP (Convention on Long-Range Transboundary Air Pollution). Hence, DCE prepares and publishes the annual submission for Denmark to the EU and UNFCCC of the National Inventory Report and the GHG inventories in the Common Reporting Format, in accordance with the UNFCCC guidelines. Furthermore, DCE is responsible for reporting the national inventory for the Kingdom of Denmark to the UNFCCC. DCE is also the body designated with overall responsibility for the national inventory under the Kyoto Protocol for Greenland and Denmark.

The work concerning the annual greenhouse gas emission inventory is carried out in cooperation with Danish ministries, research institutes, organisations and companies. The Government of Greenland is responsible for finalising and transferring the inventory for Greenland to DCE. The Faroe Islands Environmental Agency is responsible for finalising and transferring the inventory for the Faroe Islands to DCE.

There are now data agreements in place with both Greenland and the Faroe Islands ensuring the data delivery. These agreements contain deadlines for when DCE is to receive the data and documentation.

DCE has been and is engaged in work in connection with meetings of the Conference of the Parties (COP) to the UNFCCC and the Conference of the Parties serving as the Meeting of the Parties (COP/MOP) to the Kyoto protocol and its subsidiary bodies, where the reporting rules are negotiated and settled. Furthermore, DCE participates in the EU Monitoring Mechanism, Working Group 1 (WG1), where the guidelines, methodologies etc. on inventories to be prepared by the EU Member States are regulated.

The main experts responsible for the sectoral inventories and the corresponding chapters and annexes in this report are:

Project leader		Ole-Kenneth Nielsen (okn@dmu.dk)
Sector	Sub-sector	Expert name
Energy	Stationary combustion:	Malene Nielsen
	Transport and other mobile sources	Morten Winther
	Fugitive emissions:	Marlene Plejdrup
Industrial processes		Leif Hoffmann
Solvent and other product use		Patrik Fauser, Katja Hjelgaard
Agriculture		Mette Hjorth Mikkelsen, Rikke Albrektsen & Steen Gyldenkærne
LULUCF	Forestry	Vivian Kvist Johannsen, Thomas Nord-Larsen, Inge Stupak Møller & Lars Vesterdal
LULUCF	Cropland, grassland, wetlands, settlements	Steen Gyldenkærne
Waste	Solid waste disposal on land	Marianne Thomsen, Katja Hjelgaard
	Wastewater handling	Marianne Thomsen
	Waste incineration & Other waste	Katja Hjelgaard
Greenland		Lene Baunbæk
Faroe Islands		Maria Gunnleivsdóttir Hansen

The work concerning the annual greenhouse emission inventory is carried out in cooperation with other Danish ministries, research institutes, organisations and companies:

Danish Energy Agency, the Ministry of Climate, Energy and Building: Annual energy statistics in a format suitable for the emission inventory work and fuel-use data for the large combustion plants. Company reports submitted under EU ETS.

Danish Environmental Protection Agency, the Ministry of the Environment: Database on waste and emissions of F-gases.

Danish Nature Agency, the Ministry of the Environment: Database on Danish waste water quality parameters.

Statistics Denmark, the Ministry of Economic Affairs and the Interior: Statistical yearbook, sales statistics for manufacturing industries and agricultural statistics.

Danish Centre for Food and Agriculture (DCA), Aarhus University: Data on use of mineral fertiliser, feeding stuff consumption and nitrogen turnover in animals.

Department of Transport, Technical University of Denmark: Number of vehicles grouped in categories corresponding to the EU classification, mileage (urban, rural, highway), trip speed (urban, rural, highway).

Danish Centre for Forest, Landscape and Planning, University of Copenhagen: Background data for Forestry and CO₂ uptake by forest. Responsible for preparing estimates of emissions/removals for reporting under KP article 3.3 and for reporting FM under article 3.4.

Civil Aviation Agency of Denmark, the Ministry of Transport: City-pair flight data (aircraft type and origin and destination airports) for all flights leaving major Danish airports.

Danish Railways, the Ministry of Transport: Fuel-related emission factors for diesel locomotives.

Danish companies: Audited green accounts and direct information gathered from producers and agency enterprises.

Formerly, the provision of data was on a voluntary basis, but more formal agreements are now prepared. This is the case for e.g. the Danish Energy Agency, where the data agreement specifies the data needed and the deadlines for when DCE is to receive the data.

Additionally DCE receives data from Greenland and the Faroe Islands in order to report for the Kingdom of Denmark:

Statistics Greenland: Complete CRF tables for Greenland and documentation for the inventory process.

The Faroe Islands Environmental Agency: Complete CRF tables for the Faroe Islands and documentation for the inventory process.

The complete emission inventories for the three different submissions (EU, Kyoto Protocol and UNFCCC) by Denmark are compiled by DCE and along with the documentation report (NIR) sent for official approval. In recent years the responsibility for official approval has changed. Previously it was the Danish Environmental Protection Agency (Ministry of the Environment) now it is the Danish Energy Agency (Ministry of Climate, Energy and Building). This means that the emission inventory is finalised no later than March 15, whereupon the official approval is done prior to the reporting deadlines under the UNFCCC and the Kyoto Protocol.

1.3 Brief description of the process of inventory preparation. Data collection and processing, data storage and archiving

The background data (activity data and emission factors) for estimation of the Danish emission inventories is collected and stored in central databases located at the Department of Environmental Science (ENVS), Aarhus University. The databases are in Access format and handled with software developed by the European Environmental Agency and developed originally by the former National Environmental Research Institute (NERI), but is now maintained and further developed by ENVS. As input to the databases, various sub-models are used to estimate and aggregate the background data in order to fit the format and level in the central databases. The methodologies and data sources used for the different sectors are described in Chapter 1.4 and Chapters 3 to 9. As part of the QA/QC plan (Chapter 1.6), the data structure for data processing supports the pathway from collection of raw data to data compilation, modelling and final reporting.

For each submission, databases and additional tools and submodels are frozen together with the resulting CRF-reporting format. This material is placed on central institutional servers, which are subject to routine back-up services. Material, which has been backed up, is archived safely. A further documentation and archiving system is the official journal for DCE. In this journal system, correspondence, both in-going and out-going, is registered, which in this case involves the registration of submissions and communica-

tion on inventories with the UNFCCC Secretariat, the European Commission, review teams, etc.

Figure 1.1 shows a schematic overview of the process of inventory preparation. The figure illustrates the process of inventory preparation from the first step of collecting external data to the last step, where the reporting schemes are generated for the UNFCCC and EU (in the CRF format (Common Reporting Format)) and to the United Nations Economic Commission for Europe/Cooperative Programme for Monitoring and Evaluation of the Long-range Transmission of Air Pollutants in Europe (UNECE/EMEP) (in the NFR format (Nomenclature For Reporting)). For data handling, the software tool is CollectER (Pulles et al., 1999) and for reporting the software tool is the CRF reporter tool developed by the UNFCCC Secretariat together with additional tools originally developed by NERI, but now maintained and further developed by ENVS. Data files and programme files used in the inventory preparation process are listed in Table 1.1.

Table 1.1 List of current data structure; data files and programme files in use.

QA/QC Level	Name	Application type	Path	Type	Input sources
4 store	CFR Submissions (UNFCCC and EU)	External report	U:\ST_ENVS-Luft-Emi\Inventory\AllYears\8_AllSectors\Level_4a_Storage\	MS Excel, xml	CRF Reporter
4 store	NFR Report	External report	U:\ST_ENVS-Luft-Emi\Inventory\AllYears\8_AllSectors\Level_4a_Storage\	xls	NRF Report N8 Process
3 process	CRF Reporter	Management tool	Working path: local machine Archive path: U:\ST_ENVS-Luft-Emi\Inventory\AllYears\8_AllSectors\Level_3b_Processes	(exe + mdb)	National Compiler and Importer2CRF(xml) and IDAtoCRF(xml)
3 process	NRF Report N8 Process	Help tool	U:\ST_ENVS-Luft-Emi\Inventory\AllYears\8_AllSectors\Level_3b_Processes\NFR	Excel	NERIRep and Report Template (xls)
3 process	Importer2CRF	Help tool	U:\ST_ENVS-Luft-Emi\Inventory\AllYears\8_AllSectors\Level_3b_Processes	MS Access	CRF Reporter, CollectEr2CRF, and excel files
3 process	CollectER2CRF	Help tool	U:\ST_ENVS-Luft-Emi\Inventory\AllYears\8_AllSectors\Level_3b_Processes	MS Access	NERIRep
3 process	IDA2CRF	Help tool	U:\ST_ENVS-Luft-Emi\Inventory\AllYears\8_AllSectors\Level_3b_Processes	MS Access	IDA_backend
2 process 3 store	NERIRep	Help tool	Working path: I:\ROSPROJ\LUFT_EMIDMURep	MS Access	CollectER databases; dk1972.mdb..dkxxxx.mdb and IDA_backend
2 process	CollectER	Management tool	Working path: local machine Archive path: U:\ST_ENVS-Luft-Emi\Inventory\AllYears\8_AllSectors\Level_2b_Processes	(exe +mdb)	Sector Expert
2 store	dk1980.mdb.dkxxxDatastore x.mdb		U:\ST_ENVS-Luft-Emi\Inventory\AllYears\8_AllSectors\Level_2a_Storage	MS Access	CollectER
1 process	IDA	Management	U:\ST_ENVS-Luft-Emi\Agriculture\InventoryAgricultureData	MS Access	Sector Expert
1 store	IDA_Backend	Datastore	U:\ST_ENVS-Luft-Emi\Agriculture\InventoryAgricultureData	MS Access	IDA




Figure 1.1 Schematic diagram of the process of inventory preparation.

Denmark has different geographical definitions for different submissions. Under the European Union only mainland Denmark is included. For the reporting under the Kyoto Protocol the submission includes Denmark and Greenland, while the reporting under the UNFCCC includes Denmark, Greenland and the Faroe Islands.

Due to the different geographical scopes of the Danish inventory submissions it is necessary to operate three independent installations of the CRF Reporter software on different virtual computers.

For the preparation of the Danish submission under the Kyoto Protocol the full Danish CRF is aggregated with the Greenlandic CRF and for the UNFCCC reporting this is also aggregated with the CRF of the Faroe Islands. The process of aggregation requires additional software tools and two additional installations of CRF Reporter. The process of aggregating the KP inventory is described in Chapter 17.

1.4 Brief general description of methodologies and data sources used

Denmark's air emission inventories are based on the Revised 1996 Intergovernmental Panel on Climate Change (IPCC) Guidelines for National Greenhouse Gas Inventories (IPCC, 1996), the Good Practice Guidance and Uncertainty Management in National Greenhouse Gas Inventories (IPCC, 2000), the Good Practice Guidance for Land Use, Land-Use Change and Forestry (IPCC, 2003) and the CORINAIR methodology. CORINAIR (COoRdination of INformation on AIR emissions) is a European air emission inventory programme for national sector-wise emission estimations, harmonised with the IPCC guidelines. To ensure estimates are as timely, consistent, transparent, accurate and comparable as possible, the inventory programme has developed calculation methodologies for most subsectors and software for storage and further data processing (EMEP-/CORINAIR, 2007).

A thorough description of the CORINAIR inventory programme used for Danish emission estimations is given in Illerup et al. (2000). The CORINAIR calculation principle is to calculate the emissions as activities multiplied by emission factors. Activities are numbers referring to a specific process generating emissions, while an emission factor is the mass of emissions per unit activity. Information on activities to carry out the CORINAIR inventory is largely based on official statistics. The most consistent emission factors have been used, either as national values or default factors proposed by international guidelines.

A list of all subsectors at the most detailed level is given in Illerup et al. (2000) together with a translation between CORINAIR and IPCC codes for sector classifications.

1.4.1 Stationary Combustion Plants

Stationary combustion plants are part of the CRF emission sources *1A1 Energy Industries*, *1A2 Manufacturing Industries* and *1A4 Other sectors*.

The Danish emission inventory for stationary combustion plants is based on the CORINAIR system described in Illerup et al. (2000). The emission inventory for stationary combustion is based on activity rates from the Danish energy statistics. General emission factors for various fuels, plants and sectors have been determined. Some large plants, such as power plants, are registered individually as large point sources and plant-specific emission data are used.

The fuel consumption rates are based on the official Danish energy statistics prepared by the Danish Energy Agency (DEA). DCE aggregates fuel consumption rates to SNAP categories. The fuel consumption of the NFR category 1A4 Manufacturing industries and construction is disaggregated to subsectors according to the DEA data prepared and reported to Eurostat.

For each of the fuel and SNAP categories (sector and e.g. type of plant), a set of general emission factors has been determined. Some emission factors refer to the EMEP/EEA guidebook and some are country specific and refer to Danish legislation, Danish research reports or calculations based on emission data from a considerable number of plants.

Some of the large plants, such as e.g. power plants and municipal waste incineration plants are registered individually as large point sources and emission data from the actual plants are used. This enables use of plant specific emission factors that refer to emission measurements stated in annual environmental reports, etc. At present, the emission factors for CH₄ and N₂O are, however, not plant-specific, whereas emission factors for SO₂ and NO_x often are. For CO₂ it was possible to use data reported under the EU-ETS in the emission inventory from 2006. Therefore it was possible to derive some plant specific CO₂ emission factors for coal and oil fired power plants.

The CO₂ from incineration of the plastic part of municipal waste is included in the Danish inventory.

In addition to the detailed emission calculation in the national approach, CO₂ emission from fuel combustion is aggregated using the reference approach. In 2012, the CO₂ emission inventory based on the reference approach and the national approach, respectively, differ by 0.97 %.

Please refer to Chapter 3.2 and Annex 3A for further information on the emission inventory for stationary combustion plants.

1.4.2 Transport

The emissions from transport, referring to SNAP category 07 (road transport) and the sub-categories in 08 (other mobile sources), are made up in the IPCC categories: 1A2f (Industry-other), 1A3a (Civil aviation), 1A3b (road transport), 1A3c (Railways), 1A3d (Navigation), 1A4a (Commercial and Institutional), 1A4b (Residential), 1A4c (Agriculture/forestry/fisheries) and 1A5 (Other).

An internal DCE model with a structure similar to the European COPERT IV emission model (EMEP/EEA, 2009) is used to calculate the Danish annual emissions for road traffic. The emissions are calculated for operationally hot engines, during cold start and fuel evaporation. The model also includes the emission effect of catalyst wear. Input data for vehicle stock and mileage is obtained from DTU Transport and Statistics Denmark, and is grouped according to average fuel consumption and emission behaviour. For each group, the emissions are estimated by combining vehicle type and annual mileage figures with hot emission factors, cold:hot ratios and evaporation factors (Tier 2 approach).

For air traffic, from 2001 onwards estimates are made on a city-pair level, using flight data provided by the Danish Civil Aviation Agency (CAA-DK) for flights between Danish airports and flights between Denmark and Greenland/Faroe Islands), and LTO and distance-related emission factors from the CORINAIR guidelines (Tier 2 approach). For previous years, the background data consists of LTO/aircraft type statistics from Copenhagen Airport and total LTO numbers from CAA-DK. With appropriate assumptions, consistent time series of emissions are produced back to 1990 and include the findings from a Danish city-pair emission inventory in 1998.

Off-road working machines and equipment are grouped in the following sectors: inland waterways (pleasure craft), agriculture, forestry, industry, and household and gardening. The sources for stock and operational data are various branch organisations and key experts. In general, the emissions are calculated by combining information on the number of different machine types and their respective load factors, engine sizes, annual working hours and emission factors (Tier 2 approach).

The inventory for navigation consists of regional ferries, local ferries and other national sea transport (sea transport between Danish ports and between Denmark and Greenland/Faroe Islands). For regional ferries, the fuel consumption and emissions are calculated as a product of number of round trips per ferry route (Statistics Denmark), sailing time per round trip, share of round trips per ferry, engine size, engine load factor and fuel consumption/emission factor. The estimates take into account the changes in emission factors and ferry specific data during the inventory period.

For the remaining navigation categories, the emissions are calculated simply as a product of total fuel consumption and average emission factors. For each inventory year, this emission factor average comprises the emission factors for all present engine production years, according to engine life times.

Please refer to Chapter 3.3 and Annex 3B for further information on emissions from transport.

1.4.3 Fugitive emissions from fuels

Fugitive emissions from oil (1.B.2.a)

Fugitive emissions from oil are estimated according to the methodology described in the Emission Inventory Guidebook (EMEP/EEA, 2009). The sources include offshore extraction of oil and gas, onshore oil tanks, onshore and offshore loading of ships, and gasoline distribution. Activity data is given in the Danish Energy Statistics by the Danish Energy Agency. The emission factors are based on the figures given in the guidebook except in the case of onshore oil tanks and gasoline distribution where national values are included.

The VOC emissions from petroleum refinery processes cover non-combustion emissions from feed stock handling/storage, petroleum products processing, and product storage/handling. SO₂ is also emitted from non-combustion processes and includes emissions from product processing and sulphur-recovery plants. The emission calculations are based on information from the Danish refineries.

Fugitive emissions from natural gas (1.B.2.b)

Inventories of NMVOC emission from transmission and distribution of natural gas and town gas are based on annual environmental reports from the Danish gas transmission company and annual reports for the gas distribution companies. The annual gas composition is based on Energinet.dk.

Fugitive emissions from flaring (1.B.2.c)

Emissions from flaring offshore, in gas treatment and storage plants, and in refineries are included in the inventory. Emissions calculations are based on annual reports from the Danish Energy Agency and environmental reports from gas storage and treatment plants and the refineries. Calorific values are based on the reports for the EU ETS for offshore flaring, on annual gas quality data from Energinet.dk, and on additional data from the refineries. Emission factors are based on the Emission Inventory Guidebook (EMEP/EEA, 2009).

Please refer to Chapter 3.5 for further information on fugitive emissions from fuels.

1.4.4 Industrial processes

Energy consumption associated with industrial processes and the emissions thereof are included in the Energy sector of the inventory. This is due to the overall use of energy balance statistics for the inventory.

There is only one producer of cement in Denmark, Aalborg Portland Ltd. The activity data for the production of cement clinker is obtained from the company and the CO₂ emission is from the company report to EU-ETS. The methodology is approved by the Danish Energy Agency and the yearly emission estimate is in accordance with the methodology.

The reference for the activity data for production of lime, hydrated lime, expanded clay products and bricks is the production statistics from the manufacturing industries, published by Statistics Denmark.

Limestone is used for the refining of sugar as well as for wet flue gas cleaning at power plants and waste incineration plants. The reference for the activity data is Statistics Denmark for sugar, Energinet.dk for gypsum from power plants combined with specific information on consumption of CaCO_3 at specific power plants and National Waste Statistics for gypsum from waste incineration. The emission factors are based on stoichiometric relations between consumption of CaCO_3 and gypsum generation as well as consumption of lime for sugar refining and precipitation with CO_2 . This information is supplemented with company reports to EU-ETS.

The reference for the activity data for asphalt roofing is Statistics Denmark for consumption of roofing materials, combined with technical specifications for roofing materials produced in Denmark. The emission factors are default factors.

For road paving with asphalt the reference for the activity data is Statistics Denmark for consumption of asphalt and cut-back asphalt. The emission factors are default factors for consumption of asphalt and an estimated emission factor for cut-back asphalt based on the statistics on the emission of NMVOC compiled by the industrial organisations in question.

The reference for activity data for the production of glass and glass wool are obtained from the producers published in their environmental reports. Emission factors are based on stoichiometric relations between raw materials and CO_2 emissions. This information is supplemented with company reports to EU-ETS.

The production of lime and yellow bricks gives rise to CO_2 emissions. The emission factors are based on stoichiometric relations, assumption on CaCO_3 content in clay as well as a default emission factor for expanded clay products. This information is supplemented with company reports to EU-ETS.

There was one producer of nitric acid in Denmark. The data in the inventory relies on information from the producer. The producer reported emissions of NO_x and NH_3 as measured emissions and emissions of N_2O for 2003 as estimated emissions. The emission of N_2O in 2005 and forward is not occurring as the nitric acid production was closed down in the middle of 2004.

There is one producer of catalysts in Denmark. The data in the inventory relies on information published by the producer in environmental reports.

There was one steelwork in Denmark. The activity data as well as data on consumption of raw materials (coke) has been published by the producer in environmental reports. Emission factors are based on stoichiometric relations between raw materials and CO_2 emission. The electro steelwork was closed in 2005.

The inventory on F-gases (HFCs, PFCs and SF_6) is based on work carried out by the Danish Consultant Company "Planmiljø". Their yearly report (DEPA, 2014) documents the inventory data up to the year 2012. The methodology is implemented for the whole time series 1990-2012, but full information on activities only exists since 1995.

Please refer to Chapter 4 for further information on industrial processes.

1.4.5 Solvents

The approach for calculating the emissions of Non-Methane Volatile Organic Carbon (NMVOC) from industrial and household use in Denmark focuses on single chemicals rather than activities. This leads to a clearer picture of the influence from each specific chemical, which enables a more detailed differentiation on products and the influence of product use on emissions. The procedure is to quantify the use of the chemicals and estimate the fraction of the chemicals that is emitted as a consequence of use.

The detailed approach in EMEP/EEA Guidebook (2009) is used. Here all relevant consumption data on all relevant solvents must be inventoried or at least those together representing more than 90 % of the total NMVOC emission. Simple mass balances for calculating the use and emissions of chemicals are set up 1) use = production + import - export, 2) emission = use x emission factor. Production, import and export figures are extracted from Statistics Denmark, from which a list of more than 400 single chemicals, a few groups and products is generated. For each of these, a "use" amount in tonnes per year (from 1990 to 2012) is calculated. For some chemicals and/or products, e.g. propellants used in aerosol cans and ethanol used in wind-screen washing agents, use amounts are obtained from the industry as the information from Statistics Denmark does not comply with required specificity. It is found that approx. 40 different NMVOCs comprise over 95 % of the total use and it is these 40 chemicals that are investigated further. The "use" amounts are distributed across industrial activities according to the Nordic SPIN (Substances in Preparations in Nordic Countries) database, where information on industrial use categories is available in a NACE coding system. The chemicals are also related to specific products according to the Use Category (UCN) system. Emission factors are obtained from regulators, literature or the industry.

Outputs from the inventory are: a list where the approximately 40 most predominant NMVOCs are ranked according to emissions to air; specification of emissions from industrial sectors and from households - contribution from each chemical to emissions from industrial sectors and households; tidal (annual) trend in NMVOC emissions, expressed as total NMVOC and single chemical, and specified in industrial sectors and households.

This emission inventory includes N₂O emissions from the use of anaesthesia for 2000 onwards. Five companies sell N₂O in Denmark and only one company produces N₂O. Due to confidentiality no data on produced amount are available and thus the emissions related to N₂O production are unknown. An emission factor of one is assumed for all use, which equals the sold amount to the emitted amount.

Emissions from other product use such as fireworks, tobacco and charcoal for grilling are included in the inventory. Activity data on consumption of fireworks, tobacco and charcoal are obtained from Statistics Denmark. The emission factors used refer to international literature.

Please refer to Chapter 5 and Annex 3D for further information on the emission inventory for solvent and other product use.

1.4.6 Agriculture

The calculation of emissions from the agricultural sector is based on methods described in the IPCC Guidelines (IPCC, 1996) and the Good Practice Guidance (IPCC, 2000). Activity data for livestock is on a one-year average basis from the agricultural statistics published by Statistics Denmark (2013). Data concerning the land use and crop yield is also from the agricultural statistics. Data concerning the feed consumption and nitrogen excretion is based on information from the Danish Centre for Food and Agriculture (Aarhus University). The CH₄ Implied Emission Factors for Enteric Fermentation and Manure Management are based on a Tier 2/CS approach for all animal categories except for poultry which are based on a Tier 1 approach. All livestock categories in the Danish emission inventory are based on an average of certain subgroups separated by differences in animal breed, age and weight class. The emissions from enteric fermentation for fur farming are estimated to be not applicable.

Emission of N₂O is closely related to the nitrogen balance. Thus, quite a lot of the activity data is related to the Danish calculations for ammonia emission (Mikkelsen et al., 2011). National standards are used to estimate the amount of ammonia emission. When estimating the N₂O emission the IPCC standard value is used for all emission sources. The emission of CO₂ from Agricultural Soils is included in the LULUCF sector.

A model-based system is applied for the calculation of the emissions in Denmark. This model (IDA – Integrated Database model for Agricultural emissions) is used to estimate emission from both greenhouse gases and ammonia. A more detailed description is published in Mikkelsen et al. (2011). The emissions from the agricultural sector are mainly related to livestock production. IDA works on a detailed level and includes around 38 livestock categories, and each category is subdivided according to housing type and manure type. The emissions are calculated from each subcategory and the emissions are aggregated in accordance with the livestock category given in the CRF.

To ensure data quality, both data used as activity data and background data used to estimate the emission factor are collected, and discussed in cooperation with specialists and researchers in different institutions. Thus, the emission inventory will be evaluated continuously according to the latest knowledge. Furthermore, time series of both emission factors and emissions in relation to the CRF categories are prepared. Any considerable variations in the time series are explained.

The uncertainties for assessment of emissions from enteric fermentation, manure management, agricultural soils and field burning of agricultural residue have been estimated based on a Tier 1 and Tier 2 approach. The most significant uncertainties are related to the emissions of N₂O from agricultural soils.

A more detailed description of the methodology for the agricultural sector is given in Chapter 6 and Annex 3E.

1.4.7 Forestry, Land Use and Land Use Change

A complete Land Use Change matrix based on satellite imaging of the whole Danish land area together with cadastral information has been prepared for

the six major area classes. This has improved the coverage and the quality of the inventory substantially.

CO₂ emissions from cropland and grassland are based on census data from Statistics Denmark as regards size of area and crop yield combined with GIS-analysis on land use from the EU agricultural subsidiary system. This gives a very high accuracy for land use. All applicable pools are reported for Cropland and Grassland. The emission from mineral soils for cropland is estimated with a three-pooled dynamical soil carbon model (C-TOOL). C-TOOL was initialised in 1980. The model is run for each region corresponding to former counties in Denmark. Emissions from organic soils in cropland are based on new nationally developed emission factors. For grassland IPCC Tier 1b values are used. National models have been developed for wooden perennial crops in cropland based on land use statistics from Statistic Denmark. These are of minor importance. Sinks in hedgerows are calculated based on a nationally developed model. The area with hedgerows is estimated from information on hedgerows established with financial support from the Danish Government and aerial photos. Emissions from liming are calculated from annual sales data collected by the Danish Agricultural Advisory Centre, combined with the acid neutralisation capacity for each lot produced.

For wetlands emissions are reported from peat extraction areas. Natural wetlands are not reported. A comprehensive programme for restoration of wetlands is implemented in Denmark. Other land uses converted to wetlands is therefore reported.

For the purpose of having estimates for the KP accounting other land uses converted to settlements is reported but not settlements remaining as settlements.

No estimates are made for other land remaining other land and no conversion of land to other land is occurring. For the purpose of having estimates for the KP accounting estimates for living biomass are provided for land converted from other land to other land uses.

1.4.8 Waste

For 6.A Solid Waste Disposal on Land, only managed waste disposal sites are of importance and registered; i.e. unmanaged and illegal disposal of waste is considered to play a negligible role in the context of this category. The CH₄ emission at the Danish SWDSs is based on a First Order Decay (FOD) model according to an IPCC tier 2 approach (IPCC 1997, 2000 and 2006). Data on waste types and amounts deposited at solid waste disposal sites is according to the official registration collected by the Danish Environmental Protection Agency (DEPA, 2013). The model calculations are performed using landfill site characteristics and statistics on the amounts of waste fractions deposited each year. Improved documentation of the methodology, input parameter data including uncertainty analysis is described in Chapter 8.2.

For 6.B Waste Water Handling, country-specific methodologies are used for calculating the emissions of CH₄ and N₂O at wastewater treatment plants (WWTPs). Recent expert review teams (ERTs) in the UNFCCC review have requested better documentation of derived EF and national activity data, and improvements has been performed with respect to dividing the contri-

butions to the net methane emission into specific treatment processes. Fugitive methane releases from the municipal and private WWTPs have been divided into contributions from 1) the sewer system, primary settling tank and biological N and P removal processes, 2) from anaerobic treatment processes in closed systems with biogas extraction and combustion for energy production and 3) septic tanks. N₂O formation and releases during the treatment processes at the WWTPs and from discharged effluent waste water are included. Documentation of the improved methodology, emission factors and activity data are described in Chapter 8.3.

Regarding 6.C Waste Incineration, all municipal, industrial, hazardous and medical waste incinerated is used for energy and heat production. This production is included in the energy statistics, hence emissions are included in the CRF under fuel combustion activities (CRF sector 1A), and more specifically waste incineration takes place in CRF sectors 1A1a, 1A2f and 1A4a. For the 2011 submission reporting in this category covers incineration of corpses and carcasses. The activity data are obtained from the National Association of Danish Crematoria and the three facilities incinerating carcasses.

In CRF category 6.D Other small emissions due to gasification of waste are included for the years 1994-2005. In 2006 onwards these emissions do not occur. In the 2011 submission emissions from accidental fires have been reallocated from category 6C to category 6D.

Please refer to Chapter 8 and Annex 3F for further information on emission inventories for waste.

1.4.9 KP-LULUCF

Regarding the possibility of including in the first commitment period emissions and removals associated with land use, land-use change and forestry activities under Article 3.4 of the Kyoto Protocol, Denmark decided to include emissions and removals from Forest Management (FM), Cropland Management (CM) and Grazing land Management (GM).

The national system has identified land areas associated with the activities under Article 3.4 of the Kyoto Protocol in accordance with definitions, modalities, rules and guidelines relating to land use, land-use change and forestry activities under the protocol by satellite monitoring, use of the EU Land Parcel Information System (LPIS), detailed crop information data on field level, soil mapping and sample plots from the National Forest Inventory (NFI). All land converted from other activities into cropland and grassland is accounted for. No land can leave elected areas under art. 3.4.

The forest definition adopted in the NFI is identical to the FAO definition (TBFRA, 2000). It includes "wooded areas larger than 0.5 ha, that are able to form a forest with a height of at least 5 m and crown cover of at least 10 %". The minimum width is 20 m. For afforestation the carbon stock change in the period 1990 - 2011 is calculated based on the area of afforestation, the information on species composition from the Forest Census 2000 and from the NFI. In the afforestation a steady increase in carbon stock is found. The estimates for the carbon pools in the afforestation are similar to previous estimates, with a slight increase due to the new knowledge on species composition, average carbon stock in those areas based on the NFI data and new data on the carbon stock in soils. Carbon stock change caused by deforestation is estimated based on the deforested area and the mean values of carbon

stock in the total forest area. This is due to the fact that no specific knowledge is available on the carbon pools of the deforested areas. For Forest Management census and NFI data are used.

For cropland and grassland the same methodology is used in the KP reporting as used in the Convention reporting.

Please see Chapter 11 for further details.

1.4.10 Use of EU Emission Trading Scheme data

In 2004 the first guidelines for the monitoring and reporting of greenhouse gas emissions pursuant to the EU Emission Trading Scheme (ETS) Directive (2003/87/EC) were implemented (EU Commission, 2004). These were updated in 2007 and are available from the EU Commission website (EU Commission, 2007).

The Danish emission inventory only includes data from plants using higher tier methods as defined in the EU decision establishing guidelines for monitoring and reporting (EU Commission, 2007). In the Guidelines the specific methods for determining carbon contents, oxidation factor and calorific value are specified.

In the Danish inventory plant or activity based CO₂ emission factors have been derived for power plants combusting coal and oil, refinery gas and flare gas in refineries, fuel gas and flare gas at off-shore installations, cement production, production of brick and tiles and lime production. For all these sources the EU ETS reports are only used in the Danish inventory for plants using high tier methods. The EU ETS data have been applied for the years 2006 onwards.

The EU ETS reporting guidelines emphasizes the need for a high quality reporting through ensuring completeness, consistency, accuracy, transparency and faithfulness. The quality criteria as defined under the EU ETS reporting guidelines are in complete agreement with the principles in the IPCC good practice guidance. For all activities covered by the EU ETS installations are divided into three categories (A, B and C) depending on the annual CO₂ emission. A category A installation has an annual emission of less than 50 Gg CO₂, a category B installation has an annual emission of between 50 and 500 Gg CO₂ and a category C installation has an annual emission of more than 500 Gg CO₂. For each activity Table 1 of the EU ETS guidelines (EU Commission, 2007) specifies the minimum tier level for the different calculation parameters. An example for combustion installations is shown in Table 1.2, the full list for all activities is available in the EU ETS guidelines (EU Commission, 2007).

Table 1.2 Example of minimum requirements in EU ETS guidelines (EU Commission, 2007).

	Activity data						Emission factor			Oxidation factor		
	Fuel flow			Net calorific value								
Activity	A	B	C	A	B	C	A	B	C	A	B	C
Commercial standard fuels	2	3	4	2a/2b	2a/2b	2a/2b	2a/2b	2a/2b	2a/2b	1	1	1
Other gaseous and liquid fuels	2	3	4	2a/2b	2a/2b	3	2a/2b	2a/2b	3	1	1	1
Solid fuels	1	2	3	2a/2b	3	3	2a/2b	3	3	1	1	1

The determination of the variables needed for the emission calculation has to be done in accordance with international standards. It is not possible to list all the relevant standards here, but an overview is available in annex 1, chapter 13 of the EU ETS guidelines. There are also demands concerning sampling methods and frequency of analysis.

As an example the tier 3 regarding fuel flow for fuel combustion, corresponds to a determination of the fuel consumption with an maximum uncertainty of 2.5 % taking into account possible effects of stock change. Tier 4 has a maximum uncertainty of 1.5 %. These uncertainties are very low and are in line with what could be expected from a well-functioning energy statistics system. More information regarding the use of EU ETS data in the specific subsectors of the inventory is included in Chapter 3.2.5 (CHP plants), Chapter 3.5.2 (Refineries and off-shore installations) and Chapter 4.2.2 (Cement production and other mineral products).

The operators shall establish, document, implement and maintain effective data acquisition and handling activities. This means assigning responsibilities for the quality process, as well as quality assurance, reviews and validation of data. Furthermore an independent verification ensuring that emissions have been monitored in accordance with the EU ETS guidelines and that reliable and correct emission data are reported. There are also demands that records and documentation of the control activities must be stored for at least 10 years. The demands for the QA/QC system in the EU ETS guidelines are fully comparable to the requirements in the IPCC good practice guidance. Even so, DCE also performs QC checks of the data received as part of company reporting under EU ETS. This includes comparing the reported parameters with previous years, identifying outliers etc. In case DCE detects what is considered to be outliers DCE contacts the Danish Energy Agency, which is the regulating authority for the EU ETS system in Denmark.

1.5 Brief description of key categories

The key category analysis described in this section covers only Denmark. The aggregation used for the analysis is not directly suited for emissions from Greenland. If Greenlandic emissions were included in the analysis, they would not affect the overall results of the key category analysis. For a key category analysis covering Greenland refer to Chapter 16 and for Denmark and Greenland refer to Chapter 17.

All KCA have been carried out in accordance with Good Practice Guidance (GPG) and IPCC Guidelines.

The KCA for Denmark includes a total of 12 different analyses:

- Base year, reporting year and trend
- Including and excluding LULUCF
- Tier 1 and tier 2 approach

The KCA is based on 153 emission source categories including 22 LULUCF source categories.

The 12 different KCA for Denmark point out 25-35 key source categories each and a total of 52 different key source categories. The number of key cat-

egories in each of the main sectors is: energy 28, industrial processes 3, solvents and other product use 0, agriculture 11, LULUCF 7 and waste 3.

The tier 1 approach point out mainly the large emission sources as key categories and thus CO₂ emission from stationary and mobile combustion are important key categories. The tier 2 approach point out some of the sources with larger uncertainty rates.

Table 1.3 shows the 50 source categories that are key categories in at least one of the six key category analysis including LULUCF. The table includes ranking in the analysis. A similar table for the KCAs excluding LULUCF is included in Annex 1.

The categorisation and detailed results of each of the KCAs are included in Annex 1.

Table 1.3 Key categories for KCAs including LULUCF. The numbers show the ranking in each of the KCAs.

IPCC Source Categories (LU-LUCF included)			Level Tier 1	Level Tier 1	Trend Tier 1	Level Tier 2	Level Tier 2	Trend Tier 2
			1990	2012	1990-2012	1990	2012	1990-2012
Energy	Stationary Combustion, Coal	CO ₂	1	2	1	22	31	27
Energy	Stationary Combustion, Fossil waste	CO ₂	23	8	7		25	20
Energy	Stationary Combustion, Petroleum coke	CO ₂	25	19	19			
Energy	Stationary Combustion, Residual oil	CO ₂	7	21	6			
Energy	Stationary Combustion, Gas oil	CO ₂	3	16	5	23		21
Energy	Stationary Combustion, Kerosene	CO ₂	26		22			
Energy	Stationary Combustion, Refinery gas	CO ₂	18	14	18			
Energy	Stationary Combustion, Natural gas	CO ₂	4	3	2			31
Energy	Natural gas fuelled engines, GAS	CH ₄			28			
Energy	Stationary Combustion, BIOMASS	CH ₄					27	28
Energy	Stationary Combustion, SOLID	N ₂ O				20	30	23
Energy	Stationary Combustion, LIQUID	N ₂ O				11	29	12
Energy	Stationary Combustion, GAS	N ₂ O					23	22
Energy	Stationary Combustion, BIOMASS	N ₂ O				12	5	2
Energy	Transport, Road transport	CO ₂	2	1	3	10	8	9
Energy	Transport, Railways	CO ₂	30	29				
Energy	Transport, Navigation (large vessels)	CO ₂	19	24				
Energy	Transport, Fisheries	CO ₂	22	22				
Energy	Transport, Agriculture	CO ₂	12	9	13	17	15	19
Energy	Transport, Industry (mobile)	CO ₂	17	13	14	14	11	13
Energy	Transport, Commercial/institutional	CO ₂			26			
Energy	Transport, Navigation (large vessels)	N ₂ O				26		
Energy	Transport, Agriculture	N ₂ O				25	24	25
Energy	Transport, Industry (mobile)	N ₂ O					28	30
Energy	1.B.2 Flaring off-shore	CO ₂	29	32				
Energy	1.B.2 Refinery processes	CH ₄						32
Industrial Proc.	2A1 Cement production	CO ₂	16	15	21			
Industrial Proc.	2B2 Nitric acid production	N ₂ O	14		8	21		14
Industrial Proc.	2F Consumption of HFC	HFC		18	11		14	10
Agriculture	4A Enteric Fermentation	CH ₄	5	5	9	7	9	15
Agriculture	4B Manure Management	CH ₄	15	10	10	24	18	17
Agriculture	4.B Manure Management	N ₂ O	21	25		8	12	
Agriculture	4.D1.1 Synthetic Fertilizer	N ₂ O	9	12	12	3	4	3
Agriculture	4.D1.2 Animal waste applied to soils	N ₂ O	13	11	16	5	3	4
Agriculture	4.D1.3 N-fixing crops	N ₂ O		28		19	19	26
Agriculture	4.D1.4 Crop Residue	N ₂ O	27	26		13	16	29
Agriculture	4.D1.5 Cultivation of histosols	N ₂ O		31		18	22	
Agriculture	4.D.2 Grassing animals	N ₂ O	28	30		15	21	
Agriculture	4.D3 Atmospheric deposition	N ₂ O	24	27		9	17	33
Agriculture	4.D3 Leaching	N ₂ O	8	7	23	2	2	8
LULUCF	5.A.1 Forest remaining forest	CO ₂		4	4		7	1
LULUCF	5.B Cropland, Living biomass	CO ₂		33	20		13	5
LULUCF	5.B Cropland, Mineral soils	CO ₂	10	20	17	6	10	6
LULUCF	5.B Cropland, Organic soils	CO ₂	6	6		1	1	
LULUCF	5.C Grassland, Living biomass	CO ₂		23	15		20	11
LULUCF	5.D Wetlands, Soils	CO ₂						34
LULUCF	5(IV) Cropland Limestone	CO ₂	20		25	16		18
Waste	6 A. Solid Waste Disposal on Land	CH ₄	11	17	24	4	6	7
Waste	6.D Compost production	CH ₄					32	24
Waste	6.D Compost production	N ₂ O			27		26	16

1.5.1 KP-LULUCF

See Chapter 11.9.1 for discussion on the key category analysis of KP-LULUCF.

1.6 Information on QA/QC plan including verification and treatment of confidential issues where relevant

1.6.1 Introduction

This section outlines the Quality Control (QC) and Quality Assurance (QA) plan for greenhouse gas emission inventories performed by DCE (Sørensen et al., 2005; Nielsen et al., 2013). The plan is in accordance with the guidelines provided by the IPCC (IPCC, 1996), and the Good Practice Guidance and Uncertainty Management in National Greenhouse Gas Inventories (IPCC, 2000). The ISO 9000 standards are also used as important input for the plan.

The QA/QC plan also covers Greenland. DCE receives the data corresponding to data processing level 3 and data storage level 4 and the data undergoes the same QA/QC procedure as the Danish data, some further QC checks are described in Chapter 17. The QA/QC specific to the Greenlandic emission inventory is described in Chapter 16.

1.6.2 Concepts of quality work

The quality planning is based on the following definitions as outlined by the ISO 9000 standards as well as the Good Practice Guidance (IPCC, 2000):

- Quality management (QM) Coordinates activity to direct and control with regard to quality.
- Quality Planning (QP) Defines quality objectives including specification of necessary operational processes and resources to fulfil the quality objectives.
- Quality Control (QC) Fulfils quality requirements.
- Quality Assurance (QA) Provides confidence that quality requirements will be fulfilled.
- Quality Improvement (QI) Increases the ability to fulfil quality requirements.

The activities are considered inter-related in this report as shown in Figure 1.2.




Figure 1.2 Interrelation between the activities with regard to quality. The arrows are explained in the text below this figure.

1: The QP sets up the objectives and, from these, measurable properties valid for the QC.

- 2: The QC investigates the measurable properties that are communicated to QA for assessment in order to ensure sufficient quality.
3. The QP identifies and defines measurable indicators for the fulfilment of the quality objectives. This yields the basis for the QA and has to be supported by the input coming from the QC.
- 4: The result from QC highlights the degree of fulfilment for every quality objective. It is thus a good basis for suggestions for improvements to the inventory to meet the quality objectives.
- 5: Suggested improvements in the quality may induce changes in the quality objectives and their measurability.
- 6: The evaluation carried out by external authorities is important input when improvements in quality are being considered.

1.6.3 Definition of quality

A solid definition of quality is essential. Without such a solid definition, the fulfilment of the objectives will never be clear and the process of quality control and assurance can easily turn out to be a fuzzy and unpleasant experience for the people involved. On the contrary, in case of a solid definition and thus a clear goal, it will be possible to make a valid statement of “good quality” and thus form constructive conditions and motivate the inventory work positively. A clear definition of quality has not been given in the UN-FCCCC guidelines. In the Good Practice Guidance, Chapter 8.2, however, it is mentioned that:

“Quality control requirements, improved accuracy and reduced uncertainty need to be balanced against requirements for timeliness and cost effectiveness.” The statement of balancing requirements and costs is not a solid basis for QC as long as this balancing is not well defined.

The resulting standard of the inventory is defined as being composed of accuracy and regulatory usefulness. The goal is to maximise the standard of the inventory and the following statement defines the quality objective:

The quality objective is only inadequately fulfilled if it is possible to make an inventory of a higher standard without exceeding the frame of resources.

1.6.4 Definition of Critical Control Points (CCP)

A Critical Control Point (CCP) is defined in this submission as an element or an action which needs to be taken into account in order to fulfil the quality objectives. Every CCP has to be necessary for the objectives and the CCP list needs to be extended if other factors, not defined by the CCP list, are needed in order to reach at least one of the quality objectives.

The objectives for the QM, as formulated by IPCC (2000), are to improve elements of transparency, consistency, comparability, completeness and confidence. In the IPCC guidelines (IPCC, 1996), the element “confidence” is replaced by “accuracy” and in this plan “accuracy” is used.

The objectives for the *QM* are used as *CCPs*, including the elements mentioned above. The following explanation is given by IPCC guidelines (IPCC, 1996) for each *CCP*:

Transparency means that the assumptions and methodologies used for an inventory should be clearly explained to facilitate replication and assessment of the inventory by users of the reported information. The transparency of the inventories is fundamental to the success of the process for communication and consideration.

Consistency means that an inventory should be internally consistent in all its elements with inventories of other years. An inventory is consistent if the same methodologies are used for the base and for all subsequent years and if consistent datasets are used to estimate emissions or removals from source or sinks. Under certain circumstances, an inventory using different methodologies for different years can be considered to be consistent if it has been recalculated in a transparent manner in accordance with the Intergovernmental Panel on Climate Change (IPCC) guidelines and good practice guidance.

Comparability means that estimates of emission and removals reported by Annex I Parties in inventories should be comparable among Annex I parties. For this purpose, Annex I Parties should use the methodologies and formats agreed upon by the COP for estimating and reporting inventories. The allocation of different source/sink categories should follow the split of *Revised 1996 IPCC Guidelines for national Greenhouse Gas Inventories* (IPCC, 1996) at the level of its summary and sectoral tables.

Completeness means that an inventory covers all sources and sinks, as well as all gases, included in the IPCC guidelines as well as other existing relevant source/sink categories, which are specific to individual Annex I Parties and, therefore, may not be included in the IPCC guidelines. Completeness also means full geographic coverage of sources and sinks of an Annex I Party.

Accuracy is a relative measure of the exactness of an emission or removal estimate. Estimates should be accurate and should systematically neither over- nor underestimate emissions nor removals. Uncertainties on estimates should be reduced if possible. Appropriate methodologies should be used in accordance with the *IPCC good practice guidance*, to promote data accuracy in inventories.

The robustness against unexpected disturbance of the inventory work has to be high in order to secure high quality, which is not covered by the *CCPs* above. The correctness of the inventory is formulated as an independent objective. This is so because the correctness of the inventory is a condition for all other objectives to be effective. A large part of the Tier 1 procedure given by the Good Practice Guidance (IPCC, 2000) is actually checks for miscalculations and, thus, supports the objective of correctness. Correctness, as defined here, is not similar to accuracy, because the correctness takes into account miscalculations, while accuracy relates to minimizing the always present data-value uncertainty.

Robustness implies arrangement of inventory work as regards e.g. inventory experts and data sources in order to minimize the consequences of any unexpected disturbance due to external and internal conditions. A change in an external condition could be interruption of access to an external data source

and an internal change could be a sudden reduction in qualified staff, where a skilled person suddenly leaves the inventory work.

Correctness has to be secured in order to avoid uncontrollable occurrence of uncertainty directly due to errors in the calculations.

The different CCPs are not independent and represent different degrees of generality. E.g. deviation from *comparability* may be accepted if a high degree of *transparency* is applied. Furthermore, there may even be a conflict between the different CCPs. E.g. new knowledge may suggest improvements in calculation methods for better *completeness*, but the same improvements may to some degree violate the *consistency* and *comparability* criteria with regard to earlier years' inventories and the reporting from other nations. It is, therefore, a multi-criteria problem of optimisation to apply the set of CCPs in the aim for good quality.

1.6.5 Process-oriented QC

The strategy is based on a process-oriented principle (ISO 9000 series) and the first step is, thus, to set up a system for the process of the inventory work. The product specification for the inventory is a dataset of emission figures and the process, thereby, equates with the data flow in the preparation of the inventory.

The data flow needs to support the QC/QA in order to facilitate a cost-effective procedure. The flow of data has to take place in a transparent way by making the transformation of data detectable. It should be easy to find the original background data for any calculation and to trace the sequence of calculations from the raw data to the final emission result. Computer programming for automated calculations and checking will enhance the accuracy and minimize the number of miscalculations and flaws in input value settings. Especially manual typing of numbers needs to be minimized. This assumes, however, that the quality of the programming has been verified to ensure the correctness of the automated calculations. Automated value control is also one of the important means to secure accuracy. Realistic uncertainty estimates are necessary for securing accuracy, but they can be difficult to produce due to the uncertainty related to the uncertainty estimates themselves. It is, therefore, important to include the uncertainty calculation procedures into the data structure as far as possible. The QC/QA needs to be supported as far as possible by the data structure; otherwise the procedures can easily become troublesome and subject to frustration.

Both data processing and data storage form the data structure. The data processing is carried out using mathematical operations or models. The models may be complicated where they concern human activity or be simple summations of lower aggregated data. The data storage includes databases and file systems of data that are either calculated using the data processing at the lower level, using input to new processing steps or even using both output and input in the data structure. The measure for quality is basically different for processing and storage, so these need to be kept separate in a well-designed quality manual. A graphical display of the data flow is seen in Figure 1.3 and explained in the following.

The data storage takes place for the following types of data:

External Data: a single numerical value of a parameter coming from an external source. These data govern the calculation of *Emission calculation input*.

Emission calculation input: Data for input to the final emission calculation in terms of data for release source strength and activity. The data is directly applicable for use in the standardized forms for calculation. These data are calculated using external data or represent a direct use of *External Data* when they are directly applicable for *Emission Calculations*.

Emission Data: Estimated emissions based on the *emission calculation input*.

Emission Reporting: Reporting of emission data in requested formats and aggregation level.




Figure 1.3 The general data structure for the emission inventory.

Key levels are defined in the data structure as:

Data storage Level 1, External data

Collection of external data for calculation of emission factors and activity data. The activity data are collected from different sectors and statistical surveys, typically reported on a yearly basis. The data consist of raw data, having an identical format to the data received and gathered from external sources. Level 1 data acts as a base-set, on which all subsequent calculations are based. If alterations in calculation procedures are made, they are based on the same dataset. When new data are introduced they can be implemented in accordance with the QA/QC structure of the inventory.

Data storage Level 2, Data directly usable for the inventory

This level represents data that have been prepared and compiled in a form that is directly applicable for calculation of emissions. The compiled data are structured in a database for internal use as a link between more or less raw data and data that are ready for reporting. The data are compiled in a way that elucidates the different approaches in emission assessment: (1) directly on measured emission rates, especially for larger point sources, (2) based on activities and emission factors, where the value setting of these factors are stored at this level.

Data storage Level 3 Emission data

The emission calculations are reported by the most detailed figures and divided in sectors. The unit at this level is typically mass pr yr for the country. For sources included in the SNAP system, the SNAP level 3 is relevant. Internal reporting is performed at this level to feed the external communication of results.

Data storage Level 4 Final reports for all subcategories

The complete emission inventory is reported to UNFCCC at this level by summing up the results from every subcategory.

Data processing Level 1 Compilation of external data

Preparation of input data for the emission inventory based on the external data sources. Some external data may be used directly as input to the data processing at level 2, while other data needs to be interpreted using more or less complicated models, which takes place at this level. The interpretation of activity data is to be seen in connection with availability of emission factors and vice versa. These models are compiled and processed as an integrated part of the inventory preparation.

Data processing Level 2 Calculation of inventory figures

The emission for every subcategory is calculated, including the uncertainty for all sectors and activities. The summation of all contributions from sub-sources makes up the inventory.

Data processing Level 3 Calculation aggregated parameters

Some aggregated parameters need to be reported as part of the final reporting. This does not involve complicated calculations but important figures, e.g. implied emission factors at a higher aggregated level to be compared in time series and with other countries.

1.6.6 Definition of Point of Measurements (PM)

The CCPs have to be based on clear measurable factors - otherwise the *QP* will end up being just a loose declaration of intent. Thus, in the following, a series of *Points for Measuring (PM)* is identified as building blocks for a solid QC. Table 8.1 in Good Practice Guidance is a listing of such *PMs*. However, the listing in Table 1.2 is an extended and modified listing, in comparison to Table 8.1 in the Good Practice Guidance supporting all the *CCPs*. The *PMs* will be routinely checked in the QC reporting and, when external reviews take place, the reviewers will be asked to assess the fulfilment of the *PMs* using a checklist system. The list of *PMs* is continually evaluated and modified to offer the best possible support for the *CCPs*. The actual list used is seen in Table 1.2.

Table 1.2 The list of *PMs* as used.

Level	CCP	Id	Description	
Data Storage level 1	1. Accuracy	DS.1.1.1	General level of uncertainty for every dataset including the reasoning for the specific values	Sectoral
		DS.1.1.2	Quantification of the uncertainty level of every single data value, including the reasoning for the specific values.	Sectoral
	2. Comparability	DS.1.2.1	Comparability of the data values with similar data from other countries, which are comparable with Denmark, and evaluation of the discrepancy.	Sectoral
	3. Completeness	DS.1.3.1	Documentation showing that all possible national data sources are included, by setting down the reasoning behind the selection of datasets.	Sectoral
	4. Consistency	DS.1.4.1	The origin of external data has to be preserved whenever possible without explicit arguments (referring to other PMs)	Sectoral
	6. Robustness	DS.1.6.1	Explicit agreements between the external institution holding the data and DCE about the conditions of delivery	Sectoral
		DS.1.6.2	At least two employees must have a detailed insight into the gathering of every external dataset.	General
	7. Transparency	DS.1.7.1	Summary of each dataset including the reasoning behind the selection of the specific dataset	Sectoral
		DS.1.7.2	The archiving of datasets needs to be easily accessible for any person in the emission inventory	General
		DS.1.7.3	References for citation for any external dataset have to be available for any single number in any dataset.	Sectoral
		DS.1.7.4	Listing of external contacts for every dataset	Sectoral
Data Processing level 1	1. Accuracy	DP.1.1.1	Uncertainty assessment for every data source as input to Data Storage level 2 in relation to type of variability. (Distribution as: normal, log normal or other type of variability)	Sectoral
		DP.1.1.2	Uncertainty assessment for every data source as input to Data Storage level 2 in relation to scale of variability (size of variation intervals)	Sectoral
		DP.1.1.3	Evaluation of the methodological approach using international guidelines	Sectoral
		DP.1.1.4	Verification of calculation results using guideline values	Sectoral
	2. Comparability	DP.1.2.1	The inventory calculation has to follow the international guidelines suggested by UNFCCC and IPCC.	Sectoral
	3. Completeness	DP.1.3.1	Assessment of the most important quantitative knowledge which is lacking.	Sectoral
		DP.1.3.2	Assessment of the most important cases where access is lacking with regard to critical data sources that could improve quantitative knowledge.	Sectoral
	4. Consistency	DP.1.4.1	In order to keep consistency at a high level, an explicit description of the activities needs to accompany any change in the calculation procedure	Sectoral
		DP.1.4.2	Identification of parameters (e.g. activity data, constants) that are common to multiple source categories and confirmation that there is consistency in the values used for these parameters in the emission calculations	General
	5. Correctness	DP.1.5.1	Shows at least once, by independent calculation, the correctness of every data manipulation	Sectoral
		DP.1.5.2	Verification of calculation results using time series	Sectoral
		DP.1.5.3	Verification of calculation results using other measures	Sectoral

Level	CCP	Id	Description	
	6.Robustness 7.Transparency	DP.1.5.4	Show one-to-one correctness between external data sources and the databases at Data Storage level 2	Sectoral
		DP.1.6.1	Any calculation must be anchored to two responsible persons who can replace each other in the technical issue of performing the calculations.	General
		DP.1.7.1	The calculation principle and equations used must be described	Sectoral
		DP.1.7.2	The theoretical reasoning for all methods must be described	Sectoral
		DP.1.7.3	Explicit listing of assumptions behind all methods	Sectoral
		DP.1.7.4	Clear reference to dataset at Data Storage level 1	Sectoral
		DP.1.7.5	A manual log to collect information about recalculations	Sectoral
Data Storage level 2	2.Comparability	DS.2.2.1	Comparison with other countries that are closely related to Denmark and explanation of the largest discrepancies	General
	5.Correctness	DS.2.5.1	Documentation of a correct connection between all data types at level 2 to data at level 1	Sectoral
		DS.2.5.2	Check if a correct data import to level 2 has been made	Sectoral
	6.Robustness	DS.2.6.1	All persons in the inventory work must be able to handle and understand all data at level 2.	General
	7.Transparency	DS.2.7.1	The time trend for every single parameter must be graphically available and easy to map	General
Data Processing level 2	1. Accuracy	DP.2.1.1	Documentation of the methodological approach for the uncertainty analysis	General
		DP.2.1.2	Quantification of uncertainty	General
	2.Comparability	DP.2.2.1	The inventory calculation has to follow the international guidelines suggested by UNFCCC and IPCC	General
	6.Robustness	DP.2.6.1	Any calculation at level 4 must be anchored to two responsible persons who can replace each other in the technical issue of performing the calculations.	General
	7.Transparency	DP.2.7.1	Reporting of the calculation principle and equations used	General
		DP.2.7.2	The reasoning for the choice of methodology for uncertainty analysis needs to be written explicitly.	General
Data Storage level 3	1. Accuracy	DS.3.1.1	Quantification of uncertainty	General
	5.Correctness	DS.3.5.1	Comparison with inventories of the previous years on the level of the categories of the CRF as well as on SNAP source categories. Any major changes are checked, verified, etc.	General
		DS.3.5.2	Total emissions, when aggregated to CRF source categories, are compared with totals based on SNAP source categories (control of data transfer).	General
		DS.3.5.3	Checking of time series of the CRF and SNAP source categories as they are found in the Corinair databases. Considerable trends and changes are checked and explained.	General
	7. Transparency	DS.3.7.1	The databases and other software used shall be clearly documented. The documentation should include a description that the appropriate data processing steps are correctly represented in the database; that data relationships are correctly represented in the database and that data fields are properly labelled and have the correct design specifications.	General

Level	CCP	Id	Description	
		DS.3.7.2	The documentation referred to under DS.3.7.1 should be archived at the same network folder as the program is located in.	General
Data Processing level 3	6. Robustness	DP.3.6.1	The process of generating the official submissions must be anchored by at least two responsible persons who can replace each other in the technical issue of generating CRF tables including of the aggregation of submissions for Denmark and Greenland.	General
	7. Transparency	DP.3.7.1	The databases and other software used shall be clearly documented. The documentation should include a description that the appropriate data processing steps are correctly represented in the database; that data relationships are correctly represented in the database and that data fields are properly labelled and have the correct design specifications.	General
	7. Transparency	DP.3.7.2	The documentation referred to under DP.3.7.1 should be archived at the same network folder as the program is located in.	General
Data Storage level 4	2.Comparability	DS.4.2.1	Description of similarities and differences in relation to other countries' inventories for the methodological approach.	General
	3.Completeness	DS.4.3.1	National and international verification including explanation of the discrepancies.	General
		DS.4.3.2	Check that the no sources where a methodology exists in the IPCC guidelines are reported as NE.	General
	4.Consistency	DS.4.4.1	The inventory reporting must follow the international guidelines suggested by UNFCCC and IPCC.	General
		DS.4.4.2	Check time series consistency of the reporting by Greenland and the Faroe Islands prior to aggregating the final submissions.	General
		DS.4.4.3	The IEFs from the CRF are checked both regarding level and trend. The level is compared to relevant emission factors to ensure correctness. Large dips/jumps in the time series are explained.	Sectoral
	5.Correctness	DS.4.5.1	Check that the aggregated submissions for Denmark under the Kyoto Protocol and the UNFCCC match the sum of the individual submissions.	General
		DS.4.5.2	Check that additional information and information related to land-use changes has been correctly aggregated compared to the individual submissions of Denmark and Greenland.	Sectoral
	6. Robustness	DS.4.6.1	The reporting to the UNFCCC must be anchored to two responsible persons who can replace each other in the technical issue of reporting to and communicating with the UNFCCC secretariat.	General
	7.Transparency	DS.4.7.1	Perform QA on the documentation report provided by the Government of Greenland.	General

1.6.7 Plan for the quality work

The IPCC uses the concept of a tiered approach, i.e. a stepwise approach, where complexity, advancement and comprehensiveness increase. Generally, more detailed and advanced methods are recommended in order to give guidance to countries which have more detailed datasets and more capacity, as well as to countries with less available data and manpower. The tiered approach helps to focus attention on the areas of the inventories that are relatively weak, rather than investing effort in irrelevant areas. Furthermore, the IPCC guidelines recommend using higher tier methods for key categories.

ries in particular. Therefore, the identification of key categories is crucial for planning quality work. However, there exist several issues regarding the listing of priority categories: (1) The contribution to the total emission figure (key source listing); (2) The contribution to the total uncertainty; (3) Most critical categories in relation to implementation of new methodologies and thus highest risk for miscalculations. All the points listed are necessary for different aspects of producing high quality work. These listings will be used to secure implementation of the full quality scheme for the most relevant categories. Verification in relation to other countries has been undertaken for priority categories.

1.6.8 Implementation of the QA/QC plan

The PMs listed in Table 1.2 are described for each sector in the QA/QC sections of Chapters 3-8, where a status with regard to implementation is also given. Some of the PMs are the same for all sectors and a common description for these PMs is given in Section 1.6.10, below. The focus has been on level 1 for both data storage and data processing as this is the most labour-intensive part. The quality system will be evaluated and adjusted continuously.

1.6.9 Archiving of data and documentations

The QA/QC work is supported by an inventory file system, where all data, models and QA/QC procedures and checks are stored as files in folders (Figure 1.4).



Figure 1.4 Schematic diagram of the folder structure in the inventory file system.

The inventory file system consists of the following levels: year, sector and the level for the process of the inventory work, as illustrated in Figure 1.4. The first level in the file system is year, which here means the inventory year

and not the calendar year. The sector level contains the PMs relevant for the individual sectors i.e. the first levels (DS1 and DP1) (except the PMs described in Section 1.6.10), while the rest of the PMs (DS2-4 and DP2-3), are common for all sectors.

All data, models and other QA/QC related files are stored in the inventory file system and are accessible for all staff involved in the inventory work.

1.6.10 Common QA/QC PMs

The following PMs are common for all the sectors:

Data storage Level 1

Data Storage level 1	6. Robustness	DS.1.6.2	At least two employees must have a detailed insight into the gathering of every external dataset.
----------------------	---------------	----------	---

For all sectors: energy, industrial processes, solvent and other product use, agriculture, LULUCF and waste, two persons have detailed insight in data gathering and processing. A strong effort is continuously made to ensure the robustness of the inventory process.

Data Storage level 1	7. Transparency	DS.1.7.2	The archiving of datasets needs to be easily accessible for any person involved in the emission inventory.
----------------------	-----------------	----------	--

All data, models and other QA/QC related files are stored in the inventory file system and are accessible for all inventory staff members. Refer to Section 1.6.9.

Data processing Level 1

Data Processing level 1	4. Consistency	DP.1.4.2	Identification of parameters (e.g. activity data, constants) that are common to multiple source categories and confirmation that there is consistency in the values used for these parameters in the emission calculations.
-------------------------	----------------	----------	---

This PM is supported by the inventory file system where it is possible to compare and harmonise parameters that are common to multiple source categories.

Data Processing level 1	6. Robustness	DP.1.6.1	Any calculation must be anchored to two responsible persons who can replace each other in the technical issue of performing the calculations.
-------------------------	---------------	----------	---

All data, models and other QA/QC related files are stored in the inventory file system and are accessible for all inventory staff members. Refer to Section 1.6.9.

Data storage Level 2

Data Storage level 2	2. Comparability	DS.2.2.1	Comparison with other countries that are closely related to Denmark and explanation of the largest discrepancies.
----------------------	------------------	----------	---

Systematic inter-country comparison has only been made on data storage level 4. Refer to DS 4.3.2.

Data Storage level 2	6.Robustness	DS.2.6.1	All persons in the inventory work must be able to handle and understand all data at level 2.
----------------------	--------------	----------	--

This PM is fulfilled for all sectors. The PM is supported by the inventory file system. Refer to Section 1.6.9.

Data Storage level 2	7.Transparency	DS.2.7.1	The time trend for every single parameter must be graphically available and easy to map.
----------------------	----------------	----------	--

Programs exist to make time series for all parameters. A tool for graphically showing time series has not yet been developed.

Data Processing Level 2

Data Processing level 2	1. Accuracy	DP.2.1.1	Documentation of the methodological approach for the uncertainty analysis
-------------------------	-------------	----------	---

Refer to Chapter 1.7.

Data Processing level 2	1. Accuracy	DP.2.1.2	Quantification of uncertainty
-------------------------	-------------	----------	-------------------------------

Refer to Chapter 1.7 and the uncertainty sections in the sectoral chapters (Chapter 3-8).

Data Processing level 2	2.Comparability	DP.2.2.1	The inventory calculation has to follow the international guidelines suggested by UN-FCCC and IPCC.
-------------------------	-----------------	----------	---

The emission calculations follow the international guidelines.

Data Processing level 2	6.Robustness	DS.2.6.1	All persons in the inventory work must be able to handle and understand all data at level 2.
-------------------------	--------------	----------	--

At present the emission calculations are carried out using applications developed at DCE. The software development and programme runs are anchored to two inventory staff members.

Data Processing level 2	7.Transparency	DP.2.7.1	Reporting of the calculation principle and equations used.
-------------------------	----------------	----------	--

Due to the uniform treatment of input data in the calculation routines used by the DCE software programmes, a central documentation of calculation principles, equations, theoretical reasoning and assumptions must be given, treating all national emission sources. This documentation still remains to be made, but is planned to be carried out in the future.

Data Processing level 2	7. Transparency	DP.2.7.2	The reasoning for the choice of methodology for uncertainty analysis needs to be written explicitly.
-------------------------	-----------------	----------	--

Refer to Chapter 1.7 and the QA/QC sections in the sectoral chapters.

Data storage Level 3

Data Storage level 3	1. Accuracy	DS.3.1.1	Quantification of uncertainty
----------------------	-------------	----------	-------------------------------

Refer to Chapter 1.7 and the QA/QC sections in the sector chapters.

Data Storage level 3	5. Correctness	DS.3.5.1	Comparison with inventories of the previous years on the level of the categories of the CRF as well as on SNAP source categories. Any major changes are checked, verified, etc.
----------------------	----------------	----------	---

Time series is prepared and checked, any major change is closely examined with the purpose of verifying and explaining changes from earlier inventories.

Data Storage level 3	5. Correctness	DS.3.5.2	Total emissions when aggregated to CRF source categories are compared with totals based on SNAP source categories (control of data transfer).
----------------------	----------------	----------	---

Total emission, when aggregated to IPCC and LRTAP reporting tables, is compared with totals based on SNAP source categories (control of data transfer).

Data Storage level 3	5. Correctness	DS.3.5.3	Checking of time series of the CRF and SNAP source categories as they are found in the Corinair databases. Considerable trends and changes are checked and explained.
----------------------	----------------	----------	---

Time series are prepared and checked, any major change is closely examined with the purpose of verifying and explaining fluctuations.

Data Storage level 3	7. Transparency	DS.3.7.1	The databases and other software used shall be clearly documented. The documentation should include a description that the appropriate data processing steps are correctly represented in the database; that data relationships are correctly represented in the database and that data fields are properly labelled and have the correct design specifications.
----------------------	-----------------	----------	--

The databases used at data storage level 3 are documented. The documentation includes description of the queries and programming code used in the data processing. The documentation further includes information on all data fields in the database and the design specifications. Part of the detailed documentation is built into the database while the overall documentation is prepared as a separate documentation note.

Data Storage level 3	7. Transparency	DS.3.7.2	The documentation referred to under DS.3.7.1 should be archived at the same network folder as the program is located in.
----------------------	-----------------	----------	--

The documentation prepared as part of DS.3.7.1 is archived in the same folder as the program is stored. For information on the file structure, please see Chapter 1.6.9.

Data Processing Level 3

Data Processing level 3	6. Robustness	DP.3.6.1	The process of generating the official submissions must be anchored by at least two responsible persons who can replace each other in the technical issue of generating CRF tables including of the aggregation of submissions for Denmark and Greenland.
-------------------------	---------------	----------	---

The process of generating the official submissions including the aggregation of submissions to the UNFCCC and the Kyoto Protocol is currently anchored by two people within the team. In the future the goal is to have three team members capable of completing this task.

Data Processing level 3	7. Transparency	DP.3.7.1	The databases and other software used shall be clearly documented. The documentation should include a description that the appropriate data processing steps are correctly represented in the database; that data relationships are correctly represented in the database and that data fields are properly labelled and have the correct design specifications.
-------------------------	-----------------	----------	--

The databases used at data storage level 3 are documented. The documentation includes description of the queries and programming code used in the data processing. The documentation further includes information on all data fields in the database and the design specifications. Part of the detailed documentation is built into the database while the overall documentation is prepared as a separate documentation note.

Data Processing level 3	7. Transparency	DP.3.7.2	The documentation referred to under DS.3.7.1 should be archived at the same network folder as the program is located in.
-------------------------	-----------------	----------	--

The documentation prepared as part of DS.3.7.1 is archived in the same folder as the program is stored. For information on the file structure, please see Chapter 1.6.9.

Data Storage Level 4

Data Storage level 4	2. Comparability	DS.4.2.1	Description of similarities and differences in relation to other countries' inventories for the methodological approach
----------------------	------------------	----------	---

For each key source category, a comparison has been made between Denmark and the EU-15 countries (Fauser et al., 2007 & 2013). This is performed by comparing emission density indicators, defined as emission intensity value divided by a chosen indicator. The indicators are identical to the ones identified in the Norwegian verification inventory (Holtskog et al., 2000). The correlation between emissions and an independent indicator does not necessarily imply cause and effect, but in cases where the indicator is direct-

ly associated with the emission intensity value, such as for the energy sector, the emission density indicator is a measure of the implied emission factor and a direct comparison can be made. A qualitative verification of implied emission factors can, furthermore, be made when a measured or theoretical value of the CO₂ content in the respective fuel type (or other relevant parameter) is available. For the energy sector, all countries are, in principle, comparable and inter-country deviations arise from variations in fuel purities and fuel combustion efficiencies. A comparison of national emission density indicators, analogous to the implied emission factors, will give valuable information on the quality and efficiency of the national energy sectors.

Furthermore, the inter-country comparison of emission density indicators and comparison of theoretical values gives a methodological verification of the derivation of emission intensity values, and of the correlation between emission intensity values and activity values.

When emissions are compared with non-dependent parameters, similarities with regard to geography, climate, industry structure and level of economic development may be necessary for obtaining comparable emission density indicators.

Data Storage level 4	3.Completeness	DS.4.3.1	National and international validation including explanation of the discrepancies.
----------------------	----------------	----------	---

Refer to DS 4.2.1

Data Storage level 4	3.Completeness	DS.4.3.2	Check that the no sources where a methodology exists in the IPCC guidelines are reported as NE.
----------------------	----------------	----------	---

It is verified both by DCE experts and by EU consistency checks that no sources where methodologies and default parameters exist have been reported as NE. If methodologies do exist efforts are made to estimate and report emissions.

Data Storage level 4	4.Consistency	DS.4.4.1	The inventory reporting must follow the international guidelines suggested by UNFCCC and IPCC.
----------------------	---------------	----------	--

The inventory reporting is in accordance with the UNFCCC guidelines on reporting and review (UNFCCC, 2007). The present report includes detailed and complete information on the inventories for all years from the base year to the year of the current annual inventory submission, in order to ensure the transparency of the inventory. The annual emission inventory for Denmark is reported in the Common Reporting Format (CRF) as requested in the reporting guidelines. The CRF-spreadsheets contain data on emissions, activity data and implied emission factors for each year. Emission trends are given for each greenhouse gas and for total greenhouse gas emissions in CO₂ equivalents. The link to complete sets of CRF-files and more information on the Danish emission inventories are on the ENVS homepage (<http://envs.au.dk/videnudveksling/luft/emissioner/emissioninventory>).

Data Storage level 4	4.Consistency	DS.4.4.2	Check time series consistency of the reporting of Greenland and the Faroe Islands prior to aggregating the final submissions
----------------------	---------------	----------	--

The time series for all pollutants in the submissions from Greenland and the Faroe Islands are checked at the CRF 3 level for large variations in the time series. Any large variations are explained or corrected in cooperation with the authorities in Greenland and the Faroe Islands.

Data Storage level 4	5. Correctness	DS.4.5.1	Check that the aggregated submissions for Denmark under the Kyoto Protocol and the UNFCCC matches the sum of the individual submissions
----------------------	----------------	----------	---

To ensure that the submission for Denmark under the Kyoto Protocol matches the sum of the submissions of Denmark and Greenland a spreadsheet check has been implemented to ensure complete correctness of the submitted inventory. The same procedure is followed for the submission under the UNFCCC, where it is ensured that the submitted emissions equate to the sum of Denmark, Greenland and the Faroe Islands. Special attention is paid to the additional information provided in the CRF, e.g. for the agricultural sector. Certain parameters cannot simply be added, e.g. animal weights. In these cases a weighted average is reported in the CRF tables.

Data Storage level 4	6. Robustness	DS.4.6.1	The reporting to the UNFCCC must be anchored to two responsible persons who can replace each other in the technical issue of reporting to and communicating with the UNFCCC secretariat.
----------------------	---------------	----------	--

The reporting to the UNFCCC secretariat is currently anchored by two team members. All official correspondence between the secretariat and DCE involves both the responsible team members.

Data Storage level 4	7. Transparency	DS.4.7.1	Perform QA on the documentation report provided by the Government of Greenland
----------------------	-----------------	----------	--

The documentation report is received by DCE from the Government of Greenland in the early spring every year. The documentation report is included in the NIR as Chapter 16. DCE experts read and provide comments on the report to the Government of Greenland, so that any questions are resolved prior to the UNFCCC reporting deadline of April 15.

1.7 General uncertainty evaluation, including data on the overall uncertainty for the inventory totals

1.7.1 Tier 1 uncertainties

The uncertainty estimates are based on the Tier 1 methodology in the IPCC Good Practice Guidance (GPG) (IPCC, 2000). Uncertainty estimates for the following sectors are included in the current year: stationary combustion plants, mobile combustion, fugitive emissions from fuels, industry, solid waste and wastewater treatment, CO₂ from solvents, agriculture and LU-LUCF. The sources included in the uncertainty estimate cover 100 % of the total net Danish greenhouse gas emissions and removals.

The uncertainties for the activity rates and emission factors are shown in Table 1.3.

Table 1.3 Summary of base year and 2012 emissions in Gg CO₂ eqv. and activity data and emission factor uncertainties. Calculated Tier 1 and Tier 2 uncertainties for each emission source are given as % of the total 2012 emission. The base year for F-gases is 1995 and for all other gases the base year is 1990. Tier 2 uncertainty is not calculated for LULUCF.

IPCC Source category	Gas	Base year emission Gg CO ₂ eqv.	2012 emission Gg CO ₂ eqv.	Activity data uncer- tainty %	Emission factor uncertain- ty %	Tier 1 Combined uncertainty % of total emissions	Tier 2 uncertainty % of total emissions
Stationary Combustion, Coal	CO ₂	23 834	10 005	1	1	0.203	0.190
Stationary Combustion, BKB	CO ₂	11	1	3	5	0.0002	0.0002
Stationary Combustion, Coke	CO ₂	138	74	2	5	0.008	0.0075
Stationary Combustion, Fossil waste	CO ₂	573	1397	5	10	0.307	0.309
Stationary Combustion, Petroleum coke	CO ₂	415	628	2	5	0.067	0.064
Stationary Combustion, Residual oil	CO ₂	2496	571	1	2	0.026	0.025
Stationary Combustion, Gas oil	CO ₂	4547	742	2	4	0.067	0.064
Stationary Combustion, Kerosene	CO ₂	366	2	2	5	0.0002	0.0002
Stationary Combustion, LPG	CO ₂	184	91	2	5	0.010	0.0092
Stationary Combustion, Refinery gas	CO ₂	816	906	1	2	0.040	0.087
Stationary Combustion, Natural gas	CO ₂	4335	8293	1	0.4	0.182	0.176
Stationary Combustion, SOLID	CH ₄	13	3	1	100	0.006	0.0090
Stationary Combustion, LIQUID	CH ₄	3	1	1	100	0.002	0.0032
Stationary Combustion, GAS	CH ₄	3	6	1	100	0.011	0.0168
Natural gas fuelled engines, GAS	CH ₄	5	120	1	2	0.005	0.0051
Stationary Combustion, WASTE	CH ₄	1	2	5	100	0.003	0.0044
Stationary Combustion, BIOMASS	CH ₄	102	135	16	100	0.269	0.392
Biogas fuelled engines, BIOMASS	CH ₄	1	31	4	10	0.007	0.0066
Stationary Combustion, SOLID	N ₂ O	68	27	1	400	0.216	0.307
Stationary Combustion, LIQUID	N ₂ O	44	11	1	1000	0.218	0.291
Stationary Combustion, GAS	N ₂ O	16	24	1	750	0.361	0.476
Stationary Combustion, WASTE	N ₂ O	7	16	5	400	0.124	0.170
Stationary Combustion, BIOMASS	N ₂ O	38	91	2	1000	1.789	1.025
Transport, Road transport	CO ₂	9284	11 224	2	5	1.190	1.173
Transport, Military	CO ₂	119	116	2	5	0.012	0.0120
Transport, Railways	CO ₂	297	249	2	5	0.026	0.0257
Transport, Navigation (small boats)	CO ₂	48	99	41	5	0.080	0.0947
Transport, Navigation (large vessels)	CO ₂	748	399	11	5	0.095	0.127
Transport, Fisheries	CO ₂	591	479	2	5	0.051	0.0493
Transport, Agriculture	CO ₂	1272	1343	24	5	0.648	0.719
Transport, Forestry	CO ₂	36	17	30	5	0.010	0.0113
Transport, Industry (mobile)	CO ₂	839	1021	41	5	0.830	0.955
Transport, Residential	CO ₂	39	62	35	5	0.043	0.0494
Transport, Commercial/institutional	CO ₂	74	171	35	5	0.119	0.134
Transport, Civil aviation	CO ₂	243	133	10	5	0.029	0.0289
Transport, Road transport	CH ₄	47	11	2	40	0.009	0.0103
Transport, Military	CH ₄	0.1	0.1	2	100	0.0001	0.0002
Transport, Railways	CH ₄	0.3	0.2	2	100	0.0003	0.0005
Transport, Navigation (small boats)	CH ₄	0.3	1	41	100	0.001	0.0017
Transport, Navigation (large vessels)	CH ₄	0.3	0.2	11	100	0.0004	0.0004
Transport, Fisheries	CH ₄	0.3	0.2	2	100	0.000	0.0007
Transport, Agriculture	CH ₄	2	2	24	100	0.004	0.0060
Transport, Forestry	CH ₄	0.4	0.05	30	100	0.0001	0.0001
Transport, Industry (mobile)	CH ₄	1	1	41	100	0.002	0.0024
Transport, Residential	CH ₄	1	1	35	100	0.003	0.0044
Transport, Commercial/institutional	CH ₄	2	3	35	100	0.007	0.0101
Transport, Civil aviation	CH ₄	0.1	0.05	10	100	0.0001	0.0001
Transport, Road transport	N ₂ O	91	116	2	50	0.115	0.140

IPCC Source category	Gas	Base year emission Gg CO ₂ eqv.	2012 emission Gg CO ₂ eqv.	Activity data uncer- tainty %	Emission factor uncertain- ty %	Tier 1 Combined uncertainty % of total emissions	Tier 2 uncertainty % of total emissions
Transport, Military	N ₂ O	1	1	2	1000	0.025	0.0396
Transport, Railways	N ₂ O	3	2	2	1000	0.042	0.0606
Transport, Navigation (small boats)	N ₂ O	0.4	1	41	1000	0.021	0.0315
Transport, Navigation (large vessels)	N ₂ O	15	8	11	1000	0.153	0.171
Transport, Fisheries	N ₂ O	11	9	2	1000	0.185	0.244
Transport, Agriculture	N ₂ O	15	18	24	1000	0.349	0.511
Transport, Forestry	N ₂ O	0.2	0.2	30	1000	0.003	0.0049
Transport, Industry (mobile)	N ₂ O	11	13	41	1000	0.265	0.368
Transport, Residential	N ₂ O	0.2	0.3	35	1000	0.007	0.0100
Transport, Commercial/institutional	N ₂ O	0.3	1	35	1000	0.016	0.0215
Transport, Civil aviation	N ₂ O	3	2	10	1000	0.041	0.0591
1.B.2 Flaring in refinery	CO ₂	23	22	11	2	0.005	0.0049
1.B.2 Flaring off-shore	CO ₂	302	195	8	2	0.030	0.0300
1.B.2 Land based activities	CO ₂	0.003	0.01	2	40	4.2E-06	0.000005
1.B.2 Off-shore activities	CO ₂	2	4	2	30	0.002	0.002
1.B.2 Transmission of natural gas	CO ₂	0.003	0.0005	15	2	1.4E-07	0.0000004
1.B.2 Distribution of natural gas	CO ₂	0.005	0.005	25	10	2.5E-06	0.000003
1.B.2 Venting in gas storage	CO ₂	0.001	0.001	15	2	3.7E-07	0.0000001
1.B.2. Flaring in refinery	CH ₄	0.2	0.2	11	15	0.00006	0.00006
1.B.2. Flaring off-shore	CH ₄	1	0.3	8	125	0.001	0.0016
1.B.2 Refinery processes	CH ₄	1	47	1	125	0.115	0.186
1.B.2 Land based activities	CH ₄	17	18	2	40	0.014	0.0156
1.B.2 Off-shore activities	CH ₄	15	37	2	30	0.022	0.0241
1.B.2 Transmission of natural gas	CH ₄	4	0.3	15	2	0.000	0.0001
1.B.2 Distribution of natural gas	CH ₄	5	3	25	10	0.002	0.0017
1.B.2 Venting in gas storage	CH ₄	1	1	15	2	0.0004	0.0004
1.B.2 Flaring in refinery	N ₂ O	0.1	0.1	11	1000	0.001	0.0016
1.B.2 Flaring off-shore	N ₂ O	1	0.5	8	1000	0.009	0.0122
2A1 Cement production	CO ₂	882	871	1	2	0.038	0.0375
2A2 Lime production	CO ₂	116	40	5	5	0.006	0.0056
2A3 Limestone and dolomite use	CO ₂	14	26	5	5	0.004	0.0036
2A5 Asphalt roofing	CO ₂	0.02	0.02	5	25	0.00001	0.0000
2A6 Road paving with asphalt	CO ₂	2	2	5	25	0.001	0.0010
2A7a Glass and Glass wool	CO ₂	17	10	5	2	0.001	0.0010
2A7b Yellow bricks	CO ₂	23	17	5	2	0.002	0.0018
2A7c Expanded clay	CO ₂	15	6	5	2	0.001	0.0006
2B5 Catalysts/Fertilizers, Pesticides and Sulphuric acid	CO ₂	1	1	5	5	0.0002	0.0002
2C1 Iron and steel production	CO ₂	28	0	5	5	0.0E+00	0
2D2 Food and Drink	CO ₂	4	2	5	5	0.0003	0.0003
2G Lubricants	CO ₂	50	32	2	5	0.003	0.0032
2B2 Nitric acid production	N ₂ O	1043	0	2	25	0.000	0
2F Consumption of HFC	HFC	218	657	10	50	0.660	0.558
2F Consumption of PFC	PFC	1	9	10	50	0.009	0.0073
2F Consumption of SF ₆	SF ₆	107	118	10	50	0.118	0.105
3A Paint application	CO ₂	13	7	10	15	0.003	0.0027
3B Degreasing and dry cleaning	CO ₂	0.00004	0.000001	10	15	5.2E-10	0
3C Chemical products, manufacturing and processing	CO ₂	19	12	10	15	0.004	0.0045
3D5 Other	CO ₂	61	44	10	20	0.019	0.0207
3D5 Consumption of fireworks	CO ₂	0.1	0.2	8	100	0.0003	0.0004
3D5 Use of candles	CO ₂	22	81	10	20	0.036	0.0378

IPCC Source category	Gas	Base year emission Gg CO ₂ eqv.	2012 emission Gg CO ₂ eqv.	Activity data uncer- tainty %	Emission factor uncertain- ty %	Tier 1 Combined uncertainty % of total emissions	Tier 2 uncertainty % of total emissions
3D1 Other - Use of N ₂ O for Anaesthesia	N ₂ O	0	9	5	5	0.001	0.0013
3D5 Use of tobacco	N ₂ O	0.3	0.2	20	30	0.0001	0.0001
3D5 Use of charcoal for BBQ	N ₂ O	0.1	0.1	10	100	0.0003	0.0004
3D5 Consumption of fireworks	N ₂ O	1	2	8	100	0.004	0.0063
3D5 Use of candles	N ₂ O	0.1	0.2	10	20	0.0001	0.0001
4A Enteric Fermentation	CH ₄	3247	2904	2	20	1.149	1.192
4B Manure Management	CH ₄	985	1297	5	20	0.526	0.560
4F Field burning of agricultural residues	CH ₄	2	2	25	50	0.003	0.0032
4.B Manure Management	N ₂ O	600	391	22	100	0.788	0.525
4.D1.1 Synthetic Fertilizer	N ₂ O	2354	1103	25	100	2.240	3.428
4.D1.2 Animal waste applied to soils	N ₂ O	1112	1161	30	100	2.387	3.650
4.D1.3 N-fixing crops	N ₂ O	269	256	20	100	0.514	0.750
4.D1.4 Crop Residue	N ₂ O	361	311	20	100	0.625	0.956
4.D1.5 Cultivation of histosols	N ₂ O	290	198	20	100	0.398	0.599
4.D.2 Grassing animals	N ₂ O	334	211	25	100	0.428	0.648
4.D3 Atmospheric deposition	N ₂ O	496	295	19	100	0.590	0.885
4.D3 Leaching	N ₂ O	2447	1430	20	100	2.870	4.289
4.D1.6 Sewage sludge and Industrial waste used as fertiliser	N ₂ O	28	39	20	100	0.078	0.114
4.F Field Burning of Agricultural Residues	N ₂ O	1	1	25	50	0.001	0.0012
5.A.1 Forest remaining forest	CO ₂	50	-4491	15	2	-1.338	
5.A.2 Land converted to forest	CO ₂	77	38	15	9	0.013	
5(II) Forest Land.	N ₂ O	16	12	30	10	0.008	
5.B Cropland, Living biomass	CO ₂	-74	193	10	50	0.194	
5.B Cropland, Dead organic matter	CO ₂	3	12	10	50	0.012	
5.B Cropland, Mineral soils	CO ₂	1415	577	10	75	0.860	
5.B Cropland, Organic soils	CO ₂	2887	1981	10	90	3.531	
5(III) Disturbance, Land converted to cropland	N ₂ O	0.3	1	50	75	0.0018	
5.C Grassland, Living biomass	CO ₂	75	463	10	50	0.465	
5.C Grassland, Dead organic matter	CO ₂	2	7	10	50	0.007	
5.C Grassland, Mineral soils	CO ₂	0.2	5	10	75	0.007	
5.C Grassland, Organic soils	CO ₂	107	79	10	90	0.140	
5.D Wetlands, Living biomass	CO ₂	3	-0.2	10	50	0.000	
5.D Wetlands, Dead organic matter	CO ₂	1	0	10	100	0.000	
5.D Wetlands, Soils	CO ₂	85	2	10	100	0.005	
5(II) Wetlands	N ₂ O	0.1	0.1	10	100	0.000	
5.E Settlements, Living biomass	CO ₂	11	56	10	50	0.056	
5.E Settlements, Dead organic matter	CO ₂	1	0.03	10	50	0.000	
5.E Settlements, Soils	CO ₂	1	35	10	50	0.035	
5(IV) Cropland Limestone	CO ₂	623	192	5	50	0.190	
5(V) Biomass Burning	CH ₄	1	0.02	50	30	0.000	
5(V) Biomass Burning	N ₂ O	0.4	0.03	50	30	0.00004	
6 A. Solid Waste Disposal on Land	CH ₄	1366	698	10	118	1.626	2.536
6 B. Wastewater Handling	CH ₄	65	74	24	32	0.058	0.0743
6 B. Wastewater Handling - Direct	N ₂ O	23	41	22	50	0.043	0.0543
6 B. Wastewater Handling - Indirect	N ₂ O	82	32	59	42	0.046	0.0624
6.D Accidental fires, buildings	CO ₂	11	11	10	300	0.064	0.151
6.D Accidental fires, vehicles	CO ₂	6	6	10	500	0.055	0.151
6.C Incineration of corpses	CH ₄	0.01	0.01	1	150	0.00003	0.0001
6.C Incineration of carcasses	CH ₄	0.001	0.005	40	150	0.00001	0
6.D Compost production	CH ₄	29	90	40	100	0.192	0.293

IPCC Source category	Gas	Base year emission Gg CO ₂ eqv.	2012 emission Gg CO ₂ eqv.	Activity data uncer- tainty %	Emission factor uncertain- ty %	Tier 1 Combined uncertainty % of total emissions	Tier 2 uncertainty % of total emissions
6.D Accidental fires, buildings	CH ₄	1	1	10	500	0.013	0.0354
6.D Accidental fires, vehicles	CH ₄	0.3	0.2	10	700	0.003	0.0105
6.C Incineration of corpses	N ₂ O	0.2	0.2	1	150	0.001	0.0010
6.C Incineration of carcasses	N ₂ O	0.0	0.1	40	150	0.0003	0.0005
6.D Compost production	N ₂ O	13	127	40	100	0.269	0.413

1.7.2 Results of the tier 1 uncertainty estimation

The estimated uncertainties for total GHG and for CO₂, CH₄, N₂O and F-gases are shown in Table 1.4. The base year for F-gases is 1995 and for all other sources the base year is 1990. The total Danish net GHG emission is estimated with an uncertainty of ± 6.9 % and the trend in net GHG emission since the base year has been estimated to be -31.6 % ± 2.6 %-age points. The GHG uncertainty estimates do not take into account the uncertainty of the GWP factors.

The uncertainty on N₂O emission from stationary biomass combustion, CH₄ emission from solid waste disposal, N₂O emission from leaching and run-off and N₂O emission from animal waste applied to soil and synthetic fertiliser are the largest sources of uncertainty for the Danish GHG inventory (excluding LULUCF). For LULUCF the largest sources of uncertainty are forest land remaining forest land and soil emissions from cropland.

The uncertainty of the GHG emission from combustion (sector 1A) is 2.2 % and the trend uncertainty is -25.3 % ± 2.2 %-age points.

Table 1.4 Uncertainties 1990-2012.

	Uncertainty [%]	Trend [%]	Uncertainty in trend [%-age points]
GHG	6.9	-31.6	± 2.6
CO ₂	5.6	-33.7	± 2.4
CH ₄	19	-7.3	± 11.8
N ₂ O	42	-39.1	± 12.6
F-gases	43	141	± 41
CO ₂ excl. LULUCF	2.1	-25.5	± 1.7
GHG excl. LULUCF	5.6	-25.1	± 2.5

1.7.3 Tier 2 uncertainties

On the recommendation of the UNFCCC expert review team (ERT) in 2009 Denmark has undertaken a tier 2 uncertainty analysis. Please see the sectoral chapters for the sectoral results of the tier 2 uncertainty analysis. Below is a description on the theoretical basis for the tier 2 uncertainty calculations. For the overall result please refer to Chapter 1.7.4.

When to use Tier 2

When the activity data and emission factors cannot fulfil the criteria for using the error propagation equations in Tier 1 an alternative stochastic simulation, i.e. Monte Carlo method, can be employed. The Monte Carlo method constitutes Tier 2 and Approach 2 in IPCC (2000 and 2006) and is suitable for

estimating uncertainty in emission rates, from uncertainties in activity data and emission factors, when:

- Uncertainties are large.
- Their distribution is non-normal.
- The algorithms are complex function and not only simple multiplication of activity data with emission factors.
- Correlations occur between some of the activity data sets, emission factors, or both.

Uncertainties found in inventory source categories can vary widely from a few per cent to orders of magnitude. When using a normal distribution for a parameter with large uncertainty there is a risk of having a certain probability for negative values, which is not possible in reality. Furthermore large uncertainty gives a certain probability of having extremely large values, i.e. values orders of magnitude larger than the mean value. Extreme values are an often occurring quality for the distribution of realistic activity data and emission factors. However, in some cases the extreme values are unrealistic and here the method allows for upper and lower truncation of input parameters. This implies applying a lower and/or upper boundary for the distribution function of input parameters. A logarithmic plot of data with large uncertainties will transform a skewed distribution probability function (a) into a bell-shaped log-normal distribution function (b), cf. Figure 1.5. The latter can be defined by a mean value, α , and standard deviation, σ , respectively. The log-normal distribution is selected as standard in the first version of the Tier 2 and Approach 2 uncertainty assessment for year 2009. A further feature of applying truncation boundaries is that a probability distribution will converge towards a box distribution when narrowing the truncation interval.




Figure 1.5 Log-normal distribution (\log_{10}), both on original (a) and log scale (b). The median (α') is 100 and the multiple standard deviation (σ') is 2. The resulting median (equal mean) and the standard deviation in the \log_{10} distribution is respectively $\alpha = \log_{10}(100) = 2$ and $\sigma = \log_{10}(2) = 0.301$ (Limbert et al., 2001).

In case the uncertainty is much smaller than the mean value, then the normal and log-normal distributions will not differ much, cf. Figure 1.6, where the relationship between normal and log-normal distributions are illustrated (Limbert et al., 2001).




Figure 1.6 Comparison between the normal distribution (green area, median 100, standard deviation 20) the different degrees of variability (described by σ^*) for log-normal distributions that all have the same median value, i.e. α^* on original scale, of 100 (Limbert et al., 2001).

The difference in shape between a normal and log-normal distribution is seen in Figure 1.6 for different values of σ^* . The standard deviation for the normal distribution is 20 and thus equal to 20 % of the mean value and the log-normal distribution having a σ^* value of 1.2 reflects the same level of “deviation” as in the normal distribution. So, the discrepancy between the green area and the curve for $\sigma^*=1.2$ illustrates the difference in interpretation of a 20 % deviation as measured by respectively the normal and log-normal distribution. This discrepancy is so limited that it is overruled by the vagueness related to empirical quantification of the uncertainty level based on expert knowledge and data and the fact that any assumed distribution function is an approximation. Therefore, by using log-normal distributions as standard description of all uncertainty input it will in reality include normal distributions when the magnitude of uncertainty is limited to a minor fraction of the mean value.

A way of calculating the intervals of confidence, expressed by the median (α^*) and standard deviation (σ^*), for a log-normal distribution on original scale, cf. Figure 1a, is presented in Limbert et al. (2001). For normally distributed data, the interval [median \pm standard deviation] covers a probability of 68.3 %, while [median \pm 2*standard deviation] covers 95.5 %. Correspondingly for log normal data on original scale, cf. Figure 1a, the interval [α^* / σ^* , $\alpha^* * \sigma^*$] covers 68.3 % and the interval [$\alpha^* / (\sigma^*)^2$, $\alpha^* * (\sigma^*)^2$] covers 95.5 %.

Often the default uncertainty values in IPCC (2000) e.g. for emission factors, are expressed as a percentage, e.g. 30 %. When this represents a standard deviation (68.3 %) on original scale we will proceed using $\sigma^* = 1.3$ in the uncertainty analysis. When it represents a 95 % interval of confidence, we will use $\sigma^* = (1.3)^{0.5} = 1.14$ in the uncertainty analysis. When the 95 % interval of confidence on original scale is below approximately 300 % the standard deviation for a log-normal distribution on original scale, can be approximated by dividing with a factor of 2, i.e. $0.3/2 = 0.15$, and thus $\sigma^* = 1.15$.

Procedure of Tier 2 (Monte Carlo method)

The procedure of the Tier 2 (MC) analysis consists of four steps where only Step 1 requires effort from the user:

- Step 1: Estimation of activity data and emission factors, their associated mean values, uncertainties such as standard deviation, probability density functions and any correlations.
- Step 2: Selection of random values of activity data and emission factors.
- Step 3: Calculate emissions from selected random values.
- Step 4: The calculated result in step 3 is stored and the process is repeated from step 2.

Repetition of steps 2 and 3 are continued until the calculated mean value and error intervals are sufficiently determined (typically 10,000 times). Each single repetition is denoted a “single sample” in the following and one execution of steps 2 and 3 is denoted a “MC sample”.

The software is developed in excel VBA programming by a scientist associated with the sector experts, which enables a transparent and accurate transfer and interpretation of emission factors and activity data (input) and calculated emissions with uncertainties (output).

Different criteria and guidelines for estimation of value uncertainty for activity data and emission factors are outlined in the next section. Whether they are based on information from models, empirical data or expert judgement, they form lines of evidence towards the most appropriate estimate. The basic paradigm for a MC analysis is outlined in Figure 1.7.




Figure 1.7 Methodological principle in compiling and quantifying input data for input parameters, e.g. emission factors, which are to be used in Tier 2 (MC) uncertainty analysis. Each evidence is formed from assessment of information from models, empirical data or expert judgement. The upper dotted box represents step 1 in the MC analysis, which is performed for each input parameter. The lower dotted box represents steps 2 to 4, and is performed in the emission modelling with all input parameters.

The principle of the MC method is to generate many “possible” calculations and thus map the resulting “possible” results. The possible calculations are made based on the “realistic” variability (uncertainty) related to the input

parameter values. This variability needs to be described as a distribution function. The MC method is considered in two parts: (1) A distribution estimation part, where the variabilities of the input parameters are parameterised; (2) A technical part that makes the simulation based on the estimated distributions. The first part is highly critical and requires high attention. The second part is a question of programming and therefore mostly a technical issue. The MC method is a model for how uncertainty of input parameters influences the calculation results, so the MC also involves uncertainty in the prediction of uncertainty. It is therefore important to predict the variability of the input parameters as correctly as possible. The MC method does not include the validity of the calculations as estimators of reality but only the uncertainty of the input parameter values. Consequently, there are many fundamental types of uncertainty that are not included in the MC method.

The method is based on single samples, where the mean is unity and where the variability is determined by the uncertainty of the parameter as discussed above, see Figure 1.8. This sampled value is subsequently multiplied with the best estimate of the parameter value to yield a sampled value for this parameter. The reason for this two stage sampling is that it makes it possible directly to include correlation in uncertainty between years as explained below.




Figure 1.8 The principle in a single MC draw of the value x_d , where the median (α) is unity and where the standard deviation (σ) determines the variation around 1.

Correlation in the uncertainty may occur between years, e.g. when the same sources are responsible for uncertainties in several years. This takes place because many sources of uncertainty are dependent between years, so if a parameter is over-estimated for one year then this parameter may also tend to be overestimated other years. This implies that when the uncertainty is high one year the uncertainty will also be high the other year(s). The principle of performing a MC analysis with an emission factor and activity data that have uncertainties that are correlated between one or more years is illustrated in Figure 1.9.

The principle in Figure 1.9 is to sample a value (x) as shown in Figure 4, where the median value is unity and subsequently multiply the sampled value with the estimated median value (e.g. $AD_{s1} = AD_1 x$). This two-step approach makes it possible to include correlating uncertainty between different years. If two years are correlated then a deviation from the estimated mean value is assumed to be the same in relative terms for the two years. By

sampling, using the median of unity once, and subsequently use this value to estimate the value for the two years, using the two medians for each year, this will yield the correlation between the two years as a simple consequence and thereby be directly simulated in the MC sampling.

The MC sampling is illustrated in Figure 1.9 for a single source, where s is the sampling number index, counting up to e.g. 10,000. In Figure 5 there will be a strong correlation between year 2 and 3, because both the uncertainty of EF and AD is correlated, for year 1 there will be a partial correlation with respectively year 2 and 3 because the uncertainty of the EF value is correlated, but the uncertainty is independent for AD . Year 4 is completely independent of the other years. The figure is only illustrating a single source and typically the emission estimates includes several sources each having some more or less correlated uncertainty. The final emission estimates are thus more or less correlated between years in a highly complex way.

Performing MC analysis for correlated parameters corresponds to the calculation scheme for MC analysis of emissions and the trend of a category as shown in Appendix A (IPCC, 2006) (Figure 3.7 pp. 3.36). The scheme shows calculations for correlated and non-correlated parameters.




Figure 1.9 The principle of a MC sample for draws of random numbers and generation of any emission factor and activity data for a four year period. The upper half illustrates the sampling of any emission factor for year 1 to year 4. The uncertainty associated to the emission factor is correlated for year 1, 2 and 3 and therefore the same random number is used for generating EF1, EF2 and EF3. The lower half illustrates the sampling of activity data. The uncertainty associated to the activity data is correlated for year 2 and 3 and therefore the same random number is used for generating AD2 and AD3. In the middle row the emission factor and activity data are multiplied for each year.

In some cases there exists additional a priori information about categories of activity data, where the total sum is known with high certainty, but where the sub categories are more uncertain. In this case the single samples within one year are adjusted so all sub sources together adds up to the correct total number and the single sampling in this case will describe the uncertainty between the single categories.

MC analyses for emissions

When a 95% confidence interval has been entered as percentages of median values of the input parameters, i.e. emission factors and activity data, for source categories and sub-categories, the above MC procedure is executed 10,000 times. The output of the MC analysis is reported as in Table 1.5 where the median emissions are shown together with the 95% confidence interval (2.5% - 97.5%).

Two basic questions are important to answer: (1) What is the uncertainty for a time trend estimate; (2) What is the uncertainty within the same year of the single sub-categories, source categories and the total estimate. The first question takes correlation of uncertainty between years into account and the second question considers one year at a time and correlation between years is not relevant.

In the ideal case it will be possible to answer the two questions based on the same MC samples, where every single sample is stored for every source and for every year. However, this is not possible in the VBA programming due to limitations in variable table on a normal pc. Thus two MC samplings take place: (1) The total emission is calculated for every year and every MC sample, so for 10,000 MC samples and 20 years, this needs storage of 200,000 numbers; (2) Every year is analysed separately where only results for one year is stored at a time, so for 10,000 MC samples and 50 sources this yields 500,000 numbers to be stored. Using this two-stage approach it is easily possible to run the MC analysis in Excel. Consequently, the exact value for the median analysed for a specific year (question 2 above) is not similar with the medians in the time trend analysis (question 1 above) due to a finite number of MC samples, but this is not a real problem. If this discrepancy is considered as critical then it simply tells that the number of MC samples should be increased and that the analysis thus has to be redone.

Table 1.5 Example of output scheme for tier 2 MC uncertainty analysis. Median emissions and 95 % confidence intervals are calculated for total emission, emissions for source categories and emissions for sub-categories. Calculated 95% confidence intervals are furthermore calculated for activity data and emission factors.

Source category	Sub-categories	Activity			EF			Emissions			
		< 2.5%	>97.5%	Interval	< 2.5%	> 97.5%	Interval	Median	< 2.5%	> 97.5%	Interval
all	all	-	-	-	-	-	-				
A	all	-	-	-	-	-	-				
B	all	-	-	-	-	-	-				
C	all	-	-	-	-	-	-				
A	1										
A	2										
A	3										
B	1										
B	2										
C	1										
C	2										
C	3										
C	4										

Results for each row can also be reported as:

Median emission [- (median - <2.5%)/median/100%, + (>97.5% - median)/median/100%]

MC trend analysis

The trend analysis is performed by comparing emissions from two individual years at a time. The probability for Year 1 (base year) to be above Year 2 (latest year) is calculated using the equation:

$$P_{Year1>Year2} = \frac{N_{year1>year2}}{N_{total}},$$

where $N_{year1>year2}$ is the number of MC samples where year 1 is estimated to have higher emission compared to year 2 and N_{total} is the total number of MC runs. In case of $P_{year1>year2} \approx 1$ it is strongly significant to conclude that year 1 has higher emission than year 2, and reverse for $P_{year1>year2} \approx 0$. This is a comparison between years in pairs that can be filled in to a matrix, where all years are compared with all other years.

Table 1.6 Comparison of emissions between years in trend analysis.

	Year 1	Year 2	Year 3	Year 4
Year 1	0			
Year 2		0		
Year 3			0	
Year 4				0

Results for trend analysis of emissions between two years, year 1 and year 2, can be reported as median difference, <2.5% and >97.5%, or as:

Median difference [- (median difference - <2.5%)/median difference/100%,
+ (>97.5% - median difference)/median difference/100%]

Quantifying uncertainties in Tier 2

In order to perform the four steps of a Tier 2 (MC) uncertainty analysis as described in the previous paragraph the user has to gather the information stated in step 1. It is essential to establish the best possible estimate, and the following guide sets up a procedure for assessing, quantifying and compiling uncertainties for the parameters that are entered in the emission models. The guide is based on IPCC guidelines (IPCC, 2000 & 2006) and NUSAP and expert elicitation in van der Sluijs et al. (2004).

The uncertainty of a parameter, e.g. activity data and emission factor, is considered to be proportional to the associated parameter. This means that the uncertainty is expressed as a percentage of the parameter value. The median value is used and the uncertainties represent the parameter standard deviation, σ^* . We assume log-normal distributions, which equals normal distributions at low uncertainty values. Although van der Sluijs et al. (2004) suggest different probability distribution functions depending on the level of knowledge on input parameters we will use log-normal distributions for all parameters, as argued in the previous section.

The methodology offers a possibility for correlating the uncertainties of two or more parameters. When uncertainties of two or more parameters are assumed to be correlated they will be attributed the same random number in any MC sample, as explained in the previous paragraph.

Uncertainties will be reported according to the IPCC General Reporting Table for Uncertainty. Uncertainties will be reported for:

- Total uncertainty of the entire sector
- Key source categories
- Aggregated CRF levels
- Most differentiated CRF category levels that are entered by the user

IPCC guideline - Sources of data

Quantifying uncertainties is dependent on the source of data, and in general there are three broad sources of data and information (IPCC, 2000 & 2006):

Information contained in models

A model is a representation of the real world and does therefore not exactly mimic real-world systems. The structure of a model is often thought of in terms of the equations used. The key considerations in model uncertainty are; has the correct, most relevant real-world system been identified and are the model equations accurate representations of the chosen system. Typically the model equations are the product of activity data and emission factors, cf. Eq 1, but there may also be more complex model equations for emissions and also for derivation of activity data and emission factors.

In some cases, model uncertainty can be significant. It is typically poorly characterised and may not be characterised at all. The inventory expert must consider the parameters that are used and assess if there are model assumptions that are imprecise or inaccurate. For the most critical models an effort can be made to evaluate and quantify the size of the potential error that occurs from using the model. There are at least three approaches for estimating

the model uncertainty: 1) comparison of a model result with independent data, 2) comparison of a model result with the result of alternative models, and 3) expert judgement regarding the magnitude of the model uncertainty. These approaches can be used in combination.

Empirical data for sources and sinks and activity

This implies empirical data associated with measurements of emissions, emission factors and activity data from surveys and censuses. When estimating uncertainty from measured emissions data, considerations include; representativeness of the data and potential for bias, precision and accuracy of the measurements, sample size and inter-individual variability in measurements and their implications for uncertainty in mean annual emissions, inter-annual variability in emissions and whether estimates are based on an average of several years or on the basis of a particular year.

Quantification of uncertainties and defining the probability distribution function (PDF) for empirical data can be summarised as follows: 1) Compilation of activity data, emission factors and other parameters. These data typically represent variability, 2) Visualisation of data by plotting empirical distribution functions for each parameter; horizontally according to numerical value or interval and vertically by frequency, 3) Fitting, evaluation and selection of PDFs for representing variability of data, 4) Characterisation of mean value and of uncertainty in the mean of the distributions for variability. If the standard error of the mean is small, a normality assumption can be made regardless of the sample size or skewness of data. If the standard error of the mean is large, then typically a log-normality assumption can be made, 5) Once mean values, uncertainties and standard errors have been specified, these can be used as input to Tier 2 MC analysis for estimating uncertainties in total emissions, 6) Sensitivity analysis can be used to determine which parameters induce highest uncertainties in the total uncertainty, and prioritise efforts to develop good estimates of these key uncertainties.

Expert judgement as a source of information

In many situations, relevant empirical data are not available for activity data, emission factors etc. to an inventory. In such situations, a practical solution is to obtain well informed judgements from domain experts regarding best estimates and uncertainties of input data.

Commonly used methods for converting an expert's judgement regarding uncertainty into a quantitative PDF are: 1) Fixed value; Estimate the probability of being higher (or lower) than an arbitrary value and repeat, three or five times. For example, what is the probability that an emission factor would be less than 100? 2) Fixed probability; Estimate the value associated with a specified probability of being higher (or lower). For example, what is the emission factor such that there is only a 2.5% probability that the emission factor could be lower (or higher) than that value, 3) Interval methods; For example, choose a value of the emission factor such that it is equally likely that the true emission factor would be higher or lower than that value. This yields the median. Then divide the lower range into two bins such that there is assumed to be equally likely (25% probability) that the emission factor could be in either bin. Repeat this for the other end of the distribution. Finally, either fixed probability or fixed value methods could be used to get judgements for extreme values, 4) Graphing; the expert draws a distribution. This should be used cautiously because some experts are overconfident about their knowledge of PDFs.

Sometimes the only available expert judgement consists of a range, maybe quoted together with a most likely value. Under these circumstances the following rules are considered good practice: Where experts only provide an upper and a lower value, assume that the PDF is uniform and that the range corresponds to the 95 per cent confidence interval. Where experts also provide a most likely value (point estimate), assume a triangular PDF using the most likely values as the mode and assume that the upper and lower values each exclude 2.5% of the population. The distribution needs not to be symmetrical. Normal or log-normal distributions can be used given appropriate justifications.

Concluding remarks and planned improvements

Tier 2 uncertainties are typically found to be greater than Tier 1 uncertainties. When large input uncertainties, e.g. > 10%, are used, the deviation becomes pronounced. For smaller input uncertainties, e.g. < 1%, Tier 1 approximates Tier 2 calculations.

The Log-normal distribution was selected due the likely conditions for the distribution as being close to a normal distribution for smaller uncertainties on one hand and close to the understanding of larger uncertainties on the other hand. However, in case of larger uncertainty the outcome of the MC analysis includes rather extreme values that in some cases are unrealistic. The method therefore allows for truncation of input uncertainties, either a lower boundary, upper boundary or both, depending of which truncation are most realistic.

1.7.4 Results of the tier 2 uncertainty estimation

Tier 2 uncertainty results for sectors and categories are shown in Table 1.3. The input uncertainties for activity data and emission factors stated in Table 1.3 are used both in Tier 1 and Tier 2 uncertainty calculations. The total Danish net GHG emission for 2012 is estimated with an uncertainty of +6.7 % and -4.5 % and the trend in net GHG emission since 1990 is estimated with an uncertainty of +9.2 and -7.4 %-age points.

Tier 2 uncertainties are typically larger than Tier 1 uncertainties when input uncertainties are larger than approximately 25%, which corresponds to the model domain of Tier 1 method. This implies that the Tier 2 method is more reliable for large input uncertainties.

1.8 General assessment of the completeness

The present Danish greenhouse gas emission inventory includes all major sources identified by the Revised 1996 IPPC Guidelines. Please see Annex 5 for detailed discussion on minor sources that are not included.

1.9 References

DEPA, 2014: The greenhouse gases HFCs, PFCs and SF₆. Danish consumption and emissions 2012. Tomas Sander Poulsen & Mads Werge, PlanMiljø. In press.

DEPA, 2013: Waste Statistics 2011. Affaldsstatistik 2011. Danish Environmental Protection Agency. (In Danish). Available: http://www.mst.dk/Virksomhed_og_myndighed/Affald/Tal_for_affald/Statistikker_og_ISAG-dataudtraek/ (21-02-2014).

EMEP/CORINAIR, 2007: EMEP/CORINAIR Atmospheric Emission Inventory Guidebook – 2007, prepared by the UNECE/EMEP Task Force on Emissions Inventories and Projections. Technical Report No 16/2007. Available: <http://www.eea.europa.eu/publications/EMEP-CORINAIR5> (21-02-2014).

EMEP/EEA, 2009: EMEP/EEA air pollutant emission inventory guidebook 2009. Technical guidance to prepare national emission inventories. EEA Technical Report 9/2009. Available: <http://www.eea.europa.eu/publications/emep-eea-emission-inventory-guidebook-2009> (21-02-2014).

EU Commission, 2004: COMMISSION DECISION of 29 January 2004 establishing guidelines for the monitoring and reporting of greenhouse gas emissions pursuant to Directive 2003/87/EC of the European Parliament and of the Council. Available: <http://eurlex.europa.eu/LexUriServ/LexUriServ.do?uri=OJ:L:2004:059:0001:0074:EN:PDF> (21-02-2014).

EU Commission, 2007: COMMISSION DECISION of 18 July 2007 establishing guidelines for the monitoring and reporting of greenhouse gas emissions pursuant to Directive 2003/87/EC of the European Parliament and of the Council. Available: <http://eurlex.europa.eu/LexUriServ/LexUriServ.do?uri=OJ:L:2007:229:0001:0085:EN:PDF> (21-02-2014).

Fausser, P., Thomsen, M., Nielsen, O-K., Winther, M., Gyldenkærne, S., Hoffmann, L., Lyck, E. & Illerup, J.B. 2007: Verification of the Danish emission inventory data by national and international data comparisons. National Environmental Research Institute, University of Aarhus, Denmark. 53 pp. – NERI Technical Report no. 627. Available: http://www2.dmu.dk/pub/fr627_final.pdf

Fausser, P., Nielsen, M., Winther, M., Plejdrup, M., Gyldenkærne, S., Mikkelsen, M.H., Albrektsen, R., Hoffmann, L., Thomsen, M., Hjelgaard, K. & Nielsen, O.-K., 2013: Verification of the Danish 1990, 2000 and 2010 emission inventory data. Aarhus University, DCE – Danish Centre for Environment and Energy, 85 pp. Scientific Report from DCE – Danish Centre for Environment and Energy No. 79. <http://dce2.au.dk/pub/SR79.pdf>

Holtskog, S., Haakonsen, G., Kvingedal, E., Rypdal, K. & Tornsjø, B., 2000: Verification of the Norwegian emission inventory – Comparing emission intensity values with similar countries. The Norwegian Pollution Control Authority in cooperation with Statistics Norway. SFT report 1736/2000.

Illerup, J.B., Lyck, E., Winther, M. & Rasmussen, E. 2000: Denmark's National Inventory Report – Submitted under the United Nations Framework Convention on Climate Change. Emission Inventories. Research Notes from National Environmental Research Institute, Denmark no. 127, 326 pp. Available http://www.dmu.dk/1_viden/2_Publikationer/3_arbrapporter/rapporter/ar127.pdf

IPCC, 1996: Revised 1996 IPCC Guidelines for National Greenhouse Gas Inventories. Available: <http://www.ipcc-nggip.iges.or.jp/public/gl/invs6.htm> (21-02-2014).

- IPCC, 2000: Good Practice Guidance and Uncertainty Management in National Greenhouse Gas Inventories. Available: <http://www.ipcc-nggip.iges.or.jp/public/gp/english/> (21-02-2014).
- IPCC, 2003: Good Practice Guidance for Land Use, Land-Use Change and Forestry Available: <http://www.ipcc-nggip.iges.or.jp/public/gp/gluglucf/gpuglucf.html> (21-02-2014).
- IPCC, 2006: 2006 IPCC Guidelines for National Greenhouse Gas Inventories, Prepared by the National Greenhouse Gas Inventories Programme, Eggleston H.S., Buendia L., Miwa K., Ngara T. & Tanabe K. (eds). Published: IGES, Japan. Available at: <http://www.ipcc-nggip.iges.or.jp/public/2006gl/index.html> (21-02-2014)
- IPCC, 2007: Climate Change 2007. The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment report of the Intergovernmental Panel on Climate Change [Solomon, S., D. Qin, M. Manning, Z.Chen. M. Marquis, K.B. Averyt, M. Tignor & H.L. Miller (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA 996 pp.
- Limpert, E., Stahel, W.A., Abbt, M., (2001), Log-normal distributions across the Sciences: Keys abd Clues, BioScience, Vol. 51, No. 5.
- Mikkelsen, M.H. Albrektsen, R. & Gyldenkærne, S. 2011: Danish emission inventories for agriculture. Inventories 1985 - 2009. National Environmental Research Institute, Aarhus University. 136 pp. - NERI Technical Report No. 810. Available: <http://www2.dmu.dk/Pub/FR810.pdf>
- Nielsen, O.-K., Plejdrup, M.S., Winther, M., Gyldenkærne, S., Thomsen, M., Fauser, P., Nielsen, M. Mikkelsen, M.H., Albrektsen, R., Hjelgaard, K., Hoffmann, L. & Bruun, H.G., 2013: Quality manual for the Danish greenhouse gas inventory. Version 2. Aarhus University, DCE - Danish Centre for Environment and Energy, 44 pp. Scientific Report from DCE - Danish Centre for Environment and Energy No. 47. Available: <http://www.dmu.dk/Pub/SR47.pdf>
- Pulles, T., Mareckova, K., Svetlik, J., Linek, M., & Skakala, J. 1999: CollectER -Installation and User Guide, EEA Technical Report No 31.
- Statistics Denmark, 2013: Agriculture Statistics 2012. Denmark. Available at: <http://www.statistikbanken.dk/statbank5a/default.asp?w=1280> (21-02-2014).
- Sørensen, P.B., Illerup, J.B., Nielsen, M., Lyck, E., Bruun, H.G., Winther, M., Mikkelsen, M.H. & Gyldenkærne, S. 2005: Quality manual for the greenhouse gas inventory. Version 1. National Environmental Research Institute. - Research Notes from NERI 224: 25 pp. (electronic). Available: http://www2.dmu.dk/1_viden/2_Publikationer/3_arbrapporter/rapporter/AR224.pdf
- UNFCCC, 2007: Updated UNFCCC reporting guidelines on annual inventories following incorporation of the provisions of decision 14/CP.11 Available: <http://unfccc.int/resource/docs/2006/sbsta/eng/09.pdf> (21-02-2014).

Van der Sluijs, J.P., Janssen, P.H.M., Petersen, A.C., Kloprogge, P., Risbey, J.S., Tuinstra, W., Ravetz, J.R., 2004. RIVM/MNP Guidance for Uncertainty Assessment and Communication: Tool Catalogue for Uncertainty assessment. Netherlands Environmental Assessment Agency, RIVM. Report no. NMS-E-2004-37.

2 Trends in greenhouse gas emissions

The trends presented in this Chapter cover the emissions from Denmark. Due to the small emissions originating from Greenland the trends are very similar in fact close to identical. A trend discussion of the aggregated greenhouse gas emissions from Denmark and Greenland is included in Chapter 17.1.

2.1 Description and interpretation of emission trends for aggregated greenhouse gas emissions

2.1.1 Greenhouse Gas Emissions

The greenhouse gas emissions are estimated according to the IPCC guidelines and are aggregated into seven main sectors. The greenhouse gases include CO₂, CH₄, N₂O, HFCs, PFCs and SF₆. Figure 2.1 shows the estimated total greenhouse gas emissions in CO₂ equivalents from 1990 to 2012. The emissions are not corrected for electricity trade or temperature variations. CO₂ is the most important greenhouse gas contributing in 2012 to the national total in CO₂ equivalents excluding LULUCF (Land Use and Land Use Change and Forestry) with 76.3 % followed by N₂O with 11.5 %, CH₄ 10.6 % and F-gases (HFCs, PFCs and SF₆) with 1.5 %. Seen over the time-series from 1990 to 2012 these percentages have been increasing for CH₄ and F-gases, and decreasing for N₂O. The percentages for CO₂ show larger fluctuations during the time series. Stationary combustion plants, Transport and Agriculture represent the largest contributing categories to emissions of greenhouse gases, followed by Industrial processes, Waste, fugitive emissions and Solvents, see Figure 2.1. The net CO₂ uptake by LULUCF in 2012 is 1.6 % of the total emission in CO₂ equivalents excl. LULUCF. The national total greenhouse gas emission in CO₂ equivalents excluding LULUCF has decreased by 24.8 % from 1990 to 2012 and decreased 31.3 % including LULUCF. From 2011 to 2012 the total greenhouse gas emission excluding LULUCF decreased by 8.6 %. The decrease is mainly caused by decreasing emissions from the energy sector due to increasing import of electricity and increasing production of wind power. Comments on the overall trends etc. seen in Figure 2.1 are given in the sections below on the individual greenhouse gases.




Figure 2.1 Greenhouse gas emissions in CO₂ equivalents distributed on main sectors for 2012 (excluding LULUCF) and time series for 1990 to 2012 (including LULUCF).

2.2 Description and interpretation of emission trends by gas

2.2.1 Carbon dioxide

The largest source of the emission of CO₂ is the energy sector, which includes the combustion of fossil fuels such as oil, coal and natural gas (Figure 2.2). Energy Industries contribute with 42.0 % of the emissions (excl. LULUCF). About 31 % come from the transport sector. The CO₂ emission (excl. LULUCF) decreased by 10.5 % from 2011 to 2012. The main reason for this decrease owe to decreasing fuel consumption, mainly for coal and natural gas. The decrease partly owe to increasing import of electricity and increasing production of wind power. In 2012, the actual CO₂ emission (incl. LULUCF) was 33.7 % less than the emission in 1990.




Figure 2.2 CO₂ emissions. Distribution according to the main sectors (2012) and time series for 1990 to 2012.

2.2.2 Nitrous oxide

Agriculture is the most important N₂O emission source in 2012 contributing 90.7 % (Figure 2.3) of which N₂O from agricultural soils accounts for 92.8 %. N₂O is emitted as a result of microbial processes in the soil. Substantial emissions also come from drainage water and coastal waters where nitrogen is converted to N₂O through bacterial processes. However, the nitrogen converted in these processes originates mainly from the agricultural use of manure and nitrogen fertilisers. The main reason for the drop in the emissions of N₂O in the agricultural sector of 34.9 % from 1990 to 2012 is legislation to improve the utilisation of nitrogen in manure. The legislation has resulted in less nitrogen excreted per unit of livestock produced and a considerable reduction in the use of nitrogen fertilisers. The basis for the N₂O emission is then reduced. Combustion of fossil fuels in the energy sector, both stationary and mobile sources, contributes 5.8 %. The N₂O emission from transport contributed with 2.5 % in 2012. This emission has increased during the nineties because of the increase in the use of catalyst cars. Production of nitric acid stopped in 2004 and the emissions from industrial processes is therefore not occurring from 2005 onwards. The sector Solvent and Other Product Use covers N₂O from e.g. anaesthesia.




Figure 2.3 N₂O emissions. Distribution according to the main sectors (2012) and time series for 1990 to 2012.

2.2.3 Methane

The largest sources of anthropogenic CH₄ emissions are agricultural activities contributing in 2012 with 76.5 %, waste (15.8 %), public power and energy industries (2.5 %), see Figure 2.4. The emission from agriculture derives from enteric fermentation and management of animal manure contributing with 52.9 % and 23.6 % of the national CH₄ emission excl. LULUCF in 2012. The CH₄ emission from public power and district heating plants increased in the nineties, mainly 1992-1996, due to the increasing use of gas engines in the decentralised cogeneration plant sector. Up to 3 % of the natural gas in the gas engines is not combusted. The deregulation of the electricity market has made production of electricity in gas engines less favourable, therefore the fuel consumption has decreased and hence the CH₄ emission has decreased. Over the time series from 1990 to 2012, the emission of CH₄ from enteric fermentation has decreased 10.6 % due to the decrease in the number of cattle. However, the emission from manure management has in the same period increased 31.7 % due to a change in traditional stable systems towards an increase in slurry-based stable systems. Altogether, the emission of CH₄ from the agriculture sector has decreased by 0.7 % from 1990 to 2012. The emission of CH₄ from solid waste disposal has decreased 48.9 % since 1990 due to an increase in the incineration of waste and hence a decrease in the waste being deposited at landfills and a ban on depositing waste fit for incineration.




Figure 2.4 CH₄ emissions. Distribution according to the main sectors (2012) and time series for 1990 to 2012.

2.2.4 HFCs, PFCs and SF₆

This part of the Danish inventory only comprises a full data set for all substances from 1995. From 1995 to 2000, there has been a continuous and sub-

stantial increase in the contribution from the range of F-gases as a whole, calculated as the sum of emissions in CO₂ equivalents, see Figure 2.5. This increase is simultaneous with the increase in the emission of HFCs. For the time series 2000-2012, the increase is lower than for the years 1995 to 2000. The increase from 1995 to 2012 for the total F-gas emission is 140.7 %, while emissions decreased from 2011 to 2012 by 7.0 % mainly due to decreasing emissions of HFCs. SF₆ contributed considerably to the F-gas sum in earlier years, with 33 % in 1995. Environmental awareness and regulation of this gas under Danish law has reduced its use in industry, see Figure 2.5. A further result is that the contribution of SF₆ to F-gases in 2012 was only 15.0 %. The use of HFCs has increased several folds. HFCs have, therefore, become even more dominant, comprising 66.9 % in 1995, but 83.9 % in 2012. HFCs are mainly used as a refrigerant. Danish legislation regulates the use of F-gases, e.g. since January 1, 2007, new HFC-based refrigerant stationary systems are forbidden. Refill of old systems is still allowed. The use of air conditioning in mobile systems and the amount of HFC for this purpose increases.




Figure 2.5 F-gas emissions. Time series for 1990 to 2012.

2.3 Description and interpretation of emission trends by source

2.3.1 Energy

The emission of CO₂ from Energy Industries has decreased by 36.8 % from 1990 to 2012. The relatively large fluctuation in the emission is due to inter-country electricity trade. Thus, the high emissions in 1991, 1994, 1996, 2003 and 2006 reflect a large electricity export and the low emissions in 1990, 1992, 2005, 2008, 2011 and 2012 are due to a large import of electricity. The large decrease from 2011 to 2012 owe to decreasing fuel consumption, mainly for coal and natural gas. The decrease is mainly caused by decreasing emissions from fuel combustion due to increasing import of electricity and increasing production of wind power.

The increasing emission of CH₄ during the nineties is due to the increasing use of gas engines in decentralised cogeneration plants. The CH₄ emissions from this sector have been decreasing since 2001 due to the liberalisation of the electricity market. The CO₂ emission from the transport sector increased

by 14.0 % from 1990 to 2012, mainly due to increasing emissions from road traffic.

2.3.2 Industrial processes

The emissions from industrial process, i.e. emissions from processes other than fuel combustion, amount in 2012 to 2.6 % of the total emission in CO₂ equivalents (excl. LULUCF). The main sources are cement production, refrigeration, foam blowing and calcination of limestone. The CO₂ emission from cement production – which is the largest source contributing in 2012 with 1.7 % of the national total – decreased by 1.3 % from 1990 to 2012. The second largest source has previously been N₂O from the production of nitric acid. However, the production of nitric acid/fertiliser ceased in 2004 and therefore the emission of N₂O also ceased. The total emission in CO₂ equivalents has decreased by 3.3 % from 2011 to 2012 due to decreasing F-gas emissions, especially of HFCs. This follows the recent trend in emissions of HFCs caused by the legislation banning certain uses.

2.3.3 Agriculture

The agricultural sector contributes in 2012 with 18.6 % of the total greenhouse gas emission in CO₂ equivalents (excl. LULUCF) and is the most important sector regarding the emissions of N₂O and CH₄. In 2012, the contribution of N₂O and CH₄ to the total emission of these gases was 90.5 % and 76.6 %, respectively. The N₂O emission from agriculture decreased by 34.9 % and the CH₄ emission including field burning and reduction of biogas decreased by 0.7 % from 1990 to 2012. The CH₄ emission has decreased by 1.3 % from 2011 to 2012, while the N₂O emission has increased by 0.3 %.

2.3.4 Forestry

The trend in CO₂ uptake from forests varies greatly due to several factors both relating to weather and other effects. The calculations for 2012 estimates the Danish forests to be a large sink of 4 441 Gg CO₂ equivalents.

The area of forests established before 1990 has decreased by 1 % from 1990 to 2012, leading to a decrease of the carbon stock. From 2011 to 2012 the area has decreased by 364 ha, leading to an emission of 4 491 Gg. The area of afforestation for the years 1990-2012 is 94 kha and the area of deforestation for the years 1990-2012 is 5.8 kha.

2.3.5 Cropland, Grassland and Wetlands

The carbon stock change for organic soils has been estimated to 540 Gg C in 2012. For the same year the total emission from Cropland is estimated to 2995 Gg CO₂.

Grassland is estimated to have an emission of 554 Gg CO₂ equivalents in 2012, which is an increase since 1990, where the emission from grassland is estimated to 184 Gg CO₂ equivalents.

Emissions from managed wetlands with peat extraction are at a low level for all years in the time series. In 2012 the emission from wetlands was 2.3 Gg CO₂ equivalents. In 1990 the estimated emission is 88 Gg equivalents.

The overall trend in the LULUCF sector without Forestry is a decrease of 30 % since 1990.

2.3.6 Waste

The waste sector contributes in 2012 with 2.1 % to the national total of greenhouse gas emissions (excl. LULUCF), 15.7 % of the total CH₄ emission and 3.4 % of the total N₂O emission. In 2012 the contribution of CH₄ from SWDS was 12.7 % of the total CH₄ emission. The CH₄ emission from WW amounts in 2012 to 1.3 % of the total CH₄ emissions. The emission of N₂O from WW in 2012 is 1.2 % of national total of N₂O. Since all incinerated waste is used for power and heat production, the emissions are included in the 1A CRF category.

The emission in CO₂ equivalents has decreased by 2.9 % from 2011 to 2012 mainly due to decreasing CH₄ emissions from solid waste disposal on land. The GHG emission from the sector has decreased by 32.4 % from 1990 to 2012. This decrease is a result of (1) a decrease in the CH₄ emission from solid waste disposal sites (SWDS) by 48.9 % due to the increasing use of waste for power and heat production, and (2) a decrease in emission of N₂O from wastewater (WW) handling systems of 30.6 % due to upgrading of WW treatment plants. These decreases are counteracted by an increase in CH₄ from WW of 13.0 % due to increasing industrial load to WW systems. The emission in CO₂ equivalents has decreased by 2.9 % from 2011 to 2012.

2.4 Description and interpretation of emission trends for indirect greenhouse gases and SO₂

2.4.1 NO_x

The largest sources of emissions of Nitrogen oxide (NO_x) are road transport followed by other mobile sources and combustion in energy industries (mainly public power and district heating plants). The transport sector is the sector contributing the most to the emission of NO_x and, in 2012, 49 % of the Danish emissions of NO_x stems from road transport, national navigation, railways and civil aviation. Also emissions from national fishing and off-road vehicles contribute significantly to the NO_x emission. For non-industrial combustion plants, the main sources are combustion of gas oil, natural gas and wood in residential plants. The emissions from energy industries have decreased by 76.9 % from 1990 to 2012. In the same period, the total emission decreased by 58.3 %. The reduction is due to the increasing use of catalyst cars and installation of low-NO_x burners and denitrifying units in power plants and district heating plants.




Figure 2.6 NO_x emissions. Distribution according to the main sectors (2012) and time series for 1990 to 2012.

2.4.2 CO

Non-industrial combustion plants and transport are by far the major contributors to the total emission of this pollutant with 63.5 % and 28.4 % of the total CO emission. The total CO emission decreased by 50.8 % from 1990 to 2012, largely because of decreasing emissions from road transportation due to the introduction of private catalyst cars in 1990 and the introduction of even more emission efficient private cars in the following years.




Figure 2.7 CO emissions. Distribution according to the main sectors (2012) and time series for 1990 to 2012.

2.4.3 NMVOC

The emissions of NMVOC originate from many different sources and can be divided into two main groups: incomplete combustion and evaporation.

In the nineties road vehicles and other mobile sources such as national navigation vessels and off-road machinery were the main sources of NMVOC emissions from incomplete combustion processes. Road transportation is still a main contributor even though the emissions have declined since the introduction of catalyst cars in 1990, but since 2006 emissions from non-industrial combustion, mainly residential wood combustion, has been the largest source. In 2012 non-industrial combustion and transport contribute 27.5 % and 14.5 % of the total NMVOC emission.

The evaporative emissions mainly originate from the use of solvents and the extraction, handling and storage of oil and natural gas. The emissions from the energy industries have increased during the nineties due to the increasing use of stationary gas engines, which have much higher emissions of NMVOC than conventional boilers.

The total NMVOC emissions have decreased by 58.7 % from 1990 to 2012, mainly due to the increased use of catalysts in cars and reduced emissions from use of solvents.




Figure 2.8 NMVOC emissions. Distribution according to the main sectors (2012) and time series for 1990 to 2012.

2.4.4 SO₂

The main part of the SO₂ emission originates from combustion of fossil fuels, i.e. mainly coal and oil, in public power and district heating plants. From 1990 to 2012, the total emission decreased by 93.0 %. The large reduction is mainly due to installation of desulphurisation plant and use of fuels with lower content of sulphur in public power and district heating plants. Despite the large reduction of the SO₂ emissions, these plants make up 24.6 % of the total emission in 2012. Also emissions from industrial combustion plants, non-industrial combustion plants and other mobile sources are important. National sea traffic (navigation and fishing) contributes with about 13.1 % of the total SO₂ emission. This is due to the use of residual oil with high sulphur content.




Figure 2.9 SO₂ emissions. Distribution according to the main sectors (2012) and time series for 1990 to 2012.

2.5 Description and interpretation of emission trends for KP-LULUCF inventory in aggregate, by activity and by gas

Coverage relating to reporting of activities under Article 3.3 and selected activities under Article 3.4 are listed in Table 2.1 for reporting concerning change in carbon pool and for greenhouse gas sources. All pools are reported. Carbon stock change in below-ground biomass for Cropland Management and Grazing Land Management under Article 3.4 are included under Above-ground biomass for the same area categories. Fertilisation of forests

and other land is negligible and all fertiliser consumption is therefore reported in the agricultural sector. All liming is reported under Cropland because only very limited amounts are used in forestry and on permanent grassland. Field burning of wooden biomass is prohibited in Denmark and therefore reported as not occurring. Wildfires are very seldom and if occurring very small in Denmark.

CO₂ is by far the most important greenhouse gas relating to activities under Article 3.3 and Article 3.4. There is however a minor contribution of N₂O due to Deforestation (0.9 % of GHG from Deforestation in 2012) and Forest Management (0.3 % of GHG from Forest Management in 2012). Additionally, there is a small emission of greenhouse gases from biomass burning of 0.05 Gg CO₂ equivalents in 2012.

Table 2.1 Coverage of reporting of change of carbon pools relating to activities under Article 3.3 and elected activities under Article 3.4.

		Change in carbon pool reported					Greenhouse gas sources reported							
		Above-ground bio-mass	Below-ground bio-mass	Litter	Dead wood	Soil	Fertilization	Drainage of soils under forest manage- ment	Disturbance associated with land- use conver- sion to croplands	Liming	Biomass burning			
							N ₂ O	N ₂ O	N ₂ O	CO ₂	CO ₂	CH ₄	N ₂ O	
Article 3.3	Afforestation and Reforestation	R	R	R	R	R	IE			IE	NO	IE	IE	
	Deforestation	R	R	R	R	R			R	IE	NO	IE	IE	
Article 3.4	Forest Management	R	R	R	R	R	IE	R		IE	NO	R	R	
	Cropland Management	R	IE	NO	NO	R			R	R	NO	NO	NO	
	Grazing Land Management	R	IE	NO	NO	R				IE	NO	R	R	
	Revegetation	NA	NA	NA	NA	NA				NA	NA	NA	NA	

R: reported, NR: not reported, IE: included elsewhere, NO: not occurring, NA: not applicable. Biomass burning does not occur in all years and therefore sometimes reported as NO in the CRF.

2.5.1 Forest

The trends in forest in the first commitment period are dependent on both the current structure of the forests and the management actions in the coming years. If similar management is applied as in the previous 15 years a decline in the total carbon stock in the forest is expected. However, for 2008 to 2012 a sink in forest is reported. For the afforested areas a steady increase in carbon stocks is expected also in the future years. The rate of increase of area will depend on both availability of land and on possible subsidies for afforestation. Deforestation occurs mainly in relation to other specific projects e.g. for nature restoration or test areas for wind turbines.

2.5.2 Cropland, Grassland and Wetlands

The trend for the Cropland Management and Grazing Land Management under KP-LULUCF indicates that there has been a stabilisation of the loss of

carbon from agricultural soils compared to previous due to an increased input of organic matter in the soil. However, the loss depends much of the climatic conditions. As a consequence of the global warming, where 18 years out of the last 20 years has been above the average for 1961-1990, it is difficult to avoid substantial losses of carbon from the agricultural soils in the future. The changes in Cropland Management since 1990 have undoubtedly prevented further losses of soil carbon. A further increase in the actual temperature will affect the ability to prevent further losses of soil carbon.

The reestablishment of wetlands on agricultural land is especially targeted towards organic soils, which leads to a decreased emission from these soils. Further reestablishments are expected to take place in the future.

3 Energy (CRF sector 1)

3.1 Overview of the sector

The data presented in Chapter 3 relates to Denmark only, whereas information for Greenland is included in Chapter 16 and for the Faroe Islands in Annex 8.

The energy sector has been reported in four main chapters:

- 3.2 Stationary combustion plants (CRF sector 1A1, 1A2 and 1A4)
- 3.3 Transport (CRF sector 1A2, 1A3, 1A4 and 1A5)
- 3.4 Additional information on fuel combustion (Reference approach)
- 3.5 Fugitive emissions (CRF sector 1B)

Though industrial combustion is part of stationary combustion, detailed documentation for some of the specific industries is discussed in the industry chapters. Table 3.1.1 shows detailed source categories for the energy sector and plant category in which the sector is discussed in this report.

Table 3.1.1 CRF energy sectors and relevant NIR chapters.

IPCC id	IPCC sector name	NIR chapter
1	Energy	
1A	Fuel Combustion Activities	
1A1	Energy Industries	
1A1a	Electricity and Heat Production	Stationary combustion
1A1b	Petroleum Refining	Stationary combustion, Fugitive
1A1c	Solid Fuel Transf./Other Energy Industries	Stationary combustion
1A2	Fuel Combustion Activities/Industry (ISIC)	
1A2a	Iron and Steel	Stationary combustion, Industrial processes
1A2b	Non-Ferrous Metals	Stationary combustion, Industrial processes
1A2c	Chemicals	Stationary combustion, Industrial processes
1A2d	Pulp, Paper and Print	Stationary combustion
1A2e	Food Processing, Beverages and Tobacco	Stationary combustion, Industrial processes
1A2f	Other (please specify)	Stationary combustion, Transport, Industrial processes
1A3	Transport	
1A3a	Civil Aviation	Transport
1A3b	Road Transportation	Transport
1A3c	Railways	Transport
1A3d	Navigation	Transport
1A3e	Other (please specify)	NO
1A4	Other Sectors	
1A4a	Commercial/Institutional	Stationary combustion, Transport
1A4b	Residential	Stationary combustion, Transport
1A4c	Agriculture/Forestry/Fishing	Stationary combustion, Transport
1A5	Other (please specify)	
1A5a	Stationary	NO
1A5b	Mobile	Transport
1B	Fugitive Emissions from Fuels	
1B1	Solid Fuels	
1B1a	Coal Mining	NO
1B1a1	Underground Mines	NO
1B1a2	Surface Mines	NO
1B1b	Solid Fuel Transformation	NO
1B1c	Other (please specify)	NO
1B2	Oil and Natural Gas	
1B2a	Oil	Fugitive
1B2a2	Production	Fugitive
1B2a3	Transport	Fugitive
1B2a4	Refining/Storage	Fugitive
1B2a5	Distribution of oil products	Fugitive
1B2a6	Other	NO
1B2b	Natural Gas	Fugitive
1B2b1	Production/processing	Fugitive

<i>Continued</i>		
1B2b2	Transmission/distribution	Fugitive
1B2c	Venting and Flaring	Fugitive
1B2c1	Venting and Flaring Oil	Fugitive
1B2c2	Venting and Flaring Gas	Fugitive
1B2d	Other	NO

Summary tables for the energy sector are shown below.

Table 3.1.2 CO₂ emissions from the energy sector.

Greenhouse gas source categories	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
(Gg)										
1. Energy	51,630	62,133	56,287	58,596	62,548	59,342	72,554	63,058	59,015	56,436
A. Fuel Combustion (Sectoral Approach)	51,303	61,487	55,621	58,017	61,979	58,893	72,061	62,363	58,500	55,343
1. Energy Industries	26,146	35,015	30,086	31,662	35,723	32,168	44,432	35,320	31,677	28,588
2. Manufacturing Industries and Construction	5,444	5,975	5,837	5,702	5,786	5,909	6,058	6,110	6,129	6,219
3. Transport	10,619	11,002	11,201	11,321	11,803	11,940	12,189	12,381	12,353	12,373
4. Other Sectors	8,976	9,209	8,355	9,094	8,414	8,623	9,206	8,382	8,137	7,980
5. Other	119	287	141	237	252	252	176	171	204	182
B. Fugitive Emissions from Fuels	327	645	666	580	570	449	494	695	515	1,094
1. Solid Fuels	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO
2. Oil and Natural Gas	327	645	666	580	570	449	494	695	515	1,094

Continued	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
(Gg)										
1. Energy	52,083	53,756	53,349	58,605	53,015	49,430	57,362	52,594	49,406	47,508
A. Fuel Combustion (Sectoral Approach)	51,363	52,986	52,702	57,936	52,263	48,888	56,830	52,050	49,014	47,243
1. Energy Industries	25,554	26,852	27,071	31,815	25,933	22,731	30,644	26,006	23,890	23,823
2. Manufacturing Industries and Construction	6,016	6,107	5,816	5,780	5,836	5,551	5,672	5,519	5,061	4,111
3. Transport	12,173	12,184	12,282	12,738	13,047	13,166	13,544	14,161	13,860	13,135
4. Other Sectors	7,509	7,745	7,444	7,511	7,209	7,168	6,843	6,188	6,096	6,014
5. Other	111	97	89	92	239	271	126	175	108	160
B. Fugitive Emissions from Fuels	720	770	647	670	752	543	532	544	392	265
1. Solid Fuels	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO
2. Oil and Natural Gas	720	770	647	670	752	543	532	544	392	265

Continued	2010	2011	2012
(Gg)			
1. Energy	47,891	42,856	38,244
A. Fuel Combustion (Sectoral Approach)	47,533	42,600	38,023
1. Energy Industries	23,625	19,746	16,531
2. Manufacturing Industries and Construction	4,524	4,494	4,235
3. Transport	13,068	12,703	12,103
4. Other Sectors	6,209	5,465	5,038
5. Other	107	193	116
B. Fugitive Emissions from Fuels	357	256	221
1. Solid Fuels	NA,NO	NA,NO	NA,NO
2. Oil and Natural Gas	357	256	221

Table 3.1.3 CH₄ emissions from the energy sector.

Greenhouse gas source categories	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
(Gg)										
1. Energy	10.82	12.34	12.81	15.12	18.66	24.98	29.09	29.05	30.45	30.74
A. Fuel Combustion (Sectoral Approach)	8.71	9.71	10.32	12.40	15.65	21.52	25.76	25.47	26.92	26.85
1. Energy Industries	0.68	1.03	1.44	3.05	6.13	11.47	14.64	13.97	15.34	15.45
2. Manufacturing Industries and Construction	0.36	0.38	0.36	0.37	0.37	0.42	0.79	0.79	0.89	0.88
3. Transport	2.28	2.38	2.39	2.38	2.36	2.28	2.21	2.14	2.07	1.96
4. Other Sectors	5.38	5.90	6.11	6.59	6.78	7.33	8.12	8.55	8.61	8.55
5. Other	0.01	0.02	0.01	0.01	0.01	0.02	0.01	0.01	0.01	0.01
B. Fugitive Emissions from Fuels	2.11	2.63	2.49	2.73	3.01	3.46	3.33	3.58	3.53	3.89
1. Solid Fuels	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO
2. Oil and Natural Gas	2.11	2.63	2.49	2.73	3.01	3.46	3.33	3.58	3.53	3.89
Continued	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
(Gg)										
1. Energy	30.42	31.45	30.90	30.62	31.22	29.58	29.59	27.83	27.25	24.88
A. Fuel Combustion (Sectoral Approach)	26.41	27.17	26.59	26.21	25.98	24.37	23.07	21.53	21.21	19.24
1. Energy Industries	14.74	15.64	15.22	14.49	14.18	12.55	11.64	9.72	10.25	8.98
2. Manufacturing Industries and Construction	1.08	1.11	0.99	0.97	0.96	0.84	0.76	0.55	0.60	0.55
3. Transport	1.83	1.72	1.62	1.54	1.44	1.32	1.22	1.12	0.96	0.82
4. Other Sectors	8.75	8.70	8.76	9.20	9.39	9.65	9.45	10.13	9.40	8.88
5. Other	0.01	0.01	0.00	0.00	0.01	0.01	0.01	0.01	0.00	0.01
B. Fugitive Emissions from Fuels	4.02	4.27	4.30	4.41	5.24	5.21	6.52	6.30	6.05	5.64
1. Solid Fuels	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO
2. Oil and Natural Gas	4.02	4.27	4.30	4.41	5.24	5.21	6.52	6.30	6.05	5.64
Continued	2010	2011	2012							
(Gg)										
1. Energy	26.92	23.82	20.18							
A. Fuel Combustion (Sectoral Approach)	21.60	18.54	15.14							
1. Energy Industries	11.17	9.42	6.61							
2. Manufacturing Industries and Construction	0.62	0.58	0.44							
3. Transport	0.74	0.66	0.58							
4. Other Sectors	9.07	7.87	7.50							
5. Other	0.00	0.01	0.00							
B. Fugitive Emissions from Fuels	5.32	5.28	5.05							
1. Solid Fuels	NA,NO	NA,NO	NA,NO							
2. Oil and Natural Gas	5.32	5.28	5.05							

Table 3.1.4 N₂O emissions from the energy sector.

Greenhouse gas source categories	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
(Gg)										
1. Energy	1.05	1.18	1.14	1.16	1.20	1.19	1.35	1.28	1.23	1.23
A. Fuel Combustion (Sectoral Approach)	1.04	1.17	1.14	1.16	1.19	1.19	1.35	1.28	1.22	1.22
1. Energy Industries	0.28	0.36	0.32	0.35	0.38	0.36	0.49	0.42	0.38	0.38
2. Manufacturing Industries and Construction	0.18	0.19	0.19	0.17	0.16	0.15	0.15	0.15	0.15	0.15
3. Transport	0.36	0.37	0.39	0.40	0.42	0.44	0.46	0.47	0.47	0.47
4. Other Sectors	0.23	0.24	0.23	0.24	0.23	0.23	0.23	0.22	0.21	0.21
5. Other	0.00	0.01	0.00	0.01	0.01	0.01	0.01	0.00	0.01	0.01
B. Fugitive Emissions from Fuels	0.00	0.00	0.01	0.00	0.00	0.00	0.00	0.01	0.00	0.01
1. Solid Fuels	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO
2. Oil and Natural Gas	0.00	0.00	0.01	0.00	0.00	0.00	0.00	0.01	0.00	0.01
Continued	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
(Gg)										
1. Energy	1.20	1.22	1.23	1.29	1.24	1.22	1.31	1.28	1.25	1.20
A. Fuel Combustion (Sectoral Approach)	1.19	1.22	1.23	1.29	1.23	1.22	1.31	1.27	1.24	1.19
1. Energy Industries	0.36	0.37	0.38	0.42	0.37	0.33	0.40	0.34	0.33	0.33
2. Manufacturing Industries and Construction	0.15	0.14	0.14	0.13	0.13	0.12	0.15	0.14	0.14	0.11
3. Transport	0.46	0.46	0.46	0.47	0.48	0.47	0.47	0.48	0.47	0.44
4. Other Sectors	0.23	0.24	0.24	0.26	0.25	0.28	0.29	0.30	0.31	0.30
5. Other	0.00	0.00	0.00	0.00	0.01	0.01	0.00	0.01	0.00	0.01
B. Fugitive Emissions from Fuels	0.01	0.01	0.00	0.01	0.01	0.00	0.00	0.00	0.00	0.00
1. Solid Fuels	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO
2. Oil and Natural Gas	0.01	0.01	0.00	0.01	0.01	0.00	0.00	0.00	0.00	0.00
Continued	2010	2011	2012							
(Gg)										
1. Energy	1.23	1.17	1.11							
A. Fuel Combustion (Sectoral Approach)	1.23	1.17	1.10							
1. Energy Industries	0.35	0.31	0.28							
2. Manufacturing Industries and Construction	0.13	0.12	0.12							
3. Transport	0.44	0.44	0.42							
4. Other Sectors	0.31	0.29	0.28							
5. Other	0.00	0.01	0.00							
B. Fugitive Emissions from Fuels	0.00	0.00	0.00							
1. Solid Fuels	NA,NO	NA,NO	NA,NO							
2. Oil and Natural Gas	0.00	0.00	0.00							

3.2 Stationary combustion (CRF sector 1A1, 1A2 and 1A4)

Stationary combustion is the largest source of CO₂ emission in Denmark accounting for 58 % of the national total CO₂ emissions (excl. LULUCF) in 2012. The CO₂ emission from stationary combustion has decreased by 15 % since 2011 and decreased by 40 % since 1990. The decreased emission since 1990 is a result of a change of fuels; the consumption of coal has decreased whereas the consumption of natural gas and biomass has increased since 1990. The relatively large fluctuations in the CO₂ emission time series from 1990 to 2012 are due to inter-country electricity trade fluctuations caused mainly by variation in hydropower generation in Norway and Sweden. The

CO₂ emission in 2012 was lower than in 2011 due to a higher electricity import in 2012 than in 2011.

The methane (CH₄) emission from stationary combustion plants accounted for 5 % of the national CH₄ emission in 2012. The CH₄ emission from stationary combustion has increased by a factor of 2.3 since 1990. This results from the considerable number of lean-burn gas engines installed in CHP plants in Denmark during the 1990s. The deregulation of the electricity market has made production of electricity in gas engines less favourable, therefore the fuel consumption and CH₄ emission has decreased since 2004. The CH₄ emission in 2012 was 19 % lower than in 2011 due to lower fuel consumption in gas engines.

The nitrous oxide (N₂O) emission from stationary combustion plants accounted for 3 % of the national N₂O emission in 2012. The N₂O emission from stationary combustion has increased by 2 % since 1990, but as for CO₂, fluctuations in emission level due to electricity import/export are considerable. The emission in 2012 was 7 % lower than in 2011 as a result of the higher electricity import in 2012.

3.2.1 Source category description

Source category definition

Stationary combustion plants are included in the emission source subcategories to *Energy, Fuel combustion*:

- 1A1 Energy Industries.
- 1A2 Manufacturing Industries and Construction.
- 1A4 Other Sectors.

However, the emission source categories *1A2 Manufacturing Industries and Construction* and *1A4 Other Sectors* also include emissions from mobile combustion. The emission source category *Manufacturing Industries and Construction* includes emissions from non-road machinery reported separately in the CRF. The emission source *1A4* also includes non-road machinery and in the CRF, the emissions from stationary and mobile sources have been reported together.

The emission and fuel consumption data included in tables and figures in Chapter 3.2 only include emissions originating from stationary combustion plants of a given CRF source category. The CRF source category codes have been applied unchanged, but some source category names have been changed to reflect the stationary combustion element of the source.

In the Danish emission database all activity rates and emissions are defined in SNAP sector categories (Selected Nomenclature for Air Pollution) according to the CORINAIR system. The emission inventories are prepared from a complete emission database based on the SNAP source categories. Danish Centre for Environment and Energy, Aarhus University (DCE) has modified the SNAP categorisation to enable direct reporting of the disaggregated data for manufacturing industries and construction. Aggregation to the IPCC source category codes is based on a correspondence list enclosed in Annex 3A-1. Stationary combustion is defined as combustion activities in the SNAP sectors 01 – 03, not including SNAP 0303.

The CO₂ emission from calcinations is not part of the source category *Energy*. This emission is included in the source category *Industrial Processes*.

Methodology overview, tier

The type of emission factor and the applied tier level for each emission source are shown in Table 3.2.1 below. The tier level has been determined based on the 1996 IPCC Guidelines (IPCC, 1997).

The fuel consumption data for transformation are technology specific. For end-use of fuels, the disaggregation to specific technologies is less detailed. However, for residential wood combustion the technology disaggregation is technology specific.

The distinction between tier 2 and 3 has been based on the emission factor. The tier definitions have been interpreted as follows:

- Tier 1: The emission factor is an IPCC default tier 1 value.
- Tier 2: The emission factors are country specific and based on a limited number of emission measurements or IPCC tier 2 emission factors.
- Tier 3: Based on plant specific emission data or on a country specific emission factor based on a considerable number of plant specific emission measurements and detailed technology knowledge.

Table 3.2.1 gives an overview of the calculation methods and type of emission factor. The table also shows which of the source categories are key in any of the key category analysis¹ (including LULUCF, tier 1/tier 2, level/trend).

Table 3.2.1 Methodology and type of emission factor.

		Tier	EMF ¹⁾	Key category ²⁾
Stationary Combustion, Coal	CO ₂	Tier 3 ² (Tier 3) / Tier 1 ³	PS (CS) / D	Yes
Stationary Combustion, BKB	CO ₂	Tier 1	D	No
Stationary Combustion, Coke	CO ₂	Tier 1	D	No
Stationary Combustion, Fossil waste	CO ₂	Tier 3	CS	Yes
Stationary Combustion, Petroleum coke	CO ₂	Tier 3	PS/CS	Yes
Stationary Combustion, Residual oil	CO ₂	Tier 3 / Tier 3 / Tier 1	PS / CS / D ⁴	Yes
Stationary Combustion, Gas oil	CO ₂	Tier 2 / Tier 3	CR / PS	Yes
Stationary Combustion, Kerosene	CO ₂	Tier 1	D	Yes
Stationary Combustion, LPG	CO ₂	Tier 1	D	No
Stationary Combustion, Refinery gas	CO ₂	Tier 3	PS / CS	Yes
Stationary Combustion, Natural gas	CO ₂	Tier 3	CS / PS ⁵	Yes
Stationary Combustion, SOLID	CH ₄	Tier 2 / Tier 1	D(2) / D	No
Stationary Combustion, LIQUID	CH ₄	Tier 2 / Tier 2 / Tier 1	D(2) / CS / D	No
Stationary Combustion, GAS	CH ₄	Tier 2 / Tier 3	D(2) / CS	No
Natural gas fuelled engines, GAS	CH ₄	Tier 3	CS	Yes
Stationary Combustion, WASTE	CH ₄	Tier 2	CS	No
Stationary Combustion, BIOMASS	CH ₄	Tier 2 / Tier 1	D(2) / CS / D	Yes
Biogas fuelled engines, BIOMASS	CH ₄	Tier 3	CS	No
Stationary Combustion, SOLID	N ₂ O	Tier 2 / Tier 1	CS / D	Yes
Stationary Combustion, LIQUID	N ₂ O	Tier 2 / Tier 1	D(2) / D / CS	Yes
Stationary Combustion, GAS	N ₂ O	Tier 1 / Tier 2	D / CS / D(2)	Yes
Stationary Combustion, WASTE	N ₂ O	Tier 2	CS	No
Stationary Combustion, BIOMASS	N ₂ O	Tier 1 / Tier 2	D / CS / D(2)	Yes

¹⁾ D: IPCC tier 1, D(2): IPCC tier 2/3, CR: Corinair default, CS: Country specific, PS: Plant specific.

²⁾ KCA tier 1 or tier 2 for Denmark (excluding Greenland and Faroe Islands), including LULUCF, level 1990/ level 2012/ trend.

¹ Key category according to the KCA tier 1 or tier 2 for Denmark (excluding Greenland and Faroe Islands), including LULUCF, level 1990/ level 2012/ trend.

² For 2006 onwards. Country specific emission factors and tier 2 have been applied for 1990-2005.

³ Tier 1 and IPCC default emission factor is only applied for 2 % of the coal consumption in 2012.

⁴ Residual oil not applied in source category 1A1a.

⁵ Off shore gas turbines and some power plants.

The CO₂ emission factor for kerosene is a default emission factor referring to guidelines. Since the source category is a key category the emission factor will have to be revised prior to the 2015 emission inventory.

Key Categories

Key Category Analysis (KCA) tier 1 and 2 for year 1990, 2012 and trend for Denmark has been carried out in accordance with the IPCC Good Practice Guidance (IPCC, 2000) and IPCC Guidelines (IPCC, 2006). Table 3.2.2 shows which of the stationary combustion source categories that are key categories. The table is based on the analysis including LULUCF. Detailed key category analysis is shown in NIR Chapter 1.5 and Annex 1.

The CO₂ emissions from stationary combustion are key for all the major fuels. In addition, CH₄ from combustion of biomass emission is a key category in the tier 2 KCA. Due to the relatively high uncertainty for N₂O, emission factors the N₂O emission from solid, liquid, gaseous and biomass fuels are key categories in the tier 2 analysis.

Table 3.2.2 Key categories⁶, stationary combustion.

			Tier 1			Tier 2		
			1990	2012	1990-2012	1990	2012	1990-2012
Energy	Stationary Combustion, Coal	CO ₂	Level	Level	Trend	Level	Level	Trend
Energy	Stationary Combustion, BKB	CO ₂						
Energy	Stationary Combustion, Coke	CO ₂						
Energy	Stationary Combustion, Fossil waste	CO ₂	Level	Level	Trend		Level	Trend
Energy	Stationary Combustion, Petroleum coke	CO ₂	Level	Level	Trend			
Energy	Stationary Combustion, Residual oil	CO ₂	Level	Level	Trend			
Energy	Stationary Combustion, Gas oil	CO ₂	Level	Level	Trend	Level		Trend
Energy	Stationary Combustion, Kerosene	CO ₂	Level		Trend			
Energy	Stationary Combustion, LPG	CO ₂						
Energy	Stationary Combustion, Refinery gas	CO ₂	Level	Level	Trend			
Energy	Stationary Combustion, Natural gas	CO ₂	Level	Level	Trend			Trend
Energy	Stationary Combustion, SOLID	CH ₄						
Energy	Stationary Combustion, LIQUID	CH ₄						
Energy	Stationary Combustion, GAS	CH ₄						
Energy	Natural gas fuelled engines, GAS	CH ₄						
Energy	Stationary Combustion, WASTE	CH ₄						
Energy	Stationary Combustion, BIOMASS	CH ₄					Level	Trend
Energy	Biogas fuelled engines, BIOMASS	CH ₄						
Energy	Stationary Combustion, SOLID	N ₂ O				Level	Level	Trend
Energy	Stationary Combustion, LIQUID	N ₂ O				Level	Level	Trend
Energy	Stationary Combustion, GAS	N ₂ O					Level	Trend
Energy	Stationary Combustion, WASTE	N ₂ O						
Energy	Stationary Combustion, BIOMASS	N ₂ O				Level	Level	Trend

3.2.2 Fuel consumption data


In 2012, the total fuel consumption for stationary combustion plants was 439 PJ of which 310 PJ was fossil fuels and 129 PJ was biomass.

Fuel consumption distributed according to the stationary combustion sub-categories is shown in Figure 3.2.1 and Figure 3.2.2. The majority - 57 % - of all fuels is combusted in the source category, *Public electricity and heat pro-*

⁶ For Denmark not including Greenland and Faroe Island.

duction. Other source categories with high fuel consumption are *Residential* and *Industry*.

Fuel consumption including biomass



Fuel consumption, fossil fuels




Figure 3.2.1 Fuel consumption of stationary combustion source categories, 2012. Based on DEA (2013a).

Coal, natural gas and wood are the most utilised fuels for stationary combustion plants. Coal is mainly used in power plants and natural gas is used in power plants and decentralised combined heating and power (CHP) plants, as well as in industry, district heating, residential plants and off-shore gas turbines (see Figure 3.2.2). Wood is mainly applied for public electricity and heat production and in residential plants.

Detailed fuel consumption rates are shown in Annex 3A-2.




Figure 3.2.2 Fuel consumption of stationary combustion 2012, disaggregated to fuel type. Based on DEA (2013a).

Fuel consumption time series for stationary combustion plants are presented in Figure 3.2.3. The fuel consumption for stationary combustion was 12 % lower in 2012 than in 1990, while the fossil fuel consumption was 32 % lower and the biomass fuel consumption 3.2 times the level in 1990.

The consumption of natural gas, waste and biomass has increased since 1990 whereas coal and oil consumption has decreased.




Figure 3.2.3 Fuel consumption time series, stationary combustion. Based on DEA (2013a).

The fluctuations in the time series for fuel consumption are mainly a result of electricity import/export, but also of outdoor temperature variations from year to year. This, in turn, leads to fluctuations in emission levels. The fluctuations in electricity trade, fuel consumption, CO₂ and NO_x emission are illustrated and compared in Figure 3.2.4. In 1990, the Danish electricity import was large causing relatively low fuel consumption, whereas the fuel consumption was high in 1996 due to a large electricity export. In 2012, the net electricity import was 18.8, whereas there was a 4.7 PJ electricity import in 2011. The large electricity export that occurs some years is a result of low rainfall in Norway and Sweden causing insufficient hydro-power production in both countries.

The Danish electricity production is highly dependent on the electricity trade with especially Sweden and Norway. Denmark has a number of central coal-fuelled power plants that consists of a number of blocks. These do not under normal conditions operate at max load, i.e. there is free capacity for peak situations. In addition, there are blocks, which are mothballed but can be reopened in situations where there is a significant increase in the electricity demand.

To be able to follow the national energy consumption as well as for statistical and reporting purposes, the Danish Energy Agency (DEA) produces a correction of the actual fuel consumption and CO₂ emission without ran-

dom variations in electricity import/export and in ambient temperature. This fuel consumption trend is also illustrated in Figure 3.2.4. The corrections are included here to explain the fluctuations in the time series for fuel rate and emission.




Figure 3.2.4 Comparison of time series fluctuations for electricity trade, fuel consumption, CO₂ emission and NO_x emission. Based on DEA (2013a).

Fuel consumption time series for the subcategories to stationary combustion are shown in Figure 3.2.5, 3.2.6 and 3.2.7.

Fuel consumption for *Energy industries* fluctuates due to electricity trade as discussed above. The fuel consumption in 2012 was 7 % lower than in 1990. The fluctuation in electricity production is based on fossil fuel consumption in the subcategory *Public electricity and heat production*. The energy consumption in *Other energy industries* is mainly natural gas used in gas turbines in the off-shore industry. The biomass fuel consumption in *Energy industries* in 2012 added up to 79 PJ, which is 4.8 times the level in 1990 and a 5 % increase since 2011.

The fuel consumption in *Industry* was 17 % lower in 2012 than in 1990 (Figure 3.2.6). The fuel consumption in industrial plants has decreased considerably as a result of the financial crisis. However, the fuel consumption is unchanged since 2009. The biomass fuel consumption in *Industry* in 2012 added up to 9 PJ, which is a 53 % increase since 1990.

The fuel consumption in *Other Sectors* has decreased 24% since 1990 and decreased 7 % since 2011 (Figure 3.2.7). The biomass fuel consumption in *Other sectors* in 2012 added up to 41 PJ which is 2.2 times the consumption in 1990 and a 1 % decrease since 2011. Wood consumption in residential plants in 2012 was 2.2 times the consumption in year 2000.

Time series for subcategories are shown in Chapter 3.2.4.




Figure 3.2.5 Fuel consumption time series for subcategories - 1A1 Energy Industries.




Figure 3.2.6 Fuel consumption time series for subcategories - 1A2 Industry.




Figure 3.2.7 Fuel consumption time series for subcategories - 1A4 Other Sectors.

3.2.3 Emissions

Greenhouse gas emission

The greenhouse gas emissions from stationary combustion are listed in Table 3.2.3. The emission from stationary combustion accounted for 45 % of the national greenhouse gas emission (excluding LULUCF) in 2012.

The CO₂ emission from stationary combustion plants accounts for 58 % of the national CO₂ emission (excluding LULUCF). The CH₄ emission accounts for 5 % of the national CH₄ emission and the N₂O emission for 3 % of the national N₂O emission.

Table 3.2.3 Greenhouse gas emission, 2012 ¹⁾.

	CO ₂ Gg CO ₂ equivalent	CH ₄	N ₂ O
1A1 Fuel Combustion, Energy industries	16531	139	88
1A2 Fuel Combustion, Manufacturing Industries and Construction ¹⁾	3214	9	23
1A4 Fuel Combustion, Other sectors ¹⁾	2965	151	58
Emission from stationary combustion plants	22710	298	170
Emission share for stationary combustion	58%	5%	3%

¹⁾ Only stationary combustion sources of the category is included.

CO₂ is the most important greenhouse gas accounting for 98.0 % of the greenhouse gas emission (CO₂ eq.) from stationary combustion. CH₄ accounts for 1.3 % and N₂O for 0.7 % of the greenhouse gas emission (CO₂ eq.) from stationary combustion (Figure 3.2.8).




Figure 3.2.8 Stationary combustion - Greenhouse gas emission (CO₂ equivalent), contribution from each pollutant.

Figure 3.2.9 depicts the time series of greenhouse gas emissions (CO₂ eq.) from stationary combustion and it can be seen that the greenhouse gas emission development follows the CO₂ emission development very closely. Both the CO₂ and the total greenhouse gas emission are lower in 2012 than in 1990, CO₂ by 40 % and greenhouse gas by 39 %. However, fluctuations in the GHG emission level are large.




Figure 3.2.9 GHG emission time series for stationary combustion.

The fluctuations in the time series are largely a result of electricity import/export, but also of outdoor temperature variations from year to year. The fluctuations follow the fluctuations in fuel consumption discussed in Chapter 3.2.2. As mentioned in Chapter 3.2.2, the Danish Energy Agency estimates a correction of the actual CO₂ emission without random variations in electricity imports/exports and in ambient temperature. The greenhouse gas emission corrected for electricity import/export and ambient temperature has decreased by 38 % since 1990, and the CO₂ emission by 39 %. These data are included here to explain the fluctuations in the emission time series.

CO₂

The carbon dioxide (CO₂) emission from stationary combustion plants is one of the most important sources of greenhouse gas emissions. Thus, the CO₂ emission from stationary combustion plants accounts for 58 % of the national CO₂ emission. Table 3.2.4 lists the CO₂ emission inventory for stationary combustion plants for 2012. *Public electricity and heat production* accounts for 62 % of the CO₂ emission from stationary combustion. This share is somewhat higher than the fossil fuel consumption share for this category, which is 56 % (Figure 3.2.1). This is due to a large share of coal in this category. Other large CO₂ emission sources are *Industry* and *Residential* plants. These are the source categories, which also account for a considerable share of fuel consumption.

Table 3.2.4 CO₂ emission from stationary combustion plants, 2012¹⁾.


	CO ₂ Gg	
1A1a Public electricity and heat production	14111	1A4a Commercial / Institutional 3%
1A1b Petroleum refining	984	1A4b Residential 9%
1A1c Other energy industries	1437	1A4c Agriculture / Forestry / Fisheries 1.1%
1A2 Industry	3214	1A2 Industry 14%
1A4a Commercial/Institutional	653	1A1c Other energy industries 6%
1A4b Residential	2062	1A1b Petroleum refining 5%
1A4c Agriculture/Forestry/Fisheries	249	
Total	26393	

¹⁾ Only emission from stationary combustion plants in the categories is included.

In the Danish inventory, the source category *Public electricity and heat production* is further disaggregated. The CO₂ emission from each of the subcategories is shown in Table 3.2.5. The largest subcategory is power plant boilers >300MW.


Table 3.2.5 CO₂ emission from subcategories to 1A1a Public electricity and heat production.

SNAP	SNAP name	CO ₂ , Gg
0101	Public power	
010101	Combustion plants ≥ 300MW (boilers)	10204
010102	Combustion plants ≥ 50MW and < 300 MW (boilers)	1029
010103	Combustion plants <50 MW (boilers)	315
010104	Gas turbines	1021
010105	Stationary engines	578
0102	District heating plants	
010202	Combustion plants ≥ 50MW and < 300 MW (boilers)	71
010203	Combustion plants <50 MW (boilers)	894



CO₂ emission from combustion of biomass fuels is not included in the total CO₂ emission data, because biomass fuels are considered CO₂ neutral. The CO₂ emission from biomass combustion is reported as a memo item in the Climate Convention reporting. In 2012, the CO₂ emission from biomass combustion was 18 763 Gg.

In Figure 3.2.10, the fuel consumption share (fossil fuels) is compared to the CO₂ emission share disaggregated to fuel origin. Due to the higher CO₂ emission factor for coal than oil and gas, the CO₂ emission share from coal combustion is higher than the fuel consumption share. Coal accounts for 34 % of the fossil fuel consumption and for 44 % of the CO₂ emission. Natural gas accounts for 47 % of the fossil fuel consumption but only 37 % of the CO₂ emission.






Figure 3.2.10 CO₂ emission, fuel origin.

The time series for CO₂ emission is provided in Figure 3.2.11. Despite a decrease in fuel consumption of 12 %⁷ since 1990, the CO₂ emission from stationary combustion has decreased by 40 % because of the change of fuel type used.

The fluctuations in total CO₂ emission follow the fluctuations in CO₂ emission from *Public electricity and heat production* (Figure 3.2.11) and in coal consumption (Figure 3.2.4). The fluctuations are a result of electricity import/export as discussed in Chapter 3.2.2.





Figure 3.2.11 CO₂ emission time series for stationary combustion plants.

CH₄

The methane (CH₄) emission from stationary combustion plants accounts for 5 % of the national CH₄ emission. Table 3.2.6 lists the CH₄ emission inventory for stationary combustion plants in 2012. *Public electricity and heat production* accounts for 46 % of the CH₄ emission from stationary combustion. The emission from residential plants adds up to 39 % of the emission.

⁷ The consumption of fossil fuels has decreased 32 %.

Table 3.2.6 CH₄ emission from stationary combustion plants, 2012¹⁾.

¹⁾ Only emission from stationary combustion plants in the source categories is included.

The CH₄ emission factor for reciprocating gas engines is much higher than for other combustion plants due to the continuous ignition/burn-out of the gas. Lean-burn gas engines have an especially high emission factor. A considerable number of lean-burn gas engines are in operation in Denmark and in 2012, these plants accounted for 51 % of the CH₄ emission from stationary combustion plants (Figure 3.2.12). Most engines are installed in CHP plants and the fuel used is either natural gas or biogas. Residential wood combustion is also a large emission source accounting for 30 % of the emission in 2012.




Figure 3.2.12 CH₄ emission share for gas engines and residential wood combustion, 2012.

Figure 3.2.13 shows the time series for CH₄ emission. The CH₄ emission from stationary combustion has increased by a factor of 2.3 since 1990. This results from the considerable number of lean-burn gas engines installed in CHP plants in Denmark during the 1990s. Figure 3.2.14 provides time series for the fuel consumption rate in gas engines and the corresponding increase of CH₄ emission. The decline in later years is due to structural changes in the Danish electricity market, which means that the fuel consumption in gas engines has been decreasing.

The CH₄ emission from residential plants has increased since 1990 due to increased combustion of biomass in residential plants. Combustion of wood accounted for 76 % of the CH₄ emission from residential plants in 2012.




Figure 3.2.13 CH₄ emission time series for stationary combustion plants.





Figure 3.2.14 Time series for a) fuel consumption in gas engines and b) CH₄ emission from gas engines, residential wood combustion and other plants.


N₂O

The nitrous oxide (N₂O) emission from stationary combustion plants accounts for 3 % of the national N₂O emission. Table 3.2.7 lists the N₂O emission inventory for stationary combustion plants in the year 2012. *Public electricity and heat production* accounts for 46 % of the N₂O emission from stationary combustion.

Table 3.2.7 N₂O emission from stationary combustion plants, 2012¹⁾.

¹⁾ Only emission from stationary combustion plants in the source categories is included.

Figure 3.2.15 shows the time series for N₂O emission. The N₂O emission from stationary combustion has decreased by 2 % from 1990 to 2012, but again fluctuations in emission level due to electricity import/export are considerable.

Figure 3.2.15 N₂O emission time series for stationary combustion plants.

SO₂, NO_x, NMVOC and CO

The emissions of sulphur dioxide (SO₂), nitrogen oxides (NO_x), non-volatile organic compounds (NMVOC) and carbon monoxide (CO) from Danish stationary combustion plants 2012 are presented in Table 3.2.8.

SO₂ from stationary combustion plants accounts for 64 % of the national emission. NO_x, CO and NMVOC account for 28 %, 35 % and 20 % of national emissions, respectively.

Table 3.2.8 SO₂, NO_x, NMVOC and CO emission, 2012¹⁾.

Pollutant	NO _x Gg	CO Gg	NMVOC Gg	SO ₂ Gg
1A1 Fuel consumption, Energy industries	21.8	11.0	1.5	3.1
1A2 Fuel consumption, Manufacturing Industries and Construction ¹⁾	5.3	4.2	0.3	2.9
1A4 Fuel consumption, Other sectors ¹⁾	5.2	109.6	14.0	2.1
Emission from stationary combustion plants	32.3	124.8	15.7	8.1
Emission share for stationary combustion, %	28	35	20	64

¹⁾ Only emissions from stationary combustion plants in the source categories are included.

SO₂


Stationary combustion is the most important emission source for SO₂ accounting for 64 % of the national emission. Table 3.2.9 presents the SO₂ emission inventory for the stationary combustion subcategories.

Public electricity and heat production is the largest emission source accounting for 35 % of the emission. However, the SO₂ emission share is lower than the fuel consumption share for this source category, which is 57 %. This is a result of effective flue gas desulphurisation equipment installed in power plants combusting coal. In the Danish inventory, the source category *Public electricity and heat production* is further disaggregated. Figure 3.2.16 shows the SO₂ emission from *Public electricity and heat production* on a disaggregated level. Power plants >300MW_{th} are the main emission source, accounting for 41 % of the emission.

The SO₂ emission from industrial plants adds up to 36 % of the emission from stationary combustion, a remarkably high emission share compared to fuel consumption. The main emission sources in the industrial category are combustion of coal and residual oil, but emissions from the cement industry is also a considerable emission source. Until year 2000, the SO₂ emission from the industrial category only accounted for a small part of the emission from stationary combustion, but as a result of reduced emissions from power plants, the share has now increased.

The time series for SO₂ emission from stationary combustion are shown in Figure 3.2.17. The SO₂ emission from stationary combustion plants has decreased by 95 % since 1990. The large emission decrease is mainly a result of the reduced emission from *Public electricity and heat production*, made possible due to installation of desulphurisation units and due to the use of fuels with lower sulphur content. Despite the considerable reduction in emission from public electricity and heat production plants, these still account for 35 % of the emission from stationary combustion, as mentioned above. The emission from other source categories also decreased considerably since 1990. Time series for subcategories are shown in Chapter 3.2.4.

Table 3.2.9 SO₂ emission from stationary combustion plants, 2012¹⁾



¹⁾ Only emission from stationary combustion plants in the source categories is included.




Figure 3.2.16 Disaggregated SO₂ emissions from 1A1a Energy and heat production.




Figure 3.2.17 SO₂ emission time series for stationary combustion.

NO_x

Stationary combustion accounts for 28 % of the national NO_x emission. Table 3.2.10 shows the NO_x emission inventory for stationary combustion subcategories.

Public electricity and heat production is the largest emission source accounting for 43 % of the emission from stationary combustion plants. The emission from public power boilers > 300 MWth accounts for 29 % of the emission in this subcategory.

Industrial combustion plants are also an important emission source accounting for 17 % of the emission. The main industrial emission source is cement production, which accounts for 30 % of the emission.


Residential plants account for 12 % of the NO_x emission. The fuel origin of this emission is mainly wood accounting for 61 % of the residential plant emission.

Other energy industries, which is mainly off-shore gas turbines accounts for 19 % of the NO_x emission.

Time series for NO_x emission from stationary combustion are shown in Figure 3.2.18. NO_x emission from stationary combustion plants has decreased by 72 % since 1990. The reduced emission is largely a result of the reduced emission from public electricity and heat production due to installation of low NO_x burners, selective catalytic reduction (SCR) units and selective non-catalytic reduction (SNCR) units. The fluctuations in the time

series follow the fluctuations in public electricity and heat production, which, in turn, result from electricity trade fluctuations.

Table 3.2.10 NO_x emission from stationary combustion plants, 2012¹⁾.



1) Only emission from stationary combustion plants in the source categories is included.




Figure 3.2.18 NO_x emission time series for stationary combustion.

NMVOC

Stationary combustion plants accounted for 20 % of the national NMVOC emission in 2012. Table 3.2.11 presents the NMVOC emission inventory for the stationary combustion subcategories.

Residential plants are the largest emission source accounting for 85 % of the emission from stationary combustion plants. For residential plants NMVOC is mainly emitted from wood and straw combustion, see Figure 3.2.19.

Public electricity and heat production is also a considerable emission source, accounting for 9 % of the emission. Lean-burn gas engines have a relatively high NMVOC emission factor and are the most important emission source in this subcategory (see Figure 3.2.19). The gas engines are either natural gas or biogas fuelled.


Time series for NMVOC emission from stationary combustion are shown in Figure 3.2.20. The emission has increased by 12 % from 1990. The increased emission is mainly a result of the increasing wood consumption in residential plants and of the increased use of lean-burn gas engines in CHP plants.

The emission from residential plants increased 15 % since 1990. The NMVOC emission from wood combustion in 2012 was 2.3 times the 1990 level due to increased wood consumption. However, the emission factor has decreased since 1990 due to installation of modern stoves and boilers with improved combustion technology. Further the emission from straw combustion in farmhouse boilers has decreased (75 %) over this period due to both a decreasing emission factor and decrease in straw consumption in this source category.

The use of wood in residential boilers and stoves was relatively low in 1998-99 resulting in a lower emission level.

The increasing consumption of wood in residential plants ceased in 2007. The improved technology that has been implemented in residential wood combustion have led to lower emission factors and thus decreasing NMVOC emission since 2007.

Table 3.2.11 NMVOC emission from stationary combustion plants, 2012¹⁾.



1) Only emission from stationary combustion plants in the categories is included.




Figure 3.2.19 NMVOC emission from Residential plants and from Public electricity and heat production, 2012.




Figure 3.2.20 NMVOC emission time series for stationary combustion.

CO

Stationary combustion accounted for 35 % of the national CO emission in 2012. Table 3.2.12 presents the CO emission inventory for stationary combustion subcategories.

Residential plants are the largest emission source, accounting for 80 % of the emission. Wood combustion accounts for 88 % of the emission from residential plants, see Figure 3.2.21. This is in spite of the fact that the fuel consumption share is only 47 %. Combustion of straw is also a considerable emission source whereas the emission from other fuels used in residential plants is almost negligible.

Time series for CO emission from stationary combustion are shown in Figure 3.2.22. The emission has increased by 5 % from 1990. The time series for CO from stationary combustion plants follows the time series for CO emission from residential plants.

The increase of wood consumption in residential plants in 1999-2007 is reflected in the time series for CO emission. The consumption of wood in residential plants in 2012 was 3.6 times the 1990 level. The decreased emission in 2007-2012 is a result of implementation of improved residential wood combustion technologies and the fact that the rapid increase of wood consumption until 2007 have stopped.

Both straw consumption and CO emission factor for residential plants have decreased since 1990.

Table 3.2.12 CO emission from stationary combustion plants, 2012¹⁾.

	CO, Mg	
1A1a Public electricity and heat production	10728	1A4c Agriculture / Forestry / Fisheries 7%
1A1b Petroleum refining	126	1A1a Public electricity and heat production 9%
1A1c Other energy industries	120	1A1b Petroleum refining 0.1%
1A2 Industry	4201	1A1c Other energy industries 0.1%
1A4a Commercial/Institutional	715	1A2 Industry 3%
1A4b Residential	99781	1A4a Commercial/Institutional 0.6%
1A4c Agriculture/Forestry/Fisheries	9131	1A4b Residential 80%
Total	124802	

¹⁾ Only emission from stationary combustion plants in the source categories is included.





Figure 3.2.21 CO emission sources, residential plants, 2012.

Stationary combustion



1A4b Residential plants, fuel origin




Figure 3.2.22 CO emission time series for stationary combustion.

3.2.4 Trend for subsectors

In addition to the data for stationary combustion, this chapter presents and discusses data for each of the subcategories in which stationary combustion is included. Time series are presented for fuel consumption and emissions.

1A1 Energy industries

The emission source category *1A1 Energy Industries* consists of the subcategories:

- 1A1a Public electricity and heat production.
- 1A1b Petroleum refining.
- 1A1c Other energy industries.

Figure 3.2.23 – 3.2.25 present time series for the *Energy Industries*. *Public electricity and heat production* is the largest subcategory accounting for the main part of all emissions. Time series are discussed below for each subcategory.




Figure 3.2.23 Time series for fuel consumption, 1A1 Energy industries.




Figure 3.2.24 Time series for greenhouse gas emission, 1A1 Energy industries.




Figure 3.2.25 Time series for SO₂, NO_x, NMVOC and CO emission, 1A1 Energy industries.

1A1a Public electricity and heat production

Public electricity and heat production is the largest source category regarding both fuel consumption and greenhouse gas emissions for stationary combustion. Figure 3.2.26 shows the time series for fuel consumption and emissions.

The fuel consumption in public electricity and heat production was 13 % lower in 2012 than in 1990. As discussed in Chapter 3.2.2 the fuel consumption fluctuates mainly as a consequence of electricity trade. Coal is the fuel that is affected the most by the fluctuating electricity trade. Coal is the main fuel in the source category even in years with electricity import. The coal consumption in 2012 was 57 % lower than in 1990. Natural gas is also an important fuel and the consumption of natural gas has increased since 1990, but decreased since 2003. A considerable part of the natural gas is combusted in gas engines (Figure 3.2.23). The consumption of waste and biomass has increased.

The CO₂ emission was 43 % lower in 2012 than in 1990. This decrease – in spite of only a 13 % decrease in fuel consumption – is a result of the change of fuels used as discussed above.

The CH₄ emission has increase until the mid-nineties as a result of the considerable number of lean-burn gas engines installed in CHP plants in Denmark in this period. The decline in later years is due to structural changes in the Danish electricity market, which means that the fuel consumption in gas engines has been decreasing (Figure 3.2.23). The emission in 2012 was 10 times the 1990 emission level.

The N₂O emission in 2012 was equal to the 1990 emission level. The emission fluctuates similar to the fuel consumption.

The SO₂ emission has decreased 98 % from 1990 to 2012. This decrease is a result of both lower sulphur content in fuels and installation and improved performance of desulphurisation plants. The emission was higher in 2012 than in 2011, but the emission has however decreased 64 % since 2005.

The NO_x emission has decreased 85 % since 1990 due to installation of low NO_x burners, selective catalytic reduction (SCR) units and selective non-catalytic reduction (SNCR) units. The fluctuations in time series follow the fluctuations in fuel consumption and electricity trade.

The emission of NMVOC in 2012 was 3.2 times the 1990 emission level. This is a result of the large number of gas engines that has been installed in Danish CHP plants. The decreasing emission in 2004-2012 is results of the time series for natural gas consumption in gas engines (Figure 3.2.23). In addition, the emission of NMVOC from engines decreased in 1995-2007 as a result of introduction of an emission limits for unburned hydrocarbon⁸ (DEPA, 2005).

The CO emission was 35 % higher in 2012 than in 1990. The fluctuations follow the fluctuations of the fuel consumption. In addition, the emission from gas engines is considerable.

⁸ Including methane.




Figure 3.2.26 Time series for 1A1a Public electricity and heat production.

1A1b Petroleum refining

Petroleum refining is a small source category regarding both fuel consumption and greenhouse gas emissions for stationary combustion. There are presently only two refineries operating in Denmark. Figure 3.2.27 shows the time series for fuel consumption and emissions.

The significant decrease in both fuel consumption and emissions in 1996 is a result of the closure of a third refinery.

The fuel consumption has increased 9 % since 1990 and the CO₂ emission also increased 9 %.

The CH₄ emission has increased 11 % since 1990 and 6 % since 2011. The reduction in CH₄ emission from 1995 to 1996 is caused by the closure of a refinery.

The N₂O emission was 71 % higher in 2012 than in 1990. The emission increased in 1990 – 1993 as a result of the installation of a gas turbine in one of the refineries. The gas turbine was installed in 1993 (DEA, 2013b).

The N₂O emission factor for the refinery gas fuelled gas turbine has been assumed equal to the emission factor for natural gas fuelled turbines and thus the emission factor have been decreasing since 1994. This causes the decreasing trend in the time series since 1994.

The emission of SO₂ has shown a pronounced decrease (74 %) since 1990, mainly due to the decreased consumption of residual oil (38 %) also shown in Figure 3.2.27. The NO_x emission in 2012 was 10 % lower than in 1990. Since 2005, data for both SO₂ and NO_x are plant specific data stated by the refineries.

The NMVOC emission time series follows the time series for fuel consumption.

Emissions from refineries are further discussed in Chapter 3.5.




Figure 3.2.27 Time series for 1A1b Petroleum refining.

1A1c Other energy industries

The source category *Other energy industries* comprises natural gas consumption in the off-shore industry and in addition a small consumption in the Danish gas treatment plant⁹. Gas turbines are the main plant type. Figure 3.2.28 shows the time series for fuel consumption and emissions.

The fuel consumption in 2012 was 2.6 times the consumption in 1990. The CO₂ emission follows the fuel consumption and the emission in 2012 was also 2.6 times the emission in 1990.

⁹ Nybro.

The time series for CO and N₂O will be recalculated in the next inventory. The decreasing emission factor time series will be corrected in the next inventory. The current emission factor time series for CO and N₂O follow the emission factors for gas turbines applied in CHP plants.

The emissions from all other pollutants follow the increase of fuel consumption.




Figure 3.2.28 Time series for 1A1c Other energy industries.

1A2 Industry

Manufacturing industries and construction (Industry) consists of both stationary and mobile sources. In this chapter, only stationary sources are included.

The emission source category *1A2 Industry* consists of the subcategories:

- 1A2a Iron and steel
- 1A2b Non-ferrous metals
- 1A2c Chemicals
- 1A2d Pulp, Paper and Print
- 1A2e Food processing, beverages and tobacco
- 1A2f i Industry-Other

Figures 3.2.29-3.2.31 show the time series for fuel consumption and emissions. The subsector *Industry – Other* is the main subsector for fuel consumption and emissions. *Food processing, beverages and tobacco* is also an important subsector.

The total fuel consumption in industrial combustion was 17 % lower in 2012 than in 1990. The consumption of natural gas has increased since 1990 whereas the consumption of coal has decreased. The consumption of residual oil has decreased, but the consumption of petroleum coke increased. The biomass consumption has increased 53 % since 1990.

The greenhouse gas emission and the CO₂ emission are both rather stable until 2006 following the small fluctuations in fuel consumption. After 2006, the fuel consumption has decreased. Due to change of applied fuels, the greenhouse gas and CO₂ emissions have decreased more than the fuel consumption since 1990; both emissions have decreased 30 %.

The CH₄ emission has increased from 1994-2001 and decreased again from 2001 - 2007. In 2012, the emission was 34 % higher than in 1990. The CH₄ emission follows the consumption of natural gas in gas engines (Figure 3.2.29). Most industrial CHP plants based on gas engines came in operation in the years 1995 to 1999. The decrease in later years is a result of the liberalisation of the electricity market.

The N₂O emission has decreased 47 % since 1990, mainly due to the decreased residual oil consumption. In recent years, combustion of wood is a considerable emission source.

The SO₂ emission has decreased 82 % since 1990. This is mainly a result of lower consumption of residual oil in the industrial sector. Further, the sulphur content of residual oil and several other fuels has decreased since 1990 due to legislation and tax laws.

The NO_x emission has decreased 57 % since 1990 due to the reduced emission from industrial boilers in general. Cement production is the main emission source accounting for more than 50 % of the industrial emission in 1990-2009¹⁰. In 2012, the NO_x emission from cement industry was 33 % of the industrial emission. The NO_x emission from cement production was reduced 75 % since 1990. The reduced emission is a result of installation of

¹⁰ More than 65 % of sector 1A2f i.

SCR on all production units at the cement production plant in 2004-2007¹¹ and improved performance of the SCR units in recent years. A NO_x tax was introduced in 2010 (DMT, 2008).

The NMVOC emission has decreased 75 % since 1990. The decrease is a mainly result of decreased emission factor for combustion of wood in industrial boilers. The emission from gas engines has however increased considerably after 1995 due to the increased fuel consumption that is a result of the installation of a large number of industrial CHP plants (Figure 3.35). The NMVOC emission factor for gas engines is much higher than for boilers regardless of the fuel.

The CO emission in 2012 was 11 % lower than in 1990. The main source of emission is combustion in *Industry – Other*, primarily in wood and cement production. The CO emission from mineral wool production is included in the industry sector (2A7d).




Figure 3.2.29 Time series for fuel consumption, 1A2 Industry.

¹¹ To meet emission limit.




Figure 3.2.30 Time series for greenhouse gas emission, 1A2 Industry.




Figure 3.2.31 Time series for SO₂, NO_x, NMVOC and CO emission, 1A2 Industry.

1A2a Iron and steel

Iron and steel is a very small emission source category. Figure 3.2.32 shows the time series for fuel consumption and emissions.

Natural gas is the main fuel in the subsector.




Figure 3.2.32 Time series for 1A2a Iron and steel.

1A2b Non-ferrous metals

Non-ferrous metals is a very small emission source category. Figure 3.2.33 shows the time series for fuel consumption and emissions.

Natural gas is the main fuel in the subsector. The consumption of residual oil has decreased and the SO₂ emission follows this fuel consumption. The emissions of NO_x, NMVOC and CO follow the fuel consumption.




Figure 3.2.33 Time series for 1A2b Non-ferrous metals.

1A2c Chemicals

Chemicals is a minor emission source category. Figure 3.2.34 shows the time series for fuel consumption and emissions.

Natural gas is the main fuel in this subsector. The consumption of residual oil has decreased and the SO₂ emission follows this fuel consumption. The time series for CH₄, NMVOC and CO is related to consumption of natural gas in gas engines.




Figure 3.2.34 Time series for 1A2c Chemicals.

1A2d Pulp, paper and print

Pulp, paper and print is a minor emission source category. Figure 3.2.35 shows the time series for fuel consumption and emissions.

Natural gas and - since 2007 - also wood are the main fuels in the subsector. The increased consumption of wood in 2007 onwards is reflected in the CH₄, N₂O, NMVOC and CO emission time series.

The consumption of coal and residual oil has decreased and this is reflected in the SO₂ emission time series.




Figure 3.2.35 Time series for 1A2d Pulp, paper and print.

1A2e Food processing, beverages and tobacco

Food processing, beverages and tobacco is a considerable industrial subsector. Figure 3.2.36 shows the time series for fuel consumption and emissions.

Natural gas, residual oil and coal are the main fuels in the subsector. The consumption of coal and residual oil has decreased whereas the consumption of natural gas has increased. This is reflected in the SO₂ emission time series. The time series for CH₄ and NMVOC emission reflects the consumption of natural gas in gas engines.




Figure 3.2.36 Time series for 1A2e Food processing, beverages and tobacco.

1A2f Industry - other

Industry - other is a considerable industrial subsector. Figure 3.2.37 shows the time series for fuel consumption and emissions. The subsector includes cement production that is a major industrial emission source in Denmark.

Natural gas is the main fuels in the subsector in recent years. The consumption of coal and residual oil has decreased.

The NO_x time series is discussed above.




Figure 3.2.37 Time series for 1A2f Industry - other.


1A4 Other Sectors

The emission source category *1A4 Other Sectors* consists of the subcategories:


- 1A4a Commercial/Institutional plants.
- 1A4b Residential plants.
- 1A1c Agriculture/Forestry.

Figure 3.2.38-40 present time series for this emission source category. Residential plants is the dominant subcategory accounting for the largest part of all emissions. Time series are discussed below for each subcategory.


1A4 Other Sectors, subsectors




1A4 Other Sectors, fuel category




Gas engines, biogas (subsectors to Other Sectors)



Gas engines, natural gas (subsectors to Other Sectors)



Combustion of wood in Other Sectors



Combustion of straw in Other Sectors




Figure 3.2.38 Time series for fuel consumption, 1A4 Other Sectors.




Figure 3.2.39 Time series for greenhouse gas emission, 1A4 Other Sectors.




Figure 3.2.40 Time series for SO_2 , NO_x , NMVOC and CO emission, 1A4 Other Sectors.

1A4a Commercial and institutional plants

The subcategory *Commercial and institutional plants* consists of both stationary and mobile sources. In this chapter, only stationary sources are included. Figure 3.2.41 shows the time series for fuel consumption and emissions.

The subcategory *Commercial and institutional plants* has low fuel consumption and emissions compared to the other stationary combustion emission

source categories. Figure 3.2.35 shows the time series for fuel consumption and emissions.

The fuel consumption in commercial/institutional plants has decreased 41 % since 1990 and there has been a change of fuel type. The fuel consumption consists mainly of gas oil and natural gas. The consumption of gas oil has decreased whereas the consumption of natural gas has increased since 1990. The consumption of wood and biogas has also increased. The wood consumption in 2012 was 4.3 times the consumption in 1990.

The CO₂ emission has decreased 53 % since 1990. Both the decrease of fuel consumption and the change of fuels – from gas oil to natural gas – contribute to the decreased CO₂ emission.

The CH₄ emission in 2012 was 4.4 times the 1990 level. The increase is mainly a result of the increased emission from natural gas fuelled engines. The emissions from biogas fuelled engines and from combustion of wood also contribute to the increase. The time series for consumption of natural gas and biogas are shown in Figure 3.2.32.

The N₂O emission in 2012 was 4 % lower than in 1990. The fluctuations of the N₂O emission follow the fuel consumption.

The SO₂ emission has decreased 92 % since 1990. The decrease is a result of both the change of fuel from gas oil to natural gas and of the lower sulphur content in gas oil and in residual oil. The lower sulphur content (0.05 % for gas oil since 1995 and 0.7 % for residual oil since 1997) is a result of Danish tax laws (DMT, 1998). New boilers and abatement equipment was installed in a large wastewater treatment plant in 2002, but the efficiency of the abatement equipment was not as expected in the first months. Thus, an increased emission from this plant has caused the increased SO₂ emission in 2002.

The NO_x emission was 49 % lower in 2012 than in 1990. The decrease is mainly a result of the lower fuel consumption but also the change from gas oil to natural gas has contributed to the decrease. The emission from gas engines and wood combustion has increased.

The NMVOC emission in 2012 was 54 % higher than the 1990 emission level. The large increase is a result of the increased combustion of wood that is the main source of emission. The increased consumption of natural gas in gas engines (Figure 3.2.32) also contribute to the increased NMVOC emission.

The CO emission has decreased 25 % since 1990. The emission from wood and from natural gas fuelled engines and boilers have increased, whereas the emission from gas oil has decreased. This is a result of the change of fuels applied in the sector.




Figure 3.2.41 Time series for 1A4a Commercial/institutional.

1A4b Residential plants

The emission source category *Residential plants* consists of both stationary and mobile sources. In this chapter, only stationary sources are included. Figure 3.2.42 shows the time series for fuel consumption and emissions.

For residential plants, the total fuel consumption was 17 % lower in 2012 than in 1990. The large decrease from 2010 to 2011 is caused by higher temperature in the winter season of 2012. The consumption of gas oil has decreased since 1990 whereas the consumption of wood has increased considerably (3.6 times the 1990 level). The consumption of natural gas has also increased since 1990.

The CO₂ emission has decreased by 58 % since 1990. This decrease is mainly a result of the considerable change in fuels used from gas oil to wood and natural gas.

The CH₄ emission from residential plants has increased 43 % since 1990 due to the increased combustion of wood in residential plants, which is the main source of emission. The increased emission from gas engines also contributes to the increased emission.

The change of fuel from gas oil to wood has resulted in a 64 % increase of N₂O emission since 1990 due to a higher emission factor for wood than for gas oil.

The large decrease (85 %) of SO₂ emission from residential plants is mainly a result of a change of sulphur content in gas oil since 1995. The lower sulphur content (0.05 %) is a result of Danish tax laws (DMT, 1998). In addition, the consumption of gas oil has decreased and the consumption of natural gas that results in very low SO₂ emissions has increased.

The NO_x emission has increased by 15 % since 1990 due to the increased emission from wood combustion. The emission factor for wood is higher than for gas oil.

The emission of NMVOC has increased 15 % since 1990 as a result of the increased combustion of wood. The emission factor for wood has decreased since 2000, due to improved technology, but not as much as the increase in consumption of wood. The emission factor for wood and straw is higher than for liquid or gaseous fuels.

The CO emission has increased 16 % due to the increased use of wood that is the main source of emission. The emission factor for wood has decreased since 2000, due to improved technology, but not as much as the increase in consumption of wood. The emission from combustion of straw has decreased since 1990.




Figure 3.2.42 Time series for 1A4b Residential plants.

1A4c Agriculture/forestry

The emission source category *Agriculture/forestry* consists of both stationary and mobile sources. In this chapter, only stationary sources are included. Figure 3.2.43 shows the time series for fuel consumption and emissions.

For plants in agriculture/forestry, the fuel consumption has decreased 42 % since 1990. A considerable decrease in the fuel consumption has taken place since year 2000.

The type of fuel that has been applied has changed since 1990. In the years 1994-2004, the consumption of natural gas was high, but in recent years, the consumption decreased again. A large part of the natural gas consumption has been applied in gas engines (Figure 3.2.32). Most CHP plants in agriculture/forestry based on gas engines came in operation in 1995-1999. The decrease in later years is a result of the liberalisation of the electricity market.

The consumption of straw has decreased since 1990. The consumption of both residual oil and gas oil has increased after 1990 but has decreased again in recent years.

The CO₂ emission in 2012 was 57 % lower than in 1990. The CO₂ emission increased from 1990 to 1996 due to increased fuel consumption. Since 1996, the CO₂ emission has decreased in line with the decrease in fuel consumption.

The CH₄ emission in 2012 was 2 % higher than the emission in 1990. The emission follows the time series for natural gas combusted in gas engines (Figure 3.2.32). The emission from combustion of straw has decreased as a result of the decreasing consumption of straw in the sector.

The emission of N₂O has decreased by 36 % since 1990. The decrease is a result of the lower fuel consumption as well as the change of fuel. The decreasing consumption of straw contributes considerably to the decrease of emission.

The SO₂ emission was 68 % lower in 2012 than in 1990. The emission decreased mainly in the years 1996-2002. The main emission sources are coal, residual oil and straw.

The emission of NO_x was 50 % lower in 2012 than in 1990.

The emission of NMVOC has decreased 42 % since 1990. The major emission source is combustion of straw. The consumption of straw has decreased since 1990. The emission from gas engines has increased mainly due to increased fuel consumption.

The CO emission has decreased 72 % since 1990. The major emission source is combustion of straw. In addition to the decrease of straw consumption, the emission factor for straw has also decreased since 1990.




Figure 3.2.43 Time series for 1A4c Agriculture/Forestry.

3.2.5 Methodological issues

The Danish emission inventory is based on the CORINAIR (CORE INven-tory on AIR emissions) system, which is a European program for air emission inventories. CORINAIR includes methodology structure and software for inventories. The methodology is described in the EMEP/CORINAIR Emission Inventory Guidebook 2009 update, prepared by the UNECE/EMEP Task Force on Emissions Inventories and Projections (EEA, 2009). Emission data are stored in an Access database, from which data are transferred to the reporting formats.

In the Danish emission database all activity rates and emissions are defined in SNAP sector categories (Selected Nomenclature for Air Pollution) according the CORINAIR system. The emission inventories are prepared from a complete emission database based on the SNAP source categories. Aggregation to the source category codes used in CRF is based on a correspondence list enclosed in Annex 3A-1.

The emission inventory for stationary combustion is based on activity rates from the Danish energy statistics. General emission factors for various fuels, plants and sectors have been determined. Some large plants, such as power plants, are registered individually as large point sources and plant-specific emission data are used.

Tiers

The type of emission factor and the applied tier level for each emission source are shown in Table 3.2.13 below. The tier levels have been determined based on the IPCC Guidelines (IPCC, 1997).

The fuel consumption data for transformation are technology specific. For end-use of fuels, the disaggregation to specific technologies is less detailed. However, for residential wood combustion technology specific fuel consumption rates have been estimated.

Distinguishing between tier level 2 and tier 3 has been based on the emission factor. The tier level definitions have been interpreted as follows:

- Tier 1: The emission factor is an IPCC default tier 1 value.
- Tier 2: The emission factors are country specific and based on either a few emission measurements or IPCC tier 2 emission factors.
- Tier 3: Based on plant specific emission data or on a country specific emission factor based on a considerable number of plant specific emission measurements and detailed technology knowledge.

Table 3.2.13 gives an overview of the calculation methods and type of emission factor. The table also shows which of the source categories are key in any of the key source analysis¹².

¹² Key category according to the KCA tier 1 or tier 2 for Denmark (excluding Greenland and Faroe Islands), including LULUCF, level 1990/ level 2012/ trend.

Table 3.2.13 Methodology and type of emission factor.

		Tier	EMF ¹⁾	Key category ²⁾
Stationary Combustion, Coal	CO ₂	Tier 3 ¹³ (Tier 3) /Tier 1 ¹⁴	PS (CS) / D	Yes
Stationary Combustion, BKB	CO ₂	Tier 1	D	No
Stationary Combustion, Coke	CO ₂	Tier 1	D	No
Stationary Combustion, Fossil waste	CO ₂	Tier 3	CS	Yes
Stationary Combustion, Petroleum coke	CO ₂	Tier 3	PS/CS	Yes
Stationary Combustion, Residual oil	CO ₂	Tier 3 / Tier 3 / Tier 1	PS / CS / D ¹⁵	Yes
Stationary Combustion, Gas oil	CO ₂	Tier 2 / Tier 3	CR / PS	Yes
Stationary Combustion, Kerosene	CO ₂	Tier 1	D	Yes
Stationary Combustion, LPG	CO ₂	Tier 1	D	No
Stationary Combustion, Refinery gas	CO ₂	Tier 3	PS / CS	Yes
Stationary Combustion, Natural gas	CO ₂	Tier 3	CS / PS ¹⁶	Yes
Stationary Combustion, SOLID	CH ₄	Tier 2 / Tier 1	D(2) / D	No
Stationary Combustion, LIQUID	CH ₄	Tier 2 / Tier 2 / Tier 1	D(2) / CS / D	No
Stationary Combustion, GAS	CH ₄	Tier 2 / Tier 3	D(2) / CS	No
Natural gas fuelled engines, GAS	CH ₄	Tier 3	CS	Yes
Stationary Combustion, WASTE	CH ₄	Tier 2	CS	No
Stationary Combustion, BIOMASS	CH ₄	Tier 2 / Tier 1	D(2) / CS / D	Yes
Biogas fuelled engines, BIOMASS	CH ₄	Tier 3	CS	No
Stationary Combustion, SOLID	N ₂ O	Tier 2 / Tier 1	CS / D	Yes
Stationary Combustion, LIQUID	N ₂ O	Tier 2 / Tier 1	D(2) / D / CS	Yes
Stationary Combustion, GAS	N ₂ O	Tier 1 / Tier 2	D / CS / D(2)	Yes
Stationary Combustion, WASTE	N ₂ O	Tier 2	CS	No
Stationary Combustion, BIOMASS	N ₂ O	Tier 1 / Tier 2	D / CS / D(2)	Yes

¹⁾ D: IPCC tier 1, D(2): IPCC tier 2/3, CR: Corinair default, CS: Country specific, PS: Plant specific.

²⁾ KCA tier 1 or tier 2 for Denmark (excluding Greenland and Faroe Islands), including LULUCF, level 1990/level 2012/ trend

Large point sources

Large emission sources such as power plants, industrial plants and refineries are included as large point sources in the Danish emission database. Each point source may consist of more than one part, e.g. a power plant with several units. By registering the plants as point sources in the database, it is possible to use plant-specific emission factors.

In the inventory for the year 2012, 62 stationary combustion plants are specified as large point sources. These point sources include:

- Power plants and decentralised CHP plants (combined heat and power plants).
- Waste incineration plants.
- Large industrial combustion plants.
- Petroleum refining plants.

The criteria for selection of point sources consist of the following:

- All centralized power plants, including smaller units.
- All units with a capacity above 25 MW_e.
- All district heating plants with an installed effect of 50 MW_{th} or above and significant fuel consumption.
- All waste incineration plants obligated to report environmental data annually according to Danish law (DEPA, 2010a).

¹³ For 2006 onwards. Country specific emission factors and tier 2 have been applied for 1990-2005.

¹⁴ Tier 1 and IPCC default emission factor is only applied for 2 % of the coal consumption in 2012.

¹⁵ Residual oil not applied in source category 1A1a.

¹⁶ Off shore gas turbines and some power plants.

- Industrial plants,
 - with an installed effect of 50 MW_{th} or above and significant fuel consumption.
 - with a significant process related emission.

The fuel consumption of stationary combustion plants registered as large point sources in the 2012 inventory was 225 PJ. This corresponds to 51 % of the overall fuel consumption for stationary combustion.

A list of the large point sources for 2012 and the fuel consumption rates is provided in Annex 3A-5. The number of large point sources registered in the databases increased from 1990 to 2012.

The emissions from a point source are based either on plant specific emission data or, if plant specific data are not available, on fuel consumption data and the general Danish emission factors. Annex 3A-5 shows which of the emission data for large point sources are plant-specific and the corresponding share of the emission from stationary combustion.

The emission shares from point sources with plant specific data are shown in Table 3.2.14.

Table 3.2.14 Emission share, plant specific data.

Pollutant	Share from plant specific data
CO ₂	63 %
CH ₄	0 %
N ₂ O	0 %
SO ₂	52%
NO _x	43%
NM VOC	0.02%
CO	2.7%

CO₂ emission factors are plant specific for the major power plants, refineries, off shore gas turbines and for cement production. SO₂ and NO_x emissions from large point sources are often plant-specific based on emission measurements. Emissions of CO and NM VOC are also plant-specific for some plants. Plant-specific emission data are obtained from:

- CO₂ data reported under the EU Emission Trading Scheme (ETS).
- Annual environmental reports / environmental reporting available on the Danish EPA home page¹⁷
- Annual plant-specific reporting of SO₂ and NO_x from power plants >25MW_e prepared for the Danish Energy Agency and Energinet.dk.
- Emission data reported by DONG Energy and Vattenfall, the two major electricity suppliers.
- Emission data reported from industrial plants.

The EU ETS data are discussed in the chapter Emission factors.

Annual environmental reports for the plants include a considerable number of emission data sets. Emission data from annual environmental reports are, in general, based on emission measurements, but some emissions have potentially been calculated from general emission factors.

¹⁷ <http://www3.mst.dk/Miljoeoplysninger/PrtrPublicering/Index>

If plant-specific emission factors are not available, general area source emission factors are used.

Emissions of the greenhouse gases CH₄ and N₂O from the large point sources are all based on the area source emission factors.

Area sources

Fuels not combusted in large point sources are included as source category specific area sources in the emission database. Plants such as residential boilers, small district heating plants, small CHP plants and some industrial boilers are defined as area sources. Emissions from area sources are based on fuel consumption data and emission factors. Further information on emission factors is provided below in the chapter Emission factors (see page 157).

Activity rates, fuel consumption

The fuel consumption rates are based on the official Danish energy statistics prepared by the Danish Energy Agency (DEA). DCE aggregates fuel consumption rates to SNAP categories. Some fuel types in the official Danish energy statistics are added to obtain a less detailed fuel aggregation level cf. Annex 3A-3. The calorific values on which the energy statistics are based are also enclosed in Annex 3A-3. The correspondence list between the energy statistics and SNAP categories is enclosed in Annex 3A-9.

The fuel consumption of the CRF category *Manufacturing industries and construction* (corresponding to SNAP category 03) is disaggregated into industrial subsectors based on the DEA data set aggregated for the Eurostat reporting (DEA, 2013c).

The fuel consumption data flow is shown in Figure 3.2.44.




Figure 3.2.44 Fuel consumption data flow.

Both traded and non-traded fuels are included in the Danish energy statistics. Thus, for example, estimation of the annual consumption of non-traded wood is included.

Petroleum coke purchased abroad and combusted in Danish residential plants (border trade of 628 TJ in 2012) is not included in the Danish inventory. This is in agreement with the IPCC Guidelines (IPCC, 1997).

The fuel consumption data for large point sources refer to the EU Emission Trading Scheme (EU ETS) data for plants for which the CO₂ emission also refer to EU ETS, see page 157.

For all other large point sources, the fuel consumption refers to a DEA database (DEA, 2013b). The DEA compiles a database for the fuel consumption of each district heating and power-producing plant, based on data reported by plant operators. The consistency between EU ETS reporting and the DEA database (DEA, 2013b) is checked by the DEA and any discrepancies are corrected prior to the use in the emission inventory.

The fuel consumption of area sources is calculated as total fuel consumption in the energy statistics minus fuel consumption of large point sources.

The Danish national energy statistics includes three fuels used for non-energy purposes; bitumen, white spirit and lubricants. The total consumption for non-energy purposes is relatively low, e.g. 11.5 PJ in 2012. The use of white spirit is included in the inventory in *Solvent and other product use*. The emissions associated with the use of bitumen and lubricants are included in *Industrial Processes*. The non-energy use of fuels is included in the reference approach for Climate Convention reporting and appropriately corrected in line with the Revised 1996 IPCC Guidelines (IPCC, 1997).

In Denmark, all waste incineration are utilised for heat and power production. Thus, incineration of waste is included as stationary combustion in the source category *Fuel combustion* (subcategories 1A1, 1A2 and 1A4).

Fuel consumption data are presented in Chapter 3.2.2.

Town gas

Town gas has been included in the fuel category natural gas. The consumption of town gas in Denmark is very low, e.g. 0.6 PJ in 2012. In 1990, the town gas consumption was 1.6 PJ and the consumption has been steadily decreasing throughout the time series.

In Denmark, town gas is produced based on natural gas. The use of coal for town gas production ceased in the early 1980s.

An indicative composition of town gas according to the largest supplier of town gas in Denmark is shown in Table 3.2.15 (KE, 2014).

Table 3.2.15 Composition of town gas, 2012 (KE, 2014).

Component	Town gas, % (mol.)
Methane	43.9
Ethane	2.9
Propane	1.1
Butane	0.5
Carbon dioxide	0.4
Nitrogen	40.5
Oxygen	10.7

The lower heating value of the town gas currently used is 19.3 MJ per Nm³ and the CO₂ emission factor 56.1 kg per GJ. This is very close to the emission factor used for natural gas of 56.97 kg per GJ. According to the supplier, both the composition and heating value will change during the year. It has not been possible to obtain a yearly average.

In earlier years, the composition of town gas was somewhat different. Table 3.2.16 shows data for town gas composition in 2000-2005. These data are constructed with the input from Københavns Energi (KE) (Copenhagen Energy) and Danish Gas Technology Centre (DGC), (Jeppesen, 2008; Kristensen, 2007). The data refer to three measurements performed several years apart; the first in 2000 and the latest in 2005.

Table 3.2.16 Composition of town gas, information from the period 2000-2005.

Component	Town gas, % (mol.)
Methane	22.3-27.8
Ethane	1.2-1.8
Propane	0.5-0.9
Butane	0.13-0.2
Higher hydrocarbons	0-0.6
Carbon dioxide	8-11.6
Nitrogen	15.6-20.9
Oxygen	2.3-3.2
Hydrogen	35.4-40.5
Carbon monoxide	2.6-2.8

The lower calorific value has been between 15.6 and 17.8 MJ per Nm³. The CO₂ emission factors - derived from the few available measurements - are in the range of 52-57 kg per GJ.

The Danish approach includes town gas as part of the fuel category natural gas and thus indirectly assumes the same CO₂ emission factor. This is a conservative approach ensuring that the CO₂ emissions are not underestimated.

Due to the scarce data available and the very low consumption of town gas compared to consumption of natural gas (< 0.5 %), the methodology will be applied unchanged in future inventories.

Waste

All waste incineration in Denmark is utilised for heat and/or power production and thus included in the energy sector. The waste incinerated in Denmark for energy production consists of the waste fractions shown in

Figure 3.2.45. In 2009¹⁸, 3 % of the incinerated waste was hazardous waste¹⁹.




Figure 3.2.45 Waste fractions (weight) for incinerated waste in 2009 and the corresponding time series 1994-2009 (ISAG, 2012).

In connection to the project estimating an improved CO₂ emission factor for waste (Astrup et al., 2012), the fossil energy fraction was calculated. The fossil fraction was not measured or estimated as part of the project, but the flue gas measurements combined with data from Fellner & Rechberger (2010) indicated a fossil energy part of 45 %. The energy statistics also applies this fraction in the national statistics.

Biogas

Biogas includes landfill gas, sludge gas and manure/organic waste gas²⁰. The Danish energy statistics specifies production and consumption of each of the biogas types. In 2012, 75 % of the applied biogas was based on manure /organic waste.

¹⁸ Currently, data are only available for 1994-2009.

¹⁹ In 2001 onwards, health-care risk waste is included in hazardous waste in the ISAG database.

²⁰ Based on manure with addition of other organic waste.




Figure 3.2.46 Biogas types 2012 and the corresponding time series 1990-2012 (DEA, 2013a).

Emission factors

For each fuel and SNAP category (sector and e.g. type of plant), a set of general area source emission factors has been determined. The emission factors are either nationally referenced or based on the international guidebooks: EMEP/EEA Guidebook (EEA, 2013)²¹ and IPCC Guidelines (IPCC, 1997).

An overview of the type of CO₂ emission factor is shown in Table 3.2.13. A complete list, of emission factors including time series and references, is provided in Annex 3A-4.

EU ETS data for CO₂

The CO₂ emission factors for some large power plants and for combustion in the cement industry and refineries are plant specific and based on the reporting to the EU Emission Trading Scheme (EU ETS). In addition, emission factors for offshore gas turbines and refinery gas is based on EU ETS data²². The EU ETS data have been applied for the years 2006 - 2012.

The EU ETS data are also applied for other source categories and are further discussed in Chapter 1.4.10.

ETS data, methodology, criteria for implementation and QA/QC

The Danish emission inventory for stationary combustion only includes data from plants using higher tier methods as defined in the EU decision (EU Commission, 2007), where the specific methods for determining carbon contents, oxidation factor and calorific value are specified. The EU decision includes rules for measuring, reporting and verification.

²¹ And former editions of the EMEP/EEA and EMEP/Corinair Guidebook.

²² See page 134 and 134.

For each of the plants included individually in the Danish inventory all applied methodologies are specified in individual monitoring plans that are approved by Danish authorities (DEA) prior to the reporting of the emissions. The plants/fuels included individually in the Danish inventory all apply the Tier 3 methodology for calculating the CO₂ emission factor. This selection criteria results in a dataset for which the emission factor values are based on fuel quality measurements²³, not default values from the Danish UNFCCC reporting. All fuel analyses are performed according to ISO 17025.

The power plants/fuels selected based on emission factor methodology apply the tiers for activity data, net calorific value (NCV), emission factor and oxidation factor listed below.

Coal

The CO₂ emission factor for coal is based on analysis of C content of the coal (g C per kg) and coal weight measurements. However, NCV values are also measured according to high tier methods in spite of the fact that this value is not input data for the calculation of total CO₂ emission.

- Fuel flow: Tier 4 methodology (± 1.5 %). For coal, the activity data (weight) is based on measurements on belt conveyor scale. The uncertainty is below the required ± 1.5 %.
- NCV: Tier 3 methodology. Data are based on measurements according to ISO 13909 / ISO 18283 (sampling) and ISO 1928 (NCV). The uncertainty for data is below ± 0.5 %.
- Emission factor: The emission factor is C-content of the coal. Tier 3 methodology (± 0.5 %) is applied and the measurements are performed according to ISO 13909 (sampling) and ISO/TS 12902 (C-content).
- Oxidation factor: Based on Tier 3 methodology except for one plant that applies Tier 1 methodology²⁴. The Tier 3 methodology is based on measurements of C-content in bottom ash and fly ash according to ISO/TS 12902 or on burning loss measurements according to ISO 1171. The uncertainty has been estimated to 0.5 %. For Tier 1 the oxidation factor is assumed to be 1.

Residual oil

- Fuel flow: Tier 4 methodology (± 1.5 %) for most plants. However, a few of the included plants apply Tier 3 methodology (± 2.5 %).
- NCV: Tier 3 methodology. Data are based on sampling according to API Manual of Petroleum Measurement Standards / ASTM D 270 and fuel analysis (NCV) according to ASTM D 240 / ISO 1928 / data stated by the fuel supplier.
- Emission factor: Tier 3 methodology according to API Manual of Petroleum Measurement Standards / ASTM D 4057 (sampling) and ISO 12902 / ASTM D 5291 (C-content).
- Oxidation factor: Based on Tier 2 or Tier 3 methodology, both resulting in the oxidation factor 1 with an uncertainty of 0.8 %.

²³ Applying specific methods defined in the EU decision

²⁴ In addition, DCE have assumed the oxidation factor to be 1 for a plant for which the stated oxidation factor was rejected in the QC work.

For coal and residual oil fuel analyses are required for each 20,000 tonnes or at least six times each year. The fuel analyses are performed by accredited laboratories²⁵.

QC of EU ETS data

DCE performs QC checks on the reported emission data, see Chapter 1.4.10. Based on the QC checking DCE excluded the oxidation factor for coal for one stationary combustion plant for 2012.

EU ETS data presentation

The EU ETS data include plant specific emission factors for coal, residual oil, gas oil, natural gas, refinery gas, petroleum coke and fossil waste. The EU ETS data account for 63 % of the CO₂ emission from stationary combustion.

EU ETS data for coal

EU ETS data for 2012 were available from 15 coal fired plants. The plant specific information accounts for 98 % of the Danish coal consumption and 43 % of the total (fossil) CO₂ emission from stationary combustion plants. The average CO₂ emission factor for coal for these 15 units was 94.25 kg per GJ (Table 3.2.17). The plants all apply bituminous coal.

Data from two plants have been excluded in the calculation of the implied emission factor for coal because the coal applied in those plants is not representative for the coal applied in Denmark. However, both data sets have been included in the inventories and in Table 3.2.17 below. The estimated IEF for CO₂ is however 94.25 kg/GJ in both cases.

Table 3.2.17 EU ETS data for 15 coal fired plants, 2012.

	Average	Min	Max
Heating value, GJ per tonne ²⁶	24.11	23.5	26.6
CO ₂ implied emission factor, kg per GJ ¹⁾	94.25	91.4	96.4
Oxidation factor	0.996	0.988	1.000

¹⁾ Including oxidation factor.

Table 3.2.18 CO₂ implied emission factor time series for coal fired plants based on EU ETS data.

Year	CO ₂ implied emission factor, kg per GJ ¹⁾
2006	94.4
2007	94.3
2008	94.0
2009	93.6
2010	93.6
2011	94.7
2012	94.3

¹⁾ Including oxidation factor.

EU ETS data for residual oil

EU ETS data for 2012 based on higher tier methodologies were available from 13 plants combusting residual oil. Aggregated data and time series are shown in Table 3.2.19 and Table 3.2.20. The EU ETS data accounts for 53 % of the residual oil consumption in stationary combustion.

²⁵ EN ISO 17025.

²⁶ One data set has been excluded as part of the QC work.

Table 3.2.19 EU ETS data for 13 plants combusting residual oil.

	Average	Min	Max
Heating value, GJ per tonne	40.5	40.2	40.7
CO ₂ implied emission factor, kg per GJ	79.21	77.30	80.48
Oxidation factor	1.000	1.000	1.000

Table 3.2.20 CO₂ implied emission factor time series for residual oil fired power plant units based on EU ETS data.

Year	CO ₂ implied emission factor, kg per GJ ¹⁾
2006	78.2
2007	78.1
2008	78.5
2009	78.9
2010	79.2
2011	79.25
2012	79.21

¹⁾ Including oxidation factor

EU ETS data for gas oil combusted in power plants or refineries

EU ETS data for 2012 based on higher tier methodologies were included from 4 plants combusting gas oil. Aggregated data and time series are shown in Table 3.2.21 and Table 3.2.22. The EU ETS data accounts for 10 % of the gas oil consumption in stationary combustion.

Table 3.2.21 EU ETS data for gas oil applied in power plants/refineries.

	Average	Min	Max
Heating value, GJ per tonne	35.9	34.1	35.9
CO ₂ implied emission factor, kg per GJ	73.92	73.73	75.15
Oxidation factor	1.000	1.000	1.000

Table 3.2.22 CO₂ implied emission factor time series for gas oil based on EU ETS data.

Year	CO ₂ implied emission factor, kg per GJ ¹⁾
2006	75.1
2007	74.9
2008	73.7
2009	75.1
2010	74.8
2011	74.7
2012	73.9

¹⁾ Including oxidation factor.

EU ETS data for petroleum coke and waste applied in industrial plants

Plant specific CO₂ emission factors from EU ETS have also been applied for the some industrial plants including cement industry, sugar production, glass wood production, lime production, and vegetable oil production. The implemented EU ETS data set also includes CO₂ emission factors for petroleum coke and waste applied in industrial plants.

EU ETS data for natural gas applied in offshore gas turbines

EU ETS data have been applied to estimate an average CO₂ emission factor for natural gas applied in offshore gas turbines, see page 166.

EU ETS data for refinery gas

EU ETS data are also applied for the two refineries in Denmark. The emission factor for refinery gas is based on EU ETS data, see page 165.

CO₂, other emission factors

The CO₂ emission factors that are not included in EU ETS data or that are included but based on lower tier methodologies are not plant specific in the Danish inventory. The emission factors that are not plant specific accounts for 37 % of the fossil CO₂ emission.

The CO₂ emission factors applied for 2012 are presented in Table 3.2.23. Time series have been estimated for:

- Coal applied for production of electricity and district heating
- Residual oil applied for production of electricity and district heating
- Refinery gas
- Natural gas applied in off shore gas turbines
- Natural gas, other
- Industrial waste, biomass part

For all other fuels, the same emission factor has been applied for 1990-2012.

In the reporting to the UNFCCC, the CO₂ emission is aggregated to five fuel types: Solid fuels, Liquid fuels, Gaseous fuels, Biomass and Other fuels. The correspondence list between the DCE fuel categories and the IPCC fuel categories is also provided in Table 3.2.23.

Only emissions from fossil fuels are included in the total national CO₂ emission. The biomass emission factors are also included in the table, because emissions from biomass are reported to the UNFCCC as a memo item.

The CO₂ emission from incineration of waste (37 + 75.1 kg per GJ) is divided into two parts: The emission from combustion of the fossil content of the waste, which is included in the national total, and the emission from combustion of the rest of the waste – the biomass part, which is reported as a memo item. In the IPCC reporting, the fuel consumption and emissions from the fossil content of the waste is reported in the fuel category, *Other fuels*.

Table 3.2.23 CO₂ emission factors, 2012.

Fuel	Emission factor kg per GJ		Reference type	IPCC fuel category
	Bio-mass	Fossil fuel		
Coal, source category 1A1a Public electricity and heat production	94.25 ¹⁾		Country specific	Solid
Coal, Other source categories	94.6 ³⁾		IPCC (1997)	Solid
Brown coal briquettes	94.6		IPCC (1997)	Solid
Coke oven coke	108		IPCC (1997)	Solid
Other solid fossil fuels ⁶⁾	108 ¹⁾		IPCC (1997)	Solid
Fly ash fossil (from coal)	93.6		Country specific	Solid
Petroleum coke	93 ³⁾		Country specific	Liquid
Residual oil, source category 1A1a Public electricity and heat production	79.21 ¹⁾		Country specific	Liquid
Residual oil, other source categories	77.4 ³⁾		IPCC (1997)	Liquid
Gas oil	74 ¹⁾		EEA (2007)	Liquid
Kerosene	71.9		IPCC (1997)	Liquid
Orimulsion	80 ²⁾		Country specific	Liquid
LPG	63.1		IPCC (1997)	Liquid
Refinery gas	58.099		Country specific	Liquid
Natural gas, off shore gas turbines	57.423		Country specific	Gas
Natural gas, other	57.03		Country specific	Gas
Waste	75.1 ^{3/4)}	+ 37 ^{3/4)}	Country specific	Biomass and Other fuels
Straw	110		IPCC (1997)	Biomass
Wood	110		IPCC (1997)	Biomass
Bio oil	74		Country specific	Biomass
Biogas	83.6		Country specific	Biomass
Biomass producer gas	142.9 ⁵⁾		Country specific	Biomass

1) Plant specific data from EU ETS incorporated for individual plants.

2) Not applied in 2012. Orimulsion was applied in Denmark in 1995 – 2004.

3) Plant specific data from EU ETS incorporated for cement industry and sugar, lime and glass wool production.

4) The emission factor for waste is (37+75.1) kg CO₂ per GJ waste. The fuel consumption and the CO₂ emission have been disaggregated to the two IPCC fuel categories *Biomass* and *Other fuels* in CRF. The IEF²⁷ for CO₂, Other fuels is 82.22 kg CO₂ per GJ fossil waste.

5) Includes a high content of CO₂ in the gas.

6) Anodic carbon.

Coal

As mentioned above²⁸, EU ETS data have been utilised for the years 2006 - 2012 in the emission inventory. The emission factor for coal is the implied emission factor for plants that report EU ETS data that are based on fuel analysis. Data for industrial plants have been included. Data from two plants have been excluded in the calculation of the implied emission factor for coal because the coal applied in those plants is not representative for the coal applied in Denmark. However, it turns out that the consumption in these two plants is sufficiently low to not influence the resulting implied emission factor. In 2012, the implied emission factor (including oxidation factor) was 94.25 kg per GJ. The implied emission factor values were between 92.81 and 96.36 kg per GJ.

In 2012, only 2 % of the CO₂ emission from coal consumption was based on the emission factor, whereas 98 % of the coal consumption was covered by EU ETS data. All coal applied in Denmark is bituminous coal (DEA, 2013c).

²⁷ Not including cement production.

²⁸ EU ETS data for CO₂ on page 60.

The emission factors for coal combustion in *Public electricity and heat production* in the years 2006-2012 refer to the implied emission factors of the EU ETS data estimated for each year. For the years 1990-2005, the emission factor for coal in *Public electricity and heat production* refer to the average IEF for 2006-2009.

Time series for net calorific value (NCV) of coal are available in the Danish energy statistics. NCV for *Electricity plant coal* fluctuates in the interval 24.23-25.8 GJ per tonne.

The correlation between NCV and CO₂ IEF (including the oxidation factor) in the EU ETS data (2006-2009) have been analysed and the results are shown in Annex 3A-10. However, a significant correlation between NCV and IEF have not been found in the dataset and thus an emission factor time series based on the NCV time series was not relevant. In addition, the correlation of NCV and CO₂ emission factors has been analysed. This analysis is also shown in Annex 3A-10. As expected, the correlation was better in this dataset, but still insufficient for estimating a time series for the CO₂ emission factor based on the NCV time series.

As mentioned above all coal applied in Denmark is bituminous coal and within the range of coal qualities applied in the plants reporting data to EU ETS a correlation could not be documented.

For other sectors apart from 1A1a, the applied emission factor 94.6 kg per GJ refers to IPCC Guidelines (IPCC, 1997). This emission factor has been applied for all years.

Time series for the CO₂ emission factor are shown in Table 3.2.24.

Table 3.2.24 CO₂ emission factors for coal, time series.

Year	1A1a Public electricity and heat production kg per GJ	Other source categories kg per GJ
1990-2005	94.0	94.6
2006	94.4	94.6
2007	94.3	94.6
2008	94.0	94.6
2009	93.6	94.6
2010	93.6	94.6
2011	93.73	94.6
2012	93.25	94.6

Brown coal briquettes

The emission factor for brown coal briquettes, 94.6 kg per GJ, is based on a default value from the IPCC Guidelines (IPCC, 1997) assuming full oxidation. The default value in the IPCC Guidelines is 25.8 t C per TJ, corresponding to $25.8 \cdot (12+2 \cdot 16)/12 = 94.6$ kg CO₂ per GJ assuming full oxidation. The same emission factor has been applied for 1990-2012.

Coke oven coke

The emission factor for coke oven coke, 108 kg per GJ, is based on a default value from the IPCC Guidelines (IPCC, 1997) assuming full oxidation. The default value in the IPCC guidelines is 29.5 t C per TJ, corresponding to $29.5 \cdot (12+2 \cdot 16)/12 = 108$ kg CO₂ per GJ assuming full oxidation. The same emission factor has been applied for 1990-2012.

Other solid fossil fuels (Anodic carbon)

Anodic carbon has been applied in Denmark in 2009-2012 in two mineral wool production units. EU ETS data are available for both plants and thus the area source emission factor (108 kg/GJ) have not been applied.

Fly ash fossil (from coal)

Fly ash from coal combustion is applied in some power plants. The emission factor 93.6 kg/GJ have been applied. This is the emission factor for coal consumption in power plants in 2009-2010. The emission factor for 1990-2005 will be applied in future inventories (94 kg/GJ).

The emission factor have however not been applied due to the fact that plant specific data are available from the EU ETS dataset.

Petroleum coke

The emission factor for petroleum coke has been recalculated in this inventory. The improved emission factor 93 kg per GJ is based on EU ETS data for 2006-2010. The data includes one power plant and the cement production plant.

Plant specific EU ETS data have been utilised for the cement production for the years 2006 - 2012. This consumption represents more than 98 % of the consumption of petroleum coke in Denmark.

Residual oil

The emission factor for residual oil applied in public electricity and heat production is based on EU ETS data.

As mentioned above²⁹ EU ETS data have been utilised for the 2006 - 2012 emission inventories. In 2012, the implied emission factor (including oxidation factor) for the power plants and refineries combusting residual oil was 79.21 kg per GJ. The implied emission factor values were between 77.30 and 80.48 kg per GJ.

In 2012, 47 % of the CO₂ emission from residual oil consumption was based on the emission factor, whereas 53 % of the residual oil consumption was covered by EU ETS data³⁰.

The emission factors for residual oil combustion in *Public electricity and heat production* in the years 2006-2012 refer to the implied emission factors of the EU ETS data estimated for each year. For the years 1990-2005, the emission factor for residual oil in *Public electricity and heat production* refer to the average IEF for 2006-2009.

For other source categories apart from 1A1a, the applied emission factor 77.4 kg per GJ refers to the IPCC Guidelines (IPCC, 1997). This emission factor has been applied for all years.

Time series for the CO₂ emission factor are shown in Table 3.2.25.

²⁹ EU ETS data for CO₂ on page 61

³⁰ Including EU ETS data for cement production.

Table 3.2.25 CO₂ emission factors for residual oil, time series.

Year	Source category 1A1a Public electricity and heat production kg per GJ	Other source categories kg per GJ
1990-2005	78.4	77.4
2006	78.2	77.4
2007	78.1	77.4
2008	78.5	77.4
2009	78.9	77.4
2010	79.2	77.4
2011	79.25	77.4
2012	79.21	77.4

Gas oil

The emission factor for gas oil, 74 kg per GJ, refers to EEA (2007). The emission factor is consistent with the IPCC default emission factor for gas oil (74.1 kg per GJ assuming full oxidation). The CO₂ emission factor has been confirmed by the two major power plant operators in 1996 (Christiansen, 1996 and Andersen, 1996). The same emission factor has been applied for 1990-2012.

Plant specific EU ETS data have been utilised for a few plants in the 2006 - 2012 emission inventories. In 2012, the implied emission factor for the power plants using gas oil was 73.9 kg per GJ. The EU ETS CO₂ emission factors were in the interval 73.7 - 75.1 kg per GJ. In 2012, 10 % of the CO₂ emission from gas oil consumption was based on EU ETS data.

Kerosene

The emission factor for kerosene, 71.9 kg per GJ, refers to IPCC Guidelines (IPCC, 1997). The same emission factor has been applied for 1990-2012. CO₂ emission from combustion of kerosene is a key category and thus a country specific emission factor should be implemented in future inventories.

Orimulsion

The emission factor for orimulsion, 80 kg per GJ, refers to the Danish Energy Agency (DEA, 2013a). The IPCC default emission factor is almost the same: 80.7 kg per GJ assuming full oxidation. The CO₂ emission factor has been confirmed by the only major power plant operator using orimulsion (Andersen, 1996). The same emission factor has been applied for all years. Orimulsion was used in Denmark in 1995-2004.

LPG

The emission factor for LPG, 63.1 kg per GJ, refers to IPCC Guidelines (IPCC, 1997). The same emission factor has been applied for 1990-2012.

Refinery gas

The emission factor applied for refinery gas refers to EU ETS data for the two refineries in operation in Denmark. Since 2006, implied emission factors for Denmark have been estimated annually based on the EU ETS data. The average implied emission factor (57.6 kg per GJ) for 2006-2009 have been applied for the years 1990-2005. This emission factor is consistent to the emission factor stated in the 2006 IPCC Guidelines (IPCC, 2006). The time series is shown in Table 3.2.26.

Table 3.2.26 CO₂ emission factors for refinery gas, time series.

Year	CO ₂ emission factor, kg per GJ
1990-2005	57.6
2006	57.812
2007	57.848
2008	57.948
2009	56.814
2010	57.134
2011	57.881
2012	58.099

Natural gas, offshore gas turbines

EU ETS data for the fuel consumption and CO₂ emission for offshore gas turbines are available for the years 2006-2012. Based on data for each oil-field implied emission factors have been estimated for 2006-2012. The average value for 2006-2009 has been applied for the years 1990-2005. The time series is shown in Table 3.2.27.

Table 3.2.27 CO₂ emission factors for offshore gas turbines, time series.

Year	CO ₂ emission factor, kg per GJ
1990-2005	57.469
2006	57.879
2007	57.784
2008	56.959
2009	57.254
2010	57.314
2011	57.379
2012	57.423

Natural gas, other source categories

The emission factor for natural gas is estimated by the Danish gas transmission company, Energinet.dk³¹. The calculation is based on gas analysis carried out daily by Energinet.dk at Egtved.

In 2012, the natural gas import was 33 PJ, the natural gas export 112 PJ and a consumption that added up to 145 PJ. Before 2010, only natural gas from the Danish gas fields was utilised in Denmark. If the import of natural gas increases further, the methodology for estimating the CO₂ emission factor might be revised based on an on-going dialog with the Danish Energy Agency and Energinet.dk. However, Energinet.dk have stated that the difference between the emission factor for 2011 based on measurements at Egtved and the average value at Froeslev very close to the border differed less than 0.3 % for 2011 (Bruun, 2012).

Energinet.dk and the Danish Gas Technology Centre have calculated emission factors for 2000-2012. The emission factor applied for 1990-1999 refers to Fenhann & Kilde (1994). This emission factor was confirmed by the two major power plant operators in 1996 (Christiansen, 1996 and Andersen, 1996). The time series for the CO₂ emission factor is provided in Table 3.2.28.

³¹ Former Gastra and before that part of DONG. Historical data refer to these companies.

Table 3.2.28 CO₂ emission factor time series for natural gas.

Year	CO ₂ emission factor, kg per GJ
1990-1999	56.9
2000	57.1
2001	57.25
2002	57.28
2003	57.19
2004	57.12
2005	56.96
2006	56.78
2007	56.78
2008	56.77
2009	56.69
2010	56.74
2011	56.97
2012	57.03

Waste

The CO₂ emission from incineration of waste is divided into two parts: The emission from combustion of the fossil content of the waste, which is included in the national total, and the emission from combustion of the rest of the waste – the biomass part, which is reported as a memo item.

The CO₂ emission factor is based on the project, *Biogenic carbon in Danish combustible waste* that included emission measurements from five Danish waste incineration plants (Astrup et al., 2012). The average fossil emission factors for waste have estimated to be 37 kg/GJ waste and the interval for the five plants was 25 – 51 kg/GJ. The five plants represent 44 % of the incinerated waste in 2010. The emission factor 37 kg/ GJ waste corresponds to 82.22 kg/GJ fossil waste.

The total CO₂ emission factor for waste refers to a Danish study (Jørgensen & Johansen, 2003). Based on emission measurements on five waste incineration plants the total CO₂ emission factor for waste incineration has been determined to 112.1 kg per GJ. Thus, the biomass emission factor has been determined to 75.1 kg/GJ waste.

Plant specific EU ETS data have been utilised for cement production in the 2006 - 2012 emission inventories.

Wood

The emission factor for wood, 110 kg per GJ, refers IPCC (1997). The same emission factor has been applied for 1990-2012.

Straw

The emission factor for wood, 110 kg per GJ, refers IPCC (1997). The same emission factor has been applied for 1990-2012.

Bio oil

The emission factor is assumed to be the same as for gas oil – 74 kg per GJ. The consumption of bio oil is below 2 PJ.

Biogas

In Denmark, 3 different types of biogas are applied: Manure/organic waste based biogas, landfill based biogas and wastewater treatment biogas (sludge gas). Manure / organic waste based biogas represent 75 % of the consumption, see page 156.

The emission factor for biogas, 83.6 kg per GJ, is based on a biogas with 65 % (vol.) CH₄ and 35 % (vol.) CO₂. Danish Gas Technology Centre has stated that this is a typical manure-based biogas as utilised in stationary combustion plants (Kristensen, 2001). The same emission factor has been applied for 1990-2012.

Biomass producer gas

Biomass producer gas applied in Denmark is based on wood. The gas composition is known for three different plants and the applied emission factor have been estimated by Danish Gas Technology Centre (Kristensen, 2010) based on the gas composition measured on the plant that with the highest consumption.

The consumption of biomass producer gas is below 0.4 PJ for all years.

CH₄

The CH₄ emission factors applied for 2012 are presented in Table 3.2.29. In general, the same emission factors have been applied for 1990-2012. However, time series have been estimated for both natural gas fuelled engines and biogas fuelled engines, residential wood combustion, natural gas fuelled gas turbines³² and waste incineration plants³².

Emission factors for CHP plants < 25 MW_e refer to emission measurements carried out on Danish plants (Nielsen et al., 2010; Nielsen & Illerup, 2003; Nielsen et al., 2008). The emission factors for residential wood combustion are based on technology dependent data.

Emission factors that are not nationally referenced all refer to the IPCC Guidelines (IPCC, 1997).

Gas engines combusting natural gas or biogas account for more than half the CH₄ emission from stationary combustion plants. The relatively high emission factor for gas engines is well-documented and further discussed below.

³² A minor emission source.

Table 3.2.29 CH₄ emission factors, 2012.

Fuel group	Fuel	CRF source category	CRF source category	SNAP	Emission factor, g pr GJ	Reference
SOLID	COAL	1A1a	Public electricity and heat production	01010102	0.9	IPCC (1997), Tier 2, Table 1-15, Utility Boiler, Pulverised Bituminous Combustion, Wet bottom.
		1A2 a-f	Industry	03	10	IPCC (1997), Tier 2, Table 1-19, Commercial coal boilers.
		1A4b i	Residential	0202	300	IPCC (1997), Tier 1, Table 1-7, Residential, coal.
		1A4c i	Agriculture/ Forestry	0203	10	IPCC (1997), Tier 2, Table 1-19, Commercial coal boilers. ¹⁾
	BROWN COAL BRI.	1A4b i	Residential	0202	300	IPCC (1997), Tier 1, Table 1-7, Residential, coal.
	COKE OVEN COKE	1A2 A-f	Industry	03	10	IPCC (1997), Tier 2, Table 1-19, Commercial coal boilers.
		1A4b i	Residential	0202	300	IPCC (1997), Tier 1, Table 1-7, Residential, coal.
	ANODIC CARBON	1A2a-f	Industry	03	10	IPCC (1997), Tier 2, Table 1-19, Commercial coal boilers.
	FOSSIL FLY ASH	1A1a	Public electricity and heat production	0101	0.9	IPCC (1997), Tier 2, Table 1-15, Utility Boiler, Pulverised Bituminous Combustion, Wet bottom.
	LIQUID PETROLEUM COKE	1A2a-f	Industry	03	2	IPCC (1997), Tier 1, Table 1-7, Industry, oil.
	RESIDUAL OIL	1A1a	Public electricity and heat production	010101	0.9	IPCC (1997), Tier 2, Table 1-15, Utility Boiler, Residual fuel oil.
				010102	1.3	Nielsen et al. (2010)
				010103		
				010104	3	IPCC (1997), Tier 1, Table 1-7, Energy industries, oil.
				010203	0.9	IPCC (1997), Tier 2, Table 1-15, Utility Boiler, Residual fuel oil.
		1A1b	Petroleum refining	010306	3	IPCC (1997), Tier 1, Table 1-7, Energy industries, oil.
		1A2 a-f	Industry	03	1.3	Nielsen et al. (2010)
				En-gines	4	IPCC (1997), Tier 2, Table 1-15, Utility, Large diesel engines
		1A4a	Commercial/ Institutional	0201	1.4	IPCC (1997), Tier 2, Table 1-19, Commercial, residual fuel oil ¹⁾ .
		1A4b	Residential	0202	1.4	IPCC (1997), Tier 2, Table 1-19, Commercial, residual fuel oil ¹⁾ .
		1A4c	Agriculture/ Forestry	0203	1.4	IPCC (1997), Tier 2, Table 1-19, Commercial, residual fuel oil ¹⁾ .
	GAS OIL	1A1a	Public electricity and heat production	010101	0.9	IPCC (1997), Tier 2, Table 1-15, Utility Boiler, distillate fuel oil.
				010102		
				010103		
				010104	3	IPCC (1997), Tier 1, Table 1-7, Energy industries, oil.
				010105	24	Nielsen et al. (2010)
				010202	0.9	IPCC (1997), Tier 2, Table 1-15, Utility Boiler, distillate fuel oil.
				010203		
		1A1b	Petroleum refining	010306	3	IPCC (1997), Tier 1, Table 1-7, Energy industries, oil.
		1A1c	Other energy industries	010504	3	IPCC (1997), Tier 1, Table 1-7, Energy industries, oil
		1A2 a-f	Industry	03	0.2	IPCC (1997), Tier 2, Table 1-16, Industry, distillate fuel oil.
				Tur-bines	2	IPCC (1997), Tier 1, Table 1-7, Industry, oil.
				En-gines	24	Nielsen et al. (2010)
		1A4a	Commercial/ Institutional	0201	0.7	IPCC (1997), Tier 2, Table 1-19, Commercial, distillate fuel oil.
				020105	24	Nielsen et al. (2010)
				0202	0.7	IPCC (1997), Tier 2, Table 1-18,

Fuel group	Fuel	CRF source category	CRF source category	SNAP	Emission factor, g pr GJ	Reference
		1A4c	Agriculture/ Forestry	0203	0.7	Residential, distillate fuel oil. IPCC (1997), Tier 2, Table 1-19, Commercial, distillate fuel oil.
	KEROSENE	1A2 a-f	Industry	all	0.2	IPCC (1997), Tier 2, Table 1-16, Industry, distillate fuel oil.
		1A4a	Commercial/ Institutional	0201	0.7	IPCC (1997), Tier 2, Table 1-19, Commercial, distillate fuel oil.
		1A4b i	Residential	0202	0.7	IPCC (1997), Tier 2, Table 1-18, Residential, distillate fuel oil.
		1A4c i	Agriculture/ Forestry	0203	0.7	IPCC (1997), Tier 2, Table 1-19, Commercial, distillate fuel oil ¹⁾ .
	LPG	1A1a	Public electricity and heat production	0101 0102	3	IPCC (1997), Tier 1, Table 1-7, Energy Industries, oil.
		1A1b	Petroleum refining	0103	3	IPCC (1997), Tier 1, Table 1-7, Energy Industries, oil.
		1A2 a-f	Industry	03	2	IPCC (1997), Tier 1, Table 1-7, Industry, oil
		1A4a	Commercial/ Institutional	0201	10	IPCC (1997), Tier 1, Table 1-7, Commercial, oil.
		1A4b i	Residential	0202	1.1	IPCC (1997), Tier 2, Table 1-18, Residential propane/butane furnaces.
		1A4c i	Agriculture/ Forestry	0203	10	IPCC (1997), Tier 1, Table 1-7, Agriculture, oil.
	REFINERY GAS	1A1b	Petroleum refining	010304	1.7	Assumed equal to natural gas fuelled gas turbines. Nielsen et al. (2010)
				010306	1	Assumed equal to natural gas fuelled plants. IPCC (1997), Tier 1, Table 1-7, Natural gas
GAS	NATURAL GAS	1A1a	Public electricity and heat production	010101 010102 010103	0.1	IPCC (1997), Tier 2, Table 1-15, Utility Boiler, natural gas.
				010104	1.7	Nielsen et al. (2010)
				010105	481	Nielsen et al. (2010)
				010202 010203	0.1	IPCC (1997), Tier 2, Table 1-15, Utility Boiler, natural gas.
		1A1c	Other energy industries	010504	1.7	Nielsen et al. (2010)
		1A2 a-f	Industry	Other	1.4	IPCC (1997), Tier 2, Table 1-16, Industry, natural gas boilers.
				Gas turbines	1.7	Nielsen et al. (2010)
				En-gines	481	Nielsen et al. (2010)
		1A4a	Commercial/ Institutional	0201	1.2	IPCC (1997), Tier 2, Table 1-19, Commercial, natural gas boilers.
				020105	481	Nielsen et al. (2010)
		1A4b i	Residential	0202	5	IPCC (1997), Tier 1, Table 1-7, Residential, natural gas.
				020204	481	Nielsen et al. (2010)
		1A4c i	Agriculture/ Forestry	0203	1.2	IPCC (1997), Tier 2, Table 1-19, Commercial, natural gas boilers ¹⁾ .
				020304	481	Nielsen et al. (2010)
WASTE	WASTE	1A1a	Public electricity and heat production	0101 0102	0.34	Nielsen et al. (2010)
		1A2a-f	Industry	03	30	IPCC (1997), Tier 1, Table 1-7, Industry, wastes.
		1A4a	Commercial/ Institutional	0201	30	IPCC (1997), Tier 1, Table 1-7, Industry, wastes.
	INDUSTRIAL WASTE	1A2f	Industry	0316	30	IPCC (1997), Tier 1, Table 1-7, Industry, wastes.
BIO-MASS	WOOD	1A1a	Public electricity and heat production	0101 0102	3.1	Nielsen et al. (2010) IPCC (1997), Tier 1, Table 1-7, Energy industries, wood
		1A2 a-f	Industry	03	15	IPCC (1997), Tier 2, Table 1-16, Industry, wood stoker boilers.

Fuel group	Fuel	CRF source category	CRF source category	SNAP	Emission factor, g pr GJ	Reference
		1A4a	Commercial/ Institutional	0201	30	IPCC (1997), Tier 1, Table 1-7, Industry, wood ²⁾ .
		1A4b i	Residential	0202	129.3	DCE estimate based on technology distribution ³⁾
		1A4c i	Agriculture/ Forestry	0203	30	IPCC (1997), Tier 1, Table 1-7, Industry, wood ²⁾ .
STRAW		1A1a	Public electricity and heat production	0101	0.47	Nielsen et al. (2010)
				0102	30	IPCC (1997), Tier 1, Table 1-7, Energy industries, other biomass
		1A4b i	Residential	0202	300	IPCC (1997), Tier 1, Table 1-7, Residential, other biomass.
		1A4c i	Agriculture/ Forestry	020300	300	IPCC (1997), Tier 1, Table 1-7, Agriculture, other biomass.
				020302	30	IPCC (1997), Tier 1, Table 1-7, Energy industries, other biomass
BIO OIL		1A1a	Public electricity and heat production	010102	0.9	IPCC (1997), Tier 2, Table 1-15, Utility Boiler, distillate fuel oil.
				010105	24	Nielsen et al. (2010) assumed same emission factor as for gas oil fuelled engines.
				0102	0.7	IPCC (1997), Tier 2, Table 1-19, Commercial, distillate fuel oil.
		1A2 a-f	Industry	03	0.7	Assumed equal to district heating plants.
		1A4b i	Residential	0202	0.7	IPCC (1997), Tier 2, Table 1-18, Residential, distillate fuel oil.
BIOGAS		1A1a	Public electricity and heat production	0101	1	IPCC (1997), Tier 1, Table 1-7, Energy industries, natural gas. Assumed similar to natural gas (DCE assumption).
				010105	434	Nielsen et al. (2010)
				0102	1	IPCC (1997), Tier 1, Table 1-7, Energy industries, natural gas. Assumed similar to natural gas (DCE assumption).
		1A2 a-f	Industry	03	5	IPCC (1997), Tier 1, Table 1-7, Industry, natural gas. Assumed similar to natural gas (DCE assumption).
				En-gines	434	Nielsen et al. (2010)
		1A4a	Commercial/ Institutional	0201	5	IPCC (1997), Tier 1, Table 1-7, Commercial, natural gas. Assumed similar to natural gas (DCE assumption).
				020105	434	Nielsen et al. (2010)
		1A4c i	Agriculture/ Forestry	0203	5	IPCC (1997), Tier 1, Table 1-7, Agriculture, natural gas. Assumed similar to natural gas (DCE assumption).
				020304	434	Nielsen et al. (2010)
BIO PROD GAS		1A1a	Public electricity and heat production	010105	13	Nielsen et al. (2010)
		1A4a	Commercial/Institutional	020105	13	Nielsen et al. (2010)

- 1) Assumed same emission factors as for commercial plants. Plant capacity and technology are similar for Danish plants.
- 2) Assumed same emission factor as for industrial plants. Plant capacity and technology is similar to industrial plants rather than to residential plants.
- 3) Aggregated emission factor based on the technology distribution in the sector (DEPA, 2013) and technology specific emission factors that refer to: Paulrud et al. (2005), Johansson et al. (2004) and Olsson & Kjällstrand (2005).

CHP plants

A considerable part of the electricity production in Denmark is based on decentralised CHP plants, and well-documented emission factors for these plants are, therefore, of importance. In a project carried out for the electricity transmission company, Energinet.dk, emission factors for CHP plants <25MW_e have been estimated. The work was reported in 2010 (Nielsen et al., 2010).

The work included waste incineration plants, CHP plants combusting wood and straw, natural gas and biogas-fuelled (reciprocating) engines, natural gas fuelled gas turbines, gas oil fuelled engines, gas oil fuelled gas turbines, steam turbines fuelled by residual oil and engines fuelled by biomass producer gas. CH₄ emission factors for these plants all refer to Nielsen et al. (2010). The estimated emission factors were based on existing emission measurements as well as on emission measurements carried out within the project. The number of emission data sets was comprehensive. Emission factors for subgroups of each plant type were estimated, e.g. the CH₄ emission factor for different gas engine types has been determined.

Time series for the CH₄ emission factors are based on a similar project estimating emission factors for year 2000 (Nielsen & Illerup, 2003).

Natural gas, gas engines

SNAP 010105, 030905, 030705, 031005, 031205, 031305, 031405, 031605, 020105, 020204 and 020304

The emission factor for natural gas engines refers to the Nielsen et al. (2010). The emission factor includes the increased emission during start/stop of the engines estimated by Nielsen et al. (2008). Emission factor time series for the years 1990-2007 have been estimated based on Nielsen & Illerup (2003). These three references are discussed below.

Nielsen et al. (2010):

CH₄ emission factors for gas engines were estimated for 2003-2006 and for 2007-2012. The dataset was split in two due to new emission limits for the engines from October 2006. The emission factors were based on emission measurements from 366 (2003-2006) and 157 (2007-2010) engines respectively. The engines from which emission measurements were available for 2007-2010 represented 38 % of the gas consumption. The emission factors were estimated based on fuel consumption for each gas engine type and the emission factor for each engine type. The majority of emission measurements that were not performed within the project related solely to the emission of total unburned hydrocarbon (CH₄ + NMVOC). A constant disaggregation factor was estimated based on 9 emission measurements including both CH₄ and NMVOC.

Nielsen & Illerup (2003):

The emission factor for natural gas engines was based on 291 emission measurements in 114 different plants. The plants from which emission measurements were available represented 44 % of the total gas consumption in gas engines in year 2000.

Nielsen et al. (2008):

This study calculated a start/stop correction factor. This factor was applied to the time series estimated in Nielsen & Illerup (2003). Further, the correction factors were applied in Nielsen et al. (2010).

The emission factor for lean-burn gas engines is relatively high, especially for pre-chamber engines, which account for more than half the gas consumption in Danish gas engines. However, the emission factors for different pre-chamber engine types differ considerably.

The installation of natural gas engines in decentralised CHP plants in Denmark has taken place since 1990. The first engines installed were relatively small open-chamber engines but later mainly pre-chamber engines were installed. As mentioned above, pre-chamber engines have a higher emission factor than open-chamber engines; therefore, the emission factor has increased during the period 1990-1995. After that technical improvements of the engines have been implemented as a result of upcoming emission limits that most installed gas engines had to meet in late 2006 (DEPA, 2005).

The time series were based on:

- Full load emission factors for different engine types in year 2000 (Nielsen & Illerup, 2003), 2003-2006 and 2007-2012 (Nielsen et al., 2010).
- Data for year of installation for each engine and fuel consumption of each engine 1994-2002 from the Danish Energy Agency (DEA, 2003).
- Research concerning the CH₄ emission from gas engines carried out in 1997 (Nielsen & Wit, 1997).
- Correction factors including increased emission during start/stop of the engines (Nielsen et al., 2008).

Table 3.2.30 Time series for the CH₄ emission factor for natural gas fuelled engines.

Year	Emission factor, g per GJ
1990	266
1991	309
1992	359
1993	562
1994	623
1995	632
1996	616
1997	551
1998	542
1999	541
2000	537
2001	522
2002	508
2003	494
2004	479
2005	465
2006	473
2007-2012	481

Gas engines, biogas

SNAP 010105, 030905, 020105 and 020304

The emission factor for biogas engines was estimated to 434 g per GJ in 2012. The emission factor is lower than the factor for natural gas, mainly because most biogas fuelled engines are lean-burn open-chamber engines - not prechamber engines.

Time series for the emission factor have been estimated. The emission factors for biogas engines were based on Nielsen et al. (2010) and Nielsen & Illerup (2003). The two references are discussed below. The time series are shown in Table 3.2.31.

Nielsen et al. (2010):

CH₄ emission factors for gas engines were estimated for 2006 based on emission measurements performed in 2003-2012. The emission factor was based on emission measurements from 10 engines. The engines from which emission measurements were available represented 8 % of the gas consumption. The emission factor was estimated based on fuel consumption for each gas engine type and the emission factor for each engine type. The majority of emission measurements that were not performed within the project related solely to the emission of total unburned hydrocarbon (CH₄ + NMVOC). A constant disaggregation factor was estimated based on 3 emission measurements including both CH₄ and NMVOC.

Nielsen & Illerup (2003):

The emission factor for natural gas engines was based on 18 emission measurements from 13 different engines. The engines from which emission measurements were available represented 18 % of the total biogas consumption in gas engines in year 2000.

Table 3.2.31 Time series for the CH₄ emission factor for biogas fuelled engines.

Year	Emission factor, g per GJ
1990	239
1991	251
1992	264
1993	276
1994	289
1995	301
1996	305
1997	310
1998	314
1999	318
2000	323
2001	342
2002	360
2003	379
2004	397
2005	416
2006	434
2007-2012	434

Gas turbines, natural gas

SNAP 010104, 010504, 030604 and 031104

The emission factor for gas turbines was estimated to be below 1.7 g per GJ in 2005 (Nielsen et al., 2010). The emission factor was based on emission measurements on five plants. The emission factor in year 2000 was 1.5 g per GJ (Nielsen & Illerup, 2003). A time series have been estimated.

CHP, wood

SNAP 010101, 010102, 010103 and 010104

The emission factor for CHP plants combusting wood was estimated to be below 3.1 g per GJ (Nielsen et al., 2010) and the emission factor 3.1 g per GJ has been applied for all years. The emission factor was based on emission measurements on two plants.

CHP, straw*SNAP 010101, 010102, 010103 and 010104*

The emission factor for CHP plants combusting straw was estimated to be below 0.47 g per GJ (Nielsen et al., 2010) and the emission factor 0.47 g per GJ has been applied for all years. The emission factor was based on emission measurements on four plants.

CHP, waste*SNAP 010102, 010103, 010104 and 010203*

The emission factor for CHP plants combusting waste was estimated to be below 0.34 g per GJ in 2006 (Nielsen et al., 2010) and 0.59 g per GJ in year 2000 (Nielsen & Illerup, 2003). A time series have been estimated. The emission factor was based on emission measurements on nine plants.

The emission factor has also been applied for district heating plants.

Residential wood combustion*SNAP 020200, 020202 and 020204*

The emission factor for residential wood combustion is based on technology specific data. The emission factor time series is shown in Table 3.2.32.

Table 3.2.32 CH₄ emission factor time series for residential wood combustion.

Year	Emission factor, g per GJ
1990-2000	224.0
2001	199.3
2002	189.6
2003	187.2
2004	184.5
2005	175.5
2006	165.2
2007	166.5
2008	156.5
2009	143.8
2010	136.6
2011	129.3
2012	123.7

The emission factors for each technology and the corresponding reference are shown in Table 3.2.33. The emission factor time series are estimated based on time series (2000-2012) for wood consumption in each technology (DEPA, 2013). The time series for wood consumption in the ten different technologies are illustrated in Figure 3.2.47. The consumption in pellet boilers and new stoves has increased.

Table 3.2.33 Technology specific CH₄ emission factors for residential wood combustion.

Technology	Emission factor, g per GJ	Reference
Old stove	430	Methane emissions from residential biomass combustion, Paulrud et al. (2005) (SMED report, Sweden)
New stove	215	Assumed ½ the emission factor for old stoves.
Stove according to resent Danish legislation (2008)	125	Estimated based on the emission factor for new stoves and the emission factors for NMVOC.
Eco labelled stove	2	Low emissions from wood burning in an ecolabelled residential boiler. Olsson & Kjällstrand (2005).
Other stove	430	Assumed equal to old stove.
Old boilers with hot water storage	211	Methane emissions from residential biomass combustion, Paulrud et al., 2005 (SMED report, Sweden)
Old boilers without hot water storage	256	Methane emissions from residential biomass combustion, Paulrud et al., 2005 (SMED report, Sweden)
New boilers with hot water storage	50	Emission characteristics of modern and old-type residential boilers fired with wood logs and wood pellets. Johansson et al. (2004)
New boilers without hot water storage	50	Emission characteristics of modern and old-type residential boilers fired with wood logs and wood pellets. Johansson et al. (2004)
Pellet boilers/stoves	3	Methane emissions from residential biomass combustion, Paulrud et al., 2005 (SMED report, Sweden)




Figure 3.2.47 Technology specific wood consumption in residential plants.

Other stationary combustion plants

Emission factors for other plants refer to the IPCC Guidelines (IPCC, 1997).

N₂O

The N₂O emission factors applied for the 2012 inventory are listed in Table 3.2.34. Time series have been estimated for natural gas fuelled gas turbines

and refinery gas fuelled turbines. All other emission factors have been applied unchanged for 1990-2012.

Emission factors for natural gas fuelled reciprocating engines, natural gas fuelled gas turbines, CHP plants < 300 MW combusting wood, straw or residual oil, waste incineration plants, engines fuelled by gas oil and gas engines fuelled by biomass producer gas all refer to emission measurements carried out on Danish plants, Nielsen et al. (2010).

The emission factor for coal-powered plants in public power plants refers to research conducted by Elsam (now part of DONG Energy).

The emission factor for natural gas has been applied for refinery gas. Denmark uses two different N₂O emission factors for refinery gas, one when the gas is utilised in gas turbines and one for use in boilers. The emission factor for gas turbines is nationally referenced while the emission factor for boilers is based on the Revised 1996 IPCC Guidelines (IPCC, 1997). Refinery gas has similar properties as natural gas, i.e. similar nitrogen content in the fuel, which means that N₂O formation will be similar under similar combustion conditions. This is the reasoning behind choosing the emission factor for natural gas rather than for liquid fuel for both turbines and boilers.

All emission factors that are not nationally referenced refer to the IPCC Guidelines (IPCC, 1997).

Table 3.2.34 N₂O emission factors 2012.

Fuel group	Fuel	CRF source category	CRF source category	SNAP	Emission factor, g per GJ	Reference
SOLID	COAL	1A1a	Public electricity and heat production	0101	0.8	Elsam (2005)
				0102	1.6	IPCC (1997), Tier 2, Table 1-15, Utility boiler, pulverised bituminous coal.
		1A2 a-f	Industry	03	1.4	IPCC (1997), Tier 1, Table 1-8, Industry, coal
		1A4b i	Residential	0202	1.4	IPCC (1997), Tier 1, Table 1-8, Residential, coal
		1A4c i	Agriculture/ Forestry	0203	1.4	IPCC (1997), Tier 1, Table 1-8, Commercial, coal
	BROWN COAL BRI.	1A4b i	Residential	0202	1.4	IPCC (1997), Tier 1, Table 1-8, Residential, coal
	COKE OVEN COKE	1A2 a-f	Industry	03	1.4	IPCC (1997), Tier 1, Table 1-8, Industry, coal
		1A4b i	Residential	020200	1.4	IPCC (1997), Tier 1, Table 1-8, Residential, coal
	ANODIC CARBON	1A2 a-f	Industry	03	1.4	IPCC (1997), Tier 1, Table 1-8, Industry, coal
LIQ-UID	PETROLEUM COKE	1A2a-f	Industry – other	03	0.6	IPCC (1997), Tier 1, Table 1-8, Industry, oil
	RESIDUAL OIL	1A1a	Public electricity and heat production	010101	0.3	IPCC (1997), Tier 2, Table 1-15, Utility, residual fuel oil
				010102	5	Nielsen et al. (2010)
				010103		
				010104	0.6	IPCC (1997), Tier 1, Table 1-8, Energy industries, oil
				010203	0.3	IPCC (1997), Tier 2, Table 1-15, Utility, residual fuel oil
		1A1b	Petroleum refining	010306	0.6	IPCC (1997), Tier 1, Table 1-8, Energy industries, oil
		1A2 a-f	Industry	03	5	Nielsen et al. (2010)
				Engines	0.6	IPCC (1997), Tier 2, Table 1-15, Utility, residual fuel oil
		1A4a	Commercial/ Institutional	0201	0.3	IPCC (1997), Tier 2, Table 1-19, Commercial, fuel oil
		1A4b i	Residential	0202	0.6	IPCC (1997), Tier 1, Table 1-8, Residential, oil
		1A4c i	Agriculture/ Forestry	0203	0.3	IPCC (1997), Tier 2, Table 1-19, Commercial, fuel oil
	GAS OIL	1A1a	Public electricity and heat production	010101	0.4	IPCC (1997), Tier 2, Table 1-15, Utility, distillate fuel oil
				010102		
				010103		
				010104	0.6	IPCC (1997), Tier 1, Table 1-8, Energy industries, oil
				010105	2.1	Nielsen et al. (2010)
				0102	0.4	IPCC (1997), Tier 2, Table 1-15, Utility, distillate fuel oil
		1A1b	Petroleum refining	010306	0.6	IPCC (1997), Tier 1, Table 1-8, Energy industries, oil
		1A2 a-f	Industry	03	0.4	IPCC (1997), Tier 2, Table 1-16, Industry, distillate fuel oil boilers
				Turbines	0.6	IPCC (1997), Tier 1, Table 1-8, Industry, oil
				Engines	2.1	Nielsen et al. (2010)
		1A4a	Commercial/ Institutional	0201	0.4	IPCC (1997), Tier 2, Table 1-19, Commercial, distillate fuel oil
				Engines	2.1	Nielsen et al. (2010)
		1A4b i	Residential	0202	0.6	IPCC (1997), Tier 1, Table 1-8, Residential,

Fuel group	Fuel	CRF source category	CRF source category	SNAP	Emission factor, g per GJ	Reference
GAS	KEROSENE	1A4c	Agriculture/ Forestry	0203	0.4	oil IPCC (1997), Tier 2, Table 1-19, Commercial, distillate fuel oil
		1A2 a-f	Industry	03	0.4	IPCC (1997), Tier 2, Table 1-16, Industry, distillate fuel oil boilers
		1A4a	Commercial/ Institutional	0201	0.4	IPCC (1997), Tier 2, Table 1-19, Commercial, distillate fuel oil
		1A4b i	Residential	0202	0.6	IPCC (1997), Tier 1, Table 1-8, Residential, oil
		1A4c i	Agriculture/ Forestry	0203	0.4	IPCC (1997), Tier 2, Table 1-19, Commercial, distillate fuel oil ¹⁾
		1A1a	Public electricity and heat production	0101 0102	0.6	IPCC (1997), Tier 1, Table 1-8, Energy industries, oil
		1A1b	Petroleum refining	010306	0.6	IPCC (1997), Tier 1, Table 1-8, Energy industries, oil
		1A2 a-f	Industry	03	0.6	IPCC (1997), Tier 1, Table 1-8, Industry, oil
		1A4a	Commercial/ Institutional	0201	0.6	IPCC (1997), Tier 1, Table 1-8, Commercial, oil
		1A4b i	Residential	0202	0.6	IPCC (1997), Tier 1, Table 1-8, Residential, oil
	REFINERY GAS	1A1b	Petroleum refining	010304	1	Assumed equal to natural gas fuelled turbines. Based on Nielsen et al. (2010).
				010306	0.1	IPCC (1997), Tier 1, Table 1-8, Energy industries, natural gas
	NATURAL GAS	1A1a	Public electricity and heat production	010101	0.1	IPCC (1997), Tier 1, Table 1-8, Energy industries, natural gas
				010102		
				010103		
				010104	1	Nielsen et al. (2010)
				010105	0.58	Nielsen et al. (2010)
		1A1c	Other energy industries	010504	1	Nielsen et al. (2010)
				03	0.1	IPCC (1997), Tier 1, Table 1-8, Industry, natural gas
				Gas turbines	1	Nielsen et al. (2010)
		1A4a	Commercial/ Institutional	Engines	0.58	Nielsen et al. (2010)
				020100 020103	2.3	IPCC (1997), Tier 2, Table 1-19, Commercial, natural gas boilers
				Engines	0.58	Nielsen et al. (2010)
		1A4b i	Residential	0202	0.1	IPCC (1997), Tier 1, Table 1-8, Residential, natural gas
				Engines	0.58	Nielsen et al. (2010)
		1A4c i	Agriculture/ Forestry	0203	2.3	IPCC (1997), Tier 2, Table 1-19, Commercial, natural gas boilers ¹⁾
				Engines	0.58	Nielsen et al. (2010)
WASTE	WASTE	1A1a	Public electricity and heat production	0101 0102	1.2	Nielsen et al. (2010)
		1A2 a-f	Industry	03	4	IPCC (1997), Tier 1, Table 1-8, Industry, wastes
		1A4a	Commercial/ Institutional	0201	4	IPCC (1997), Tier 1, Table 1-8, Commercial, wastes
	INDUSTR. WASTE	1A2a-f	Industry	03	4	IPCC (1997), Tier 1, Table 1-8, Industry, waste
BIO-MASS	WOOD	1A1a	Public electricity and heat production	0101	0.8	Nielsen et al. (2010)

Fuel group	Fuel	CRF source category	CRF source category	SNAP	Emission factor, g per GJ	Reference
				0102	4	IPCC (1997), Tier 1, Table 1-8, Energy industries, wood
		1A2 a-f	Industry	03	4	IPCC (1997), Tier 1, Table 1-8, Industry, wood
		1A4a	Commercial/ Institutional	0201	4	IPCC (1997), Tier 1, Table 1-8, Commercial, wood
		1A4b i	Residential	0202	4	IPCC (1997), Tier 1, Table 1-8, Residential, wood
		1A4c i	Agriculture/ Forestry	0203	4	IPCC (1997), Tier 1, Table 1-8, Agriculture, wood
	STRAW	1A1a	Public electricity and heat production	0101	1.1	Nielsen et al. (2010)
				0102	4	IPCC (1997), Tier 1, Table 1-8, Energy industries, other biomass
		1A4b i	Residential	0202	4	IPCC (1997), Tier 1, Table 1-8, Residential, other biomass
		1A4c i	Agriculture/ Forestry	0203	4	IPCC (1997), Tier 1, Table 1-8, Agriculture, other biomass
	BIO OIL	1A1a	Public electricity and heat production	0101 0102 Engines	0.4 2.1	IPCC (1997), Tier 2, Table 1-15, Utility, distillate fuel oil Assumed equal to gas oil. Based on Nielsen et al. (2010)
		1A2 a-f	Industry	03	0.4	IPCC (1997), Tier 1, Table 1-8, Industry, oil
		1A4b i	Residential	0202	0.6	IPCC (1997), Tier 1, Table 1-8, Residential, oil
	BIOGAS	1A1a	Public electricity and heat production	0101 0102 Engines	0.1 1.6	IPCC (1997), Tier 1, Table 1-8, Energy industries, natural gas Nielsen et al. (2010)
		1A2 a-f	Industry	03 Engines	0.1 1.6	IPCC (1997), Tier 1, Table 1-8, Industry, natural gas Nielsen et al. (2010)
		1A4a	Commercial/ Institutional	0201 Engines	0.1 1.6	IPCC (1997), Tier 1, Table 1-8, Commercial, natural gas Nielsen et al. (2010)
		1A4c i	Agriculture/ Forestry	0203 Engines	0.1 1.6	IPCC (1997), Tier 1, Table 1-8, Agriculture, natural gas Nielsen et al. (2010)
	BIO PROD GAS	1A1a	Public electricity and heat production	010105	2.7	Nielsen et al. (2010)
		1A4a	Commercial/ Institutional	020105	2.7	Nielsen et al. (2010)

¹⁾ In Denmark, plants in Agriculture/Forestry are similar to Commercial plants.

SO₂ and NO_x emission factors

Emission factors for SO₂ and NO_x listed in Annex 3A-4. The appendix includes references and time series. Further details about the references, additional references, assumptions and discussions are included in Nielsen et al. (2014).

The emission factors refer to:

- The EMEP/EEA Guidebook: EEA (2013) and former editions.
- The IPCC Guidelines, Reference Manual (IPCC, 1997).
- Danish legislation
- Danish research reports including:
 - Two emission measurement programs for decentralised CHP plants (Nielsen et al. 2010; Nielsen & Illerup, 2003).

- Research and emission measurements programs for biomass fuels:
- Research and environmental data from the gas sector:
- Aggregated emission factors for residential wood combustion based on technology distribution (DEPA, 2013) and technology specific emission factors (EEA, 2013; DEPA, 2010b).
- Calculations based on plant-specific emissions from a considerable number of power plants.
- Calculations based on plant-specific emission data from a considerable number of waste incineration plants. These data refer to annual environmental reports published by plant operators.
- Sulphur content data from oil companies and the Danish gas transmission company, Energinet.dk.
- Additional personal communication.

Emission factor time series have been estimated for a considerable number of the emission factors. These are provided in Annex 3A-4.

NMVOC emission factors

Emission factors for NMVOC are listed in Annex 3A-4. The annex includes references and time series. The emission factors for NMVOC refer to:

- An emission measurement program for decentralised CHP plants (Nielsen et al., 2010).
- The EMEP/EEA Guidebook (EEA, 2009).
- Aggregated emission factor based on the technology distribution for residential wood combustion and guidebook (EEA, 2013) emission factors. Technology distribution refers to DEPA (2013).
- DGC Danish Gas Technology Centre 2001, Naturgas – Energi og miljø (DGC, 2001).
- Gruijthuijsen L.v. & Jensen J.K., 2000. Energi- og miljøoversigt, Danish Gas Technology Centre 2000 (In Danish).

CO emission factors

Emission factors for CO are listed in Annex 3A-4. The annex includes references and time series. The emission factors for CO refer to:

- The EMEP/EEA Guidebook (EEA, 2009) and the former update (EEA, 2007).
- IPCC Guidelines (IPCC, 1997)
- An emission measurement program for decentralised CHP plants (Nielsen et al., 2010).
- Danish legislation (DEPA, 2001)
- Aggregated emission factor based on the technology distribution for residential wood combustion and guidebook (EEA, 2013) emission factors. The technology distribution refer to DEPA (2013).
- DCE estimate based on annual environmental reports for Danish waste incineration plants without power production, year 2000.
- Nikolaisen et al. (1998)
- Jensen & Nielsen (1990)
- Bjerrum (2002)
- Sander (2002)
- Gruijthuijsen & Jensen (2000)

Technology specific emission factors for residential wood combustion, NO_x, NMVOC and CO

For the pollutants NO_x, NMVOC and CO emission factors have been based on fuel consumption data and emission factors for 10 different technologies. Technology categories, emission factors and implied emission factors for 2012 are shown in Table 3.2.35. References for the technology specific emission factors are included in Annex 3A-4.

Technology specific emission factors for CH₄ including references are shown on page 175.

For SO₂, time series have not been estimated and the emission factors are shown in Annex 3A-4.

Table 3.2.35 Technology specific emission factors for residential wood combustion.

Technology	NO _x , g/GJ	NMVOC, g/GJ	CO, g/GJ
Old stove	50	1200	8000
New stove	50	600	4000
Stove according to resent Danish legislation (2008)	80	350	4000
Eco labelled stove	95	175	1117
Other stove	50	600	4000
Old boilers with hot water storage	80	350	4000
Old boilers without hot water storage	80	350	4000
New boilers with hot water storage	95	175	1117
New boilers without hot water storage	95	350	2234
Pellet boilers/stoves	80	1200	8000
IEF residential wood combustion, 2012	74	363	2676

3.2.6 Uncertainty

Uncertainty estimates include uncertainty with regard to the total emission inventory as well as uncertainty with regard to trends.

Methodology

Greenhouse gases

The uncertainty for greenhouse gas emissions have been estimated according to the IPCC Good Practice Guidance (IPCC, 2000). The uncertainty has been estimated by two approaches; tier 1 and tier 2. Both approaches are further described in Chapter 1.7.

The **tier 1** approach is based on a normal distribution and a confidence interval of 95 %.

The input data for the tier 1 approach are:

- Emission data for the base year and the latest year.
- Uncertainties for emission factors
- Uncertainty for fuel consumption rates.

The emission source categories applied are listed in Table 3.2.36.

The **tier 2** approach is a Monte Carlo approach based on a lognormal distribution. The input data for the model is also based on 95 % confidence interval. The input data for the tier 2 approach are:

- Fuel consumption data for the base year and the latest year.

- Emission factors or implied emission factors (IEF) for the base year and the latest year
- Uncertainties for emission factors for the base year and the latest year. If the same uncertainty is applied for both years, the data can be indicated as statistically dependent or independent.
- Uncertainties for fuel consumption rates in the base year and the latest year. If the same uncertainty is applied for both years, the data can be indicated as statistically dependent or independent.

The same emission source categories and emission data have been applied for both approaches. The separate uncertainty estimation for gas engine CH₄ emission and CH₄ emission from other plants does not follow the recommendations in the IPCC Good Practice Guidance. Disaggregation is applied, because in Denmark, the CH₄ emission from gas engines is much larger than the emission from other stationary combustion plants, and the CH₄ emission factor for gas engines is estimated with a much smaller uncertainty level than for other stationary combustion plants.

In general, the same uncertainty levels have been applied for both approaches. However, the tier 2 approach allows different uncertainty levels for 1990 and 2012 and this is relevant to a few uncertainties as discussed below. The 2012 uncertainty levels have been applied in the tier 1 approach.

Most of the applied uncertainty estimates for activity rates and emission factors are default values from the IPCC Reference Manual or aggregated by DCE based on the default values. Some of the uncertainty estimates are, however, based on national estimates.

In general, the uncertainty of the fuel consumption data has been assumed to be the same in 1990 and 2012 and the uncertainty has been assumed to be statistically independent. However, a considerable part of the residential wood consumption is non-traded and the uncertainty of biomass consumption has been assumed statistically dependent.

Fuel consumption data for waste are more uncertain for 1990 than for 2012.

For coal and refinery gas combustion, the uncertainty of the CO₂ emission factor is lower in 2012 than in 1990 due to availability of EU ETS data. Further, the CO₂ emission factor for the fossil part of waste is less uncertain for 2012 than for 1990.

The uncertainty of the CH₄ emission factors for gas engines have been assumed higher in 1990 than in 2012 due to the emission measurement programmes on which the emission factors in later years are based.

All other uncertainty levels for emission factors have been assumed equal in 1990 and 2012 and statistically dependent.

Table 3.2.36 Uncertainty rates for fuel consumption and emission factors, 2012.

IPCC Source category	Gas	Fuel consumption uncertainty, %		Emission factor uncertainty, %	
		1990	2012	1990	2012
Stationary Combustion, Coal, CO ₂	CO ₂	0.9 ²⁾	0.9 ⁷⁾	4 ¹⁰⁾	0.5 ⁷⁾
Stationary Combustion, BKB, CO ₂	CO ₂	2.9 ²⁾	3.0 ²⁾		5 ¹⁾
Stationary Combustion, Coke ³³ , CO ₂	CO ₂	1.8 ²⁾	1.8 ²⁾		5 ¹⁾
Stationary Combustion, Fossil waste, CO ₂	CO ₂	10.0 ²⁾	5.0 ²⁾	20 ⁵⁾	10 ⁵⁾
Stationary Combustion, Petroleum coke, CO ₂	CO ₂	1.7 ²⁾	2.0 ²⁾		5 ¹⁾
Stationary Combustion, Residual oil, CO ₂	CO ₂	1.2 ²⁾	1.2 ²⁾	2 ⁴⁾	2 ⁷⁾
Stationary Combustion, Gas oil, CO ₂	CO ₂	2.9 ²⁾	2.2 ²⁾		4 ¹⁰⁾
Stationary Combustion, Kerosene, CO ₂	CO ₂	3.0 ²⁾	2.0 ²⁾		5 ¹⁾
Stationary Combustion, LPG, CO ₂	CO ₂	1.7 ²⁾	1.8 ²⁾		5 ¹⁾
Stationary Combustion, Refinery gas, CO ₂	CO ₂	1.0 ²⁾	1.0 ²⁾	5 ¹⁾	2 ¹²⁾
Stationary Combustion, Natural gas, CO ₂	CO ₂	1.2 ²⁾	1.0 ²⁾		0.4 ⁸⁾
Stationary Combustion, SOLID, CH ₄	CH ₄	0.9 ²⁾	1.0 ²⁾		100 ¹⁾
Stationary Combustion, LIQUID, CH ₄	CH ₄	1.5 ²⁾	0.8 ²⁾		100 ¹⁾
Stationary Combustion, GAS, CH ₄	CH ₄	1.0 ⁸⁾	1.0 ⁸⁾		100 ¹⁾
Natural gas fuelled engines, GAS, CH ₄	CH ₄	1.0 ⁹⁾	1.0 ⁹⁾	10 ¹¹⁾	2 ³⁾
Stationary Combustion, WASTE, CH ₄	CH ₄	10.0 ⁵⁾	5.0 ⁵⁾		100 ¹⁾
Stationary Combustion, BIOMASS, CH ₄	CH ₄	14.9 ²⁾	15.7 ²⁾		100 ¹⁾
Biogas fuelled engines, BIOMASS, CH ₄	CH ₄	6.8 ²⁾	3.9 ²⁾	20 ¹¹⁾	10 ¹¹⁾
Stationary Combustion, SOLID, N ₂ O	N ₂ O	0.9 ²⁾	1.0 ²⁾		400 ^{6) 13)}
Stationary Combustion, LIQUID, N ₂ O	N ₂ O	1.5 ²⁾	0.8 ²⁾		1000 ^{1) 13)}
Stationary Combustion, GAS, N ₂ O	N ₂ O	1.0 ⁸⁾	1.0 ⁸⁾		750 ^{6) 13)}
Stationary Combustion, WASTE, N ₂ O	N ₂ O	10.0 ⁵⁾	5.0 ⁵⁾		400 ^{6) 13)}
Stationary Combustion, BIOMASS, N ₂ O	N ₂ O	14.7 ²⁾	15.3 ²⁾		400 ^{6) 13)}

1) IPCC Good Practice Guidance, default value (IPCC, 2000).

2) Estimated by DCE based on default uncertainty levels in IPCC Good Practice Guidance, Table 2.6 (IPCC, 2000).

3) Jørgensen et al. (2010). Uncertainty data for NMVOC + CH₄.

4) Jensen & Lindroth (2002).

5) Estimated by DCE based on Astrup et al. (2012).

6) DCE, rough estimate based on a default value of 400 % when the emission factor is based on emission measurements from plants in Denmark.

7) Emission data based on EU ETS data.

8) Lindgren (2010). Personal communication, Tine Lindgren, Energinet.dk, e-mail 2010-03-16.

9) Equal to natural gas total. DCE assumption.

10) DCE assumption based on EU ETS data interval and IPCC Guidelines (IPCC, 1997) data interval.

11) DCE estimate based on Nielsen et al. (2010).

12) DCE assumption based on the fact that data are based on EU ETS data.

13) With a truncation of twice the uncertainty rate. The truncation is relevant for the very large uncertainty rates for N₂O emission factors due to the log-normal distribution applied in the tier 2 model.

Other pollutants

With regard to other pollutants, IPCC methodologies for uncertainty estimates have been adopted for the LRTAP Convention reporting activities (Pulles & Aardenne, 2004). The Danish uncertainty estimates are based on the simple Tier 1 approach.

The uncertainty estimates are based on emission data for the base year and year 2012 as well as on uncertainties for fuel consumption and emission factors for each of the main SNAP source categories. The applied uncer-

³³ Including anodic carbon.

tainties for activity rates and emission factors are default values referring to Pulles & Aardenne (2004). The default uncertainties for emission factors are given in letter codes representing an uncertainty range. It has been assumed that the uncertainties were in the lower end of the range for all sources and pollutants. The applied uncertainties for emission factors are listed in Table 3.2.37. The uncertainty for fuel consumption in stationary combustion plants is assumed to be 2 %.

Table 3.2.37 Uncertainty rates for emission factors, %.

SNAP source category	SO ₂	NO _x	NMVOC	CO
01	10	20	50	20
02	20	50	50	50
03	10	20	50	20

Results

The tier 1 uncertainty estimates for stationary combustion emission inventories are shown in Table 3.2.38. Detailed calculation sheets are provided in Annex 3A-7. The tier 2 uncertainty estimates are shown in Table 3.2.39 and detailed results are provided in Annex 3A-7.

The tier 1 uncertainty interval for greenhouse gas is estimated to be ± 2.2 % and trend in greenhouse gas emission is -39 % ± 1.1 %-age points. The main sources of uncertainty for greenhouse gas emission 2012 are the N₂O emission from combustion of biomass and gaseous fuels, CH₄ emission from biomass combustion and CO₂ emission from fossil waste combustion. The main sources of uncertainty in the trend in greenhouse gas emission are the CO₂ emission from coal and fossil waste combustion and the N₂O emission from combustion of biomass and liquid fuels.

The total emission uncertainty is 7.5 % for SO₂, 16 % for NO_x, 45 % for NMVOC and 44 % for CO.

The tier 2 approach points out N₂O and CH₄ emissions from combustion of biomass and CO₂ from fossil waste combustion as the main contributors to the total uncertainty for greenhouse gas emission from stationary combustion.

Table 3.2.38 Danish uncertainty estimates, tier 1 approach, 2012.

Pollutant	Uncertainty Total emission, %	Trend 1990-2012, %	Uncertainty trend, %-age points
GHG	$\pm 2.2\%$	-39%	$\pm 1.1\%$
CO ₂	$\pm 0.9\%$	-40%	$\pm 0.7\%$
CH ₄	$\pm 46\%$	134%	$\pm 86\%$
N ₂ O	$\pm 260\%$	-2.0%	$\pm 244\%$
SO ₂	$\pm 7.5\%$	-95%	$\pm 0.3\%$
NO _x	$\pm 16\%$	-72%	$\pm 1.8\%$
NMVOC	$\pm 45\%$	12%	$\pm 6\%$
CO	$\pm 44\%$	-5.2%	$\pm 2.7\%$

Table 3.2.39 Danish uncertainty estimates, tier 2 approach, 2012.

Pollutant	Uncertainty of total emission, %		Trend 1990-2012, %	Uncertainty of trend, %-age points	
GHG	-1.4	+3.9	-38.9	-2.9	+2.7
CO ₂	-0.9	+1.9	-39.8	-2.8	+2.6
CH ₄	-28	+138	131	-19	+41
N ₂ O	-73	+1096	-2.8	-171	+119

The results are illustrated and compared in figure 3.2.48. The uncertainties are in the same level for each pollutant. The emission data shown for the tier 1 approach are the CRF emission data. The tier 2 emission levels are median values based on the Monte Carlo approach.




Figure 3.2.48 Uncertainty level, the two approaches are compared for 2012.

3.2.7 Source specific QA/QC and verification

An updated quality manual for the Danish emission inventories has been published in 2013 (Nielsen et al., 2013). The quality manual describes the concepts of quality work and definitions of sufficient quality, critical control points and a list of Point for Measuring (PM).

Documentation concerning verification of the Danish GHG emission inventories has been published by (Fauser et al., 2013). In addition, the IPCC reference approach for CO₂ emission is an important verification of the CO₂ emission from the energy sector. The reference approach for the energy sector is shown in Chapter 3.4.

Information on the Danish QA/QC plan is included in Chapter 1.6. Source specific QA/QC and PM's are shown below.

National external review

The 2004, 2006 and 2009 updates of the sector report for stationary combustion has been reviewed by external experts (Nielsen & Illerup, 2004; Nielsen & Illerup, 2006; Nielsen et al., 2009). An updated sector report for stationary combustion will be published in 2014 (Nielsen et al., 2014). The national external review forms a vital part of the QA activities for stationary combustion.

The 2004, 2006 and 2009 updates of this report were reviewed by Jan Erik Johnsson from the Technical University of Denmark, Bo Sander from Elsam Engineering and Annemette Geertinger from FORCE Technology.

Data storage, level 1

Table 3.2.40 lists the sector specific PM's for data storage level 1.

Table 3.2.40 List of PM, data storage level 1.

Level	CCP	Id	Description	Sectoral/general	Stationary combustion
Data Storage level 1	1. Accuracy	DS.1.1.1	General level of uncertainty for every dataset including the reasoning for the specific values.	Sectoral	Uncertainties are estimated and references given in NIR chapter 3.2.6.
	2. Comparability	DS1.2.1	Comparability of the emission factors/calculation parameters with data from international guidelines, and evaluation of major discrepancies.	Sectoral	In general if national referenced emission factors differ considerably from IPCC Guideline/EEA Guidebook values this is discussed in NIR chapter 3.2.5. This documentation is improved annually based on reviews. At CRF level, a project has been carried out comparing the Danish inventories with those of other countries (Fauser et al., 2013).
	3.Completeness	DS.1.3.1	Ensuring that the best possible national data for all sources are included, by setting down the reasoning behind the selection of datasets.	Sectoral	A list of external data are shown and discussed below.
	4.Consistency	DS.1.4.1	The original external data has to be archived with proper reference.	Sectoral	It is ensured that all external data are archived at DCE. Subsequent data processing takes place in other spreadsheets or databases. The datasets are archived annually in order to ensure that the basic data for a given report are always available in their original form.
	6.Robustness	DS.1.6.1	Explicit agreements between the external institution holding the data and DCE about the conditions of delivery	Sectoral	For stationary combustion, a data delivery agreement is made with the DEA. NERI (now DCE) and DEA have renewed the data delivery agreement in 2010. An update of the data delivery agreement will be signed in 2014. Most of the other external data sources are available due to legislative requirements. See Table 3.2.41.
	7.Transparency	DS.1.7.1	Listing of all archived datasets and external contacts.	Sectoral	A list of external datasets and external contacts is shown in Table 3.2.41 below.

Table 3.2.41 List of external data sources.

Dataset	Description	AD or Emf.	Reference	Contact(s)	Data agreement/ Comment
Energiproducenttællingen.xls	Data set for all electricity and heat producing plants.	Activity data	The Danish Energy Agency (DEA)	Kaj Stærkind	Data agreement 2010. An updated data agreement will be signed in 2014.
Gas consumption for gas engines and gas turbines 1990-1994	Historical data set for gas engines and gas turbines.	Activity data	The Danish Energy Agency (DEA)	Kaj Stærkind	No data agreement. Historical data
Basic data (Grunddata.xls)	The Danish energy statistics. Data set applied for both the reference approach and the national approach.	Activity data	The Danish Energy Agency (DEA)	Jane Rusbjerg	Data agreement 2010. However, the data set is also published as part of national energy statistics An updated data agreement will be signed in 2014.
Energy statistics for industrial sub-sectors	Disaggregation of the industrial fuel consumption.	Activity data	The Danish Energy Agency (DEA)	Jane Rusbjerg	Only informal data delivery agreement. The data set will be included in the next update of the data delivery agreement with DEA.
SO ₂ & NO _x data, plants > 25 MW _e	Annual emission data for all power plants > 25 MW _e . Includes information on methodology: measurements or emission factor.	Emissions	Energinet.dk	Christian F.B. Nielsen	No data agreement.
Emission factors	Emission factors refer to a large number of sources.	Emission factors	See chapter regarding emission factors		Some of the annually updated CO ₂ emission factors are based on EU ETS data, see below. For the other emission factors no formal data delivery agreement.
Annual environmental reports / environmental data	Emissions from plants defined as large point sources	Emissions	Various plants		No data agreement necessary. Plants are obligated by law (DEPA 2010a) and data published on the Danish EPA homepage.
EU ETS data	Plant specific CO ₂ emission factors	Emission factors and fuel consumption	The Danish Energy Agency (DEA)	Dorte Maimann Helen Falster	Plants are obligated by law. The availability of detailed information is part of the renewed data agreement with DEA.

Energiproducenttaellingen - statistic on fuel consumption from district heating and power plants (DEA)

The data set includes all plants producing power or district heating. The spreadsheet from DEA is listing fuel consumption of all plants included as large point sources in the emission inventory. The statistic on fuel consumption from district heating and power plants is regarded as complete and with no significant uncertainty since the plants are bound by law to report their fuel consumption and other information.

Gas consumption for gas engines and gas turbines 1990-1994 (DEA)

For the years 1990-1994, DEA has estimated consumption of natural gas and biogas in gas engines and gas turbines (DEA, 2003). Estimated fuel consumption data for 1990-1993 was based on engine specific data for year of installation and for fuel consumption in 1994. DCE assesses that the DEA estimate is the best available data.

Basic data (DEA)

The Danish Energy statistics. The spreadsheet from DEA is used for the CO₂ emission calculation in accordance with the IPCC reference approach and is also the first data set applied in the national approach. The data set is included in the data delivery agreement with DEA, but it is also published annually on DEA's homepage.

Energy statistics for industrial subsectors (DEA)

The data includes disaggregation of the fuel consumption for industrial plants. The data set is estimated for the reporting to Eurostat. The data delivery agreement is informal at this time, but the dataset will be included in the 2014 update of the agreement with DEA.

SO₂ and NO_x emission data from electricity producing plants > 25MW_e (Energinet.dk)

Plants larger than 25 MW_e are obligated to report emission data for SO₂ and NO_x to the DEA annually. Data are collected by Energinet.dk and forwarded to DEA. Data are on production unit level and classified. The data on plant level are part of the plants annually environmental reports. DCE's QC of the data consists of a comparison with data from previous years and with data from the plants' annual environmental reports.

Emission factors

For specific references, see the Chapter 3.2.5 regarding emission factors. Some of the annually updated CO₂ emission factors are based on EU ETS data, see below.

Annual environmental reports (DEPA)

A large number of plants are obligated by law to report annual environmental data including emission data. DCE compares the data with those from previous years and large discrepancies are checked.

EU ETS data (DEA)

EU ETS data includes information on fuel consumption, heating values, carbon content of fuel, oxidation factor and CO₂ emissions. DCE receives the verified reports for all plants which utilises a detailed estimation methodology. DCE's QC of the received data consists of comparing to calculation using standard emission factors as well as comparing reported values with those for previous years. The data delivery agreement is informal at this

time, but the dataset will be included in the 2014 update of the agreement with DEA.

Data processing, level 1

Table 3.2.42 lists the sector specific PM's for data processing level 1.

Table 3.2.42 List of PM, data processing level 1.

Level	CCP	Id	Description	Sectoral / general	Stationary combustion
Data Processing level 1	1. Accuracy	DP.1.1.1	Uncertainty assessment for every data source not part of DS.1.1.1 as input to Data Storage level 2 in relation to type and scale of variability.	Sectoral	Uncertainties are estimated and references given in NIR chapter 3.2.6.
	2.Comparability	DP.1.2.1	The methodologies have to follow the international guidelines suggested by UNFCCC and IPCC.	Sectoral	The methodological approach is consistent with international guidelines. An overview of tiers is given in NIR Chapter 3.2.5
	3.Completeness	DP.1.3.1	Identification of data gaps with regard to data sources that could improve quantitative knowledge.	Sectoral	The energy statistics is considered complete.
	4.Consistency	DP.1.4.1	Documentation and reasoning of methodological changes during the time series and the qualitative assessment of the impact on time series consistency.	Sectoral	The two main methodological changes in the time series; implementation of Energiproducenttaeligen (plant specific fuel consumption data) from 1994 onwards and implementation of EU ETS data from 2006 onwards is discussed in NIR chapter 3.2.5.
	5.Correctness	DP.1.5.2	Verification of calculation results using time series	Sectoral	Time series for activity data on SNAP and CRF source category level are used to identify possible errors. Time series for emission factors and the emission from CRF subcategories are also examined.
		DP.1.5.3	Verification of calculation results using other measures	Sectoral	The IPCC reference approach validates the fuel consumption rates and CO ₂ emission. Both differ less than 2.0 % (1990-2012). The reference approach is further discussed in NIR Chapter 3.4.
	7.Transparency	DP.1.7.1	The calculation principle, the equations used and the assumptions made must be described.	Sectoral	This is included in NIR chapter 3.2.5.
		DP.1.7.2	Clear reference to dataset at Data Storage level 1	Sectoral	This is included in NIR chapter 3.2.5.
		DP.1.7.3	A manual log to collect information about recalculations.	Sectoral	-

Data storage, level 2

Table 3.2.43 lists the sector specific PM's for data storage level 2.

Table 3.2.43 List of PM, data storage level 2.

Level	CCP	Id	Description	Sectoral / general	Stationary combustion
Data Storage level 2	5. Correctness	DS.2.5.1	Check if a correct data import to level 2 has been made	Sectoral	To ensure a correct connection between data on level 2 and level 1, different controls are in place, e.g. control of sums and random tests.

Data storage level 4

Table 3.2.44 lists the sector specific PM's for data storage level 4.

Table 3.2.44 List of PM, data storage level 4.

Level	CCP	Id	Description	Sectoral / general	Stationary combustion
Data Storage level 4	4. Consistency	DS.4.4.3	The IEFs from the CRF are checked both regarding level and trend. The level is compared to relevant emission factors to ensure correctness. Large dips/jumps in the time series are explained.	Sectoral	Large dips/jumps in time series are discussed and explained in NIR chapter 3.2.3 and 3.2.4.

Other QC procedures

Some automated checks have been prepared for the emission databases:

- Check of units for fuel rate, emission factors and plant-specific emissions.
- Check of emission factors for large point sources. Emission factors for pollutants that are not plant-specific should be the same as those defined for area sources.
- Additional checks on database consistency.
- Emission factor references are included in this report (Chapter 3.2.5 and Appendix 3A-4).
- Annual environmental reports are kept for subsequent control of plant-specific emission data.
- QC checks of the country-specific emission factors have not been performed, but most factors are based on input from companies that have implemented some QA/QC work. The major power plant owner/operators in Denmark, DONG Energy and Vattenfall have obtained the ISO 14001 certification for an environmental management system. The Danish Gas Technology Centre and Force Technology both run accredited laboratories for emission measurements.
- The emission from each large point source is compared with the emission reported the previous year.

3.2.8 Source specific recalculations and improvements

Recalculations for stationary combustion 2011 are shown in Table 3.2.45. The main calculations are discussed below.

Table 3.2.45 Recalculations for stationary combustion, 2011.

	CO ₂ , Gg CO ₂	CH ₄ , Gg CO ₂ eqv.	N ₂ O Gg CO ₂ eqv.	CO ₂ , %	CH ₄ , %	N ₂ O %
1.A.1. Energy Industries	7.72	2.13	0.02	0%	1%	0%
Liquid Fuels	0.41	0.03	0.03	0%	6%	2%
Solid Fuels	9.95	0.00	0.05	0%	0%	0%
Gaseous Fuels	-2.63	1.59	-0.08	0%	1%	0%
Biomass	4.43	0.50	0.02	0%	2%	0%
Other Fuels	0.00	0.00	0.00	0%	0%	0%
a. Public Electricity and Heat Production	28.79	2.14	0.14	0%	1%	0%
Liquid Fuels	0.10	0.03	0.03	0%	29%	3%
Solid Fuels	9.95	0.00	0.05	0%	0%	0%
Gaseous Fuels	18.74	1.61	0.03	1%	1%	0%
Biomass	4.43	0.50	0.02	0%	2%	0%
Other Fuels	0.00	0.00	0.00	0%	0%	0%
b. Petroleum Refining	0.30	0.00	0.00	0%	0%	0%
Liquid Fuels	0.30	0.00	0.00	0%	0%	0%
Solid Fuels	0.00	0.00	0.00	0%	0%	0%
Gaseous Fuels	0.00	0.00	0.00	0%	0%	0%
Biomass	0.00	0.00	0.00	0%	0%	0%
Other Fuels	0.00	0.00	0.00	0%	0%	0%
c. Manufacture of Solid Fuels and Other Energy Industries	-21.37	-0.01	-0.12	-1%	-1%	-1%
Liquid Fuels	0.00	0.00	0.00	0%	0%	0%
Solid Fuels	0.00	0.00	0.00	0%	0%	0%
Gaseous Fuels	-21.37	-0.01	-0.12	-1%	-1%	-1%
Biomass	0.00	0.00	0.00	0%	0%	0%
Other Fuels	0.00	0.00	0.00	0%	0%	0%
1.A.2 Manufacturing Industries and Construction	132.50	-0.76	3.26	3%	-6%	9%
Liquid Fuels	127.36	0.05	2.46	7%	4%	13%
Solid Fuels	3.32	0.01	0.02	1%	1%	1%
Gaseous Fuels	5.80	-0.99	-0.01	0%	-14%	-1%
Biomass	72.48	0.20	0.86	8%	6%	8%
Other Fuels	-3.98	-0.03	-0.06	-5%	-5%	-5%
a. Iron and Steel	-11.35	-0.01	0.00	-13%	-13%	-3%
Liquid Fuels	0.23	0.00	0.00	86%	43%	608%
Solid Fuels	0.00	0.00	0.00	0%	0%	0%
Gaseous Fuels	-11.60	-0.01	-0.01	-13%	-13%	-13%
Biomass	0.00	0.00	0.00	0%	0%	0%
Other Fuels	0.00	0.00	0.00	0%	0%	0%
b. Non-Ferrous Metals	-2.57	0.00	0.01	-39%	-42%	355%
Liquid Fuels	0.73	0.00	0.01	17180%	9090%	116882%
Solid Fuels	0.00	0.00	0.00	0%	0%	0%
Gaseous Fuels	-3.30	0.00	0.00	-50%	-50%	-50%
Biomass	0.00	0.00	0.00	0%	0%	0%
Other Fuels	0.00	0.00	0.00	0%	0%	0%
c. Chemicals	97.74	0.04	0.34	58%	38%	100%
Liquid Fuels	15.84	0.01	0.32	10570%	6489%	80732%
Solid Fuels	0.00	0.00	0.00	0%	0%	0%
Gaseous Fuels	82.45	0.04	0.04	49%	47%	14%
Biomass	0.00	0.00	0.00	0%	0%	0%
Other Fuels	0.00	0.00	0.00	0%	0%	0%
d. Pulp, Paper and Print	-14.01	0.55	2.30	-10%	108%	121%
Liquid Fuels	2.40	0.00	0.05	358%	163%	2763%
Solid Fuels	0.00	0.00	0.00	0%	0%	0%
Gaseous Fuels	-14.88	-0.01	-0.01	-11%	-10%	-3%
Biomass	201.02	0.57	2.27	140%	136%	141%
Other Fuels	-1.53	-0.01	-0.02	-74%	-74%	-74%
e. Food Processing, Beverages and Tobacco	106.81	-0.14	1.31	9%	-3%	20%

	CO ₂ , Gg CO ₂	CH ₄ , Gg CO ₂ eqv.	N ₂ O Gg CO ₂ eqv.	CO ₂ , %	CH ₄ , %	N ₂ O %
Liquid Fuels	68.42	0.02	1.38	30%	29%	31%
Solid Fuels	-40.85	-0.09	-0.19	-18%	-18%	-18%
Gaseous Fuels	77.09	0.05	0.04	10%	1%	5%
Biomass	1.37	-0.14	0.04	4%	-60%	23%
Other Fuels	2.16	0.02	0.03	80%	80%	80%
f. Other (please specify)(4)	-44.13	-1.21	-0.69	-2%	-15%	-3%
Cement production	0.00	0.00	0.00	0%	0%	0%
Liquid Fuels	0.00	0.00	0.00	0%	0%	0%
Solid Fuels	0.00	0.00	0.00	0%	0%	0%
Gaseous Fuels	0.00	0.00	0.00	0%	0%	0%
Biomass	0.00	0.00	0.00	0%	0%	0%
Other Fuels	0.00	0.00	0.00	0%	0%	0%
Non-road machinery	0.00	0.00	0.00	0%	0%	0%
Liquid Fuels	12.19	0.00	0.16	1%	1%	1%
Solid Fuels	0.00	0.00	0.00	0%	0%	0%
Gaseous Fuels	0.00	0.00	0.00	0%	0%	0%
Biomass	0.00	0.00	0.00	0%	0%	0%
Other Fuels	0.00	0.00	0.00	0%	0%	0%
Other non-specified	0.00	0.00	0.00	0%	0%	0%
Liquid Fuels	27.55	0.01	0.53	120%	81%	929%
Solid Fuels	44.15	0.10	0.20	39%	42%	42%
Gaseous Fuels	-123.38	-0.96	-0.08	-16%	-32%	-18%
Biomass	-128.79	-0.22	-1.44	-19%	-11%	-19%
Other Fuels	-4.06	-0.03	-0.06	-39%	-39%	-39%
1.A.4 Other Sectors	17.66	15.25	-1.00	0%	10%	-1%
Liquid Fuels	-56.21	0.07	-1.39	-2%	1%	-4%
Solid Fuels	3.31	0.01	0.02	3%	2%	3%
Gaseous Fuels	66.65	0.17	0.28	3%	1%	3%
Biomass	5.53	14.97	0.04	0%	13%	0%
Other Fuels	3.90	0.03	0.06	140%	140%	140%
a. Commercial/Institutional	-5.87	-0.19	0.09	-1%	-1%	1%
Liquid Fuels	-15.57	-0.06	-0.05	-4%	-2%	-4%
Solid Fuels	0.00	0.00	0.00	0%	0%	0%
Gaseous Fuels	5.80	0.02	0.07	1%	0%	1%
Biomass	3.03	-0.19	0.01	1%	-3%	0%
Other Fuels	3.90	0.03	0.06	140%	140%	140%
b. Residential	57.75	15.15	0.05	2%	14%	0%
Liquid Fuels	11.72	0.01	0.04	1%	0%	2%
Solid Fuels	0.00	0.00	0.00	0%	0%	0%
Gaseous Fuels	46.03	0.12	0.03	3%	1%	3%
Biomass	-1.58	15.03	-0.02	0%	16%	0%
Other Fuels	0.00	0.00	0.00	0%	0%	0%
c. Agriculture/Forestry/Fisheries	-34.22	0.29	-1.13	-2%	1%	-3%
Liquid Fuels	-52.36	0.12	-1.38	-3%	5%	-5%
Solid Fuels	3.31	0.01	0.02	3%	3%	3%
Gaseous Fuels	14.82	0.03	0.18	12%	0%	16%
Biomass	4.08	0.13	0.05	1%	1%	2%
Other Fuels	0.00	0.00	0.00	0%	0%	0%

For stationary combustion plants, the emission estimates for the years 1990-2011 have been updated according to the latest energy statistics published by the Danish Energy Agency. The update included both end use and transformation sectors as well as a source category update. The changes in the energy statistics are largest for the years 2009, 2010 and 2011.

For CO₂ the largest recalculation is in source category Manufacturing industries and constructions. The recalculation is related to liquid fuels and is a

result of correction of an error. The consumption of residual oil was underestimated in the former inventories. The CO₂ emission from liquid fuels applied in manufacturing industries and constructions 2011 is 7% higher in the 2014 reporting than in the 2013 reporting.

The CH₄ emission from residential wood combustion has been recalculated based on improved emission factors for stoves. This has caused a 16 % increase of the CH₄ emission reported for biomass fuels in residential plants for 2011.

For N₂O the largest recalculation is in source category Manufacturing industries and constructions. This recalculation is also related to the former underestimate for residual oil. The N₂O emission from liquid fuels applied in manufacturing industries and constructions 2011 is 13% higher in the 2014 reporting than in the 2013 reporting.

Improvements related to reviews

The text in the CRF documentation box in Table 1.A(c) has been improved and a reference to the relevant NIR chapter have been added.

Data for lubricants have been added in Table 1.C.

3.2.9 Source specific planned improvements

The next emission inventory will implement the 2006 IPCC Guidelines (IPCC, 2006).

3.2.10 References for chapter 3.2 and Annex 3A

Andersen, M.A. 1996: Elkraft, personal communication letter 07-05-1996.

Astrup, T., Larsen, A.W., Fuglsang, K. & Pedersen, N.H. 2012: PSO-0213, Biogenic Carbon in Danish Combustible Waste. DTU 2012.

Bjerrum, M., 2002: Danish Technological Institute, personal communication 09-10-2002.

Bruun, J. 2012: Energinet.dk, personal communication, email 03-01-2012.

Christiansen, M., 1996: Elsam, personal communication, letter 07-05-1996.

Christiansen (2003): Personal communication, Jørgen Christiansen, Energigruppen Jylland, 2003

Danish Energy Agency (DEA), 2003: Data for installation and fuel consumption for each gas engine 1994-2002. Estimated fuel consumption data for gas engines in 1990-1993. Unpublished.

Danish Energy Agency (DEA), 2013a: The Danish energy statistics, Available at:

<http://www.ens.dk/en/info/facts-figures/energy-statistics-indicators-energy-efficiency/annual-energy-statistics> (2014-02-14)

Danish Energy Agency (DEA), 2013b: The Danish energy statistics, Energiproducenttællingen 1990-2012. Unpublished.

Danish Energy Agency (DEA), 2013c: The Danish energy statistics, International reporting. Available at:

<http://www.ens.dk/en/info/facts-figures/energy-statistics-indicators-energy-efficiency/annual-reporting-ieaeurostatun> (2014-02-14).

Danish Environmental Protection Agency (DEPA), 1990: Bekendtgørelse om begrænsning af emissioner af svovl-dioxid, kvælstofoxider og støv fra store fyringsanlæg, Bekendtgørelse 689 af 15/10/1990, (Danish legislation). Available at: <https://www.retsinformation.dk/Forms/R0710.aspx?id=48656> (2014-02-14)

Danish Environmental Protection Agency (DEPA), 2001(a): Luftvejledningen, Begrænsning af luftforurening fra virksomheder, Vejledning fra Miljøstyrelsen nr. 2, 2001 (Danish legislation). Available at:

<https://www.retsinformation.dk/forms/R0710.aspx?id=12728> (2014-02-14)

Danish Environmental Protection Agency (DEPA), 2001b: Bekendtgørelse om begrænsning af svovlindholdet i visse flydende og faste brændstoffer. Bekendtgørelse 532 af 14/12/2006. Danish legislation. Available at:

<https://www.retsinformation.dk/Forms/R0710.aspx?id=12458> (2013-08-08).

Danish Environmental Protection Agency (DEPA), 2005: Bekendtgørelse om begrænsning af emission af nitrogenoxider, uforbrændte carbonhydrider og carbonmonooxid mv. fra motorer og turbiner. Bekendtgørelse 621 af 23/06/2005. (Danish legislation). Available at:

<https://www.retsinformation.dk/Forms/R0710.aspx?id=12836&exp=1> (2013-02-13).

Danish Environmental Protection Agency (DEPA), 2010a: Bekendtgørelse om visse virksomheders afgivelse af miljøoplysninger. Bekendtgørelse 210 af 03/03/2010. (Danish legislation). Available at:

<https://www.retsinformation.dk/Forms/R0710.aspx?id=128761> (2013-02-08).

Danish Environmental Protection Agency (DEPA), 2010b: Emissioner fra træfyrede brændeovne/kedler (Emissions from wood fired stoves/boilers). Danish Environmental Protection Agency, 2010. Available at:

<http://www.mst.dk/Publikationer/Publikationer/2012/04/978-87-92617-85-9.htm> (2014-02-14).

Danish Environmental Protection Agency (DEPA), 2010c: Bekendtgørelse om svovlindholdet i faste og flydende brændstoffer. Bekendtgørelse 1098 af 19/09/2010. Danish legislation. Available at:

<https://www.retsinformation.dk/Forms/R0710.aspx?id=133318> (2014-02-14).

Danish Environmental Protection Agency (DEPA) , 2013. Forudsætninger for fremskrivning af emissioner fra brændeovne (In Danish). Hessberg & Justesen Miljøteknologi, J.nr. MST-5261-00007, Ref. stus, Den 5. november 2013 . (Related to e-mail correspondence 2011-04-28 and 2013-11-05).

Danish Gas Technology Centre (DGC), 2001: Naturgas – Energi og miljø (In Danish).

Danish Ministry of Taxation (DMT), 1998: Bekendtgørelse af lov om afgift af svovl, Bekendtgørelse 688 af 17/09/1998 (Danish legislation). Available at: <https://www.retsinformation.dk/forms/r0710.aspx?id=13266> (2014-02-14)

Danish Ministry of Taxation (DMT), 2008: Lov om afgift af kvælstofoxider. Lov nr 472 af 17/06/2008. Danish legislation. Available at: <https://www.retsinformation.dk/Forms/R0710.aspx?id=120340> (2013-01-24)

Elsam 2005: e-mail from Arne Henriksen, Elsam Engineering, 2005-08-11. Unpublished emission measurement data for N₂O from 3 power plants combusting coal.

EU Commission, 2007: COMMISSION DECISION of 18 July 2007 establishing guidelines for the monitoring and reporting of greenhouse gas emissions pursuant to Directive 2003/87/EC of the European Parliament and of the Council. Available at: <http://eurlex.europa.eu/LexUriServ/LexUriServ.do?uri=OJ:L:2007:229:0001:0085:EN:PDF> (2014-02-14).

European Environment Agency (EEA), 2007: EMEP/CORINAIR Atmospheric Emission Inventory Guidebook – 2007, prepared by the UNECE/-EMEP Task Force on Emissions Inventories and Projections. Technical Report No 16/2007. Available at: <http://www.eea.europa.eu/publications/EMEPCORINAIR5> (2013-02-13).

European Environment Agency (EEA), 2009: EMEP/EEA air pollutant emission inventory guidebook 2009. Technical guidance to prepare national emission inventories. EEA Technical Report 9/2009 Available at: <http://www.eea.europa.eu/publications/emep-eea-emission-inventory-guidebook-2009> (2012-01-27).

European Environment Agency (EEA), 2013: EMEP/EEA air pollutant emission inventory guidebook 2013. Available at: <http://www.eea.europa.eu/publications/emep-eea-guidebook-2013> (2014-02-14)

Fausser, P., Nielsen, M., Winther, M., Plejdrup, M., Gyldenkerne, S., Mikkelsen, M.H., Albrechtsen, R., Hoffmann, L., Thomsen, M., Hjelgaard, K. & Nielsen, O.-K. 2013. Verification of the Danish 1990, 2000 and 2010 emission inventory data. Aarhus University, DCE – Danish Centre for Environment and Energy, 85 pp. Scientific Report from DCE – Danish Centre for Environment and Energy No. 79. <http://dce2.au.dk/pub/SR79.pdf> (2014-02-14)

Fellner, J. & Rechberger, H. 2010: Biomass Content in the Waste Feed of Danish Waste-to-energy Plant, TU Wien, 2010. (Enclosed as Annex 10 in Astrup et al., 2012).

Fenhann, J. & Kilde, N.A. 1994: Inventory of Emissions to the air from Danish Sources 1972-1992, RISØ

Gruijthuisen, L.v. & Jensen, J.K., 2000: Energi- og miljøoversigt, Danish Gas Technology Centre 2000 (In Danish). Available at: <http://www.dgc.dk/publikation/2000/energi-og-miljoeversigt> (2014-02-14).

IPCC, 1997: Revised 1996 IPCC Guidelines for National Greenhouse Gas Inventories: Reference Manual (Volume 3). Available at: <http://www.ipcc-nggip.iges.or.jp/public/gl/invs6.html> (2014-02-14).

IPCC, 2000: Penman, J., Kruger, D., Galbally, I., Hiraishi, T., Nyenzi, B., Emmanuel, S., Buendia, L., Hoppaus, R., Martinsen, T., Meijer, J., Miwa, K. & Tanabe, K. (Eds). Good Practice Guidance and Uncertainty Management in National Greenhouse Gas Inventories. Published: IPCC/OECD/IEA/IGES, Hayama, Japan. Available at: <http://www.ipcc-nggip.iges.or.jp/public/gp/english/index.html> (2014-02-14).

IPCC, 2006: Revised 2006 IPCC Guidelines for National Greenhouse Gas Inventories: Available at: <http://www.ipcc-nggip.iges.or.jp/public/2006gl/index.html> (2014-02-14).

ISAG, 2012: Danish waste statistics for 2009, ISAG database. Available at: http://www.mst.dk/Virksomhed_og_myndighed/Affald/Tal_for_affald/S_tatistikker_og_ISAG-dataudtraek/ (2014-02-14).

Jensen, B.G. & Lindroth, M. 2002: Kontrol af indberetning af CO₂-udledning fra el-producenter i 2001, Carl Bro for Energistyrelsens 6. Kontor (in Danish).

Jensen, L. & Nielsen, P.A. 1990: Emissioner fra halm- og flisfyr, dk-Teknik & Levnedsmiddelstyrelsen 1990 (In Danish).

Jeppesen, J.S. 2008: København energi (Copenhagen Energy), Jørgen Steen Jeppesen, personal communication.

Johansson, L.S., Leckner, B., Gustavsson, L., Cooper, D., Tullin, C. & Potter, A. 2004: Emission characteristics of modern and old-type residential boilers fired with wood logs and wood pellets, Atmospheric Environment 38 (2004) 4183-4195.

Jørgensen, L. & Johansen, L.P. 2003: Eltra PSO 3141, Kortlægning af emissioner fra decentrale kraftvarmeværker, Anlæg A1-A5, dk-Teknik (In Danish). Available at: <http://energinet.dk/DA/KLIMA-OG-MILJOE/Emissionskortlægning/Sider/Maalerapporter-fra-emissionskortlaegningsprojektet.aspx> (2014-02-14).

Jørgensen, L., Andersen, S. D., Lovett, D. M. & Kristensen, P. G., 2010: Emissionskortlægning for decentral kraftvarme 2007, Måleprogram og analyse af emissioner fra gas- og oliefyrede decentrale kraftvarmeanlæg < 25 MW_e. Danish Gas Technology Centre. Available at: <http://www.energinet.dk/SiteCollectionDocuments/Danske%20dokumenter/Klimaogmiljo/Emissionsprojektet%20-%20Emissionsmålinger%20og%20analyse%20-%20Gas%20og%20olie.pdf> (2013-08-08)

Københavns Energi (KE), 2014: Københavns energi (Copenhagen Energy) fact sheet on town gas. Available at: <http://www.ke.dk/portal/pls/portal/docs/346012.PDF> (2014-02-14)

Kristensen, P.G. 2001: Personal communication, e-mail 10-04-2001, Danish Gas Technology Centre.

- Kristensen, P.G. 2007: Danish Gas Technology Centre, Per Gravers Kristensen, personal communication.
- Kristensen, P.G. 2010: Danish Gas Technology Centre, Per Gravers Kristensen, email 2010-02-10.
- Lindgren, 2010: Energinet.dk, Tine Lindgren, email 2010-03-16.
- Nielsen, M. & Wit, J. 1997: Emissionsforhold for gasdrevne kraftvarmeanlæg < 25MW_e, Miljøstyrelsen, Arbejdsrapport Nr. 17 1997 (In Danish). Available at: <http://www2.mst.dk/Udgiv/publikationer/1997/87-7810-759-8/pdf/87-7810-759-8.pdf> (2013-02-13).
- Nielsen, M. & Illerup, J.B. 2003: Emissionsfaktorer og emissionsopgørelse for decentral kraftvarme. Eltra PSO projekt 3141. Kortlægning af emissioner fra decentrale kraftvarmeværker. Delrapport 6. Danmarks Miljøundersøgelser. 116 s. –Faglig rapport fra DMU nr. 442. (In Danish, with an English summary). Available at: http://www2.dmu.dk/1_viden/2_Publikationer/3_fagrapporter/rapporter/FR442.pdf (2014-02-14).
- Nielsen, M. & Illerup, J.B. 2004: Danish emission inventories for stationary combustion plants. Inventories until year 2002. National Environmental Research Institute, Denmark. 127pp. – Research Notes from NERI no. 200. Available at: http://www2.dmu.dk/1_viden/2_Publikationer/3_arbrapporter/rapporter/AR200.pdf (2013-08-08)
- Nielsen, M. & Illerup, J.B. 2006: Danish emission inventories for stationary combustion plants. Inventories until year 2003. National Environmental Research Institute, Denmark. 162 pp. – Research Notes from NERI no. 229. Available at: http://www2.dmu.dk/1_viden/2_Publikationer/3_arbrapporter/rapporter/AR229.pdf (2014-02-14)
- Nielsen, O.-K., Nielsen, M. & Illerup, J.B. 2007: Danish Emission Inventories for Stationary Combustion Plants. Inventories until year 2004. National Environmental Research Institute, University of Aarhus, Denmark. 176 pp. – NERI Technical Report no. 628. Available at: http://www2.dmu.dk/Pub/FR628_Final.pdf (2014-02-14)
- Nielsen, M., Illerup, J.B. & Birr-Petersen, K., 2008: Revised emission factors for gas engines including start/stop emissions. Sub-report 3 (NERI). National Environmental Research Institute, University of Aarhus. 69 pp. – NERI Technical Report No. 672. Available at: <http://www.dmu.dk/Pub/FR672.pdf> (2014-02-14).
- Nielsen, M., Nielsen, O.-K., Plejdrup, M. & Hjelgaard, K., 2009: Danish Emission Inventories for Stationary Combustion Plants. Inventories until year 2007. National Environmental Research Institute, Aarhus University, Denmark. 216 pp. – NERI Technical Report no. 744. Available at: <http://www.dmu.dk/Pub/FR744.pdf> (2014-02-14).
- Nielsen, M., Nielsen, O.-K. & Thomsen, M. 2010(a): Emissions from decentralised CHP plants 2007 - Energinet.dk Environmental project no. 07/1882.

Project report 5 – Emission factors and emission inventory for decentralised CHP production. National Environmental Research Institute, Aarhus University. 113 pp. – NERI Technical report No. 786. Available at: <http://www.dmu.dk/Pub/FR786.pdf> (2014-02-14).

Nielsen, M., Nielsen, O.-K., Plejdrup, M. & Hjelgaard, K., 2010b: Danish Emission Inventories for Stationary Combustion Plants. Inventories until 2008. National Environmental Research Institute, Aarhus University, Denmark. 236 pp. – NERI Technical Report No. 795. Available at: <http://www.dmu.dk/Pub/FR795.pdf>. (2014-02-14)

Nielsen, O.-K., Plejdrup, M.S., Winther, M., Gyldenkerne, S., Thomsen, M., Fauser, P., Nielsen, M. Mikkelsen, M.H., Albrechtsen, R., Hjelgaard, K., Hoffmann, L. & Bruun, H.G. 2013. Quality manual for the Danish greenhouse gas inventory. Version 2. Aarhus University, DCE – Danish Centre for Environment and Energy, 44 pp. Scientific Report from DCE – Danish Centre for Environment and Energy No. 47. Available at: <http://www.dmu.dk/Pub/SR47.pdf> (2014-02-14)

Nielsen, M., Nielsen, O.-K. & Plejdrup, M., 2014: Danish Emission Inventories for Stationary Combustion Plants. Inventories until 2011. Aarhus University, DCE – Danish Centre for Environment and Energy.

Nikolaisen, L., Nielsen, C., Larsen, M.G., Nielsen, V. Zielke, U., Kristensen, J.K. & Holm-Christensen, B. 1998: Halm til energiformål, Teknik – Miljø – Økonomi, 2. udgave, 1998, Videncenter for halm og flisfyring (In Danish).

Olsson, M. & Kjällstrand, J. 2005: Low emissions from wood burning in an ecolabelled residential boiler, Atmospheric Environment 40 (2006) 1148-1158.

Paulrud, S., Kindbom, K., Cooper, K. & Gustafsson, T., 2005: Methane emissions from residential biomass combustion, SMED report 2005-10-14, IVL Swedish Environmental Research Institute and SCB, Statistics Sweden SMHI. Assignment for Swedish Environmental Protection Agency.

Pulles, T. & Aardenne, J.v., 2004: Good Practice Guidance for CLRTAP Emission Inventories, 24. Juni 2004. Available at: <http://www.eea.europa.eu/publications/EMEPCORINAIR4/BGPG.pdf> (2014-02-14).

Sander, B., 2002: Personal communication, email from Bo Sander, Elsam, 2002-05-17

Serup, H., Falster, H., Gamborg, C., Gundersen, P., Hansen, L. Heding, N., Jacobsen, H.H., Kofman, P., Nikolaisen, L. & Thomsen, I.M. 1999: Træ til energiformål, Teknik – Miljø – Økonomi, 2. udgave, 1999, Videncenter for halm og flisfyring (In Danish).

3.3 Transport and other mobile sources (CRF sector 1A2, 1A3, 1A4 and 1A5)

The emission inventory basis for mobile sources is fuel consumption information from the Danish energy statistics. In addition, background data for road transport (fleet and mileage), air traffic (aircraft type, flight numbers, origin and destination airports), national sea transport (fuel surveys, ferry technical data, number of return trips, sailing time) and non-road machinery (engine no., engine size, load factor and annual working hours) are used to make the emission estimates sufficiently detailed. Emission data mainly comes from the EMEP/EEA Air Pollutant Emission Inventory Guidebook (EMEP/EEA, 2013). However, for railways, measurements specific to Denmark are used.

In the Danish emissions database, all activity rates and emissions are defined in SNAP sector categories (Selected Nomenclature for Air Pollution) according to the CORINAIR system. The emission inventories are prepared from a complete emission database based on the SNAP sectors. The aggregation to the sector codes used for both the UNFCCC and UNECE Conventions is based on a correspondence list between SNAP and IPCC classification codes (CRF), shown in Table 3.3.1 (mobile sources only).

Table 3.3.1 SNAP – CRF correspondence table for transport.

SNAP classification	CRF/NFR classification
07 Road transport	1A3b Transport-Road
0801 Military	1A5 Other
0802 Railways	1A3c Railways
0803 Inland waterways	1A3d Transport-Navigation
080402 National sea traffic	1A3d Transport-Navigation
080403 National fishing	1A4c Agriculture/forestry/fisheries
080404 International sea traffic	1A3d Transport-Navigation (international)
080501 Dom. airport traffic (LTO < 1000 m)	1A3a Transport-Civil aviation
080502 Int. airport traffic (LTO < 1000 m)	1A3a Transport-Civil aviation (international)
080503 Dom. cruise traffic (> 1000 m)	1A3a Transport-Civil aviation
080504 Int. cruise traffic (> 1000 m)	1A3a Transport-Civil aviation (international)
0806 Agriculture	1A4c Agriculture/forestry/fisheries
0807 Forestry	1A4c Agriculture/forestry/fisheries
0808 Industry	1A2f Industry-Other
0809 Household and gardening	1A4b Residential
0811 Commercial and institutional	1A4a Commercial and institutional

Military transport activities (land and air) refer to the CRF/NFR sector Other (1A5), while the Transport-Navigation sector (1A3d) comprises national sea transport (SNAP code 080402, ship movements between two Danish ports) and recreational craft (SNAP code 0803).

For aviation, LTO (Landing and Take Off)¹ refers to the part of flying which is below 1000 m. This part of the aviation emissions (SNAP codes 080501 and 080502) are included in the national emissions total as prescribed by the UNECE reporting rules. According to UNFCCC the national emissions for aviation comprise the emissions from domestic LTO (080501) and domestic

¹ A LTO cycle consists of the flying modes approach/descent, taxiing, take off and climb out. In principle the actual times-in-modes rely on the actual traffic circumstances, the airport configuration, and the aircraft type in question.

cruise (080503). The fuel consumption and emission development explained in the following are based on these latter results.

Agricultural and forestry non-road machinery (SNAP codes 0806 and 0807) is accounted for in the Agriculture/forestry/fisheries (1A4c) sector together with fishing activities (SNAP code 080403).

For mobile sources, internal database models for road transport, air traffic, sea transport and non road machinery have been set up at Department of Environmental Science (ENVS)/Danish Centre for Environment and Energy (DCE), Aarhus University (former NERI), in order to produce the emission inventories. The output results from the DCE models are calculated in a SNAP format, as activity rates (fuel consumption) and emission factors, which are then exported directly to the central Danish CollectER database.

Apart from national inventories, the DCE models are used also as a calculation tool in research projects, environmental impact assessment studies, and to produce basic emission information which requires various aggregation levels.

3.3.1 Source category description

The following description of source categories explains the development in fuel consumption and emissions for road transport and other mobile sources.

Fuel consumption

Table 3.3.2 Fuel consumption (PJ) for domestic transport in 2012 in CRF sectors.

CRF ID	Fuel consumption (PJ)
Industry-Other (1A2f)	13,9
Civil Aviation (1A3a)	1,8
Road (1A3b)	160,9
Railways (1A3c)	3,4
Navigation (1A3d)	6,6
Comm./Inst. (1A4a)	2,3
Residential (1A4b)	0,9
Agri./for./fish. (1A4c)	24,9
Military (1A5)	1,6
Total	216.4

Table 3.3.2 shows the fuel consumption for domestic transport based on DEA statistics for 2012 in CRF sectors. The fuel consumption figures in time series 1990-2012 are given in Annex 3.B.16 (CRF format) and are shown for 1990 and 2012 in Annex 3.B.15 (CollectER format). Road transport has a major share of the fuel consumption for domestic transport. In 2012 this sector's fuel consumption share is 74 %, while the fuel consumption shares for Agriculture/forestry/fisheries and Industry-Other are 11 and 6 %, respectively. For the remaining sectors the total fuel consumption share is 9 %.

From 1990 to 2012, diesel (sum of diesel and biodiesel) and gasoline (sum of gasoline and E5) fuel consumption has changed by 43 % and - 12 %, respectively (Figure 3.3.1), and in 2012 the fuel consumption shares for diesel and gasoline were 66 % and 28 %, respectively. Other fuels only have a 6 % share of the domestic transport total (Figures 3.3.2). Almost all gasoline is used in road transportation vehicles. Gardening machinery and recreational craft are

merely small consumers. Regarding diesel, there is considerable fuel consumption in most of the domestic transport categories, whereas a more limited use of residual oil and jet fuel is being used in the navigation sector and by aviation (civil and military flights), respectively².




Figure 3.3.1 Fuel consumption pr fuel type for domestic transport 1990-2012.




Figure 3.3.2 Fuel consumption share pr fuel type for domestic transport in 2012.

Road transport

As shown in Figure 3.3.3, the fuel consumption for road transport³ has generally increased until 2007, except from a small fuel consumption decline noted in 2000. The impact of the global financial crisis on fuel consumption for road transport becomes visible for 2008 and 2009. The fuel consumption development is due to a decreasing trend in the use of gasoline fuels from 1999 onwards combined with a steady growth in the use of diesel until 2007. Within sub-sectors, passenger cars represent the most fuel-consuming vehicle category, followed by heavy-duty vehicles, light duty vehicles and 2-wheelers, in decreasing order (Figure 3.3.4).

² Biofuels are sold at gas filling stations and are assumed to be used by road transport vehicles.

³ The sum share of bioethanol and biodiesel in the gasoline and diesel fuel blends for road transport is 5.3 %, in 2012.




Figure 3.3.3 Fuel consumption pr fuel type and as totals for road transport 1990-2012.




Figure 3.3.4 Total fuel consumption pr vehicle type for road transport 1990-2012.

As shown in Figure 3.3.5, fuel consumption for gasoline passenger cars dominates the overall gasoline consumption trend. The development in diesel fuel consumption in recent years (Figure 3.3.6) is characterised by increasing fuel consumption for diesel passenger cars, while declines in the fuel consumption for trucks and buses (heavy-duty vehicles) and light duty vehicles are noted for 2008-2009, 2012, and 2008-2012, respectively.




Figure 3.3.5 Gasoline fuel consumption pr vehicle type for road transport 1990-2012.




Figure 3.3.6 Diesel fuel consumption pr vehicle type for road transport 1990-2012.

In 2012, fuel consumption shares for gasoline passenger cars, heavy-duty vehicles, diesel passenger cars, diesel light duty vehicles and gasoline light duty vehicles were 34, 26, 25, 14 and 1 %, respectively (Figure 3.3.7).




Figure 3.3.7 Fuel consumption share (PJ) pr vehicle type for road transport in 2012.

Other mobile sources

It must be noted that the fuel consumption figures behind the Danish inventory for mobile equipment in the agriculture, forestry, industry, household and gardening (residential), and inland waterways (part of navigation) sectors, are less certain than for other mobile sectors. For these types of machinery, the DEA statistical figures do not directly provide fuel consumption information, and fuel consumption totals are subsequently estimated from activity data and fuel consumption factors. For recreational craft the latest historical year is 2004.

As seen in Figure 3.3.8, classified according to CRF the most important sectors are Agriculture/forestry/fisheries (1A4c), Industry-other (mobile machinery part of 1A2f) and Navigation (1A3d). Minor fuel consuming sectors are Civil Aviation (1A3a), Railways (1A3c), Other (military mobile fuel consumption: 1A5), Commercial/institutional (1A4a) and Residential (1A4b).

The 1990-2012 time series are shown per fuel type in Figures 3.3.9-3.3.12 for diesel, gasoline and jet fuel, respectively.




Figure 3.3.8 Total fuel consumption in CRF sectors for other mobile sources 1990-2012.




Figure 3.3.9 Diesel fuel consumption in CRF sectors for other mobile sources 1990-2012.




Figure 3.3.10 Gasoline fuel consumption in CRF sectors for other mobile source 1990-2012.




Figure 3.3.11 Residual oil fuel consumption in CRF sectors for other mobile sources 1990-2012.




Figure 3.3.12 Jet fuel consumption in CRF sectors for other mobile sources 1990-2012.

In terms of diesel, the fuel consumption decreases for agricultural machines until 2000, due to fewer numbers of tractors and harvesters. After that, the increase in the engine sizes of new sold machines has more than outbalanced the trend towards smaller total stock numbers. The fuel consumption for in-

dustry has increased from the beginning of the 1990's, due to an increase in the activities for construction machinery. The fuel consumption increase has been very pronounced in 2005-2008, for 2009, however, the global financial crisis has a significant impact on the building and construction activities. For fisheries, the development in fuel consumption reflects the activities in this sector.

The Navigation sector comprises national sea transport (fuel consumption between two Danish ports including sea travel directly between Denmark and Greenland/Faroe Islands) and recreational craft. For the latter category, fuel consumption has increased significantly from 1990 to 2004 due to the rising number diesel-fuelled private boats. For national sea transport, the diesel fuel consumption curve reflects the combination of traffic and ferries in use for regional ferries. From 1998 to 2000, a significant decline in fuel consumption is apparent. The most important explanation here is the closing of ferry service routes in connection with the opening of the Great Belt Bridge in 1997. The fuel consumption decreases in 2011 and 2012 are due to reductions in the number of round trips made by regional ferries. For railways, the gradual shift towards electrification explains the lowering trend in diesel fuel consumption and the emissions for this transport sector. The fuel consumed (and associated emissions) to produce electricity is accounted for in the stationary combustion part of the Danish inventories.

The largest gasoline fuel consumption is found for household and gardening machinery in the Commercial/Institutional (1A4a) and Residential (1A4b) sectors. Especially from 2001-2006, a significant fuel consumption increase is apparent due to considerable growth in the machinery stock. The decline in gasoline fuel consumption for Agriculture/forestry/fisheries (1A4c) is due to the gradual phasing out of gasoline-fuelled agricultural tractors.

In terms of residual oil there has been a substantial decrease in the fuel consumption for regional ferries. The fuel consumption decline is most significant from 1990-1992 and from 1997-1999.

The considerable variations from one year to another in military jet fuel consumption are due to planning and budgetary reasons, and the passing demand for flying activities. Consequently, for some years, a certain amount of jet fuel stock-building might disturb the real picture of aircraft fuel consumption. Civil aviation has decreased until 2004, since the opening of the Great Belt Bridge in 1997, both in terms of number of flights and total jet fuel consumption. After 2004 an increase in the consumption of jet fuel is noted until 2007/2008.

Bunkers

The residual oil and diesel oil fuel consumption fluctuations reflect the quantity of fuel sold in Denmark to international ferries, international warships, other ships with foreign destinations, tank vessels and foreign fishing boats. For jet petrol, the sudden fuel consumption drop in 2002 is explained by the recession in the air traffic sector due to the events of September 11, 2001 and structural changes in the aviation business. In 2009, the impact of the global financial crisis on flying activities becomes very visible.




Figure 3.3.13 Bunker fuel consumption 1990-2012.

Emissions of CO₂, CH₄ and N₂O

In Table 3.3.3 the CO₂, CH₄ and N₂O emissions for road transport and other mobile sources are shown for 2012 in CRF sectors. The emission figures in time series 1990-2012 are given in Annex 3.B.16 (CRF format) and are shown for 1990 and 2012 in Annex 3.B.15 (CollectER format).

From 1990 to 2012 the road transport emissions of CO₂ and N₂O have increased by 21 and 28 %, respectively, whereas the emissions of CH₄ have decreased by 76 % (from Figures 3.3.14 - 3.3.16). From 1990 to 2012 the other mobile CO₂ emissions have decreased by 5 %, (from Figures 3.3.18 - 3.3.20).

Table 3.3.3 Emissions of CO₂, CH₄ and N₂O in 2012 for road transport and other mobile sources.

CRF Sector	CH ₄	CO ₂	N ₂ O
	Tonnes	Tonnes	Tonnes
Industry-Other (1A2f)	36	1 021	43
Civil Aviation (1A3a)	2	133	7
Railways (1A3c)	7	249	7
Navigation (1A3d)	34	498	29
Comm./Inst. (1A4a)	151	171	3
Residential (1A4b)	65	62	1
Ag./for./fish. (1A4c)	111	1 839	88
Military (1A5)	3	116	4
Total other mobile	409	4 090	181
Road (1A3b)	538	11 224	376
Total mobile	947	15 314	557

Road transport

CO₂ emissions are directly fuel consumption dependent and, in this way, the development in the emission reflects the trend in fuel consumption. As shown in Figure 3.3.14, the most important emission source for road transport is passenger cars, followed by heavy-duty vehicles, light-duty vehicles and 2-wheelers in decreasing order. In 2012, the respective emission shares were 60, 25, 15 and 0 %, respectively (Figure 3.3.17).

The majority of CH₄ emissions from road transport come from gasoline passenger cars (Figure 3.3.15). The emission drop from 1992 onwards is explained by the penetration of catalyst cars into the Danish fleet. The 2012 emission shares for CH₄ were 62, 18, 17 and 3 % for passenger cars, heavy-

duty vehicles, 2-wheelers and light-duty vehicles, respectively (Figure 3.3.17).




Figure 3.3.14 CO₂ emissions (k-tonnes) pr vehicle type for road transport 1990-2012.




Figure 3.3.15 CH₄ emissions (tonnes) pr vehicle type for road transport 1990-2012.

An undesirable environmental side effect of the introduction of catalyst cars is the increase in the emissions of N₂O from the first generation of catalyst cars (Euro 1) compared to conventional cars. The emission factors for later catalytic converter technologies are considerably lower than the ones for Euro 1, thus causing the emissions to decrease from 1998 onwards (Figure 3.3.16). In 2012, emission shares for passenger cars, heavy and light-duty vehicles were 54, 31 and 14 %, of the total road transport N₂O, respectively (Figure 3.3.17).

Referring to the second IPCC assessment report, 1 g CH₄ and 1 g N₂O has the greenhouse effect of 21 and 310 g CO₂, respectively. In spite of the relatively large CH₄ and N₂O global warming potentials, the largest contribution to the total CO₂ emission equivalents for road transport comes from CO₂, and the CO₂ emission equivalent shares per vehicle category are almost the same as the CO₂ shares.




Figure 3.3.16 N₂O emissions (tonnes) pr vehicle type for road transport 1990-2012.




Figure 3.3.17 CO₂, CH₄ and N₂O emission shares and GHG equivalent emission distribution for road transport in 2012.

Other mobile sources

For other mobile sources, the highest CO₂ emissions in 2012 come from Agriculture/forestry/fisheries (1A4c), Industry-other (1A2f) and Navigation (1A3d), with shares of 49, 25 and 12 %, respectively (Figure 3.3.21). The 1990-2012 emission trend is directly related to the fuel consumption development in the same time-period. Minor CO₂ emission contributors are sectors such as Commercial/Institutional (1A4a), Residential (1A4b), Railways (1A3c), Civil Aviation (1A3a) and Military (1A5).

For CH₄, far the most important sources are the gasoline fuelled gardening machinery in the Commercial/Institutional (1A4a) and Residential (1A4b) sectors, see Figure 3.3.21. The emission shares are 37 % and 16 %, respectively in 2012. The 2012 emission shares for Agriculture/forestry/fisheries (1A4c), Industry (1A2f) and Navigation (1A3d) are 27, 9 and 8 %, respectively, whereas the remaining sectors have emission shares of 3 % or less.




Figure 3.3.18 CO₂ emissions (ktonnes) in CRF sectors for other mobile sources 1990-2012.




Figure 3.3.19 CH₄ emissions (tonnes) in CRF sectors for other mobile sources 1990-2012.

For N₂O, the emission trend in sub-sectors is the same as for fuel consumption and CO₂ emissions (Figure 3.3.20).

As for road transport, CO₂ alone contributes with by far the most CO₂ emission equivalents in the case of other mobile sources, and per sector the CO₂ emission equivalent shares are almost the same as those for CO₂, itself (Figure 3.3.21).




Figure 3.3.20 N₂O emissions (tonnes) in CRF sectors for other mobile sources 1990-2012.




Figure 3.3.21 CO₂, CH₄ and N₂O emission shares and GHG equivalent emission distribution for other mobile sources in 2012.

Emissions of SO₂, NO_x, NMVOC and CO

In Table 3.3.4 the SO₂, NO_x, NMVOC and CO emissions for road transport and other mobile sources are shown for 2011 in CRF sectors. The emission figures in the time series 1990-2011 are given in Annex 3.B.16 (CRF format) and are shown for 1990 and 2011 in Annex 3.B.15 (CollectER format).

From 1990 to 2011, the road transport emissions of NMVOC, CO and NO_x emissions have decreased by 84, 77 and 58 %, respectively (Figures 3.3.23-3.3.25).

For other mobile sources, the emissions of NO_x decreased by 21 % from 1990 to 2011 and for SO₂ the emission drop is as much as 82 %. In the same period, the emissions of NMVOC have declined by 31 %, whereas the CO emissions have increased by 11 % (Figures 3.3.27-3.3.30).

Table 3.3.4 Emissions of SO₂, NO_x, NMVOC and CO in 2012 for road transport and other mobile sources.

CRF ID	SO ₂	NO _x	NMVOC	CO
	Tonnes	Tonnes	Tonnes	Tonnes
Industry-Other (1A2f)	6	7 597	1 076	6 239
Civil Aviation (1A3a)	42	579	98	607
Railways (1A3c)	2	2 531	190	425
Navigation (1A3d)	1 334	8 692	749	5 030
Comm./Inst. (1A4a)	1	219	3 636	72 587
Residential (1A4b)	0	92	1 953	26 236
Ag./for./fish. (1A4c)	311	18 163	2 137	19 246
Military (1A5)	17	492	33	270
Total other mobile	1 713	38 366	9 873	130 639
Road (1A3b)	71	44 316	10 439	96 022
Total mobile	1 784	82 682	20 312	226 661

Road transport

The step-wise lowering of the sulphur content in diesel fuel has given rise to a substantial decrease in the road transport emissions of SO₂ (Figure 3.3.22). In 1999, the sulphur content was reduced from 500 ppm to 50 ppm (reaching gasoline levels), and for both gasoline and diesel the sulphur content was reduced to 10 ppm in 2005. Since Danish diesel and gasoline fuels have the same sulphur percentages, at present, the 2012 shares for SO₂ emissions and fuel consumption for passenger cars, heavy-duty vehicles, light-duty vehicles and 2-wheelers are the same in each case: 60, 25, 15 and 0 %, respectively (Figure 3.3.26).




Figure 3.3.22 SO₂ emissions (tonnes) pr vehicle type for road transport 1990-2012.

Historically, the emission totals of NMVOC and CO have been very dominated by the contributions coming from private cars, as shown in Figures 3.3.24-3.3.25. However, the NMVOC and CO (and NO_x) emissions from this vehicle type have shown a steady decreasing tendency since the introduction of private catalyst cars in 1990 (EURO I) and the introduction of even more emission-efficient EURO II, III, IV and V private cars (introduced in 1997, 2001, 2006 and 2011, respectively).

In the case of NO_x, the real traffic emissions for heavy duty vehicles do not decline as intended by the EU emission legislation. This is due to the so-called engine cycle-beating effect. Outside the legislative test cycle stationary

measurement points, the electronic engine control for heavy duty Euro II and III engines switches to a fuel efficient engine running mode, thus leading to increasing NO_x emissions. However, the reduction in transport activities due to the global financial crisis causes the NO_x emissions for heavy duty vehicles to decrease significantly in 2008 and 2009.

The 2012 emission shares for heavy-duty vehicles, passenger cars, light-duty vehicles and 2-wheelers for NO_x (41, 44, 15 and 0 %), NMVOC (4, 62, 7 and 13 %) and CO (6, 80, 6 and 8 %), are also shown in Figure 3.3.26.




Figure 3.3.23 NO_x emissions (tonnes) pr vehicle type for road transport 1990-2012.




Figure 3.3.24 NMVOC emissions (tonnes) pr vehicle type for road transport 1990-2012




Figure 3.3.25 CO emissions (tonnes) pr vehicle type for road transport 1990-2012.




Figure 3.3.26 SO₂, NO_x, NMVOC and CO emission shares pr vehicle type for road transport in 2012

Other mobile sources

For SO₂ the trends in the Navigation (1A3d) emissions shown in Figure 3.3.27 mainly follow the development of the heavy fuel oil consumption (Figure 3.25). Though, from 1993 to 1995 relatively higher contents of sulphur in the fuel (estimated from sales) cause a significant increase in the emissions of SO₂. The SO₂ emissions for Fisheries (1A4c) correspond with the development in the consumption of marine gas oil. The main explanation for the development of the SO₂ emission curves for Railways (1A3c) and non-road machinery in Agriculture/forestry (1A4c) and Industry (1A2f), are the stepwise sulphur content reductions for diesel used by machinery in these sectors.




Figure 3.3.27 SO₂ emissions (tonnes) in CRF sectors for other mobile sources 1990-2012.

In general, the emissions of NO_x, NMVOC and CO from diesel-fuelled working equipment and machinery in agriculture, forestry and industry have decreased slightly since the end of the 1990s due to gradually strengthened emission standards given by the EU emission legislation directives. For industry, the emission impact from the global financial crisis becomes very visible for 2009.

NO_x emissions mainly come from diesel machinery, and the most important sources are Agriculture/forestry/fisheries (1A4c), Navigation (1A3d), Industry (1A2f) and Railways (1A3c), as shown in Figure 3.3.20. The 2012 emission shares are 48, 23, 20 and 7 %, respectively (Figure 3.3.23). Minor emissions come from the sectors, Civil Aviation (1A3a), Military (1A5) and Residential (1A4b).

The NO_x emission trend for Navigation, Fisheries and Agriculture is determined by fuel consumption fluctuations for these sectors, and the development of emission factors. For ship engines the emission factors tend to increase for new engines until mid-1990s. After that, the emission factors gradually reduce until 2000, bringing them to a level comparable with the emission limits for new engines in this year. For agricultural machines, there have been somewhat higher NO_x emission factors for 1991-stage I machinery, and an improved emission performance for stage I and II machinery since the late 1990s.




Figure 3.3.28 NO_x emissions (tonnes) in CRF sectors for other mobile sources 1990-2012.

The emission development for industry NO_x is the product of a fuel consumption increase from 1985 to 2008, most pronounced from 2005-2008, and a development in emission factors as explained for agricultural machinery. For railways, the gradual shift towards electrification explains the declining trend in diesel fuel consumption and NO_x emissions for this transport sector until 2001.

The 1990-2012 time series of NMVOC and CO emissions are shown in Figures 3.3.29 and 3.3.30 for other mobile sources. The 2012 sector emission shares are shown in Figure 3.3.31. For NMVOC, the most important sectors are Commercial/Institutional (1A4a), Agriculture/forestry/-fisheries (1A4c), Residential (1A4b), Industry (1A2f) and Navigation (1A3d) with 2012 emission shares of 37, 22, 20, 11 and 7 %, respectively. The same five sectors also contribute with most of the CO emissions. For Commercial/Institutional (1A4a), Residential (1A4b), Agriculture/forestry/fisheries (1A4c), Industry (1A2f) and Navigation (1A3d) the emission shares are 56, 20, 15, 5 and 4 %, respectively. Minor NMVOC and CO emissions come from Railways (1A3c), Civil Aviation (1A3a) and Military (1A5).

For NMVOC and CO, the significant emission increases for the commercial/-institutional and residential sectors after 2000 are due to the increased number of gasoline working machines. Improved NMVOC emission factors for diesel machinery in agriculture and gasoline equipment in forestry (chain saws) are the most important explanations for the NMVOC emission decline in the Agriculture/forestry/fisheries sector. This explanation also applies for the industrial sector, which is dominated by diesel-fuelled machinery. From 1997 onwards, the NMVOC emissions from Navigation decrease due to the gradually phase-out of the 2-stroke engine technology for recreational craft. The main reason for the significant 1985-2006 CO emission decrease for Agriculture/forestry/-fisheries is the phasing out of gasoline tractors.




Figure 3.3.29 NMVOC emissions (tonnes) in CRF sectors for other mobile sources 1990-2012.




Figure 3.3.30 CO emissions (tonnes) in CRF sectors for other mobile sources 1990-2012.




Figure 3.3.31 SO₂, NO_x, NMVOC and CO emission shares pr vehicle type for other mobile sources in 2012.

Bunkers

The most important emissions from bunker fuel consumption (fuel consumption for international transport) are SO₂, NO_x and CO₂ (and TSP, not shown). However, compared with the Danish national emission total (all sources), the greenhouse gas emissions from bunkers are small. The bunker emission totals are shown in Figure 3.3.7 for 2012, split into sea transport and civil aviation. All emission figures in the 1990-2012 time series are given in Annex 3.B.16 (CRF format). In Annex 3.B.15, the emissions are also given in CollectER format for the years 1990 and 2012.

Table 3.3.5 Emissions in 2012 for international transport.

CRF sector	SO ₂	NO _x	NMVOC	CH ₄	CO	CO ₂	N ₂ O
	Tonnes	Tonnes	Tonnes	Tonnes	Tonnes	K-tonnes	Tonnes
Navigation int. (1A3d)	5 136	36 524	1 199	37	3 955	1 505	95
Civil Aviation int. (1A3a)	801	10 628	379	12	1 776	2 510	86
International total	5 937	47 153	1 578	49	5 731	4 015	181

The differences in emissions between navigation and civil aviation are much larger than the differences in fuel consumption (and derived CO₂ emissions), and display a poor emission performance for international sea transport. In broad terms, the emission trends shown in Figure 3.3.32 are similar to the fuel consumption development.

However, for navigation minor differences occur for the emissions of SO₂, NO_x and CO₂ due to varying amounts of marine gas oil and residual oil, and for SO₂ and NO_x the development in the emission factors also have an impact on the emission trends. For civil aviation, apart from the annual consumption of jet fuel, the development of the NO_x emissions is also due to yearly variations in LTO/aircraft type (earlier than 2001) and city-pair statistics (2001 onwards).




Figure 3.3.32 CO₂, SO₂ and NO_x emissions for international transport 1990-2012.

3.3.2 Methodological issues

The description of methodologies and references for the transport part of the Danish inventory is given in two sections: one for road transport and one for the other mobile sources.

Methodology and references for Road Transport

For road transport, the detailed methodology is used to make annual estimates of the Danish emissions, as described in the EMEP/EEA Air Pollutant Emission Inventory Guidebook (EMEP/EEA, 2013). The actual calculations are made with a model developed by DCE, using the European COPERT IV model methodology explained by (EMEP/EEA, 2013). In COPERT, fuel consumption and emission simulations can be made for operationally hot engines, taking into account gradually stricter emission standards and emission degradation due to catalyst wear. Furthermore, the emission effects of cold-start and evaporation are simulated.

Vehicle fleet and mileage data

Corresponding to the COPERT IV fleet classification, all present and future vehicles in the Danish fleet are grouped into vehicle classes, sub-classes and layers. The layer classification is a further division of vehicle sub-classes into groups of vehicles with the same average fuel consumption and emission behaviour, according to EU emission legislation levels. Table 3.3.6 gives an overview of the different model classes and sub-classes, and the layer level with implementation years are shown in Annex 3.B.1.

Table 3.3.6 Model vehicle classes and sub-classes and trip speeds.

Vehicle clas-	Fuel type	Engine size/weight	Trip speed [km pr h]		
			Urban	Rural	Highway
PC	Gasoline	< 1.4 l.	40	70	100
PC	Gasoline	1.4 – 2 l.	40	70	100
PC	Gasoline	> 2 l.	40	70	100
PC	Diesel	< 2 l.	40	70	100
PC	Diesel	> 2 l.	40	70	100
PC	LPG		40	70	100
PC	2-stroke		40	70	100
LDV	Gasoline		40	65	80
LDV	Diesel		40	65	80
LDV	LPG		40	65	80
Trucks	Gasoline		35	60	80
Trucks	Diesel	Rigid 3,5 - 7,5t	35	60	80
Trucks	Diesel	Rigid 7,5 - 12t	35	60	80
Trucks	Diesel	Rigid 12 - 14 t	35	60	80
Trucks	Diesel	Rigid 14 - 20t	35	60	80
Trucks	Diesel	Rigid 20 - 26t	35	60	80
Trucks	Diesel	Rigid 26 - 28t	35	60	80
Trucks	Diesel	Rigid 28 - 32t	35	60	80
Trucks	Diesel	Rigid >32t	35	60	80
Trucks	Diesel	TT/AT 14 - 20t	35	60	80
Trucks	Diesel	TT/AT 20 - 28t	35	60	80
Trucks	Diesel	TT/AT 28 - 34t	35	60	80
Trucks	Diesel	TT/AT 34 - 40t	35	60	80
Trucks	Diesel	TT/AT 40 - 50t	35	60	80
Trucks	Diesel	TT/AT 50 - 60t	35	60	80
Trucks	Diesel	TT/AT >60t	35	60	80
Urban buses	Gasoline		30	50	70
Urban buses	Diesel	< 15 tonnes	30	50	70
Urban buses	Diesel	15-18 tonnes	30	50	70
Urban buses	Diesel	> 18 tonnes	30	50	70
Coaches	Gasoline		35	60	80

<i>Continued</i>					
Coaches	Diesel	< 15 tonnes	35	60	80
Coaches	Diesel	15-18 tonnes	35	60	80
Coaches	Diesel	> 18 tonnes	35	60	80
Mopeds	Gasoline		30	30	-
Motorcycles	Gasoline	2 stroke	40	70	100
Motorcycles	Gasoline	< 250 cc.	40	70	100
Motorcycles	Gasoline	250 – 750 cc.	40	70	100
Motorcycles	Gasoline	> 750 cc.	40	70	100

Fleet and annual mileage data are provided by DTU Transport for the vehicle categories present in COPERT IV (Jensen, 2013). DTU Transport use data from the Danish vehicle register kept by Statistics Denmark. The vehicle register data consist of vehicle type (passenger cars, vans, trucks, buses, mopeds, motorcycles), fuel type, vehicle weight, gross vehicle weight, engine size (passenger cars registered from 2005+), Euro class (trucks and buses registered from 1997+), NEDC type approval fuel efficiency value (passenger cars registered from 1997+) and vehicle first registration year.

In order to establish engine size data for passenger cars registered before 2005, a weight class-engine size transformation key is used examined by Cowi (2008) for new Danish cars from 1998. For the years before 1998, data for 1998 is used, and for the years 1999-2004 a linear interpolation between 1998 and 2005 weight class-engine size relations is used. For trucks, truck driver registration notes gathered by Statistics Denmark are used to split the fleet figures of ordinary trucks into number of solo trucks and truck-trailer combinations. Further the registration notes make it possible to assume the average total vehicle weight of the truck trailer combination. For articulated trucks also, the registration notes make it possible to assume the average total vehicle weight of the full articulated truck.

Danish mileage data comes from the Danish Road Directorate based on the Danish vehicle inspection programme. Total mileage per year and vehicle category are derived for the years 1985-2012, together with a more detailed mileage matrix examined for the year 2008 (based on detailed vehicle inspection data analysis). The detailed mileage matrix contains annual mileage per vehicle subcategory for new vehicles and for every vintage back in time, which determines the yearly mileage reduction percentages as a function of vehicle age. In a first step, the detailed mileage matrix is combined with corresponding fleet numbers in order to estimate intermediate total mileages for each year on a detailed fleet level. Next, each year's detailed (intermediate) mileage figures are scaled according to the difference between true and intermediate total mileage per vehicle subcategory.

DTU Transport (Jensen, 2013) also provides information of the mileage split between urban, rural and highway driving based on traffic monitoring data. The respective average speeds come from The Danish Road Directorate (Ekman, 2005). Additional data for the moped fleet and motorcycle fleet disaggregation is given by The National Motorcycle Association (Markamp, 2013).

In addition data from a survey made by the Danish Road Directorate (Hansen, 2010) has given information of the total mileage driven by foreign trucks on Danish roads in 2009. This mileage contribution has been added to the total mileage for Danish trucks on Danish roads, for trucks > 16 tonnes of gross vehicle weight. The data has been further processed by DTU

Transport; by using appropriate assumptions the mileage have been back-casted to 1985 and forecasted to 2012.




Figure 3.3.33 Number of vehicles in sub-classes in 1990-2012.

For passenger cars, the engine size differentiation is less certain for the years before 2005. The increase in the total number of passenger cars is mostly due to a growth in the number of gasoline cars with engine sizes between 1.4 and 2 litres (from 1990-2002) and diesel cars between 1.4 and 2 litres (from the 2000's up to now). Until 2005, there has been a decrease in the number of gasoline cars with an engine size smaller than 1.4 litres. These cars, however, have also increased in numbers during the later years. Since the late 1990's small cars (< 0.8 l gasoline and < 1.4 l. diesel) has slowly begun to penetrate the fleet.

There has been a considerable growth in the number of diesel light-duty vehicles from 1985 to 2006; the number of vehicles has, however, decreased somewhat after 2006.

For the truck-trailer and articulated truck combinations there is a tendency towards the use of increasingly larger trucks throughout the time period. The decline in fleet numbers for many of the truck categories in 2007/2008 and until 2009 is caused by the impact of the global financial crisis and the reflagging of Danish commercial trucks to companies based in the neighbouring countries.

The number of urban buses has been almost constant between 1985 and 2011. The sudden change in the level of coach numbers from 1994 to 1995 is due to uncertain fleet data.

The reason for the significant growth in the number of mopeds from 1994 to 2002 is the introduction of the so-called Moped 45 vehicle type. From 2004 onwards there is a gradual switch from 2-stroke to 4-stroke in new sales for this vehicle category. For motorcycles, the number of vehicles has grown in general throughout the entire 1985-2012 period. The increase is, however, most visible from the mid-1990s and onwards.

The vehicle numbers are summed up in EU emission layers for each year (Figure 3.3.34) by using the correspondence between layers and first year of registration:

$$N_{j,y} = \sum_{i=FYear(j)}^{LYear(j)} N_{i,y} \quad (1)$$

Where N = number of vehicles, j = layer, y = year, i = first year of registration.

Weighted annual mileages pr layer are calculated as the sum of all mileage driven pr first registration year divided by the total number of vehicles in the specific layer.

$$M_{j,y} = \frac{\sum_{i=FYear(j)}^{LYear(j)} N_{i,y} \cdot M_{i,y}}{\sum_{i=FYear(j)}^{LYear(j)} N_{i,y}} \quad (2)$$

Since 2006 economical incitements have been given to private vehicle owners to buy Euro 5 diesel passenger cars and vans in order to bring down the particulate emissions from diesel vehicles. The estimated sales between 2006 and 2010 have been examined by the Danish EPA and are included in the fleet data behind the Danish inventory (Winther, 2011).

For heavy duty trucks, there is a slight deviation from the strict correspondence between EU emission layers and first registration year.

In this case, specific Euro class information for most of the vehicles from 2001 onwards is incorporated into the fleet and mileage data model developed by Jensen (2012). For inventory years before 2001, and for vehicles with no Euro information the normal correspondence between layers and first year of registration is used.

Vehicle numbers and weighted annual mileages per layer are shown in Annex 3.B.1 and 3.B.2 for 1990-2012. The trends in vehicle numbers per layer are also shown in Figure 3.3.34. The latter figure shows how vehicles complying with the gradually stricter EU emission levels (EURO I, II, III, IV, V etc.) have been introduced into the Danish motor fleet.




Figure 3.3.34 Layer distribution of vehicle numbers pr vehicle type in 1990-2012.

Emission legislation

The EU 443/2009 regulation sets new emission performance standards for new passenger cars as part of the Community's integrated approach to reduce CO₂ emissions from light-duty vehicles. Some key elements of the adopted text are as follows:

- **Limit value curve:** the fleet average to be achieved by all cars registered in the EU is 130 grams per kilometre (g per km). A so-called limit value curve implies that heavier cars are allowed higher emissions than lighter cars while preserving the overall fleet average.
- **Further reduction:** A further reduction of 10 g CO₂ per km, or equivalent if technically necessary, will be delivered by other technological improvements and by an increased use of sustainable biofuels.
- **Phasing-in of requirements:** in 2012, 65% of each manufacturer's newly registered cars must comply on average with the limit value curve set by the legislation. This will rise to 75% in 2013, 80% in 2014, and 100% from 2015 onwards.
- **Lower penalty payments for small excess emissions until 2018:** If the average CO₂ emissions of a manufacturer's fleet exceed its limit value in any year from 2012, the manufacturer has to pay an excess emissions premium for each car registered. This premium amounts to €5 for the first g per km of exceedance, €15 for the second g per km, €25 for the third g per km, and €95 for each subsequent g per km. From 2019, already the first g per km of exceedance will cost €95.
- **Long-term target:** a target of 95 g pr km is specified for the year 2020. The modalities for reaching this target and the aspects of its implementation including the excess emissions premium will have to be defined in a review to be completed no later than the beginning of 2013.
- **Eco-innovations:** because the test procedure used for vehicle type approval is outdated, certain innovative technologies cannot demonstrate their CO₂ reducing effects under the type approval test. As an interim procedure until the test procedure is reviewed by 2014, manufacturers can be granted a maximum of 7 g per km of emission credits on average for their fleet if they equip vehicles with innovative technologies, based on independently verified data.

The EU 510/2011 regulation sets new emission performance standards for new light commercial vehicles (vans). Some key elements of the regulation are as follows:

- **Target dates:** The EU fleet average of 175 g CO₂ per km will be phased in between 2014 and 2017. In 2014 an average of 70 % of each manufacturer's newly registered vans must comply with the limit value curve set by the legislation. This proportion will rise to 75 % in 2015, 80 % in 2016, and 100% from 2017 onwards.
- **Limit value curve:** emissions limits are set according to the mass of vehicle, using a limit value curve. The curve is set in such a way that a fleet average of 175 grams of CO₂ per kilometre is achieved. A so-called limit value curve of 100% implies that heavier vans are allowed higher emissions than lighter vans while preserving the overall fleet average. Only the fleet average is regulated, so manufacturers will still be able to make vehicles with emissions above the limit value curve provided these are balanced by other vehicles which are below the curve.
- **Vehicles affected:** the vehicles affected by the legislation are vans, which account for around 12% of the market for light-duty vehicles. This includes vehicles used to carry goods weighing up to 3.5t (vans and car-derived vans, known as N1) and which weigh less than 2610 kg when empty.
- **Long-term target:** A target of 147g CO₂ per km is specified for the year 2020. This needs to be confirmed in a review of the vans Regulation, based on an updated assessment of its costs and benefits that is to be completed no later than the beginning of 2013. The modalities for reaching this target and aspects of its implementation, including the excess emissions premium, will also be defined as part of the review.
- **Excess emissions premium for small excess emissions until 2018:** If the average CO₂ emissions of a manufacturer's fleet exceed its limit value in any year from 2014, the manufacturer has to pay an excess emissions premium for each van registered. This premium amounts to €5 for the first g per km of exceedance, €15 for the second g per km, €25 for the third g per km, and €95 for each subsequent g per km. From 2019, the first g per km of exceedance will cost €95. This value is equivalent to the premium for passenger cars.
- **Super-credits:** vehicles with extremely low emissions (below 50g per km) will be given additional incentives whereby each low-emitting van will be counted as 3.5 vehicles in 2014 and 2015, 2.5 in 2016 and 1.5 vehicles in 2017.
- **Eco-innovations:** because the test procedure used for vehicle type approval is outdated, certain innovative technologies cannot demonstrate their CO₂-reducing effects under the type approval test. As an interim procedure until the test procedure is reviewed by 2014, manufacturers can be granted a maximum of 7g per km of emission credits on average for their fleet if they equip vehicles with innovative technologies, based on independently verified data.
- **Other flexibilities:** manufacturers may group together to form a pool and act jointly in meeting the specific emissions targets. Independent manufacturers who sell fewer than 22,000 vehicles per year can also apply to the Commission for an individual target instead.

For Euro 1-4 passenger cars and light duty trucks, the chassis dynamometer test cycle used in the EU for measuring fuel is the NEDC (New European Driving Cycle), see Nørgaard and Hansen (2004). The test cycle is also used also for emissions testing. The NEDC cycle consists of two parts, the first part being a 4-time repetition (driving length: 4 km) of the ECE test cycle. The latter test cycle is the so-called urban driving cycle⁴ (average speed: 19 km pr h). The second part of the test is the run-through of the EUDC (Extra Urban Driving Cycle) test driving segment, simulating the fuel consumption under rural and highway driving conditions. The driving length of EUDC is seven km at an average speed of 63 km pr h. More information regarding the fuel measurement procedure can be found in the EU-directive 80/1268/-EØF.

For NO_x, VOC (NMVOC + CH₄), CO and PM, the emissions from road transport vehicles have to comply with the different EU directives listed in Table 3.3.7. The emission directives distinguish between three vehicle classes according to vehicle reference mass⁵: Passenger cars and light duty trucks (<1305 kg), light duty trucks (1305-1760 kg) and light duty trucks (>1760 kg). The specific emission limits are shown in Annex 3.B.3.

⁴ For Euro 3 and on, the emission approval test procedure was slightly changed. The 40 s engine warm up phase before start of the urban driving cycle was removed.

⁵ Reference mass: net vehicle weight + mass of fuel and other liquids + 100 kg.

Table 3.3.7 Overview of the existing EU emission directives for road transport vehicles.

Vehicle category	Emission layer	EU directive	First reg. date
Passenger cars (gasoline)	PRE ECE	-	-
	ECE 15/00-01	70/220 - 74/290	1972 ^a
	ECE 15/02	77/102	1981 ^b
	ECE 15/03	78/665	1982 ^c
	ECE 15/04	83/351	1987 ^d
	Euro I	91/441	1.10.1990 ^e
	Euro II	94/12	1.1.1997
	Euro III	98/69	1.1.2001
	Euro IV	98/69	1.1.2006
	Euro V	715/2007	1.1.2011
	Euro VI	715/2007	1.9.2015
Passenger cars (diesel and LPG)	Conventional	-	-
	ECE 15/04	83/351	1987 ^d
	Euro I	91/441	1.10.1990 ^e
	Euro II	94/12	1.1.1997
	Euro III	98/69	1.1.2001
	Euro IV	98/69	1.1.2006
	Euro V	715/2007	1.1.2011
	Euro VI	715/2007	1.9.2015
Light duty trucks (gasoline and diesel)	Conventional	-	-
	ECE 15/00-01	70/220 - 74/290	1972 ^a
	ECE 15/02	77/102	1981 ^b
	ECE 15/03	78/665	1982 ^c
	ECE 15/04	83/351	1987 ^d
	Euro I	93/59	1.10.1994
	Euro II	96/69	1.10.1998
	Euro III	98/69	1.1.2002
	Euro IV	98/69	1.1.2007
	Euro V	715/2007	1.1.2012
	Euro VI	715/2007	1.9.2016
Heavy duty vehicles	Euro 0	88/77	1.10.1990
	Euro I	91/542	1.10.1993
	Euro II	91/542	1.10.1996
	Euro III	1999/96	1.10.2001
	Euro IV	1999/96	1.10.2006
	Euro V	1999/96	1.10.2009
	Euro VI	595/2009	1.10.2013
Mopeds	Conventional	-	-
	Euro I	97/24	2000
	Euro II	2002/51	2004
	Euro III	2002/51	2014 ^f
	Euro IV	168/2013	2017
	Euro V	168/2013	2021
Motor cycles	Conventional	Conventional	0
	Euro I	97/24	2000
	Euro II	2002/51	2004
	Euro III	2002/51	2007
	Euro IV	168/2013	2017
	Euro V	168/2013	2021

a,b,c,d: Expert judgement suggest that Danish vehicles enter into the traffic before EU directive first registration dates. The effective inventory starting years are a: 1970; b: 1979; c: 1981; d: 1986.e: The directive came into force in Denmark in 1991 (EU starting year: 1993).

In practice, the emissions from vehicles in traffic are different from the legislation limit values and, therefore, the latter figures are considered to be too inaccurate for total emission calculations. A major constraint is that the emission approval test conditions reflect only to a small degree the large variety of emission influencing factors in the real traffic situation, such as cumulated mileage driven, engine and exhaust after treatment maintenance levels and driving behaviour.

Therefore, in order to represent the Danish fleet and to support average national emission estimates, emission factors must be chosen which derive from numerous emissions measurements, using a broad range of real world driving patterns and a sufficient number of test vehicles. It is similar important to have separate fuel consumption and emission data for cold-start emission calculations and gasoline evaporation (hydrocarbons).

For heavy-duty vehicles (trucks and buses), the emission limits are given in g per kWh and the measurements are carried out for engines in a test bench, using the EU ESC (European Stationary Cycle) and ETC (European Transient Cycle) test cycles, depending on the Euro norm and exhaust gas after-treatment system installed. A description of the test cycles is given by Nørgaard and Hansen (2004). Measurement results in g per kWh from emission approval tests cannot be directly used for inventory work. Instead, emission factors used for national estimates must be transformed into g per km, and derived from a sufficient number of measurements which represent the different vehicle size classes, Euro engine levels and real world variations in driving behaviour.

Fuel consumption and emission factors

Trip-speed dependent basis factors for fuel consumption and emissions are taken from the COPERT model using trip speeds as shown in Table 3.3.6. The factors are listed in Annex 3.B.4. For EU emission levels not represented by actual data, the emission factors are scaled according to the reduction factors given in Annex 3.B.5.

The fuel consumption and emission factors used in the Danish inventory come from the COPERT IV model. The source for these data is various European measurement programmes. In general the COPERT data are transformed into trip-speed dependent fuel consumption and emission factors for all vehicle categories and layers.

For passenger cars, real measurement results are behind the emission factors for Euro 1-4 vehicles, and those earlier. For light duty trucks the measurements represent Euro 1 and prior vehicle technologies. For mopeds and motorcycles, updated fuel consumption and emission figures are behind the conventional and Euro 1-3 technologies. For heavy-duty trucks and buses the experimental basis is computer simulated emission factors for Euro 0-V engines.

Adjustment for fuel efficient vehicles

In the Danish fleet and mileage database, the type approval fuel efficiency value based on the NEDC driving cycle (TA_{NEDC}) is registered for each single car. Further, a modified fuel efficiency value (TA_{inuse}) is calculated using TA_{NEDC} , vehicle weight and engine size as input parameters. The TA_{inuse} value better reflects the fuel consumption associated with the NEDC driving cycle under real ("inuse") traffic conditions (Emisia, 2012).

From 2006 up to last historical year represented by fleet data, the average CO₂ emission factor (by fleet number) is calculated for each year's new sold cars, based on the registered TA_{NEDC} values. Using the average CO₂ emission factor for the last historical year as starting point, the average emission factor for each year's new sold cars are linearly reduced, until the emission factor reaches 95 g CO₂ per km in 2020.

From 2006 up to last historical year, the average CO₂ emission factor (by fleet number) is also calculated for each year's new sold cars, and for each fuel type/engine size combination, based on TA_{NEDC} and TA_{inuse}.

The linear reduction of the average emission factor for each year's new sold cars is then used to reduce the CO₂ emission factors for new sold cars based on TA_{inuse}, between last historical year and 2020, for each of the fuel type/engine size fleet segments.

Subsequently for each layer and inventory year, CO₂ emission factors are calculated based on TA_{inuse} and weighted by total mileage. On the same time corresponding layer specific CO₂ factors from COPERT IV are set up valid for Euro 4+ vehicles in the COPERT model. The COPERT IV CO₂ factors are derived from fuel consumption factors assessed by the developers of COPERT IV (Emisia, 2012) to represent the COPERT test vehicles under the NEDC driving cycle in real world traffic (TA_{COPERT IV, inuse}).

In a final step the ratio between the layer specific CO₂ emission factors for the Danish fleet and the COPERT Euro IV vehicles under TA_{inuse} are used to scale the trip speed dependent fuel consumption factors provided by COPERT IV for Euro 4 layers onwards.

Adjustment for EGR, SCR and filter retrofits

In COPERT IV updated emission factors have recently been made available for Euro V heavy duty vehicles using EGR and SCR exhaust emission after-treatment systems, respectively. The estimated new sales of Euro V diesel trucks equipped with EGR and SCR during the 2006-2010 time periods has been examined by Hjelgaard and Winther (2011). These inventory fleet data are used in the Danish inventory to calculate weighted emission factors for Euro V trucks in different size categories.

During the 2000's urban environmental zones have been established in Danish cities in order to bring down the particulate emissions from diesel fuelled heavy duty vehicles. Driving in these environmental zones prescribe the use of diesel particulate filters. The Danish EPA has provided the estimated number of Euro I-III urban buses and Euro II-III trucks and tourist buses which have been retrofitted with filters during the 2000's. These retrofit data are included in the Danish inventory by assuming that particulate emissions are lowered by 80 % compared with the emissions from the same Euro technology with no filter installed (Winther, 2011).

For all vehicle categories/technology levels not represented by measurements, the emission factors are produced by using reduction factors. The latter factors are determined by assessing the EU emission limits and the relevant emission approval test conditions, for each vehicle type and Euro class.

Deterioration factors

For three-way catalyst cars the emissions of NO_x, NMVOC and CO gradually increase due to catalyst wear and are, therefore, modified as a function of total mileage by the so-called deterioration factors. Even though the emission curves may be serrated for the individual vehicles, on average, the emissions from catalyst cars stabilise after a given cut-off mileage is reached due to OBD (On Board Diagnostics) and the Danish inspection and maintenance programme.

For each year, the deterioration factors are calculated per first registration year by using deterioration coefficients and cut-off mileages, as given in EMEP/EEA (2013), for the corresponding layer. The deterioration coefficients are given for the two driving cycles: "Urban Driving Cycle" (UDF) and "Extra Urban Driving Cycle" (EUDF: urban and rural), with trip speeds of 19 and 63 km per hour, respectively.

Firstly, the deterioration factors are calculated for the corresponding trip speeds of 19 and 63 km per hour in each case determined by the total cumulated mileage less than or exceeding the cut-off mileage. The Formulas 3 and 4 show the calculations for the "Urban Driving Cycle":

$$UDF = U_A \cdot MTC + U_B, MTC < U_{MAX} \quad (3)$$

$$UDF = U_A \cdot U_{MAX} + U_B, MTC \geq U_{MAX} \quad (4)$$

where UDF is the urban deterioration factor, U_A and U_B the urban deterioration coefficients, MTC = total cumulated mileage and U_{MAX} urban cut-off mileage.

In the case of trip speeds below 19 km per hour the deterioration factor, DF, equals UDF, whereas for trip speeds exceeding 63 km per hour, DF=EUDF. For trip speeds between 19 and 63 km per hour the deterioration factor, DF, is found as an interpolation between UDF and EUDF. Secondly, the deterioration factors, one for each of the three road types, are aggregated into layers by taking into account vehicle numbers and annual mileage levels per first registration year:

$$DF_{j,y} = \frac{\sum_{i=FYear(j)}^{LYear(j)} DF_{i,y} \cdot N_{i,y} \cdot M_{i,y}}{\sum_{i=FYear(j)}^{LYear(j)} DF_{i,y} \cdot N_{i,y}} \quad (5)$$

where DF is the deterioration factor.

For N₂O and NH₃, COPERT IV takes into account deterioration as a linear function of mileage for gasoline fuelled EURO 1-4 passenger cars and light duty vehicles. The level of emission deterioration also relies on the content of sulphur in the fuel. The deterioration coefficients are given in EMEP/EEA (2013), for the corresponding layer. A cut-off mileage of 250 000 km is behind the calculation of the modified emission factors, and for the Danish situation the low sulphur level interval is assumed to be most representative.

Emissions and fuel consumption for hot engines

Emissions and fuel consumption results for operationally hot engines are calculated for each year and for layer and road type. The procedure is to combine fuel consumption and emission factors (and deterioration factors for catalyst vehicles), number of vehicles, annual mileage levels and the relevant road-type shares given in Table 3.3.7. For non-catalyst vehicles this yields:

$$E_{j,k,y} = EF_{j,k,y} \cdot S_k \cdot N_{j,y} \cdot M_{j,y} \quad (6)$$

Here E = fuel consumption/emission, EF = fuel consumption/emission factor, S = road type share and k = road type.

For catalyst vehicles the calculation becomes:

$$E_{j,k,y} = DF_{j,k,y} \cdot EF_{j,k,y} \cdot S_k \cdot N_{j,y} \cdot M_{j,y} \quad (7)$$

Extra emissions and fuel consumption for cold engines

Extra emissions of NO_x, VOC, CH₄, CO, PM, N₂O, NH₃ and fuel consumption from cold start are simulated separately. For SO₂ and CO₂, the extra emissions are derived from the cold start fuel consumption results.

Each trip is associated with a certain cold-start emission level and is assumed to take place under urban driving conditions. The number of trips is distributed evenly across the months. First, cold emission factors are calculated as the hot emission factor times the cold:hot emission ratio. Secondly, the extra emission factor during cold start is found by subtracting the hot emission factor from the cold emission factor. Finally, this extra factor is applied on the fraction of the total mileage driven with a cold engine (the β -factor) for all vehicles in the specific layer.

The cold:hot ratios depend on the average trip length and the monthly ambient temperature distribution. The Danish temperatures for 2013 are given in Cappelen et al. (2012). For previous years, temperature data are taken from similar reports available from The Danish Meteorological Institute (www.dmi.dk). The cold:hot ratios are equivalent for gasoline fuelled conventional passenger cars and vans and for diesel passenger cars and vans, respectively, see EMEP/EEA (2013). For conventional gasoline and all diesel vehicles the extra emissions become:

$$CE_{j,y} = \beta \cdot N_{j,y} \cdot M_{j,y} \cdot EF_{U,j,y} \cdot (CEr - 1) \quad (8)$$

Where CE is the cold extra emissions, β = cold driven fraction, CEr = Cold:Hot ratio.

For catalyst cars, the cold:hot ratio is also trip speed dependent. The ratio is, however, unaffected by catalyst wear. The Euro I cold:hot ratio is used for all future catalyst technologies. However, in order to comply with gradually stricter emission standards, the catalyst light-off temperature must be reached in even shorter periods of time for future EURO standards. Correspondingly, the β -factor for gasoline vehicles is reduced step-wise for Euro II vehicles and their successors.

For catalyst vehicles the cold extra emissions are found from:

$$CE_{j,y} = \beta_{red} \cdot \beta_{EUROI} \cdot N_{j,y} \cdot M_{j,y} \cdot EF_{U,j,y} \cdot (CE_{EUROI} - 1) \quad (9)$$

where β_{red} = the β reduction factor.

For CH₄, specific emission factors for cold driven vehicles are included in COPERT IV. The β and β_{red} factors for VOC are used to calculate the cold driven fraction for each relevant vehicle layer. The NMVOC emissions during cold start are found as the difference between the calculated results for VOC and CH₄.

For N₂O and NH₃, specific cold start emission factors are also proposed by COPERT IV. For catalyst vehicles, however, just like in the case of hot emission factors, the emission factors for cold start are functions of cumulated mileage (emission deterioration). The level of emission deterioration also relies on the content of sulphur in the fuel. The deterioration coefficients are given in EMEP/EEA (2013), for the corresponding layer. For cold start, the cut-off mileage and sulphur level interval for hot engines are used, as described in the deterioration factors paragraph.

Evaporative emissions from gasoline vehicles

For each year, evaporative emissions of hydrocarbons are simulated in the forecast model as hot and warm running losses, hot and warm soak loss and diurnal emissions. The calculation approach is the same as in COPERT III. All emission types depend on RVP (Reid Vapour Pressure) and ambient temperature. The emission factors are shown in EMEP/EEA (2013).

Running loss emissions originate from vapour generated in the fuel tank while the vehicle is running. The distinction between hot and warm running loss emissions depends on engine temperature. In the model, hot and warm running losses occur for hot and cold engines, respectively. The emissions are calculated as annual mileage (broken down into cold and hot mileage totals using the β -factor) times the respective emission factors. For vehicles equipped with evaporation control (catalyst cars), the emission factors are only one tenth of the uncontrolled factors used for conventional gasoline vehicles.

$$R_{j,y} = N_{j,y} \cdot M_{j,y} \cdot ((1 - \beta) \cdot HR + \beta \cdot WR) \quad (10)$$

where R is running loss emissions and HR and WR are the hot and warm running loss emission factors, respectively.

In the model, hot and warm soak emissions for carburettor vehicles also occur for hot and cold engines, respectively. These emissions are calculated as number of trips (broken down into cold and hot trip numbers using the β -factor) times respective emission factors:

$$S_{j,y}^C = N_{j,y} \cdot \frac{M_{j,y}}{l_{trip}} \cdot ((1 - \beta) \cdot HS + \beta \cdot WS) \quad (11)$$

where S^C is the soak emission, l_{trip} = the average trip length, and HS and WS are the hot and warm soak emission factors, respectively. Since all catalyst vehicles are assumed to be carbon canister controlled, no soak emissions are estimated for this vehicle type. Average maximum and minimum temperatures per month are used in combination with diurnal emission factors to estimate the diurnal emissions from uncontrolled vehicles $E^d(U)$:

$$E_{j,y}^d(U) = 365 \cdot N_{j,y} \cdot e^d(U) \quad (12)$$

Each year's total is the sum of each layer's running loss, soak loss and diurnal emissions.

Fuel consumption balance

The calculated fuel consumption in COPERT IV must equal the statistical fuel sale totals according to the UNFCCC and UNECE emissions reporting format. The statistical fuel sales for road transport are derived from the Danish Energy Agency data (see DEA, 2013). The DEA data are further processed for gasoline in order to account for e.g. non road and recreational craft fuel consumption, which are not directly stated in the statistics, please refer to paragraph 3.3.3 for further information regarding the transformation of DEA fuel data.

The standard approach to achieve a fuel balance in annual emission inventories is to multiply the annual mileage with a fuel balance factor derived as the ratio between simulated and statistical fuel figures for gasoline and diesel, respectively. This method is also used in the present model.




Figure 3.3.35 DEA:DCE Fuel ratios (mileage adjustment factors) based on DEA fuel sales data and DCE fuel consumption estimates.




Figure 3.3.36 DEA:DCE Fuel ratios (mileage adjustment factors) based on DEA fuel consumption data and DCE fuel consumption estimates.

In Figure 3.3.35 and Figure 3.3.36 the COPERT IV:DEA gasoline and diesel fuel consumption ratios are shown for fuel sales and fuel consumption from 1990-2012. The data behind the figures are also listed in Annex 3.B.8. The fuel consumption figures are related to the traffic on Danish roads.

Per fuel type, all mileage numbers are equally scaled in order to obtain fuel equilibrium, and hence the mileage factors used are the reciprocal values of the COPERT IV:DEA fuel consumption: fuel sales ratio.

The reasons for the differences between DEA sales figures and bottom-up fuel estimates are mostly due to a combination of the uncertainties related to COPERT IV fuel consumption factors, allocation of vehicle numbers in sub-categories, annual mileage, trip speeds and mileage splits for urban, rural and highway driving conditions.

The final fuel consumption and emission factors per vehicle type are shown in Annex 3.B.7 for 1990-2012. The total fuel consumption and emissions are shown in Annex 3.B.8, per vehicle category and as grand totals, for 1990-2012 (and CRF format in Annex 3.B.16). In Annex 3.B.15, fuel consumption and emission factors as well as total emissions are given in CollectER format for 1990 and 2012.

In the following Figures 3.3.37 - 3.3.40, the fuel and km related emission factors for CO₂ (km related only), CH₄ and N₂O are shown per vehicle type for the Danish road transport (from 1990-2012).

For CO₂ the neat gasoline/diesel emission factors shown in Table 3.3.8 are country specific values, and come from the DEA. In 2006 and 2008, respectively, bio ethanol and biodiesel became available from a limited number of gas filling stations in Denmark, and today bio ethanol and biodiesel is added to all fuel commercially available. Following the IPCC guideline definitions, bio ethanol is regarded as CO₂ neutral for the transport sector as such. The sulphur content for bio ethanol/biodiesel is assumed to be zero, and hence, the aggregated CO₂ (and SO₂) factors for gasoline/ diesel have been adjusted, on the basis of the energy content of neat gasoline/diesel and bio ethanol/biodiesel, respectively, in the available fuels.

At present, the Danish road transport fuels only have low biofuel (BF) shares (Table 3.3.8), and hence, no thermal efficiency changes are expected for the fuels. Consequently, the energy based fuel consumption factors (MJ/km) derived from COPERT IV are used also in this case.

As a function of the current ethanol/biodiesel energy percentage, BF%_E, (Table 4.3) the average fuel related CO₂ emission factors, emf_{CO₂,E}(BF%) become:

$$EF_{CO_2,E}(BF\%) = EF_{CO_2,E}(BF0) \cdot (100 - BF\%_E) \quad (13)$$

Where:

EF_{CO₂,E}(BF%) = average fuel related CO₂ emission factor (g MJ⁻¹) for current BF%

EF_{CO₂,E}(BF0) = fuel related CO₂ emission factor (g MJ⁻¹) for fossil fuels

The kilometre based average CO₂ emission factor is subsequently calculated as the product of the fuel related CO₂ emission factor from equation 3 and the energy based fuel consumption factor, $FC_{CO_2,E}(BF0)$, derived from COPERT IV:

$$EF_{CO_2,km}(BF\%) = EF_{CO_2,E}(BF\%) \cdot FC_E(BF0) \quad (14)$$

A literature review carried out in the Danish research project REBECA revealed no significant changes in emission factors between neat gasoline and E5 gasoline-ethanol blends for the combustion related emission components; NO_x, CO and VOC (Winther et al., 2012). Hence, due to the current low ethanol content in today's road transport gasoline, no modifications of the neat gasoline based COPERT emission factors are made in the inventories in order to account for ethanol usage.

REBECA results published by Winther (2009) have shown that the emission impact of using diesel-biodiesel blends is very small at low biodiesel blend ratios. Consequently no bio fuel emission factor adjustments are needed for diesel vehicles as well. However, adjustment of the emission factors for diesel vehicles will be made if the biodiesel content of road transport diesel fuel increases to a more significant level in the future.

The fuel related CO₂ emission factors for neat gasoline/diesel, bio ethanol/biodiesel, and aggregated CO₂ factors are shown in Table 3.3.8.

Table 3.3.8 Fuel-specific CO₂ emission factors and biofuel shares for road transport in DK.

Fuel type	Emission factors (g/MJ)							
	1990-2005	2006	2007	2008	2009	2010	2011	2012
Neat gasoline	73	73	73	73	73	73	73	73
Neat diesel	74	74	74	74	74	74	74	74
LPG	63.1	63.1	63.1	63.1	63.1	63.1	63.1	63.1
Bio ethanol	0	0	0	0	0	0	0	0
Biodiesel	0	0	0	0	0	0	0	0
Gasoline, average	73	72.9	72.8	72.8	72.8	71.7	70.5	
Diesel, average	74	74	74	74	73.9	74	71.5	
Biofuel share (BF%) of Danish road transport fuels								
	1990-2005	2006	2007	2008	2009	2010	2011	2012
	0	0.09	0.14	0.13	0.21	0.69	3.40	5.30





Figure 3.3.37 Km related CO₂ emission factors per vehicle type for Danish road transport (1990-2012).




Figure 3.3.38 Fuel and km related CH₄ emission factors per vehicle type for Danish road transport (1990-2012).







Figure 3.3.39 Fuel and km related N₂O emission factors per vehicle type for Danish road transport (1990-2012).

Methodologies and references for other mobile sources

Other mobile sources are divided into several sub-sectors: sea transport, fishery, air traffic, railways, military, and working machinery and equipment in the sectors agriculture, forestry, industry and residential. The emission calculations are made using the detailed method as described in the EMEP/EEA air pollutant emission inventory guidebook (EMEP/EEA, 2013) for air traffic, off-road working machinery and equipment, and ferries, while for the remaining sectors the simple method is used.

3.3.3 Activity data

Air traffic

The activity data for air traffic consists of air traffic statistics provided by the Danish Transport Authority and Copenhagen Airport. Fuel statistics for jet fuel consumption and aviation gasoline are obtained from the Danish energy statistics (DEA, 2013).

For 2001 onwards, per flight records are provided by the Danish Transport Authority as data codes for aircraft type, and origin and destination airports (city-pairs).

Subsequently the aircraft types are separated by DCE into larger aircraft using jet fuel (jet engines, turbo props, helicopters) and small aircraft types with piston engines using aviation gasoline. This is done by using different aircraft dictionaries, internet look-ups and by communication with the Danish Transport Authority. Each of the larger aircraft type is then matched with a representative type for which fuel consumption and emission data are available from the EMEP/EEA databank. Relevant for this selection is aircraft maximum take off mass, engine types, and number of engines. A more thorough explanation is given in Winther (2001a, b).

Annex 3.B.10 shows the correspondence table between the actual aircraft type codes and representative aircraft types behind the Danish inventory. Annex 3.B.10 also show the number of LTO's per representative aircraft type for domestic and international flights starting from Copenhagen Airport and other airports, respectively⁶, in a time series from 2001-2012. The airport split is necessary to make due to the differences in LTO emission factors (c.f. section 3.3.4).

The same type of LTO activity data for the flights for Greenland and the Faroe Islands are shown in Annex 3.B.10 also, further detailed into an origin-destination airport matrix and having flight distances attached. This level of detail satisfies the demand from UNFCCC to provide precise documentation for the part of the inventory for the Kingdom of Denmark being outside the Danish mainland.

In the later years many flights in Denmark are being made by the new aircraft types CRJ9, E70, E170 and E175. These aircraft types are not represented by data in the EMEP/EEA databank. Instead new fuel consumption and emission factors have been calculated using fuel consumption and emission indexes from the ICAO Engine Exhaust Emission Database (www.caa.co.uk) for the CFM34-8C5 engine type which is installed in CRJ9, E70, E170 and E175. For LTO the fuel consumption and emission indexes are directly avail-

⁶ Excluding flights for Greenland and the Faroe Islands. These flights are separately listed in Annex 3.B.10.

able from the ICAO database. For cruise, distance related indexes are calculated by weighting the baseline CFM34-8C5 indexes with the development in distance related emission indexes for the B737 400 representative aircraft type taken from the EMEP/EEA database.

The ideal flying distance (great circle distance) between the city-pairs is calculated by DCE in a separate database. The calculation algorithm uses a global latitude/altitude coordinate table for airports. In cases when airport coordinates are not present in the DCE database, these are looked up on the internet and entered into the database accordingly.

For inventory years prior to 2001, detailed LTO/aircraft type statistics are obtained from Copenhagen Airport (for this airport only), while information of total take-off numbers for other Danish airports is provided by the Danish Transport Authority. The assignment of representative aircraft types for Copenhagen Airport is done as described above. For the remaining Danish airports representative aircraft types are not directly assigned. Instead appropriate average assumptions are made relating to the fuel consumption and emission data part.




Figure 3.3.40 Most frequent domestic flying routes for large aircraft in Denmark.

Copenhagen Airport is the starting or end point for most of the domestic aviation made by large aircraft in Denmark (Figure 3.3.40; routes to Greenland/Faroe Islands are not shown). Even though many domestic flights not touching Copenhagen Airport are also reported in the flight statistics kept by the Danish Transport Authority, these flights, however, are predominantly made with small piston engine aircraft using aviation gasoline. Hence, the consumption of jet fuel by flights not using Copenhagen is merely marginal.




Figure 3.3.41 No. of LTO's for the most important airports in Denmark 2001-2012.

Figure 3.3.41 shows the number of domestic and international LTO's for Danish airports⁷, in a time series from 2001-2012.

Non-road working machinery and equipment

Non-road working machinery and equipment are used in agriculture, forestry and industry, for household/gardening purposes and in inland waterways (recreational craft). Information on the number of different types of machines, their respective load factors, engine sizes and annual working hours has been provided by Winther et al. (2006) for the years until 2004. For later inventory years, supplementary stock data are annually provided by the Association of Danish Agricultural Machinery Dealers and the Association of Producers and Distributors of Fork Lifts in Denmark. The stock development from 1990-2012 for the most important types of machinery are shown in Figures 3.3.42 - 3.3.49. The stock data are also listed in Annex 3.B.11, together with figures for load factors, engine sizes and annual working hours. As regards stock data for the remaining machinery types, please refer to (Winther et al., 2006).

It is important to note that from key experts in the field of industrial non road activities a significant decrease in the activities is assumed for 2009 due to the global financial crisis. This reduction is in the order of 25 % for 2009 for industrial non road in general (pers. comm. Per Stjernqvist, Volvo Con-

⁷ Flights for Greenland and the Faroe Islands are included under domestic in the figure.

struction Equipment 2010). For fork lifts, 5 % and 20 % reductions are assumed for 2008 and 2009, respectively (pers. comm. Peter H. Møller, Rocla A/S).

For agriculture, the total number of agricultural tractors and harvesters per year are shown in the Figures 3.3.42 - 3.3.43, respectively. The figures clearly show a decrease in the number of small machines, these being replaced by machines in the large engine-size ranges.




Figure 3.3.42 Total numbers in kW classes for tractors from 1990 to 2012.




Figure 3.3.43 Total numbers in kW classes for harvesters from 1990 to 2012.

The tractor and harvester developments towards fewer vehicles and larger engines, shown in Figure 3.3.44, are very clear. From 1990 to 2012, tractor and harvester numbers decrease by around 22 % and 42 %, respectively, whereas the average increase in engine size for tractors is 35 % and 169 % for harvesters, in the same time period.




Figure 3.3.44 Total numbers and average engine size for tractors and harvesters (1990 to 2012).

The most important machinery types for industrial use are different types of construction machinery and fork lifts. The Figures 3.3.45 and 3.3.46 show the 1990-2012 stock development for specific types of construction machinery and diesel fork lifts. For most of the machinery types there is an increase in machinery numbers from 1990 onwards, due to increased construction activities. It is assumed that track type excavators/wheel type loaders (0-5 tonnes), and telescopic loaders first enter into use in 1991 and 1995, respectively.





Figure 3.3.45 1990-2012 stock development for specific types of construction machinery.




Figure 3.3.46 Total numbers of diesel fork lifts in kW classes from 1990 to 2012.

The emission level shares for tractors, harvesters, construction machinery and diesel fork lifts are shown in Figure 3.3.47, and present an overview of the penetration of the different pre-Euro engine classes, and engine stages complying with the gradually stricter EU stage I and II emission limits. The average lifetimes of 30, 25, 20 and 10 years for tractors, harvesters, fork lifts and construction machinery, respectively, influence the individual engine technology turn-over speeds.

The EU emission directive Stage I and II implementation years relate to engine size, and for all four machinery groups the emission level shares for the specific size segments will differ slightly from the picture shown in Figure 3.3.47. Due to scarce data for construction machinery, the emission level penetration rates are assumed to be linear and the general technology turnover pattern is as shown in Figure 3.3.47.




Figure 3.3.47 Emission level shares for tractors, harvesters, construction machinery and diesel fork lifts (1990 to 2012).

The 1990-2012 stock development for the most important household and gardening machinery types is shown in Figure 3.3.48.

For lawn movers and cultivators, the machinery stock remains approximately the same for all years. The stock figures for chain saws, shrub clearers, trimmers and hedge cutters increase from 1990 until 2004, and for riders this increase continues also after 2004. The yearly stock increases, in most cases, become larger after 2000. The lifetimes for gasoline machinery are short and, therefore, there new emission levels (not shown) penetrate rapidly.




Figure 3.3.48 Stock development 1990-2012 for the most important household and gardening machinery types.

Figure 3.3.49 shows the development in numbers of different recreational craft from 1990-2012. The 2004 stock data for recreational craft are repeated for 2005+, since no new fleet information has been obtained.

For diesel boats, increases in stock and engine size are expected during the whole period, except for the number of motor boats (< 27 ft.) and the engine sizes for sailing boats (<26 ft.), where the figures remain unchanged. A decrease in the total stock of sailing boats (<26 ft.) by 21 % and increases in the total stock of yawls/cabin boats and other boats (<20 ft.) by around 25 % are expected. Due to a lack of information specific to Denmark, the shifting rate from 2-stroke to 4-stroke gasoline engines is based on a German non-road study (IFEU, 2004).




Figure 3.3.49 1990-2012 Stock and engine size development for recreational craft.

National sea transport

A detailed methodology is used to estimate the fuel consumption figures for national sea transport, based on fleet activity estimates for regional ferries, local ferries and other national sea transport (Winther, 2008).

Table 3.3.9 lists the most important domestic ferry routes in Denmark in the period 1990-2012. For these ferry routes and the years 1990-2005, the following detailed traffic and technical data have been gathered by Winther (2008): Ferry name, year of service, engine size (MCR), engine type, fuel type, average load factor, auxiliary engine size and sailing time (single trip).

For 2006-2012, the above mentioned traffic and technical data for specific ferries have been provided by Kristensen (2013) in the case of Mols-Linien (Sjællands Odde-Ebeltoft, Sjællands Odde-Århus, Kalundborg-Århus), by Jørgensen (2013) for Færgen A/S (Køge-Rønne, Tårs-Spødsbjerg). For Esbjerg/Hanst-holm/Hirtshals-Torshavn traffic and technical data have been provided by Dávastovu (2011).

Table 3.3.9 Domestic ferry routes comprised in the Danish inventory.

Ferry service	Service period
Esbjerg-Torshavn	1990-1995, 2009+
Halsskov-Knudshoved	1990-1999
Hanstholm-Torshavn	1991-1992, 1999+
Hirtshals-Torshavn	2010
Hundested-Grenaa	1990-1996
Kalundborg-Juelsminde	1990-1996
Kalundborg-Samsø	1990+
Kalundborg-Århus	1990+
Korsør-Nyborg, DSB	1990-1997
Korsør-Nyborg, Vognmandsruten	1990-1999
København-Rønne	1990-2004
Køge-Rønne	2004+
Sjællands Odde-Ebeltoft	1990+
Sjællands Odde-Århus	1999+
Tårs-Spødsbjerg	1990+




Figure 3.3.50 Domestic regional ferry routes in Denmark (2012).

The number of round trips per ferry route from 1990 to 2011 is provided by Statistics Denmark (2013), see Figure 3.3.51 (Esbjerg/Hanstholm/Hirtshals-Torshavn not shown). The traffic data are also listed in Annex 3.B.12, together with different ferry specific technical and operational data.

For each ferry, Annex 3.B.12 lists the relevant information as regards ferry route, name, year of service, engine size (MCR), engine type, fuel type, average load factor, auxiliary engine size and sailing time (single trip).




Figure 3.3.51 No. of round trips for the most important ferry routes in Denmark 1990-2012.

It is seen from Table 3.3.9 (and Figure 3.3.51) that several ferry routes were closed in the time period from 1996-1998, mainly due to the opening of the Great Belt Bridge (connecting Zealand and Funen) in 1997. Hundested-Grenaa and Kalundborg-Juelsminde was closed in 1996, Korsør-Nyborg (DSB) closed in 1997, and Halsskov-Knudshoved and Korsør-Nyborg (Vognmandsruten) was closed in 1998. The ferry line København-Rønne was replaced by Køge-Rønne in 2004 and from 1999 a new ferry connection was opened between Sjællands Odde and Århus.

For the local ferries, a bottom-up estimate of fuel consumption for 1996 has been taken from the Danish work in Wismann (2001). The latter project calculated fuel consumption and emissions for all sea transport in Danish waters in 1995/1996 and 1999/2000. In order to cover the entire 1990-2012 inventory period, the fuel figure for 1996 has been adjusted according to the developments in local ferry route traffic shown in Annex 3.B.12.

Fuel sold for freight transport by Royal Arctic Line between Aalborg (Denmark) and Greenland and by Eim Skip - East route between Aarhus (Denmark) and Torshavn (Faroe Islands) are included under other national sea transport in the Danish inventories. In both cases all fuel is being bought in Denmark (Rasmussen, 2013 and Thorarensen, 2013).

For the remaining part of the traffic between two Danish ports, other national sea transport, bottom-up estimates for fuel consumption have been calculated for the years 1995 and 1999 by Wismann (2007). These fuel consumption estimates are used as activity data for the inventory years until 1995 and 1999 onwards. Interpolated figures are used for the inventory years 1996-1998.

The calculations use the database set up for Denmark in the Wismann (2001) study, with actual traffic data from the Lloyd's LMIS database (not including ferries). The database was split into three vessel types: bulk carriers, container ships, and general cargo ships; and five size classes: 0-1000, 1000-3000, 3000-10000, 10000-20000 and >20000 DTW. The calculations assume that bulk carriers and container ships use heavy fuel oil, and that general cargo ships use gas oil. For further information regarding activity data for local ferries and other national sea transport, please refer to Winther (2008).

The fleet activity data for regional ferries, and the fleet activity based fuel consumption estimates for local ferries and other national sea transport replace the fuel based activity data which originated directly from the DEA statistics.

Other sectors

The activity data for military, railways, international sea transport and fishery consists of fuel consumption information from DEA (2013). For international sea transport, the basis is fuel sold in Danish ports for vessels with a foreign destination, as prescribed by the IPCC guidelines.

However, it must be noted that fuel sold for sailing activities between Denmark and Greenland/Faroe Islands are reported as international in the DEA energy statistics. Hence, for inventory purposes in order to follow the IPCC guidelines the bottom-up fuel estimates for the ferry routes Esbjerg/Hanstholm-Torshavn, and fuel reports from Royal Arctic Line and Eim Skip is being subtracted from the fuel sales figures for international sea transport prior to inventory fuel input.

For fisheries, the calculation methodology described by Winther (2008) remains fuel based. However, the input fuel data differ from the fuel sales figures previously used. The changes are the result of further data processing of the DEA reported gas oil sales for national sea transport and fisheries, prior to inventory input. For years when the fleet activity estimates of fuel consumption for national sea transport (not including trips to Greenland/Faroe Islands) are smaller than DEA reported fuel sold for national sea transport, fuel is added to fisheries in the inventory. In the opposite case, fuel is being subtracted from the original DEA fisheries fuel sales figure in order to make up the final fuel consumption input for fisheries in the inventories.

The updated fuel consumption time series for national sea transport lead, in turn, to changes in the energy statistics for fisheries (gas oil) and industry (heavy fuel oil), so the national energy balance can remain unchanged.

For all sectors, fuel consumption figures are given in Annex 3.B.15 for the years 1990 and 2012 in CollectER format.

Emission legislation

For the engines used by other mobile sources, no legislative limits exist for specific fuel consumption. And no legislative limits exist for the emissions of CO₂ which are directly fuel dependent. The engines, however, do have to comply with the emission legislation limits agreed by the EU and, except for ships, the VOC emission limits influence the emissions of CH₄, these forming part of total VOC.

For non-road working machinery and equipment, and recreational craft and railway locomotives/motor cars, the emission directives list specific emission limit values (g per kWh) for CO, VOC, NO_x (or VOC + NO_x) and TSP, depending on engine size (kW for diesel, ccm for gasoline) and date of implementation (referring to engine market date).

For diesel, the directives 97/68 and 2004/26 relate to non-road machinery other than agricultural and forestry tractors, and the directives have different implementation dates for machinery operating under transient and constant loads. The latter directive also comprises emission limits for railway machinery. For tractors the relevant directives are 2000/25 and 2005/13. For gasoline, the directive 2002/88 distinguishes between hand-held (SH) and not hand-held (NS) types of machinery.

For engine type approval, the emissions (and fuel consumption) are measured using various test cycles (ISO 8178). Each test cycle consists of a number of measurement points for specific engine loads during constant operation. The specific test cycle used depends on the machinery type in question and the test cycles are described in more details in the directives.

Table 3.3.10 Overview of EU emission directives relevant for diesel fuelled non-road machinery.

Stage/ Engine size [kW]	CO	VOC	NO _x	VOC+NO _x	PM	Diesel machinery			Tractors	
	[g pr kWh]					EU Directive	Implement. date Transient	Implement. date Constant	EU directive	Implement. date
Stage I										
37<=P<75	6.5	1.3	9.2	-	0.85	97/68	1/4 1999	-	2000/25	1/7 2001
Stage II										
130<=P<560	3.5	1	6	-	0.2	97/68	1/1 2002	1/1 2007	2000/25	1/7 2002
75<=P<130	5	1	6	-	0.3		1/1 2003	1/1 2007		1/7 2003
37<=P<75	5	1.3	7	-	0.4		1/1 2004	1/1 2007		1/1 2004
18<=P<37	5.5	1.5	8	-	0.8		1/1 2001	1/1 2007		1/1 2002
Stage IIIA										
130<=P<560	3.5	-	-	4	0.2	2004/26	1/1 2006	1/1 2011	2005/13	1/1 2006
75<=P<130	5	-	-	4	0.3		1/1 2007	1/1 2011		1/1 2007
37<=P<75	5	-	-	4.7	0.4		1/1 2008	1/1 2012		1/1 2008
19<=P<37	5.5	-	-	7.5	0.6		1/1 2007	1/1 2011		1/1 2007
Stage IIIB										
130<=P<560	3.5	0.19	2	-	0.025	2004/26	1/1 2011	-	2005/13	1/1 2011
75<=P<130	5	0.19	3.3	-	0.025		1/1 2012	-		1/1 2012
56<=P<75	5	0.19	3.3	-	0.025		1/1 2012	-		1/1 2012
37<=P<56	5	-	-	4.7	0.025		1/1 2013	-		1/1 2013
Stage IV										
130<=P<560	3.5	0.19	0.4	-	0.025	2004/26	1/1 2014		2005/13	1/1 2014
56<=P<130	5	0.19	0.4	-	0.025		1/10 2014			1/10 2014

Table 3.3.11 Overview of the EU Emission Directive 2002/88 for gasoline fuelled non-road machinery.

	Catego- ry	Engine size [ccm]	CO [g pr kWh]	HC [g pr kWh]	NO _x [g pr kWh]	HC+NO _x [g pr kWh]	Implemen- tation date
Stage I							
Hand held	SH1	S<20	805	295	5.36	-	1/2 2005
	SH2	20=<S<50	805	241	5.36	-	1/2 2005
	SH3	50=<S	603	161	5.36	-	1/2 2005
Not hand held	SN3	100=<S<225	519	-	-	16.1	1/2 2005
	SN4	225=<S	519	-	-	13.4	1/2 2005
Stage II							
Hand held	SH1	S<20	805	-	-	50	1/2 2008
	SH2	20=<S<50	805	-	-	50	1/2 2008
	SH3	50=<S	603	-	-	72	1/2 2009
Not hand held	SN1	S<66	610	-	-	50	1/2 2005
	SN2	66=<S<100	610	-	-	40	1/2 2005
	SN3	100=<S<225	610	-	-	16.1	1/2 2008
	SN4	225=<S	610	-	-	12.1	1/2 2007

For recreational craft, Directive 2003/44 comprises the Stage 1 emission legislation limits for diesel engines, and for 2-stroke and 4-stroke gasoline engines, respectively. The CO and VOC emission limits depend on engine size (kW) and the inserted parameters presented in the calculation formulas in Table 3.3.12a. For NO_x, a constant limit value is given for each of the three engine types. For TSP, the constant emission limit regards diesel engines only.

In Table 3.3.12b the Stage II emission limits are shown for recreational craft. CO and HC+NO_x limits are provided for gasoline engines depending on the rated engine power and the engine type (stern-drive vs. outboard) while CO, HC+NO_x, and particulate emission limits are defined for Compression Ignition (CI) engines depending on the rated engine power and the swept volume.

Table 3.3.12a Overview of the EU Emission Directive 2003/44 for recreational craft.

Engine type	Impl. date	CO=A+B/P ⁿ			HC=A+B/P ⁿ			NO _x	TSP
		A	B	n	A	B	n		
2-stroke gasoline	1/1 2007	150.0	600.0	1.0	30.0	100.0	0.75	10.0	-
4-stroke gasoline	1/1 2006	150.0	600.0	1.0	6.0	50.0	0.75	15.0	-
Diesel	1/1 2006	5.0	0.0	0	1.5	2.0	0.5	9.8	1.0

Table 3.3.12b Overview of the EU Emission Directive 2013/53 for recreational craft.

Diesel engines					
Swept Volume, SV l/cyl.	Rated Engine Power, P _N kW	Impl. Date	CO g/kWh	HC + NO _x g/kWh	PM g/kWh
SV < 0.9	P _N < 37				
	37 ≤ P _N < 75 (*)	18/1 2017	5	4.7	0.30
	75 ≤ P _N < 3 700	18/1 2017	5	5.8	0.15
0.9 ≤ SV < 1.2	P _N < 3 700	18/1 2017	5	5.8	0.14
1.2 ≤ SV < 2.5		18/1 2017	5	5.8	0.12
2.5 ≤ SV < 3.5		18/1 2017	5	5.8	0.12
3.5 ≤ SV < 7.0		18/1 2017	5	5.8	0.11
Gasoline engines					
Engine type	Rated Engine Power, P _N kW		CO g/kWh	HC + NO _x g/kWh	PM g/kWh
Stern-drive and inboard engines	P _N ≤ 373	18/1 2017	75	5	-
	373 ≤ P _N ≤ 485	18/1 2017	350	16	-
	P _N > 485	18/1 2017	350	22	-
Outboard engines and PWC engines (**)	P _N ≤ 4.3	18/1 2017	500 – (5.0 x P _N)	15.7 + (50/PN ^{0.9})	-
	4.3 ≤ P _N ≤ 40	18/1 2017	500 – (5.0 x P _N)	15.7 + (50/PN ^{0.9})	-
	P _N > 40	18/1 2017	300		-

(*) Alternatively, this engine segment shall not exceed a PM limit of 0.2 g/kWh and a combined HC + NO_x limit of 5.8 g/kWh

(**) Small and medium size manufacturers making outboard engines ≤ 15 kW have until 18/1 2020 to comply

Table 3.2.13 Overview of the EU Emission Directive 2004/26 for railway locomotives and motorcars.

Engine size [kW]		CO [g pr kWh]	HC [g pr kWh]	NO _x [g pr kWh]	HC+NOX [g pr kWh]	PM [g pr kWh]	Implement. date
Locomotives Stage IIIA							
130 ≤ P < 560	RL A	3.5	-	-	4	0.2	1/1 2007
560 < P	RH A	3.5	0.5	6	-	0.2	1/1 2009
2000 ≤ P and piston displacement ≥ 5 l/cyl.	RH A	3.5	0.4	7.4	-	0.2	1/1 2009
Stage IIIB	RB	3.5	-	-	4	0.025	1/1 2012
Motor cars Stage IIIA							
130 < P	RC A	3.5	-	-	4	0.2	1/1 2006
Stage IIIB							
130 < P	RC B	3.5	0.19	2	-	0.025	1/1 2012

Aircraft engine emissions of NO_x, CO, VOC and smoke are regulated by ICAO (International Civil Aviation Organization). The engine emission certification standards are contained in Annex 16 – Environmental Protection, Volume II – Aircraft Engine Emissions to the Convention on International Civil Aviation (ICAO Annex 16, 1993). The emission standards relate to the total emissions (in grams) from the so-called LTO (Landing and Take Off) cycle divided by the rated engine thrust (kN). The ICAO LTO cycle contains the idealised aircraft movements below 3000 feet (915 m) during approach, landing, airport taxiing, take off and climb out.

For smoke all aircraft engines manufactured from 1 January 1983 have to meet the emission limits agreed by ICAO. For NO_x, CO, VOC The emission legislation is relevant for aircraft engines with a rated engine thrust larger

than 26.7 kN. In the case of CO and VOC, the ICAO regulations apply for engines manufactured from 1 January 1983.

For NO_x , the emission regulations fall in four categories:

- a) For engines of a type or model for which the date of manufacture of the first individual production model is on or before 31 December 1995, and for which the production date of the individual engine is on or before 31 December 1999.
- b) For engines of a type or model for which the date of manufacture of the first individual production model is after 31 December 1995, or for individual engines with a production date after 31 December 1999.
- c) For engines of a type or model for which the date of manufacture of the first individual production model is after 31 December 2003.
- d) For engines of a type or model for which the date of manufacture of the first individual production model is after 31 December 2007.

The regulations published by ICAO are given in the form of the total quantity of pollutants (D_p) emitted in the LTO cycle divided by the maximum sea level thrust (F_{oo}) and plotted against engine pressure ratio at maximum sea level thrust.

The limit values for NO_x are given by the formulas in Table 3.3.14.

Table 3.3.14 Current certification limits for NO_x for turbo jet and turbo fan engines.

	Engines first produced before 31.12.1995 & for engines manufactured up to 31.12.1999	Engines first produced after 31.12.1995 & for engines manufactured after 31.12.1999	Engines for which the date of manufacture of the first individual production model was after 31 December 2003	Engines for which the date of manufacture of the first individual production model was after 31 December 2007
Applies to engines >26.7 kN	$D_p/F_{oo} = 40 + 2\pi_{oo}$	$D_p/F_{oo} = 32 + 1.6\pi_{oo}$		
Engines of pressure ratio less than 30				
Thrust more than 89 kN			$D_p/F_{oo} = 19 + 1.6\pi_{oo}$	$D_p/F_{oo} = 16.72 + 1.4080\pi_{oo}$
Thrust between 26.7 kN and not more than 89 kN			$D_p/F_{oo} = 37.572 + 1.6\pi_{oo} - 0.208F_{oo}$	$D_p/F_{oo} = 38.54862 + (1.6823\pi_{oo}) - (0.2453F_{oo}) - (0.00308\pi_{oo}F_{oo})$
Engines of pressure ratio more than 30 and less than 62.5				
Thrust more than 89 kN			$D_p/F_{oo} = 7 + 2.0\pi_{oo}$	$D_p/F_{oo} = -1.04 + (2.0\pi_{oo})$
Thrust between 26.7 kN and not more than 89 kN			$D_p/F_{oo} = 42.71 + 1.4286\pi_{oo} - 0.4013F_{oo} + 0.00642\pi_{oo}F_{oo}$	$D_p/F_{oo} = 46.1600 + (1.4286\pi_{oo}) - (0.5303F_{oo}) - (0.00642\pi_{oo}F_{oo})$
Engines with pressure ratio 82.6 or more			$D_p/F_{oo} = 32 + 1.6\pi_{oo}$	$D_p/F_{oo} = 32 + 1.6\pi_{oo}$

Source: International Standards and Recommended Practices, Environmental Protection, ICAO Annex 16 Volume II Part III Paragraph 2.3.2, 2nd edition July 1993, plus amendments: Amendment 3 (20 March 1997), Amendment 4 (4 November 1999), Amendment 5 (24 November 2005).

where:

D_p = the sum of emissions in the LTO cycle in g

F_{oo} = thrust at sea level take-off (100 %)

π_{oo} = pressure ratio at sea level take-off thrust point (100 %)

The equivalent limits for HC and CO are $D_p/F_{oo} = 19.6$ for HC and $D_p/F_{oo} = 118$ for CO (ICAO Annex 16 Vol. II paragraph 2.2.2). Smoke is limited to a regulatory smoke number = $83 (F_{oo})^{-0.274}$ or a value of 50, whichever is the lower.

A further description of the technical definitions in relation to engine certification as well as actual engine exhaust emission measurement data can be found in the ICAO Engine Exhaust Emission Database. The latter database is accessible from <http://www.caa.co.uk>, hosted by the UK Civil Aviation Authority.

For seagoing vessels, NO_x emissions are regulated as explained in Marpol 73/78 Annex VI, formulated by IMO (International Maritime Organisation). The legislation is relevant for diesel engines with a power output higher than 130 kW, which are installed on a ship constructed on or after 1 January 2000 and diesel engines with a power output higher than 130 kW which undergo major conversion on or after 1 January 2000.

The NO_x emission limits for ship engines in relation to their rated engine speed (n) given in RPM (Revolutions Per Minute) are the following:

- 17 g pr kWh, $n < 130$ RPM.
- $45 \times n - 0.2$ g pr kWh, $130 \leq n < 2000$ RPM.

- 9,8 g pr kWh, $n \geq 2000$ RPM.

Further, the Marine Environment Protection Committee (MEPC) of IMO has approved proposed amendments to MARPOL Annex VI to be agreed by IMO in October 2008 in order to strengthen the emission standards for NO_x and the sulphur contents of heavy fuel oil used by ship engines.

For NO_x emission regulations, a three tiered approach is considered, which comprises the following:

- Tier I: Diesel engines (> 130 kW) installed on a ship constructed on or after 1 January 2000 and prior to 1 January 2011.
- Tier II: Diesel engines (> 130 kW) installed on a ship constructed on or after 1 January 2011.
- Tier III⁸: Diesel engines (> 130 kW) installed on a ship constructed on or after 1 January 2016.

As for the existing NO_x emission limits, the new Tier I-III NO_x legislation values rely on the rated engine speeds. The emission limit equations are shown in Table 3.3.15.

Table 3.3.15 Tier I-III NO_x emission limits for ship engines (amendments to MARPOL Annex VI).

	NO _x limit	RPM (n)
Tier I	17 g pr kWh	$n < 130$
	$45 \times n - 0.2$ g pr kWh	$130 \leq n < 2000$
	9,8 g pr kWh	$n \geq 2000$
Tier II	14.4 g pr kWh	$n < 130$
	$44 \times n - 0.23$ g pr kWh	$130 \leq n < 2000$
	7.7 g pr kWh	$n \geq 2000$
Tier III	3.4 g pr kWh	$n < 130$
	$9 \times n - 0.2$ g pr kWh	$130 \leq n < 2000$
	2 g pr kWh	$n \geq 2000$

The Tier I emission limits are identical with the existing emission limits from MARPOL Annex VI.

Also to be agreed by IMO in October 2008, the NO_x Tier I limits are to be applied for existing engines with a power output higher than 5000 kW and a displacement per cylinder at or above 90 litres, installed on a ship constructed on or after 1 January 1990 but prior to 1 January 2000.

In relation to the sulphur content in heavy fuel and marine gas oil used by ship engines, Table 3.3.16 shows the current legislation in force, and the amendment of MARPOL Annex VI to be agreed by IMO in October 2008.

⁸ For ships operating in a designated Emission Control Area. Outside a designated Emission Control Area, Tier II limits apply.

Table 3.3.16 Current legislation in relation to marine fuel quality.

Legislation		Heavy fuel oil		Gas oil	
		S- %	Impl. date (day/month/year)	S- %	Impl. date
EU-directive 93/12		None		0.2 ¹	1.10.1994
EU-directive 1999/32		None		0.2	1.1.2000
EU-directive 2005/33 ²	SECA - Baltic sea	1.5	11.08.2006	0.1	1.1.2008
	SECA - North sea	1.5	11.08.2007	0.1	1.1.2008
	Outside SECA's	None		0.1	1.1.2008
MARPOL Annex VI	SECA – Baltic sea	1.5	19.05.2006		
	SECA – North sea	1.5	21.11.2007		
	Outside SECA	4.5	19.05.2006		
MARPOL Annex VI amendments	SECA's	1	01.03.2010		
	SECA's	0.1	01.01.2015		
	Outside SECA's	3.5	01.01.2012		
	Outside SECA's	0.5	01.01.2020 ³		

¹ Sulphur content limit for fuel sold inside EU.

² From 1.1.2010 fuel with a sulphur content higher than 0.1 % must not be used in EU ports for ships at berth exceeding two hours

³ Subject to a feasibility review to be completed no later than 2018. If the conclusion of such a review becomes negative the effective date would default 1 January 2025.

For non road machinery, the EU directive 2003/17/EC gives a limit value of 10 ppm sulphur in diesel (from 2011).

Emission factors

The CO₂ emission factors are country-specific and come from the DEA. The N₂O emission factors are taken from the EMEP/EEA guidebook (EMEP/EEA, 2013).

For military ground material, aggregated CH₄ emission factors for gasoline and diesel are derived from the road traffic emission simulations. The CH₄ emission factors for railways are derived from specific Danish VOC measurements from the Danish State Railways (Delvig, 2013) and a NMVOC/CH₄ split, based on expert judgement.

For agriculture, forestry, industry, household gardening and inland waterways, the VOC emission factors are derived from various European measurement programmes and the current EU emission legislation; see IFEU (2004) and Winther et al. (2006). The NMVOC/CH₄ split is taken from USEPA (2004). The baseline emission factors are shown in Annex 3.B.10.

For national sea transport and fisheries, the VOC emission factors come from Trafikministeriet (2000). Specifically for the ferries used by Mols Linjen new VOC emission factors are provided by Kristensen (2008), originating from measurement results by Hansen et al. (2004), Wismann (1999) and PHP (1996). Kristensen (2013) has provided complimentary emission factor data for new ferries.

For ship engines VOC/CH₄ splits are taken from EMEP/EEA (2013), and all emission factors are shown in Annex 3.B.13.

The CH₄ emission factors for domestic aviation come from the EMEP/EEA (2013). For a number of different representative aircraft types, the

EMEP/EEA guidebook comprises fuel flow and NO_x, CO and VOC emission indices for the four LTO modes and distance based emission factors for cruise.

For all sectors, emission factors for the years 1990 and 2012 are given in CollectER format in Annex 3.B.15.

Table 3.3.17 shows the aggregated emission factors for CO₂, CH₄ and N₂O in 2012 used to calculate the emissions from other mobile sources in Denmark.

Table 3.3.17 Fuel-specific emission factors for CO₂, CH₄ and N₂O for other mobile sources in Denmark.

				Emission factors ⁹			
SNAP ID	CRF ID	Category	Fuel type	CH ₄ g pr GJ	CO ₂ g pr GJ	N ₂ O g pr GJ	
1A5	080100	Military	AvGas	21.90	73.00	2.00	
1A5	080100	Military	Diesel	1.14	74.00	2.81	
1A5	080100	Military	Gasoline	7.39	73.00	1.46	
1A5	080100	Military	Jet fuel	2.65	72.00	2.30	
1A3c	080200	Railways	Diesel	2.17	74.00	2.04	
1A3d	080300	Inland waterways	Diesel	2.55	74.00	2.97	
1A3d	080300	Inland waterways	Gasoline	64.41	73.00	1.55	
1A3d	080402	National sea traffic	Diesel	1.66	74.00	4.68	
1A3d	080402	National sea traffic	LPG	20.26	63.10	0.00	
1A3d	080402	National sea traffic	Residual oil	1.95	78.00	4.89	
1A4c	080403	Fishing	Diesel	1.79	74.00	4.68	
1A4c	080403	Fishing	LPG	20.26	63.10	0.00	
Memo item	080404	International sea traffic	Diesel	1.78	74.00	4.68	
Memo item	080404	International sea traffic	Residual oil	1.96	78.00	4.89	
1A3a	080501	Air traffic, Dom. < 3000 ft.	Other airports	AvGas	21.90	73.00	2.00
1A3a	080501	Air traffic, Dom. < 3000 ft.	Other airports	Jet fuel	1.82	72.00	9.24
Memo item	080502	Air traffic, Int. < 3000 ft.	Other airports	AvGas	21.90	73.00	2.00
Memo item	080502	Air traffic, Int. < 3000 ft.	Other airports	Jet fuel	4.24	72.00	7.24
1A3a	080503	Air traffic, Dom. > 3000 ft.	Other airports	Jet fuel	0.00	72.00	2.30
Memo item	080504	Air traffic, Int. > 3000 ft.	Other airports	Jet fuel	0.00	72.00	2.30
1A4c	080600	Agriculture	Diesel	0.80	74.00	3.19	
1A4c	080600	Agriculture	Gasoline	159.58	73.00	1.71	
1A4c	080700	Forestry	Diesel	0.42	74.00	3.21	
1A4c	080700	Forestry	Gasoline	30.97	73.00	0.46	
1A2f	080800	Industry	Diesel	0.88	74.00	3.10	
1A2f	080800	Industry	Gasoline	108.83	73.00	1.49	
1A2f	080800	Industry	LPG	7.69	63.10	3.50	
1A4b	080900	Household and gardening	Gasoline	75.45	73.00	1.26	
1A4a	081100	Commercial and institutional	Gasoline	64.42	73.00	1.13	
1A3a	080501	Air traffic, Dom. < 3000 ft.	Copenhagen airport	AvGas	21.90	73.00	2.00
1A3a	080501	Air traffic, Dom. < 3000 ft.	Copenhagen airport	Jet fuel	2.07	72.00	5.66
Memo item	080502	Air traffic, Int. < 3000 ft.	Copenhagen airport	AvGas	21.90	73.00	2.00
Memo item	080502	Air traffic, Int. < 3000 ft.	Copenhagen airport	Jet fuel	3.76	72.00	3.87
1A3a	080503	Air traffic, Dom. > 3000 ft.	Copenhagen airport	Jet fuel	0.00	72.00	2.30
Memo item	080504	Air traffic, Int. > 3000 ft.	Copenhagen airport	Jet fuel	0.00	72.00	2.30

⁹ References. CO₂: Country-specific. N₂O: EMEP/EEA. CH₄: Railways: DSB/DCE; Agriculture/Forestry/Industry/Household-Gardening: IFEU/USEPA; National sea traffic/Fishing/International sea traffic: Trafikministeriet/EMEP/EEA; domestic and international aviation: EMEP/EEA.

Factors for deterioration, transient loads and gasoline evaporation for non road machinery

The emission effects of engine wear are taken into account for diesel and gasoline engines by using the so-called deterioration factors. For diesel engines alone, transient factors are used in the calculations, to account for the emission changes caused by varying engine loads. The evaporative emissions of NMVOC are estimated for gasoline fuelling and tank evaporation. The factors for deterioration, transient loads and gasoline evaporation are taken from IFEU (2004), and are shown in Annex 3.B.10. For more details regarding the use of these factors, please refer to paragraph 3.3.4 or Winther et al. (2006).

3.3.4 Calculation method

Air traffic

For aviation, the domestic and international estimates are made separately for landing and take-off (LTOs < 3000 ft), and cruising (> 3000 ft).

By using the LTO mode specific fuel flow and emission indices from EMEP/EEA (2013), the fuel consumption and emission factors for the full LTO cycle can be estimated for each of the representative aircraft types used in the Danish inventory.

The fuel consumption for one LTO cycle is calculated according to the following sum formula:

$$FC_{LTO}^a = \sum_{m=1}^4 t_m \cdot ff_{a,m} \quad (15)$$

Where FC = fuel consumption (kg), m = LTO mode (approach/landing, taxiing, take off, climb out), t = times in mode (s), ff = fuel flow (kg per s), a = representative aircraft type.

The emissions for one LTO cycle are estimated as follows:

$$E_{LTO}^a = \sum_{m=1}^4 FC_{a,m} \cdot EI_{a,m} \quad (16)$$

Due to lack of specific airport data, for approach/descent, take off and climb out, standardised times-in-modes of 4, 0.7 and 2.2 minutes are used as defined by ICAO (ICAO, 1995), whereas for taxiing the appropriate time interval is 13 minutes in Copenhagen Airport and 5 minutes in other airports present in the Danish inventory.

For each representative aircraft type, the calculated fuel consumption and emission factors per LTO are shown in Annex 3.B.10 for Copenhagen Airport and other airports.

The calculations for cruise use the distance specific fuel consumption and emissions given by EMEP/EEA (2013) per representative aircraft type. Data interpolations or extrapolations are made – in each case determined by the great circle distance between the origin and the destination airports.

If the great circle distance, y, is smaller than the maximum distance for which fuel consumption and emission data are given in the EMEP/EEA data bank the fuel consumption or emission E (y) becomes:

$$E(y) = E_{x_i} + \frac{(y - x_i)}{x_{i+1} - x_i} \cdot (E_{x_{i+1}} - E_{x_i}) \quad y < x_{\max}, i = 0, 1, 2, \dots, \max-1 \quad (17)$$

In (15) x_i and x_{\max} denominate the separate distances and the maximum distance, respectively, with known fuel consumption and emissions. If the flight distance y exceeds x_{\max} the maximum figures for fuel consumption and emissions must be extrapolated and the equation then becomes:

$$E(y) = E_{x_{\max}} + \frac{(y - x_{\max})}{x_{\max} - x_{\max-1}} \cdot (E_{x_{\max}} - E_{x_{\max-1}}) \quad y > x_{\max} \quad (18)$$

Total results are summed up and categorised according to each flight's destination airport code in order to distinguish between domestic and international flights.

Annex 3.B.10 shows the average fuel consumption and emission factors per representative aircraft type for cruise flying, as well as total distance flown, for 2012¹⁰. The factors are split between Copenhagen Airport and other airports and distinguish between domestic and international flights.

Specifically for flights between Denmark and Greenland or the Faroe Islands, for each representative aircraft type, the flight distances are directly shown in Annex 3.B.10, which go into the cruise calculation expressions 17 and 18.

The overall fuel precision in the model is around 0.8, derived as the fuel ratio between model estimates and statistical sales. The fuel difference is accounted for by adjusting cruising fuel consumption and emissions in the model according to domestic and international cruising fuel shares.

Prior to 2001, the calculation procedure was first to estimate each year's fuel consumption and emissions for LTO. Secondly, total cruising fuel consumption was found year by year as the statistical fuel consumption total minus the calculated fuel consumption for LTO. Lastly, the cruising fuel consumption was split into a domestic and international part by using the results from a Danish city-pair emission inventory in 1998 (Winther, 2001a). For more details of this latter fuel allocation procedure, see Winther (2001b).

Non-road working machinery and recreational craft

Prior to adjustments for deterioration effects and transient engine operations, the fuel consumption and emissions in year X , for a given machinery type, engine size and engine age, are calculated as:

$$E_{Basis}(X)_{i,j,k} = N_{i,j,k} \cdot HRS_{i,j,k} \cdot P \cdot LF_i \cdot EF_{y,z} \quad (19)$$

where E_{Basis} = fuel consumption/emissions in the basic situation, N = number of engines, HRS = annual working hours, P = average rated engine size in kW, LF = load factor, EF = fuel consumption/emission factor in g pr kWh, i = machinery type, j = engine size, k = engine age, y = engine-size class and z = emission level. The basic fuel consumption and emission factors are shown in Annex 3.B.11.

The deterioration factor for a given machinery type, engine size and engine age in year X depends on the engine-size class (only for gasoline), y , and the

¹⁰ Excluding flights for Greenland and the Faroe Islands.

emission level, z. The deterioration factors for diesel and gasoline 2-stroke engines are found from:

$$DF_{i,j,k}(X) = \frac{K_{i,j,k}}{LT_i} \cdot DF_{y,z} \quad (20)$$

where DF = deterioration factor, K = engine age, LT = lifetime, i = machinery type, j = engine size, k = engine age, y = engine-size class and z = emission level.

For gasoline 4-stroke engines the deterioration factors are calculated as:

$$DF_{i,j,k}(X) = \sqrt{\frac{K_{i,j,k}}{LT_i}} \cdot DF_{y,z} \quad (21)$$

The deterioration factors inserted in (20) and (21) are shown in Annex 3.B.11. No deterioration is assumed for fuel consumption (all fuel types) or for LPG engine emissions and, hence, DF = 1 in these situations.

The transient factor for a given machinery type, engine size and engine age in year X, relies only on emission level and load factor, and is denominated as:

$$TF_{i,j,k}(X) = TF_z \quad (22)$$

Where i = machinery type, j = engine size, k = engine age and z = emission level.

The transient factors inserted in (20) are shown in Annex 3.B.11. No transient corrections are made for gasoline and LPG engines and, hence, $TF_z = 1$ for these fuel types.

The final calculation of fuel consumption and emissions in year X for a given machinery type, engine size and engine age, is the product of the expressions 17-20:

$$E(X)_{i,j,k} = E_{Basis}(X)_{i,j,k} \cdot TF(X)_{i,j,k} \cdot (1 + DF(X)_{i,j,k}) \quad (23)$$

The evaporative hydrocarbon emissions from fuelling are calculated as:

$$E_{Evap, fueling, i} = FC_i \cdot EF_{Evap, fueling} \quad (24)$$

Where $E_{Evap, fueling, i}$ = hydrocarbon emissions from fuelling, i = machinery type, FC = fuel consumption in kg, $EF_{Evap, fueling}$ = emission factor in g NMVOC pr kg fuel.

For tank evaporation, the hydrocarbon emissions are found from:

$$E_{Evap, tank, i} = N_i \cdot EF_{Evap, tank, i} \quad (25)$$

Where $E_{Evap, tank, i}$ = hydrocarbon emissions from tank evaporation, N = number of engines, i = machinery type and $EF_{Evap, fueling}$ = emission factor in g NMVOC pr year.

Ferries, other national sea transport and fisheries

The fuel consumption and emissions in year X, for regional ferries are calculated as:

$$E(X) = \sum_i N_i \cdot T_i \cdot S_{i,j} \cdot P_i \cdot LF_j \cdot EF_{k,l,y} \quad (26)$$

Where E = fuel consumption/emissions, N = number of round trips, T = sailing time pr round trip in hours, S = ferry share of ferry service round trips, P = engine size in kW, LF = engine load factor, EF = fuel consumption/emission factor in g pr kWh, i = ferry service, j = ferry, k = fuel type, l = engine type, y = engine year.

For the remaining navigation categories, the emissions are calculated using a simplified approach:

$$E(X) = \sum_i EC_{i,k} EF_{k,l,y} \quad (27)$$

Where E = fuel consumption/emissions, EC = energy consumption, EF = fuel consumption/emission factor in g per kg fuel, i = category (local ferries, other national sea, fishery, international sea), k = fuel type, l = engine type, y = average engine year.

The emission factor inserted in (27) is found as an average of the emission factors representing the engine ages which are comprised by the average lifetime in a given calculation year, X:

$$EF_{k,l,y} = \frac{\sum_{year=X-LT}^{year=X} EF_{k,l}}{LT_{k,l}} \quad (28)$$

Other sectors

For military and railways, the emissions are estimated with the simple method using fuel-related emission factors and fuel consumption from the DEA:

$$E = FC \cdot EF \quad (29)$$

where E = emission, FC = fuel consumption and EF = emission factor. The calculated emissions for other mobile sources are shown in CollectER format in Annex 3.B.16 for the years 1990 and 2012 and as time series 1990-2012 in Annex 3.B.15 (CRF format).

Fuel balance between DEA statistics and inventory estimates

Following convention rules, the DEA statistical fuel sales figures are the basis for the full Danish inventory. However, in some cases for mobile sources the DEA statistical sectors do not fully match the inventory sectors. This is the case for non road machinery, where relevant DEA statistical sectors also include fuel consumed by stationary sources.

In other situations, fuel consumption figures estimated by DCE from specific bottom-up calculations are regarded as more reliable than DEA reported sales. This is the case for national sea transport.

In the following the transferral of fuel consumption data from DEA statistics into inventory relevant categories is explained for national sea transport and fisheries, non road machinery and recreational craft, and road transport. A full list of all fuel consumption data, DEA figures as well as intermediate

fuel consumption data, and final inventory input figures is shown in Annex 3.B.14.

National sea transport and fisheries

For national sea transport in Denmark, the fuel consumption estimates obtained by DCE (see 3.3.3 Activity data – national sea transport) are regarded as much more accurate than the DEA fuel sales data, since the large fluctuations in reported fuel sales cannot be explained by the actual development in the traffic between different national ports. As a consequence, the new bottom-up estimates replace the previous fuel based figures for national sea transport.

There are different potential reasons for the differences between estimated fuel consumption and reported sales for national sea transport in Denmark. According to the DEA, the latter fuel differences are most likely explained by inaccurate customer specifications made by the oil suppliers. This inaccuracy can be caused by a sector misallocation in the sales statistics between national sea transport and fisheries for gas oil, and between national sea transport and industry for heavy fuel oil (Peter Dal, DEA, personal communication, 2007). Further, fuel sold for vessels sailing between Denmark and Greenland/Faroe Islands are reported as international in the DEA statistics, and this fuel categorisation is different from the IPCC guideline definitions (see following paragraph “Bunkers”).

Following this, for fisheries and industry the updated fuel consumption time series for national sea transport lead, in turn, to changes in the fuel activity data for fisheries (gas oil), industry (heavy fuel oil) and international sea transport, so the national energy balance can remain unchanged.

For fisheries, fuel investigations made prior to the initiation of the work made by Winther (2008) have actually pointed out a certain area of inaccuracy in the DEA statistics. No engines installed in fishing vessels use heavy fuel oil, even though a certain amount of heavy fuel oil is listed in the DEA numbers for some statistical years (H. Amdissen, Danish Fishermen's Association, personal communication, 2006). Hence, for fisheries small amounts of fuel oil are transferred to national sea transport, and in addition small amounts of gasoline and diesel are transferred to recreational craft.

Non road machinery and recreational craft

For diesel and LPG, the non-road fuel consumption estimated by DCE is partly covered by the fuel consumption amounts in the following DEA sectors: agriculture and forestry, market gardening, and building and construction. The remaining quantity of non-road diesel and LPG is taken from the DEA industry sector.

For gasoline, the DEA residential sector, together with the DEA sectors mentioned for diesel and LPG, contribute to the non-road fuel consumption total. In addition, a certain amount of fuel from road transport is needed to reach the fuel consumption goal.

The amount of diesel and LPG in DEA industry not being used by non-road machinery is included in the sectors, “Combustion in manufacturing industry” (0301) and “Non-industrial combustion plants” (0203) in the Danish emission inventory.

For recreational craft, the calculated fuel consumption totals for diesel and gasoline are subsequently subtracted from the DEA fishery sector. For gasoline, the DEA reported fuel consumption for fisheries is far too small to fill the fuel gap, and hence the missing fuel amount is taken from the DEA road transport sector.

Bunkers

The distinction between domestic and international emissions from aviation and navigation should be in accordance with the Revised 1996 IPCC Guidelines for National Greenhouse Gas Inventories. For the national emission inventory, this, in principle, means that fuel sold (and associated emissions) for flights/sea transportation starting from a seaport/airport in the Kingdom of Denmark, with destinations inside or outside the Kingdom of Denmark, are regarded as domestic or international, respectively.

Aviation

As prescribed by the IPCC guidelines, for aviation, the fuel consumption and emissions associated with flights inside the Kingdom of Denmark are counted as domestic.

This report includes flights from airports in Denmark and associated jet fuel sales. Hence, the flights between airports in Denmark and flights from Denmark to Greenland and the Faroe Islands are classified as domestic and flights from Danish airports with destinations outside the Kingdom of Denmark are classified as international flights.

In Greenland and in the Faroe Islands, the jet fuel sold is treated as domestic. This decision becomes reasonable when considering that almost no fuel is bunkered in Greenland/the Faroe Islands by flights other than those going to Denmark.

Navigation

In DEA statistics, the domestic fuel total consists of fuel sold to Danish ferries and other ships sailing between two Danish ports. The DEA international fuel total consists of the fuel sold in Denmark to international ferries, international warships, other ships with foreign destinations, transport to Greenland and the Faroe Islands, tank vessels and foreign fishing boats.

In order to follow the IPCC guidelines the bottom-up fuel estimates for the ferry routes between Denmark and the Faroe Islands, and freight transport between Denmark and Greenland/Faroe Islands are being subtracted from the fuel sales figures for international sea transport prior to inventory fuel input.

In Greenland, all marine fuel sales are treated as domestic. In the Faroe Islands, fuel sold in Faroese ports for Faroese fishing vessels and other Faroese ships is treated as domestic. The fuel sold to Faroese ships bunkering outside Faroese waters and the fuel sold to foreign ships in Faroese ports or outside Faroese waters is classified as international (Lastein and Winther, 2003).

Conclusively, the domestic/international fuel split (and associated emissions) for navigation is not determined with the same precision as for aviation. It is considered, however, that the potential of incorrectly allocated fuel

quantities is only a small part of the total fuel sold for navigational purposes in the Kingdom of Denmark.

3.3.5 Uncertainties and time series consistency

Uncertainty estimates for greenhouse gases on Tier 1 and Tier 2 levels, are made for road transport and other mobile sources using the guidelines formulated in the Good Practice Guidance and Uncertainty Management in National Greenhouse Gas Inventories (IPCC, 2000). For road transport, railways and fisheries, these guidelines provide uncertainty factors for activity data that are used in the Danish situation. For other sectors, the factors reflect specific national knowledge (Winther et al., 2006 and Winther, 2008). These sectors are (SNAP categories): Inland Waterways (a part of 1A3d: Navigation), Agriculture and Forestry (parts of 1A4c: Agriculture/forestry/fisheries), Industry (mobile part of (1A2f: Industry-other), Residential (1A4b) and National sea transport (a part of 1A3d: Navigation).

The activity data uncertainty factor for civil aviation is based on expert judgement.

The calculations for Tier 1 are shown in Annex 3.B.17 for all emission components. Please refer to Chapter 1.7 for further information regarding the calculation procedure for Tier 2 uncertainty calculations.

Table 3.3.18 Tier 1 Uncertainties for activity data, emission factors and total emissions in 2012 and as a trend.

Category	Activity data	CO ₂	CH ₄	N ₂ O
	%	%	%	%
Road transport	2	5	40	50
Military	2	5	100	1000
Railways	2	5	100	1000
Navigation (small boats)	41	5	100	1000
Navigation (large vessels)	11	5	100	1000
Fisheries	2	5	100	1000
Agriculture	24	5	100	1000
Forestry	30	5	100	1000
Industry (mobile)	41	5	100	1000
Residential	35	5	100	1000
Commercial/Institutional	35	5	100	1000
Civil aviation	10	5	100	1000
Overall uncertainty in 2012		5.3	26.6	152.0
Trend uncertainty		6.0	5.4	66.8

Table 3.3.19 Tier 2 Uncertainty factors for activity data and emission factors in 2012.

Category	Activity data	CO ₂	CH ₄	N ₂ O
	%	%	%	%
Road transport	2	5	40	500
Military	2	5	100	1000
Railways	2	5	100	1000
Pleasure craft	41	5	100	1000
Regional ferries	20	5	100	1000
Local ferries	20	5	100	1000
Fisheries	2	5	100	1000
Greenland & Faroe Islands	20	5	100	1000
Other national sea transport	20	5	100	1000
Civil aviation	10	5	100	1000
Agriculture	24	5	100	1000
Forestry	30	5	100	1000
Industry	41	5	100	1000
Household and gardening	35	5	100	1000
Commercial and institutional	35	5	100	1000

Table 3.3.20 Tier 2 Uncertainty estimates for CO₂, CH₄, N₂O and CO₂-eq. in 2012.

		1990			2012			1990-2012		
		Median		Uncertainty	Median		Uncertainty	Median		Uncertainty
				(%)			(%)			(%)
		Emission	Lower	Upper	Emission	Lower	Upper	Emission	Lower	Upper
			(-)	(+)		(-)	(+)		(-)	(+)
CO ₂	ktonnes	13624	5	5	15354	5	5	13	9	9
CH ₄	tonnes	2695	28	39	997	26	37	-63	33	47
N ₂ O	Tonnes	694	46	202	753	43	183	10	299	521
CO ₂ eq.	Ktonnes	13924	5	6	15635	5	6	12	10	11

As regards time series consistency, background flight data cannot be made available on a city-pair level prior to 2000. However, aided by LTO/aircraft statistics for these years and the use of proper assumptions, a good level of consistency is, in any case, obtained for this part of the transport inventory.

The time series of emissions for mobile machinery in the agriculture, forestry, industry, household and gardening (residential) and inland waterways (part of navigation) sectors are less certain than time series for other sectors, since DEA statistical figures do not explicitly provide fuel consumption information for working equipment and machinery.

3.3.6 Quality assurance/quality control (QA/QC)

The intention is to publish every second year a sector report for road transport and other mobile sources. The last sector report prepared concerned the 2010 inventory (Winther, 2012).

The QA/QC descriptions of the Danish emission inventories for transport follow the general QA/QC description for DCE in Section 1.6, based on the prescriptions given in the IPCC Good Practice Guidance and Uncertainty Management in National Greenhouse Gas Inventories (IPCC, 2000). A general QA/QC plan for the Danish greenhouse gas inventory has been elaborated by Nielsen et al. (2012).

An overview diagram of the Danish emission inventory system is presented in Figure 1.2 (Data storage and processing levels), and the exact definitions of Critical Control Points (CCP) and Points of Measurements (PM) are given in Section 1.6. The status for the PMs relevant for the mobile sector are given in the following text and the result of this investigation indicates a need for future QA/QC activities in order to fulfil the QA/QC requirements from the IPCC GPG.

Data storage level 1

Data Storage level 1	3.Completeness	DS.1.3.1	Documentation showing that all possible national data sources are included by setting down the reasoning behind the selection of datasets.
----------------------	----------------	----------	--

The following external data sources are used in the mobile part of the Danish emission inventories for activity data and supplementary information:

- Danish Energy Agency: Official Danish energy statistics.
- National sea transport (Royal Arctic Line, Eim Skip): Annual fuel consumption data.
- DTU Transport: Road traffic vehicle fleet and mileage data.
- Civil Aviation Agency of Denmark: Flight statistics.
- Non-road machinery: Information from statistical sources, research organisations, different professional organisations and machinery manufacturers.
- Ferries (Statistics Denmark): Data for annual return trips for Danish ferry routes.
- Ferries (Danish Ferry Historical Society): Detailed technical and operational data for specific ferries.
- Ferries (Mols Linjen, Bornholmstrafikken, Langelandstrafikken, Smyril Line): Detailed technical and operational data for specific ferries.
- Danish Meteorological Institute (DMI): Temperature data.
- The National Motorcycle Association: 2-wheeler data.

The emission factors come from various sources:

- Danish Energy Agency: CO₂ emission factors and lower heating values (all fuel types).
- COPERT IV: Road transport (all exhaust components, except CO₂, SO₂).
- Danish State Railways: Diesel locomotives (NO_x, VOC, CO and TSP).
- EMEP/EEA guidebook: Civil aviation and supplementary.
- Non road machinery: References given in NERI reports.
- National sea transport and fisheries: TEMA2000 (NO_x, VOC, CO and TSP) and MAN Diesel (sfc, NO_x).

Table 3.3.21 to follow contains Id, File/Directory/Report name, Description, Reference and Contacts. As regards File/Directory/Report name, this field refers to a file name for Id when all external data (time series for the existing inventory) are stored in one file. In other cases, a computer directory name is given when the external data used are stored in several files, e.g. each file contains one inventory year's external data or each file contains time series of external data for sub-categories of machinery. A third situation occurs when the external data are published in publicly available reports; here the aim is to obtain electronic copies for internal archiving.

Table 3.3.21 Overview table of external data and contact persons for transport.

Id no	File/- Directory/- Report name	Description	Activity data or emission factor	Reference	Contacts	Data agreement
T1	Transport energy¹	Dataset for all transport energy use	Activity data	The Danish Energy Agency (DEA)	Jane Rusbjerg	Yes
T2	Fleet and mileage data¹	Road transport fleet and mileage data	Activity data	DTU Transport	Thomas Jensen	Yes
T3	Flight statistics²	Data records for all flights	Activity data	Danish Transport Authority	Jess Nørgaard	Yes
T4	Non road machinery²	Stock and operational data for non-road machinery	Activity data	Non road Documentation report		No
T5	Emissions from ships³	Data for ferry traffic	Activity data	Statistics Denmark	Bo Henry Eriksen	No
T6	Emissions from ships³	Technical and operational data for Danish ferries	Activity data	Navigation emission documentation report	Hans Otto Kristensen	No
T7	Temperature data³	Monthly average of daily max/min temperatures	Other data	Danish Meteorological Institute	Danish Meteorological Institute	No
T8	Fleet and mileage data¹	Stock data for mopeds and motorcycles	Activity data	The National Motorcycle Association	Henrik Markamp	No
T9	CO₂ emission factors¹	DEA CO ₂ emission factors (all fuel types)	Emission factor	The Danish Energy Agency (DEA)	Jane Rusbjerg	No
T10	COPERT IV emission factors³	Road transport emission factors	Emission factor	Laboratory of applied thermodynamics Aristotle University Thessaloniki	Leonidas Ntziachristos	No
T11	Railways emission factors¹	Emission factors for diesel locomotives	Emission factor	Danish State Railways	Per Delvig	Yes
T12	EMEP/EEA guidebook³	Emission factors for navigation, civil aviation and supplementary	Emission factor	European Environment Agency	European Environment Agency	No
T13	Non road emission factors³	Emission factors for agriculture, forestry, industry and household/gardening	Emission factor	Non road Documentation report		No
T14	Emissions from ships³	Emission factors for national sea transport and fisheries	Emission factor	Navigation emission documentation report		No

¹⁾ File name; ²⁾ Directory in the DCE data library structure; ³⁾ Reports available on the internet.

Danish Energy Agency (energy statistics)

The official Danish energy statistics are provided by the Danish Energy Agency (DEA) and are regarded as complete on a national level. For most

transport sectors, the DEA subsector classifications fit the SNAP classifications used by DCE.

For non-road machinery, this is however not the case, since DEA do not distinguish between mobile and stationary fuel consumption in the subsectors relevant for non-road mobile fuel consumption.

Here, DCE calculates a bottom-up non-road fuel consumption estimate and for diesel (land based machinery only) and LPG, the residual fuel quantities are allocated to stationary consumption. For gasoline (land-based machinery) the relevant fuel consumption quantities for the DEA are smaller than the DCE estimates, and the amount of fuel consumption missing is subtracted from the DEA road transport total to account for all fuel sold. For recreational craft, no specific DEA category exists and, in this case, the gasoline and diesel fuel consumption is taken from road transport and fisheries, respectively.

In the case of Danish national sea transport, fuel consumption estimates are obtained by DCE (Winther, 2008), since they are regarded as much more accurate than the DEA fuel sales data. For the latter source, the large fluctuations in reported fuel sales cannot be explained by the actual development in the traffic between different national ports.

In order to maintain the national energy balance, the updated fuel consumption time series for national sea transport lead, in turn, to changes in the fuel activity data for fisheries (gas oil) and industry (heavy fuel oil).

The DCE fuel modifications, thus, give DEA-SNAP differences for road transport, national sea transport and fisheries.

A special note must be made for the DEA civil aviation statistical figures. The domestic/international fuel consumption division derives from bottom-up fuel consumption calculations made by DCE.

DTU Transport

Figures for fleet numbers and mileage data are provided by DTU Transport on behalf of the Danish Ministry of Transport. Following the data deliverance contract between DCE and the Danish Ministry of Transport, it is a basic task for DTU Transport to possess comprehensive information on Danish road traffic. The fleet figures are based on data from the Car Register, kept by Statistics Denmark and are, therefore, regarded as very precise. Annual mileage information is obtained by DTU Transport from the Danish Vehicle Inspection and Maintenance Programme.

Danish Transport Authority (Civil Aviation Agency of Denmark)

The Danish Transport Authority monitors all aircraft movements in Danish airspace and, in this connection, possesses data records for all take-offs and landings at Danish airports. The dataset from 2001 onwards, among others consisting of aircraft type and origin and destination airports for all flights leaving major Danish airports, are, therefore, regarded as very complete. For inventory years before 2001, the most accurate data contain Transport Authority total movements from major Danish airports and detailed aircraft type distributions for aircraft using Copenhagen Airport, provided by the airport itself.

Non-road machinery (stock and operational data)

A great deal of new stock and operational data for non road machinery was obtained in a research project carried out by Winther et al. (2006) for the 2004 inventory. The source for the agricultural machinery stock of tractors and harvesters is Statistics Denmark. Sales figures for tractors, harvesters and construction machinery, together with operational data and supplementary information, are obtained from The Association of Danish Agricultural Machinery Dealers. IFAG (The Association of Producers and Distributors of Fork Lifts in Denmark) provides fork-lift sale figures, whereas total stock numbers for gasoline equipment are obtained from machinery manufacturers with large Danish market shares, with figures validated through discussions with KVL. Stock information disaggregated into vessel types for recreational craft was obtained from the Danish Sailing Association. A certain part of the operational data comes from previous Danish non-road research projects (Dansk Teknologisk Institut, 1992 and 1993; Bak et al., 2003).

No statistical register exists for non-road machinery types and this affects the accuracy of stock and operational data. For tractors and harvesters, Statistics Denmark provide total stock data based on information from questionnaires and the registers of crop subsidy applications kept by the Ministry of Food, Agriculture and Fishery. In combination with new sales figures per engine size from The Association of Danish Agricultural Machinery Dealers, the best available stock data are obtained. In addition, using the sources for construction machinery and fork lift sale figures are regarded as the only realistic approach for consolidated stock information for these machinery types. Use of this source-type also applies in the case of machinery types (gasoline equipment, recreational craft) where data is even scarcer.

To support the 2012 inventory, new 2012 stock data for tractors, harvesters, fork lifts and construction machinery was obtained from the same sources as in Winther et al. (2006). For non-road machinery in general, it is, however, uncertain if data in such a level can be provided annually in the future.

Ferries (Statistics Denmark)

Statistics Denmark provides information of annual return trips for all Danish ferry routes from 1990 onwards. The data are based on monthly reports from passenger and ferry shipping companies in terms of transported vehicles passengers and goods. Thus, the data from Statistics Denmark are regarded as complete. Most likely the data can be provided annually in the future.

Ferries (Danish Ferry Historical Society, DFS)

No central registration of technical and operational data for Danish ferries and ferry routes is available from official statistics. However, one valuable reference to obtain data and facts about construction and operation of Danish ferries, especially in the recent 20 - 30 years is the archives of Danish Ferry Historical Society. Pure technical data has not only been obtained from this society's archives, but some of the knowledge has been obtained through the personal insight about ferries from some of the members of the society, which have been directly involved in the ferry business for example consultants, naval architects, marine engineers, captains and superintendents. However, until recently no documentation of the detailed DFS knowledge was established in terms of written reports or a central database system.

To make use of all the ferry specific data for the Danish inventories, DSF made a data documentation for the years 1990-2005 as a specific task of the research project carried out by Winther (2008).

Ferries (Mols Linjen, Bornholmstrafikken, Langelandstrafikken, Smyril Line)

For the years 2006+, the major Danish ferry companies are contacted each year in order to obtain ferry technical data, relating to specific ferries in service, annual share of total round trips and other technical information. The relevant annual information is given as personal communication, a method which can be repeated in the future.

National sea transport (Royal Arctic Line, Eim Skip)

For the years 2006+, the major shipping companies with frequent sailing activities between Denmark and Greenland/Faroe Islands are contacted each year in order to obtain data for fuel sold in Denmark used for these vessel activities. The relevant annual information is given as personal communication, a method which can be repeated in the future.

Danish Meteorological Institute

The monthly average max/min temperature for Denmark comes from DMI. This source is self explanatory in terms of meteorological data. Data are publicly available for each year on the internet.

The National Motorcycle Association

Road transport: 2-wheeler stock information (The National Motorcycle Association). Given that no consistent national data are available for mopeds in terms of fleet numbers and distributions according to new sales per year, The National Motorcycle Association is considered to be the professional organisation, where most expert knowledge is available. The relevant annual information is given as personal communication, a method which can be repeated in the future.

Danish Energy Agency (CO₂ emission factors and lower heating values)

The CO₂ emission factors and net calorific values (NCV) are fuel-specific constants. The country-specific values from the DEA are used for all inventory years.

COPERT IV

COPERT IV provides factors for fuel consumption and for all exhaust emission components which are included in the national inventory. For several reasons, COPERT IV is regarded as the most appropriate source of road traffic fuel consumption and emission factors. First of all, very few Danish emission measurements exist, so data are too scarce to support emission calculations on a national level. Secondly, most of the fuel consumption and emission information behind the COPERT model are derived from different large European research activities, and the formulation of fuel consumption and emission factors for all single vehicle categories has been made by a group of road traffic emission experts. A large degree of internal consistency is, therefore, achieved. Finally, the COPERT model is regularly updated with new experimental findings from European research programmes and, apart from updated fuel consumption and emission factors, the use of COPERT IV by many European countries ensures a large degree of cross-national consistency in reported emission results.

Danish State Railways

Aggregated emission factors of NO_x, VOC, CO and TSP for diesel locomotives are provided annually by the Danish State Railways. Taking into account available time resources for subsector emission calculations, the use of data from Danish State Railways is sensible. This operator accounts for around 90 % of all diesel fuel consumed by railway locomotives in Denmark and the remaining diesel fuel is used by various private railways companies. Setting up contacts with the private transport operators is considered to be a rather time consuming experience taking time away from inventory work in areas of greater emission importance.

EMEP/EEA guidebook

Fuel consumption and emission data from the EMEP/EEA guidebook is the prime and basic source for the aviation and navigation part of the Danish emission inventories. For aviation, the guidebook contains the most comprehensive list of representative aircraft types available for city-pair fuel consumption and emission calculations. The data have been evaluated specifically for detailed national inventory use by a group of experts representing civil aviation administration, air traffic management, emission modellers and inventory compilers.

In addition, the EMEP/EEA guidebook is the source of non-exhaust TSP, PM₁₀ and PM_{2.5} emission factors for road transport, and the primary source of emission factors for some emission components – typically N₂O, NH₃ and PAH – for other mobile sources.

Non-road machinery (fuel consumption and emission factors)

The references for non-road machinery fuel consumption and emission factors are listed in Winther et al. (2006). The fuel consumption and emission data is regarded as the most comprehensive data collection on a European level, having been thoroughly evaluated by German emission measurement and non-road experts within the framework of a German non-road inventory project.

National sea transport and fisheries

Emission factors for NO_x, VOC, CO and TSP are taken from the TEMA2000 model developed for the Ministry of Transport. To a large extent the emission factors originate from the exhaust emission measurement programme carried out by Lloyd's (1995). For NO_x, additional information of emission factors in a time series going back to 1949, and PM₁₀ and PM_{2.5} fractions of total TSP was provided by the engine manufacturer MAN Diesel.

Specifically for the ferries used by Mols Linjen new NO_x, VOC and CO emission factors are provided by Kristensen (2008), originating from measurement results by Hansen et al. (2004), Wismann (1999) and PHP (1996). Kristensen (2013) has provided complimentary emission factor data for new ferries.

The experimental work by Lloyd's is still regarded as the most comprehensive measurement campaign with results publicly available. The additional NO_x and PM₁₀/PM_{2.5} information comes from the world's largest ship engine manufacturer and data from this source is consistent with data from Lloyd's. Consequently the data used in the Danish inventories for national sea transport is regarded as the best available for emission calculations.

Data Storage level 1	1. Accuracy	DS.1.1.1	General level of uncertainty for every dataset, including the reasoning for the specific values
----------------------	-------------	----------	---

The uncertainty involved in the DEA fuel consumption information (except civil aviation) and the Danish Transport Authority flight statistics is negligible, as such, and this is also true for DMI temperature data. For civil aviation, some uncertainty prevails, since the domestic fuel consumption figures originate from a division of total jet-fuel sales figures into domestic and international fuel quantities, derived from bottom-up calculations. A part of the fuel consumption uncertainties for non-road machines is due to the varying levels of stock and operational data uncertainties, as explained in DS 1.3.1.

As regards emission factors, the CO₂ factors (and NCVs) from the DEA are considered to be very precise, since they relate only to fuel. For the remaining emission factor sources, the SO₂ (based on fuel sulphur content), NO_x, NMVOC, CH₄, CO, TSP, PM₁₀ and PM_{2.5} emission factors are less accurate. Though many measurements have been made, the experimental data rely on the individual measurement and combustion conditions. The uncertainties for N₂O and NH₃ emission factors increase even further due to the small number of measurements available. For heavy metals and PAH, experimental data are so scarce that uncertainty becomes very high.

A special note, however, must be made for energy. The uncertainties due to the subsequent treatment of DEA data for road transport, national sea transport, fisheries and the non-road relevant sectors, explained in DS 1.3.1, trigger some uncertainties in the fuel consumption figures for these sectors. This point is, though, more relevant for QA/QC description for data processing, Level 1.

Data Storage level 1	2.Comparability	DS.1.2.1	Comparability of the emission factors/calculation parameters with data from international guidelines, and evaluation of major discrepancies.
----------------------	-----------------	----------	--

Work has been carried out to compare Danish figures with corresponding data from other countries in order to evaluate discrepancies. The comparisons have been made on a CRF level, mostly for implied emission factors (Fauser et al., 2007, 2013).

Data Storage level 1	4.Consistency	DS.1.4.1	The origin of external data has to be archived with proper reference.
----------------------	---------------	----------	---

It is ensured that the original files from external data sources are archived internally at DCE. Subsequent raw data processing is carried out either in the DCE database models or in spreadsheets (data processing level 1).

Data Storage level 1	6.Robustness	DS.1.6.1	Explicit agreements between the external institution holding the data and DCE about the condition of delivery
----------------------	--------------	----------	---

For transport, DCE has made formal agreements with regard to external data deliverance with (Table 3.3.21 external data source Id's in brackets): DEA

(T1), the Danish Transport Authority (T3), Danish State Railways (T9) and DTU Transport (T2).

Data Storage level 1	7. Transparency	DS.1.7.1	Listing of all archived datasets and external contacts
----------------------	-----------------	----------	--

The listing of all archived datasets and external contact persons are given in Table 3.3.21.

Data Processing Level 1

Data Processing level 1	1. Accuracy	DP.1.1.1	Uncertainty assessment for every data source not part of DS.1.1.1 as input to Data Storage level 2 in relation to type and scale of variability.
-------------------------	-------------	----------	--

The general uncertainties of the DEA fuel consumption information, DMI temperature data, road transport stock totals and the Danish Transport Authority flight statistics are zero. For domestic aviation fuel consumption, the uncertainty is based on own judgement. For road transport, military and railways the fuel consumption uncertainties are taken from the IPCC Good Practice Guidance manual. It is noted that for road transport, it is not possible to quantify in-depth the uncertainties (1) of stock distribution into COPERT IV-relevant vehicle subsectors and (2) of the national mileage figures, as such.

In the mobile part of the Danish emission inventories, uncertainty assessments are made at Data Processing Level 1 for non-road machinery, recreational craft and national sea transport. For these types of mobile machinery, the stock and operational data variations are assumed to be normally distributed (Winther et al., 2006; Winther, 2008). Tier 1 uncertainty calculations produce final fuel consumption uncertainties ready for Data Storage Level 2 (SNAP level 2: Inland waterways, agriculture, forestry, industry and household-gardening). The sizes of the variation intervals are given for activity data and emission factors in the present report.

For non-road machinery stock and operational data, the uncertainty figures are given in Winther et al. (2006). For navigation, the uncertainty figures are given in Winther (2008).

For emission factors, the uncertainties for mobile sources are determined as suggested in the IPCC and UNECE guidelines. The uncertainty figures are listed in Paragraph 1.1.5 for greenhouse gases, and in Winther et al. (2006) and Winther (2008, 2012) for the remaining emission components.

Data Processing level 1	1. Accuracy	DP.1.2.1	The methodologies have to follow the international guidelines suggested by UNFCCC and IPCC.
-------------------------	-------------	----------	---

An evaluation of the methodological inventory approach has been made, which proves that the emission inventories for transport are made according to the international guidelines (Winther, 2005: Kyoto notat, in Danish). This paper will be translated into English and the conclusions will be implemented in the future national inventory reports.

Data Processing level 1	1. Accuracy	DP.1.1.4	Verification of calculation results using guideline values
-------------------------	-------------	----------	--

It has been checked that the greenhouse gas emission factors used in the Danish inventory are within margin of the IPCC guideline values.

Data Processing level 1	3.Completeness	DP.1.3.1	Identification of data gaps with regard to data sources that could improve quantitative knowledge.
-------------------------	----------------	----------	--

No important areas can be identified.

Data Processing level 1	4.Consistency	DP.1.4.1	Documentation and reasoning of methodological changes during the time series and the qualitative assessment of the impact on time series consistency.
-------------------------	---------------	----------	---

Se DP 1.7.5.

Data Processing level 1	5.Correctness	DP.1.5.2	Verification of calculation results using time series
-------------------------	---------------	----------	---

Data Processing level 1	5.Correctness	DP.1.5.3	Verification of calculation results using other measures
-------------------------	---------------	----------	--

For road transport, aviation, navigation and non-road machinery, whether all external data are correctly put into the DCE transport models is checked. This is facilitated by the use of sum queries which sum up stock data (and mileages for road transport) to input aggregation levels. However, spreadsheet or database manipulations of external data are, in some cases, included in a step prior to this check.

This is carried out in order to produce homogenous input tables for the DCE transport models (road, civil aviation, non-road machinery/recreational craft, navigation/fisheries). The sub-routines perform operations, such as the aggregation/disaggregation of data into first sales year (Examples: Fleet numbers and mileage for road transport, stock numbers for tractors, harvesters and fork lifts) or simple lists of total stock pr year (per machinery type for e.g. household equipment and for recreational craft). For civil aviation, additional databases control the allocation of representative aircraft to real aircraft types and the cruise distance between airports. A more formal description of the sub-routines will be made.

Regarding fuel data, it is checked for road transport and civil aviation that DEA totals (modified for road) match the input values in the DCE models. For the transport modes military and railways, the DEA fuel consumption figures go directly into Data Storage Level 2. This is also the case for the railway emission factors obtained from Danish State Railways and, generally, for the emission factors, which are kept constant over the years.

The DCE model simulations of fuel consumption and emission factors for road transport, civil aviation and non-road machinery refer to Data Processing Level 1.

When DCE transport model changes are made relating to fuel consumption, it is checked that the calculated fuel consumption sums correspond to the expected fuel consumption levels in the time series. The fuel consumption check also includes a time series comparison with fuel consumption totals calculated in the previous model version. The checks are performed on a SNAP level and, if appropriate, detailed checks are made for vehicle/machinery technology splits.

As regards model changes in relation to derived emission factors (and calculated emissions), the time series of emission factors (and emissions) are compared to previous model figures. A part of this evaluation includes an assessment, if the development corresponds to the underlying assumptions given by detailed input parameters. Among other things, the latter parameters depend on emission legislation, new technology phase-in, deterioration factors, engine operational conditions/driving modes, gasoline evaporation (hydrocarbons) and cold starts. For methodological issues, please refer to Section 3.3.2.

Data Processing level 1	7.Transparency	DP.1.7.1	The calculation principle, the equations used and the assumptions made must be described
-------------------------	----------------	----------	--

The DCE model calculation principles and basic equations are thoroughly described in the present report, together with the theoretical model reasoning and assumptions. Documentation is also given e.g. in Winther (2001a, 2008, 2012) and Winther et al. (2006). Further formal descriptions of DCE model sub routines are given in internal notes, and flow maps show the interrelations between tables and calculation queries in the models.

During model development it has been checked that all mathematical model relations give exactly the same results as independent calculations.

Data Processing level 1	7.Transparency	DP.1.7.2	Clear reference to dataset at Data Storage level 1.
-------------------------	----------------	----------	---

In the different documentation reports for transport in the Danish emission inventories, there are explicit references for the different external data used.

Data Processing level 1	7.Transparency	DP.1.7.3	A manual log to collect information about recalculations
-------------------------	----------------	----------	--

Recalculation changes in the emission inventories are described in the NIR and IIR reports as a standard. These descriptions take into account changes in emission factors, activity data and calculation methods.

Data Storage Level 2

Data Storage level 2	5.Correctness	DS.2.5.1	Check if a correct data import to level 2 has been made.
----------------------	---------------	----------	--

At present, a DCE software programme imports data from prepared input data tables (SNAP fuel consumption figures and emission factors) into the CollectER database.

Tables for CollectER fuel consumption and emission results are prepared by a special DCE database (NERIrep.mdb). The results relevant for mobile

sources are copied into a database containing all the official inventory results for mobile sources (Data2012 NIR-UNECE.mdb). By the use of database queries, the results from this latter database are aggregated into the same formats as being used by the relevant DCE transport models in their results calculation part. The final comparison between CollectER and DCE transport model results are set up in a spreadsheet.

Data Storage Level 4

Data Storage level 4	4.Consistency	DS.4.4.3	The IEFs from the CRF are checked both regarding level and trend. The level is compared to relevant emission factors to ensure correctness. Large dips/jumps in the time series are explained.
----------------------	---------------	----------	--

A spreadsheet “Check CRF 2011.xls” has been set up to check that the fuel consumption and emission totals from CollectER imported in Data2012 NIR-UNECE.mdb are identical to the fuel consumption and emission totals from the CRF.

3.3.7 Recalculations and improvements

The following recalculations and improvements of the emission inventories have been made since the emission reporting in 2012.

Road transport

Based on the updated version of COPERT IV launched in 2013, new vehicle sub categories have been introduced in the emission inventories for mopeds and passenger cars. For mopeds a division is now made between 2-stroke and 4-stroke engine technologies and for passenger cars small engine sizes below 0.8 l. for gasoline and below 1.4 l. for diesel have been included. Also NO_x emission factors for euro 5 diesel passenger cars have been updated in the model based on the new COPERT IV version.

Small errors in input gasoline fuel consumption for the years 2009-2011 and for input diesel fuel consumption in the years 2010-2011 have been corrected.

The percentage emission change interval and year of largest percentage differences (low %; high %, year) for the different emission components are: CO₂ (-0.5 %; -0.05 %, 2008), CH₄ (-0.2 %; 2.4 %, 2011) and N₂O (-0.4 %; 0.3 %, 2008).

Navigation

Minor changes in ferry input data has been made for the years 2008-2011 causing minor emission changes for domestic navigation. The following largest percentage differences (in brackets) for domestic navigation are noted for: CO₂ (-1.0 %), CH₄ (-0.8 %) and N₂O (-1.1 %).

Agriculture/forestry/fisheries

The number and engine size of machine pool tractors has been updated for the years 2007-2011. The number of ATV's has been changed for the years 2009-2011.

Errors in the fuel consumption for fisheries in 2000, 2010 and 2011 have been corrected.

In 2000 the following percentage differences (in brackets) for agriculture/forestry/fisheries are noted for: CO₂ (9.1 %), CH₄ (3.7 %) and N₂O (11.9 %) due to fuel consumption changes in fisheries. For other years than 2000, the following largest percentage differences (in brackets) are noted for: CO₂ (-3.7 %), CH₄ (5 %) and N₂O (-4.9 %).

Industry

The number of mini loaders has been updated for the years 2004-2011.

The following largest percentage differences (in brackets) for industrial non road machinery are noted for: CO₂ (1.0 %), CH₄ (1.6 %) and N₂O (1.6 %).

Civil aviation

An error in the CH₄ emission factor has been corrected for the years 1985-2000. The emission factors are now in line with the factors proposed by the EMEP/EEA emission inventory guidebook. The CH₄ emission percentage differences are between -31 % and -42 %.

Military

Emission factors derived from the new road transport simulations have caused some emission changes from 1985-2010. The following largest percentage differences (in brackets) for military are noted for: CO₂ (0 %), CH₄ (0.6 %) and N₂O (0.2 %).

3.3.8 Planned improvements

No planned improvements are envisaged to be made.

QA/QC

Future improvements regarding this issue are dealt with in Section 3.1.4.

3.3.9 References for Chapter 3.3

Cappelen, J. 2013: The Climate of Denmark 2012 - with English summary, Technical report No 13-01, pp. 68, Danish Meteorological Institute.

Cowi, 2008: Model til beregning af vej og banetransportens CO₂-æquivalent emissioner, Technical note, pp. 42, Ministry of Transport (in Danish).

Dávastovu, J. 2011: Unpublished data material from Smyril Line.

Danish Energy Agency, 2013: The Danish energy statistics, Available at: <http://www.ens.dk/sites/ens.dk/files/info/tal-kort/statistik-noegletal/aarlig-energistatistik/energistatistik2012.pdf> (21-01-2014).

Delvig, P. 2013: Unpublished data material from the Danish State Railways.

Ekman, B. 2005: Unpublished data material from the Danish Road Directorate.

EMEP/EEA, 2013: Air Pollutant Emission Inventory Guidebook, prepared by the UNECE/EMEP Task Force on Emissions Inventories and Projections (TFEIP). Available at: <http://www.eea.europa.eu/publications/emep-eea-guidebook-2013> (02-09-2013).

Fausser, P., Thomsen, M., Nielsen, O.K., Winther, M., Gyldenkerne, S., Hoffmann, L., Lyck, E. & Illerup, J.B. 2007: Verification of the Danish emis-

sion inventory data by national and international data comparisons. National Environmental Research Institute, University of Aarhus. - NERI Technical Report 627: 53 pp. (electronic). Available at:
http://www2.dmu.dk/Pub/FR627_Final.pdf.

Fausser, P., Nielsen, M., Winther, M., Plejdrup, M., Gyldenkerne, S., Mikkelsen, M.H., Albrektsen, R., Hoffmann, L., Thomsen, M., Hjelgaard, K. & Nielsen, O.-K. 2013. Verification of the Danish 1990, 2000 and 2010 emission inventory data. Aarhus University, DCE – Danish Centre for Environment and Energy, 85 pp. Scientific Report from DCE – Danish Centre for Environment and Energy No. 79. <http://dce2.au.dk/pub/SR79.pdf>

Hansen, C.O. 2010: Estimation af udenlandske bilers trafikarbejde i Danmark i 2009, Notatnr. 12032-005, 12 pp. Danish Road Directorate, 2010.

Hansen, K.F. & Jensen, M.G. 2004: MÅLING AF EMISSIONER FRA FREMDRIVNINGSSANLÆG PÅ MADSMOLS. Ruston 20RK270, Sagsnr.: 1076868, Documentation note, 5 pp. (in Danish).

Hjelgaard, K. & Winther, M. 2011: The estimated new sales of Euro V diesel trucks equipped with EGR and SCR during the 2006-2010 time periods. Internal DCE note (unpublished). 2 pp. (in Danish).

ICAO Annex 16: "International standards and recommended practices", Volume II "Aircraft Engine Emissions", 2th ed. (1993), plus amendments: Amendment 3, 20th March 1997 and Amendment 4, 4th November 1999.

IFEU, 2004: Entwicklung eines Modells zur Berechnung der Luftschadstoffemissionen und des Kraftstoffverbrauchs von Verbrennungsmotoren in mobilen Geräten und Maschinen - Endbericht, UFOPLAN Nr. 299 45 113, pp. 122, Heidelberg.

IPCC, 2000: Good Practice Guidance and Uncertainty Management in National Greenhouse Gas Inventories. Available at:
<http://www.ipcc-nggip.iges.or.jp/public/gp/english/> (15-04-2007).

Jensen, T.C., O. 2013: Dokumentation af konvertering af trafiktal til emissionsopgørelser, arbejdsnotat, 22 pp. DTU Transport, 2013.

Jørgensen, H. 2013: Unpublished data material from Færgen A/S.

Kristensen, F. 2008: Unpublished data material from Mols-Linjen.

Kristensen, F. 2013: Unpublished data material from Mols-Linjen.

Lastein, L. & Winther, M. 2003: Emission of greenhouse gases and long-range transboundary air pollutants in the Faroe Islands 1990-2001. National Environmental Research Institute. - NERI Technical Report 477: 62 pp. (electronic). Available at:
http://www.dmu.dk/1_viden/2_Publikationer/3_fagrapporter/FR477.PDF

Markamp, H. 2013: Personal communication, Henrik Markamp, The National Motorcycle Association.

Marpol 73/78 Annex VI: Regulations for the prevention of air pollution from ships, technical and operational implications, DNV, 21 February 2005.

Ministry of Transport, 2000: TEMA2000 - et værktøj til at beregne transporters energiforbrug og emissioner i Danmark (TEMA2000 - a calculation tool for transport related fuel use and emissions in Denmark). Technical report. Available at: <http://www.trm.dk/sw664.asp>

Ntziachristos, L. & Samaras, Z. 2000: COPERT III Computer Programme to Calculate Emissions from Road Transport - Methodology and Emission Factors (Version 2.1). Technical report No 49. European Environment Agency, November 2000, Copenhagen. Available at: http://re-ports.eea.eu.int/Technical_report_No_49/en (June 13, 2003).

Nørgaard, T. & Hansen, K.F. 2004: Chiptuning af køretøjer - miljømæssig effekt, Miljøprojekt nr. 888, Miljøstyrelsen.

Nielsen, O.-K., Plejdrup, M.S., Winther, M., Gyldenkerne, S., Thomsen, M., Fauser, P., Nielsen, M., Mikkelsen, M.H., Albrechtsen, R., Hjelgaard, K., Hoffmann, L. & Bruun, H.G. 2012. Quality manual for the Danish greenhouse gas inventory. Version 2. Aarhus University, DCE – Danish Centre for Environment and Energy, 44 pp. Scientific Report from DCE – Danish Centre for Environment and Energy No. 47. Available at: <http://www.dmu.dk/Pub/SR47.pdf>

PHP, 1996: Research Report – Emission tests at Alpha, Mols 2 and Mols 4, 9L25MC mk6 engines #35031 and #35033, 22-23/10 1995 and 16/1 1996, DOK, PHP Basic Research, October 1996, 20 pp.

Pulles, T., Aardenne J.v., Tooly, L. & Rypdal, K. 2001: Good Practice Guidance for CLRTAP Emission Inventories, Draft chapter for the UNECE CORINAIR Guidebook, 7 November 2001, 42 pp.

Rasmussen, H. 2013: Unpublished data material from Royal Arctic Line.

Statistics Denmark, 2013: Data from Statbank Denmark. Available at: <http://www.statistikbanken.dk/statbank5a/default.asp?w=1364>

Thorarensen, B. 2013: Unpublished data material from Eim Skip.

USEPA, 2004: Conversion Factors for Hydrocarbon Emission Components. EPA420-P-04-001, US Environmental Protection Agency, 5 pp.

Winther, M. 2001a: 1998 Fuel Use and Emissions for Danish IFR Flights. Environmental Project no. 628, 2001. 112 pp. Danish EPA. Prepared by the National Environmental Research Institute, Denmark. Available at: <http://www.mst.dk/udgiv/Publications/2001/87-7944-661-2/html/>.

Winther, M. 2001b: Improving fuel statistics for Danish aviation. National Environmental Research Institute, Denmark. 56 pp. – NERI Technical Report No. 387.

Winther, M. 2005: Kyoto notat - Transport. Internal NERI note (unpublished). 4 pp. (in Danish).

Winther, M. & Nielsen, O.K. 2006: Fuel use and emissions from non-road machinery in Denmark from 1985-2004 - and projections from 2005-2030. The Danish Environmental Protection Agency. - Environmental Project 1092: 238 pp. Available at:

<http://www.dmu.dk/Udgivelser/-Arbejdsrapporter/Nr.+200-249/>

Winther, M. 2008: Fuel consumption and emissions from navigation in Denmark from 1990-2005 - and projections from 2006-2030. Technical Report from NERI no. 650. 109 pp. Available at:

<http://www2.dmu.dk/Pub/FR650.pdf>

Winther, M. 2009: Emission Differences between Petroleum based Diesel and different Biodiesel Blend Ratios for Road Transport Vehicles. / Winther, Morten. 2009. Transport and Air Pollution Symposium - 3rd Environment and Transport Symposium, nr. 17, Toulouse, France, 2.- 4. june 2009.

Winther, M. 2011: Ændring i partikelemission som følge af fremskyndet salg af Euro 5 biler og retrofit af filtre på tunge køretøjer. Internal DCE note (unpublished). 2 p. (in Danish).

Winther, M. 2012: Danish emission inventories for road transport and other mobile sources. Inventories until the year 2010. National Environmental Research Institute, University of Aarhus. 283 pp. - DCE Scientific Report No. 24. <http://www.dmu.dk/Pub/SR24.pdf>.

Wismann, T. 1999: MOLS-LINIEN, Mai Mols - Måling af emissioner fra fra hovedturbiner, dk-RAPPORT 14.901, 9 pages (in Danish).

Wismann, T. 2001: Energiforbrug og emissioner fra skibe i farvandene omkring Danmark 1995/1996 og 1999/2000 (Fuel consumption and emissions from ships in Danish coastal waters 1995/1996 and 1999/2000). The Danish Environmental Protection Agency. - Environmental Project 597: 88 pp. Available at:

<http://www2.mst.dk/common/Udgivramme/Frame.asp?pg=http://www2.mst.dk/Udgiv/publikationer/2001/87-7944-505-5/html/default.htm>.

Wismann, T. 2007: Energiforbrug for skibe i fart mellem danske havne (Fuel consumption by ships sailing between Danish ports), Internal note, September 2007, 3 pp.

3.4 Additional information, CRF sector 1A Fuel combustion

Reference approach, feedstocks and non-energy use of fuels

In addition to the sector specific CO₂ emission inventories (the national approach), the CO₂ emission is also estimated using the reference approach described in the IPCC Reference Manual (IPCC, 1997). The reference approach is based on data for fuel production, import, export and stock change. The CO₂ emission inventory based on the reference approach is reported to the Climate Convention and used for verification of the official data in the national approach.

Data for import, export and stock change used in the reference approach originate from the annual “basic data” table prepared by the Danish Energy Agency (DEA) and published on their home page (DEA, 2013). The fraction of carbon oxidised has been assumed to be 1.00. The carbon emission factors are default factors originating from the IPCC Reference Manual (IPCC, 1997). The country-specific emission factors are not used in the reference approach, the approach being for the purposes of verification. The emission factor for fossil waste is, however, based on the emission factor applied in the national approach.

The Climate Convention reporting tables include a comparison of the national approach and the reference approach estimates. To make results comparable, the incineration of fossil waste and the corresponding CO₂ emission have been added in the reference approach. Furthermore, consumption for non-energy purposes is subtracted in the reference approach, because non-energy use of fuels is included in other sectors (Industrial processes and Solvent use) in the Danish national approach. The consumption for non-energy purposes have been subtracted in the reference approach by setting the fraction of carbon stored equal to 1.00 for fuels used for non-energy purposes.

Three fuels are used for non-energy purposes: lubricants, bitumen and white spirit. The total consumption for non-energy purposes is relatively low – 11.5 PJ in 2012.

The CO₂ emission from oxidation of lube oil during use was 32 Gg in 2012 and this emission is reported in the industrial processes sector. The reported emission corresponds to 20 % of the CO₂ emission from lube oil consumption assuming full oxidation. This is in agreement with the 2006 IPCC Guidelines (IPCC, 2006) methodology for lube oil emissions. Methodology and emission data for lube oil are shown in NIR Chapter 4.8.

For white spirit the CO₂ emission is indirect as the emissions occur as NMVOC emissions from the use of white spirit as a solvent. The indirect CO₂ emission from white spirit was 17 Gg in 2012 corresponding to 62 % of the CO₂ emission from white spirit assuming full oxidation. The NMVOC emission data for white spirit are shown in NIR Chapter 5, Table 5.4.

The CO₂ emission from bitumen is included as part of the emission from the source sectors 2A5 *Asphalt roofing* and 2A6 *Road paving with asphalt*.

According to IPCC Good Practice Guidance (IPCC, 2000) the difference should be within 2 %. A comparison of the national approach and the reference approach is illustrated in Figure 3.4.1.

In 2012, the fuel consumption rates in the two approaches differ by 0.97 % and the CO₂ emission differs by 0.97 %. In the period 1990-2012 both the fuel consumption and the CO₂ emission differ by less than 2.0 %. The differences are below 1% for all years except 1998, 2001, 2009 and 2011. The results from the national approach and the reference approach are compared in Figure 3.4.1.




Figure 3.4.1 Comparison of the reference approach and the national approach.

The fluctuations in figure 3.4.1 follow the fluctuations of the statistical difference in the Danish energy statistics shown in figure 3.4.2. The large differences in certain years, e.g. in 1998 and 2009 are due to high statistical differences in the Danish energy statistics in these years.




Figure 3.4.2 Statistical difference in the Danish energy statistics (DEA, 2013).

References for Chapter 3.4

Danish Energy Agency (DEA), 2013: The Danish energy statistics, Available at: <http://www.ens.dk/en/info/facts-figures/energy-statistics-indicators-energy-efficiency/annual-energy-statistics> (2014-02-03)

IPCC, 1997: Revised 1996 IPCC Guidelines for National Greenhouse Gas Inventories: Reference Manual (Volume 3). Available at: <http://www.ipcc-nggip.iges.or.jp/public/gl/invs6.html> (2013-02-13).

IPCC, 2000: Penman, J., Kruger, D., Galbally, I., Hiraishi, T., Nyenzi, B., Emmanuel, S., Buendia, L., Hoppaus, R., Martinsen, T., Meijer, J., Miwa, K. & Tanabe, K. (Eds). Good Practice Guidance and Uncertainty Management in National Greenhouse Gas Inventories. Published: IPCC/OECD/IEA/IGES, Hayama, Japan. Available at: <http://www.ipcc-nggip.iges.or.jp/public/gp/english/index.html> (2014-02-03).

IPCC, 2006: Revised 2006 IPCC Guidelines for National Greenhouse Gas Inventories: Available at: <http://www.ipcc-nggip.iges.or.jp/public/2006gl/index.html> (2014-02-03).

3.5 Fugitive emissions (CRF sector 1B)

Fugitive emissions from fuels include emissions from production, storage, refining and transport of oil and natural gas. Emissions from solid fuels are not occurring in Denmark. Most fugitive emission sources are of minor importance compared to the total Danish emissions. Fugitive and national total emissions are given in Table 3.5.1 with the fugitive emissions share of national total emission.

Table 3.5.1 National and fugitive emissions of CO₂, CH₄ N₂O and GHG in 2012, and the fugitive emissions share of national total emissions.

Compound	National emission		Fugitive emission		Fugitive/national emission	
CO ₂	39 412	Gg	221	Gg	0.6	%
CH ₄	261	Gg	5.0	Gg	1.9	%
N ₂ O	19	Gg	0.002	Gg	0.01	%
GHG	51 637	Gg CO ₂ eqv.	327	Gg CO ₂ eqv.	0.6	%

The key category analysis shows that CO₂ from offshore flaring is a Tier 1 Level key category in 1990 and 2012. Further CH₄ from Refinery processes is a Tier 2 Trend key source in 1990-2012 (Table 3.5.2).

Table 3.5.2 Key categories in the fugitive emission sector.

CRF table	Pollutant	Emission source	Key category identification	
			Tier 1	Tier 2
1.B.2	CO ₂	Flaring in refinery	-	-
1.B.2	CO ₂	Flaring off-shore	Level 1990 and 2012	-
1.B.2	CO ₂	Land based activities	-	-
1.B.2	CO ₂	Off-shore activities	-	-
1.B.2	CO ₂	Transmission of natural gas	-	-
1.B.2	CO ₂	Distribution of natural gas	-	-
1.B.2	CO ₂	Venting in gas storage	-	-
1.B.2	CH ₄	Flaring in refinery	-	-
1.B.2	CH ₄	Flaring off-shore	-	-
1.B.2	CH ₄	Refinery processes	-	Trend 1990-2012
1.B.2	CH ₄	Land based activities	-	-
1.B.2	CH ₄	Off-shore activities	-	-
1.B.2	CH ₄	Transmission of natural gas	-	-
1.B.2	CH ₄	Distribution of natural gas	-	-
1.B.2	CH ₄	Venting in gas storage	-	-
1.B.2	N ₂ O	Flaring in refinery	-	-
1.B.2	N ₂ O	Flaring off-shore	-	-

Calculations of fugitive emissions are mainly based on Tier 3 methodologies. Only calculation of emissions from Onshore loading of ships and distribution of oil products are using Tier 2 and Tier 1 methodologies. In accordance with the IPCC Good Practice Guidance (2000) emission calculations for fugitive key sources are done using higher methodological tiers than Tier 1. The applied methodologies and the level of detail for the applied emission factors in are listed in (Table 3.5.3).

Table 3.5.3 Applied methodology for fugitive emission sources.

CRF	Source	Pollutant	Method	Emission factor
1B2a ii	Oil – Production , Offshore activities	CO ₂	Tier 3	D
		CH ₄	Tier 3	OTH (EMEP/EEA 2013)
		NM VOC	Tier 3	OTH (EMEP/EEA 2013)
1B2a ii	Oil – Production , Onshore activities	CO ₂	Tier 2, Tier 3	D
		CH ₄	Tier 2, Tier 3	PS, OTH (EMEP/EEA 2013)
		NM VOC	Tier 2, Tier 3	PS, OTH (EMEP/EEA 2013)
1B2a iv	Oil - Refining /Storage	CH ₄	Tier 3	PS
		NM VOC	Tier 3	PS
		SO ₂	Tier 3	PS
1B2a v	Oil - Distribution of oil products	NM VOC	Tier 1	CS
1B2b	Natural gas	CO ₂	Tier 3	CS
		CH ₄	Tier 3	CS
		NM VOC	Tier 3	CS
1B2c	Venting /flaring	CO ₂	Tier 3	PS *
		CH ₄	Tier 3	PS, OTH (EMEP/EEA 2009)
		N ₂ O	Tier 3	D, OTH (OLF, 1993)
		NO _x	Tier 3	PS **, CS, OTH (EMEP/EEA 2009)
		CO	Tier 3	OTH (EMEP/EEA 2009)
		NM VOC	Tier 3	PS, CS, OTH (EMEP/EEA 2009)
		SO ₂	Tier 3	CS

PS: plant specific. CS: country specific, D: default (IPCC, 2000), OTH: other.

* Plant specific emission factors are available from the EU ETS from 2006 and forward. For earlier years country specific emission factors are applied.

** Plant specific emission factors are available for one refinery.

3.5.1 Source category description

According to the IPCC sector definitions the category *fugitive emissions* is a sub-category under the main-category Energy (Sector 1). The category *fugitive emissions* (Sector 1B) is segmented into sub-categories covering emissions from solid fuels (coal mining and handling (1B1a), solid fuel transformation (1B1b) and other (1B1c)) and from oil and natural gas (oil (1B2a), natural gas (1B2b), venting and flaring (1B2c) and other (1B2d)). The sub-categories relevant for the Danish emission inventory are shortly described below according to Danish conditions:

- 1B1a: Fugitive emission from solid fuels: Coal mining is not occurring in Denmark. Therefore only emissions of particulate matter from storage and handling of coal are considered.
- 1B2a: Fugitive emissions from oil include emissions from extraction, storage, and transmission of crude oil, distribution of oil products and fugitive emissions from refining. Emission data for offshore extraction of oil and gas are not available separately, and consequently emissions from gas extraction are included in 1B2a
- 1B2b: Fugitive emissions from natural gas include emissions from transmission and distribution of natural gas. Emissions from gas extraction are included in 1B2a.
- 1B2c: Venting and flaring include activities onshore and offshore. Flaring occur both offshore and onshore in gas treatment and storage plants and in refineries. Venting occurs in gas storage plants. Venting of gas is assumed to be negligible in extraction and in refineries as controlled venting enters the gas flare system.

Activity data, emission factors and emissions are stored in the Danish emission database on SNAP sector categories (Selected Nomenclature for Air Pollution). In Table 3.5.4 the corresponding SNAP codes and IPCC sectors relevant to fugitive emissions are shown. Further, the table holds the SNAP names for the SNAP codes and the overall activity (e.g. oil and natural gas).

Table 3.5.4 List of the IPCC sectors and corresponding SNAP codes for the categories included in the Danish emission inventory model.

IPCC sectors	SNAP code	SNAP name	Activity
1 B 1 a	050103 *	Storage of solid fuel	Coal mining and handling
1 B 2 a ii	050201	Land-based activities	Oil
1 B 2 a ii	050202 **	Offshore activities	Oil
1 B 2 a iv	040101	Petroleum products processing	Oil
1 B 2 a iv	040103	Other	Oil
		Service stations (including refuelling of cars)	
1 B 2 a v	050503		Oil
1 B 2 b iii	050601	Pipelines	Natural gas
1 B 2 b iv	050603	Distribution networks	Natural gas
1 B 2 c 2 i	090203	Flaring in oil refinery	Flaring
1 B 2 c 2 ii	050699	Venting in gas storage	Venting
1 B 2 c 2 ii	090206	Flaring in oil and gas extraction	Flaring

*Only relevant for emissions of particulate matter from storage and handling of coal.

**In the Danish inventory emissions from extraction of gas are united under "Extraction, 1st treatment and loading of liquid fossil fuels/offshore activities" (IPCC 1B2a / SNAP 050202).

Table 3.5.5 summarizes the Danish fugitive emissions in 2012. The methodologies, activity data and emission factors used for calculation are described in the following chapters.

Table 3.5.5 Summary of the Danish fugitive emissions 2012. P refers to point source and A to area source.

IPCC code	SNAP code	Source	Pollutant	Emission	Unit
1B2a i	050201	A	NMVOC	1 948	Mg
1B2a i	050201	A	CH ₄	837	Mg
1B2a i	050201	A	CO ₂	<0.1	Gg
1B2a i	050202	A	NMVOC	1 891	Mg
1B2a i	050202	A	CH ₄	1 744	Mg
1B2a i	050202	A	CO ₂	4	Gg
1B2a iv	040101	P	SO ₂	0 *	Mg
1B2a iv	040101	P	NMVOC	3 932	Mg
1B2a iv	040101	P	CH ₄	2 227	Mg
1B2a iv	040103	P	SO ₂	889	Mg
1B2a v	050503	A	NMVOC	977	Mg
1B2b iii	050601	A	NMVOC	4	Mg
1B2b iii	050601	A	CH ₄	14	Mg
1B2b iii	050601	A	CO ₂	< 0.1	Gg
1B2b iv	050603	A	NMVOC	57	Mg
1B2b iv	050603	A	CH ₄	147	Mg
1B2b iv	050603	A	CO ₂	< 0.1	Gg
1B2c	050699	P	NMVOC	20	Mg
1B2c	050699	P	CH ₄	57	Mg
1B2c	050699	P	CO ₂	< 0.1	Gg
1B2c	090203	P	SO ₂	167	Mg
1B2c	090203	P	NO _x	23	Mg
1B2c	090203	P	NMVOC	32	Mg
1B2c	090203	P	CH ₄	7	Mg
1B2c	090203	P	CO	74	Mg
1B2c	090203	P	CO ₂	22	Gg
1B2c	090203	P	N ₂ O	0.2	Mg
1B2c	090206	A	SO ₂	1	Mg
1B2c	090206	A	NO _x	87	Mg
1B2c	090206	A	NMVOC	8	Mg
1B2c	090206	A	CH ₄	15	Mg
1B2c	090206	A	CO	75	Mg
1B2c	090206	A	CO ₂	192	Gg
1B2c	090206	A	N ₂ O	1	Mg
1B2c	090206	P	SO ₂	<0.1	Mg
1B2c	090206	P	NO _x	9	Mg
1B2c	090206	P	NMVOC	0.3	Mg
1B2c	090206	P	CH ₄	0.5	Mg
1B2c	090206	P	CO	1	Mg
1B2c	090206	P	CO ₂	2	Gg
1B2c	090206	P	N ₂ O	<0.1	Mg

* From 2001 SO₂ emissions from oil refining are included in stationary combustion.

3.5.2 Methodological issues

The following chapters give descriptions on the methods of calculation used in the Danish emission inventory. Further, the activity data and emission

factors that form the basis for the calculations are described according to data source and values.

Test Use of EU ETS data

Reporting to the European Union Emission Trading Scheme (EU ETS) are available in the annual EU ETS reports for refineries, offshore oil and gas extraction facilities and the natural gas treatment plant, concerning fugitive emissions. EU ETS data are only included in the national emission inventory if higher tier methodologies are applied. The EU ETS data used are fully in line with the requirements in the IPCC good practice guidance and are considered the best data source on CO₂ emission factors due to the legal obligation for the relevant companies to make the accounting following the specified EU decisions. The EU ETS data are thereby a source of consistent data with low uncertainties. For further information on EU ETS please refer to Chapter 1.4.10. Unfortunately, corresponding data do not exist before the commencement of EU ETS in 2006 and therefore it is not possible to set up time series based on EU ETS. In these cases appropriate methods from the IPCC good practice guidance have been selected to ensure time series consistency. This is described in the specific sections.

Refineries:

Activity data are measured with flow meters and amounts are reported with high accuracy and the oxidation factor is set to 1. CO₂ emission factors are calculated according to the relevant Tier given in the EU Commission Decision of 18 July 2007 (EU Commission, 2007). For combustion of fuel gas, the Tier 2b methodology based on yearly density and calorific values is applied, while the activity specific Tier 3 methodology is applied for diesel. CO₂ emissions factors for flaring are calculated using the Tier 3 methodology based on the measured carbon contents of flare gas.

Offshore installations:

Activity data are measured with flow meters and amounts are reported with high accuracy (± 1.5 % for combustion and $\pm 7.5 - \pm 17.5$ % for flare). The oxidation factor is set to 1. CO₂ emission factors are calculated according to the relevant Tier given in the EU Commission Decision of 18 July 2007 (EU Commission, 2007). For combustion of fuel gas the Tier 3 methodology, which is activity specific, is applied, while the country specific Tier 2a methodology is applied for diesel. CO₂ emissions factors for flaring are calculated using the Tier 3 methodology based on the measured carbon contents of flare gas.

Fugitive emissions from oil (1B2a)

The emissions from oil derive from offshore activities (extraction of oil and gas, and offshore loading of ships), onshore activities (storage and handling at the raw oil terminal, and onshore loading of ships), service stations and refineries. In the case of service stations emissions from reloading of tankers and refuelling of vehicles are included. The emissions from refineries derive from petroleum products processing (oil refining). Emissions from flaring in refineries are included in the chapters concerning flaring.

Activities

Fugitive emissions from oil include emissions from extraction, the raw oil terminal, and onshore and offshore loading of ships.

The total emission can be expressed as:

$$E_{total} = E_{extraction} + E_{shiploading} + E_{oil terminal} \quad (\text{Eq. 3.5.1})$$

Fugitive emissions from extraction

According to the EMEP/EEA Guidebook (EMEP/EEA, 2009) the total fugitive emissions of volatile organic compounds (VOC) from extraction of oil and gas can be estimated by means of equation 3.5.2.

$$E_{extraction,VOC} = 40.2 \cdot N_p + 1.1 \cdot 10^{-2} P_{gas} + 8.5 \cdot 10^{-6} \cdot P_{oil} \quad (\text{Eq. 3.5.2})$$

where $E_{extraction,VOC}$ is the emission of VOC in Mg pr year, N_p is the number of platforms, P_{gas} is the production of gas, 10^6 Nm^3 and P_{oil} is the production of oil, 10^6 tonnes.

It is assumed that the VOC contains 75 % methane (CH_4) and 25 % NMVOC (EMEP/EEA, 2013), and in consequence the total emission of CH_4 and NMVOC for extraction of oil and gas can be calculated as:

$$E_{extraction,CH_4} = 0.75 \cdot E_{extraction,VOC} \quad (\text{Eq. 3.5.3})$$

$$E_{extraction,NMVOC} = 0.25 \cdot E_{extraction,VOC} \quad (\text{Eq. 3.5.4})$$

Loading of ships

Fugitive emissions of CH_4 and NMVOC from loading of ships include the transfer of oil from storage tanks or directly from the well into ships. The activity also includes losses during transport. When oil is loaded hydrocarbon vapour will be displaced by oil and new vapour will be formed, both leading to emissions. The emissions from ships are calculated by equation 3.5.5.

$$E_{ships} = EF_{ships,onshore} \cdot L_{oil,onshore} + EF_{ships,offshore} \cdot L_{oil,offshore} \quad (\text{Eq. 3.5.5})$$

where EF_{ships} is the emission factor for loading of ships offshore and onshore and L_{oil} is the amount of oil loaded.

Raw oil terminal

The CH_4 and NMVOC emissions from storage and handling of oil are given in the environmental reports from DONG Oil Pipe A/S for 2012 (DONG Oil Pipe A/S, 2013). An implied emission factor is calculated for use in the reporting template on the basis of the amount of oil transported in pipelines according to equation 3.5.6.

$$IEF_{tanks} = \frac{E_{tanks}}{T_{oil}} \quad (\text{Eq. 3.5.6})$$

where IEF_{tanks} is the implied emission factor for storage of raw oil in tanks, E_{tanks} is the emission and T_{oil} is the amount of oil transported in pipelines.

H5 Service stations

NMVOC emissions from service stations are estimated as outlined in equation 3.5.7.

$$E_{service stations} = (EF_{reloading} \cdot T_{fuel}) + (EF_{refuelling} \cdot T_{fuel}) \quad (\text{Eq. 3.5.7})$$

where $EF_{reloading}$ is the emission factor for reloading of tankers to underground storage tanks at the service stations, $EF_{refuelling}$ is the emission factor for refuelling of vehicles and T_{fuel} is the amount of gasoline used for road transport.

Oil refining

When oil is processed in the refineries, part of the volatile organic compounds (VOC) is emitted to the atmosphere. The VOC emissions from the petroleum refinery process include non-combustion emissions from handling and storage of feedstock (raw oil), from the petroleum product processing and from handling and storage of products. Emissions from flaring in refineries are included under "Flaring". Emissions related to process furnaces in refineries are included in stationary combustion with the relevant emission factors. When only the total VOC emission is given by the refinery the emission of CH₄ and NMVOC is estimated due to the assumption that 10 % of VOC is CH₄ and the remaining 90 % is NMVOC (Hjerrild, 1997).

Both the non-combustion processes including product processing and sulphur recovery plants emit SO₂. The SO₂ emissions are calculated by the refineries and implemented in the emission inventory without further calculation.

Fugitive emissions from gas (1B2b)

Transmission and distribution of gas

The fugitive emission from transmission, storage and distribution is based on information from the gas companies. The transmission and distribution companies give data on the transported amount and length and material of the pipeline systems. The fugitive losses from pipelines are provided by the transmission company. The natural gas and town gas distribution companies don't include fugitive losses in their annual reports, but only the total gas loss including measure uncertainty, erroneous debited customers and displacements between production periods and settlement period. The share of the total gas loss owing to fugitive losses is estimated based on further information from one of four Danish distribution companies and used for the remaining companies. From the fugitive losses of natural gas from transmission and distribution pipelines the emissions of CH₄ and NMVOC are calculated based on the gas quality measured by Energinet.dk. The same approach is used for town gas, which is natural gas admixed ~ 50 % ambient air.

Flaring

Emissions from flaring are estimated from the amount of gas flared offshore, in gas treatment/storage plants and in refineries and from the corresponding emission factors. From 2006 data on offshore flaring (flared amounts, calorific values and CO₂ emission factors) are given in the reports under the EU ETS and thereby flaring can be split to the individual production units. For previous years only the total flared amount are available.

3.5.3 Activity data

Extraction of oil and gas and loading of ships

Activity data used in the calculations of the emissions from oil and gas production and loading of ships are shown in Table 3.5.6. Data are based on information from the Danish Energy Agency (2013a) and from the environmental reports from DONG Oil Pipe A/S (DONG Oil Pipe A/S, 2013).

Table 3.5.6 Activity data for 2012.

Activity	Symbols	Amounts	Data source
Number of platforms	N_p	54	Danish Energy Agency, 2013a
Produced gas, 10^6 Nm^3	P_{gas}	5 617	Danish Energy Agency, 2013a
Produced oil, 10^3 m^3	$P_{\text{oil,vol}}$	11 728	Danish Energy Agency, 2013a
Produced oil, 10^3 tonnes	P_{oil}	10 086	Danish Energy Agency, 2013a
Oil loaded, 10^3 m^3	$L_{\text{oil offshore}}$	1 549	Danish Energy Agency, 2013a
Oil loaded, 10^3 tonnes	$L_{\text{oil offshore}}$	1 332	Danish Energy Agency, 2013a
Oil loaded, 10^3 m^3	$L_{\text{oil on-shore}}$	10 500	DONG Oil Pipe A/S, 2013
Oil loaded, 10^3 tonnes	$L_{\text{oil on-shore}}$	9 030	DONG Oil Pipe A/S, 2013

Density of crude oil = 0.86 tonnes per m^3

As seen in Figure 3.5.1 the production of oil and gas in the North Sea has generally increased in the years 1990-2004. Since 2004 the production has decreased. The number of platforms is yet still increasing (Figure 3.5.2). Five major platforms were completed in 1997-1999, which is the main reason for the great increase in the oil production in the years 1998-2000.




Figure 3.5.1 Production of oil and gas in the Danish part of the North Sea.




Figure 3.5.2 The number of platforms in the Danish part of the North Sea.

The amounts of oil loaded offshore on ships roughly follow the trend of the oil and gas production (Figure 3.5.3). In case of onshore loading of ships the trend is more smoothed.



Figure 3.5.3 Onshore and offshore loading of ships.

Oil refining

Data on the amount of crude oil processed in the two Danish refineries are given by the refineries in their annual environmental report (A/S Dansk Shell, 2013 and Statoil A/S, 2013). Until 1996 a third refinery was in operation, leading to a decrease in the processed crude oil amount from 1996 to 1997. Data are shown in Figure 3.6.4. The amount of crude oil being processed was 7 894 Gg in 2012.




Figure 3.5.4 Oil refineries. Processed crude oil in Danish refineries.

Service stations

The Danish Energy statistics contains data on the sale of gasoline that are the basis for estimating emissions of NMVOC from service stations. The gasoline sales show an increase from 1990-1998 and a decreasing trend since 1999 as shown in Figure 3.5.5. In 2012 the gasoline sale was 1 389 Gg.




Figure 3.5.5 Gasoline sales in Denmark.

Transmission, storage and distribution of gas

The activity data used in the calculation of the emissions from natural gas are shown in Table 3.5.7. Transmission rates for 1990-1998 refer to annual environmental reports of DONG Energy. In 1999-2006 transmission rates refer to the Danish Gas Technology Centre (Karll 2003, Karll 2005, Oertenblad 2006, Oertenblad 2007). From 2008 onwards transmission rates refer to Energinet.dk. Transmission losses for 1991-1999 are based on annual environmental report of DONG Energy. The average for 1991-1995 is applied for 1990. From 2005 onwards transmission losses are given by Energinet.dk. The average for 2005-2010 is applied for the years 2000-2004.

Distribution rates for 1990-1998 are estimated from the Danish energy statistics. Distribution rates are assumed to equal total Danish consumption rate minus the consumption rates of sectors that receive the gas at high pressure. The following consumers are assumed to receive high pressure gas: town gas production companies, production platforms and power plants. In 1999-2006 distribution rates refer to DONG Energy/Danish Gas Technology Centre/Danish gas distribution companies (Karll 2003, Karll 2005, Oertenblad 2006, Oertenblad 2007). Since 2007 the distribution rates are given by the companies. Distribution rates for town gas is based on the available data from the Danish town gas distribution companies of which more are closed down today. Distribution losses for 1990-2000 are based on annual environmental report of DONG Energy. For 2000-2006 the average losses-% for the gas distribution companies are used. From 2007 data on distribution losses available from the companies are used.

Table 3.5.7 Activity data on transmission and distribution of gas for selected years of the time series. Town gas is included in distribution.

	1990	1995	2000	2005	2008	2009	2010	2011	2012
Transmission, Mm ³ *	2 739	4 689	7 079	7 600	7 565	6 500	7 462	6 181	5 365
Distribution of natural gas, Mm ³ **	1 714	3 054	3 181	3 265	3 113	2 870	3 416	2 933	2 728
Distribution of town gas, Mm ³ ***	35	35	34	32	22	20	22	21	24

* Transmission rates for 1990-1998 refer to the annual environmental report of DONG Energy. In 1999-2006 transmission rates refer to the Danish Gas Technology Centre (Karll 2003, Karll 2005, Oertenblad 2006, Oertenblad 2007). From 2008 onwards transmission rates refer to Energinet.dk.

**) In 1990-98 distribution rates are estimated from the Danish energy statistics. Distribution rates are assumed to equal total Danish consumption rate minus the consumption rates of sectors that receive the gas at high pressure. The following consumers are assumed to receive high pressure gas: town gas production companies, production platforms and power plants. In 1999-2006 distribution rates refer to DONG Energy / Danish Gas Technology Centre / Danish gas distribution companies (Karll 2003, Karll 2005, Oertenblad 2006, Oertenblad 2007). Since 2007 the distribution rates are given by the companies.

***) The distribution of town gas is based on the available data from the Danish town gas distribution companies of which more are closed down today.

In 2012 the gas transmission was 5 365 Mm³ and the distribution rate is 2 752 Mm³, hereof 24 Mm³ town gas (Figure 3.5.6). The variation over the time series owes mainly to variations in the winter temperature and to the variation of import/export of electricity from Norway and Sweden. The transmission rate is less than the production rate, as part of the produced natural gas is exported through the NOGAT pipeline system.




Figure 3.5.6 Rates for transmission and distribution of gas. Distribution cover both natural gas and town gas.

Data on the transmission pipelines excluding offshore pipelines and on the distribution network are given by Energinet.dk, DGC and the distribution companies concerning length and material. The length of the transmission pipelines is approximately 900 km. Because the distribution system in Denmark is relatively new most of the distribution network is made of plastic (PE). In 2012 the length of the distribution network was around 20 000 km. The major part is made of plastic (approximately 90 %) and the remaining part is made of steel. For this reason the fugitive emission is negligible under normal operating conditions as the distribution system is basically tight with no fugitive losses. However, the plastic pipes are vulnerable and therefore most of the fugitive emissions from the pipes are caused by losses due to excavation damages and construction and maintenance activities performed by the gas companies. These losses are either measured or estimated by cal-

culuation in each case by the gas companies. About 5 % of the distribution network is used for town gas. This part of the network is older and the fugitive losses are greater. The fugitive losses from this network are associated with more uncertainty as it is estimated as a percentage (15 %) of the meter differential. This assumption is based on expert judgement from one of the town gas companies. It must be noted that two town gas distribution companies have been closed in recent years (one in 2004 and another in 2006). There are only two town gas distribution companies left, and therefore the data availability is scarce.

Venting and flaring

Venting

In Denmark there are two natural gas storage facilities. Both are obligated to make an environmental report on annual basis. Data on gas input and withdrawal are included and were 445 Mm³ and 634 Mm³ in 2012, respectively. Venting and flaring at the gas storage plants are included in the inventory. Venting of gas is assumed to be not occurring in extraction and in refineries as controlled venting enters the gas flare system. Venting rates in gas storage facilities are shown in Figure 3.4.8. As venting rates are not available for the years 1990-1994, the average for 1995-1998 is used.




Figure 3.5.7 Amount vented in gas treatment and storage plants.

Flaring

Flaring rates for the two Danish refineries are given in their environmental reports and in additional data provided by the refineries directly to DCE. From 2006 flaring amounts are given in the EU ETS reporting. Activity data for flaring in refineries are shown in Figure 3.5.8. Data are not available for the years 1990-1993. The flaring amount for 1994 has been adopted for the previous years. Use of a mean value for the following five or ten years as applied for e.g. flaring in storage and treatment plants are not appropriate in this case, as one of three refineries was closed down in 1996.



Figure 3.5.8 Amount flared in refineries (annual environmental reports from A/S Dansk Shell and Statoil A/).

Offshore flaring amounts are given in Denmark's oil and gas production (Danish Energy Agency, 2013a) while flaring in treatment/storage plants are given in environmental reports by Dong Energy and Energinet.dk (Dong Energy, 2013a,b; Energinet.dk, 2013a). Flaring rates are shown in Figure 3.5.9 and 3.5.10.

Flaring rates in gas treatment and gas storage plants are not available until 1994. The mean value for the following five years (1994 to 1998) has been adopted as basis for the emission calculation for the years 1990-1993. The large amount of flared gas in 2007 owe to a larger maintenance work at the gas treatment plant.




Figure 3.5.9 Amount flared in gas treatment and storage plants.

Offshore flaring amounts have been decreasing over the last 10 years period in accordance with the decrease in production as seen in Figure 3.5.1. Further, there is focus on reduction of the amount being flared for environmental reasons.




Figure 3.5.10 Amounts of gas flared in offshore exploration.

3.5.4 Emission factors

Extraction of oil and gas and loading of ships

Emissions of CH₄ and NMVOC from extraction of oil and gas are calculated from standard formula as listed in equation 3.5.3 and 3.5.4. Emissions of CO₂ from extraction of oil and gas and from transport of oil in pipelines are based on the emission factor from IPCC Good Practice Guidance (IPCC, 2000). IEF for extraction and pipeline oil are shown in Table 3.5.8.

Table 3.5.8 Implied emission factors for CO₂, CH₄ and NMVOC for extraction and storage/transport of oil.

	CO ₂ , g/Mg crude oil	CH ₄ , g/Nm ³ crude oil transported via pipeline	NMVOC, g/Nm ³ crude oil transported via pipeline
Extraction *	367	24.0	8.0
Pipeline oil **	0.53	70.3	54.5

* Based on EMEP/EEA, 2009 and Danish Energy Agency, 2012a.

** Default EF for CO₂ from IPCC, 2000, and IEF for CH₄ and NMVOC from DONG Oil Pipe A/S, 2013.

In the EMEP/EEA Guidebook standard emission factors for different countries are given (EMEP/EEA, 2013). In the Danish emission inventory the Norwegian emission factors are used for estimation of fugitive emissions from loading of ships onshore and offshore for the years 1990-2009. During 2009 new emission reducing technologies (degassing unit) were installed at the crude oil terminal. Measurements were carried out at the terminal before and after installation show a decrease of 21 % of the CH₄ emission and 25 % of the NMVOC emission from loading of ships. The reduced emission factors used for 2010 onwards are listed in Table 3.5.9.

Table 3.5.9 Emission factors for loading of ships onshore and offshore.

	CH ₄ , fraction of loaded		NMVOC, fraction of loaded	
	1990-2009	2010 onwards	1990-2009	2010 onwards
Ships offshore *	0.00005	0.00005	0.001	0.001
Ships on-shore **	0.00001	0.0000079	0.0002	0.00015

* EMEP/EEA, 2013.

** EMEP/EEA, 2013; Miljøcenter Odense, 2010.

Oil refining

The refineries deliver information on consumption of fuel gas and fuel oil. The calorific values are given by the refineries in the reporting to the EU ETS from 2006. Before 2006 the calorific values given by the refineries were used when available. When not available standard calorific values given in the basic data tables from the Danish Energy Agency combined with the conversion factor between fuel gas and fuel oil given by the refinery were used for calculation.

Emissions of SO₂, NO_x and VOC are given by the refineries. Only one of the two refineries has made a split between NMVOC and CH₄. For the other refinery it is assumed that 10 % of the VOC emission is CH₄ and the remaining 90 % is NMVOC (Hjerrild, 1997).

Service stations

The NMVOC emission from service stations is calculated by use of different emission factors for the time series as shown in Table 3.5.10. In 1994 the emission factors for NMVOC from service stations were investigated by Fenhann and Kilde (1994) for the years 1990, 1991 and 1992, individually. The emission factors reported for reloading and refuelling for 1990 were used for the years 1985-1990, while the emission factors for 1991 was used for that year only. For the years 1992-1995 only emission factor for refuelling reported by Fenhann and Kilde (1994) was used in the Danish emission inventory. For reloading of tankers the British emission factor - as given in the UK Emission Factor Database - was adopted for the years 1992-2000. From 2008 the emission factors from the EMEP/EEA guidebook 2009 are used for reloading and refuelling. For the years 2001-2007 and 1996-2007 the emission factors for reloading and refuelling, respectively, are estimated by using interpolation.

Table 3.5.10 Emission factors used for estimating NMVOC from service stations.

Year	Reloading of tankers, kg NMVOC pr tonnes gasoline	Refuelling of vehicles, kg NMVOC pr tonnes gasoline	Sum of reloading and refuelling, kg NMVOC pr tonnes gasoline	Source (reloading/refuelling)
1985-1990	1.28	1.52	2.80	Fenhann & Kilde,1994 / Fenhann & Kilde,1994
1991	0.64	1.52	2.16	Fenhann & Kilde,1994 / Fenhann & Kilde,1994
1992-1995	0.08	1.52	1.60	UK emf. database / Fenhann & Kilde,1994
1996	0.08	1.45	1.53	UK emf. database / interpolation 1995-2008
1997	0.08	1.39	1.47	UK emf. database / interpolation 1995-2008
1998	0.08	1.32	1.40	UK emf. database / interpolation 1995-2008
1999	0.08	1.25	1.33	UK emf. database / interpolation 1995-2008
2000	0.08	1.19	1.27	UK emf. database / interpolation 1995-2008
2001	0.077	1.12	1.20	Interpolation 2000-2008 / 1995-2008
2002	0.073	1.05	1.13	Interpolation 2000-2008 / 1995-2008
2003	0.070	0.99	1.05	Interpolation 2000-2008 / 1995-2008
2004	0.067	0.92	0.98	Interpolation 2000-2008 / 1995-2008
2005	0.063	0.85	0.91	Interpolation 2000-2008 / 1995-2008
2006	0.060	0.78	0.84	Interpolation 2000-2008 / 1995-2008
2007	0.056	0.72	0.77	Interpolation 2000-2008 / 1995-2008
2008 onwards	0.053	0.65	0.70	EMEP/EEA 2013 / EMEP/EEA 2013

Transmission, storage and distribution of gas

The fugitive emissions from transmission, storage and distribution of natural gas are based on data on gas losses from the companies and on the aver-

age annual natural gas composition given by Energinet.dk (Table 3.5.11). For distribution of town gas the emission factor is reduced due to the admixture of 50 % atmospheric air to the natural gas.

Table 3.5.11 Annual gas composition, lower heating value and density for Danish natural gas (Energinet.dk).

		Unit	1990	2000	2005	2008	2009	2010	2011	2012
Methane	CH ₄	molar-%	90.92	86.97	88.97	89.80	90.08	89.95	89.10	88.84
Ethane	C ₂ H ₆	molar-%	5.08	6.88	6.14	5.77	5.70	5.71	5.98	6.11
Propane	C ₃ H ₈	molar-%	1.89	3.17	2.50	2.26	2.17	2.19	2.36	2.44
i-Butane	i-C ₄ H ₁₀	molar-%	0.36	0.43	0.40	0.37	0.37	0.37	0.37	0.37
n-Butane	n-C ₄ H ₁₀	molar-%	0.50	0.61	0.55	0.53	0.52	0.54	0.55	0.54
i-Petane	i-C ₅ H ₁₂	molar-%	0.14	0.11	0.11	0.13	0.13	0.13	0.13	0.13
n-Petane	n-C ₅ H ₁₂	molar-%	0.10	0.08	0.08	0.08	0.08	0.08	0.09	0.08
n-Hexane and heavier hydrocarbons	C ⁶⁺	molar-%	0.09	0.06	0.05	0.06	0.06	0.06	0.06	0.06
Nitrogen	N ₂	molar-%	0.31	0.34	0.29	0.30	0.29	0.31	0.37	0.36
Carbon dioxide	CO ₂	molar-%	0.60	1.35	0.90	0.71	0.59	0.66	0.98	1.06
Lower heating value	H _n	MJ/m ³ _n	39.176	40.154	39.671	39.485	39.459	39.461	39.507	39.548
Density	pp	kg/m ³ _n	0.808	0.846	0.825	0.817	0.814	0.816	0.824	0.827

Venting and flaring

Venting

CH₄ and NMVOC emissions from venting are given in the environmental reports for the gas storage facilities (DONG Energy, 2013a; Energinet.dk, 2013a). CO₂ emissions from venting are calculated from country specific emission factors based on annual natural gas composition published by Energinet.dk.

Flaring in refineries

The composition of fuel gas is given for 2008 by one of the two refineries. As the composition for fuel gas is marked different than the composition of natural gas, which has been used in earlier year's calculations, the same fuel gas composition is used in calculations for the other Danish refinery.

The new emission factors for CH₄ and NMVOC based on fuel gas composition data from one refinery have been included in the inventory for all years from 1990 and onwards. The CO₂ emission factor is based on the refineries reporting to the EU ETS for the years 2006 and onwards. Before 2006 corresponding data are not available and the CO₂ emission factors are calculated from the annual natural gas composition given by Energinet.dk. NO_x emissions are provided by one refinery. Emission factors from the EMEP/EEA Guidebook (2013) are used to calculate emissions of CO and for NO_x where plant specific data are not available. The emission factor applied for N₂O is based on the OLF (1993) for flaring in oil and gas extraction as no value are given for flaring in refineries. The emission factors are listed in Table 3.5.12.

Table 3.5.12 Emission factors for flaring in refineries.

Pollutant	Emission factor	Unit
NO _x *	32.2	g per GJ
NMVOC	76.4	g per GJ
CH ₄	18.1	g per GJ
CO	177	g per GJ
CO ₂ **	51.83 / 58.59	kg per GJ
N ₂ O	0.47	g per GJ

* Direct measured emission of NO_x is available for one refinery and the emission factor is used for the remaining refinery only.

** The CO₂ emission is based on the refineries reports for ETS and is source specific.

Flaring offshore

The emission factors for offshore flaring are shown in Table 3.5.13. Since 2006 the CO₂ emission factor is calculated according to the reporting for EU ETS. Corresponding data are not available for earlier years and therefore the CO₂ emission factor is assumed to follow the same time series as for natural gas combusted in stationary combustion plants.

The NO_x emission factor is based on the conclusion in a Danish study of NO_x emissions from offshore flaring carried out by the Danish Environmental Protection Agency (2008). The recommended NO_x emission factor (31 g per GJ or 0.0015 tonnes NO_x per tonnes gas) corresponds well with the emission factors used to estimate NO_x emission in other countries with oil production in the North Sea (Netherlands: approximately 0.0014 tonnes NO_x per tonnes gas and United Kingdom: approximately 0.0013 tonnes NO_x per tonnes gas).

Emission factors for N₂O are based on IPCC (2000) and emission factors for NMVOC and CO are based on the EMEP/EEA Guidebook (2009). The emission factor for CH₄ is based on OLF (1993).

Emissions from flaring in gas treatment and storage plants are calculated from the same emission factors which are used for offshore flaring. Only difference is the CO₂ emission factor for the years from 2006. The emission factor used for the plants are based on the same data source, the reporting for EU ETS, but the values are different than for offshore flaring. The gas that are flared in the treatment and storage plants are natural gas with the same composition as natural gas distributed in Denmark. Therefore, the emission factors in the EU ETS reports are the same as the one calculated on basis of the gas composition given by Energinet.dk.

Table 3.5.13 Emission factors for offshore flaring in 2012.

Pollutant	Emission factor	Unit
SO ₂	0.014	g per Nm ³
NO _x	1.227	g per Nm ³
NMVOC	0.105	g per Nm ³
CH ₄	0.211	g per Nm ³
CO	1.055	g per Nm ³
CO ₂	2.707	kg per Nm ³
N ₂ O	0.021	g per Nm ³

3.5.5 Emissions

Extraction of oil and gas and loading of ships

From the activity data in Table 3.5.6, equation 3.5.3 and equation 3.5.4 the fugitive emissions of CH₄ and NMVOC from extraction are calculated. Corresponding emissions from loading of ships can be estimated by Table 3.5.6 and equation 3.5.5. The emissions are listed in Table 3.5.14 to Table 3.5.17 along with the emissions from oil pipelines and storage tanks given in the environmental reports from DONG Oil Pipe A/S (2013). CO₂ emissions from oil pipeline and storage tanks, as well as from extraction are calculated from standard emission factors (IPCC 2000).

Table 3.5.14 Emissions of CO₂, CH₄ and NMVOC from onshore loading of ships.

	1990	1995	2000	2005	2008	2009	2010	2011	2012
CO ₂ , Gg	NA	NA	NA	NA	NA	NA	NA	NA	NA
CH ₄ , Mg	34	62	109	125	96	86	63	56	71
NMVOC, Mg	678	1 249	2 183	2 494	1 926	1 720	1 187	1 071	1 355

Table 3.5.15 Emissions of CO₂, CH₄ and NMVOC from Oil pipeline and storage tanks.

	1990	1995	2000	2005	2008	2009	2010	2011	2012
CO ₂ , Gg	< 0.1	< 0.1	< 0.1	< 0.1	< 0.1	< 0.1	< 0.1	< 0.1	< 0.1
CH ₄ , Mg	783	1 209	1 700	2 100	1 648	1 300	984	822	766
NMVOC, Mg	1 726	2 664	4 000	4 500	3 625	2 098	763	638	594

Table 3.5.16 Emissions of CO₂, CH₄ and NMVOC from Fugitive emissions from extraction.

	1990	1995	2000	2005	2008	2009	2010	2011	2012
CO ₂ , Gg	2	4	7	7	5	5	5	4	4
CH ₄ , Mg	708	990	1 365	1 608	1 684	1 703	1 698	1 685	1 677
NMVOC, Mg	236	330	455	536	561	568	566	562	559

Table 3.5.17 Emissions of CO₂, CH₄ and NMVOC from Offshore loading of ships.

	1990	1995	2000	2005	2008	2009	2010	2011	2012
CO ₂ , Gg	NA	NA	NA	NA	NA	NA	NA	NA	NA
CH ₄ , Mg	NA	NA	201	167	93	73	83	76	67
NMVOC, Mg	NA	NA	4 021	3 337	1 856	1 451	1 658	1 525	1 332

Oil refining

In Table 3.5.18 the activity data and emissions of VOC from the Danish refineries are listed for selected years in the time series. Further, the emissions of SO₂ from oil refining and sulphur recovery in refineries are shown. The emission of SO₂ has shown a pronounced decrease since 1990 because of technical improvements at the refineries. Note that SO₂ from refining and recovery prior to 1994 was aggregated and reported as an area. Note also that SO₂ from oil refining from 2001 are included in stationary combustion.

The large fluctuations in the SO₂ emission from sulphur recovery owe to outage and shut downs of the sulphur recovery units. The SO₂ emission from sulphur recovery has shown a gradual decrease from 1995 to 2000.

Table 3.5.18 Oil Refineries. Emissions of NMVOC and SO₂ from oil refining and SO₂ from sulphur recovery.

	1990	1995	2000	2005	2008	2009	2010	2011	2012
Crude oil, 1000 Mg	7 263	9 802	8 508	8 033	7 933	7 978	7 414	6 997	7 414
VOC, Mg	3 667	5 815	4 845	3 442	3 588	3 877	3 867	3 868	3 932
SO ₂ , oil refining, Mg ^{1,2}	3 335	585	178	IE	IE	IE	IE	IE	IE
SO ₂ , sulphur recovery, Mg ¹		2 437	803	390	987	481	1 019	1 179	889

¹⁾ Prior to 1994 SO₂ emissions from oil refining and sulphur recovery are aggregated and reported as area.

²⁾ From 2001 SO₂ emissions from oil refining are included in stationary combustion.

Service stations

Emissions from service stations are calculated using the emission factors in Table 3.5.10 and the sold amounts of gasoline given by the Danish energy statistics (Danish Energy Agency, 2013b). The NMVOC emissions are listed in Table 3.5.19.

Table 3.5.19 Emissions of NMVOC from service stations for selected years of the time series.

	1990	1995	2000	2005	2008	2009	2010	2011	2012
NMVOC, Mg	4 856	3 016	2 616	1 742	1 216	1 158	1 106	1 027	977

Transmission, storage and distribution of gas

The gas transmission company gives emissions of CH₄ for the years 1999 and onwards, based on registered loss in the transmission grid and the emission from the natural gas consumption in the pressure regulating stations (Oertenblad, 2007). For the years 1991-1998 the CH₄ emissions for transmission are estimated on the basis of registered loss provided by the transmission company and the annual composition of Danish natural gas given by Energinet.dk. Transmission loss is not available for 1990, why the average for 1991-1995 is applied.

The distribution companies provide emissions of CH₄ for the years 1997 and onwards. For the years 1995-1996 CH₄ emissions are calculated from the registered loss from distribution and the annual composition of Danish natural gas given by Energinet.dk. As distribution losses are not available for the years 1990-1994, the percentage loss for 1995 is used.

The emissions of NMVOC are calculated on the basis of the CH₄ emission according to the gas quality measured by Energinet.dk (equation 3.5.8).

$$E_{NMVOC} = E_{CH_4} \times (w_{NMVOC} / w_{CH_4}) \quad (\text{Eq.3.5.8})$$

where w_{NMVOC} is the weight-% NMVOC and w_{CH_4} is the weight-% CH₄ according to the gas quality of the current year.

Emissions of CH₄ and NMVOC from transmission of natural gas (including storage) and distribution of natural gas and town gas are shown in Table 3.5.20 and Table 3.5.21, respectively. Emissions of CO₂ from transmission and distribution are very limited amounts and therefore not included in the tables.

As the pipelines in Denmark are relatively new and made of plastic, emissions other than from construction and maintenance are considered not occurring. The decrease in emission from transmission in 2007 is caused by the

completion of a greater construction work and rerouting of a major pipeline. In preparation for construction work on a new compressor station, there has been laid a number of new line valve stations in 2011. Before this work could be done, larger amounts of natural gas were vented to drain the pipes. Therefore emissions from transmission of natural gas are significantly higher in 2011 than in the previous years.

Emissions from distribution of gas mainly owe to excavations and maintenance of the pipelines, but also difference between the calendar year and the meter reading year might influence the annual variations. As the town gas distribution network is significant older the gas losses and thus the emissions are larger than for the natural gas distribution network, even though the distribution rates for natural gas far exceeds the rates for town gas.

Data on distribution of town gas is limited as more distribution companies are closed. For the active companies, data are available for part of the time series only. Gap filling and extrapolation of distribution and loss amounts to early years are done in collaboration with the individual companies to the extent possible.

Table 3.5.20 CH₄ emission from transmission of natural gas and distribution of natural gas and town gas for selected years.

CH ₄ , Mg	1990	1995	2000	2005	2008	2009	2010	2011	2012
Transmission	190	597	86	141	16	9	26	171	14
Distribution, natural gas	56	99	49	66	51	49	37	49	71
Distribution, town gas	202	211	178	176	95	85	106	106	75

Table 3.5.21 NMVOC emission from transmission of natural gas and distribution of natural gas and town gas for selected years.

NMVOC emission, Mg	1990	1995	2000	2005	2008	2009	2010	2011	2012
Transmission	41	135	26	36	4	2	6	43	4
Distribution, natural gas	12	22	15	17	12	12	9	12	18
Distribution, town gas	683	700	680	587	56	48	53	50	39

Venting and Flaring

Venting

Venting is limited to the gas storage plants and thus the emissions are of minor importance (Table 3.5.5). The emissions of CH₄ and NMVOC from venting are given in the environmental reports for the gas storage plants (DONG Energy, 2012a,b; Energinet.dk, 2012a). Calculation of CO₂ emissions are based on venting amounts and annual composition of Danish natural gas. Venting emissions are included in Figure 3.5.12 and Figure 3.5.13.

Flaring

As shown in Figure 3.5.11 there was a marked increase in the amount of offshore flaring in 1997 and especially in 1999. The increase in 1997 was due to the new Dan field and the completion of the Harald field. The increase in 1999 was due to the opening of the three new fields Halfdan, Siri and Syd Arne.

The time series for the emission of CO₂ from offshore flaring fluctuates due to the fluctuations in the fuel rate and to a minor degree due to the CO₂ emission factor. The latter is based on gas quality measurements. From 2006 the calorific values for flare gas are given at installation level in the EU ETS.

This information is incorporated in the inventory for the years 2006-2007 for part of the offshore installations, and from 2008 and onwards for all installations. This has led to an increase of the CO₂ emission factor. The average of the emission factors for 2008-2010 is adopted for 1990-2007. Fuel rate and CO₂ emission are shown in Figure 3.5.11.




Figure 3.5.11 Fuel rate and CO₂ emission from offshore flaring of gas.

The emissions of other pollutants than CO₂ from offshore flaring are estimated from the same emission factors for all years and the variations reflect only the variations in the flared amounts. Emissions of CH₄ and NMVOC from flaring and venting are shown in Figure 3.5.12 and 3.5.13.




Figure 3.5.12 CH₄ emissions from venting and flaring of gas.




Figure 3.5.13 NMVOC emissions from venting and flaring of gas.

The decrease in the emissions from 2009 to 2010 for gas storage and treatment plants owes to a change from continuous to regulating power operation of the power producing gas turbine at the gas storage plant.

Table 3.5.22 Emissions from flaring offshore and in gas treatment/storage plants.

Year	1990	1995	2000	2005	2008	2009	2010	2011	2012
	Mg	Mg	Mg	Mg	Mg	Mg	Mg	Mg	Mg
CO ₂	302	423	695	513	360	244	333	232	195
SO ₂	2	2	4	3	2	1	2	1	1
NO _x	134	188	310	231	168	111	155	103	96
NMVOC	15	20	30	23	20	23	13	9	8
CO	115	161	265	195	141	91	127	83	76

Flaring also occurs in refineries. Flaring in refineries is a significant fugitive emission source for SO₂. In 1990-1993, emissions from petroleum product processing were included in emissions from flaring in refineries (1B2c). From 1994 the data delivery format was changed, which made it possible to split the emissions into contributions from flaring and processing, respectively. Emissions from processing are from 1994 included in the category Oil, Refining/Storage.

The decreasing emissions of SO₂ from 1995 to 1998 are due to technical improvements of the sulphur recovery system at one of the two Danish refineries (Table 3.5.23). The increase in SO₂ from flaring in refineries in 2005 and 2007 was due to planned shutdowns due to inspection and maintenance at one of the two refineries. Further, in 2007-2009 the same refinery has had problems with the ATS system leading to an increased SO₂ emission from flaring.

Table 3.5.23 Emissions from flaring in refineries.

	1990*	1995	2000	2005	2008	2009	2010	2011	2012
	Mg	Mg	Mg	Mg	Mg	Mg	Mg	Mg	Mg
CO ₂ , Gg	23	23	19	23	27	16	19	19	22
SO ₂ *, Mg	943	203	51	296	380	453	288	242	167
NO _x , Mg	13	13	11	26	26	17	19	18	23
NMVOC, Mg	31	31	26	32	38	23	27	25	32
CO, Mg	73	73	60	73	88	53	62	59	74

*In 1990-1993 emissions from petroleum product processing were included in flaring in refineries due to the data delivery form.

3.5.6 Uncertainties and time series consistency

Two set of uncertainty estimates are made for the Danish emission inventory for greenhouse gases based on Tier 1 and Tier 2 methodology, respectively. The uncertainty models follow the methodology in IPCC Good Practise Guidance (IPCC, 2000). Tier 1 is based on the simplified uncertainty analysis (error propagation method) and Tier 2 is based on Monte Carlo simulations.

Uncertainty estimates are made for total emissions in the base year (only Tier 2), in the latest inventory year and for the emission trend for the corresponding time series. Uncertainty estimates are made for the GHGs separately and summarized.

Input data

The Tier 1 uncertainty model is based on emission data, uncertainty levels for activity data and uncertainty levels for emission factors for base year and latest inventory year. The Tier 2 model is based on activity data and emission factors for the same years and the same uncertainty levels as in Tier 1. Emission data, activity data and emission factors are described in Chapter 3.5.3, 3.5.4 and 3.5.5.

The uncertainty levels used in the uncertainty models are based on different sources, e.g. IPCC Good Practice Guidance, EMEP/EEA Guidebook and reports under the EU ETS. Further, a number of the uncertainty levels are given as NERI assumptions. DCE assumptions are based on source and/or plant specific uncertainty levels for part of the SNAP category and assumptions for the remaining sources and/or plants in the category.

Input data are aggregated on SNAP level. Estimates are made for the greenhouse gases CO₂, CH₄ and N₂O both separately and summarized (GHG). Uncertainty levels for activity data and emission factors are listed in Table 3.5.24. Uncertainty levels are given in percentage related.

Table 3.5.24 Uncertainty levels for activity rates and emission factors.

Pollutant	Source	Activity data uncertainty level, %	Emission factor uncertainty level, %
CO ₂	Land based activities	2N	40S
CO ₂	Offshore activities	2N	30I
CO ₂	Gas transmission	15G	2Q
CO ₂	Gas distribution	25G, N	10Q, N
CO ₂	Venting in gas storage plants	25G, N	2Q
CO ₂	Flaring in refineries	11E	2E
CO ₂	Flaring offshore	7.5E	2E
CH ₄	Petroleum product processing	1E, N	125N
CH ₄	Land based activities	2N	40S
CH ₄	Offshore activities	2N	30I
CH ₄	Gas transmission	15G	2Q
CH ₄	Gas distribution	25G, N	10Q, N
CH ₄	Venting in gas storage plants	15G, N	2Q
CH ₄	Flaring in refineries	11E	15H, N
CH ₄	Flaring offshore	7.5E	125G
N ₂ O	Flaring in refineries	11E	1 000I
N ₂ O	Flaring offshore	7.5E	1 000I

N: DCE assumption.

I: IPCC Good Practice Guidance (default value).

S: Statistisk Sentralbyrå, Statistics Norway, 2008.

E: EU Emission Trading Scheme (EU ETS).

G: EMEP/EEA Guidebook, 2009.

H: Holst, 2009 and Statoil A/S, 2010.

Q: Annual gas quality, Energinet.dk.

The CO₂ emission factors for flaring offshore and in refineries and the CO₂ and CH₄ emission factors for natural gas transmission, distribution and venting, are the most accurate as they are calculated on basis of gas composition measurements. Emissions factors for flare gas are available in the EU-ETS reporting while emissions factors for natural gas are published by Energinet.dk.

The CO₂ emission factor for offshore activities is based on standard emission factors from IPCC (2000). Source specific uncertainty levels are not included in IPCC (2000), but it is written that the uncertainty levels must be assumed to be in the range of $\pm 25\%$ to $\pm 50\%$ for most gases. For onshore activities, the emission factor uncertainty corresponds to the uncertainty for onshore loading by Statistics Norway (2008), and the same uncertainty level is assumed for the CH₄ emission factor for onshore activities.

Data from the Danish operators (one year only) indicate that the VOC emissions in the Danish inventory have an uncertainty around 30 %, which has been used as uncertainty level for CH₄ in the uncertainty model. The EMEP/CORINAIR Guidebook (2007) suggests an error of 65 % for the standard equation used to estimate fugitive emissions of VOC from extraction, noting that the error could be much higher when the equation is used for other fields than the ones in USA, which it has been based on. Further the EMEP/EEA Guidebook (2009) says that the uncertainty level of 65 % seems to be in reasonable agreement with estimates for Norway and UK – countries expected to have more similar conditions to Danish than the USA. The EMEP/EEA uncertainty level is in the same order of magnitude as the Danish uncertainty level, which support the assumption that the Danish estimate are applicable for all years in the time series.

The uncertainty level for the emission factor for fugitive CH₄ emissions from refineries is dominated by a large uncertainty for one refinery. Further, measurements of fugitive emissions from the refineries are only available for one and two years, respectively, and these measurements indicate larger emissions than earlier estimates. As more measurements become available the uncertainty level is expected to decrease significantly.

According to IPCC (2000) the emission factor for N₂O is the least reliable. An uncertainty level of 1 000 % is adopted in the Danish uncertainty model.

The Tier 2 uncertainty model is based on Monte Carlo simulations and the input uncertainty levels are given for the 95 % confidence interval assuming a log-normal distribution. The input uncertainty levels are the same as those used in the Tier 1 uncertainty model (Table 3.5.20). For more information on the Tier 2 methodology, please see Chapter 1.7.

Results

The results of the Tier 1 uncertainty model for 2012 are shown in Table 3.5.25. In 2012 N₂O has the largest uncertainty for the total emission followed by CH₄ and CO₂. Due to the emission trend CH₄ has the largest uncertainty followed by N₂O and CO₂. The estimated uncertainty for the total GHG emission is 19 % and the GHG emission trend is -12 % \pm 17 %-point.

Table 3.5.25 Uncertainty estimates for total emissions and emission trends from the Tier 1 uncertainty model.

	Emission, Gg CO ₂ eqv Base year	Emission, Gg CO ₂ eqv 2011	Uncertainty, % Lower and upper (±)	Trend 1990-2012, %	Uncertainty, % Lower and upper (±)
CO ₂	325	217	8	-33	6
CH ₄	44	106	57	139	128
N ₂ O	1	1	892	-31	37
GHG	372	327	19	-12	17

Table 3.5.26 show the results from the Tier 2 uncertainty model for 1990 and 2012. The overall emission uncertainty in 2012 is -12/+30 %. The Tier 2 trend estimate is -11 % -20/+29 %-point.

Table 3.5.26 Uncertainty estimates for total emissions in 1990 and 2012 and for the emission trends from the Tier 2 uncertainty model.

	1990			2012			1990-2012		
	Median emission Gg CO ₂ eqv	Uncertainty, % Lower (-) Upper (+)		Median emission Gg CO ₂ eqv	Uncertainty, % Lower (-) Upper (+)		Median trend, %	Uncertainty, % Lower (-) Upper (+)	
CO ₂	328	15	17	221	7	7	-240	139	116
CH ₄	44	19	28	108	34	91	19	11	30
N ₂ O	1	94	1 193	1	93	1 024	-26	413	26
GHG	374	13	16	330	12	27	-11	20	29

Tier 1 and Tier 2 emissions and uncertainties are shown together in Figure 3.5.14. The figures show that the emissions and median emissions from Tier 2 are very similar. Further, the uncertainty estimates are in the same range for Tier 1 and Tier 2. The N₂O uncertainty is leaved out of Figure 3.5.14 b as the N₂O uncertainties are much higher than for CO₂ and CH₄. It must be noted that the uncertainty models, especially the Tier 1 model, are not suitable for very large uncertainty levels and therefore the uncertainty estimates for N₂O may only be seen as an indicator for a large uncertainties while the values are less accurate. The Tier 2 model has been developed to be more suitable for very large uncertainties, as it is possible to apply truncation for uncertainties. This has been included in the uncertainty calculation for fugitive emissions in case of N₂O, as the uncertainty level for the emission factors is 1 000 %. A truncation of 2 000 % has been applied to ensure that the emission factor interval is within an order of magnitude as given in IPCC Good Practice Guidance.




Figure 3.5.14 Emissions and uncertainty estimates from the Tier 1 and Tier 2 models; a) GHG, CH₄, CO₂ and N₂O, b) as figure a, but without N₂O

3.5.7 Source specific QA/QC and verification

The elaboration of a formal QA/QC plan started in 2004 and has recently been updated (Nielsen et al., 2013). The plan describes the concepts of quality work and definitions of sufficient quality, critical control points and a list of Points of Measuring, PM (Figure 3.5.15). Please refer to the general Chapter 1.6 for further information.




Figure 3.5.15 The general data structure for the Danish emission inventory (Nielsen et al., 2013).

Data storage level 1

Data storage level 1 refers to the data collected by DCE before any processing or preparing. Table 3.5.27 lists the external data deliveries used for the inventory of fugitive emissions. Further the table holds information on the contacts at the data delivery companies.

Table 3.5.27 List of external data sources.

Category	Data description	Activity data, emission factors or emissions	Reference	Contact(s)	Data agreement/ Comment
Offshore activities	Gas and oil production. Dataset for Activity data production of oil, gas and number of platforms. Amounts of offshore loading of ships	Activity data	The Danish Energy Agency	Jan H. Andersen	Not necessary due to obligation by law
Offshore flaring	Flaring offshore in oil and gas extraction	Activity data	The Danish Energy Agency	Dorte Maimann	Data agreement
Service stations	Data on gasoline sales from the Danish energy statistics.	Activity data	The Danish Energy Agency	Jane Rusbjerg	Data agreement
Gas transmission	Natural gas from the transmission	Activity data	Energinet.dk	Christian Friberg B.	Not necessary due

	company, sales and losses (meter differences)			Nielsen	to obligation by law
Onshore activities	Amounts of oil transport in pipeline and onshore loading to ships. Emissions from storage of raw oil in the terminal.	Activity data and emission data	DONG Oil Pipes A/S	Stine B. Bergmann	No formal data agreement.
Gas distribution	Natural gas from the distribution company, sales and losses (meter differences)	Activity data	Naturgas Fyn, HMN DONG Energy	Hanne Mochau Erik Wolf-Petersen Finn Adser	No formal data agreement.
Air emissions from refinery	Fuel consumption and emission data.	Activity data and emission data	Statoil A/S, A/S Danish Shell	Anette Holst, Lis Rønnow Rasmussen	No formal data agreement.
Storage and treatment of gas	Environmental reports from plants defined as large point sources (Lille Torup, Stenlille, Nybro)	Activity data	Various plants		Not necessary due to obligation by law
CO ₂ emission factors for different sources	Reports according to the CO ₂ emission trading scheme (ETS)	Activity data	Various plants		Not necessary due to obligation by law
Emission factors	Emission factors origin from a large number of sources	Emission factors	See chapter regarding emission factors		

Data Storage level 1	1. Accuracy	DS.1.1.1	General level of uncertainty for every dataset including the reasoning for the specific values.
----------------------	-------------	----------	---

The uncertainty for every dataset included in the inventory of fugitive emissions are evaluated and included in the Tier 1 and Tier 2 uncertainty calculations with short descriptions of the reasoning that underlie the specific values.

The general levels of uncertainty are relatively low. The largest uncertainties are expected for emissions from refineries and distribution of town gas, the latter being of minor importance to the total fugitive emissions.

For further comments regarding uncertainties, see Chapter 3.5.6.

Data Storage level 1	2. Comparability	DS.1.2.1	Comparability of the emission factors/calculation parameters with data from international guidelines, and evaluation of major discrepancies.
----------------------	------------------	----------	--

Systematic inter-country comparison has only been made on Data Storage Level 4. Refer to DS 4.3.2.

Data Storage level 1	3. Completeness	DS.1.3.1	Ensuring that the best possible national data for all sources are included, by setting down the reasoning behind the selection of datasets.
----------------------	-----------------	----------	---

External data sources are the Danish Energy Agency, EU ETS reports and annual environmental reports from plants which are obligated to publish environmental reports. Further, annual reports from the gas distribution companies and the raw oil terminal are used. Some environmental reports and annual reports are supplemented with data and information from the given companies.

Only one national data set is found for most fugitive sources, and all data set is expected to be complete and include all activities/emissions from the source. Data on flaring offshore, in refineries and in gas storage and treatment plants are available both in annual environmental reports and in EU-ETS reports. Data are compared and if any differences occur, this is checked with the data supplier. Minor differences may owe to the allocation of fuels, e.g. if pilot gas are included in the flare gas or the fuel gas amount.

Energy statistics

The Danish Energy Agency reports fuel consumption statistics on the SNAP level based on a correspondence table developed in co-operation with DCE. Both traded and non-traded fuels are included in the Danish energy statistics. Data on offshore extraction, offshore flaring and gasoline sales are used for estimation of fugitive emissions.

Environmental reports

A large number of plants are obligated by law to publish an environmental report annually with information on fuel consumption and emissions, among other things. DCE compares data with those from previous years, discrepancies are checked and large fluctuations are verified.

Annual reports

The gas distribution companies and the raw oil terminal are not obligated to publish environmental reports. Instead the self-regulation reports, annual reports and/or additional data and information are used. All information is compared with previous years.

Reports for the European Union Greenhouse Gas Emission Trading System (EU ETS)

CO₂ emission factors for flaring offshore and in refineries are taken from the EU ETS reports since 2006 when the EU ETS reports became available. EU ETS reports are available for the individual Danish oil/gas production fields and for the refineries.

Emission factors from a wide range of sources

For specific references, see Chapter 3.5.4 regarding emission factors.

Data Storage level 1	4.Consistency	DS.1.4.1	The original external data has to be archived with proper reference.
----------------------	---------------	----------	--

All external data are stored in the inventory file system and are accessible for all inventory staff members. Data processing is carried out in separate spread sheets or databases to ensure that the external data are always available in the original form. Refer to Section 1.3.

Data Storage level 1	6.Robustness	DS.1.6.1	Explicit agreements between the external institution holding the data and DCE about the conditions of delivery
----------------------	--------------	----------	--

Formal agreements are made with the Danish Energy Agency. Annual environmental reports are available due to legal requirements in this regard. The remaining data are published or delivered by the companies on voluntary basis. See Table. 3.5.26

Data Storage	7.Transparency	DS.1.7.1	Listing of all archived datasets and external
--------------	----------------	----------	---

level 1			contacts.
---------	--	--	-----------

See DS 1.3.1 and Table 3.5.26

Data Processing Level 1

Data Processing level 1	1. Accuracy	DP.1.1.1	Uncertainty assessment for every data source not part of DS.1.1.1 as input to Data Storage level 2 in relation to type and scale of variability.
-------------------------	-------------	----------	--

Refer to Section 1.7 in the Danish NIR and the QA/QC Section 3.5.7.

Data Processing level 1	2. Comparability	DP.1.2.1	The methodologies have to follow the international guidelines suggested by UNFCCC and IPCC.
-------------------------	------------------	----------	---

The methodologies in the inventory follow the principles in international guidelines by UNFCCC and IPCC.

Data Processing level 1	3. Completeness	DP.1.3.1	Identification of data gaps with regard to data sources that could improve quantitative knowledge.
-------------------------	-----------------	----------	--

Data gaps are found for distribution of town gas, as more companies are closed before the source was included in the Danish inventory. Emissions, which account for only a limited part of the total fugitive emissions, are calculated on a scarce data foundation.

More detailed data on emissions from exploration of oil and gas would be preferred, even though emissions calculated by use of the standard IPCC formula is in good agreement with an inventory made by the operators for one emission year. Unfortunately the inventory by the operators is not available for other years.

Regarding the VOC emissions from refineries, more detailed data material would be preferred.

Data Processing level 1	4. Consistency	DP.1.4.1	Documentation and reasoning of methodological changes during the time series and the qualitative assessment of the impact on time series consistency.
-------------------------	----------------	----------	---

Since 2006 the EU-ETS data have been available for a number of sources. In all cases the new data replace use of data assumed to be less accurate. Therefore the CO₂ emission factors have been updated for all years, and no methodological change occur in the time series.

A change in the calculating procedure would entail elaboration of an updated description in Chapter 3.5.

Data Processing level 1	5. Correctness	DP.1.5.2	Verification of calculation results using time series
-------------------------	----------------	----------	---

Time series for activity data on SNAP level as well as emission factors is used to identify possible errors in the calculation procedure.

Data Processing level 1	5.Correctness	DP.1.5.3	Verification of calculation results using other measures
-------------------------	---------------	----------	--

Emissions from offshore activities are compared to an inventory made by the operators and the emissions are found in good agreement.

For the remaining sources only one data set is available for calculation, and no verification using other measures are possible.

Data Processing level 1	7.Transparency	DP.1.7.1	The calculation principle, the equations used and the assumptions made must be described.
-------------------------	----------------	----------	---

Descriptions are included in the NIR in Chapter 3.5.

Data Processing level 1	7.Transparency	DP.1.7.2	Clear reference to dataset at Data Storage level 1
-------------------------	----------------	----------	--

Notes on data sources are included in the calculation files for all input data.

Data Processing level 1	7.Transparency	DP.1.7.3	A manual log to collect information about recalculations.
-------------------------	----------------	----------	---

A log holding information on recalculations are included in the national inventory system. Further, a log is prepared annually holding information on status of the inventory work and recalculations for each source in the fugitive sector.

Data storage level 2

Data Storage level 2	5.Correctness	DS.2.5.1	Check if a correct data import to level 2 has been made
----------------------	---------------	----------	---

To ensure a correct connection between data on level 2 to data on level 1, different controls are in place, e.g. control of sums and random tests.

Data storage level 4

Data Storage level 4	4.Consistency	DS.4.4.3	The IEFs from the CRF are checked both regarding level and trend. The level is compared to relevant emission factors to ensure correctness. Large dips/jumps in the time series are explained.
----------------------	---------------	----------	--

Time series for IEFs are checked to identify large fluctuations, which are afterwards investigated and explained. The level of the IEFs are compared to other relevant EFs, e.g. in standard EFs in guidebooks and guidelines.

The IEFs for transmission and distribution of natural gas are low compared to other countries as the Danish distribution network is relatively new and made of plastics, leading to negligible fugitive losses under normal circumstances. Only fugitive losses are due to excavations and maintenance and construction work.

Other QC procedures

A list of QA/QC tasks are performed directly in relation to the fugitive emission part of the Danish emission inventories. The following procedures are carried out to ensure the data quality:

- The emission from the large point sources (refineries, gas treatment and gas storage plants) is compared with the emission reported the previous year.
- Annual environmental reports are kept for subsequent control of plant-specific emission data.
- Checks of data transfer are incorporated in the fugitive emission models, e.g. sum checks.
- Verification of activity data from external data when data are available through more data sources (offshore fuel and flaring rates).
- Data sources are incorporated in the fugitive emission models
- A manual log table in the emission databases is applied to collect information about recalculations.
- Comparison with the inventory of the previous year. Any major changes are verified.
- Total emission, when aggregated to reporting tables, is compared with totals based on SNAP source categories (control of data transfer).
- Checking of time series in the NFR and SNAP source categories. Significant dips and jumps are controlled and explained.

The QC work will continue in future years.

National external review

In 2009 a sector report for fugitive emissions from fuels was published (Plejdrup et al. 2009). The report was reviewed by Anette Holst from the Statoil A/S refinery.

3.5.8 Recalculations

The following recalculations regarding fugitive emissions from fuels have been applied for the time series:

Natural gas transmission and distribution

Activity data and IEF for the time series 1990-2011 has been updated for transmission and distribution according to annual environmental reports and the latest national energy statistics, respectively. Further, the CH₄ EFs are updated for one town gas distribution company following an update of the estimated fugitive losses per distribution. The recalculation has changed the CO₂ emission by 0.01 ktonnes and CH₄ emission by (-0.09) - 0.07 ktonnes, corresponding to < 0.003% and (-2) % - 2% of the total fugitive CO₂ and CH₄ emission.

Venting

EFs for CH₄ have been added for the years 1990-1993 for one gas storage plant. In these years the plant is treated as an area source in the national system, while it is treated separately as a point source in the following years. EFs are based on data from annual reports for 1995-1999, as no data are available for the years 1990-1994. Further, a minor error has been applied for venting in 2011, according to the annual report from one of the natural gas storage facilities. The recalculation has changed the CH₄ emission by 0.06 ktonnes, corresponding to 2% of the total fugitive CH₄ emission.

Flaring in gas storage and treatment plants

EFs for CO₂ and CH₄ have been added for the years 1990-1993 for the gas treatment plant. In these years the plant is treated as an area source in the national system, while it is treated separately as a LPS in the following years. EFs are based on data from annual reports for 1995-1999, as no data are available for the years 1990-1994. The recalculation has changed the CO₂ emission by 2.2 ktonnes and the CH₄ emission by 0.01 ktonnes, corresponding to 0.4 % and 0.4 % of the total fugitive CO₂ and CH₄ emission.

Flaring in refineries

The CO₂ EF for one refinery has been updated for the years 1994-2006 and now reflects the average EF from the first five EU-ETS reports (2007-2011). The NO_x EF for two refineries has been changed to the standard EF in the EMEP/EEA Guidebook, as references to the previously applied EFs are outdated or not existing.

The CO₂ EF for a refinery that was shut down in 1996 has been changed for the years 1990-1996 to match the existing refineries, as no better data is available. CO₂ EF for 2010-2011 has been updated and now corresponds to the EU-ETS reporting. The recalculation has changed the CO₂ emission by (-0.05) - 0.49 ktonnes and CH₄ emission by (-0.06) ktonnes, corresponding to (-0.01) - 0.1 % and (-2) % of the total fugitive CO₂ and CH₄ emission.

3.5.9 Source specific planned improvements

The following future improvements are suggested.

- **Emissions from storage of fuels in tank facilities:** The current edition of the Danish emission inventory holds emissions from storage and refining of crude oil and from service stations. To make the inventory complete emissions from storage of fuels outside the refineries in tank facilities will be included in the future if data are available. Work is on-going to locate large tank facilities in Denmark and collect the available data. In cases where no emission estimates or measurements are available a set of emission factors have to be set up.

3.5.10 References

A/S Dansk Shell, 2013: Annual environmental report 2012.

Danish Energy Agency, 2013a: Oil and Gas Production in Denmark 2012.

Danish Energy Agency, 2013b: The Danish energy statistics (Energistatistik) (in Danish). Available at:
http://www.ens.dk/en-US/Info/FactsAndFigures/Energy_statistics_and_indicators/Annual%20Statistics/Sider/Forside.aspx (31-01-2014).

Danish Environmental Protection Agency, 2008: Emissionsfaktorer for NO_x-emissioner fra flaring fra platforme i Nordsøen (in Danish). Not published.

DONG Energy, 2013a: Annual environmental reports from DONG Energy for Stenlille gas storage plants (in Danish). Available at:
http://www.dongenergy.com/da/ansvarlighed/downloads/pages/groene_regnskaber.aspx (31-01-2014).

DONG Energy, 2013b: Annual environmental reports from DONG Energy for Nybro gas treatment plant (in Danish). Available at:
http://www.dongenergy.com/da/ansvarlighed/downloads/pages/groene_regnskaber.aspx (31-01-2014).

DONG Oil Pipe A/S, 2013: Dong Energy Raw Oil Terminal, Fredericia – Self-regulation Report 2011 (in Danish).

EMEP/EEA, 2009: Air emission inventory guidebook, prepared by the UNECE/EMEP Task Force on Emission Inventories and Projections, 2009 update. Available at:
<http://www.eea.europa.eu/publications/emep-eea-emission-inventory-guidebook-2009> (31-01-2014).

EMEP/EEA, 2013: Air emission inventory guidebook, prepared by the UNECE/EMEP Task Force on Emission Inventories and Projections, 2009 update. Available at:
<http://www.eea.europa.eu/publications/emep-eea-guidebook-2013> (31-01-2014).

Energinet.dk, 2013a: Annual report 2012 for Lille Torup gas storage plant (in Danish: "GRØNT REGNSKAB 2012 for Energinet.dk Gaslager A/S Ll. Torup Naturgaslager").

Energinet.dk, 2013b: Environmental Impact from transmission of gas. Available at:
<http://www.energinet.dk/DA/KLIMA-OG-MILJOE/Energinetdks-miljoepaavirkninger/Miljoepaavirkninger-ved-transport-af-gas/Sider/Miljoepaavirkninger-ved-transport-af-gas.aspx> (31-01-2014).

Energinet.dk, 2012c: Gas quality 2011. Available at:
<http://energinet.dk/DA/GAS/Gasdata-og-kvalitet/Gaskvalitet/Sider/Vis-gaskvalitet.aspx?Visning=aarsgennemsnit> (Gas quality for earlier years available in the background reports of the annual environmental reports - available at:
<http://energinet.dk/DA/KLIMA-OG-MILJOE/Miljoerapportering/Sider/default.aspx> (in Danish)) (31-01-2014).

EU Commission, 2007: COMMISSION DECISION of 18 July 2007 establishing guidelines for the monitoring and reporting of greenhouse gas emissions pursuant to Directive 2003/87/EC of the European Parliament and of the Council. Available at:
<http://eur-lex.europa.eu/LexUriServ/LexUriServ.do?uri=OJ:L:2007:229:0001:0085:EN:PDF> (31-01-2014).

Fenhann, J. & Kilde, N.A. 1994: Inventory of Emissions to the Air from Danish Sources 1972-1992. Risø National Laboratory, Roskilde, Denmark.

Hjerrild, F., 1997: Personal communication. Letter 31-07-1997. A/S Dansk Shell.

Holst, A. 2009: Personal communication, e-mail 18-11-2009, Statoil A/S.

IPCC, 2000: Good Practice Guidance and Uncertainty Management in National Greenhouse Gas Inventories. Available at: <http://www.ipcc-nggip.iges.or.jp/public/gp/english/> (31-01-2014).

Karll, B. 2003: Personal communication, e-mail 17-11-2003, Danish Gas Technology Centre.

Karll, B. 2005: Personal communication, e-mail 09-11-2005, Danish Gas Technology Centre.

Nielsen, O.-K., Plejdrup, M.S., Winther, M., Gyldenkerne, S., Thomsen, M., Fauser, P., Nielsen, M. Mikkelsen, M.H., Albrektsen, R., Hjelgaard, K., Hoffmann, L. & Bruun, H.G. 2013. Quality manual for the Danish greenhouse gas inventory. Version 2. Aarhus University, DCE – Danish Centre for Environment and Energy, 44 pp. Scientific Report from DCE – Danish Centre for Environment and Energy No. 47 Available at: <http://www.dmu.dk/Pub/SR47.pdf> (31-01-2014).

Oertenblad, M. 2006: personal communication, e-mail 2006, Danish Gas Technology Centre.

Oertenblad, M. 2007: personal communication, e-mail 2007, Danish Gas Technology Centre.

The Norwegian Oil industry Association, OLF environmental program. Phase II. Summary report. 1993.

Statistisk Sentralbyrå, Statistics Norway, 2008: The Norwegian Emission Inventory 2008. Documentation of methodologies for estimating emissions of greenhouse gases and long-range transboundary air pollutants. 252 pp. Reports 2008/48. Available at: http://www.ssb.no/english/subjects/01/sa_nrm/nrm2008/sa109.pdf (31-01-2014).

Statoil A/S, 2010: Ubestemtheden på massestrømmen ved blændemåling og ultralyd. Report in Danish.

Statoil A/S, 2013: Annual environmental report 2012 (in Danish). Available at: <http://www.statoil.com/en/About/Worldwide/Denmark/Downloads/De%20gr%C3%B8nne%20regnskab%202012.pdf> (31-01-2014).

4 Industrial Processes (CRF sector 2)

4.1 Overview of the sector

4.1.1 Emission overview

The aim of this chapter is to present industrial emissions of greenhouse gases, not related to generation of energy. The data presented in Chapter 4 relates to Denmark only, whereas information for Greenland is included in Chapter 16 and for the Faroe Islands in Annex 8.

An overview of the sources identified is presented in Table 4.1 with an indication of the contribution to the industrial part of the emission of greenhouse gases in 2012. The emissions are extracted from the CRF tables.

Table 4.1 Overview of industrial greenhouse gas sources (2012).

Process	IPCC Code	Substance	Emission ktonne CO ₂ eq.	Percent
Cement	2A1		871	48.6
Refrigeration	2F1	HFCs+PFCs	579	32.4
Other (laboratories, double glaze windows)	2F9	SF ₆	105	5.84
Foam blowing	2F2	HFCs	66.2	3.70
Lime	2A2		40.2	2.24
Other (lubricants)	2G		31.7	1.77
Limestone and dolomite use	2A3		25.6	1.43
Other (yellow bricks)	2A7		17.5	0.98
Aerosols / Metered dose inhalers	2F4	HFCs	15.8	0.88
Electrical equipment	2F8	SF ₆	13.3	0.74
Other (container glass, glass wool)	2A7		9.69	0.54
Other (expanded clay products)	2A7		5.89	0.33
Other (fibre optics)	2F9	HFCs+PFCs	4.31	0.24
Food and Drink	2D2		2.24	0.13
Road paving	2A6		1.77	0.10
Catalysts / fertilisers	2B5		1.41	0.08
Asphalt roofing	2A5		0.019	0.001
Metal production	2C1		NO	NO
Nitric acid	2B2	N ₂ O	NO	NO
Total			1 791	100

The subsectors *Mineral products* (2A) constitutes 54 %, *Chemical industry* (2B) constitutes below 1 %, *Metal production* (2C) constitutes 0 %, *Consumption of halocarbons and SF₆* (2F) constitutes 44 %, *Other, Food and Drink* (2D) constitutes below 1 %, and *Other, Lubricants* (2G) constitutes 1.8 % of the industrial emission of greenhouse gases. The total emission of greenhouse gases (excl. LULUCF) in Denmark is estimated to 51.6 Mt CO₂ equivalents, of which industrial processes contribute with 1.79 Mt CO₂ equivalents (3.5 %). The emission of greenhouse gases from industrial processes from 1990-2012 are presented in Figure 4.1.




Figure 4.1 Emission of greenhouse gases from industrial processes (CRF Sector 2) from 1990-2012.

The key categories in the industrial processes sector - cement and refrigeration - constitute 1.69 % and 1.12 % of the total emission of greenhouse gases. The trends in greenhouse gases from the industrial sector/subsectors are presented in Table 4.2 and they will be discussed subsector by subsector below. The emissions are extracted from the CRF tables.

Table 4.2 Emission of greenhouse gases from industrial processes in different subsectors from 1990-2012.

Year	1990	1995	2000	2005	2007	2008	2009	2010	2011	2012
CO₂ (kt CO₂)										
A. Mineral Products	1 069	1 405	1 616	1 544	1 606	1 320	881	796	972	972
B. Chemical Industry	0.80	0.80	0.79	1.06	1.35	1.20	1.42	2.12	1.39	1.41
C. Metal Production	28.4	38.6	40.7	15.6	NA	NA	NA	NA	NA	NA
D. Food and Drink	4.45	3.91	3.90	4.46	1.72	2.67	1.92	1.56	2.01	2.24
G. Other	49.7	48.8	39.7	37.6	37.9	34.0	31.2	33.2	33.2	31.7
Total	1 152	1 497	1 701	1 602	1 647	1 358	915	833	1 008	1 007
CH₄										
	-	-	-	-	-	-	-	-	-	-
N₂O (kt N₂O)										
B. Chemical Industry	3.36	2.92	3.24	NO	NO	NO	NO	NO	NO	NO
HFCs (kt CO₂ eqv.)										
F. Consumption of Halocarbons and SF ₆	NE	218	607	802	850	853	799	804	759	657
PFCs (kt CO₂ eqv.)										
F. Consumption of Halocarbons and SF ₆	NE	0.50	17.9	13.9	15,4	12,8	14,2	13,3	11,1	8,54
SF₆ (kt CO₂ eqv.)										
F. Consumption of Halocarbons and SF ₆	44.5	107	58.8	21.8	30,3	31,6	36,7	37,9	73,2	118

4.1.2 Methodology overview

Table 4.3 gives a brief overview over methodologies applied for industrial processes. Further description of the applied methodologies can be found in the following chapters.

Table 4.3 Methodology overview.

			Tier	EF	Key category
Industrial Proc.	2A1 Cement production	CO ₂	T3	PS	Yes
Industrial Proc.	2A2 Lime production	CO ₂	T2	T1	No
Industrial Proc.	2A3 Limestone and dolomite use	CO ₂	T2	T1,PS	No
Industrial Proc.	2A5 Asphalt roofing	CO ₂	T1	CS	No
Industrial Proc.	2A6 Road paving with asphalt	CO ₂	T1	CS	No
Industrial Proc.	2A7 Glass and Glass wool	CO ₂	T2,T3	T1	No
Industrial Proc.	2B5 Catalysts/Fertilizers, Pesticides and Sulphuric acid	CO ₂	T2	PS	No
Industrial Proc.	2C1 Iron and steel production	CO ₂	T1	T1	No
Industrial Proc.	2D2 Food and Drink	CO ₂	T2	T2,PS	No
Industrial Proc.	2G Lubricants	CO ₂	T1	T1	No
Industrial Proc.	2B2 Nitric acid production	N ₂ O	T1	PS	Yes
Industrial Proc.	2F Consumption of HFC	HFC	T1,T2	CS	Yes
Industrial Proc.	2F Consumption of PFC	PFC	T1,T2	CS	No
Industrial Proc.	2F Consumption of SF ₆	SF ₆	T1,T2	CS	No

4.1.3 Key categories

Key Category Analysis (KCA) for the years 1990 and 2012 as well as for the trend has been carried out. Table 4.4 present the result. A detailed KCA is presented in Chapter 1.5 and Annex 1.

Cement production is identified as key category (level and trend) according to Tier 1 for 1990 and 2012. Nitric acid production is identified as key category in 1990 and the trend is also identified as key according to Tier 1 and 2. Consumption of HFC is identified as key category in 2012 and the trend is also identified as key according to Tier 1 and 2.

Table 4.4 Key Category Analysis for Industrial processes.

			Tier 1			Tier 2		
			1990	2012	1990-2012	1990	2012	1990-2012
Industrial Proc.	2A1 Cement production	CO ₂	Level	Level	Trend			
Industrial Proc.	2A2 Lime production	CO ₂						
Industrial Proc.	2A3 Limestone and dolomite use	CO ₂						
Industrial Proc.	2A5 Asphalt roofing	CO ₂						
Industrial Proc.	2A6 Road paving with asphalt	CO ₂						
Industrial Proc.	2A7a Glass and Glass wool	CO ₂						
Industrial Proc.	2A7b Yellow bricks	CO ₂						
Industrial Proc.	2A7c Expanded clay	CO ₂						
Industrial Proc.	2B5 Catalysts/Fertilizers, Pesticides and Sulphuric acid	CO ₂						
Industrial Proc.	2C1 Iron and steel production	CO ₂						
Industrial Proc.	2D2 Food and Drink	CO ₂						
Industrial Proc.	2G Lubricants	CO ₂						
Industrial Proc.	2B2 Nitric acid production	N ₂ O	Level		Trend	Level		Trend
Industrial Proc.	2F Consumption of HFC	HFC		Level	Trend		Level	Trend
Industrial Proc.	2F Consumption of PFC	PFC						
Industrial Proc.	2F Consumption of SF ₆	SF ₆						

4.2 Mineral products (2A)

4.2.1 Source category description

The subsector *Mineral products (2A)* cover the following processes:

- Cement production
- Lime production
- Limestone and dolomite use (for fluegas cleaning at power plants and waste incineration plants and for production of stonewool).
- Asphalt roofing
- Road paving with asphalt
- Production of bricks, tiles and expanded clay products
- Production of container glass/glass wool

Production of cement is identified as a key category; see *Annex 1: Key Category Analyses*.

The time series for the emission of CO₂ from *Mineral products (2A)* are presented in Table 4.5. The emissions are extracted from the CRF tables and the values are rounded.

Table 4.5 Time series for emission of CO₂ (kt) from Mineral products (2A).

	1990	1995	2000	2005	2007	2008	2009	2010	2011	2012
1. Production of Cement	882	1 204	1 385	1 363	1 407	1 155	764	672	862	871
2. Production of Lime	116	87.7	76.7	63.5	66.9	65.6	43.2	45.6	29.8	40.2
3. Limestone and dolomite use	13.7	53.7	89.7	56.2	49.6	38.7	37.9	45.6	41.7	25.6
5. Asphalt roofing	0.019	0.020	0.032	0.024	0.025	0.025	0.016	0.016	0.021	0.019
6. Road paving	1.76	1.77	1.72	1.84	2.00	1.92	1.64	1.73	1.88	1.77
7. Other										
Glass and Glass wool	17.4	14.1	15.9	12.6	15.0	15.1	10.8	9.33	9.46	9.69
Yellow Bricks	23.0	28.8	32.6	32.2	38.0	28.4	16.5	15.8	20.4	17.5
Expanded Clay	14.9	15.3	14.2	14.0	26.9	16.1	6.48	6.00	6.63	5.89
Total	1 069	1 405	1 616	1 544	1 606	1 320	881	796	972	972

The increase in CO₂ emission is most significant for the production of cement until 2007; however, in the latest years the emission has been decreasing. The overall development in the CO₂ emission from 1990 to 2012 shows a slightly decreasing trend from 882 to 871 kt CO₂, i.e. by 1.3 %. The maximum emission occurred in 2004 and constituted 1 459 kt CO₂; see Figure 4.2.

The increase from 1990 to 1997 can be explained by the increase in the annual cement production. The emission factor has only changed slightly as the distribution between types of cement especially grey/white cement has been almost constant from 1990-1997. The decrease during the latest years may be explained by the decrease in the construction activity.




Figure 4.2 Emission of CO₂ from cement production.

4.2.2 Methodological issues

2A1 Cement production

Public statistics for cement production, import and export are presented in Table 4.6. However, the dataset is incomplete probably due to confidentiality in some of the mass flows. The statistics do not add new information to the inventory as information on total amount of produced cement clinker (i.e. clinker for cement production and clinker for sale) is not available. Activity data from Aalborg Portland and implied emission factors for cement production are presented in Table 4.7. Table 4.6 and Table 4.7 show a difference between produced amount of cement (grey and white) according to Statistics Denmark and the amount produced according to Aalborg Portland. This difference may be explained by change in clinker stocks. The most comprehensive activity data is assumed to be the information on yearly produced amount of cement clinker obtained from the Danish producer.

Table 4.6 Production, import, export and supply of cement (Statistics Denmark, 2013).

		1990	1995	2000	2005	2007	2008	2009	2010	2011	2012
Cement clinker ¹	Prod.	NI	NI	102 719	42 691	21 209	16 092	122	3 819	25	24 256
	Import	404	8	NI	31 331	39 776	42 480	32 444	21 391	29 106	24 442
	Export	17 038	186 979	NI	NI	NI	NI	NI	NI	NI	NI
	Supply	-16 634	-186 971	102 719	74 022	60 985	58 572	32 566	25 210	29 131	48 698
Portland cement, white	Prod.	411 653	531 215	551 072	714 626	722 310	607 143	462 377	481 507	514 389	495 886
	Import	NI	24	8 299	15 438	18 670	33 082	29 423	24 152	28 414	24 645
	Export	91 132	NI	NI	NI	NI	NI	NI	NI	NI	NI
	Supply	320 521	531 239	559 371	730 064	740 980	640 225	491 800	505 659	542 803	520 531
Portland cement, grey	Prod.	1 244 256	2 052 705	1 984 976	2 165 928	2 149 092	1 931 679	1 116 237	1 085 117	1 338 415	1 321 447
	Import	115 452	NI	NI	NI	NI	NI	NI	159 923	213 930	182 473
	Export	19 387	332 390	NI	NI	NI	NI	NI	NI	NI	NI
	Supply	1 340 321	1 720 315	1 984 976	2 165 928	2 149 092	1 931 679	1 116 237	1 245 040	1 552 345	1 503 920

1. Cement clinker for sale.

NI No information.

Table 4.7 Activity data, emission factors, and CO₂ emission for cement production.

	1990	1995	2000	2005	2007	2008	2009	2010	2011	2012
Activity										
Tonnes TCE ⁶	1 619 976	2 273 775	2 612 721	2 706 371	2 946 294	2 551 346	1 663 126	1 454 043	1 766 561	1 818 293
Tonnes clinker	NI	NI	2 452 394	2 520 788	2 706 048	2 269 687	1 493 230	1 313 654	1 582 023	1 628 506
Tonnes clinker + white cement ¹	1 406 212	2 353 123	-	-	-	-	-	-	-	-
Emission factors										
EF tonnes CO ₂ per tonnes TCE ²	0.545	0.529	-	-	-	-	-	-	-	-
EF tonnes CO ₂ per tonnes TCE ³	-	-	0.530	-	-	-	-	-	-	-
EF tonnes CO ₂ per tonnes TCE ⁴	-	-	-	0.504	0.478	-	-	-	-	-
EF tonnes CO ₂ per tonnes clinker ^{4,5}	0.628	0.512	0.565	0.541	0.520	0.509	0.512	0.512	0.545	0.535
Emission										
Tonnes CO ₂	882 402	1 203 777	1 384 742	1 363 000	1 407 086	1 154 749	764 407	672 224	861 805	871 083

1. 1990-1997: Amount of clinker produced has not been measured as for 1998-2008. Therefore, the amount of GLK-, FHK-, SKL-/RKL-clinker and white cement is used as estimate of total clinker production.

2. 1990-1997: EF based on information provided by Aalborg Portland.

3. 1998-2004: EF based on information provided by Aalborg Portland (Aalborg Portland, 2008).

4. 2005-2012: EF based on emissions reported to EU-ETS (Aalborg Portland, 2013a).

5. 1998-2012: EF based on clinker production statistics provided by Aalborg Portland (Aalborg Portland, 2013c).

6. Aalborg Portland (2013b).

NI No information.

1990-1997

The emission factor has been estimated from the loss on ignition determined for the different kinds of clinkers produced, combined with the volumes of grey and white cements produced.

The emission of CO₂ depends on the ratio: white/grey cement and the ratio between three types of clinker used for grey cement: GLK-clinker/FHK-clinker/SKL-RKL-clinker. The ratio white/grey cement is known from 1990-1997 with maximum in 1990 and thereafter decreasing. The ratio: GLK-clinker/FHK-clinker/SKL-RKL-clinker is known from 1990-1997. The production of SKL/RKL-clinker peaks in 1991 and decreases hereafter. FKH-clinker is introduced in 1992 and increase to 35 % in 1997.

$$M_{CO_2} = M_{grey} * \frac{M_{GLK} * EF_{GLK} + M_{FKH} * EF_{FKH} + M_{SKL/RKL} * EF_{SKL/RKL}}{M_{GLK} + M_{FKH} + M_{SKL/RKL}} + M_{white} * EF_{white}$$

Table 4.8 Types of cement and emission factors

M _{grey}	Grey cement	Tonne
M _{white}	White cement	Tonne
M _{GLK}	GLK clinker	Tonne
M _{FKH}	FHK clinker	Tonne
M _{SKL/RKL}	SKL/RKL clinker	Tonne
EF _{white}	CO ₂ emission factor	0.669 tonne CO ₂ /tonne white cement
EF _{GLK}	CO ₂ emission factor	0.477 tonne CO ₂ /tonne GLK clinker
EF _{FKH}	CO ₂ emission factor	0.459 tonne CO ₂ /tonne FKH clinker
EF _{SKL/RKL}	CO ₂ emission factor	0.610 tonne CO ₂ /tonne SKL/RKL clinker

The company has at the same time stated that data until 1997 cannot be improved as they are not available anymore.

1998-2004

From 1998-2004 carbonate content of the raw materials has been determined by loss on ignition methodology. Determination of loss on ignition takes into account all the potential raw materials leading to release of CO₂ and omits the Ca-sources leading to generation of CaO in cement clinker without CO₂ release. The applied methodology is in accordance with EU guidelines on calculation of CO₂ emissions (Aalborg Portland, 2008).

2005-2012

However, from the year 2005 the CO₂ emission determined by Aalborg Portland for EU-ETS is used in the inventory (Aalborg Portland, 2013a). The reporting to EU-ETS also provides detailed information of alternative fuels used in the production of clinker; see Table 4.9.

Table 4.9 Alternative fuels used in production of cement clinker (Aalborg Portland 2013a).

Fuel type	Biomass fraction, %
Cemmiljø fuel	30-56
Paper residues	79
Dry wastewater sludge	100
Meat and bone meal	100
Tyre residues	15
Textile residues from tyres	100
Wood waste	100
Garden waste	100
Glycerine	100

The Aalborg Portland report to EU-ETS applies the EF for the most important raw materials as shown in Table 4.10 (Aalborg Portland, 2013a).

Table 4.10 Emission factors for different raw materials in cement production, 2012.

Raw material	Tonne CO ₂ per tonne raw material
Limestone	0.44
Magnesium carbonate	0.522
Sand	0.0088-0.0367
Fly ash	0.140
CKD	0.396-0.525

The EFs for limestone and magnesium carbonate are in accordance with the stoichiometric factors and the EFs for the remaining raw materials and CKD are determined by individual analysis.

2A2 Lime production.

The CO₂ emission from the production of burnt lime (quicklime) as well as hydrated lime (slaked lime) has been estimated from the annual production figures, registered by Statistics Denmark – see Table 4.11 and emission factors.

Table 4.11 Statistics for production of lime and slaked lime (tonnes) (Faxekalk, 2013b, Statistics Denmark, 2013).

Year	1990	1995	2000	2005	2007	2008	2009	2010	2011	2012
Lime	127 978	100 789	92 002	71 239	75 504	74 981	46 202	50 397	35 824	48 999
Slaked lime	27 686	15 804	8 159	13 839	14 028	12 326	12 842	11 173	13 264	13 388
Produced (burnt) in Denmark	155 664	116 593	100 161	85 078	89 532	87 307	59 044	61 570	49 088	62 387
Imported within company									23 606	20 137
Total amount sold from										
Danish producers	155 664	116 593	100 161	85 078	89 532	87 307	59 044	61 570	72 694	82 524


The emission factors applied are 0.785 kg CO₂ per kg CaO as recommended by IPCC (IPCC, 1997, vol. 3, p. 2.8) and 0.541 kg CO₂ per kg hydrated lime (calculated from company information on composition of hydrated lime: product purity: 91 % Ca(OH)₂ in product (Faxekalk, 2003)). One Danish company – Faxekalk – is covered by the EU-ETS, however, the company do only account for approximately 75 % of the Danish production of lime and hydrated lime (average from 1998-2012). A number of small companies accounts for the remaining of the Danish production.

The company reporting to EU-ETS apply the following EF: 0.7874867 tonne CO₂ per tonne lime produced (Faxekalk, 2013a). This EF is in accordance with the stoichiometric factor.

2A3 Limestone and dolomite use

In the production of stonewool a number of raw materials contributing to CO₂ emission are used: bottom ash from coal-fired CHP, stonewool binder, stonewool waste, limestone, and dolomite. Information on emission of CO₂ has been obtained from confidential company reports to EU-ETS for 2006-2012. For the previous years the emission has been extrapolated.

The CO₂ emission from consumption of limestone for flue gas cleaning has been estimated from statistics on generation of gypsum (wet flue gas cleaning processes) and the stoichiometric relations between gypsum and release of CO₂:



and the emission factor is: 0.2325 tonnes CO₂ per tonne gypsum.

Statistics on the generation of gypsum from power plants are compiled by Energinet.dk (2008). However, for 2006 - 2012 information on consumption of CaCO₃ at the relevant power plants has been compiled (from environmental reports) and used in the calculation of CO₂ emission from flue gas cleaning.

Information on the generation of gypsum at waste incineration plants does not explicitly appear in the Danish waste statistics (Miljøstyrelsen, 2012). However, the total amount of waste products generated can be found in the statistics. The amount of gypsum is calculated by using information on flue gas cleaning systems at Danish waste incineration plants (Illerup et al., 1999; Nielsen & Illerup, 2002) and waste generation from the different flue gas cleaning systems (Hjelmar & Hansen, 2002). However, for 2011 and 2012 information of CaCO₃ at the relevant plants has been compiled from environmental reports and used in the calculation of CO₂ emission from flue gas cleaning.

2A5 Asphalt roofing and 2A6 Road paving with asphalt

The indirect emission of CO₂ from asphalt roofing and road paving has been estimated from production statistics compiled by Statistics Denmark and default emission factors presented by IPCC (1997) and EMEP/CORINAIR (2004). The default emission factors, together with the calculated emission factor for CO₂, are presented in Table 4.12.

Table 4.12 Default emission factors for application of asphalt products.

		Road paving with asphalt	Use of cutback asphalt	Asphalt roofing
CH ₄	g pr tonnes	5	0	0
CO	g pr tonnes	75	0	10
NM VOC	g pr tonnes	15	64 935	80
Carbon content fraction of NM VOC	%	0.667	0.667	0.8
Indirect CO ₂	Kg pr tonnes	0.168	159	0.250

2A7 Other

The category *Other* cover the following processes:

- Production of yellow bricks
- Expanded clay products
- Production of glass
- Production of glass woll

The CO₂ emission from the production of bricks and tiles has been estimated from information on annual production registered by Statistics Denmark, corrected for amount of yellow bricks and tiles. This amount is unknown and, therefore, is assumed to be 50 %; see Table 4.13.

Table 4.13 Statistics for production of yellow bricks and expanded clay products (tonnes) (Statistics Denmark, 2013).

Year	1990	1995	2000	2005	2007	2008	2009	2010	2011	2012
Bricks (1000 pieces.)	291 348	362 711	414 791	407 940	348 928	322 137	226 363	212 051	222 144	185 398
Yellow bricks ¹	291 348	362 711	414 791	407 940	348 928	322 137	226 363	212 051	222 144	185 398
Expanded clay products	331 760	340 881	316 174	310 901	504 925	303 948	140 915	157 378	172 263	153 305

1. Assumptions: Brick weight: 2 kg/brick, 50 % yellow bricks.

The content of CaCO₃ and a number of other factors determine the colour of bricks and tiles and, in the present estimate, the average content of CaCO₃ in clay has been assumed to be 18 % (the CaCO₃ content in clay for bricks varies from e.g. 0.5 % in clay for red bricks and 19.8 % in clay for yellow bricks (Tegl Info, 2004). The emission factor for lime (0.44 kg CO₂ per kg CaCO₃) has been used to calculate the emission factor for yellow bricks: 0.079 tonne CO₂ per tonne yellow bricks. For verification of this approach, see Figure 4.3.

For 2006-2012 emission factors have been derived from CO₂ emissions reported by the brickworks to EU-ETS (confidential reports from approximately 20 brickworks) and production statistics (Statistics Denmark, 2013). The emission factors for 2006-2012 are calculated to 0.0728-0.1089 tonne CO₂ per tonne yellow bricks (average 0.086).

The company reporting to EU-ETS apply the following EF: 0.44 tonne CO₂ per tonne limestone consumed (different brickworks). This EF is in accordance with the stoichiometric factor.

The CO₂ emission from the production of expanded clay products has been estimated from production statistics compiled by Statistics Denmark and an emission factor of 0.045 tonne CO₂ per tonne product. During the preliminary work to establishment of EU-ETS in Denmark the relevant companies reported energy and process related CO₂ emissions for the years 1998-2002 (DEA, 2004). Based on the process CO₂ emissions and statistical information (Statistics Denmark, 2013) the actual emission factors were determined to respectively 0.0345, 0.0349, 0.0415, 0.0445, and 0.0358 tonne CO₂ per tonne product. As a conservative estimate 0.045 tonne CO₂ per tonne product was chosen as national emission factor.

For 2006-2012 emission factors have been derived from CO₂ emissions reported to EU-ETS (Damolin, 2013; Maxit, 2013) and production statistics (Statistics Denmark, 2012). The emission factors for 2006-2012 are calculated to 0.0381-0.0532 tonne CO₂ per tonne product (average: 0.045).

The companies producing expanded clay products reporting to EU-ETS apply the EF shown in Table 4.14 (different producers):

Table 4.14 Emission factors for production of expanded clay products (multiple producers), 2012.

Raw material	tonne CO ₂ per tonne raw material
Bentonite	0.003364
Clay (moler)	0.03002
Clay (moler)	0.02163
Clay, dried (moler)	0.00691
Limestone	0.44
C-source	3.67
C-source	3.14631

The EFs for the clay fractions are determined at the different localities whereas the EFs for limestone and one of the C-sources are in accordance with the stoichiometric factors.

The CO₂ emission from the production of container glass/glass wool has been estimated from production statistics published in environmental reports from the producers (Rexam Glass Holmegaard, 2007; Ardagh Glass Holmegaard, 2013b; Saint-Gobain Isover, 2013b) and emission factors based on release of CO₂ from specific raw materials (stoichiometric determination). For 2006-2012 CO₂ emissions have been derived from company reports to EU-ETS (Ardagh Glass Holmegaard, 2013a; Saint-Gobain Isover, 2013a) combined with information in environmental reports.

The companies producing glass and glass wool reporting to EU-ETS apply the EF shown in Table 4.15 (Ardagh Glass Holmegaard, 2013a, Saint-Gobain Isover, 2013a)

Table 4.15 Emission factors used in glass and glass wool production, 2012.

Raw material	Tonne CO ₂ per tonne raw material
Limestone	0.44
Magnesium carbonate	0.522
Dolomite	0.478
Soda ash	0.415
C-source	3.666

The EFs are in accordance with the stoichiometric factors.

EU-ETS (EU Emission Trading Scheme)

Guidelines for calculating company specific CO₂ emissions are developed by the EU (EU, 2007). The guidelines present standard methods for minor companies and methods for developing individual plans for major companies. The standard methods include default emission factors similar to the default emission factors presented by IPCC (e.g. for limestone), whereas, the major companies has to use individual methods to determine the actual composition of raw materials (e.g. purity of limestone or Ca per Mg ratio in dolomite) or the actual CO₂ emission from the specific process.

4.2.3 Uncertainties and time series consistency

The time series are presented in Table 4.5. The methodology applied for the years 1990-2012 is considered to be consistent. The emission factor has only changed slightly as the distribution between types of cement, especially grey/white cement, has been almost constant from 1990-1997. Furthermore, the activity data originates from the same company for all years.

For the production of lime and bricks, as well as container glass and glass wool, the same methodology has also been applied for all years. The emission factors are based either on stoichiometric relations or on a standard assumption of CaCO₃-content of clay used for bricks. The source for the activity data is, for all years, Statistics Denmark.

The source specific uncertainties for mineral products are presented in Chapter 4.9. The overall uncertainty estimate is presented in Chapter 1.7.

4.2.4 Verification

The estimation of CO₂ release from the production of bricks based on an assumption of 50 % yellow bricks has been verified by comparing the estimate with actual information on emission of CO₂ from calcination of lime compiled by the Danish Energy Agency (DEA) (DEA, 2004). The information from the companies (tile-/brickworks; based on measurements of CaCO₃ content of raw material) has been compiled by DEA in order to allocate a CO₂ quota to Danish companies with the purpose of future reductions. The result of the comparison is presented in Figure 4.3. From 2006 the information obtained from the EU-ETS reports was implemented directly.




Figure 4.3 Estimated and “measured” CO₂ emission from tile-/brickworks; “measured” means information provided to the Danish Energy Agency by the individual companies (DEA, 2004).

Figure 4.3 shows a reasonable correlation between the estimated and measured CO₂ emission.

The ERT has recommended Denmark to develop a national EF based on the IEF for the years 2006-2010 i.e. emissions based on company reports to EU-ETS. Figure 4.4 presents three scenarios for yellow bricks and expanded clay products:

- Applied methodology from 1990-2005. The EF is based on the assumption that clay for yellow bricks contains 18% CaCO₃.
- CO₂ emission based on company reporting to EU ETS for the years 2006-2010.
- Methodology recommended by UNFCCC ERT. The national EF is based on an average of IEF from the years 2006-2011.

Expanded clay products are also a mix of different products with different CaCO₃ addition or content. The actual mix is unknown.




Figure 4.4 CO₂ emission from production of yellow bricks and expanded clay products. The scenarios present the applied methodology (Estimated), available EU-ETS data, and emissions based on a national EF (Average EF).

The recommended methodologies have weaknesses as the (average) EFs are based on actual reported emissions from brickworks and an assumed amount of yellow bricks as well as actual reported emissions from producers of expanded clay products and an assumed product mix. The reliability of the EFs therefore depends on non-verifiable assumptions.

Changes in the methodologies do not change/improve the base year estimates and the best and most precise estimate for the recent years is considered to be the estimates made by the companies for EU-ETS.

4.2.5 Recalculations and improvements

Information on import/export of cement clinker has been included.

4.2.6 Source specific planned improvements

Company specific information on consumption of CaCO_3 for flue gas cleaning at waste incineration plants will be included for 2010 and previous years as far as possible. A challenge to solve is to distinguish between limestone and burnt lime as some plants use the term limestone for limestone as well as lime in their environmental reports. The mass balance for production of lime/slaked lime will be investigated as some of the slaked lime may be based on imported lime.

4.3 Chemical industry (2B)

4.3.1 Source category description

The subsector *Chemical industry* (2B) covers the following processes:

- Production of nitric acid/fertiliser.
- Production of catalysts/fertilisers.

Production of nitric acid is identified as a key category due to the trend. However this is due to the closing of the lone plant producing nitric acid in Denmark in 2004.

The time series for emission of CO_2 and N_2O from *Chemical industry* (2B) are presented in Table 4.16.

Table 4.16 Time series for emission of greenhouse gasses from Chemical industry (2B).

2B	1990	1995	2000	2005	2007	2008	2009	2010	2011	2012
2. Nitric acid production (kt N_2O)	3.36	2.92	3.24	NO	NO	NO	NO	NO	NO	NO
2. Nitric acid production (kt CO_2 eqv.)	1 043	904	1 004	NO	NO	NO	NO	NO	NO	NO
5. Other (kt CO_2)	0.80	0.80	0.65	3.01	1.35	1.20	1.42	2.12	1.39	1.41
Total (kt CO_2 eq.)	1 044	905	1 004	1.06	1.35	1.20	1.42	2.12	1.39	1.41

The emissions are extracted from the CRF tables and the values are rounded.

The emission of N_2O from nitric acid production is the most considerable source of GHG from the chemical industry. The trend for N_2O from 1990 to 2003 shows a decrease from 3.36 to 2.89 kt, i.e. -14 %, and a 40 % decrease from 2003 to 2004. However, the activity and the corresponding emission show considerable fluctuations in the period considered and the decrease from 2003 to 2004 can be explained by the closing of the plant in the middle of 2004.

From 1990 to 2012, the emission of CO_2 from the production of catalysts/fertilisers has increased from 0.80 to 2.12 kt with maximum in 2010, due to an increase in the activity as well as changes in raw material consumption.

4.3.2 Methodological issues

The N₂O emission from the production of nitric acid/fertiliser is based on measurement data for 2002. For the previous years, the N₂O emission has been estimated from annual production statistics from the company and an emission factor of 7.5 kg N₂O per tonne nitric acid, based on the 2002 measurements (Kemira Growhow, 2004). Table 4.17 presents default EF for N₂O emission from nitric acid production.

Table 4.17 Default EF nitric acid production (IPCC, 1997, vol. 3, p. 2.18).

Geographical area	kg N ₂ O per tonne nitric acid
USA	2 - 9
Norway - modern, integrated plants	< 2
- atmospheric plants	4 - 5
- medium pressure plants	6 - 7.5
Japan	2.2 - 5.7

Specific information on applied technology is not available, however, the EF measured by the Danish nitric acid plant is in accordance with the default emission factors presented by IPCC (1997).

The production of nitric acid in Denmark ceased in the middle of 2004 and the company relocated the production to a more modern facility in another country.

The CO₂ emission from the production of catalysts/fertilisers is based on information in an environmental report from the company (Haldor Topsøe, 2013b), combined with personal communication. In the environmental report, the company has estimated the amount of CO₂ from the process and the amount from energy conversion. Based on information from the company, the emission of CO₂ has been calculated from the composition of raw materials used in the production (for the years 1990 and 1996-2004). The raw materials are e.g.: CaCO₃, CoCO₃, CsCO₃, Cu₂CO₃, K₂CO₃, MnCO₃, and ZnCO₃; however, the actual composition is confidential. For 2005 the EF is assumed to be the same as in 2004 based on the same activity (produced amount). For the years 1991-1995, the production, as well as the CO₂ emission, has been assumed to remain the same as in 1990.

The producer of catalysts/fertilisers is obliged to report energy related CO₂ emissions to EU-ETS but not the process related emissions. For the years 2006-2012 the process related emissions have been estimated as the difference between the total CO₂ emission reported by the company in yearly environmental reports (Haldor Topsøe, 2013b) and energy related CO₂ emission reported by the company to EU-ETS (Haldor Topsøe, 2013a).

4.3.3 Uncertainties and time series consistency

The time series are presented in Table 4.9. The applied methodology regarding N₂O is considered to be consistent. The activity data is based on information from the specific company. The emission factor applied has been constant for the whole time series and is based on measurements in 2002. The production equipment has not been changed during the period.

The estimated CO₂ emissions from production of catalysts/fertilisers are considered to be consistent as they are based on stoichiometric relations combined with company assumptions for the years 1991-1995.

The source specific uncertainties for the chemical industry are presented in Section 4.9. The overall uncertainty estimate is presented in Chapter 1.7.

4.3.4 Recalculations and improvements

The process related CO₂ emission from production of catalysts/fertilisers has recalculated for the years 2006-2012 by inclusion of available company reports to EU-ETS.

4.3.5 Source specific planned improvements

No improvements are planned for this sector.

4.4 Metal production (2C)

4.4.1 Source category description

The subsector *Metal production* (2C) covers the following process:

- Steelwork

The time series for emission of CO₂ from *Metal production* (2C) is presented in Table 4.18. The emissions are extracted from the CRF tables and the values presented are rounded.

Table 4.18 Time series for emission of CO₂ (ktonne) from Metal production (2C).

2C	1990	1995	2000	2005	2007	2008	2009	2010	2011	2012
1. Iron and steel production	28.4	38.6	40.7	15.6	NO	NO	NO	NO	NO	NO

From 1990 to 2001, the CO₂ emission from the electro-steelwork increased from 28 to 47 ktonnes, i.e. by 68 %. The increase in CO₂ emission is similar to the increase in the activity as the consumption of metallurgical coke per amount of steel sheets and bars produced has almost been constant during the period. The electro-steelwork reopened and closed down again in 2005.

4.4.2 Methodological issues

The CO₂ emission from the consumption of metallurgical coke at steelworks has been estimated from the annual production of steel sheets and steel bars combined with the consumption of metallurgical coke per produced amount (Stålvalseværket, 2002). The carbon source is assumed to be coke and all the carbon is assumed to be converted to CO₂ as the carbon content in the products is assumed to be the same as in the iron scrap. The emission factor (consumption of metallurgical coke per tonnes of product) has been almost constant from 1994 to 2001; steel sheets: 0.012-0.018 tonne metallurgical coke per tonne and steel bars: 0.011-0.017 tonne metallurgical coke per tonne. The emission factor (3.6 tonnes CO₂ per tonne metallurgical coke) is based on values in the IPCC-guidelines (IPCC (1997), vol. 3, p. 2.26). The CO₂ emission has been calculated from amounts of final products but related to amount of steel scrap handled at the electro steelwork. Emissions of CO₂ for 1990-1991 and for 1993 have been determined with extrapolation and interpolation, respectively.

4.4.3 Uncertainties and time series consistency

The time series (see Table 4.18) is considered to be consistent as the same methodology has been applied for the whole period. The activity, i.e. amount of steel sheets and bars produced as well as consumption of metal-

lurgical coke, has been published in environmental reports. In 2002, production stopped. For 2005 the production has been assumed to be one third the production in 2001 as the steelwork was operating between 4 and 6 months in 2005.

The source specific uncertainties for the metal production are presented in Section 4.9. The overall uncertainty estimate is presented in Chapter 1.7.

4.4.4 Recalculations and improvements

No source specific recalculations have been performed regarding emissions from the metal production.

4.4.5 Source specific planned improvements

Consumption of reductants in secondary metallurgical processes (e.g. lead) will be investigated.

4.5 Other production (2D)

4.5.1 Source category description

The subsector *Other production*, Food and Drink (2D2) cover the following processes:

- Bread production
- Beer production
- Spirits production
- Sugar production
- Meat (fish etc. frying/curing)
- Margarine and solid cooking fats
- Coffee roasting

Sugar production is the only process contributing to emission of CO₂ whereas all processes contribute to emission of NMVOC. The time series for emission of CO₂ from *Other production*, Food and Drink (2D) is presented in Table 4.19.

Table 4.19 Time series for emission of CO₂ (kt) from Other production, Food and Drink (2D2).

2D	1990	1995	2000	2005	2007	2008	2009	2010	2011	2012
2. Food and Drink	4.45	3.91	3.90	4.46	1,72	2,67	1,92	1,56	2,01	2,24

The emissions are extracted from The CRF tables and the values are rounded.

4.5.2 Methodological issues

The CO₂ emission from the refining of sugar is estimated from production statistics for sugar and a number of assumptions: consumption of 0.02 tonne CaCO₃ per tonne sugar and precipitation of 90 % CaO resulting in an emission factor at 0.0088 tonne CO₂ per tonne sugar (consumption: 2 % CaCO₃ per tonne sugar beets, 10 % sugar in sugar beets). The assumptions are based on environmental reports covering the year 2002. However, from the year 2006-2012 the CO₂ emission compiled by the company for EU-ETS is used in the inventory (Nordic Sugar, 2013). During the preliminary work to establishment of EU-ETS in Denmark the relevant companies reported energy

and process related CO₂ emissions for the years 1998-2002 (DEA, 2004). Based on the process CO₂ emissions and statistical information (Statistics Denmark, 2013) the actual emission factors were determined to respectively 0.0048, 0.0038, 0.0046, 0.0039, and 0.0051 tonne CO₂ per tonne product (average: 0.0044).

The time series for sugar production is presented in Table 4.20.

Table 4.20 Production of sugar (tonne) (Statistics Denmark, 2013).

2D	1990	1995	2000	2005	2007	2008	2009	2010	2011	2012
Sugar	505 709	444 143	487 107	488 934	329 811	400 261	428 446	262 072	218 065	262 026

4.5.3 Uncertainties and time series consistency

The time series is presented in Table 4.19. The same methodology has been applied for 1990-2005. From 2006-2012 data from EU-ETS has been available and therefore included in the inventory.

The source specific uncertainties for the chemical industry are presented in Section 4.9. The overall uncertainty estimate is presented in Chapter 1.7.

4.5.4 Recalculations and improvements

No source specific recalculations have been performed for the sector *Food and Drink*.

4.5.5 Source specific planned improvements

Implementation of an EF based on information in the company reporting to EU-ETS will be investigated.

4.6 Production of Halocarbons and SF₆ (2E)

There is no production of Halocarbons or SF₆ in Denmark.

4.7 Metal production (2C) and consumption of Halocarbons and SF₆ (2F)

4.7.1 Overview of the sector

The sub-sector *Consumption of halocarbons and SF₆* (2F) includes the following source categories and the following f-gases of relevance for Danish emissions:

- 2C4: SF₆ used in Magnesium Foundries: SF₆
- 2F1: Refrigeration: HFC-32, -125, -134a, -152a, -143a, PFC (C₃F₈)
- 2F2: Foam blowing: HFC-134a, -152a
- 2F4: Aerosols/Metered dose inhalers: HFC-134a
- 2F8: Production of electrical equipment: SF₆
- 2F9: Other processes (laboratories, double glaze windows, fibre optics): SF₆, HFC-23, CF₄, C₃F₈, C₄F₈

The description of consumption and emission of f-gases given below is based on an inventory published as (Poulsen & Musaeus, 2014). For further details refer to this report.

General trends

A quantitative overview is given below for each of these source categories and each f-gas, showing their emissions in tonnes through the times-series. The data are extracted from the CRF tables that form part of this submission and the data presented are rounded values. It must be noted that the inventories for the years 1990-1994 might not cover emissions of these gases in full. The choice of base-year for these gases under the Kyoto Protocol is 1995 for Denmark.

Figure 4.5 present the emission trends for f-gases within the different subsectors.




Figure 4.5 Time series for the sub-sectors (ktonne CO₂ equivalents).

The sub-sectors will be described below in the following sections:

- Refrigeration (2F1)
- Foam blowing (2F2)
- Aerosols/metered dose inhalers (2F4)
- Other processes (2C4, 2F8, 2F9)

The emission of SF₆ has been decreasing in recent years due to the fact that activities under Magnesium Foundry no longer exist and due to a decrease in the use of electric equipment. Also, a decrease in "other" occurs until 2010, which for SF₆ is used in window plate production use, laboratories and in the production of running shoes. The increase in 2011 and 2012 is explained in section 4.7.5.

The emission of HFCs increased rapidly in the 1990s and, thereafter, increased more modestly due to a modest increase in the use of HFCs as a refrigerant and a decrease in foam blowing. The f-gases have been regulated in two ways since 1 March 2001. For some types of use there is a ban on use of the gases in new installations and for other types of use, taxation is in place. These regulations seem to have influenced emissions so that they now only increase modestly.

The phase out of f-gasses has in particular been effective within the foam blowing sector and refrigeration installations. With respect to foam blowing, there was a stepwise phase-out of HFC-134a used for foam blowing in hard and soft foam production, during the period 2001-2004. In 2006, all foam production plants in Denmark had substituted HFCs (substitutes have not

been included in the inventory). Especially the phase-out of HFCs in soft foam is significant for the GWP emission in this period.

With respect to HFC from refrigeration, it is not possible to determine a stable decreasing trend yet. Since the introduction of taxes on HFC's in 2001, the consumption decreased in 2002-2003, but then the consumption of HFCs for refrigeration purposes increased again. Especially HFC-404a and HFC-134a increased. This increase is explained with other initiatives in Danish legislation, where new refrigeration systems containing HCFC-22 (ODP) was banned from 2001. It caused a boom in refrigeration systems using HFCs during 2002-2004, because the HFC technology was cheap and well proven. Thus, the consumption of HFCs for refrigeration has changed after 1 January 2007, where new larger HFC installations with stocks exceeding 10 kg are banned. Alternative refrigeration technologies based on CO₂, propane/butane and ammonia is now introduced and available for customers. Butane/propane and ammonia used in refrigeration have not been included in the inventory.

PFCs are emitted in small amounts from refrigeration systems and from optics fibre production, whereas, the consumption as cleaning agent and for new refrigeration systems has been phased out in 2002.

Table 4.21 and Figure 4.6 quantify an overview of the emissions of the gases in CO₂ equivalents. The reference is the trend table as included in the CRF table for year 2011.

Table 4.21 Time series for emission of HFCs, PFCs and SF₆ (kt CO₂ equivalents.).

	1990	1995	2000	2005	2007	2008	2009	2010	2011	2012
HFCs	-	218	607	802	850	853	799	804	759	657
PFCs	-	0.50	17.9	13.9	15.4	12.8	14.2	13.3	11.1	8.54
SF ₆	44.5	107	59.2	21.8	30.3	31.6	36.7	37.9	73.2	118
Total	44.5	326	684	838	896	897	850	855	843	784




Figure 4.6 Time series for emission of HFCs, PFCs and SF₆ (kt CO₂ equivalents.).

The decrease in the SF₆ emission has brought its emissions in CO₂ equivalents down to the level of PFC. Overall, and for all uses, the most dominant group by far is HFCs. In this grouping, HFCs constitute a key category, both with regard to the level and trend analysis.

General methodology

The data for emissions of HFCs, PFCs, and SF₆ have been obtained in continuation on work on inventories for previous years. The determination includes the quantification and determination of any import and export of HFCs, PFCs, and SF₆ contained in products and substances in stock form. This is in accordance with the IPCC guidelines (IPCC (1997), vol. 3, p. 2.43ff), as well as the relevant decision trees from the IPCC Good Practice Guidance (IPCC, 2000) p. 3.53ff).

For the Danish inventories of f-gases, a Tier 2 bottom-up approach is basically used. As for verification using import/export data, a Tier 2 top-down approach is applied. In an annex to the f-gas inventory report 2011 (Poulsen & Musaeus, 2014)), there is a specification of the approach applied for each sub-source category.

The following sources of information have been used:

- Importers, agency enterprises, wholesalers and suppliers.
- Consuming enterprises, and trade and industry associations.
- Recycling enterprises and chemical waste recycling plants.
- Statistics Denmark.
- Danish Refrigeration Installers' Environmental Scheme (KMO).
- Previous evaluations of HFCs, PFCs and SF₆.

Suppliers and/or producers provide consumption data of f-gases. Emission factors are primarily defaults from the GPG, which are assessed to be applicable in a national context. In case of commercial refrigerants and Mobile Air Condition (MAC), information from Danish suppliers has been used. The actual amount of f-gas used for refilling is used as an estimate on the actual emission.

Import/export data for sub-source categories where import/export is relevant (MAC, fridge/freezers for household) are quantified on estimates from import/export statistics of products + default values of the amount of gas in the product. The estimates are transparent and described in the annex to the report referred to above.

The Tier 2 bottom-up analysis used for determination of emissions from HFCs, PFCs, and SF₆ covers the following activities:

- Screening of the market for products in which f-gases are used.
- Determination of averages for the content of f-gases per product unit.
- Determination of emissions during the lifetime of products and disposal.
- Identification of technological development trends that have significance for the emission of f-gases.
- Calculation of import and export on the basis of defined key figures, and information from Statistics Denmark on foreign trade and industry information.

The determination of emissions of f-gases is based on a calculation of the actual emission. The actual emission is the emission in the evaluation year, accounting for the time lapse between consumption and emission. The actual emission includes Danish emissions from production, from products during their lifetimes and from waste products.

Consumption and emissions of f-gases are, whenever possible, determined for individual substances, even though the consumption of certain HFCs has been very limited. This has been carried out to ensure transparency of evaluation in the determination of GWP values. However, the continued use of a category for *Other HFCs* has been necessary since not all importers and suppliers have specified records of sales for individual substances.

The potential emissions have been calculated as follows:

Potential emission = import + production - export - destruction/treatment.

Table 4.22 Content (w/w%)¹ of "pure" HFC in HFC-mixtures, used as trade names.

HFC mixtures	HFC-32	HFC-125	HFC-134a	HFC-143a	HFC-152a	HFC-227ea
	%	%	%	%	%	%
HFC-365						8
HFC-401a					13	
HFC-402a		60				
HFC-404a		44	4	52		
HFC-407a	23	25	52			
HFC-410a	50	50				
HFC-507a		50		50		

1. The mixtures do also contain substances that do not have GHG potential and therefore, the substances do not sum up to 100 %.

The substances have been accounted for in the survey according to their trade names, which are mixtures of HFCs used in the CRF, etc. In the transfer to the "pure" substances used in the CRF reporting schemes, the ratios provided in Table 4.22 have been used.

The national inventories for f-gases are provided and documented in a yearly report (Poulsen & Musaeus, 2014). Furthermore, detailed data and calculations are available and archived in an electronic version. The report contains summaries of methods used and information on sources as well as further details on methodologies.

4.7.2 2F1 Refrigeration

Source category description

2F1 Refrigeration consists of the following processes:

- Household fridges/freezers
- Commercial refrigeration
- Transport refrigeration
- Mobile air conditioning
- Stationary air conditioning.

Table 4.23 presents the emissions of f-gases from consumption of HFCs and PFC in refrigeration and air conditioning systems.

Table 4.23 Emission of f-gases from consumption of HFCs and PFC in refrigeration and air condition systems (2F1 Refrigeration) (t).

	1990	1995	2000	2005	2007	2008	2009	2010	2011	2012
HFC-32	NO	0,11	5,75	13,7	15.4	16.8	17.6	17.5	17.1	16.2
HFC-125	NO	2,58	43,1	67,7	73.6	75.3	74.7	72.8	66.7	60.0
HFC-134a	NO	14,3	112	181	198	198	167	187	193	160
HFC-152a	NO	NO	0,58	0,26	0.17	0.14	0.11	0.093	0.077	0.077
HFC-143a	NO	2,43	39,6	60,3	65.6	66.0	64.6	62.5	56.0	49.3
PFC (C ₃ F ₈)	NO	0,072	2,29	1,99	1.51	1.29	1.13	1.00	0.90	0.80

Methodological issues

The data collection is described in the Section 4.7.1 Overview of the sector, General methodology.

The activity data expressed as total amount of HFCs and PFC filled into new products and present in operating systems are presented in Table 4.24 and Table 4.25 respectively.

Table 4.24 Consumption (filled into new products) of HFCs and PFC in refrigeration and air condition systems (2F1 Refrigeration) (t per year).

	1990	1995	2000	2005	2007	2008	2009	2010	2011	2012
Domestic refrigeration										
HFC-125	NO	0.88	3.96	1.61	1.32	0.86	0.53	0.64	0.79	0.75
HFC-134a	NO	6.16	2.62	65.7	33.6	37.7	17.6	6.82	9.28	9.49
HFC-143a	NO	1.04	4.68	1.90	1.56	1.02	0.62	0.76	0.93	0.89
Commercial Refrigeration ¹										
C ₃ F ₈	NO	1.50	6.30	0.45	0.090	0.06	NO	NO	NO	NO
HFC-125	NO	66.3	118	91.9	91.8	79.5	65.8	67.9	71.0	69.6
HFC-134a	NO	4.68	203	151	106	127	136	106	117	124
HFC-143a	NO	60.8	108	81.4	71.3	55.4	55.4	54.5	53.3	53.4
HFC-152a	NO	NO	1.30	NO	NO	NO	NO	NO	NO	NO
HFC-32	NO	7.00	22.3	21.4	29.6	30.9	17.4	20.2	24.6	22.6
Transport Refrigeration										
HFC-125	NO	1.92	7.92	3.28	0.37	3.24	2.57	2.69	2.90	2.97
HFC-134a	NO	0.12	0.72	0.79	0.44	0.79	0.75	0.74	0.86	0.57
HFC-143a	NO	1.56	9.36	3.87	0.43	3.83	3.04	3.18	3.43	3.51
Mobile Air-Conditioning										
HFC-125	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
HFC-134a	NO	NO	24,0	33,3	35.2	35.7	43.8	67.3	74.1	58.6
HFC-143a	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO

1. Including stationary A/C.

Table 4.25 HFCs and PFC present in operating refrigeration and air condition systems (2F1 Refrigeration) (t).

	1990	1995	2000	2005	2007	2008	2009	2010	2011	2012
Domestic refrigeration										
HFC-125	NO	0.86	25.9	35.1	37.5	38.0	38.1	37.5	36.4	33.8
HFC-134a	NO	165	625	824	854	855	839	811	717	748
HFC-143a	NO	1.02	30.6	41.5	44.3	44.9	45.0	44.4	43.1	40.0
Commercial Refrigeration ¹										
C ₃ F ₈	NO	1.92	25.9	17.6	12.9	11.3	10.0	8.97	8.04	7.18
HFC-125	NO	77.7	414	659	712	707	688	626	580	517
HFC-134a	NO	79.5	690	1 076	1 157	1 109	1 088	1 074	930	901
HFC-143a	NO	71.5	366	580	619	604	582	517	468	410
HFC-152a	NO	NO	5.75	2.12	1.38	1.13	0.93	0.77	0.65	0.65
HFC-32	NO	6.90	70.7	141	164	173	172	167	164	152
Transport Refrigeration										
HFC-125	NO	2.15	26.4	15.7	13.4	14.2	14.3	14.4	14.7	11.3
HFC-134a	NO	0.14	9.87	7.06	5.80	5.57	5.33	5.13	5.08	4.42
HFC-143a	NO	1.84	28.2	17.5	15.2	16.2	16.4	16.6	17.0	13.1
Mobile Air-Conditioning										
HFC-125	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
HFC-134a	NO	NO	149	213	229	231	229	231	230	230
HFC-143a	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO

1. Including stationary A/C.

The applied EFs are presented in Table 4.26. The EFs for commercial refrigerators, mobile A/C, and transport refrigeration has been assessed and compared with national conditions (Poulsen, 2002; citation from Poulsen & Musaeus, 2014).

Table 4.26 Applied EF for refrigeration and air condition systems (2F1 Refrigeration) (Poulsen & Musaeus, 2014).

	Assembly, %	Stock, % per annum	Lifetime
Household fridges and freezers	2	1	15 years
Commercial refrigerators	1.5	10	
Mobile air conditioning systems	0.5	33	
Transport refrigeration	0.5	17	6-8 years

Detailed information on the amount of HFCs used for refilling of mobile A/C has been available for 2009 - 2011, and therefore, a new approach has been implemented in the calculation of emissions. HFCs for mobile A/C are only used for refilling, and therefore the amount used for mobile A/C is assumed to be the same as the amount emitted during use (Poulsen & Musaeus, 2014):

Consumption of HFC for MAC = refilled stock = emission

Uncertainties and time series consistency

See Section 4.7.7 Uncertainties and time series consistency.

Recalculations and improvements

No source specific recalculations have been performed regarding emissions from consumption of HFCs and PFC for refrigeration and air conditioning.

Source specific planned improvements

No improvements are planned for this sector.

4.7.3 2F2 Foam blowing

Source category description

2F2 Foam blowing consists of the following processes:

- Hard foam, refrigerators
- System foam, shoes etc.
- Soft foam

Table 4.27 present the emissions of f-gases from consumption of HFCs in foam blowing.

Table 4.27 Emission of HFCs from consumption in foam blowing (2F2) (t).

	1990	1995	2000	2005	2007	2008	2009	2010	2011	2012
HFC-32	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
HFC-125	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
HFC-134a	NO	136	127	91.2	78.9	78.6	73.2	66.7	59.2	50.4
HFC-152a	NO	43.4	16.2	1.49	2.82	3.39	3.61	4.29	4.07	4.75

Methodological issues

The data collection is described in the Section 4.7.1 Overview of the sector, General methodology.

Table 4.28 and Table 4.29 present the consumption of HFCs in foam blowing and the amount accumulated in stock, respectively.

Table 4.28 Consumption of HFCs in foam blowing (2F2) (t).

	1990	1995	2000	2005	2007	2008	2009	2010	2011	2012
Hard foam										
HFC-134a	NO	193	220	52.8	94.6	76.1	0.06	0.17	NA	NA
HFC-152a	NO	4.00	1.00	5.50	13.0	15.0	12.0	15.0	8.00	13.0
Soft foam										
HFC-125	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
HFC-134a	NO	105	43.9	11.9	NO	NO	NO	NO	NO	NO
HFC-152a	NO	43.0	15.4	NO	NO	NO	NO	NO	NO	NO
HFC-32	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO

Table 4.29 HFCs present as stock in hard foam (2F2) (t).

	1990	1995	2000	2005	2007	2008	2009	2010	2011	2012
Hard foam										
HFC-134a	NO	416	1 415	1 258	1 095	1 005	885	762	642	519
HFC-152a	NO	3.60	15.3	24.8	41.9	53.6	62.0	72.7	76.6	84.9

The applied EFs are presented in Table 4.30.

Table 4.30 Applied EFs for foam blowing (2F2) (Poulsen & Musaeus, 2014).

	Consumption; %	Stock, %	Lifetime
Foam in household fridges and freezers	10	4.5 annual	15 years
Soft foam (open cell)	100 ¹		
Joint filler	100 ¹		
Foaming of polyether (shoe soles)	15	4.5	3 years
System foam	0 ²	- ³	

1. 100% emission during the first year after production.

2. No emission during production of system foam.

3. System foam is only produced for export.

System foam is produced in a closed environment and is only produced for export. Therefore, the consumption of HFCs does not contribute to the Danish stock.

Uncertainties and time series consistency

See Section 4.7.7 Uncertainties and time series consistency.

Recalculations and improvements

A few corrections have been made in the CRF for consumption of HFC-134a to hard foam – IEF and stock, however, no methodological changes have been implemented.

Source specific planned improvements

No improvements are planned for this sector.

4.7.4 2F4 Aerosols/Metered dose inhalers

Source category description

2F4 Aerosols/Metered dose inhalers consist of HFCs used for:

- Propellant in aerosols
- Medical dose inhalers

Table 4.31 present the emissions of f-gases from consumption of HFCs in aerosols and medical dose inhalers.

Table 4.31 Emissions of HCF-134a from consumption of HFC in aerosols/medical dose inhalers (2F4) (t).

	1990	1995	2000	2005	2007	2008	2009	2010	2011	2012
HFC-134a	NO	NO	14.5	16.1	16.0	14.3	13.6	12.9	12.5	12.2

Methodological issues

The data collection is described in the section 4.7.1 Overview of the sector, General methodology.

Table 4.32 present the emissions of f-gases from consumption of HFCs in aerosols and medical dose inhalers.

Table 4.32 Consumption of HFC-134a in aerosols/medical dose inhalers (2F4) (t).

	1990	1995	2000	2005	2007	2008	2009	2010	2011	2012
Metered dose inhalers	NO	NO	1.61	5.63	7.62	7.23	7.07	7.24	6.93	6.84
Aerosols	NO	NO	11.5	15.0	7.00	7.10	6.00	5.23	5.36	5.31

The applied EF is presented in Table 4.33.

Table 4.33 Applied EF for aerosols/medical dose inhalers (Poulsen & Musaeus, 2014).

	Consumption/filling	Stock	Lifetime
Aerosols	0 %	50 % first year 50 % second year	2 years
Medical dose inhalers	0 %	50 % first year 50 % second year	2 years

Uncertainties and time series consistency

See Section 4.7.7 Uncertainties and time series consistency.

Recalculations and improvements

No source specific recalculations have been performed regarding emissions from use of HFC-134a in aerosols and medical dose inhalers.

Source specific planned improvements

No improvements are planned for this sector.

4.7.5 Other processes (2C4, 2F8 and 2F9)

Source category description

Other processes (2C4, 2F8 and 2F9) consist of the following processes:

- Consumption of SF₆ in magnesium foundries; see Table 4.28
- Consumption of SF₆ in electrical equipment; see Table 4.29
- Consumption of SF₆ in running shoes; see Table 4.30
- Consumption of SF₆ in laboratories; see Table 4.30
- Consumption of SF₆ in double glazed windows; see Table 4.30
- Consumption of HFCs and PFCs in fibre optics; see Table 4.30
- Consumption of PCFs as detergent; see Table 4.30

Table 4.34-4.36 presents the emissions of f-gases from consumption of HFCs, PFC and SF₆ in other processes.

Table 4.34 Emissions from SF₆ used in magnesium foundries (2C4) (t).

	1990	1995	2000	2005	2007	2008	2009	2010	2011	2012
SF ₆ used in magnesium foundries	1.30	1.50	0.89	NO	NO	NO	NO	NO	NO	NO

Table 4.35 Emissions from consumption of SF₆ in electrical equipment (2F8) (t).

	1990	1995	2000	2005	2007	2008	2009	2010	2011	2012
SF ₆	0.060	0.16	0.47	0.52	0.63	0.68	0.61	0.59	0.59	0.55

Table 4.36 Emissions from consumption of SF₆, HFCs, and PFCs in other processes (2F9) (t)

	1990	1995	2000	2005	2007	2008	2009	2010	2011	2012
SF ₆	0.50	2.83	1.12	0.39	0.64	0.65	0.93	0.99	2.47	4.38
HFC-23	NO	NO	NO	NO	0.24	0.12	0.24	0.36	0.36	0.12
CF ₄	NO	NO	NO	NO	0.14	0.11	0.36	0.36	0.20	0.18
C ₃ F ₈	NO	NO	0.27	NO	NO	NO	NO	NO	NO	NO
C ₄ F ₈	NO	NO	NO	NO	0.45	0.35	0.45	0.45	0.40	0.20

The increase in emission of SF₆ in 2011 is caused by disposal of double glazed windows after 20 years expected lifetime.

Methodological issues

The data collection is described in the Section 4.7.1 Overview of the sector, General methodology. Activity data are presented in Table 4.37-4.38.

Table 4.37 HFCs, PFC and SF₆ consumed in other processes (t).

	1990	1995	2000	2005	2007	2008	2009	2010	2011	2012
Electrical equipment										
SF ₆	1.20	1.40	4.00	3.58	5.11	5.59	3.75	3.18	2.87	1.86
Detergent										
C ₃ F ₈	NO	NO	0.54	NO	NO	NO	NO	NO	NO	NO
Double glaze windows										
SF ₆	NO	13.5	4.13	NO	NO	NO	NO	NO	NO	NO
Fibre optics										
c-C ₄ F ₈	NO	NO	NO	NO	0.45	0.35	0.45	0.45	0.40	0.20
CF ₄	NO	NO	NO	NO	0.14	0.11	0.36	0.36	0.20	0.18
HFC-23	NO	NO	NO	NO	0.24	0.12	0.24	0.36	0.36	0.12
Laboratories										
SF ₆	NO	0.40	NO	NO	0.26	0.27	0.55	0.64	0.74	0.73
Shoes										
SF ₆	NO	0.11	0.11	NO	NO	NO	NO	NO	NO	NO
Various										
SF ₆	0.10	NO	NO	NO	NO	NO	NO	NO	NO	NO

Table 4.38 SF₆ accumulated as stock in other processes (t).

	1990	1995	2000	2005	2007	2008	2009	2010	2011	2012
Electrical equipment	9.40	26.2	57.3	67.9	75.5	80.4	83.5	85.5	88.4	89.7
Double glaze windows	NO	25.0	38.4	36.6	75.5	80.4	83.5	85.5	88.4	89.7

The applied EFs are presented in Table 4.39. Special attention has been given to use of SF₆ as insulation in high-voltage plants (Poulsen, 2001; ELTRA, 2004; citation from Poulsen & Musaeus, 2014).

Table 4.39 Applied EFs for other processes (Poulsen & Musaeus, 2014).

	Consumption	Stock	Lifetime
	50 % first year		
Liquid cleaners/detergents	50 % second year		
	100 % in the year		
Fibre optics production	of production		
Laboratories	100 %		
Insulation gas in double glaze windows	15 %	1% annual	20 years
		0.5 % annual	
		5 % in reuse/ drawing off	? ³
Insulation gas in high voltage switches	5 %		
Shock-absorbing in Nike Air training footwear	- ¹	- ²	5 years

1. No emission from production in Denmark.

2. Yearly emission has been estimated to 0.11 tonne (Poulsen & Musaeus, 2014).

3. Lifetime unknown.

Uncertainties and time series consistency

See Section 4.7.7 Uncertainties and time series consistency.

Recalculations and improvements

No source specific recalculations have been performed regarding emissions from other use of HFCs, PFCs, and SF₆.

Source specific planned improvements

No improvements are planned for this sector.

4.7.6 QA/QC and verification

Comparison of emissions estimates using different approaches

This comparison of Tier 1 potential emissions has been used for a check on the Tier 2 actual emission estimates. This check was carried out in 1995-1997 and, for all three years, it shows a difference of approximately a factor 3 higher emission by using potential emission estimates.

This comparison of bottom-up estimates has not yet been compared with the top-down Tier 2 approach. This comparison will be developed.

National activity data check

The spreadsheets containing activity data have incorporated several data-control mechanisms, which ensure that data estimates do not contain calculation failures. A very comprehensive QC procedure on the data in the model for the whole time series has been carried out for the 2013 submission in connection with the process which provided, (1) data for the CRF background tables 2(II).F. for the years (1993)-2012 and (2) data for potential emissions in CRF tables 2(I). This procedure consisted of a check of the input data for the model for each substance. As regards the HFCs, this checking was carried out in relation to their trade names. Conversion was made to the HFC substances used in the CRF tables, etc. A QC was that emission of the substances could be calculated and checked comparing results from the substances as trade names and as the "no-mixture" substances used in the CRF.

Emission factors check

Country-specific emission factors are explained and documented for MAC and commercial refrigerants and SF₆ in electric equipment. Separate studies have been carried out and reported; see the previous chapters for references. For other sub-source categories, the country-specific emission factors are assessed to be the same as the IPCC default emission factors.

Emission check

As the f-gas inventory is developed and made available in full in spread sheets, where HFCs data relate to trade names, special procedures are performed to check the full possible correctness of the transformation to the CRF-format through Access databases.

4.7.7 Uncertainties and time series consistency

The time series for actual emissions of halocarbons and SF₆ are presented in Section 4.7.1. The time series are consistent as regards the methodology. The potential emission estimates are only included in the CRF.

Tier 1 and Tier 2 uncertainty estimates have been calculated by use of default uncertainties.

In general, uncertainty in inventories will arise through at least three different processes:

- Uncertainties from definitions (e.g. incomplete, unclear, or faulty definition of an emission or uptake);
- Uncertainties from natural variability of the process that produces an emission or uptake;
- Uncertainties resulting from the assessment of the process or quantity depending on the method used: (i) uncertainties from measuring; (ii) uncertainties from sampling; (iii) uncertainties from reference data that may be incompletely described, and (iv) uncertainties from expert judgement.

Uncertainties due to poor definitions are not expected to be an issue in the f-gas inventory. The definitions of chemicals, the factors, sub-source categories in industries etc. are well defined.

Uncertainties from natural variability are likely to occur over the short-term while estimating emissions in individual years. But over a longer time period, 10-15 years, these variabilities level out in the total emission. This is due to that input data (consumption of f-gases) is known and is valid data, and has no natural variability due to the chemicals stable nature.

Uncertainties that arise due to imperfect measurement and assessment are probably an issue for the:

- Emission from MAC (HFC-134a).
- Emission from commercial refrigerants (HFC-134a).

Due to the limited knowledge for these sources, the expert assessment of consumption of f-gases can lead to inexact values of the specific consumption of f-gases.

The uncertainty varies from substance to substance. Uncertainty is greatest for HFC-134a due to its widespread application in products that are imported and exported. The greatest uncertainty in application is expected to arise from consumption of HFC-404a and HFC-134a in commercial refrigerators and mobile refrigerators. The uncertainty involved in year-to-year data is influenced by the uncertainty associated with the rates at which the substances are released. This results in significant differences in the emission determinations in the short-term (approximately five years); differences that balance in the long-term.

The source specific uncertainties for consumption of halocarbons and SF₆ are presented in Chapter 4.9. The overall uncertainty estimate is presented in Chapter 1.7.

4.8 Other (2G)

4.8.1 Source category description

The subsector *Other* (2G) covers the following process:

- Oxidation of lubricants during use.

The time series for emission of CO₂ from *Other* (2G) is presented in Table 4.40.

Table 4.40 Time series for emission of CO₂ (kt) from Other (2G).

2G	1990	1995	2000	2005	2007	2008	2009	2010	2011	2012
Oxidation of lubricants during use	49.7	48.8	39.7	37.6	37.9	34.0	31.2	33.2	33.2	31.7

The emissions are extracted from the CRF tables and the values are rounded.

The emission of CO₂ from oxidation of lubricants during use is decreasing from 49.7 kt in 1990 to 33.2 kt in 2011.

4.8.2 Methodological issues

The emission of CO₂ from oxidation of lubricants during use is calculated according to the following formula:

$$E_{CO_2} = LC \bullet CC_{\text{lubricant}} \bullet ODU_{\text{lubricant}} \bullet 44/12$$

where:

E_{CO_2} = emission of CO₂

LC = consumption of lubricants

CC = carbon content of lubricant

ODU = amount of lubricant oxidised during use

In the calculation the following default values have been applied: CC = 20.1 kg C per kg lubricant and ODU = 0.2. The activity data applied are presented in Table 4.41.

Table 4.41 Consumption of lubricant oil (TJ) (Danish Energy Agency).

2G	1990	1995	2000	2005	2007	2008	2009	2010	2011	2012
Oxidation of lubricants during use	3 372	3 314	2 693	2 550	2 574	2 307	2 116	2 251	2 251	2 150

4.8.3 Uncertainties and time series consistency

The time series is presented in Table 4.40. The applied methodology has been the same during all the years and is therefore considered to be consistent. The activity data are based on information from Danish Energy Agency. The same emission factor has been used for all the years from 1990 to 2012.

4.8.4 Recalculations and improvements

No source specific recalculations have been performed regarding emissions from the consumption of lubricants.

4.8.5 Source specific planned improvements

No improvements are planned for this sector.

4.9 Uncertainty

4.9.1 Tier 1 uncertainty

The source specific uncertainties for industrial processes are presented in Table 4.42. The uncertainties are based on IPCC guidelines combined with assessment of the individual processes.

The producer has delivered the activity data for production of cement as well as calculated the emission factor based on quality measurements. The uncertainties on activity data and emission factors are assumed to be 1 % and 2 %, respectively.

The activity data for production of lime and bricks are based on information compiled by Statistics Denmark. Due to the many producers and the variety of products, the uncertainty is assumed to be 5 %. The emission factor is partly based on stoichiometric relations and partly on an assumption of the number of yellow bricks. The last assumption has been verified (see Table 4.42). The combined uncertainty is assumed to be 5 %.

The producers of glass and glass wool have registered the consumption of -raw materials containing carbonate. The uncertainty is assumed to be 5 %. The emission factors are based on stoichiometric relations and, therefore, uncertainty is assumed to be 2 %.

The producers have registered the production of nitric acid during many years and, therefore, the uncertainty is assumed to be 2 %. The measurement of N₂O is problematic and is only carried out for one year. Therefore, the uncertainty is assumed to be 25 %.

The uncertainty for the activity data as well as for the emission factor is assumed to be 5 % for production of catalysts/fertilisers and iron and steel production.

The emission of f-gases is dominated by emissions from refrigeration equipment and therefore, the uncertainties assumed for this sector will be used for all the f-gases. The IPCC propose an uncertainty at 30-40 % for regional estimates. However, Danish statistics have been developed over many years and, therefore the uncertainty on activity data is assumed to be 10 %. The uncertainty on the emission factor is, on the other hand, assumed to be 50 %. The base year for f-gases for Denmark is 1995.

Table 4.42 Uncertainties on activity data and emission factors as well as overall trend uncertainties for the different greenhouse gases.

Greenhouse gases	Activity data uncertainty	Emission factor uncertainty				
		CO ₂	N ₂ O ²	HFCs ³	PFCs ³	SF ₆ ³
	%	%	%	%	%	%
2A1. Production of Cement	1	2				
2A2. Production of Lime and Bricks	5	5				
2A3. Limestone and dolomite use	5	5				
2A5. Asphalt roofing	5	25				
2A6. Road paving with asphalt	5	25				
2A7.1 Glass production	5	2				
2A7.2 Yellow bricks	5	2				
2A7.3 Expanded clay	5	2				
2B2. Nitric acid production ²	2		25			
2B5. Other ¹	5	5				
2C1. Iron and Steel production	5	5				
2D. Food and Drink	5	5				
2F. Consumption of HFC	10			50		
2F. Consumption of PFC	10				50	
2F. Consumption of SF ₆	10					50
2G. Other: Lubricants	2	5				
Overall uncertainty in 2012		1.97	-	51.0	51.0	51.0
Trend uncertainty		1.17	-	42.7	240	15.5

1) Production of catalysts/fertilisers.

2) The production closed down in the middle of 2004.

3) The base year for f-gases is for Denmark 1995.

4.9.2 Tier 2 uncertainty

The tier 2 uncertainty for CO₂ emission from industrial processes and consumption of F-gases is presented in Table 4.43 and Table 4.44. The uncertainty estimates are based on the same individual uncertainties as applied for the tier 1 uncertainty estimate.

Table 4.43 Tier 2 uncertainty for industrial processes.

		1990			2012			1990-2012		
		Median Emission	Uncertainty (%)		Median Emission	Uncertainty (%)		Median Emission	Uncertainty (%)	
			Lower (-)	Upper (+)		Lower (-)	Upper (+)		Lower (-)	Upper (+)
CO ₂	ktonnes	2 195	10	13	1 007	2.0	1.9	1 182	9.2	12

Table 4.44 Tier 2 uncertainty for consumption of F-gases.

	Median Emission	1995		Median Emission	2012		1995-2012		
		Uncertainty (%)			Uncertainty (%)		Uncertainty (%)		
		Lower	Upper		Lower	Upper	Lower	Upper	
		(-)	(+)		(-)	(+)	(-)	(+)	
CO ₂ eq. ktonnes	292	22	33	793	25	37	-414	-60	-33

4.10 Quality assurance/quality control (QA/QC)

4.10.1 Internal QA/QC

The approach used for quality assurance/quality control (QA/QC) is presented in Chapter 1.6; see also Nielsen et al. (2012). The present chapter presents QA/QC considerations for industrial processes based on a series of Points of Measuring (PMs); see Chapter 1.6.

Data Storage level 1	1. Accuracy	DS.1.1.1	General level of uncertainty for every dataset including the reasoning for the specific values.
----------------------	-------------	----------	---

The uncertainty assessment has been performed on Tier 1 and Tier 2 level by using default and country specific uncertainty factors. The applied uncertainty factors are presented in Table 4.36.

Data Storage level 1	2.Comparability	DS.1.2.1	Comparability of the emission factors/calculation parameters with data from international guidelines, and evaluation of major discrepancies.
----------------------	-----------------	----------	--

Comparability of the data has not been performed at “Data Storage level 1”. However, investigation of comparability at CRF level is in progress.

The applied data sets are presented in Table 4.39.

Data Storage level 1	3.Completeness	DS.1.3.1	Ensuring that the best possible national data for all sources are included, by setting down the reasoning behind the selection of datasets.
----------------------	----------------	----------	---

The data sources - in general - can be grouped as follows:

- Company specific environmental reports.
- Personal communication with individual companies.
- Company specific information compiled by Danish Energy Agency in relation to the EU-ETS.
- Industrial organisations.
- Statistics Denmark.
- Secondary literature.
- IPCC guidelines.

The environmental reports contribute with company-specific emission factors, technical information and, in some cases, activity data. The environmental reports are primarily used for large companies and, for some companies, are supplemented with information from personal contacts, especially for completion of the time series for the years before the legal requirement to prepare environmental reports (i.e. prior to 1996).

Statistics Denmark is used as source for activity data as they are able to provide consistent data for the period 1990-2012. In the cases where the statistics do not contain transparent data, statistics from industrial organisations are used to generate to required activity data.

For many of the processes, the default emission factors are based on chemical equations and are, therefore, the best choice. In some cases, the default EF has been modified in order to reflect local conditions.

Secondary literature may be used in the interpretation or in disaggregation of the public statistics.

Data Storage level 1	4.Consistency	DS.1.4.1	The original external data has to be archived with proper reference.
----------------------	---------------	----------	--

The original data files are archived in the following folder:

U:\ST_ENVS-Luft-Emi\Inventory\2012\2_Industrial_Processes\Level_1a_Storage.

Data Storage level 1	6.Robustness	DS.1.6.1	Explicit agreements between the external institution holding the data and NERI about the condition of delivery.
----------------------	--------------	----------	---

An agreement regarding inclusion of information - compiled by Danish Energy Agency for EU-ETS - in the Danish GHG-inventory has been signed. The implementation of this information has been introduced for production of cement, bricks, expanded clay products, flue gas cleaning at CHP and sugar refining.

Data Storage level 1	7.Transparency	DS.1.7.1	Listing of all archived datasets and external contacts.
----------------------	----------------	----------	---

The datasets applied are presented in Table 4.45. For the reasoning behind their selection, see DS.1.3.1.

Table 4.45 Applied datasets (archived in: U:\ST_ENVS-Luft-

Emi\LUFT_EMI\Inventory\2012\2_Industrial_Processes\Level_1a_Storage).

\EnvironmentalReports2012\	WasteIncineration (folder)
	AalborgPortland_miljoeredegoerelse_2012
	Ardagh 2012
	Cheminova 2012
	DaniscoGrindsted_GroentRegnskab2012 (folder)
	Faxe Kalk GI Strandvej ovnanlæg Stubberup 2012
	HaldorTopsøe_GroentRegnskab2012 (folder)
	NordicSugar_Nykøbing_GroentRegnskab2012 (folder)
	Rockwool 2012
	Saint Gobain Isover 2012
	\EU-ETS2011\
	CHP (folder)
	Industry (folder)
	CO2udledning_og_energiforbrug_EDO_2012
	\Statistics2011\

894996651012458366525531977_sugar_production
1296026454406188864621548830_bread
1297316516644931165171188125_cattle
1297326501845340665022549357_pigs
1297336487270930464880805095_poultry
1301665535347807555368038512_fats
201311617158130087722KN8Y62151633712_coffee_IE
1305175019872105850216162313_beverage
2013930173742128623422VARER163480203877_cement_lime_
production
201392613847128456841KN8Y47607586821_cement_lime_IE
1311164389470475343927081523_bricks
2013111416345130441698VARER159695496883_asphalt 2
1730176216664248062176998064_asphalt 2_IE
201311617158130087722KN8Y62151633712_coffee_IE
894914413900940644148244741_expanded clay
894996651012458366525531977_sugar_production
Landbrugsstatistik_1985-90
Fiskeristatistik 2012

Data Processing level 1	1. Accuracy	DP.1.1.1	Uncertainty assessment for every data source not part of DS.1.1.1 as input to Data Storage level 2 in relation to type and scale of variability.
----------------------------	-------------	----------	--

The uncertainty assessment has been performed on Tier 1 as well as Tier 2 level, assuming a normal distribution of activity data as well as emission data, by application of default uncertainty factors. Therefore, no considerations regarding distribution or type of variability have been performed.

Data Processing level 1	2. Comparability	DP.1.2.1	The methodologies have to follow the international guidelines suggested by UNFCCC and IPCC.
----------------------------	------------------	----------	---

All methodologies follow UNFCCC and IPCC unless better national methodologies have been identified.

Data Processing level 1	3.Completeness	DP.1.3.1	Identification of data gaps with regard to data sources that could improve quantitative knowledge.
-------------------------	----------------	----------	--

This issue will be investigated further.

Data Processing level 1	4.Consistency	DP.1.4.1	Documentation and reasoning of methodological changes during the time series and the qualitative assessment of the impact on time series consistency.
-------------------------	---------------	----------	---

Recalculations are described in the NIR. A manual log is included in the tool used for data processing at Data Processing level 2. This log also includes changes on Data Processing level 1.

Data Processing level 1	5.Correctness	DP.1.5.2	Verification of calculation results using time series.
-------------------------	---------------	----------	--

The calculations are verified by checking the time series.

Data Processing level 1	5.Correctness	DP.1.5.3	Verification of calculation results using other measures.
-------------------------	---------------	----------	---

A methodology to verify calculation of results using other measures will be developed.

Data Processing level 1	7.Transparency	DP.1.7.1	The calculation principle, the equations used and the assumptions made must be described.
-------------------------	----------------	----------	---

The calculation principles and equations are based on the methodology presented by the IPCC. A detailed description can be found in the sector report for industry (in prep.).

Data Processing level 1	7.Transparency	DP.1.7.2	Clear reference to dataset at Data Storage level 1
-------------------------	----------------	----------	--

The calculation files contain links to the original data files.

Data Processing level 1	7.Transparency	DP.1.7.3	A manual log to collect information about recalculations.
-------------------------	----------------	----------	---

A log on information about recalculation is included in CollectER.

Data Processing level 2	5.Correctness	DS.2.5.1	Check if a correct data import to level 2 has been made
-------------------------	---------------	----------	---

The sector report for industry (in prep.) presents the connection between the datasets on Data Storage level 1 and Data Processing level 2. Individual calculations are used to check the output of the data processing tool used at Data Processing level 2.

Data Storage level 4	4. Consistency	DS.4.4.3	The IEFs from the CRF are checked both regarding level and trend. The level is compared to relevant emission factors to
----------------------	----------------	----------	---

			ensure correctness. Large dips/jumps in the time series are explained.
--	--	--	--

The IEFs are checked by using a tool developed especially for that purpose and outliers are explained.

Data Storage level 4	4. Correctness	DS.4.5.2	Check that additional information and information related to land-use changes has been correctly aggregated compared to the individual submissions of Denmark and Greenland.
-------------------------	----------------	----------	--

The aggregated submission for Denmark and Greenland is checked against the individual submissions for Denmark and Greenland.

4.10.2 External QA/QC

External QA/QC is described for one source: cement production.

Cement production

Aalborg Portland has an environmental management system that meets the requirements in DS/ISO 14001, EMAS etc. (Aalborg Portland, 2012b). The environmental management system is part of an integrated process management system. The system is certified according to the standards by the accredited body: Danish Standards. Information on raw material consumption as well as internal recycling is compiled in an environmental database. Some pollutants (NO_x, SO₂, CO and TSP) are measured continuously. Emission of CO₂ is calculated based on (fuel and) raw material consumption and raw material flow according to an approved CO₂ emission plan (EU-ETS). The CO₂ emission plan has to fulfil the requirements in the guidelines developed by EU (EU, 2007).

4.11 References

Ardagh Glass Holmegaard, 2013a: CO₂-opgørelse og afrapportering 2012. Ardagh Glass Holmegaard A/S. (In Danish) Confidential.

Ardagh Glass Holmegaard, 2013b: Grønt regnskab for Ardagh Glass Holmegaard A/S 2012, CVR nr. 18445042 incl. 2009-2011 (in Danish).

Damolin, 2013: CO₂-opgørelse og afrapportering 2012. Damolin Fur A/S og Damolin Mors A/S. (In Danish) Confidential.

Danish Energy Agency (DEA), 2004: Anders Baunehøj Hansen, personal communication, 15 December 2004.

Danish Energy Agency (DEA), 2013: The Danish energy statistics, Available at: <http://www.ens.dk>

ELTRA, 2004: Anvendelse og håndtering af SF₆-gas i højspændingsanlæg over 100 kV. ELTRA Memo elt2004-47a, 18 marts 2004 (In Danish).

EMEP/EEA, 2009: EMEP/EEA air pollutant emission inventory guidebook - 2009. Technical report No 9/2009. Available at: <http://www.eea.europa.eu/publications/emep-eea-emission-inventory-guidebook-2009>

Energinet.dk, 2007: Baggrundsrapport til Miljøberetning 2006 (in Danish).

EU, 2007: Commission decision of 18 July 2007 establishing guidelines for monitoring and reporting of greenhouse gas emissions pursuant to Directive 2003/87/EC of the European Parliament and of the Council. Available at: <http://eur-lex.europa.eu/>

Faxe Kalk, 2003: Diverse produktblade (in Danish).

Faxe Kalk, 2013a: CO₂-opgørelse og afrapportering 2012. Faxe Kalk, Ovnanlægget Stubberup. (In Danish) Confidential.

Faxe Kalk, 2013b: Grønt regnskab 2012, Faxe Kalk A/S. Ovnanlægget Stubberup.

Haldor Topsøe, 2013a: CO₂-opgørelse og afrapportering 2012. Haldor Topsøe A/S. (In Danish) Confidential.

Haldor Topsøe, 2013b: Miljøredegørelse for katalysatorfabrikken 2012 (17. regnskabsår); inkl. 1996-2011 (in Danish).

Hjelmar, O. & Hansen, J.B., 2002: Restprodukter fra røggasrensning på affaldsforbrændingsanlæg. Nyttiggørelse eller deponering? DHI - Institut for Vand & Miljø. Kursusmateriale fra kurset Røggasrensning 2002. IDA, Brændsels- og Energiteknisk Selskab (in Danish).

Illerup, J.B., Geertinger A.M., Hoffmann, L. & Christiansen, K., 1999: Emissionsfaktorer for tungmetaller 1990-1996. Faglig rapport fra DMU, nr. 301. Miljø- og Energiministeriet, Danmarks Miljøundersøgelse (in Danish).

IPCC, 1997: Revised 1996 IPCC Guidelines for National Greenhouse Gas Inventories. Available at: <http://www.ipcc-nggip.iges.or.jp/public/gl/invs6.htm> (15-04-2007).

IPCC, 2000: Good Practice Guidance and Uncertainty Management in National Greenhouse Gas Inventories. Available at: <http://www.ipcc-nggip.iges.or.jp/public/gp/english/> (15-04-2007).

Kemira GrowHow, 2004: Miljø & arbejdsmiljø. Grønt regnskab 2003; inkl. 1996-2002 (in Danish).

Maxit, 2013: CO₂-opgørelse og afrapportering 2012. Maxit a.s. Hinge og Maxit a.s. Ølst. (In Danish) Confidential.

Miljøstyrelsen, 2012: Affaldsstatistik 2010. Orientering fra Miljøstyrelsen (In Danish; to be published).

Nielsen, M. & Illerup, J.B., 2003: Emissionsfaktorer og emissionsopgørelse for decentral kraftvarme. Eltra PSO projekt 3141. Kortlægning af emissioner fra decentrale kraftvarmeværker. Delrapport 6. Danmarks Miljøundersøgelser. 116 s. Faglig rapport fra DMU nr. 442. Available at: <http://www.dmu.dk/udgivelser/> (in Danish).

Nielsen, O.-K., Plejdrup, M.S., Winther, M., Gyldenkerne, S., Thomsen, M., Fauser, P., Nielsen, M., Mikkelsen, M.H., Albrektsen, R., Hjelgaard, K.,

Hoffmann, L. & Bruun, H.G. 2012. Quality manual for the Danish greenhouse gas inventory. Version 2. Aarhus University, DCE – Danish Centre for Environment and Energy, 44 pp. Scientific Report from DCE – Danish Centre for Environment and Energy No. 47. Available at:
<http://www.dmu.dk/Pub/SR47.pdf>

Nordic Sugar, 2013: CO₂-opgørelse og afrapportering 2012. Nordic Sugar. Fremstilling af sukker. (In Danish) Confidential.

Poulsen, T.S., 2002: Revurdering af emissionsfaktorer for kommercielle køleanlæg, mobile A/C og køleanlæg. Miljørapport nr. 766, Miljøstyrelsen (In Danish).

Poulsen, T.S. & Musaeus M., 2014: The greenhouse gases HFCs, PFCs and SF₆. Danish consumption and emissions, 2012. Environmental Project No xx. In prep.

Poulsen, T.S. 2001: Indsamling og genanvendelse af SF₆ fra elsektoren. Miljøprojekt nr. 592. Miljøstyrelsen (In Danish).

Rexam Glass Holmegaard, 2007: Grønt regnskab for Rexam Glass Holmegaard A/S 2006, CVR nr. 18445042; incl. 1996/97-2006 (in Danish).

Rockwool, 2013: Miljøreddegørelse 2012 for fabrikkerne i Hedehusene, Vamdrup og Øster Doense; incl. 1996-2011 (in Danish).

Saint-Gobain Isover, 2013a: CO₂-opgørelse og afrapportering 2012. Saint-Gobain Isover A/S. (In Danish) Confidential.

Saint-Gobain Isover, 2013b: Miljø- og energireddegørelse 2012; incl. 1996-2011 (in Danish).

Statistics Denmark, 2013: Statbank Denmark. Available at:
www.statbank.dk

Stålvalseværket, 2002: Grønt regnskab og miljøreddegørelse 2001. Det Danske Stålvalseværk A/S; inkl. 1992, 1994-2000 (in Danish).

Tegl Info, 2004: Teglfremstilling i Danmark. Available at:
<http://www.tegl.info/>

Aalborg Portland, 2008: Henrik Møller Thomsen, Personal communication 17 September 2008.

Aalborg Portland, 2013a: CO₂-opgørelse og afrapportering 2012. Aalborg Portland A/S. Fremstilling af klinker (cement); including 2005-2011. Confidential.

Aalborg Portland, 2013b: Environmental report 2012; including 1996-2011.

Aalborg Portland, 2013c: Torben Ahlmann-Laursen, Personal communication 19 November 2013.

5 Solvent and Other Product Use (CRF sector 3)

5.1 Introduction

This section presents the Danish methodology used for calculating emissions from use of solvents and other product use in industries and households that are related to the source categories Paint application (CRF sector 3A), Degreasing and dry cleaning (CRF sector 3B), Chemical products, manufacture and processing (CRF sector 3C) and Other Use of Solvents and Products (CRF sector 3D). Covered pollutants are; NMVOC, CO₂ and N₂O.

Solvents are chemical compounds that are used on a global scale in industries and as constituents in final products to dissolve e.g. paint, cosmetics, adhesives, ink, rubber, plastic, pesticides, aerosols or are used for cleaning purposes, i.e. degreasing. NMVOCs are main components in solvents - and solvent use in industries and households is typically a dominant source of anthropogenic NMVOC emissions (UNFCCC, 2008; Pärt, 2005; Karjalainen, 2005). In industries where solvents are produced or used, NMVOC emissions to air and as liquid can be recaptured and either used or destroyed. Solvent containing products are used indoor and outdoor and the majority of solvent sooner or later evaporate. A small fraction of the solvent ends up in waste or as emissions to water and may finally also contribute to air pollution by evaporation from these compartments. Emission inventories for solvents are based on model estimates, as direct and continuous emissions are rarely measured. In addition to emissions from use of solvents emissions from use of fireworks, tobacco, candles and charcoal for barbeques are included in the source category CRF 3D.

In this section the methodology for the Danish emission inventory for Solvent and Other Product Use is presented and the results for the period 1990 - 2012 are summarised. The method is mainly based on the detailed approach and methodology described in EMEP/EEA (2009) and IPCC (1997 & 2000), and emissions are calculated for industrial sectors, households for the stated CRF sectors, as well as for individual pollutants.

The data presented in Chapter 5 relate to Denmark only, whereas information for Greenland is included in Chapter 16 and for the Faroe Islands in Annex 8.

5.2 Methodology

Emission modelling of solvents can basically be done in two ways: 1) By estimating the amount of (pure) solvents consumed, or 2) By estimating the amount of solvent containing products consumed, taking account of their solvent content (EMEP/EEA, 2009).

In 1) all relevant solvents must be estimated, or at least those together representing more than 90 % of the total pollutant emission, and in 2) all relevant source categories must be inventoried or at least those together contributing more than 90 % of the total pollutant emission. A simple approach is to use a per capita emission for each category, whereas a detailed approach is to get all relevant consumption data (EMEP/EEA, 2009).

The detailed method 1) is used in the Danish emission inventory for solvent use, thus representing a chemicals approach, where each pollutant is estimated separately. The sum of emissions of all estimated pollutants used as solvents equals the pollutant emission from solvent use.

A Tier 1 method is used for determining emissions from fireworks, tobacco, candles and charcoal for barbeques included in CRF 3D, this means a simple multiplication of activity data with emission factors.

5.2.1 Pollutant list

NMVOC is the most abundant chemical group in relation to Solvent and Other Product Use. Additionally there is also some use and/or emissions of NO₂ and CO₂.

The definitions of solvents and VOC that are used in the Danish inventory (Nielsen et al., 2012) are as defined in the solvent directive (Directive 1999/13/EC) of the EU legislation: "Organic solvent shall mean any VOC which is used alone or in combination with other agents, and without undergoing a chemical change, to dissolve raw materials, products or waste materials, or is used as a cleaning agent to dissolve contaminants, or as a dissolver, or as a dispersion medium, or as a viscosity adjuster, or as a surface tension adjuster, or a plasticiser, or as a preservative". VOCs are defined as follows: "Volatile organic compound shall mean any organic compound having at 293.15 K a vapour pressure of 0.01 kPa or more, or having a corresponding volatility under the particular condition of use".

This implies that some NMVOCs, e.g. ethylenglycol, that have vapour pressures just around 0.01 kPa at 20 °C, may only be defined as VOCs at use conditions with higher temperature. However, use conditions under elevated temperature are typically found in industrial uses. Here the capture of solvent fumes is often efficient, thus resulting in small emissions (communication with industries).

The Danish list of NMVOCs comprises approximately 30 compounds or compound groups representing more than 95 % of the total emission from solvent use, cf. Table 5.3. CO₂ conversion factors, where all carbon in a carbon-containing molecule, is converted to CO₂, are also listed in Table 5.3.

Emissions that are emitted from other product use are CO₂, N₂O and NMVOC.

5.2.2 Activity data

For each compound or product a mass balance is formulated:

$$\text{Consumption} = (\text{production} + \text{import}) - (\text{export} + \text{destruction/disposal} + \text{hold-up}) \quad (\text{Eq. 1})$$

Data concerning production, import and export amounts of solvents and solvent containing products are collected from StatBank DK (2012), which contains detailed statistical information. Manufacturing and trading industries are committed to reporting production and trade figures to the Danish Customs & Tax Authorities in accordance with the Combined Nomenclature. Import and export figures are available on a monthly basis from 1990 to present and contain trade information from approximately 200 countries world-wide. Production figures are reported quarterly as industrial commodity statistics by commodity group and unit from 1990 to present.

Destruction and disposal of solvents lower the NMVOC emissions. In principle this amount must be estimated for each compound in all industrial activities and for all uses of solvent containing products. At present the solvent inventory only considers destruction and disposal for a limited number of solvents. For some compound it is inherent in the emission factor, and for others the reduction is specifically calculated from information obtained from the industry or literature.

Hold-up is the difference in the amount in stock in the beginning and at the end of the year of the inventory. No information on solvents in stock has been obtained from industries. Furthermore, the inventory spans over several years so there will be an offset in the use and production, import and export balance over time.

In some industries the solvents are consumed in the process, e.g. in the graphics and plastic industry, whereas in the production of paints and lacquers the solvents are still present in the final product. These products can either be exported or used in the country. In order not to double count consumption amounts of pollutants it is important to keep track of total solvent use, solvents not used in products and use of solvent containing products. Furthermore some pollutants may be represented as individual pollutants and also in chemical groups, e.g. "o-xylene", "mixture of xylenes" and "xylene". Some solvents are better inventoried as a group rather than individual compounds, due to missing information on use or emission for the individual compounds. The Danish inventory considers single compounds, with a few exceptions.

Activity data for solvents are thus primarily calculated from Equation 1 with input from StatBank DK (2012). When StatBank (2012) holds no information on production, import and export or when more reliable information is available from industries, scientific reports or expert judgements the data can be adjusted or even replaced.

Activity data for Other Product Use in CRF 3D are compiled from StatBank DK.

5.2.3 Emission factors

For each pollutant the emission is calculated by multiplying the consumption with the fraction emitted (emission factor), according to:

$$\text{Emission} = \text{consumption} * \text{emission factor}$$

The present Danish method uses emission factors that represent specific industrial activities, such as processing of polystyrene, dry cleaning etc. or that represent use categories, such as paints and detergents. Some pollutants have been assigned emission factors according to their water solubility. Higher hydrophobicity yields higher emission factors, since a lower amount ends in waste water, e.g. ethanol (hydrophilic) and turpentine (hydrophobic).

Emission factors for solvents are categorised in four groups in ascending order: (1) Lowest emission factors in the chemical industry, e.g. lacquer and paint manufacturing, due to emission reducing abatement techniques and destruction of solvent containing waste, (2) Other industrial uses, e.g. graphic industry, have higher emission factors, (3) Non-industrial use, e.g. auto

repair and construction, have even higher emission factors, (4) Diffuse use of solvent containing products, e.g. painting, where practically all the pollutant present in the products will be released during or after use.

For a given pollutant the consumed amount can thus be attributed with two or more emission factors; one emission factor representing the emissions occurring at a production or processing plant and one emission factor representing the emissions during use of a solvent containing product. If the chemical is used in more processes and/or is present in several products more emission factors are assigned to the respective chemical amounts.

Emission factors can be defined from surveys of specific industrial activities or as aggregated factors from industrial branches or sectors. Furthermore, emission factors may be characteristic for the use pattern of certain products. The emission factors used in the Danish inventory also rely on the work done in the joint Nordic project (Fauser et al., 2009).

Emission factors for Other Product Use in CRF 3D are compiled from literature searches.

5.2.4 Source allocation

The Danish Working Environment Authority (WEA) is administrating the registrations of chemicals and products to the Danish product register. All manufacturers and importers of products for occupational and commercial use are obliged to register. The following products are comprised in the registration agreement:

- Chemicals and materials that are classified as dangerous according to the regulations set up by the Danish Environmental Protection Agency (EPA).
- Chemicals and materials that are listed with a limit value on the WEA "limit value list".
- Materials, containing 1 % or more of a chemical, which is listed on the WEA "limit value list".
- Materials, containing 1 % or more of a chemical, which are classified as hazardous to humans or the environment according to the EPA rules on classification.

There are the following important exceptions for products, which does not need to be registered:

- Products exclusively for private use.
- Pharmaceuticals ready for use.
- Cosmetic products.

The Danish product register does therefore not comprise a complete account of used pollutants. Source allocations of exceptions from the duty of declaration are done based on information from trade organisations, industries, scientific reports and information from the internet.

Outputs from the inventory are

- a list where the most predominant pollutants are ranked according to emissions to air,
- specification of emissions from industrial sectors and from households,

- contribution from each pollutant to emissions from industrial sectors and households,
- yearly trend in emissions, expressed as total pollutant and single pollutant, and specified in industrial sectors and households.

5.3 Emissions, activity data and emission factors

5.3.1 Solvent Use (NMVOC and CO₂ equivalent emissions)

Table 5.1 and Figure 5.1 show the emissions of NMVOC and CO₂ from 1990 to 2012, where the used amounts of single pollutants have been assigned to specific products and CRF sectors. A general increase is seen for all sectors from 1990 to 1996 followed by a decrease from 1997 to 2006 and stagnation in the period 2007 to 2012. Table 5.2 shows the used amounts of pollutants for the same period. Table 5.1 is derived from Table 5.2 by applying emission factors relevant to individual pollutants and production or use activities. Table 5.2 showing the used amount of products (activity data) is derived by assessing the amount of pollutants that is comprised within products belonging to each of the four source categories. The CO₂ conversion factor for each pollutant is shown in Table 5.3.

In Table 5.3 the emission for 2012 is split into individual pollutants. The most abundantly used solvents are ethanol, turpentine, or white spirit defined as a mixture of stoddard solvent and solvent naphtha and propylalcohol. Ethanol is used as solvent in the chemical industry and as windscreen washing agent. Turpentine is used as thinner for paints, lacquers and adhesives. Propylalcohol is used in cleaning agents in the manufacture of electrical equipment, flux agents for soldering, as solvent and thinner and as windscreen washing agent. Household emissions are dominated by propane and butane, which are used as aerosols in spray cans, primarily in cosmetics. For some pollutants the emission factors are precise but for others they are rough estimates. The division of emission factors into four categories implies that high emission factors are applicable for use of solvent containing products and lower emission factors are applicable for use in industrial use.

The full time series for NMVOC emissions (Table 5.1) and used amount of products (Table 5.2) are presented in Annex 3D; Table 3D-2a, b, c, Table 3D-3 and Table 3D-4, respectively.

Table 5.1 Emission of NMVOC and CO₂-eqv. in Gg pr year.

Total emissions Gg pr year	1990	1995	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
3A Paint Application	5,11	5,90	6,40	5,25	5,18	4,99	4,66	4,32	3,73	3,23	3,38	2,85	2,75	2,87	2,86
3B Degreasing, Dry cleaning and Electronics	7E-05	8E-05	3E-05	1E-05	3E-05	3E-05	2E-05	2E-05	1E-05	2E-05	2E-05	1E-05	1E-05	1E-05	3E-06
3C Chemical Products Manufacturing and Processing	8,14	9,32	6,96	6,28	6,58	4,96	6,06	6,25	6,02	6,12	5,91	4,99	5,05	4,81	4,87
3D Other Use of Solvents and Products	24,7	30,0	27,8	24,8	24,4	22,5	21,4	20,8	20,8	18,0	18,4	19,7	19,4	19,3	19,1
3D3 Other Product Use	0,0831	0,0789	0,0950	0,0855	0,102	0,115	0,102	0,0950	0,109	0,0838	0,0775	0,0804	0,0678	0,0601	0,0817
Total NMVOC	38,0	45,3	41,3	36,4	36,2	32,5	32,3	31,5	30,7	27,5	27,8	27,6	27,3	27,0	27,0
Total CO ₂ -eqv.	93,5	109	100	88,1	88,4	80,3	78,2	75,7	71,7	64,3	65,7	65,1	63,6	63,4	63,3

Table 5.2 Used amounts of products (activity data) in Gg pr year.

Used amounts of products Gg pr year	1990	1995	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
3A Paint Application	83,2	92,1	105	86,4	86,7	79,8	77,7	75,2	64,7	57,3	58,1	48,7	45,8	43,8	43,3
3B Degreasing, Dry cleaning and Electronics	1,41	1,53	0,586	0,251	0,597	0,578	0,481	0,365	0,292	0,433	0,299	0,263	0,247	0,224	0,054
3C Chemical Products Manufacturing and Processing	406	504	567	551	540	513	634	740	749	814	771	683	641	640	516
3D Other Use of Solvents and Products	197	247	230	206	218	185	182	204	180	162	169	179	170	169	169
3D3 Other Product Use	28,0	31,4	46,5	42,0	56,3	61,8	61,4	63,4	63,6	58,5	51,2	52,2	57,7	49,9	53,6
Total products	716	877	949	886	902	841	956	1083	1058	1093	1049	963	914	903	782




Figure 5.1 Emissions of NMVOC in kt NMVOC per year




Figure 5.2 Emissions of CO₂-eq in kt CO₂-eq per year.

Table 5.3 NMVOCs with highest emissions 2012, and CO₂ conversion factors assuming that all carbon molecules in the NMVOC molecule are converted to carbon in CO₂.

Pollutant	CAS no	Emissions 2012 (tonnes)	CO ₂ -conversion factor (g CO ₂ pr g NMVOC)
ethanol	64-17-5	7982	1.91
turpentine (white spirit: stoddard solvent and solvent naphtha)	64742-88-7 8052-41-3	6239	2.79
propyl alcohol	67-63-0	2847	2.20
cyanates	79-10-7	2260	1.83
pentane	109-66-0	1650	3.06
methanol	67-56-1	1069	1.38
acetone	67-64-1	861	2.28
propylen glycol	57-55-6	851	1.74
propane	74-98-6	654	2.86
butane	106-97-8	654	2.93
butanone	78-93-3	348	2.45
xylene	1330-20-7	270	3.32
	95-47-6		
	108-38-3		
	106-42-3		
glycol ethers	110-80-5	251	1.95
	107-98-2		
	108-65-6		
	34590-94-8		
	112-34-5		
	and others		
ethylen glycol	107-21-1	180	1.42
toluene	108-88-3	141	3.35
cyclohexanones	108-94-1	135	2.69
phenol	108-95-2	130	2.81
styrene	100-42-5	100	3.39
butanols	78-92-2	87.5	2.24
	2517-43-3		
	and others		
formaldehyde	50-00-0	72.3	1.47
acyclic aldehydes	78-84-2	29.4	2.31
	111-30-8		
	and others		
ethyl acetate	141-78-6	29.3	2.00
1-butanol	71-36-3	14.7	2.38
butyl acetate	123-86-4	14.0	2.28
tetrachloroethylene	127-18-4	0.45	0.531
acrylic acid	79-10-7	0.043	1.83
Total 2012		26,950	

5.3.2 Other Use of Solvents and Product Use (N₂O, CO₂ and CO₂ equivalent emissions)

3D1 Use of N₂O for Anaesthesia and 3D5 Other Product Use

N₂O is predominantly used as anaesthesia and a small amount is used as fuel in race cars and in chemical laboratories.

Five companies sell N₂O in Denmark and only one company produces N₂O. N₂O is primarily used in anaesthesia by dentists, veterinarians and in hospitals and in minor use as propellant in spray cans, use in laboratories, racing cars and in the production of electronics. Due to confidentiality no data on produced amount are available and thus the emissions related to N₂O production are unknown. An emission factor of 1 is assumed for all uses, which equals the sold amount to the emitted amount. Sold amounts are obtained from the respective companies and the produced amount is estimated from communication with the company.

Total sold and estimated produced N₂O for sale in Denmark, which equals the emissions, is shown in Table 5.4.

Table 5.4 N₂O emissions. EF = 1, i.e. sale equals emissions, and CO₂-eqv. in Gg per year.

Total emissions													
Gg per year	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
3D1	0,0021	0,0078	0,0086	0,0133	0,0191	0,0342	0,0356	0,0407	0,0330	0,0458	0,0344	0,0420	0.0301
Total CO ₂ -eqv.	0.65	2.42	2.66	4.12	5.93	10.6	11.0	12.6	10.2	14.2	10.7	13.0	9.33

Table 5.5 and 5.7 presents the emissions, activity data and emission factors for N₂O and CO₂ from the use of fireworks, tobacco, candles and charcoal, which are included in CRF 3D5 Other. Full time series for emissions and activity data can be found in Annex 3D-5 and 3D-6. Activity data are compiled from Statbank (2013).

Table 5.5 Emission of CO₂ and N₂O from the product use of fireworks, tobacco, candles and charcoal.

		1990	1995	2000	2005	2006	2007	2008	2009	2010	2011	2012
CO ₂ emission from												
Fireworks	Gg	0.1	0.1	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
Candles	Gg	21.7	26.5	49.3	100.2	85.1	93.4	78.1	75.0	102.7	87.8	81.1
Total	Gg	21.7	26.6	49.5	100.4	85.3	93.6	78.3	75.3	102.9	88.0	81.3
N ₂ O emission from												
Fireworks	Mg	2.5	5.8	9.4	7.1	8.1	8.7	8.5	10.4	10.5	9.2	6.7
Tobacco	Mg	0.8	0.7	0.7	0.7	0.7	0.6	0.6	0.6	0.6	0.5	0.5
Candles	Mg	0.2	0.2	0.4	0.8	0.7	0.8	0.6	0.6	0.8	0.7	0.7
Charcoal	Mg	0.2	0.2	0.4	0.4	0.6	0.4	0.3	0.3	0.2	0.2	0.4
Total	Mg	3.7	7.0	10.9	9.1	10.1	10.4	10.0	12.0	12.2	10.6	8.4
Total	Gg CO ₂ -eqv.	22.9	28.8	52.9	103.2	88.4	96.9	81.4	79.0	106.7	91.3	83.8

Table 5.6 Activity data for the product use of fireworks, tobacco, candles and charcoal.

Year		1990	1995	2000	2005	2006	2007	2008	2009	2010	2011	2012
Fireworks	Gg	1.3	3.0	4.9	3.7	4.2	4.5	4.4	5.4	5.4	4.7	3.5
Tobacco	Gg	12.7	11.4	11.4	10.4	10.3	9.8	9.6	9.4	9.2	8.3	8.2
Candles	Gg	7.4	9.1	16.9	34.4	29.2	32.1	26.8	25.8	35.3	30.2	27.9
Charcoal	Gg	7.2	7.9	13.4	14.9	19.8	12.2	10.4	11.6	7.8	6.7	14.0

Emission factors for use of fireworks, tobacco, candles and charcoal are found from literature studies and are shown in Table 5.7.

Table 5.7 Emission factors for other product use, per Mg.

Compound	Unit	Fireworks	Tobacco	Candles	BBQ
CO ₂	Mg	0.043 (a)	NA	2.915 (c)	NA
N ₂ O	kg	1.935 (a)	0.064 (b)	0.024 (d)	0.030 (e)

(a) Netherlands National Water Board (2008), (b) EFs for wood (111A) in residential plants (1A4b i), SNAP 020200, the energy content used in the calculation is the average of wood pills and wood waste (16.1 GJ/Mg), (c) Shires et al. (2004), (d) Shires et al. (2009), (e) IPCC Guidelines (1997),

5.4 Uncertainties and time series consistency

5.4.1 Solvent Use (NMVOC and CO₂ equivalent emissions)

Tier 1 and 2 uncertainties are expressed as \pm 95%-confidence interval limits in percentage relative to the calculated mean emissions for 1990 and 2012, respectively.

Table 5.8 Tier 1 uncertainties for NMVOC and CO₂ equivalents.

Pollutant	Total emission uncertainty, %	Trend 1990-2012, %
NMVOC	23	11
CO ₂ -eqv.	23	10

Table 5.9 Tier 2 uncertainties for NMVOC and CO₂ equivalents.

Pollutant	1990			2012			Trend 1990-2012		
	Median	Low-	Upper	Median	Low-	Upper	Median	Low-	Upper
	Emission (Gg)	er (%)		Emission (Gg)	er (%)		difference (Gg)	er (%)	
NMVOC	38	-13	+16	27	-14	+17	11	-3	+4
CO ₂ -eqv.	94	-14	+16	63	-14	+17	30	-4	+5

Important uncertainty issues related to the mass-balance approach are:

(i) Identification of pollutants that qualify as NMVOCs. Although a tentative list of 650 pollutants from NAEI (2000) has been used, it is possible that relevant pollutants are not included, e.g. pollutants that are not listed with their name in Statistics Denmark (StatBank DK, 2012) but as a product.

(ii) Collection of data for quantifying production, import and export of single pollutants and products where the pollutants are comprised. For some pollutants no data are available in StatBank DK (2012). This can be due to confidentiality or that the amount of pollutants must be derived from products wherein they are comprised. For other pollutants the amount is the sum of the single pollutants *and* product(s) where they are included. The data available in StatBank DK (2012) is obtained from Danish Customs & Tax Authorities and they have not been verified in this assessment.

(iii) Distribution of pollutants on products, activities, sectors and households. The present approach is based on amounts of single pollutants. To differentiate the amounts into industrial sectors it is necessary to identify and quantify the associated products and activities and assign these to the industrial sectors and households. No direct link is available between the amounts of pollutants and products or activities. From the Nordic SPIN database it is possible to make a relative quantification of products and activities used in industry, and combined with estimates and expert judgement these products and activities are differentiated into sectors. The contribution from households is also based on estimates. If the household contribution is set too low, the emission from industrial sectors will be too high and vice versa. This is due to the fact that the total amount of pollutant is constant. A change in distribution of pollutants between industrial sectors and households will, however, affect the total emissions, as different emission factors are applied in industry and households, respectively.

A number of activities are assigned as "other", i.e. activities that cannot be related to the comprised source categories. This assignment is based on expert judgement but it is possible that the assigned amount of pollutants may more correctly be included in other sectors. More detailed information from the industrial sectors is continuously being implemented.

(iv) Rough estimates and assumed emission factors are used for some pollutants. For some pollutants more reliable information has been obtained from

the literature and from communication with industrial sectors. In some cases it is more appropriate to define emission factors for sector specific activities rather than for the individual pollutants. A quantitative measure of the uncertainty has not been assessed. Single values have been used for emission factors and activity distribution ratios etc.

5.4.2 Other Use of Solvents and Product Use (N₂O, CO₂ and CO₂ equivalent emissions)

Tier 1 and 2 uncertainties for CO₂, N₂O and their respective CO₂ equivalents are shown in Table 5.10 and 5.12, respectively.

Table 5.10 Tier 1 uncertainties for CO₂, N₂O and CO₂ equivalents.

Pollutant	Total emission uncertainty, %	Trend 1990-2012, %	Uncertainty trend %age points
CO ₂	100	324	48
N ₂ O	93	230	79
GHG (CO ₂ eqv.)	88	234	73

Table 5.11 Tier 2 uncertainties for CO₂, N₂O and CO₂ equivalents.

Pollutant	1990			2012			Trend 1990-2012		
	Median			Median			Median		
	Emission	Lower	Upper	Emission	Lower	Upper	difference	Lower	Upper
	(Gg)	(%)	(%)	(Gg)	(%)	(%)	(Gg)	(%)	(%)
CO ₂	22	-20	+24	81	-20	+24	-60	-67	+55
N ₂ O	4	-42	+102	19	-22	+55	-15	-180	+73
GHG (CO ₂ eqv.)	23	-18	+24	93	-17	+22	-70	-65	+50

The main issue leading to uncertainties is:

- Collection of data for quantifying production, import and export of products. Some data, like private import (cross-border shopping) of tobacco, are not available in StatBank DK (2012).

5.5 Quality assurance/quality control (QA/QC) and verification

Table 5.12 External and internal data for NMVOC emission inventory.

File or folder name	Description	AD or Emf.	Reference	Contact(s)	Data agreement/Comment
"Emissioner NMVOC" folder	Production, import and export data from Statistics Denmark	Activity data	Statistics Denmark		Statistics Denmark are obligated by law to publish the foreign trade statistics and production statistics
NMVOC emissions.xls	Calculations, emission factors, SPIN data. For industrial branches (NACE)	Activity data and emission factors	Statistics Denmark, SPIN, reports, personal communication		
Use Category National.xls	Calculations, emission factors, SPIN data (UCN and NACE) and use amounts from Statbank.	Activity data and emission factors	Statistics Denmark, SPIN, reports, personal communication		
Emission factors solvent use.xls	Emission factors for chemicals in CRF and SNAP sub-categories. CO ₂ conversion factors.	Emission factors and CO ₂ conversion factors	Scientific reports, personal communication and expert judgement		

The QA/QC procedure is outlined in Section 1.6. In general, Critical Control Points (CCP) has been defined as elements or actions, which need to be addressed in order to fulfil the quality objectives. The CCPs have to be based on clear measurable factors, expressed through a number of Points for Measuring (PM).

Data Storage level 1	1. Accuracy	DS.1.1.1	General level of uncertainty for every data set including the reasoning for the specific values
----------------------	-------------	----------	---

The sources of data described in the methodology section and in DS.1.2.1 and DS.1.3.1 are used in this inventory. It is the accuracy of these data that define the uncertainty of the inventory calculations. Any data value obtained from StatBank DK (2012) and SPIN are given as a single point estimate and no probability range or uncertainty is associated with this value. Information from reports is sometimes given in ranges. Uncertainties are therefore assessed from expert judgement and guidebook estimates.

Data Storage level 1	2. Comparability	DS.1.2.1	Comparability of the emission factors/calculation parameters with data from international guidelines, and evaluation of major discrepancies.
----------------------	------------------	----------	--

1) Production and import/export data from StatBank DK (2012) for single chemicals can be directly compared with data from Eurostat (2012) for other countries. This has been done for a few chosen chemicals and countries. Furthermore, chosen Danish data from Eurostat (2012) have been validated with

data from StatBank DK (2012) in order to check the consistency in data transfer from national to international databases.

2) Use categories for chemicals in products are found from the Nordic SPIN database. Data for all Nordic countries are available and reported uniformly. For chosen chemicals a comparison of chemical amounts and use has been made between countries.

3) A joint Nordic project funded by the Nordic Council of Ministers has been used on methodological issues and for emission factors.

Data Storage level 1	3.Completeness	DS.1.3.1	Ensuring that the best possible national data for all sources are included, by setting down the reasoning behind the selection of datasets.
----------------------	----------------	----------	---

A number of external data sources form the basis for calculating emissions of single chemicals. The general methodology in the emission inventory is described above.

1) StatBank DK (2012) is used as the main database for collecting data on production, import and export of single chemicals, chemical groups and for some products. In order to obtain a uniform and unique set of data it is important that the data for e.g. production of single chemicals is in the same reporting format and from the same source. The amount of data is very comprehensive and is linked with the data present in Eurostat. The database covers all sectors and is regarded as complete on a national level.

2) Nordic SPIN database provides data on the use of chemicals in Norway, Sweden, Denmark and Finland. It is financed by the Nordic Council of Ministers, Chemical group, and the data is supplied by the product registries of the contributing countries. The Danish product register (PROBAS) is a joint register for the WEA and the EPA and comprises a large number of chemicals and products. The information is obtained from registration according to the EPA rules and from scientific studies and surveys and other relevant sources. The product register is the most comprehensive collection of chemical data in products for Denmark and with the availability of data from the other Nordic countries it enables an inter-country comparison. For each chemical the data is reported in a uniform way, which enhances comparability, transparency and consistency.

3) Reports from and personal contacts with industrial branches. It is fundamental to have information from the industrial branches that have direct contact with the activities, i.e. chemicals and products of interest. The information can be in the form of personal communication, but also reported surveys are of great importance. In contrast to the more generic approach of collecting information from large databases, the expert information from industrial branches may give valuable information on specific chemicals and/or products and industrial activities. By considering both sources a verification as well as optimum reliability and accuracy is obtained.

4) The present inventory procedure builds partly on information from the previous Danish solvent emission inventory, which is based on questionnaires to industrial branches. Furthermore a joint Nordic collaboration on solvent inventories has given important information on methods and data.

Data Storage level 1	4.Consistency	DS.1.4.1	The original external data has to be archived with proper reference.
----------------------	---------------	----------	--

Data are predominantly extracted from the internet (StatBank 2012 and SPIN). These are saved as original copies in their original form. Specific information from industries and experts are saved as e-mails and reports.

Data Storage level 1	6.Robustness	DS.1.6.1	Explicit agreements between the external institution holding the data and NERI about the conditions of delivery
----------------------	--------------	----------	---

As stated in DS.1.4.1 most data are obtained from the internet. No explicit agreements have been made with external institutions.

Data Storage level 1	7.Transparency	DS.1.7.1	Listing of all archived datasets and external contacts.
----------------------	----------------	----------	---

Datasets are archived as stated in Table 5.5. External contacts are stored in e-mail and documents.

Data Processing level 1	1. Accuracy	DP.1.1.1	Uncertainty assessment for every data source not part of DS.1.1.1 as input to Data Storage level 2 in relation to type and scale of variability.
-------------------------	-------------	----------	--

No data are used in addition to those included in DS.1.1.1

Data Processing level 1	2.Comparability	DP.1.2.1	The methodologies have to follow the international guidelines suggested by UNFCCC and IPCC.
-------------------------	-----------------	----------	---

The methodological approach is based on the detailed methodology as outlined in the Emission Inventory Guidebook. See also DS.1.3.1.

Data Processing level 1	3.Completeness	DP.1.3.1	Identification of data gaps with regard to data sources that could improve quantitative knowledge.
-------------------------	----------------	----------	--

In "Uncertainties and time series consistency" important uncertainty issues related to missing quantitative knowledge is stated. To summarise; (i) identification and inclusion of all relevant chemicals (and products) Identification of chemicals that qualify as NMVOCs. The definition in the solvent directive (Directive 1999/13/EC) is used. Here VOCs are defined as follows: "Volatile organic compound shall mean any organic compound having at 293,15 K a vapour pressure of 0,01 kPa or more, or having a corresponding volatility under the particular condition of use". A tentative list of 650 chemicals from the "National Atmospheric Emission Inventory" (NAI 2000) has been used, it is possible that relevant chemicals are not included. (ii) Collection of data for quantifying production, import and export of single chemicals. For some chemicals no data are available in StatBank DK (2012). This can be due to confidentiality or that the amount of chemicals must be derived from products wherein they are comprised. (iii) Distribution of chemicals on products, activities, sectors and households. No direct link is available between the amounts of chemicals and products or activities. From the Nordic SPIN database it is possible to make a relative quantification of products and activities used in industry, and combined with estimates and expert judgement

these products and activities are differentiated into sectors. More detailed information from the industrial sectors may still be required. (iv) Emission factors for single chemicals, products and industrial and household activities. For many industrial and household activities involving solvent containing products no estimates on emission factors are available. Large variations occur between industry and product groups. And given the large number of chemicals more specific knowledge regarding industrial processes and consumption is needed.

Data Processing level 1	4.Consistency	DP.1.4.1	Documentation and reasoning of methodological changes during the time series and the qualitative assessment of the impact on time series consistency.
-------------------------	---------------	----------	---

Any changes in calculation procedures are noted for each year's inventory.

Data Processing level 1	5.Correctness	DP.1.5.1	Verification of calculation results using time series
-------------------------	---------------	----------	---

No detailed guidelines or calculations are accessible for time series. These are therefore not used for verification.

Data Processing level 1	5.Correctness	DP.1.5.2	Verification of calculation results using other measures
-------------------------	---------------	----------	--

Calculations performed by IIASA using RAINS codes, which are based on a different methodological approach gives total emission values that are similar to the emissions found in the present approach.

Data Processing level 1	7.Transparency	DP.1.7.1	The calculation principle, the equations used and the assumptions made must be described.
-------------------------	----------------	----------	---

See methodological approach.

Data Processing level 1	7.Transparency	DP.1.7.2	Clear reference to dataset at Data Storage level 1
-------------------------	----------------	----------	--

See methodological approach.

Data Processing level 1	7.Transparency	DP.1.7.3	A manual log to collect information about recalculations.
-------------------------	----------------	----------	---

This is stated in documents listed in Table 5.5.

Data Storage level 2	5.Correctness	DS.2.5.1	Check if a correct data import to level 2 has been made
----------------------	---------------	----------	---

The transfer of emission data from level 1, storage and processing, to data storage level 2 is manually checked.

Data Storage level 4	4.Consistency	DS.4.4.3	The IEFs from the CRF are checked both regarding level and trend. The level is compared to relevant emission factors to ensure correctness. Large dips/jumps in the time series are explained.
----------------------	---------------	----------	--

See DP.1.5.1 and DP.1.5.2.

5.6 Recalculations

Improvements and additions are continuously being implemented due to the comprehensiveness and complexity of the use and application of solvents in industries and households. The main improvements in the 2012 reporting include the following:

- Minor changes in the activity data for tobacco, fireworks, barbeques and candles due to changes updated values from StatBank DK (2013).

5.7 Planned improvements

- Emissions of N₂O used as propellant and solvent in canned whipped cream will be estimated. It has not been possible to get access to sales data and/or use data of number of cans from manufacturers and distributors. If these activity data cannot be compiled an expert estimate will be made.

5.8 References

Directive 1999/13/EC of 11 March 1999 on the limitation of emissions of volatile organic compounds due to the use of organic solvents in certain activities and installations, Brüssel, 1999.

Directive 2001/81/EC Of The European Parliament And Of The Council of 23 October 2001 - on national emission ceilings for certain atmospheric pollutants. Official Journal of the European Communities.

Directive 2004/42/EC of the European Parliament and Council on the Limitation of Emissions of Volatile Organic Compounds due to the Use of Organic Solvents in Certain Paints and Varnishes and Vehicle Refinishing Products and Amending Directive 1999/13/EC 30.04.2004, L 143/87.

EMEP/EEA, 2009. Air pollutant emission inventory guidebook (formerly referred to as the EMEP CORINAIR emission inventory guidebook): Published by the EEA, with technical guidance by UNECE's Task Force on Emission Inventories and Projections. Available at:

<http://www.eea.europa.eu/publications/emep-eea-emission-inventory-guidebook-2009> (03-01-2013).

Eurostat on the internet. Available at:

http://epp.eurostat.ec.europa.eu/portal/page/portal/statistics/search_database

Fausser, P., Saarinen, K., Harðardóttir, K., Kittilsen, M.O., Holmengen, N. & Skårman, T. 2009: Improvement of Nordic Emission Models for Solvent Use in Selected Sectors. TemaNord 2009:556. Nordic Council Of Ministers, Copenhagen.

Fyrværkeri og miljø, Fyrværkeribrancheforeningen, Danish Pyrotechnical Association, December 8 2009. (Danish).

IPCC, 1997. Revised 1996 IPCC Guidelines for National Greenhouse Gas Inventories. Available at:

<http://www.ipcc-nggip.iges.or.jp/public/gl/invs1.html> (03-01-2013).

IPCC, 2000: Good Practice Guidance and Uncertainty Management in National Greenhouse Gas Inventories. Available at:

<http://www.ipcc-nggip.iges.or.jp/public/gp/english/> (03-01-2013).

Karjalainen, T. 2005: Commission research in Action: tackling the hormone disrupting chemicals issue, EUR report 21941.

Myatt, S., Health and Safety Laboratory, United Kingdom, CHAF Workpackage 4 Report, Deliverable D4-4, The Effect of Fireworks on Health and the Environment, Literature review, June 2003, Workpackage leader: A. von Oertzen, Bundesanstalt für Materialforschung und -prüfung, Germany. Available at:

<http://www.pyrobin.com/files/european%20firework%20testing%20reports,%20bam%20etc.pdf> (21-02-2013).

NAEI, 2007: The British National Atmospheric Emission Inventory. Available at: <http://www.naei.org.uk/emissions/selection.php> (21-02-2013).

NAEI, 2000: British National Atmospheric Inventory performed by the National Environmental Technology Centre, June 2000. Available at:

http://www.aeat.co.uk/netcen/airqual/naei/annreport/annrep99/app1_28.html (05-05-2006).

Netherlands National Water Board, 2008, Emission estimates for diffuse sources, Netherlands Emission Inventory, Letting off fireworks, Version dated June 2008, Netherlands National Water Board – Water Unit in cooperation with Deltares and TNO. Available at:

<http://www.emissieregistratie.nl/ERPUBLIEK/documenten/Water/Factsheets/English/Fireworks.pdf> (21-02-2013)

Nielsen, O.-K., Lyck, E., Mikkelsen, M.H., Hoffmann, L., Gyldenkerne, S., Winther, M., Nielsen, M., Fauser, P., Thomsen, M., Albrechtsen, R., Hjelgaard, K., Vesterdal, L., Møller, I.S. & Baunbæk, L. 2009: Denmark's National Inventory Report 2009. Emission Inventories 1990-2007 - Submitted under the United Nations Framework Convention on Climate Change. National Environmental Research Institute, Aarhus University, 2009. 826 s. (NERI Technical Report; 724). IPCC, 1997. Revised 1996 IPCC Guidelines for National Greenhouse Gas Inventories. Available at

<http://www.ipcc-nggip.iges.or.jp/public/g-l/invs6.htm> (15-04-2007).

Pärt, P. (main author) 2005: Environment and health, EEA Report No 10/2005, Copenhagen.

Report Seest: Rapport vedrørende fyrværkeriulykken i Seest den 3. november 2004, Udarbejdet af en uafhængig ekspertgruppe nedsat den 26. august 2005. Prepared by the Ministry of Economic and Business Affairs Denmark and the Ministry of Defence Denmark.

Shires, T. M., Loughran, C. J., American Petroleum Institute (API), Compendium of Greenhouse Gas Emissions Methodologies for the Oil and Gas Industry, February 2004. Available at:
<http://www.scribd.com/doc/73896273/2004-COMPENDIUM-of-Greenhouse-Gas-Emissions-Estimates> (21-02-2013).

Shires, T. M., Loughran, C. J., Jones, S., Hopkins, E., American Petroleum Institute (API), Compendium of Greenhouse Gas Emissions Methodologies for the Oil and Gas Industry, August 2009. Available at:
http://www.api.org/ehs/climate/new/upload/2009_ghg_compendium.pdf (21-02-2013).

SPIN on the Internet. Substances in Preparations in Nordic Countries. Available at: <http://www.spin2000.net/spin.html>

StatBank, 2012. Statistics Denmark, 2012: Statistics Denmark, StatBank Denmark. (Danish/English). Available at:
<http://www.statistikbanken.dk/statbank5a/default.asp?w=1024>

The Danish Safety Technology Authority, 20/07-2010.

UNFCCC, 2008: Available at:
http://unfccc.int/national_reports/annex_i_ghg_inventories/national_inventories_submissions/items/3929.php

6 Agriculture (CRF sector 4)

The data presented in Chapter 6 relates to Denmark only, whereas information for Greenland is included in Chapter 16 and for the Faroe Islands in Annex 8.

The emission of greenhouse gases from agricultural activities includes:

- CH₄ emission from enteric fermentation and manure management.
- N₂O emission from manure management and agricultural soils.
- NMVOC emission from agricultural soils.
- Emission of CH₄, N₂O, NMVOC, CO and NO_x from burning of straw on field.

Emissions from rice production and burning of savannahs do not occur in Denmark and consequently these categories have been reported as Not Occurring.

6.1 Overview of sector

In CO₂ equivalents, the agricultural sector contributes with 19 % of the overall greenhouse gas emission (GHG) in 2012 excl. LULUCF. Next to the energy sector, the agricultural sector is the largest source of GHG emission in Denmark. The majority of agricultural greenhouse gas emissions are covered by N₂O and CH₄, which contributes in 2012 with 90 % and 77 % respectively of the total Danish emissions of N₂O and CH₄.

From 1990 to 2012, the emissions decreased from 12.5 million tonnes CO₂ equivalent to 9.6 million tonnes CO₂ equivalent, which corresponds to a 23 % reduction (Table 6.1). N₂O is the largest contributor to the overall agricultural greenhouse gas emission, in 2012 accounting for 56 % in CO₂ equivalents. The decrease in the agricultural emission is caused by a decrease in N₂O emission, while the CH₄ emission is nearly unaltered.

Table 6.1 Emission of GHG in the agricultural sector in Denmark 1990 – 2012

	1990	1995	2000	2005	2008	2009	2010	2011	2012
CH ₄ , Gg CO ₂ eqv.	4 234	4 227	4 036	4 043	4 106	4 095	4 165	4 151	4 203
N ₂ O, Gg CO ₂ eqv.	8 292	7 344	6 417	5 804	5 832	5 499	5 445	5 517	5 396
Total, Gg CO ₂ eqv.	12 526	11 571	10 453	9 847	9 938	9 594	9 609	9 668	9 599

The major part of the emission is related to livestock production, which in Denmark is dominated by the production of cattle and swine.

Figure 6.1 shows the distribution of the greenhouse gas emission across the main agricultural sources. The total N₂O emission from 1990-2012 has decreased by 35 % and can largely be attributed to the decrease in N₂O emissions from agricultural soils. This reduction is due to a proactive national environmental policy over the last twenty five years to prevent loss of nitrogen from agricultural soil to the aquatic environment. These measures includes among other things a ban on manure application during autumn and winter, strict requirements to storage and application of manure, increasing area with winter-green fields to catch nitrogen, a maximum number of animals per hectare (ha) and maximum nitrogen application rates for agricul-

tural crops. A combination of these increasing environmental requirements and the efforts to obtain economic advantage, the farmers has been forced to improve the utilisation of nitrogen in manure. An improvement of feed efficiency has been one of the most important drivers to reach the objectives. This has led to a halving of nitrogen use in synthetic fertiliser and a decrease of emission per produced kg meat, which all has reduced the overall GHG emission.

The CH₄ emissions from 1990 to 2012 shown in Figure 6.1 indicate a decrease in emission from enteric fermentation, which is mainly due to a decrease in the number of cattle. A contrasting development has taken place in emission from manure management. Structural changes in the sector have led to a move towards the use of slurry-based housing systems, which have a higher emission factor than systems with solid manure. By coincidence, the decrease and the increase almost balance each other out and the total CH₄ emission from 1990 to 2012 has decreased by 1 %.




Figure 6.1 Danish greenhouse gas emissions 1990 – 2012.

6.1.1 Key category identification

The key category analysis (KCA) divides the agriculture emissions into 14 subcategories, refer Annex 1. In Table 6.2 is listed KCA covering Tier 1 and Tier 2. Tier 1 only gives key source identification based on the quantitative emission, while the Tier 2 analyse also include information on uncertainties estimates (refer to Chapter 1.5). In 1990, 9 of the 14 agricultural sources are registered as key categories and 11 sources are key categories if uncertainties are taken into account (Tier 2). In 2012, 5 of the sources are listed as key categories according to level and trend for Tier 1 and 8 sources in Tier 2. For the methodological choice Denmark uses the key categories identified using both tier 1 and tier 2 for the latest year as well as key categories identified for the trend from 1990 to the latest year.

The three most important agriculture key categories are CH₄ from enteric fermentation and N₂O emissions from nitrogen leaching and run-off and synthetic fertilisers.

Table 6.2 Key category identification Tier 1 and Tier 2 from the agricultural sector 1990 and 2012.

CRF table	Compounds	Emission source	Key category identification	
2012			Tier 1	Tier 2
4.A	CH ₄	Enteric fermentation	Level/trend	Level/trend
4.B(a)	CH ₄	Manure management	Level/trend	Level/trend
4.F	CH ₄	Field burning of agri. residues	-	-
4.B(a)	N ₂ O	Manure management	Level	Level
4.D1.1	N ₂ O	Synthetic fertilisers	Level/trend	Level/trend
4.D1.2	N ₂ O	Animal manure applied to soils	Level/trend	Level/trend
4.D1.3	N ₂ O	N-fixing crops	Level	Level/trend
4.D1.4	N ₂ O	Crop residue	Level	Level/trend
4.D1.5	N ₂ O	Cultivation of histosols	Level	Level
4.D1.6	N ₂ O	Sewage sludge and industrial waste	-	-
4.D2	N ₂ O	Pasture, range and paddock	Level	Level
4.D3.1	N ₂ O	Atmospheric deposition	Level	Level/trend
4.D3.2	N ₂ O	Nitrogen leaching and run-off	Level/trend	Level/trend
4.F	N ₂ O	Field burning of agri. residues	-	-
1990				
4.A	CH ₄	Enteric fermentation	Level	Level
4.B(a)	CH ₄	Manure management	Level	Level
4.F	CH ₄	Field burning of agri. residues	-	-
4.B(b)	N ₂ O	Manure management	Level	Level
4.D1.1	N ₂ O	Synthetic fertilisers	Level	Level
4.D1.2	N ₂ O	Animal manure applied to soils	Level	Level
4.D1.3	N ₂ O	N-fixing crops	-	Level
4.D1.4	N ₂ O	Crop residue	Level	Level
4.D1.5	N ₂ O	Cultivation of histosols	-	Level
4.D1.6	N ₂ O	Sewage sludge and industrial waste	-	-
4.D2	N ₂ O	Pasture, range and paddock	Level	Level
4.D3.1	N ₂ O	Atmospheric deposition	Level	Level
4.D3.2	N ₂ O	Nitrogen leaching and run-off	Level	Level
4.F	N ₂ O	Field burning of agri. residues	-	-

6.2 Data references

The calculated emissions are based on methods described in the IPCC Reference Manual (IPCC, 1997) and the IPCC Good Practice Guidance (IPCC, 2000).

Activity data and emission factors are collected and discussed in cooperation with specialists and researchers in various institutes with agricultural expertise, such as the DCA - Danish Centre for Food and Agriculture - Aarhus University, Statistics Denmark, the Danish Agricultural Advisory Service, the Danish AgriFish Agency and the Danish Environmental Protection Agency. In this way, both data and methods will be evaluated continually, according to the latest knowledge and information. DCE - Danish Centre for Environment and Energy, Aarhus University has established data agreements with the institutes and organisations to assure that the necessary data are available to prepare the emission inventory on time.

Table 6.3 List of institutes involved in the emission inventory for the agricultural sector.

References	Link	Abbreviation	Data/information
Statistics Denmark – Agricultural Statistics	www.dst.dk	DSt	<ul style="list-style-type: none"> - livestock production - milk yield - slaughtering data - export of live animal - poultry - land use - crop production - crop yield
Danish Centre for Food and Agriculture, Aarhus University		DCA	<ul style="list-style-type: none"> - N-excretion - feeding situation - animal growth - use of straw for bedding - N-content in crops - modelling of data regarding N-leaching/runoff - NH₃ emissions factor
The Danish Agricultural Advisory Service	http://www.vfl.dk	DAAS	<ul style="list-style-type: none"> - housing type (until 2004) - grazing situation - manure application time and methods - estimation of extent of field burning of agricultural residue
Danish Environmental Protection Agency	www.mst.dk	EPA	<ul style="list-style-type: none"> - sewage sludge used as fertiliser (until 2004) - industrial waste used as fertiliser
The Danish AgriFish Agency	http://naturerhverv.fvm.dk	DAFA	<ul style="list-style-type: none"> - synthetic fertiliser (consumption and type) - housing type (from 2005) - sewage sludge used as fertiliser (from 2005 based on the register for fertilization) - number of animals from the Central Husbandry Register
The Danish Energy Agency	www.ens.dk	DEA	<ul style="list-style-type: none"> - manure used in biogas plants

The emissions from the agricultural sector are calculated in a comprehensive agricultural model complex called IDA (Integrated Database model for Agricultural emissions). The model complex is designed in a relational database system (MS Access). Input data are stored in tables in one database called IDA_Backend and the calculations are carried out as queries in another linked database called IDA. This model complex, as shown in Figure 6.2, is implemented in great detail and is used to cover emissions of air pollutants and greenhouse gases. Thus, there is a direct coherence between the NH₃ emission and the emission of N₂O.

IDA - Integrated Database model for Agricultural emissions




Figure 6.2 IDA - Integrated Database model for Agricultural emissions

Most emissions relate to livestock production, which basically is based on information on the number of animals, the distribution of animals according to housing type and, finally, information on feed consumption and excretion.

IDA operates with 38 different livestock categories, according to livestock type, weight class and age. These categories are subdivided into housing type and manure type, which results in 252 different combinations of livestock subcategories and housing types (see Annex 3E-1). For each of these combinations, information on e.g. feed intake, digestibility, excretion and grazing days is included. The emission is calculated from each of these subcategories and then aggregated in accordance with the IPCC livestock categories given in the CRF.

Table 6.4 Livestock categories and subcategories.

CRF	Aggregated livestock categories as given in IPCC	Includes	No. of subcategories in IDA, animal type/housing system
4B 1a	Dairy Cattle ¹	Dairy Cattle	34
4B 1b	Non-dairy Cattle ¹	Calves (<½ yr), heifers, bulls, suckling cattle	120
4B 3	Sheep	Including lambs	1
4B 4	Goats	Including kids (meet, dairy and mohair)	3
4B 6	Horses	<300 kg, 300 - 500 kg, 500 - 700 kg, >700 kg	4
4B 8	Swine	Sows, weaners, fattening pigs	36
4B 9	Poultry	Hens, pullets, broilers, turkeys, geese, ducks	42
4B 13	Other	Fur animals ² , deer, ostriches, pheasant	12

¹⁾ For all subcategories, large breed and jersey cattle are distinguished from each other.

²⁾ Fur farming: Mink and foxes

It is important to point out that changes over the years, both to the national emission and the implied emission factor, are not only a result of changes in the numbers of animals, but also depend on changes in the allocation of sub-categories, changes in feed consumption and changes in housing type.

6.2.1 Number of animals

Livestock production is primarily based on the agricultural census from Statistics Denmark (DSt). The emission from bulls, fattening pigs and poultry is based on slaughter data. A certain numbers of horses, goats and sheep on small farms are added to the number in DSt because Statistics Denmark does not include farms less than 5 ha, where many of these animals are placed. Statistics Denmark is the source for the database kept by FAO (Food and Agriculture Organization of the United Nations) and Eurostat. This explains why the number of sheep, goats and horses in FAO and the Danish emission inventory differ. The largest difference is found for horses. In the agricultural census for 2012, the number of horses is estimated to be around 68 000. Including horses on small farms and riding schools, however, the number of horses rises to approximately 155 000 (Clausen, 2012). Data on the number of sheep and goats are based on the Central Husbandry Register (CHR), which is the central register of farms and animals administrated by the Ministry of Food, Agriculture and Fisheries.

Information on the number of deer, ostriches and pheasants are not included in Statistics Denmark. While the number of deer and ostriches are based on the CHR, the number of pheasants is based on expert judgement from DCE and the pheasant breeding association.

In Annex 3E-2 is provided number of animals allocated on all livestock sub-categories for selected years 1990-2011, for all years see <http://envs.au.dk/videnudveksling/luft/emissioner/reportingsectors/agriculture/>.

6.2.2 Housing type

From 2005, all farmers have to report to the Danish AgriFish Agency (DAFA) information concerning the use of housing type. Annex 3E-1 shows the housing type for each livestock category for selected years 1990 – 2011.

Before 2005 there exist no official statistics which cover the distribution of animals according to housing type. The distribution is, therefore, based on

an expert judgement from the Danish Agricultural Advisory Service (DAAS) and DCA. Approximately 90-95 % of Danish farmers are members of DAAS, which regularly collects statistical data from the farmers on different issues, as well as making recommendations with regard to farm buildings. Hence, DAAS has a good understanding of which housing types that are currently in use and also the changes over time.

6.2.3 Feed consumption and excretion

The DCA provide Danish standards related to feed consumption, excreted volumes, nutrient content of nitrogen, phosphor and potassium, dry matter in manure and contribution of different manure type. These standards are all a part of the “Danish Normative System”, which is used for fertiliser planning and control by the Danish farmers and authorities (Poulsen et al., 2001, Poulsen et al., 2013). The complexity and dynamics of the system has increased during the years to secure the development of accurate values. Furthermore, the normative system includes emission factors for NH_3 , which is based on a combination of measurements and model calculations. Emission factors for NH_3 from the housing unit and storage are given in Annex 3E-3 and 4.

The Danish normative standards are based on practical farming and thus reflect the actual Danish agricultural production conditions. DCA receive data from the Danish Agricultural Advisory Service (DAAS), which is the central office for all Danish agricultural advisory services. DAAS carries out a considerable amount of research itself, as well as collecting efficacy reports from the Danish farmers for dairy production, meat production, pig production, etc., to optimise productivity in Danish agriculture. Feeding plans are used to provide values to the Danish Normative System and for dairy cows the values are based on approximately 800 feeding plans. In total the normative standards covers feed plans from 15-18 % of the Danish dairy production, 25-30 % of the pig production, 80-90 % of the poultry production and approximately 100 % of the fur production. A high fraction of the pig production is represented, which is caused by the intensive focus on the possibilities to optimize the feed intake to increase the feed efficiency. The values covering the cattle production can be considered as reliable, even though only 15-18 % of the productions are represented. These values include mainly feeding plans from the farmers with a production efficiency corresponding to a middle level. The farmers with a high productivity level are often not users of the Danish Agricultural Advisory Service, which also is the case for farmers with a low productivity level.

Previously, the normative standards were updated and published every third or fourth year (Laursen, 1987; Laursen, 1994; Poulsen and Kristensen, 1997). From 2001 these standards are updated annually and available to download at the homepage of DCA:

<http://anis.au.dk/forskning/sektioner/husdyrernaering-og-miljoe/normtal/> (17.01.2014).

One of the reports concerning the normative data is published in English in Poulsen and Kristensen (1998) and is available at the homepage of DCA, see list of references. The normative data is adjusted over time but the methodology is the same.

6.3 CH₄ emission from enteric fermentation (CRF sector 4A)

6.3.1 Description

The major part of the agricultural CH₄ emission originates from digestive processes. In 2012, this source accounts for 30 % of the total GHG emission from agriculture. The emission is primarily related to ruminants and, in Denmark, particularly to cattle, which, in 2012, contributed with 86 % of the emission from enteric fermentation. The emission from pig production is the second largest source and covers 10 % of the emission from enteric fermentation, followed by horses (3 %) and sheep, goats, deer and poultry (1 %).

6.3.2 Methodological issues

The methodology for estimating emissions from enteric fermentation is based on the Revised 1996 IPCC Guidelines (IPCC, 1997) and the IPCC Good Practice Guidance (IPCC, 2000). The methodology for poultry, ostrich and pheasants is based on Tier 1, while the remaining animal categories are based on a Tier 2/Country Specific (CS) approach. CH₄ emission from enteric fermentation from fur farming is considered to be not applicable based on country-specific information (Hansen, 2010). Feed consumption for all animal categories is based on the Danish normative figures. Default values for the methane conversion rate (Y_m) given by the IPCC are used for all livestock categories, except for dairy cattle and heifers, where a national Y_m is used for all years. In the Danish inventory sheep and goats includes lamb and kids - an average of Y_m values for mother animal and lamb/kid is used.

Tier 1

Emission factors used for poultry, ostrich and pheasants are based on the emission factors given by Wang & Huang (2005) (see Table 6.5). EF for broilers with a life cycle of 30-56 days is scaled in proportion to 42 days for broilers given by Wang & Huang (2005). Organic broilers with a life cycle of 81 days are scaled in proportion to the Taiwan country chicken with 91 days of life cycle and pullets with a life cycle of 112-119 days are scaled in proportion to the 140 days given for pullets by Wang & Huang (2005). EF for ducks, geese, turkeys, ostrich chickens and pheasant chickens is scaled by weight in proportion to a Danish broiler with 40 days of life cycle. For laying hens, the EF for laying hens given by Wang & Huang (2005) is used and for ostrich hens and pheasant hens the EF is scaled by weight in proportion to a laying hen.

Table 6.5 EF for poultry in mg CH₄ per head per lifecycle.

	CH ₄ emission factor
Broilers, 42 days	15.87
Taiwan country chicken, 91 days	84.82
Pullets, 140 days	3 561
Laying hens, 365 days	10 610

Source: Wang & Huang, 2005.

Tier 2

The Tier 2/CS equation for EF of enteric fermentation is the sum of the feeding situation in winter and summer. The EF is based on actual feeding plans, which is provided from data for feed units (FU) for each livestock category – see below. Feeding with sugar beets is taken into account because sugar beet feeding gives a higher methane production rate compared to grass and maize due to the high content of easily convertible sugar. However, it is only

dairy cattle and heifers which have sugar beets in the feed. The parts of the equation concerning sugar beet will be left out for the remaining animal categories.

$$EF = EF_{\text{winter}} + EF_{\text{summer}}$$

$$EF_{\text{winter}} = FU \cdot \left(\left(\frac{GE_{\text{FU winter}}}{55.65} \right) \cdot Y_{\text{mexcl sugarbeet}} \cdot \left(1 - \frac{\text{grazing days}}{365} - \frac{\text{days with sugar beet}}{365} \right) + \left(\frac{GE_{\text{FU winter}}}{55.65} \right) \cdot Y_{\text{mincl sugarbeet}} \cdot \frac{\text{days with sugar beet}}{365} \right)$$

$$EF_{\text{summer}} = FU \cdot \left(\frac{GE_{\text{FU summer}}}{55.65} \right) \cdot Y_{\text{mgazing}} \cdot \frac{\text{grazing days}}{365}$$

Where:

FU = feeding units

$GE_{\text{FU, winter}}$ = gross energy per feeding unit, MJ per FU in winter

$GE_{\text{FU, summer}}$ = gross energy per feeding unit, MJ per FU in summer

Y_m = methane conversion factor, per cent of gross energy in feed converted to methane

Thus, to calculate the total gross energy (GE) intake, the GE per feed unit – defined as GE_{FU} – needs to be estimated. A feed unit in Denmark is defined as the feed value in 1.00 kg barley with a dry matter content of 85 % (Statistics Denmark, yearbook 2010). For other cereals e.g. wheat and rye one feed unit is 0.97 kg and 1.05 kg, respectively.

Gross energy intake

The calculation of $GE_{\text{FU, winter}}$ and $GE_{\text{FU, summer}}$ is based on the composition of feed intake and the energy content in proteins, fats and carbohydrates based on actual efficacy feeding controls or actual feeding plans at farm level, collected by DAAS or DCA. The data are given in Danish feed units or kg feedstuff and these values are converted to mega joule (MJ). The calculation is shown in the equation below:

The principle for estimation of $GE_{\text{FU, winter}}$ and $GE_{\text{FU, summer}}$ is the same, why the following equation only is defined as GE_{FU} .

$$GE_{\text{FU}} = \frac{\text{MJ/day}}{\text{FU/day}}$$

$$\text{FU/day} = \frac{\text{kg dm}}{\text{day}} \cdot \frac{\text{FU}}{\text{kg dm}}$$

$$\text{MJ/day} = \frac{\text{kg dm}}{\text{day}} \cdot \frac{\text{MJ}}{\text{kg dm}}$$

$$\text{MJ/kg dm} = \%_{\text{Crudeprotein}} \cdot E_{\text{Crudeprotein}} + \%_{\text{Raw fat}} \cdot E_{\text{Raw fat}} + \%_{\text{Carbohydrates}} \cdot E_{\text{Carbohydrates}}$$

$$\%_{\text{Carbohydrates}} = 100 - (\%_{\text{Crudeprotein}} + \%_{\text{Raw fat}} + \%_{\text{Raw ashes}})$$

In Annex 3E-5 and 6 are listed all parameters for winter feeding plans covering the amount of proteins, fats and carbohydrates in the feed, FU per kg, kg dry matter per day and MJ per day. Annex 3E-7 and 8 provides additional information about feed intake given in FU and grazing days for each livestock category.

Estimation of $GE_{FU, summer}$ covers the time where animals are grazing.

For dairy cows, the energy intake comes out at 18.3 MJ pr. FU in a standard winter feed regardless of whether the animal grazes or not, which is based on information from DCA. For bull calves (< ½ year), as well as bulls older than ½ year, the same energy content value is used, as for dairy cows.

For horses, heifers, suckling cattle, sheep and goats an average winter feed plan is provided based on information from DCA and DAAS on which the calculation of the GE content is based. Feeding conditions for deer is comparable with goats, why the GE for deer is based on feed plans for goats.

Table 6.6 GE per feeding unit, MJ per FU.

	GFU_{winter}	GFU_{summer}
Dairy cattle	18.3	18.3
Calves and bulls	18.3	18.8
Heifers	25.8	18.8
Suckling cattle	34.0	18.8
Sows	17.5	17.5
Weaners	16.5	16.5
Fattening pigs	17.3	17.3
Horses, sheep, goats and deer	30.0	18.8

In Annex 3E-9a, the annual average feed intake given in GE as MJ per day is shown, from 1990 to 2012, for each CRF livestock category and Annex 3E-9b shows the GE for each subcategory for non-dairy cattle and swine.

The Tier2/CS for enteric fermentation differs from the IPCC Tier 2 in the calculation of GE. A comparison between these two methods is shown in Chapter 6.3.4.

Methane conversion rate (Y_m)

Investigations from DCA have shown a change in fodder practice from use of sugar beet to maize (whole cereal). Sugar beet feeding gives a higher methane production rate compared to grass and maize due to the high content of easily convertible sugar. The development in fodder practice reflects the change in the average Y_m for dairy cattle and heifer from 6.39 in 1990 to 5.94 and 5.97 in 2012, respectively.

The estimation of the national values of Y_m is based on model “Karoline” developed by DCA based on average feeding plans for 20 % of all dairy cows in Denmark obtained from the Danish Agricultural Advisory Service DAAS (Olesen et al.; 2005). DCA have estimated the CH_4 emission for a winter feeding plan for two years, 1991 ($Y_m=6.7$) and 2002 ($Y_m=6.0$). Y_m for the years between 1991 and 2002 are estimated by interpolation and for 1990 and 2003 to 2012 by extrapolation where the actual sugar beet area is taken into account. Data for actual sugar beet area are shown in Table 6.7. Sugar beets are only included in the winter feeding plan and the Y_m is therefore also ad-

justed for days on winter and summer feeding plan. It is assumed that winter feeding plan covers 200 days. The value of the estimated average Y_m for 1991 and 2002 are, when adjusted for winter/summer, 6.35 and 5.96, respectively (see Table 6.8).

Table 6.7 Area grown with sugar beets and maize for feeding 1990-2012, ha.

Area	1990	1995	2000	2005	2008	2009	2010	2011	2012
Sugar beet for feeding	102 347	52 927	17 577	4 974	5 206	5 257	4 118	3 985	4 562
Maize for feeding	18 735	36 583	61 493	131 027	159 030	168 917	172 168	173 693	183 570

Table 6.8 CH_4 conversion rate (Y_m) – national factor used for dairy cattle and heifers > ½ year 1990 – 2012, %.

Dairy cattle + Heifers > ½ year	1990	1991	1995	2000	2002	2005	2008	2009	2010	2011	2012
Y_m excl. sugar beet	5.92	5.92	5.92	5.92	5.92	5.92	5.92	5.92	5.92	5.92	5.92
Y_m incl. sugar beet	6.78	6.70	6.36	6.06	6.00	5.96	5.96	5.96	5.95	5.95	5.95
Y_m grazing	5.92	5.92	5.92	5.92	5.92	5.92	5.92	5.92	6.00	6.00	6.00
Average Y_m											
Dairy cattle	6.39	6.35	6.16	6.00	5.94	5.94	5.94	5.94	5.94	5.94	5.94
Heifers	6.39	6.35	6.16	6.00	5.94	5.94	5.94	5.94	5.97	5.96	5.97

Implied emission factor

Table 6.9 shows the implied emission factors (IEFs) for all IPCC livestock categories. IEFs vary across the years for dairy cattle, non-dairy cattle, swine, goats and poultry due to changes for feed intake, distribution of animals in subcategories and number of grazing days. For goats new subcategories are introduced in 2005 and therefore the IEF differs from the other years. For sheep, horses, deer, ostrich and pheasants the IEF is constant. The emission from fur farming is considered to be not applicable (Hansen, 2010).

Table 6.9 Implied emission factors – Enteric Fermentation 1990 – 2012, kg CH_4 per head per year.

	1990	1995	2000	2005	2008	2009	2010	2011	2012
1. Cattle									
a. Dairy	116.62	119.45	117.16	128.12	130.67	133.76	134.35	132.91	133.91
b. Non-Dairy	34.77	34.96	35.01	36.98	40.82	40.12	39.82	40.38	40.00
3. Sheep	17.17	17.17	17.17	17.17	17.17	17.17	17.17	17.17	17.17
4. Goats	13.15	13.15	13.15	12.87	13.04	13.06	13.06	13.07	13.08
6. Horses	21.81	21.81	21.81	21.81	21.81	21.81	21.81	21.81	21.81
8. Swine	1.09	1.08	1.11	1.07	1.12	1.10	1.07	1.11	1.10
9. Poultry	0.004	0.003	0.003	0.004	0.004	0.003	0.003	0.003	0.003
10. Other									
Fur farming	NA	NA	NA	NA	NA	NA	NA	NA	NA
Deer	11.30	11.30	11.30	11.30	11.30	11.30	11.30	11.30	11.30
Ostrich	NO	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02
Pheasant	0.0003	0.0003	0.0003	0.0003	0.0003	0.0003	0.0003	0.0003	0.0003

NO = Not occurring.

NA = Not applicable.

The IEF for dairy cattle has increased from 117 kg CH_4 per cow per year in 1990 to 134 kg CH_4 in 2012. The IEF depends on milk yield and feed intake – see Figure 6.3. From 1990 to 2000 the IEF is almost unchanged but increased significant from 2000 to 2012. The development in feed intake follows the same development as the IEF, while the milk yields in percentage increases even more and especially from year 2000. This is caused by increased feed efficiency; an improvements of the feed utilization.

The milk yield has in average increased from 6 000 litre per cow in 1990 to approximately 8 400 litre per cow in 2012 (Statistics Denmark).




Figure 6.3 Comparison of feed intake, milk yield and IEF for dairy cattle (1990 = 100 %).

A comparison with IPCC Tier 2 calculation in Chapter 6.3.4 shows that the IEFs for the Danish inventory are lower. However, the national IEFs are considered more accurate because the differences can be explained by the improvement in feed efficiency, which has taken place in Danish agriculture from 2000.

The category “Non-Dairy Cattle” includes calves, heifers, bulls and suckling cattle and the IEF is a weighted average of these different subcategories. Changes in allocation of animals in subcategories can be reflected in the IEF. The development 1990 - 2008 shows a slight increase due to a higher feed consumption for heifers. From 2008 - 2012 the IEF seems stable.

The Danish IEF for non-dairy cattle is lower than the Tier 1 default value given in the IPCC Reference Manual (IPCC, 1997). This is due to a combination of lower Y_m value for heifers and lower weight/lower feed intake (Table 6.10). In Chapter 6.3.4 the national IEF is compared with IPCC Tier 2 calculation and the result shows a good correlation, which indicates the Danish estimate is correct.

Table 6.10 Subcategories for Non-Dairy Cattle 2012 – enteric fermentation.

Non Dairy Cattle – subcategories		Number of animals (DSt)	Energy intake, MJ per day	Methane conversion rate (Y_m), %	IEF, kg CH ₄ per head per yr
Calves, bull (0-6 month)	200 kg	132 893	61.67	4.00	16.18
Calves, heifer (0-6 month)	150 kg	162 286	102.23	6.00	39.69
Bulls (6 month to slaughter)	large breed: 440 kg sl. weight	132 849	116.24	4.00	29.54
	jersey: 330 kg sl. weight				
Heifers (6 month to calving)	325 kg	494 417	130.21	5.97 ^a	50.91
Suckling cattle	Up to 800 kg	97 193	163.55	6.00	63.91
Average - Non-Dairy Cattle			111.5		40.0
IPCC – default value			128.0		48.0

^a Average Y_m

The annual variations for swine primarily reflect the changes in the distribution of animals in subcategories (sows, weaners and fattening pigs). The feed intake for sows and weaners has overall increased while the feed intake for fattening pigs has decreased as a result of improved fodder efficiency (Annex 3E-7 and 9b).

In Table 6.11 the IEFs for swine subcategories are shown. The Danish IEF for swine is lower than the IPCC default value. The energy intake for fattening pigs is nearly the same as the default value, while the energy intake for weaners is significantly lower. The lower Danish IEF can be explained by the relatively high share of weaners.

Table 6.11 Subcategories for Swine 2011 – enteric fermentation.

Swine – subcategories	Number of animals (DSt)	Energy intake, MJ per day	Methane conversion rate (Y_m), %	IEF, kg CH ₄ per head per year
Sows (incl. piglets until 7.4 kg)	1 010 516	73.79	0.60	2.87
Weaners (7.4 – 32 kg)	5 847 458	14.25	0.60	0.44
Fattening pigs (32 – 107 kg)	5 472 905	40.55	0.60	1.49
Average - Swine		25.9		1.10
IPCC – default value		38	0.60	1.5

It is important to point out that the IEF for sheep and goats includes emission from lambs and kids due to the Danish normative data. This explains why the Danish IEFs are nearly twice as high as the IPCC default values. A comparison with IPCC Tier 2 which includes lamb indicates that the Danish estimates can be comparable with the IPCC default, see Chapter 6.3.4.

Activity data

In Table 6.12, the development in the number of animals from the agricultural statistics (Statistics Denmark), DAAS and CHR from 1990 to 2012 is presented (for subcategories see Annex 3E-2). The agricultural census does not include farms less than 5 ha. In the Danish emission inventory, the decision has been made to add number of sheep, goats and horses on small farms and deer, pheasants and ostriches based on information from DAAS and CHR (see Chapter 6.2.1).

Since 1990, the number of swine and poultry has increased, in contrast to the number of cattle, which has decreased. The number of cattle has decreased because the milk yield has increased while the total production of milk has been fixed by the EU milk quota. Buffalos, camels & llamas and mules & asses are not occurring in Denmark.

Table 6.12 Number of animals from 1990 to 2012, 1000 head.

CRF Table 4.A, 4.B (a) and 4.B (b)	1990	1995	2000	2005	2008	2009	2010	2011	2012
<u>IPCC livestock categories:</u>									
Dairy Cattle	753	702	636	564	558	563	568	565	587
Non-Dairy Cattle	1 486	1 388	1 232	1 006	1 006	977	1 003	1 003	1 020
Sheep*	92	81	112	126	117	116	111	94	90
Goats*	7	7	8	11	14	16	16	13	13
Horses*	135	143	150	175	190	178	165	155	155
Swine	9 497	11 084	11 922	13 534	12 738	12 369	13 173	12 932	12 331
Poultry	16 249	19 619	21 830	17 632	15 406	19 676	18 731	19 319	18 991
Other;									
Fur farming	2 264	1 850	2 199	2 552	2 810	2 721	2 699	2 757	2 948
Pheasant**	1 063	1 063	1 063	1 063	1 063	1 063	1 063	1 063	1 063
Deer**	10	10	10	10	10	9	10	8	7
Ostrich**	NO	3	9	4	0.5	0.4	0.4	0.2	0.2

* Including animals on small farms (less than 5 ha), which are not covered by Statistics Denmark.

** Not included in Statistics Denmark.

NO = Not occurring.

6.3.3 Time series consistency

The CH₄ emission from enteric fermentation is given in Table 6.13. From 1990 to 2012, the emission has decreased by 13 %, which is primarily related to a decrease in the number of cattle. The number of swine has increased from 9.5 million in 1990 to 12.3 million in 2012, but this increase is only of minor importance in relation to the total CH₄ emission from enteric fermentation.

Table 6.13 Emission of CH₄ from Enteric Fermentation 1990 – 2012, Gg CH₄.

CRF 4.A	1990	1995	2000	2005	2008	2009	2010	2011	2012
Dairy Cattle	87.83	83.91	74.46	72.29	72.91	75.32	76.34	75.11	78.63
Non-Dairy Cattle	51.66	48.52	43.14	37.20	41.08	39.21	39.93	40.49	40.79
Sheep	1.58	1.39	1.92	2.17	2.02	1.98	1.91	1.61	1.55
Goats	0.10	0.09	0.11	0.15	0.18	0.20	0.21	0.16	0.17
Horses	2.94	3.11	3.27	3.82	4.14	3.87	3.60	3.38	3.38
Swine	10.33	12.02	13.17	14.53	14.27	13.66	14.14	14.34	13.61
Poultry	0.06	0.07	0.06	0.07	0.06	0.06	0.06	0.07	0.06
Other;									
Fur farming	NA	NA	NA	NA	NA	NA	NA	NA	NA
Deer	0.11	0.11	0.11	0.11	0.11	0.11	0.11	0.09	0.08
Ostrich	NO	#	#	#	#	#	#	#	#
Pheasant	#	#	#	#	#	#	#	#	#
Total, Gg CH ₄	154.62	149.22	136.25	130.33	134.78	134.42	136.28	135.25	138.28
Total, Gg CO ₂ eqv.	3 247	3 134	2 861	2 737	2 830	2 823	2 862	2 840	2 904

NO = Not occurring.

NA = Not applicable.

- emission ≤ 0.0003.

6.3.4 Tier 2/Country Specific compared to IPCC Tier 2 method

A comparison between IPCC Tier 2 and Denmark's Tier2/Country Specific (CS) calculation method for enteric fermentation is made. In the IPCC Guidelines default values are given for dairy cattle, non-dairy cattle and sheep, therefore a comparison is made for these three groups. The comparison is not done annually and the current comparison is based on 2008 data.

Calculations of IEFs are made by IPCC Tier 2, with both default and national values for Y_m , and Denmark's Tier 2/CS method. A comparison between IEFs (Table 6.14) shows that the Danish method gives a value for dairy cattle that is up to 7 % lower than the IPCC Tier 2 method. For non-dairy cattle the result is up to 5 % higher. To compare the IEF for sheep the calculation includes lamb. The Danish method gives a 7 % lower value than the IPCC with default values for Y_m ($Y_m = 0.07$), but a 9 % higher value than IPCC with an average Y_m of values for mother sheep and lamb ($Y_m = 0.06$).

Table 6.14 IEFs for enteric fermentation calculated by different methods, 2008.

kg CH ₄ per animal per year	Tier 2 (IPCC Y_m)	Tier 2 (DK Y_m)	Tier 2/CS
Dairy cattle	140.3	137.9	130.7
Non-dairy cattle	38.8	38.3	40.8
Sheep (incl. lambs)	18.4	15.8	17.2

The three different Tier 2 calculations for non-dairy cattle all show an IEF between 38.3-40.8 kg per head per year, which indicates that the Tier 2/CS used in the Danish inventory is reasonable. However, these values are lower compared to the Tier 1 default value at 48 kg per head per year given in the Reference Manual, Table 4-4 (IPCC, 1997), which can be explained by a combination of lower Y_m for heifers and lower animal weight/lower feed intake.

The calculation of IEF for sheep indicate that the value used in the Danish inventory are reasonable. A Tier 2 calculation, where the productions of lamb are included, based on IPCC Y_m shows an IEF at the same level.

The lower value for the IEF for dairy cattle is mainly due to a lower value for GE (Table 6.15). The Danish values for feed consumption are based on the Danish normative figures and the normative data are based on actual efficacy feeding controls or actual feeding plans at farm level, more info on GE calculations is included in Chapter 6.3.2.

Table 6.15 GE for dairy cattle calculated by different methods, 2008.

MJ per animal per day	Tier 2 (IPCC Y_m and DK Y_m)	Tier 2/CS
Dairy cattle	356.4	335.3

According to Statistics Denmark dairy cattle produce 22.5 kg milk in average per animal per day in 2008. Table 6.16 shows the needed energy intake to achieve this milk production calculated by two different methods. By using the Tier 2 calculation given in the Reference Manual (IPCC, 1997) 15.8 MJ is needed to produce 22.5 kg milk per animal per day. National data for feed intake, which reflect the actual Danish agricultural conditions, show a lower need of energy intake corresponding to 14.9 MJ. This is a result of improved feeding efficiency.

Table 6.16 MJ per kg milk produced 2008.

	Kg milk per animal per day	MJ per kg milk Tier 2	MJ per kg milk Tier 2/CS
Dairy cattle	22.5	15.8	14.9

In Figure 6.4 is shown the Danish trend of MJ per kg milk for dairy cattle. It is seen that the energy intake per kg milk have overall decreased from 1996 to 2008. Around 1999 the Danish level of MJ per kg milk was at the same level as given in the IPCC Tier 2 method. Since then, feeding efficiency has

continued to rise due to the structural development, i.e. bigger farms and more intensive production. This explains the lower IEF for dairy cattle based on the Danish methodology.




Figure 6.4 The Danish trend for MJ per kg milk produced for dairy cattle, 1990-2008.

6.4 CH₄ and N₂O emission from manure management (CRF sector 4B)

6.4.1 Description

This source contributes with 18 % of the total greenhouse gas emission from the agricultural sector in 2012. The major part of the emission originates from the production of cattle (48 %) followed by swine production (39 %). The remaining part is mainly from poultry and fur farming (13 %).

6.4.2 Methodological issues

CH₄ emission

The IPCC Tier 2/CS methodology is used for the estimation of the CH₄ emission from manure management. The calculation is based on manure excretion instead of feed intake as described in IPCC Reference manual (IPCC, 1997). Default values for maximum methane producing capacity (B₀) and methane conversion factor (MCF) given by the IPCC are used. The calculation of volatile solids (VS) is based on national data.

Table 6.17 CH₄ – Manure management – use of national parameters and IPCC default values.

CH ₄ – Manure management	National parameters	IPCC default value
Volatile solids, VS	Based on amount of manure (Annex 3E-10a and 10b)	
Maximum methane producing capacity, B ₀		IPCC 1997
Methane conversion factor, MCF		IPCC 1997

The amount of manure is calculated for each combination of livestock sub-category and housing type and then aggregated to the IPCC livestock categories. In the calculation grazing days and use of straw in the housing are taken into account. Equation for CH₄ calculation:

$$CH_{4\text{manure}} = CH_{4\text{housing}} + CH_{4\text{grazing}}$$

$$CH_{4\text{housing}} = VS_{\text{housing}} \cdot MCF \cdot 0.67 \cdot B_0$$

$$CH_{4\text{grazing}} = VS_{\text{grazing}} \cdot MCF \cdot 0.67 \cdot B_0$$

Estimation of VS

VS is calculated from data concerning amount of manure, dry matter content, share of VS in dry matter, amount of bedding and grazing days. Except from grazing days for dairy cattle and heifers, all these parameters are based on Danish Normative data.

The determination of VS is country-specific, given that it is based on the amount of manure excreted.

$$VS_{\text{housing}} = \frac{m}{365} \cdot DM_M \cdot VS_{DM} \cdot (365 - g_1) + s \cdot DM_S \cdot \left(1 - \frac{\% \text{ ash}}{100}\right) \cdot (365 - g_2)$$

$$VS_{\text{grazing}} = \frac{m}{365} \cdot DM_M \cdot VS_{DM} \cdot g_1$$

Where:

VS	= volatile solids, kg animal ⁻¹ yr ⁻¹
m	= amount of manure excreted, kg animal ⁻¹ yr ⁻¹
DM	= dry matter of M manure or S straw, pct.
VS _{DM}	= volatile solids of dry matter, pct.
g ₁	= feeding days on grass, days yr ⁻¹ ¹
g ₂	= actual days on grass, days yr ⁻¹
s	= amount of straw, kg animal ⁻¹ yr ⁻¹
% ash	= ash content in straw

The ash content in straw is set to 4.5 % (DAAS, 2005). The VS of dry matter is 78 % for cattle, horses, sheep, goats and deer. For swine, poultry and fur animals the VS of dry matter is 75 % (Møller, 2003). The number of days on grass is shown in Annex 3E-8. The amount of manure excreted and straw used depends on housing type and is given in the normative figures table (Poulsen, 2013).

The VS daily excretion in average for all main livestock categories and cattle subcategories is shown in Annex 3E-10a and 10b.

MCF used for slurry

Default values provided in the IPCC guidelines for the methane production B_0 and MCF are used. For liquid systems, the MCF of 10 % in the Reference Manual (IPCC, 1997) is used.

The Revised 1996 IPCC guidelines (IPCC, 1997) provide a default MCF of 10 % for liquid/slurry, which is based on research of Hashimoto & Steed (1993) and Woodbury & Hashimoto (1993). This MCF value was changed to 39 % in the IPCC Good Practice Guidance (IPCC, 2000), without any scientific argumentation, documentation or specific references. It has to be remarked that the 2006 IPCC Guidelines (IPCC, 2006) return to a MCF value of 10 % for Danish conditions referenced to “Judgement of IPCC Expert Group in

¹ Actual days on grass are the number of days that heifers is outside. Feeding days on grass is higher than actual days on grass due to a higher feed intake during grazing compared to the period in housing. Feeding days on grass is a conversion of this higher feed intake on grass. This is only relevant for heifers.

combination with Mangino et al. (2001) and Sommer et al. (2000)" (IPCC, 2006).

The methane emission from liquid systems is very sensitive to temperature effects. Basically most of the manure is stored in Denmark under cold conditions (<5-10 degrees Celsius). The CH₄ formation practically stops at 4 °C and therefore there are no plausible arguments that 39 % of total CH₄ capacity should be released under Danish conditions. Danish studies confirm this assumption (Husted, 1994 and Sommer et al., 2000). Furthermore, investigations based on measurements in Canada, where conditions are similar to Denmark, support this value (Massé et al., 2003). Support of the value of 10 % is also found from a Swedish review (Dustan, 2002), taking into account both the cold climate and the fact that the slurry containers in Scandinavia usually have a surface cover.

Considering the agricultural conditions in Denmark and the present scientific knowledge as described above a MCF of 10 % for liquid/slurry is more appropriate under the Danish conditions. The Danish decision of using a MCF of 10 % is as demonstrated above backed by several scientific papers as well as both the Revised 1996 IPCC Guidelines (IPCC, 1997) and the 2006 IPCC Guidelines (IPCC, 2006). Therefore, Denmark intends to continue using a MCF value of 10 % until new scientific knowledge is published contradicting the above mentioned references.

It has to be remarked that countries with comparable climate use a MCF for liquid/slurry at the same level as default recommended in the Revised IPCC 1996 Guidelines (IPCC, 1997). Finland use the same value as Denmark, i.e. a MCF of 10 %, Germany uses 14 %, Belgium uses 19 % and Norway and the Sweden use a MCF below 10 %.

A lower CH₄ emission from biogas treated slurry

A study indicates a lower CH₄ emission from slurry treated in biogas plants (Sommer et al., 2001). No description on how to include biogas treated slurry in the inventories is provided in the IPCC guidelines. Therefore, the Danish inventory uses data based on a Danish study (Sommer et al., 2001).

The Danish Energy Agency (DEA) provides annually the Danish energy statistics, which for 2012 shows that the produced energy from biogas corresponds to 4383 TJ (DEA, 2013). The majority of this production (2969 TJ) takes place on biogas plants and uses agricultural products which mainly are animal manure (slurry). In Denmark exist around 80 biogas plants and the largest twenty plants produce 70 % of the total agricultural biogas production.

Table 6.18 Number of biogas plants and production of energy

Production size, TJ per year	No. of biogas plants	Energy production, TJ
<10	22	94
10-50	33	780
50-100	12	848
100-200	10	1246
Total	77	2969

The energy potential very much depends on the type of biomass input; the combination of manure and other agricultural biomasses, the storage time and the technology on the different biogas DEA assume an average energy potential where 0.83 million tonnes of slurry produce 1 PJ biogas. This

means that in 2012 around 2.5 million tonnes of slurry has been treated in biogas plants.

It is assumed that of the total amount of biogas treated slurry, cattle slurry makes up 45 % and pig slurry 55 % (Tafdrup, 2010). In 2012, the amount of biogas treated slurry is equivalent to approximately 6 % of total slurry in housing.

The lower CH₄ emission as a consequence of biogas treated slurry is calculated as the difference between non-treated slurry and treated slurry.

$$CH_{4,lower} = CH_{4,non-treated\ slurry} - CH_{4,treated\ slurry}$$

The calculation is based on the amount of volatile solids (VS) calculated as the VS percentage of dry matter (DM) which is 80 % for both cattle and pig slurry. The dry matter content is based on the Danish normative figures (Poulsen et al., 2001 and Poulsen, 2013).

The CH₄ emission from treated and non-treated slurry is calculated as:

$$CH_{4,non-treated\ slurry} = VS \cdot B_0 \cdot MCF \cdot 0.67$$

$$CH_{4,treated\ slurry} = VS \cdot B_0 \cdot MCF \cdot 0.67 \cdot E_{lower}$$

Where; CH_{4,non-treated slurry} and CH_{4,treated slurry} are the emission of non-biogas treated slurry and biogas treated slurry, respectively. VS express the total amount of volatile solid in non-biogas treated slurry and biogas treated slurry, B₀ is the maximum methane forming capacity, MCF is the methane conversion factor and the factor 0.67 express the conversion from m³ to kg. E_{lower} is the lower emission of biogas treated slurry compared to untreated slurry.

Based on results from Sommer et al. (2001) it is assumed that the emission from treated cattle slurry is reduced by 23% compared with untreated slurry. This leads to an E_{lower} for cattle of 0.77. Likewise, results from treated pig slurry show a 40 % lower emission than for untreated slurry, which leads to an E_{lower} at 0.60 (Sommer et al., 2001). Refer to Annex 3E-11.

All key model parameters for estimating the lower CH₄ emission in 2012 as a result of biogas plants are listed in Table 6.19. Data for 1990 to 2012 are shown in Annex 3E-12a.

Table 6.19 Key model parameters used to calculate the lower CH₄ emission due to biogas treated slurry, 2012.

	Slurry, DM ^a		VS of	VS in	MCF ^c	B ₀ ^c	E _{lower} ^d	CH ₄ emission	CH ₄ emission	Lower CH ₄
	biogas		DM ^b	treated				in untreated	in biogas	emission
	treated			slurry				slurry	treated slurry	
	1000 Gg	Pct.	Pct.	10 ⁶ kg VS	Pct.	m ³ CH ₄		Gg CH ₄	Gg CH ₄	Gg CH ₄
	per kg VS									
Cattle slurry	1.10	10.3	80	90.85	10	0.24	0.77	1.46	1.11	0.34
Pig slurry	1.35	6.1	80	65.76	10	0.45	0.60	1.98	1.19	0.80
Lower emission										1.14

^a Poulsen et al., 2001 and Poulsen, 2013.

^b Møller, 2003.

^c IPCC default.

^d Sommer et al., 2001.

Due to biogas treated slurry, the total emission of CH₄ in 2012 is lowered by 1.14 Gg CH₄ (Table 6.19), which correspond to a 2 % reduction of the total CH₄ emission from manure management in 2012. Calculations for the lower CH₄ emission for all years 1990 – 2012 are listed in Table 6.20.

The lower emission is subtracted in the emission related to manure management from dairy cattle and fattening pigs, which are the main sources of the production of slurry.

Table 6.20 Lower CH₄ emissions as a result of biogas treated slurry 1990 – 2012.

	1990	1995	2000	2005	2008	2009	2010	2011	2012
Amount of treated slurry, Mt									
- cattle	0.09	0.29	0.52	0.87	0.99	1.08	1.12	1.08	1.10
- swine	0.10	0.35	0.64	1.06	1.20	1.31	1.37	1.31	1.35
VS total in treated slurry									
- cattle	0.01	0.02	0.04	0.07	0.08	0.09	0.09	0.09	0.09
- swine	0.01	0.02	0.03	0.05	0.06	0.06	0.07	0.06	0.07
Total reduced emission, Gg CH ₄	0.09	0.30	0.54	0.90	1.02	1.11	1.16	1.11	1.14

CH₄ implied emission factors

Table 6.21 shows the development in the implied emission factors from 1990 to 2012. Variations between the years for dairy cattle, non-dairy cattle, poultry, swine and fur farming reflect changes in feed intake, distribution of animals in subcategories, grazing situation and changes in housing type system.

The IEFs for poultry, ostriches, pheasants and deer are unaltered because of very few changes in feed intake and grazing days. A more detailed division in subcategories for goats and horses is implemented from 2007 and 2003, respectively, and explains the small changes in IEFs.

The IEF for sheep and goats includes lambs and kids and the housing systems for these categories are defined as deep litter with a MCF of 10 %. This explains why the Danish IEF is considerably higher than the IPCC default value, which is provided for sheep and goats housed in solid storage systems. IEF for sheep and goats change in 2012 due to change in amount of manure excreted and dry matter content.

Table 6.21 Implied emission factors – Manure Management 1990 – 2012, kg CH₄ per head per year.

	1990	1995	2000	2005	2008	2009	2010	2011	2012
1a. Dairy Cattle	21.06	22.87	27.55	34.25	32.18	33.09	33.20	32.73	33.60
1b. Non-Dairy Cattle	5.71	6.73	7.40	9.23	9.71	9.66	9.49	9.72	9.19
3. Sheep	2.82	2.82	2.82	2.82	2.82	2.82	2.82	2.82	2.71
4. Goats	2.44	2.44	2.44	2.45	2.45	2.45	2.45	2.45	2.37
6. Horses	2.96	2.96	2.96	2.95	2.95	2.95	2.95	2.95	2.95
8. Swine	2.12	2.19	2.22	2.22	2.27	2.29	2.24	2.31	2.30
9. Poultry	0.03	0.03	0.02	0.03	0.03	0.03	0.03	0.03	0.03
10. Other									
Fur farming	0.57	0.59	0.65	0.79	0.92	0.97	0.99	1.03	1.04
Deer	0.30	0.30	0.30	0.30	0.30	0.30	0.30	0.30	0.30
Ostrich	NO	1.47	1.47	1.47	1.47	1.47	1.47	1.47	1.47
Pheasant	0.001	0.001	0.001	0.001	0.001	0.001	0.001	0.001	0.001

IEF for dairy cattle has increased as a result of increasing milk yield, but also because of changes in housing types. Annex 3E-1 shows the changes in housing types from 1990 to 2012. Old-style tethering systems with solid manure have been replaced by loose-housing with slurry-based systems. The MCF for liquid manure is ten times higher than that for solid manure. For non-dairy cattle the same development in IEF is seen, but here it is mainly because an increasing proportion of bull-calves are raised in housings with deep litter, where the MCF is also high.

For pigs and fur farming, there has been a similar development as for dairy cattle with a move from solid manure to slurry-based systems.

As shown in Table 6.22 the national IEF for dairy cattle is particularly high compared to the IPCC default value, which is mainly due to the fact that more cattle are housed on slurry based system than given in the IPCC assumptions. Furthermore, VS used for Danish dairy cattle are higher due to higher milk yield.

For non-dairy cattle the national VS value is nearly the same as the default, but a high proportion of the animals are housed in deep litter or liquid/slurry systems, 39 % and 31 % in 2012 respectively, which both have a MCF at 10 %.

Table 6.22 Cattle – important parameters for calculation of the average implied emission factor for manure management 2012.

	IPCC			DK		
	VS kg dm per hd per day	Liquid/slurry %	IEF kg CH ₄ per hd per yr	VS kg dm per hd per day	Liquid/slurry %	IEF kg CH ₄ per hd per yr
Dairy	5.1	40	14	6.2	88	34
Non-dairy (average)	2.7	50	6	2.7	31	9
Calves, bull				1.5	0	0
Calves, heifer				1.8	0	0
Bulls > ½ yr				3.9	40	16
Heifer > ½ yr				2.8	50	9
Suckling cattle				4.3	7	12

The category of swine in the Danish inventory operates with three subcategories. The IEF is lower compared with the IPCC default value due to a lower VS value probably because of the relatively high share of weaners. In the Reference Manual (IPCC, 1997) an average feed intake of 38 MJ per head per day is used, which is significantly higher than the average feed intake for Danish weaners and fattening pigs.

Table 6.23 Swine – important parameters for calculation of the average implied emission factor for manure management 2012.

	IPCC				DK			
	VS kg dm per hd per day	Feed intake MJ per hd per day	Pit > 1 month %	IEF Kg CH ₄ per hd per year	VS kg dm per hd per day	Feed intake MJ per hd per day	Liquid/slurry %	IEF kg CH ₄ per hd per year
Swine	0.5	38	73	3.0	0.2		98	2.3
Sows (incl. piglets until 7 kg)					0.5	74		6.9
Weaners (7-32 kg)					0.1	2		0.2
Fattening pigs (32-107 kg)					0.3	10		0.8

In Table 6.24 are shown the emission of CH₄ from manure management from 1990 to 2012. The main part of the emission originates from cattle and swine. The emission is increased from 1990 to 2012 and this is mainly due to a change in housing systems to a higher share of slurry based systems.

Table 6.24 Emission of CH₄ from Manure Management 1990 – 2012, Gg CH₄.

CRF 4.A	1990	1995	2000	2005	2008	2009	2010	2011	2012
Dairy Cattle	15.86	16.07	17.51	19.33	17.96	18.63	18.86	18.50	19.73
Non-Dairy Cattle	8.48	9.34	9.12	9.28	9.77	9.44	9.52	9.74	9.37
Sheep	0.26	0.23	0.31	0.36	0.33	0.33	0.31	0.26	0.25
Goats	0.02	0.02	0.02	0.03	0.03	0.04	0.04	0.03	0.03
Horses	0.40	0.42	0.44	0.52	0.56	0.52	0.49	0.46	0.46
Swine	20.13	24.26	26.41	30.01	28.90	28.34	29.44	29.94	28.38
Poultry	0.47	0.54	0.52	0.53	0.49	0.51	0.53	0.51	0.50
Other;									
Fur farming	1.29	1.09	1.44	2.01	2.57	2.64	2.68	2.85	3.05
Deer	0.003	0.003	0.003	0.003	0.003	0.003	0.003	0.002	0.002
Ostriches	NO	0.005	0.013	0.005	0.001	0.001	0.001	0.0003	0.0003
Pheasants	0.001	0.001	0.001	0.001	0.001	0.001	0.001	0.001	0.001
Total, Gg CH ₄	46.91	51.98	55.80	62.07	60.63	60.45	61.88	62.30	61.77
Total, Gg CO ₂ eqv.	985	1 092	1 172	1 303	1 273	1 270	1 300	1 308	1 297

No = Not occurring.

N₂O emission

The N₂O emission from manure management is based on the amount of nitrogen in the manure in housings. The emission from manure deposited on grass is included in “Pasture, Range and Paddock Manure” (Chapter 6.5.2). The IPCC default emission factors are applied, see Table 6.25.

Table 6.25 Emission factors for N₂O from manure management.

	Emission factor	
	Unit	IPCC – default values
<u>Handling of manure:</u>		
Solid manure, poultry	kg N ₂ O-N per kg N	0.005
Solid manure, other	kg N ₂ O-N per kg N	0.02
Slurry and urine	kg N ₂ O-N per kg N	0.001
Deep litter	kg N ₂ O-N per kg N	0.02
Deep litter, farmyard manure < 1 month ¹	kg N ₂ O-N per kg N	0.005

¹ Farmyard manure, which is faeces and urine mixed with large amounts of bedding (usually straw) on the floors of cattle or swine housing.

The total amount of nitrogen in manure has decreased by 12 % from 1990 to 2012 (Table 6.26), despite a significant increase in the production of pigs and poultry. This reduction is due to improvements in fodder efficiency, especially for fattening pigs. A decrease in total amount of nitrogen also means a decrease for the N₂O emission. Another reason for the decreased N₂O emission is a change from the previous more traditional tethering systems with solid manure to slurry based systems due to the lower emission factor for liquid manure than for solid manure. For example for dairy cattle 70 % is placed on slurry based housing system in 1990, which is increased to 88 % in 2012. It is important to point out that the N-excretion rates shown in Table 6.26 are values weighted for the subcategories (Table 6.4). N-excretion reflects nitrogen excreted per animal per year (per AAP). The variations in N-excretion in the time series reflect changes in feed intake, fodder efficiency and distribution of animals in subcategories.

Table 6.26 Nitrogen excretion, annual average 1990 – 2012, kg N per head per year (AAP).

CRF Table 4.B(b)	1990	1995	2000	2005	2008	2009	2010	2011	2012
<u>Livestock category</u>									
Dairy cattle	129.49	125.23	125.31	133.30	137.98	138.12	138.63	138.47	138.03
Non-dairy	35.59	36.26	36.39	40.88	45.53	44.81	43.15	44.11	43.39
Sheep	21.18	21.90	16.95	16.95	16.95	16.95	16.95	16.95	16.95
Goats	21.18	21.90	16.95	15.83	16.32	16.37	16.40	16.43	16.55
Swine	11.84	9.70	9.61	9.19	8.63	8.33	7.81	7.98	8.01
Poultry	0.63	0.62	0.55	0.73	0.74	0.55	0.60	0.55	0.54
Horses	44.15	39.56	39.56	39.56	39.56	39.56	39.56	39.56	39.56
Fur farming	4.90	4.65	4.62	5.38	5.29	5.51	5.82	5.65	5.44
Deer	16.00	16.00	16.00	16.00	16.00	16.00	16.00	16.00	16.00
Ostrich	NO	15.61	15.60	15.60	15.60	15.60	15.60	15.60	15.60
Pheasant	0.04	0.04	0.04	0.04	0.04	0.04	0.04	0.04	0.04
N-excretion, total, Gg N per year	292	274	270	277	269	260	261	260	258
N-excretion, housing, Gg N per year	258	239	235	251	246	237	239	239	237

NO – Not occurring.

Table 6.27 shows the N₂O emission from manure management distributed on the main livestock categories. The total emission decreased from 1.94 Gg N₂O in 1990 to 1.32 Gg N₂O in 2012 and this trend is particularly determined by the emission from cattle and swine. The main reason is change in housing types towards more animals on slurry based system. For swine is seen the same development as for cattle where the share of slurry based system has increased from 89 % in 1990 to 98 % in 2012. Furthermore, the N-excretion from fattening pigs and weaners has decreased significantly.

Table 6.27 Emission of N₂O from manure management 1990-2012, Gg N₂O.

	1990	1995	2000	2005	2008	2009	2010	2011	2012
Dairy Cattle	0.57	0.50	0.42	0.31	0.31	0.29	0.30	0.27	0.28
Non-Dairy Cattle	0.38	0.38	0.37	0.39	0.43	0.41	0.41	0.41	0.40
Sheep	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.01	0.01
Goats	0.001	0.001	0.001	0.002	0.002	0.002	0.002	0.002	0.002
Swine	0.54	0.46	0.46	0.46	0.31	0.26	0.26	0.24	0.22
Poultry	0.26	0.33	0.35	0.36	0.32	0.31	0.32	0.30	0.29
Horses	0.09	0.09	0.09	0.11	0.12	0.11	0.10	0.10	0.10
Fur farming	0.08	0.06	0.06	0.05	0.03	0.03	0.02	0.02	0.03
Deer	IE	IE	IE	IE	IE	IE	IE	IE	IE
Ostriches	NO	0.0013	0.0035	0.0014	0.0002	0.0001	0.0001	0.0001	0.0001
Pheasants	IE	IE	IE	IE	IE	IE	IE	IE	IE
Total excl. biogas	1.94	1.84	1.76	1.70	1.53	1.43	1.43	1.36	1.32

NO – Not Occurring.

IE - Included Elsewhere. Deer and pheasants are at pasture 365 days a year and the emissions are reported under 4.D.2 Pasture, Range and Paddock.

A lower N₂O emission from biogas treated slurry

Studies indicate a lower N₂O emission from biogas treated slurry compared to untreated slurry (Sommer et al., 2001 and Sommer et al., 2004). The lower emission is a result of displacement in allocation between the fraction of degradable and non-degradable VS. Biogas treated slurry increase the fraction of non-degradable VS, which promote the oxygen content in soil. These conditions will reduce the potential for N₂O emission, because N₂O emission

takes place in environments without oxygen or with very low concentrations of oxygen (Sommer et al., 2001).

In practice this effect of a lower N₂O emission will take place in the manure applied on soil. However, it is chosen, in the inventory, to account for the lower N₂O emission by subtracting it from the manure management emission. The biogas treatment is taken place, before the slurry is applied to soil.

No methodology is provided in the IPCC Reference Manual or GPG on how to account for this reduction. The estimation is based on a Danish study (Sommer et al., 2001). The reduced N₂O emission is calculated as:

$$N_2O-N_{lower} = N_2O-N_{non-treated\ slurry} - N_2O-N_{treated\ slurry}$$

The N₂O emission from treated and non-treated slurry is calculated as:

$$N_2O-N_{non-treated} = N_{slurry\ non-treated} \cdot N_{content} \cdot EF_{N_2O}$$

$$N_2O-N_{treated} = N_{slurry\ treated} \cdot N_{content} \cdot E_{N_2O} \cdot EF_{lower}$$

Where; N₂O-N_{non-treated} slurry and N₂O-N_{treated} slurry are the emission of non-biogas treated slurry and biogas treated slurry, respectively. N_{content} express the nitrogen content in slurry. N_{slurry} is the total amounts of N in slurry, EF_{N₂O} express the N₂O emission factor based on IPCC default (1.25 %). E_{lower} is the lower emission of biogas treated slurry compared to untreated slurry.

Based on result in Sommer et al. (2001) it is assumed that the emission from treated cattle slurry is 36 % lower compared to untreated slurry, which lead to an E_{lower} of 0.64. Result from Sommer et al. (2001) concerning the pig slurry shows a 41 % lower N₂O emission from treated slurry, which results in a E_{lower} of 0.59. Refer to Annex 3E-11.

All key model parameters for estimating the lower N₂O emission in 2012 as a result of biogas treatment of manure are listed in Table 6.28.

Table 6.28 Key model parameters used to calculate the lower N₂O emission due to biogas treated slurry, 2012

	Slurry, biogas treated 1000 Gg	Total N in treated slurry ^a Pct.	E _{lower} ^b	N ₂ O emission, untreated slurry Gg N ₂ O	N ₂ O emission, biogas treated slurry Gg N ₂ O	Lower N ₂ O emission Gg N ₂ O
Cattle slurry	1.10	0.538	0.64	0.074	0.047	0.027
Pig slurry	1.35	0.541	0.59	0.091	0.054	0.037
Lower emission						0.064

^a Poulsen et al., 2001

^b Sommer et al., 2001.

Data for 1990 to 2012 are shown in Annex 3E-12b.

Due to the biogas treatment, the emission of N₂O in 2012 is reduced by 0.06 Gg N₂O (Table 6.29) which corresponds to a 4 % reduction of the N₂O emission from manure management in 2012.

Table 6.29 Lower N₂O emissions from manure management as a result of biogas-treated slurry 1990 – 2012.

	1990	1995	2000	2005	2008	2009	2010	2011	2012
Amount of treated slurry, Mt									
- cattle	0.09	0.29	0.52	0.87	0.99	1.08	1.12	1.08	1.10
- swine	0.10	0.35	0.64	1.06	1.20	1.31	1.37	1.31	1.35
Total reduced emission, Gg N ₂ O	0.01	0.02	0.03	0.05	0.06	0.06	0.06	0.06	0.06
Total N ₂ O emission									
incl. biogas, Gg N ₂ O	1.93	1.83	1.73	1.65	1.47	1.36	1.36	1.30	1.26

6.4.3 Time series consistency

In Table 6.30, the national emission from manure management from 1990 to 2012 is shown. The N₂O emission has decreased by 35 %. The national emission from manure management has, nevertheless, increased by 7 % in CO₂ equivalents due to the increase in the CH₄ emission.

Table 6.30 Emissions of N₂O and CH₄ from Manure Management 1990 – 2012.

	1990	1995	2000	2005	2008	2009	2010	2011	2012
<u>N₂O emission</u>									
Liquid manure, Gg N ₂ O	0.30	0.27	0.25	0.26	0.26	0.24	0.24	0.25	0.25
Solid manure, Gg N ₂ O	1.01	0.81	0.64	0.50	0.36	0.30	0.29	0.24	0.23
Other manure, Gg N ₂ O	0.62	0.75	0.84	0.89	0.86	0.82	0.83	0.81	0.79
Total, Gg N ₂ O	1.93	1.83	1.73	1.65	1.47	1.36	1.36	1.30	1.26
Total, Gg CO ₂ eqv.	600	566	537	512	457	423	422	403	391
<u>CH₄ emission</u>									
Total, Gg CH ₄	46.91	51.98	55.80	62.07	60.63	60.45	61.88	62.30	61.77
Total, Gg CO ₂ eqv.	985	1 092	1 172	1 303	1 273	1 270	1 300	1 308	1 297
Total Manure Management,									
Gg CO ₂ eqv.	1 585	1 657	1 709	1 816	1 730	1 693	1 722	1 711	1 688
Incl. the reduction from biogas treated slurry.									

6.5 N₂O emission from agricultural soils (CRF sector 4D)

6.5.1 Description

The emissions from agricultural soils, contribute, in 2012 with 93 % of the N₂O emission from the agricultural sector. Figure 6.5 shows the distribution and the development from 1990 to 2012 according to different N₂O sources. The emission has overall decreased 39 %. The increase from 2007 to 2008 was due to a rise in the use of synthetic fertiliser, which can mainly be explained by stockpiling due to expectations of rising prices. In 2009 the emission has decreased again and since then nearly no changes have taken place.

The main part of the emission originates as direct emission. The largest sources are manure and synthetic fertiliser applied on agricultural soils. Another large source is the indirect N₂O emission, of which the emission from nitrogen leaching is the major part.



Figure 6.5 N₂O emissions from agricultural soils 1990 - 2012.

6.5.2 Methodological issues

To calculate the N₂O emission the IPCC Tier 1b methodology is used in combination with a country specific method (CS). Tier 1b is used in calculation of emission from N-fixing crops, crop residue and atmospheric deposition.

Emissions of N₂O are closely related to the nitrogen balance and all data concerning the evaporation of NH₃ and data for manure condition is applied from the national NH₃ emission inventory. This is described in great detail in Mikkelsen et al. (2011) and Denmark's annual inventory report to the UNECE Convention on Long-Range Transboundary Air Pollution (Nielsen et al., 2013b). For the calculation of the N₂O emission from nitrogen leaching and runoff a national model is used.

In the calculation of N₂O from agricultural soils the N₂O emission factors for all sources are based on the default values given by the IPCC (2000). A NH₃ and N₂O emission factor overview is presented in Table 6.31. The estimated emissions from the different sub-sources are described in the text, which follows.

Table 6.31 Emission factors – NH₃ and N₂O from agricultural soils 1990 – 2012.

	NH ₃ emission factor (national data) Kg NH ₃ -N per kg N	N ₂ O emission factor (IPCC default value) kg N ₂ O -N per kg N
1. Direct Soil Emissions		
Synthetic Fertiliser Applied to Soils	0.02	0.0125
Animal Wastes Applied to Soils	0.21*	0.0125
N-fixing Crops		0.0125
Crop Residue		0.0125
Cultivation of Histosols		8**
Industrial Waste Used as Fertiliser		0.0125
Sewage Sludge Used as Fertiliser	0.02	0.0125
2. Pasture, range and paddock		
	0.07	0.02
3. Indirect Soil Emissions		
Atmospheric Deposition		0.01
Nitrogen Leaching and Run-off		0.025***

*Varies from year to year, has decreased from 0.28 in 1990.

**Unit: kg N₂O-N pr ha.

***Groundwater = 0.015, rivers = 0.0075 and estuaries = 0.0025.

Direct emissions

Synthetic fertilisers

The amount of nitrogen (N) applied to soil by use of synthetic fertilisers is estimated from sales estimates from the Danish AgriFish Agency, the source for the FAO database. Table 6.32 shows the consumption of each fertiliser type. Furthermore, the NH₃ emission factor for each fertiliser is given, based on the values from the EMEP/EEA Guidebook, which has been updated in 2013. The NH₃ emission depends on fertiliser type and the major part of the Danish emission is related to the use of calcium ammonium nitrate and NPK fertiliser, where the emission factor is 0.02 and 0.04 kg NH₃-N per kg N, respectively. The Danish FracGASF is low compared to the IPCC default value. This is due to the small consumption of urea (<1%), which has a high emission factor.

Table 6.32 Synthetic fertiliser consumption 2012 and the NH₃ emission factors.

Fertiliser type	NH ₃ Emission factor ¹ kg NH ₃ -N per kg N	Consumption ² 1000 t N
Calcium and boron calcium nitrate	0.11	0.1
Ammonium sulphate	0.01	5.9
Calcium ammonium nitrate and other nitrate types	0.02	90.8
Ammonium nitrate	0.04	8.1
Liquid ammonia	0.01	7.8
Urea	0.24	0.8
Other nitrogen fertiliser	0.04	22.8
Magnesium fertiliser	0.11	0.0
NPK-fertiliser	0.04	43.0
Diammonphosphate	0.11	0.9
Other NP fertiliser types	0.11	4.7
NK fertiliser	0.04	2.1
Total consumption of N in synthetic fertiliser		187.0
National emission of NH ₃ -N, Gg	5.82	
Average NH ₃ -N emission (FracGASF)	0.03	

¹) EMEP/EEA (2013).

²) The Danish AgriFish Agency (2013).

The use of synthetic fertiliser includes fertiliser used in parks, golf courses and private gardens. 1 % of the synthetic fertiliser can be related to these uses outside the agricultural area.

As a result of increasing requirements for improved use of nitrogen in livestock manure and reduce the nitrogen loss to the environment, the consumption of nitrogen in synthetic fertiliser has more than halved from 1990 to 2012 (Table 6.33). From 2007 to 2008 the consumption increased which is due to stockpiling based on an expectation of rising prices and therefore the consumption decreased again in 2009 to 2012.

Table 6.33 Nitrogen applied as fertiliser to agricultural soils 1990 – 2012.

	1990	1995	2000	2005	2008	2009	2010	2011	2012
N content in synthetic fertiliser, Gg N	400	316	251	206	220	200	190	197	187
NH ₃ -N emission, Gg NH ₃ -N	14	13	8	6	7	6	6	6	6
N in fertiliser applied on soil, Gg N	365	303	243	200	214	194	184	191	181
N ₂ O emission, Gg N ₂ O	7.17	5.96	4.78	3.92	4.20	3.82	3.62	3.75	3.56

Manure applied to soil

The amount of nitrogen applied to soil is estimated as the N-excretion in housings minus the NH₃ emission, which occur in housings, under storage and in relation to the application of manure. These values are based on national estimations and are calculated in the NH₃ emission inventory (Table 6.34). Emission factors for NH₃ from the housing unit and storage are given in Annex 3E-3 and 4. The total N-excretion in housings from 1990 to 2012 has decreased by 8 %. Despite this reduction in N-excretion, the amount of nitrogen applied to soil has increased slightly, which is caused by a reduction in NH₃ emission and especially from application of manure.

Table 6.34 Nitrogen applied as manure to agricultural soils 1990 – 2012.

	1990	1995	2000	2005	2008	2009	2010	2011	2012
N-excretion, housing, Gg N	258	239	235	251	246	237	239	239	237
N ab Storage, Gg N	214	200	197	212	213	206	208	208	207
NH ₃ -N emission from application, Gg NH ₃ -N	31	26	23	17	17	17	17	16	16
N in manure applied on soil, Gg N	183	175	174	195	196	190	191	192	191
N ₂ O emission, Gg N ₂ O	3.59	3.43	3.42	3.83	3.84	3.73	3.75	3.77	3.75

The FracGASM express the fraction of total N-excretion (N ab animal) that is volatilised as NH₃ emission in housings, storage and application. The FracGASM has decreased from 0.25 in 1990 to 0.19 in 2012 (Table 6.35). This is the result of an active strategy to improve the utilisation of the nitrogen in manure.

Table 6.35 FracGASM 1990 – 2012.

	1990	1995	2000	2005	2008	2009	2010	2011	2012
Total N-excretion, Gg N	292	274	270	277	269	260	261	260	258
NH ₃ -N emission from manure, Gg NH ₃ -N	44	42	42	42	43	43	43	43	43
FracGASM	0.25	0.23	0.22	0.20	0.19	0.19	0.19	0.19	0.19

N-fixing crops

To estimate the emission from N-fixing crops, IPCC Tier 1b is applied. The emission calculated is based on nitrogen content, the fraction of dry matter and the yield for each harvest crop type. Data for crop yield is based on data from Statistics Denmark. For nitrogen content in the plants, the data are taken from Danish feedstuff tables (Danish Agricultural Advisory Centre). The estimates for the amount of nitrogen fixed in crops are made by the DCA (Kristensen, 2003, Høgh-Jensen et al., 1998, Kyllingsbæk, 2000).

$$N_2O - N_{N-fix} = \sum (DM_{i, yield} \cdot N_{i, pct} \cdot (1 + N_{i, pct \text{ in root and stubble}}) \cdot A_{pct fix}) \cdot EF_{N_2O}$$

Where:

DM _{i, yield}	= dry matter yield, kg per ha for crop type <i>i</i>
N _{i, pct}	= nitrogen percentage in dry matter
N _{i, pct root + stubble}	= nitrogen percentage in root and stubble
A _{pct fix}	= percentage of nitrogen which is fixed
EF _{N₂O}	= emission factor, IPCC standard value of 1.25 pct.

The Danish inventory includes emissions from clover-grass, despite the fact that this source is not mentioned in the IPCC GPG. Area with grass and clover covered approximately 20 % of the total agricultural area in 2012 and, for this reason, represents an important contributor to the national emission from N-fixing crops.

In Table 6.36 and Annex 3E-13 and 14 the background data for estimating the N-fixing is listed. The emission from N-fixing crops decreases from 1990-2012, largely due to a reduction in agricultural area (Annex 3E-15).

Table 6.36 Emissions from N-fixing crops 2012.

	Crop yield, 1000 t	N-fixing, kg N per tonnes crop yield	N-fixing total, t N fix	N ₂ O emission, Gg N ₂ O
Legumes to maturity ^a	26.5 ^b	37.3	988	0.019
Lucerne	281.2	7.7	2 154	0.042
Crops for silage	206.4 ^e	6.1	1 253	0.025
Legumes/marrow-stem kale	NO	6.1	NO	NO
Grass and clover in rotation	4 166.8 ^e	8.2	34 126	0.670
Grass not in rotation	173.7 ^e	8.2	1 422	0.028
Fields with catch crop	115.7 ^e	8.2	947	0.019
Peas for conservation ^a	9.6	37.3	359	0.007
Seeds of leguminous grass crops			825	0.016
- Red clover	441.6 ^c	200 ^d		
- White clover	4 017.8 ^c	180 ^d		
- Black medic	76.2 ^c	180 ^d		
Total N-fixed			42 075	
Total N ₂ O emission				0.826

^a Dry matter content for straw is 0.87 and the N-fraction is 0.010.

^b Yield of seed, yield of straw is 60 % of yield of seed.

^c Area, ha.

^d kg N per ha.

^e Crop yield given by DSt adjusted to only amount of N-fixing crops

Crop residue

To estimate the emission from crop residue, IPCC Tier 1b is applied. N₂O emissions from crop residues are calculated as the total aboveground quantity of crop residue returned to soil. For cereals, the aboveground residues are calculated as the amount of straw plus stubble and husks. The total amount of straw is given in the annual census and reduced by the amount used for feeding, bedding and bio fuel in power plants. Straw for feed and bedding is subtracted because this quantity of removed nitrogen returns to the soil via manure.

$$N_2O - N_{crop\ residue,j} = \sum_i^N ha_{i,j} \cdot (N_{i, stubble} + N_{i, husks} + N_{i, tops} + N_{i, leafs}) \cdot EF_{N_2O}$$

Where:

i = crop type

j = year

ha = on which the crop is grown

N_i = nitrogen derived from husks, stubble, plant tops and leaf debris, kg ha⁻¹

EF_{N₂O} = emission factor, IPCC standard value of 1.25 %

National values for nitrogen content are used provided by the DCA (Djurhuus and Hansen 2003). It is calculated based on relatively few observations, but is at present the best available data. Data for yield and area cultivated are collected from Statistics Denmark. Background data is given in Annex 3E-16 and 17.

The national emission from crop residues has decreased 14 % from 1990 to 2012 (Table 6.37). This decrease is a result of a fall in cultivated area of beets for feeding, which has been replaced by cultivation of green maize. Another reason is a fall in the agricultural area and a greater part of the straw is harvest – 52 % in 1990 and 63 % in 2012.

Table 6.37 Emissions from crop residue 1990 – 2012.

Crop residue	1990	1995	2000	2005	2008	2009	2010	2011	2012
Stubble	18.9	18.2	18.2	17.6	17.9	17.5	17.9	18.1	17.8
Husks	11.4	11.6	12.0	12.3	11.9	12.4	12.5	12.5	11.7
Top of beets and potatoes	7.1	5.8	5.3	4.9	4.4	4.3	4.3	4.5	4.7
Leafs	6.8	10.3	9.0	9.4	7.9	7.0	7.4	7.1	7.4
Straw	15.1	10.4	10.8	10.2	8.1	10.0	9.7	9.7	9.6
Crop residue, total, Gg N	59.3	56.2	55.3	54.4	50.1	51.2	51.8	51.8	51.1
N ₂ O emission, Gg	1.17	1.10	1.09	1.07	0.98	1.01	1.02	1.02	1.00
Frac _R	0.86	0.85	0.85	0.85	0.86	0.87	0.86	0.86	0.87
Frac _{NCRO}	0.018	0.017	0.017	0.017	0.017	0.017	0.017	0.017	0.017
Frac _{NCRBF}	0.04	0.04	0.04	0.04	0.04	0.04	0.04	0.04	0.04

Frac values

The fractions Frac_{NCRO}, Frac_{NCRBF} and Frac_R are calculated for all years by using the definitions given in the IPCC Reference Manual pp 4.92 – 4.94.

The Frac_{NCRO} and Frac_{NCRBF} are calculated as the N-content in harvest crops divided with the total amount of dry matter in harvest crops. Frac_{NCRBF} covers all N-fixing crops and Frac_{NCRO} all the non N-fixing crops. In Table 6.38 the national calculated fraction values are compared to default values given in the Reference Manual (IPCC, 1997). The national values differ slightly during the years. For all Frac values the Danish values are higher than the IPCC default values. For N-fixing crops the explanation could be that Denmark includes fields with clover grass, which has a high N-content. The higher national Frac_{NCRO} could be a consequence of the relatively large part of straw that is harvested and used for feeding, bedding and fuel. As provided by Statistics Denmark nearly 63 % of the straw in 2012 is harvested (see Annex 3E-18).

For the fractions Frac_R, the unit kg N per kg crop-N is used, as given in the Reference Manual (IPCC, 1997). The fraction is calculated as N-content in the whole above ground crop biomass that is removed from the field as a crop product divided with total N-content in all parts of plants above ground.

The national Frac_R is significantly higher than the IPCC default. The national value express, that 84 % to 87 % of the total N in crops above ground is removed from the field. The remaining is the N-content in straw and tops from beets and potatoes, which are left on the field. From 1990 to 2012 the Frac_R is increased as a consequence of a fall in cultivated area of feeding beets.

Table 6.38 Frac values.

Fractions	Text in CRF Table 4.Ds2 – additional information	Unit	IPCC default values	National Values 1990-2012
Fra _{NCRO}	Fraction of residue dry biomass that is N (all other crops than N-fixing crop)	kg N per kg dm	0.015	0.017-0.018
Fra _{NCRBF}	Fraction of total above-ground biomass of N-fixing crop that is N	kg N per kg dm	0.03	0.04
Fra _{CR}	Fraction of N in the hole above ground crop biomass that is removed from the field as a crop product	kg N per kg crop-N	0.50 ^a	0.84-0.87

^a) IPCC, 2000.

Cultivation of histosols

N₂O emissions from histosols are based on the area with organic soils multiplied by the default emission factor given by the IPCC, 8 kg per ha and constant for all years 1990-2012. The area of histosols is shown in Table 6.39. The area of histosols has decreased from 1990 to 2012, see more in Chapter 7.4.1.

Table 6.39 Area of histosols in ha, 1990-2012.

Year	1990	1995	2000	2005	2008	2009	2010	2011	2012
Area, ha	74 473	69 282	64 092	58 901	55 786	54 748	53 710	52 687	50 886

Other Direct Emissions

The category, "Other", includes emission from sewage sludge and sludge from industries applied to agricultural soils as fertiliser. Information about industrial waste and sewage sludge applied on agricultural soil and the content of nitrogen is obtained from a series of reports published by the Danish Environmental Protection Agency. From 2005 the amount of sewage sludge and N content is based on the information registered in the fertiliser accounts controlled by The Danish AgriFish Agency. The recent official figures regarding the amount of sludge from the industrial waste are data covering year 2002 (Petersen & Kielland, 2003). Data covering year 2008 and 2009 are received from Econet AS (Petersen, 2011) and will be included in an EPA report under preparation. Based on these data the amount in 2003-2007 is interpolated.

It is assumed that 1.9 % of N-input applied to soil volatilises as NH₃, which is based on information from the Danish Environmental Protection Agency (Bielecki, 2002). In Table 6.40 are shown N, emission of NH₃-N and emission of N₂O from sewage sludge and sludge from industries applied to agricultural soils.

Table 6.40 Emission from sludge applied on agricultural soils 1990 – 2012.

	1990	1995	2000	2005	2008	2009	2010	2011	2012
Nitrogen in sewage sludge, t N	3 115	4 635	3 625	2 173	2 394	2 979	2 692	2 592	2 470
Nitrogen in industrial waste, t N	1 529	4 500	5 147	5 509	4 185	3 993	3 942	3 942	3 942
NH ₃ -N emission, t NH ₃ -N	58	87	68	41	45	56	50	49	46
N applied as fertiliser to the soil, t N	4 586	9 048	8 705	7 641	6 534	6 916	6 583	6 485	6 365
N ₂ O emission, Gg N ₂ O	0.09	0.18	0.17	0.15	0.13	0.14	0.13	0.13	0.13

Pasture, Range and Paddock Manure

The amount of nitrogen deposited on grass is based on estimations from the NH₃ inventory. Grazing days is based on expert judgement from the Danish Agricultural Advisory Service. N-excretion on grass has decreased due to a reduction in the number of dairy cattle. Under review by ESD it has been

recommended that the emission of NH₃ from grazing animals is not subtracted before calculation of N₂O.

Table 6.41 Nitrogen excreted on grass 1990 – 2012.

	1990	1995	2000	2005	2008	2009	2010	2011	2012
N-excretion, grass, Gg N	34	36	34	26	23	22	22	21	22
N ₂ O emission, Gg	1.08	1.12	1.08	0.82	0.74	0.70	0.69	0.67	0.68
FracGRAZ	0.12	0.13	0.13	0.09	0.09	0.09	0.08	0.08	0.08

Frac_{GRAZ} is estimated as the volatile fraction from grazing animals compared with the total excreted nitrogen (N ab animal) (Table 6.41). The decrease in Frac_{GRAZ} is due to a decrease in number of dairy cattle and a decrease in grazing days for dairy cattle and heifers.

Indirect emissions

Atmospheric deposition

To estimate the emission from atmospheric deposition, IPCC Tier 1b is applied. Atmospheric deposition includes all agricultural NH₃ emission sources included in the Danish NH₃ emission inventory (Nielsen et al., 2013b). This includes the emission from livestock manure, use of synthetic fertiliser, growing crops, NH₃-treated straw used as feed, field burning of crop residues and sewage sludge plus sludge from industrial production applied to agricultural soils.

The emission from atmospheric deposition has decreased from 1990 – 2012 as a result of the reduction in the total NH₃ emission, from 101 901 tonnes of NH₃-N in 1990 to 60 475 in 2012.

Table 6.42 NH₃ emission 2012.

NH ₃ emission	2012
	t NH ₃ -N
Manure, housing	48 564
Manure, grazing	1 515
Synthetic fertilisers	5 817
Crops	4 448
NH ₃ treated straw	0
Burning of agricultural residues	85
Sewage sludge and industrial sludge	46
Emission total	60 475
N ₂ O emission, Gg	0.95

Nitrogen leaching and Run-off

Nitrogen, which is transported through the soil, can be transformed to N₂O. The IPCC recommends an N₂O emission factor of 0.025 used, of which 0.015 is for leaching to groundwater, 0.0075 for transport to watercourses (in IPCC definition called rivers) and 0.0025 for transport out to sea (in IPCC definition called estuaries). The N₂O emission from nitrogen leaching is a sum of the emission for all three parts calculated as:

$$N_2O_{\text{leaching}} = (N_{\text{leach-ground}} \cdot EF_{\text{ground}} + N_{\text{leach-rivers}} \cdot EF_{\text{rivers}} + N_{\text{leach-estuaries}} \cdot EF_{\text{estuaries}}) \cdot \frac{44}{28}$$

In the Action Plans for the Aquatic Environment, nitrogen leaching to groundwater, rivers and estuaries has been estimated, see Table 6.43. The calculation of N to the groundwater is based on two different models–SKEP/Daisy and N-LES (Børgesen & Grant, 2003) carried out by DCA and

DCE, Aarhus University (see overview of model in Annex 3E Figure 1). SKEP/DAISY is a dynamical crop growth model taking into account the growth factors, whereas N-LES is an empirical leaching model based on more than 1500 leaching studies performed in Denmark during the last 15 years. The models produce rather similar results for nitrogen leaching on a national basis (Waagepetersen et al., 2008). The SKEP/Daisy model has estimated the total N leached from 2003-2007 to be 172-159 thousand tonnes N, whereas N-LES model has estimated the total N leached to be 163-154 thousand tonnes in the same period. An average of the results from the two models is used in the emission inventory.

Data concerning the N-leaching to rivers and estuaries are based on data from NOVANA (National Monitoring program of the Water Environment and Nature) received from the department of Bioscience, Aarhus University (Windorf et al., 2011). NOVANA is a monitoring program which includes monitoring of the ecologic, physic and chemical condition of water areas and transport of water and a range of substances, including N, to lakes and the sea (Wiberg-Larsen et al., 2010). These studies include measurements from 223 monitoring stations in all parts of Denmark and have been going on from the early 1990's.

Table 6.43 N leaching to groundwater, rivers and estuaries in Gg, 1990-2012.

	1990	1995	2000	2005	2008	2009	2010	2011	2012
Groundwater	267	235	179	160	163	154	151	153	149
Rivers	102	104	95	67	80	59	68	73	74
Estuaries	100	91	81	56	65	49	55	59	59

Figure 6.6 shows leaching from groundwater estimated in relation to the nitrogen applied to agricultural soils as livestock manure, synthetic fertiliser and sludge. The average proportion of nitrogen leaching from groundwater has decreased from around 39 % in the middle of the nineties to around 33 % in 2012. The decline is due to implementation of measures to avoid the nitrogen surplus in the agricultural production by improved nitrogen in manure, to use catch crops during winter and ban application of manure in winter. The reduction in nitrogen applied is particularly due to the fall in the use of synthetic fertiliser, which has been reduced by 50 % from 1990 to 2012.

The proportion of N input to soils lost through leaching and runoff (F_{CLEACH}) used in the Danish emission inventory is higher than the default value of the IPCC (30 %). The high values are partly due to the humid Danish climate, with the precipitation surplus during winter causing a downward movement of dissolved nitrogen. F_{CLEACH} has decreased from 1990 and onwards. At the beginning of 1990s, manure was often applied in autumn. Now the main part of manure application takes place in the spring and early summer, where there is nearly no downward movement of soil water. The decrease in F_{CLEACH} over time is due to increasing environmental requirements and banning manure application after harvest. The data based on model estimates from DCA and DCE reflect the Danish conditions and are considered the best estimate.




Figure 6.6 Nitrogen applied to agricultural soils and N-leaching, groundwater 1990-2012.

6.5.3 Activity data

The calculation of the N_2O emission is based on the amount of nitrogen applied to agricultural soils, which is the activity data. Table 6.44 provides an overview on activity data from 1990 to 2012 used to calculate the N_2O emission from agricultural soils. The amount of nitrogen applied to agricultural soil has decreased from 1 081 Gg N to 702 Gg N, corresponding to a 35 % reduction, which results in a lower N_2O emission.

Table 6.44 Activity data - agricultural soils 1990 – 2012, Gg N.

CRF – Table 4.D	1990	1995	2000	2005	2008	2009	2010	2011	2012
Total amount of nitrogen applied on soil	1 081	937	810	746	752	721	708	719	702
1. Direct Emissions									
Synthetic Fertilizers	387	303	243	200	214	194	184	191	181
Animal Manure Applied to Soils	183	175	174	195	196	190	191	192	191
N-fixing Crops	44	37	38	34	35	41	39	42	42
Crop Residue	59	54	55	54	50	51	52	52	51
Industrial Waste	2	5	5	6	4	4	4	4	4
Sewage Sludge	3	5	4	2	2	3	3	3	2
2. Pasture, Range and Paddock	34	36	34	26	23	22	22	21	22
3. Indirect Emissions									
Atmospheric Deposition	102	88	77	69	65	62	62	62	60
Nitrogen Leaching and Run-off	267	235	179	160	163	154	151	153	149

6.5.4 Time series consistency

The N_2O emissions from agricultural soils have been reduced by 35 % from 1990 to 2012. This is largely due to a decrease in the use of synthetic fertiliser and a decrease in N-leaching as a result of national environmental policy, where action plans have focused on decreasing the nitrogen losses and on improving the nitrogen utilisation in manure.

Table 6.45 Emissions of N₂O from Agricultural Soils 1990 – 2012, Gg N₂O.

CRF – Table 4.D	1990	1995	2000	2005	2008	2009	2010	2011	2012
Total N ₂ O emission	24.81	21.86	18.96	17.07	17.34	16.37	16.20	16.50	16.14
<u>1. Direct Emissions</u>	14.24	12.23	11.02	10.39	10.54	10.18	9.96	10.16	9.90
Synthetic Fertilisers	7.59	5.96	4.78	3.92	4.20	3.82	3.62	3.75	3.56
Animal Manure Applied to Soils	3.59	3.43	3.42	3.83	3.84	3.73	3.75	3.77	3.75
N-fixing Crops	0.87	0.73	0.75	0.67	0.69	0.80	0.77	0.83	0.83
Crop Residue	1.17	1.06	1.09	1.07	0.98	1.01	1.02	1.02	1.00
Cultivation of Histosols	0.94	0.87	0.81	0.74	0.70	0.69	0.68	0.66	0.64
Industrial Waste	0.03	0.09	0.10	0.11	0.08	0.08	0.08	0.08	0.08
Sewage Sludge	0.06	0.09	0.07	0.04	0.05	0.06	0.05	0.05	0.05
<u>2. Pasture, Range and Paddock</u>	1.08	1.12	1.08	0.82	0.74	0.70	0.69	0.67	0.68
<u>3. Indirect Emissions</u>	9.50	8.51	6.86	5.86	6.06	5.49	5.55	5.66	5.56
Atmospheric Deposition	1.60	1.38	1.21	1.09	1.01	0.97	0.98	0.97	0.95
Nitrogen leaching and Run-off	7.89	7.13	5.66	4.77	5.04	4.52	4.57	4.70	4.61

6.6 Field burning of agricultural residues (CRF sector 4F)

Field burning of agricultural residues has in Denmark been prohibited since 1990 and may only take place in connection with production of grass seeds on fields with repeated production and in cases of wet or broken bales of straw. The amount of burnt straw from the grass seed production is estimated as 15 % of the total amount produced. The amount of burnt bales of broken or wet bales of straw is estimated as 0.1 % of total amount of straw. Both estimates are based on an expert judgement by the Danish Agricultural Advisory Service. The total amounts are based on data from Statistics Denmark.

From field burning is seen emissions of a series of different compounds and related to GHG emissions of the following compounds are estimated CH₄, N₂O, NO_x, CO, CO₂, SO₂ and NMVOC. For emission of NMVOC see Chapter 6.7. The emission of NO_x and CO is given CRF Table 4s2. Emission of CO₂ (biogenic) and SO₂ is estimated, but not reported because this is not possible in CRF tables in the present format. Equation for calculating emission of various compounds:

$$E = BB \cdot \frac{EF}{1000000} \cdot FO$$

$$BB = CP \cdot FB \cdot FR_{dm}$$

E = emission of compounds, Gg
 BB = total burned biomass, Gg dm
 CP = crop production, t
 FB = fraction burned in fields
 FR_{dm} = dry matter fraction of residue
 EF = emission factor, g per kg dm
 FO = fraction oxidized

Table 6.46 Factors for estimating emissions of CH₄ and N₂O, 2012.

		Crop production	Fraction burned in fields	Dry matter (dm) fraction of residue	Total Biomass burned	EF	Fraction oxidized	Emission
		t			Gg dm	g per kg dm		Gg
CH ₄	Mixed cereals	5 803 600	0.001	0.85	4 933	2.7	0.90	0.012
CH ₄	Straw from seeds of grass	334 675	0.15	0.85	42 671	2.7	0.90	0.104
N ₂ O	Mixed cereals	5 803 600	0.001	0.85	4 933	0.07	0.90	0.0003
N ₂ O	Straw from seeds of grass	334 675	0.15	0.85	42 671	0.07	0.90	0.003
Total CO ₂ eqv								3.36

The emission of CH₄, N₂O, NO_x, CO, CO₂ and SO₂ from field burning contributes with less than 1 % of the national emission.

The fraction value Frac_{BURN} is calculated by using the definitions as given in IPCC Reference Manual pp. 4.92 – 4.94. Frac_{BURN} is calculated as the amount of N in burned straw divided with the total amount of N in crop residue and the fractions are given in kg N per kg crop-N. For all years the value of Frac_{BURN} is around 0.01 kg N per kg crop-N, which is low, compared to IPCC default value. This is due to the prohibition of field burning in Denmark.

6.7 NMVOC emission

Around 3 % of the total NMVOC emission originates from the agricultural sector, which, in the Danish emission inventory, includes emission from agricultural soils, such as arable land crops and grassland, and field burning of agricultural residue. Activity data are obtained from Statistics Denmark. The emission factor for agricultural soils is for land with arable crops 393 g NMVOC per ha and for grassland, 2120 g NMVOC per ha (Fenhann & Kilde, 1994 and Priemé & Christensen, 1991). The IPCC default value for the emission factor for field burning of agricultural residue is used. The emission from agricultural soils contributes with 88 % and field burning with 12 % of the agricultural NMVOC emission in 2012.

No significant changes from 1990 to 2012 have taken place. A decrease in the area of arable crops offsets an increase in the area of grassland.

Table 6.47 Area and NMVOC emission from agricultural soils 1990 – 2012.

	1990	1995	2000	2005	2008	2009	2010	2011	2012
Arable crops, 1000 ha	2 322	2 064	2 043	2 086	2 107	2 103	2 096	2 102	2 102
Grassland, 1000 ha	466	446	413	446	490	498	521	516	527
NMVOC emission, Gg	1.90	1.76	1.68	1.77	1.87	1.88	1.93	1.92	1.94

Table 6.48 NMVOC emission from field burning of agricultural residue 1990 – 2012.

	1990	1995	2000	2005	2008	2009	2010	2011	2012
NMVOC emission, Gg	0.20	0.24	0.30	0.33	0.27	0.32	0.23	0.23	0.27

6.8 Uncertainties

Uncertainties are calculated using both a Tier 1 and a Tier 2 approach; see Chapter 1.7 for a description of the Tier 2 methodology. The same uncertainty values for activity data and emission factors are used for both Tier 1 and Tier 2.

6.8.1 Uncertainty values

Uncertainties regarding animal production, such as number of animals, feeding consumption, normative figures etc., are very small. The number of animals is estimated by Statistics Denmark and all cattle, sheep and goats have their own ID-number (ear tags) and, hence, uncertainty with regard to their numbers is almost non-existing. Statistics Denmark has estimated the uncertainty in the number of swine to be less than 1 %.

The Danish Normative System for animal excretions is based on data from the Danish Agricultural Advisory Service (DAAS), which is the central office for all Danish agricultural advisory services. DAAS engages in a great deal of research as well as the collection of efficacy reports from Danish farmers for dairy production, meat production, swine production, etc. to optimise productivity in Danish agriculture. In total, feeding plans from 15-18 % of Danish dairy production, 25-30 % of swine production, 80-90 % of poultry production and approximately 100 % of fur production are collected annually. These basic feeding plans are used to develop the standard values of the "Danish Normative System".

The normative figures (Poulsen et al. 2001) are comprised of arithmetic means. Based on feeding plans, the standard deviation in N-excretion rates between farms can be estimated to ± 20 % for all animal types (Poulsen, DCA). However, due to the large number of farms included in the norm figures the arithmetic mean can be assumed as a very good estimate with a low uncertainty.

Data for hectares under cultivation is estimated by Statistics Denmark and the uncertainties are based on their estimates. For the most common crops the uncertainties are below 5 %.

In the 2011 submission the uncertainty estimates for both activity data and emission factors were re-evaluated and some adjustments were made. For CH₄ emission from animals (CRF category 4.A and 4.B) the uncertainties for activity data are lowered due to the combined effect of low uncertainty in actual animal numbers, relatively low uncertainty for feed consumption and excretion rates, this gives a relatively low uncertainty in the activity data as a whole – between 2% and 22%. The uncertainties for the emission factors for CH₄ emission from animals are adjusted based on default values from the IPCC (1997 and 2000).

For the N₂O emission uncertainties, the activity data uncertainty is based on the uncertainties for NH₃ emission due to the high correlation between the NH₃ and N₂O emission (Nielsen et al, 2013b). Uncertainties related to the N₂O emission factor are based on the IPCC Good Practice Guidance. See Table 6.49 for uncertainty values for the agricultural sector.

Table 6.49 Uncertainties values for activity data and emission factors for CH₄ and N₂O.

CRF category	Emission factor	Uncertainties value for activity data, %	Uncertainties value for emission factor, %
<u>4.A Enteric Fermentation</u>	CH ₄	2	20
<u>4.B Manure Management</u>	CH ₄	5	20
	N ₂ O	22	100
<u>4.D Agricultural Soils</u>			
4.D1 Direct soil emissions			
Synthetic Fertilisers	N ₂ O	25	100
Animal Manure Applied to Soils	N ₂ O	30	100
N-fixing Crops	N ₂ O	20	100
Crop Residue	N ₂ O	20	100
Cultivation of Histosols	N ₂ O	20	100
Sewage sludge used as fertiliser	N ₂ O	20	100
Industrial waste used as fertiliser	N ₂ O	20	100
4.D2 Pasture, range and paddock	N ₂ O	25	100
4.D3 Indirect soil emissions			
Atmospheric Deposition	N ₂ O	19	100
N-Leaching and Run-off	N ₂ O	20	100
<u>4.F Field Burning of Agricultural Residue</u>			
	CH ₄	25	50
	N ₂ O	25	50

6.8.2 Result of the uncertainty calculation

Table 6.50 shows the result of the Tier 1 and Tier 2 uncertainty calculation for 2012. A calculation of 1990 gives nearly the same uncertainty values as for 2012, for all emission sources. The overall uncertainty calculation for the agricultural sector based on Tier 1 is estimated to $\pm 25\%$. Tier 2 calculation shows an uncertainty interval from -19% to $+32\%$.

For most of the emission sources the uncertainty level based on Tier 2 are nearly at the same level as for Tier 1, see Figure 6.7. The two calculations can be considered as consistent. The lowest uncertainties are seen for CH₄ emission from enteric fermentation and manure management and the highest for emission from grazing animals and this pattern is reflected in both calculations.

The biggest difference between the Tier 1 and Tier 2 uncertainty calculations is seen for N₂O from manure management.

Table 6.50 Comparison between Tier 1 and Tier 2 uncertainty calculation, 2012.

Uncertainty		Tier 1		Tier 2		
		Emission, Gg CO ₂ eqv	Uncertainty, % Lower and upper (±)	Median emission, Gg CO ₂ eqv	Uncertainty, %	
					Lower (-)	Upper (+)
4 Agriculture total	CH ₄ and N ₂ O	9 599	25	10 132	19	32
4.A Enteric Fermentation	CH ₄	2 904	20	2 911	12	14
4.B Manure Management	CH ₄ and N ₂ O					
	CH ₄	1 297	21	1 302	11	13
	N ₂ O	391	102	424	20	27
4.D Agricultural soil:	N ₂ O					
4.D1 Direct soil emissions	N ₂ O	3 069	56	3 305	40	76
4.D2 Pasture, Range and Paddock Manure	N ₂ O	211	103	212	62	161
4.D3 Indirect soil emissions	N ₂ O	1 724	86	1 766	54	130
4.F Field Burning of Agricultural Residues	CH ₄ and N ₂ O					
	CH ₄	2	56	2	42	70
	N ₂ O	1	56	1	42	70




Figure 6.7 Tier 1 and Tier 2 uncertainties for the agricultural sector, 2012.

6.9 Quality assurance and quality control (QA/QC)

A first step of development and implementation of a general QA/QC plan for all sectors started in 2004 which is described in a publicised manual (Sørensen et al., 2005). The manual describes the concepts of quality work and how to handle quality management by using Critical Control Points and a list of Point of Measurements (Nielsen et al, 2013a). For more detailed in-

formation of the structure in the general QA/QC plan refers to Chapter 1.6 for QA/QC.

A complete list Points of Measures (PM) are given in Table 1.2. PM related to the agricultural inventory is listed below in Chapter 6.9.2 and are primarily connected to data storage and data processing level 1. For PM not mentioned below please refer to Chapter 1.6.

The QA/QC work specific for the agricultural sector is still improved. The overall framework regarding a QA/QC plan for agriculture are constructed in form of six stages and each stage focus on quality assurance and quality check in different part of the inventory process. This year a more detailed set up for stage I, II and III are developed – refer to Annex 3E-19.

The QA/QC procedure is divided in six stages as listed below:

Table 6.51 Stages of QA/QC procedure.

Stage I	Check of input data
	- check of data input in IDA are consistent with data from external data suppliers
Stage II	Check of IDA data – overall
	- check of recalculations for total emissions compared with the latest submission (2012)
	- check of total emissions for the total CO ₂ eqv. and for each compound
Stage III	Check of IDA data – specific
	- check of annual changes of activity data, emission factors, IEF and other important variables as GE, Nex, housing system distribution, grazing days
Stage IV	Check by comparing calculation with estimates from other institutions
	- the total Nex for all livestock production estimated by DCA
	- the Register for fertilization controlled by the Danish AgriFish Agency
Stage V	Check of data registered in CRF
	- compare data in CRF with data from IDA
Stage VI	Check of the inventory in general (external review)
	- check that data is used correctly
	- check the methodology and the calculations

Stage I: Check of input data

At stage I, it is checked that all input data in IDA are consistent with data from the external data suppliers. Data from the Statistics Denmark have to be checked for the livestock production, slaughter data for poultry and pigs, check of land use and crop yield. Data input from the DCA have to be checked for feed intake, N-excretion, manure production, dry matter content and grazing days. Data from the Danish AgriFish Agency: distribution of housing systems and the use of nitrogen in synthetic fertiliser.

Stage II: Check of IDA data - overall

Stage II includes check of the overall calculations in IDA, where the first step is to compare the inventory with the last reported emission inventory - submission 2013. In the case where an error cover the whole time series, it can be difficult to identify this error by checking the changes in inter-annual values. Therefore, a check of recalculations is needed.

Next step in stage II is a check of total emissions of CH₄, N₂O, NMVOC and the other compounds, which are related to the field burning of agricultural residues. For each compound a check of trends of time series 1990-2012 and

inter-annual changes is provided. Significant jumps or dips from one year to another could indicate an error - otherwise it has to be explained.

Stage III: Check of IDA data - specific

At stage III, a check of specific variables in IDA is provided for both inter-annual changes and trends for the entire time series. Variables includes activity data, emission factors, IEFs and other important key variables such as feed intake, GE, Nex and housing system distribution.

Stage IV: Check by comparing calculation with estimates from other institutions

The purpose of stage IV is to verify the calculations in IDA, as far as external data estimations are available. For other purposes DCA for some years calculate the overall N excretion from the total livestock production in DK, which could be compared with the survey given in the emission inventory. Another possibility to check some of the IDA estimations is the information in the fertiliser accounts controlled by The Danish AgriFish Agency. Farmers with more than 10 animal units have to be registered and have to keep accounts of the N content in manure, received manure or other organic fertiliser. These comparisons will properly show some differences, which not necessarily indicate an error, but the most important cause of the difference has to be identified.

Stage V: Check of data registered in CRF

Stage V primarily focuses on the last reported year 2012 and the base year (1990), where all activity data, emissions and IEFs are checked. Furthermore, CRF sum emissions are checked with sum emissions in IDA. If an error is detected a more detailed check is done to find the reason for the error.

Stage VI: Check of the inventory in general

A detailed description of the methodology used to calculate the Danish agricultural emissions is published as a sectorial report for agriculture (Mikkelsen et al., 2011). General checks of the inventory include considerations of which data input is used, how they are used in the calculations and whether more accurate data are available. The review of the sectorial report addresses these issues and is a most valuable part of the QA of the agricultural sector.

Status for the QA/QC plan

The framework for working out a specific QA/QC plan for the agricultural sector is complete. Stage I-III is done as part of the process of inventory preparation, which has reduced the number of errors in the CRF and in this way meet the ERT recommendations. A more detailed list showing the checked variables of stage I – III is provided in Annex 3E-19.

Concerning the stage IV we have provide some random checks but need to provide a more systematic check. We are aware of some external calculations which can be compared with the estimations in IDA – e.g. total N-excretion in manure calculated of DCA. Furthermore, some comparisons with the Register of Fertilisation administrated by the Danish AgriFish Agency can be provided.

Stage VI is implemented. Two reports describing the methodology in calculation of agricultural emissions in details are published (Mikkelsen et al., 2006 and Mikkelsen et al., 2011). Both reports have been reviewed by experts not involved with the preparation of the emission inventory. The 2011 report was reviewed by: Nicholas J. Hutchings from the DCA, Aarhus University

and Johnny M. Andersen from the Faculty of Life Sciences, University of Copenhagen. The reviewers have reviewed all sections of the report. In the next updated version it is planned to contact relevant reviewers to focus on specific subject areas. An updated version of the methodology report is planned to take place in 2013/2014.

6.9.1 QA/QC plan expressed in Critical Control Points and Point of Measurements

Data storage level 1

Data Storage level 1	3. Completeness	DS.1.3.1	Documentation showing that all possible national data sources are included by setting down the reasoning behind the selection of datasets.
----------------------	-----------------	----------	--

The following external data are in used in the agricultural sector, in more details see Table 6.2:

- Data from the annual agricultural census made by Statistics Denmark.
- DCA, Aarhus University.
- The Danish AgriFish Agency.
- Danish Agricultural Advisory Service (DAAS).
- The Danish Energy Agency.
- Danish Environmental Protection Agency.

The emission factors come from various sources:

- IPCC guidelines.
- DCA, Aarhus University: NH₃ emission, CH₄ emission from enteric fermentation and manure management.

Statistics Denmark

The agricultural census made by Statistics Denmark is the main supply of basic agricultural data. In Denmark, all cattle, sheep and goats have to be registered individually and hence the uncertainty in the data is negligible. For all other animal types, farms having more than 10 animal units are registered.

DCA

The DCA is responsible for the delivery of N-excretion data for all animal and housing types. Data on feeding consumption on commercial farms are collected annually by DAAS from on-farm efficacy controls. For dairy cattle, data is collected from 15-20 % of all farms, for pigs, 25-30 % and for poultry and mink, 90-100 % of all farms. The farm data are used to calculate average N-excretion from different animal and housing types. Due to the large amount of farm data involved in the dataset, N-excretion is seen as a very good estimate for average N-excretion at the Danish livestock production.

Danish AgriFish Agency

Total area with the various agricultural crops is provided to the Danish AgriFish Agency via the agricultural subsidy system. For every parcel of land (via a vector-based field map with a resolution of >0.01 ha), the area planted with different crops is reported. If the total crop area within a parcel is larger than the parcel area, a manual control of the information is performed by the Agency. The area with different crops, therefore, represents a very precise estimate.

All farmers are obligated to do N-mineral accounting on a farm and field level with the N-excretion data from DCA. Data at farm level is reported annually to the Danish AgriFish Agency. The N figures also include the quantities of synthetic fertilisers bought and sold. Suppliers of synthetic fertilisers are required to report all N sales to commercial farmers to the Agency. The total sold to farmers is very close to the amount imported by the suppliers, corrected for storage. The total amount of synthetic fertiliser in Denmark is, therefore, a very precise estimate for the synthetic fertiliser consumed. This is also valid for N-excretion in animal manure.

The Danish AgriFish Agency, as the controlling authority, performs analysis of feed sold to farmers. On average, 1600 to 2000 samples are analysed every year. Uncertainty in the data is seen as negligible. The data are used when estimating average energy in feedstuffs for pigs, poultry, fur animals, etc.

From 2005 the Danish AgriFish Agency provides data for distribution of housing type.

Danish Agricultural Advisory Service (DAAS)

DAAS is the central office for all Danish agricultural advisory services. DAAS carries out a considerable amount of research itself, as well as collecting efficacy reports from the Danish farmers for dairy production, meat production, pig production, etc., to optimise productivity in Danish agriculture. From DAAS data on housing type until 2004, grazing situation and information on application of manure is received.

The Danish Energy Agency

The amount of slurry treated in biogas plants is received from the Danish Energy Agency.

Danish Environmental Protection Agency

Information on the sludge from waste water treatment and the manufacturing industry and the amount applied on agricultural soil is obtained from the Danish Environmental Protection Agency.

Data Storage level 1	1. Accuracy	DS.1.1.1	General level of uncertainty for every dataset including the reasoning for the specific values
----------------------	-------------	----------	--

The most important emission source is related to the animal production. Uncertainty for the animal data is very low due to the very strict environmental laws in Denmark. Standard deviation regarding the numbers of cattle and pigs has been estimated to <0.7 %. For poultry the standard deviation is <2.1 %. For all years, 25-35 % of all holdings are included in the census. The standard deviation for N-excretion between farms is reported as 25 % for dairy cattle and pigs, but due to the large numbers involved in the estimation of the average N-excretion, the average is assumed to be a precise estimate for the Danish agricultural efficacy level.

Regarding uncertainties for the remaining emission sources see Chapter 6.8.

Data Storage level 1	1. Accuracy	DS.1.1.2	Quantification of the uncertainty level of every single data value including the reasoning for the specific values.
----------------------	-------------	----------	---

Please, refer to Chapter 6.8 and Table 6.50.

Data Storage level 1	1. Comparability	DS.1.2.1	Comparability of the data values with similar data from other countries, which are comparable with Denmark, and evaluation of discrepancy.
----------------------	------------------	----------	--

The Danish N-excretion levels are generally lower than IPCC default values. This is due to the highly skilled, professional and trained farmers in Denmark, with access to a highly competent advisory system.

The feed consumption per animal is in line with similar data from Sweden, although they are not quite comparable because Denmark is using feeding units (FE) which cannot easily be converted to energy content. Earlier, one feeding unit was defined as one kg of barley. Today, the calculations are more complicated and depend on animal type.

Data Storage level 1	4. Consistency	DS.1.4.1	The origin of external data has to be preserved whenever possible without explicit arguments (referring to other PMs).
----------------------	----------------	----------	--

External data received are stored in the original format in quality management database system.

Data Storage level 1	6. Robustness	DS.1.6.1	Explicit agreements between the external institution holding the data and DCE about the conditions of delivery.
----------------------	---------------	----------	---

DCE has established formal data agreements with all institutes and organisations which deliver data, to assure that the necessary data is available to prepare the inventory on time.

Data Storage level 1	6. Robustness	DS.1.6.2	At least two employees must have a detailed insight into the gathering of every external data set.
----------------------	---------------	----------	--

Please refer to Chapter 1.7.

Data Storage level 1	7. Transparency	DS.1.7.1	Summary of each dataset including the reasoning for selecting the specific dataset.
----------------------	-----------------	----------	---

Please refer to DS 1.1.1.

Data Storage level 1	7. Transparency	DS.1.7.2	The archiving of data sets needs to be easy accessible for any person in the emission inventory.
----------------------	-----------------	----------	--

Please refer to Chapter 1.7.

Data Storage level 1	7. Transparency	DS.1.7.3	References for citation for any external data set have to be available for any single value in any dataset.
----------------------	-----------------	----------	---

A great deal of documentation already exists in the literature list, and also achieved in the quality management database system.

Data Storage level 1	7. Transparency	DS.1.7.4	Listing of external contacts for every dataset.
----------------------	-----------------	----------	---

Statistics Denmark:

Mrs. Mona Larsen (mla@dst.dk)

Mr. Karsten K. Larsen (kkk@dst.dk)

DCA (Aarhus University):

Mrs. Hanne Damgaard Poulsen (hannedamgaard.poulsen@agrsci.dk)

Mr. Nick Hutchings (nick.hutchings@agrsci.dk)

Mr. Christen Duus Børgesen (christen.Borgesen@agrsci.dk)

The Danish Agricultural Advisory Centre:

Mr. Ole Aaes (oes@vfl.dk)

Mr. Eric F. Clausen (efc@vfl.dk)

Mr. Barthold Feidenshans'l (baf@vfl.dk)

Danish AgriFish Agency:

Mr. Troels Knudsen (tkn@pdir.dk)

Mrs. Mette Thomsen (mth@pdir.dk)

The Danish Energy Agency:

Mr. Søren Tafdrup (st@ens.dk)

The Danish Environmental Protection Agency:

Mrs. Linda Bagge (bagge@mst.dk)

Data processing level 1

Data Processing level 1	1. Accuracy	DP.1.1.1	Uncertainty assessment for every data source as input to Data Storage level 2 in relation to type of variability. (Distribution as: normal, log normal or other type of variability).
-------------------------	-------------	----------	---

The Tier 1 methodology is used to calculate the uncertainties for the agricultural sector. The uncertainties are based on a combination of IPCC guidelines and expert judgement (Olesen et al., 2001, Poulsen et al., 2001) and a normal distribution is assumed. A Tier 2 calculation is provided, please refer to Chapter 6.8.

Data Processing level 1	1. Accuracy	DP.1.1.2	Uncertainty assessment for every data source as input to Data Storage level 2 in relation to scale of variability (size of variation intervals).
-------------------------	-------------	----------	--

Please refer to DP 1.1.1.

Data Processing level 1	1. Accuracy	DP.1.1.3	Evaluation of the methodological approach using international guidelines.
-------------------------	-------------	----------	---

Denmark has worked out a report with a more detailed description of the methodological inventory approach in Mikkelsen et al. (2006) and an updated version in Mikkelsen et al. (2011). The first report has been reviewed by the Statistics Sweden, who is responsible for the Swedish agricultural inventory and the updated report has been reviewed of qualified persons with comprehensive agricultural knowledge; Nicholas J. Hutchings from the DCA, Aarhus University and Johnny M. Andersen from the Faculty of Life Sciences, University of Copenhagen. None of the reviewers is involved in the preparation of the annual inventory.

Furthermore, data sources and calculation methodology developments are continuously discussed in cooperation with specialists and researchers in different institutes and research sections. As a consequence, both the data and methods are evaluated continually according to the latest knowledge and information.

Data Processing level 1	1. Accuracy	DP.1.1.4	Verification of calculation results using guideline values
-------------------------	-------------	----------	--

The methodological approach is consistent with the IPCC Reference Manual and the Good Practice Guidance.

Implied emission factors for CH₄ from enteric fermentations are, in general, lower than the IPCC default values, which is due to the professional way farms are managed in Denmark. There has been an increase in emission from the enteric fermentation of CH₄ due to increase in milk production. The IEF is in line with Canada, the Netherlands, Sweden and Finland, who all have agricultural conditions comparable to Denmark.

The IEF for CH₄ from manure management is, in general, higher compared with the default IPCC values for Western Europe because of the higher percentage handled as slurry. However, due to the high efficiency at farm level, energy intake is lower per head and the subsequent CH₄ emission from slurry is, thereby, lower. Denmark uses an MCF factor of 10 % as provided in the 1996 guidelines and not the 39 % in the revision to the 1996 guidelines. For further explanation, see Chapter 6.4.2.

Frac_{LEACH} is higher than the default IPCC values. Frac_{LEACH} has decreased from 1990 and onwards. In the beginning of 1990s, manure was often applied in autumn. The high values are partly due to the humid Danish climate, with the precipitation surplus during winter causing a downward moment of dissolved nitrogen. The decrease in Frac_{LEACH} over time is caused by sharpened environmental requirements, banning manure application after harvest. As a result, most manure application occurs during spring and summer, where there is a precipitation deficit. The generally accepted leaching values in Denmark are 0.3 for mineral nitrogen and 0.45 for organic-bound nitrogen. These values are based on a series of leaching studies.

Data Processing level 1	2. Comparability	DP.1.2.1	The inventory calculation has to follow the international guidelines suggested by UNFCCC and IPCC.
-------------------------	------------------	----------	--

The methodological approach is consistent with the IPCC Reference Manual and the Good Practice Guidance.

Data Processing level 1	3. Completeness	DP.1.3.1	Assessment of the most important quantitative knowledge which is lacking.
-------------------------	-----------------	----------	---

Regarding the reduction potential for biogas treated slurry, more information and investigation would be preferred. There is on-going work to increase the accuracy of this emission source.

Data Processing level 1	3. Completeness	DP.1.3.2	Assessment of the most important missing accessibility to critical data sources
-------------------------	-----------------	----------	---

All known major sources are included in the inventory. In Denmark, only very few data are restricted. Accessibility is not a key issue; it is more lack of data.

Data Processing level 1	4. Consistency	DP.1.4.1	In order to keep consistency at a high level, an explicit description of the activities needs to accompany any change in the calculation procedure
-------------------------	----------------	----------	--

The calculation procedure is consistent for all years.

Data Processing level 1	4. Consistency	DP.1.4.2	Identification of parameters (e.g. activity data, constants) that are common to multiple source categories and confirmation that there is consistency in the values used for these parameters in the emission calculations
-------------------------	----------------	----------	--

Please refer to Chapter 1.7.

Data Processing level 1	5. Correctness	DP.1.5.1	Show at least once, by independent calculation, the correctness of every data manipulation.
-------------------------	----------------	----------	---

During the development of the model, thorough checks have been made by all persons involved in preparation of the agricultural section.

Data Processing level 1	5. Correctness	DP.1.5.2	Verification of calculation results using time series.
-------------------------	----------------	----------	--

Time series for activity data, emission factors and national emission are performed to check consistency in the methodology, to avoid errors, to identify and explain considerable year to year variations.

Data Processing level 1	5. Correctness	DP.1.5.3	Verification of calculation results using other measures.
-------------------------	----------------	----------	---

A comparison between IPCC Tier 2 method for enteric fermentation and Denmark's Tier 2/CS is made, see Chapter 6.3.4.

Data Processing level 1	5. Correctness	DP.1.5.4	Show one-to-one correctness between external data sources and the databases at Data Storage level 2
-------------------------	----------------	----------	---

In the database key ids is used to identify the unique data. The data on DS level 1 is linked to the key id used in the database so a clear reference from DS level 1 to higher levels of both DP and DS is secured.

Data Processing level 1	6. Robustness	DP.1.6.1	Any calculation must be anchored to two responsible persons that can replace each other in the technical issue of performing the calculations.
-------------------------	---------------	----------	--

Please refer to Chapter 1.7.

Data Processing level 1	7. Transparency	DP.1.7.1	The calculation principle and equations used must be described.
-------------------------	-----------------	----------	---

All calculation principles are described in the NIR and the documentation report (Mikkelsen et al., 2011).

Data Processing level 1	7. Transparency	DP.1.7.2	The theoretical reasoning for all methods must be described.
-------------------------	-----------------	----------	--

All theoretical reasoning is described in the NIR and the documentation report (Mikkelsen et al., 2011).

Data Processing level 1	7. Transparency	DP.1.7.3	Explicit listing of assumptions behind methods.
-------------------------	-----------------	----------	---

All theoretical reasoning is described in the NIR and the documentation report (Mikkelsen et al., 2011).

Data Processing level 1	7. Transparency	DP.1.7.4	Clear reference to dataset at Data Storage level 1.
-------------------------	-----------------	----------	---

In the database key ids is used to identify the unique data. The data on DS level 1 is linked to the key id used in the database so a clear reference from DS level 1 to higher levels of both DP and DS is secured.

Data Processing level 1	7. Transparency	DP.1.7.5	A manual log to collect information about recalculations.
-------------------------	-----------------	----------	---

Changes compared with the last emissions report are described in the NIR and the national emission changes is given in a table under the section, "Recalculation". The text describes whether the change is caused by changes in the dataset or changes in the methodology used. Furthermore a log table is filled in when data are updated or adjusted continuously.

Data storage and processing level 2

For point of measurements not mentioned below please refer to Chapter 1.7.

Data Storage level 2	5. Correctness	DS.2.5.1	Documentation of a correct connection between all data types at level 2 to data at level 1.
----------------------	----------------	----------	---

A manual check-list is under development for correct connection between all data types at level 1 and 2.

Data Processing level 2	5. Correctness	DS.2.5.2	Check if a correct data import to level 2 has been made.
-------------------------	----------------	----------	--

A manual check-list is under development for correctness of data import to level 2.

6.10 Recalculation

Below follows an overview of improvements and recalculations implemented since the 2013 submission.

Recalculations

Some changes in calculation of agricultural emissions 1990-2011 have taken place. The recalculation has contributed to a decrease in the total agricultural emissions for the years 1990-2011 of up to 0.2 % given in CO₂ equivalent (Table 6.52).

Table 6.52 Changes in GHG emission in the agricultural sector compared with the CRF reported last year.

	1990	1995	2000	2005	2008	2009	2010	2012
<u>Previous inventory, Gg CO₂ eqv.</u>								
4.A Enteric Fermentation	3 247	3 134	2 861	2 737	2 830	2 823	2 862	2 840
4.B Manure Management	1 593	1 669	1 722	1 816	1 730	1 693	1 723	1 711
4.D Agricultural Soils	7 702	6 787	5 885	5 295	5 379	5 079	5 026	5 118
4.F Field Burning of Agricultural Residues	3	3	4	4	3	4	3	3
<u>Recalculated, Gg CO₂ eqv.</u>								
4.A Enteric Fermentation	3 247	3 134	2 861	2 737	2 830	2 823	2 862	2 840
4.B Manure Management	1 585	1 657	1 709	1 816	1 730	1 693	1 722	1 711
4.D Agricultural Soils	7 692	6 777	5 879	5 290	5 374	5 075	5 022	5 114
4.F Field Burning of Agricultural Residues	3	3	4	4	3	4	3	3
<u>Change in Gg CO₂ eqv.</u>								
4.A Enteric Fermentation	0.0	0.0	0.0	0.0	0.0	0.0	0.0	-0.1
4.B Manure Management	-8.0	-11.6	-12.4	0.0	0.0	0.0	-1.6	-0.2
4.D Agricultural Soils	-10.5	-9.8	-6.1	-4.8	-4.6	-3.5	-3.7	-3.7
4.F Field Burning of Agricultural Residues	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
<u>Change in pct.</u>								
4.A Enteric Fermentation	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
4.B Manure Management	-0.5	-0.7	-0.7	0.0	0.0	0.0	-0.1	0.0
4.D Agricultural Soils	-0.1	-0.1	-0.1	-0.1	-0.1	-0.1	-0.1	-0.1
4.F Field Burning of Agricultural Residues	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
<u>Change in total, Gg CO₂ eqv.</u>								
	-18	-21	-19	-5	-5	-4	-5	-4
<u>Change in total, pct.</u>								
	-0.15	-0.19	-0.18	-0.05	-0.05	-0.04	-0.06	-0.04

The most significant inventory changes are mentioned below:

The amount of straw used for bedding for heifers has been changed for 1990-2002 and this affects the emission of CH₄ from manure management. The number of geese has been changed for all years and the number of weaners

and fattening pigs has been changed for 2011, this affects both emissions from enteric fermentation, manure management and agricultural soils.

The amount of biogas treated manure has been changed in 2010 and this decreases the emission of manure management.

The main change in emission from agricultural soils is due to change in EF for NH_3 from synthetic fertilisers, which has been changed for all years due to the updated version of the EEA/EMEP Guidebook in 2013. This increases the emission of N_2O from atmospheric deposition due to the increase in emission of NH_3 from synthetic fertilisers while the emission of N_2O directly from synthetic fertilisers decreases.

6.11 Planned improvements

The Danish emission inventory for the agricultural sector meets the requirements given in the IPCC Good Practice Guidance. In the years to come and based on the ERT recommendations, two specific improvements have to be mentioned.

For next submission it is planned to update some of the emission calculations to follow the IPCC 2006 guidelines. The global warming potential for CH_4 and N_2O will be changed from 21 and 310 to 25 and 298, respectively. Emission factors for N_2O from agricultural soils will be revised. Furthermore will the types of housing systems be revised in proportion to the systems given in IPCC 2006.

First step is taken to improve the documentation of the biogas treated slurry. The Energy Statistics got information of the total produced biogas production in Denmark and from another statistic; Energy production census it is possible to get plant specific information regarding the energy production. We still got the challenge with the energy potential in manure – how many tonnes of slurry are used to produce the total energy as given in the Energy Statistics. Present a conversion factor based on an expert judgment from the Danish Energy Agency is used. Next step is to find information which can verify the expert judgment or even better improve the conversion factor. We still investigate the opportunity to use the register for fertilization where all farmers have to register the amount of nitrogen, but it requires a calculation from amount of nitrogen to amount of manure, which differ significant between the different animal types. Data revived from the Danish Biogas Plant Association could also be helpful to check the information of the amount of biogas treated slurry for the largest biogas plants.

Another issue which has to be investigated are improvements of the documentation regarding the emission reduction potential. This is planned to be done by a literature study. Other countries e.g. Germany also use biogas treated slurry and could have some available interesting data. Based on this knowledge it is hopefully possible to do some improvements in near future.

Besides the biogas issue, further work to document the comprehensive QC procedures is planned. Further focus will in particular be addressed to compare the calculations from our database IDA with estimates from other institutions as far as available data makes it possible (refer to “Stage V” in the QA/QC plan – see Chapter 6.9.1).

6.12 References

- Bielecki, J. 2002: Personal communication. Danish Environmental Protection Agency.
- Børgesen, C.D. & Grant, R. 2003: Vandmiljøplan II – modelberegning af kvælstofudvaskning på landsplan, 1984 til 2002. Baggrundsnotat til Vandmiljøplan II - slutevaluering. December 2003, Danmarks Jordbrugsforskning og Danmarks Miljøundersøgelser. (In Danish).
- Clausen, E. 2012: Knowledge Centre for Agriculture, Department for horses. Pers. comm.
- DAAS, 2005: Møller, J., Thøgersen, R., Helleshøj, M.E., Weisbjerg, M.R., Søegaard, K. & Hvelplund, T. Fodermiddeltabel 2005. Rapport nr. 112. The Danish Agricultural Advisory Service, Department for Cattle. (In Danish).
- Danish AgriFish Agency, 2013: Synthetic fertiliser sold in Denmark 2011/2012. Available at (in Danish):
http://naturerhverv.dk/fileadmin/user_upload/NaturErhverv/Filer/Landbrug/Handelsgoedning/Danmarks_salg_af_handelsgoedning_2011_2012.pdf (27.01.2014)
- DEA, 2013: The Danish energy statistics, Available at:
<http://www.ens.dk/node/2228>
- Djurhuus, J. & Hansen, E.M. 2003: Notat vedr. tørstof og kvælstof i efterladte planterester for landbrugsjord – af 21. maj 2003. Forskningscenter Foulum, Tjele. (In Danish).
- Dustan, A. 2002: Review of methane and nitrous oxide emission factors for manure management in cold climates. JTI – Swedish Institute of Agricultural and Environmental Engineering, Uppsala. Report 299.
- EMEP/EEA air pollutant emission inventory guidebook, 2013. Available at:
<http://www.eea.europa.eu/publications/emep-eea-guidebook-2013> (27.01.2014).
- Fenhann, J. & Kilde, N.A. 1994: Inventory of Emissions to the Air from Danish Sources 1972-1992. System Analysis Department – Risø National Laboratory.
- Hansen, N.E. 2010: Department of Basic Animal and Veterinary Sciences/Animal Nutrition, Faculty of Life Science, University of Copenhagen. Pers. comm., mail 13.12.2010.
- Hashimoto, A. & Steed, J. 1993: Methane Emissions from Typical U.S. Livestock Manure Management Systems. Draft report prepared for ICF Incorporated under contract to the Global Change Division of the Office of Air and Radiation, US Environmental Protection Agency, Washington, D.C., U.S.A.
- Husted, S. 1994: Waste Management, Seasonal Variation in Methane Emission from Stored Slurry and Solid Manures. J. Environ. Qual. 23:585-592 (1994).

Høgh-Jensen, H., Loges, R., Jensen, E.S., Jørgensen, F.V. & Vinther, F.P. 1998: Empirisk model til kvantificering af symbiotisk kvælstoffiksering i bælgeplanter. – Kvælstofudvaskning og -balancer i konventionelle og økologiske produktionssystemer (Red. Kristensen E.S. & Olesen, J.E.) s. 69-86, Forskningscenter for Økologisk Jordbrug. (In Danish).

IPCC 1997: Revised 1996 IPCC Guidelines for National Greenhouse Gas Inventories. Available at: <http://www.ipcc-nggip.iges.or.jp/public/gl/invs1.html> (29.01.2014).

IPCC 2000: Good Practice Guidance and Uncertainty Management in National Greenhouse Gas Inventories. Available at: <http://www.ipcc-nggip.iges.or.jp/public/gp/english/> (29.01.2014).

IPCC 2006: 2006 IPCC Guidelines for National Greenhouse Gas Inventories. Available at: <http://www.ipcc-nggip.iges.or.jp/public/2006gl/index.html> (29.01.2014).

Kristensen, I.S. 2003: Indirekte beregning af N-fiksering. Intern note, Danmarks JordbrugsForskning. (In Danish).

Kyllingsbæk, A. 2000: Kvælstofbalancer og kvælstofoverskud i dansk landbrug 1979-1999. DJF rapport nr. 36/markbrug, Dansk Jordbrugsforskning.

Laursen, B. 1987: Normtal for husdyrgødning. Report no. 82, Institute of Food and Resource Economics, University of Copenhagen (previously referred to as Statens Jordbrugsøkonomiske Institut). (In Danish).

Laursen, B. 1994: Normtal for husdyrgødning – revideret udgave af rapport nr. 28. Report no. 82, Institute of Food and Ressource Economics, University of Copenhagen (previously referred to as Statens Jordbrugsøkonomiske Institut). (In Danish).

Mangino, J., Bartram, D. & Brazy, A. 2001: Development of a methane conversion factor to estimate emissions from animal waste lagoons. Presented at U.S. EPA's 17th Annual Emission Inventory Conference, Atlanta GA, April 16-18, 2002.

Massé, D.I., Croteau, F., Patni, N.K. & Masse, L. 2003: Methane emissions from dairy cow and swine slurries stored at 10°C and 15°C. Agriculture and Agri-Food Canada, Canadian Biosystem Engineering, Volume 45 p. 6.1-6.6.

Mikkelsen, M.H., Albrektsen, R. & Gyldenkerne, S. 2011: Danish emission inventories for agriculture. Inventories 1985 - 2009. National Environmental Research Institute, Aarhus University. 136 pp. – NERI Technical Report No. 810. Available at: <http://www2.dmu.dk/Pub/FR810.pdf> (29.01.2014).

Mikkelsen, M.H., Gyldenkerne, S., Poulsen, H.D. Olesen, J.E. & Sommer, S.G. 2006: Emission of ammonia, nitrous oxide and methane from Danish Agriculture 1985-2002. Methodology and Estimates. NERI Research Notes Nr. 231/2006. Available at <http://www2.dmu.dk/Pub/AR231.pdf> (29.01.2014).

Møller, H.B. 2003: Pers. comm. Department of Agriculture Engineering, Research Centre Bygholm, Faculty of Agricultural Sciences.

Nielsen, O.-K., Plejdrup, M.S., Winther, M., Gyldenkerne, S., Thomsen, M., Fauser, P., Nielsen, M., Mikkelsen, M.H., Albrektsen, R., Hjelgaard, K., Hoffmann, L. & Bruun, H.G. 2013a. Quality manual for the Danish greenhouse gas inventory. Version 2. Aarhus University, DCE – Danish Centre for Environment and Energy, 44 pp. Scientific Report from DCE – Danish Centre for Environment and Energy No. 47
<http://www.dmu.dk/Pub/SR47.pdf> (29.01.2014)

Nielsen, O.-K., Winther, M., Mikkelsen, M.H., Hoffmann, L., Nielsen, M., Gyldenkerne, S., Fauser, P., Plejdrup, M.S., Albrektsen, R., Hjelgaard, K. & Bruun, H.G. 2013b: Annual Danish Informative Inventory Report to UNECE. Emission inventories from the base year of the protocols to year 2011. Aarhus University, DCE – Danish Centre for Environment and Energy, 669 pp. Scientific Report from DCE – Danish Centre for Environment and Energy Nr. 53. Available at: <http://www2.dmu.dk/Pub/SR53.pdf> (27.01.2014)

Olesen, J.E., Fenhann, J.F., Petersen, S.O., Andersen, J.M. & Jacobsen, B.H. 2001: Emission af drivhusgasser fra dansk landbrug. DJF rapport nr. 47, markbrug, Danmarks Jordbrugsforskning, 2001. (In Danish).

Olesen, J.E., Jørgensen, H., Danfær, A., Gyldenkerne, S., Mikkelsen, M.H., Asmon, W.A.H. & Petersen, S.O. 2005: Evaluering af mulige tiltag til reduktion af landbrugets metanemissioner. Arbejdsrapport fra Miljøstyrelsen Nr. 11 /2005. Chapter 1 (Allan Danfær): Methane emission from dairy cows. Available at:
<http://www2.mst.dk/Udgiv/publikationer/2005/87-7614-699-5/pdf/87-7614-700-2.pdf> (29-01-2014).

Petersen, C. 2011: Econet AS independent consulting firm. Pers comm.

Petersen, C. & Kielland, M. 2003: Statistik for jordbrugsmæssig anvendelse af affaldsprodukter fra husholdning og institutioner og virksomheder 2001. Miljøprojekt nr. 858 2003. The Danish Environmental Protection Agency.

Poulsen, H.D. 2013: Normative figures 2000-2013. DCA - Danish Centre for Food and Agriculture, Aarhus University. Available at:
<http://anis.au.dk/forskning/sektioner/husdyrernaering-og-miljoe/normtal/> (24.01.2014).

Poulsen, H.D., Børsting, C.F., Rom, H.B. & Sommer, S.G. 2001: Kvælstof, fosfor og kalium i husdyrgødning – normtal 2000. DJF rapport nr. 36 – husdyrbrug, Danmarks Jordbrugsforskning. (In Danish).

Poulsen, H.D. DCA - Danish Centre for Food and Agriculture, Aarhus University, Pers. comm.

Poulsen, H.D. & Kristensen, V.F., 1997: Normtal for husdyrgødning – en revidering af danske normtal for husdyrgødningens indhold af kvælstof, fosfor og kalium. Beretning nr. 736, Faculty of Agricultural Sciences, Aarhus University. (In Danish).

Poulsen, H.D. & Kristensen, V.F. 1998: Standards Values for Farm Manure – A revaluation of the Danish Standards Values concerning the Nitrogen, Phosphorus and Potassium Content of Manure. DJF Report No. 7 - Animal

Husbandry. Danish Institute of Agricultural Sciences. Available at: <http://web.agrsci.dk/djfpublikation/djfpdf/djfh7.pdf> (29.01.2014)

Priemé, A. & Christensen, S. 1991: Emission of methane and non-methane volatile organic compounds in Denmark – Sources related to agriculture and natural ecosystems. National Environmental Research Institute. NERI, Technical Report No. 19/1999.

Sommer, S.G., Møller, H.B. & Petersen, S.O. 2001: Reduktion af drivhusgas-emission fra gylle og organisk affald ved Biogasbehandling. DJF rapport - Husdyrbrug, 31, 53 pp. (In Danish).

Sommer, S.G., Petersen, S.O. & Møller, H.B. 2004: Algorithms for calculating methane and nitrous oxide emissions from manure management. Nutrient Cycling in Agroecosystems, Volume 69, No. 2: 143-154.

Sommer, S.G., Petersen, S.O. & Sogaard, H.T. 2000: Greenhouse gas emissions from stored livestock slurry. Journal of Environmental Quality, 29: pp. 744-751.

Statistics Denmark - Agricultural Statistic from year 1990 to 2011. Available at: <http://www.dst.dk/en.aspx> (29.01.2014).

Sørensen, P.B., Illerup, J.B., Nielsen, M., Lyck, E., Bruun, H.G., Winther, M., Mikkelsen, M.H., & Gyldenkerne, S. 2005: Quality manual for the greenhouse gas inventory Version 1. National Environmental Research Institute, Denmark. 25 pp. – Research Notes from NERI no. 224. – Available at: http://www2.dmu.dk/1_viden/2_Publikationer/3_arbrapporter/rapporter/AR224.pdf (22.01.2013).

Tafdrup, S. 2010: Danish Energy Agency, Pers. Comm.

Wang, S.-Y. & Huang, D.-J. 2005: Assessment of Greenhouse Gas Emissions from Poultry Enteric Fermentation, Asian-Aust. J. Anim. Sci. Vol 18, No. 6: 873-878.

Wiberg-Larsen, P., Windolf, J., Baattrup-Pedersen, A., Bøgestrand, J., Ovesen, N.B., Larsen, S.E., Thodsen, H., Sode, A., Kristensen, E., Kronvang, B. & Kjeldgaard, A. 2010: Vandløb 2009. NOVANA. Danmarks Miljøundersøgelser, Aarhus Universitet. 100 s. – Faglig rapport fra DMU nr. 804. Available at (In Danish): <http://www2.dmu.dk/Pub/FR804.pdf> (29.01.2014).

Windorf, J., Thodsen, H., Trolborg, L., Larsen, S.E., Bøgestrand, J., Ovesen, N.B. & Kronvang, B. 2011: A distributed modelling system for simulation of monthly runoff and nitrogen sources, loads and sinks for ungauged catchments in Denmark. Journal of Environmental Monitoring 2011, 13, 2645-2658. Available at: <http://pubs.rsc.org/en/Content/ArticleLanding/2011/EM/c1em10139k> (29.01.2014).

Waagepetersen, J., Grant, R., Børgesen, C.D. & Iversen, T.M. 2008: Midtvejs-evaluering af Vandmiljøplan III. Available at (In Danish): http://pure.agrsci.dk:8080/fbspretrieve/2617161/VMPIII_midtvejs_2008.pdf (29.01.2014).

Woodbury, J.W. & Hashimoto, A. 1993: "Methane Emissions from Livestock Manure." In International Methane Emissions, US Environmental Protection Agency, Climate Change Division, Washington, D.C., U.S.A.

7 LULUCF (CRF sector 5)

7.1 Overview of the sector

This chapter covers only the territory of Denmark without the Faroe Islands and Greenland. Greenland is submitting a separate NIR and the corresponding CRF tables for the Greenlandic territory to UNFCCC. This can be found as Chapter 16 in this NIR.

Denmark (Capital: Copenhagen) is situated around 56°N and 13°E and covers 43,098 km². No permanent ice is occurring and only very small insignificant areas with rocks. The climate is according to IPCC GPG 2003 cold and wet. Denmark is an intensive agricultural country where most of the area is affected by agriculture. The average temperature in the standard 30 year, 1961-1990 was 7.7 °C with a minimum temperature in February of 0.3 °C and a maximum in July of 17.0 °C. Year 2012 had an average mean temperature of only 8.3 °C which is 0.6 °C above the average. 2012 was slightly cooler than 2011. 2011 was the ninth warmest recorded year since 1873 and slightly cooler than 2008, which had an average temperature of 9.4 °C (www.dmi.dk).

All land is classified into Forest, Cropland, Grassland, Wetlands, Settlements or Other Land.

7.1.1 Abbreviations

The following abbreviations are used in accordance with definitions in the IPCC guidelines:

- A: Afforestation, areas with forest established after 1990 under article 3.3.
- R: Reforestation, areas which have temporarily been unstocked for less than 10 years - included under article 3.4.
- D: Deforestation, areas where forests are permanently removed to allow for other land use, included under article 3.3.
- FF: Forest remaining Forest, areas remaining forest after 1990.
- FL: Forest Land meeting the definition of forests.
- CL: Cropland.
- GL: Grassland.
- SE: Settlements.
- OL: Other land, unclassified land.
- FM: Forest Management, areas managed under article 3.4.
- CM: Cropland Management, areas managed under article 3.4.
- GM: Grazing land Management, areas managed under article 3.4.

The LULUCF sector differs from the other sectors in that it contains both sources and sinks of carbon dioxide. Removals are given as negative figures and emissions are reported as positive figures according to the guidelines. For 2012 emissions from LULUCF were estimated to be a net sink of approximately 838 Gg CO₂ equivalents or 1.7 % of the total reported Danish emission.

7.1.2 Methodology overview

Tier

The type of emission factor and the applied tier level for each emission source are shown in Table 7.1 below. The tier level has been determined based on the 1996 Guidebook (IPCC 1997).

Distinguishing between tier level 2 and 3 have been based on the emission factor. The tier levels definitions have been interpreted as follows:

- Tier 1: The emission factor is an IPCC default tier 1 value.
- Tier 2: The emission factors are country specific and based on either a few emission measurements or IPCC tier 2 emission factors.
- Tier 3: Based on models which include carbon stock changes methodologies.

Table 7.1 shows which of the source categories are key in any of the key source analysis¹ (including LULUCF, tier 1/tier 2, level/trend).

Table 7.1 Methodology and type of emission factor.

		Tier	EF	Key category
5.A.1 Forest	CO ₂	Tier 3, Tier 1	CS, D	Yes
5.A.2 Forest	CO ₂	Tier 3, Tier 1	CS, D	No
5(II) Forest Land.	N ₂ O	Tier 2	CS	No
5.B Cropland, Living biomass	CO ₂	Tier 2	CS	No
5.B Cropland, Mineral soils	CO ₂	Tier 3	CS, D	Yes
5.B Cropland, Organic soils	CO ₂	Tier 2	CS, D	Yes
5(III) Disturbance, Land converted to cropland	N ₂ O	Tier 2	CS, D	No
5.C Grassland, Living biomass	CO ₂	Tier 2	CS, D	Yes
5.C Grassland, Mineral soils	CO ₂	Tier 2	CS, D	No
5.C Grassland, Organic soils	CO ₂	Tier 2	CS, D	No
5.D Wetlands, Living biomass	CO ₂	Tier 2	CS, D	No
5.D Wetlands, Dead organic matter	CO ₂	Tier 2	CS, D	No
5.D Wetlands, Soils	CO ₂	Tier 2	CS, D	Yes
5(II) Wetlands	N ₂ O	Tier 2	CS, D	No
5.E Settlements, Living biomass	CO ₂	Tier 2	CS, D	No
5(IV) Cropland Limestone	CO ₂	Tier 2	CS, D	Yes
5(V) Biomass Burning	CH ₄	Tier 2, Tier 1	CS, D	No
5(V) Biomass Burning	N ₂ O	Tier 2, Tier 1	CS, D	No

7.1.3 Key categories

Key Category Analysis (KCA) tier 1 and 2 for year 1990, 2012 and trend for Denmark has been carried out in accordance with the IPCC Good Practice Guidance / IPCC Guidelines (2006). Table 7.2 shows which of the LULUCF categories are identified as key categories. The table is based on the analysis including LULUCF. Detailed key category analysis is shown in NIR Chapter 1.5 and Annex 1.

¹ Key category according to the KCA tier 1 or tier 2 for Denmark (excluding Greenland and Faroe Islands), including LULUCF, level 1990/ level 2012/ trend.

The CO₂ emissions from forests are key for forest remaining forest both on the level and the trend. For Cropland both mineral and organic soils are key sources as well as lime stone.

Table 7.2 Key categories, LULUCF.

			Tier 1			Tier 2		
			1990	2012	1990-2012	1990	2012	1990-2012
LULUCF	5.A.1 Forest	CO ₂		Level	Trend		Level	Trend
LULUCF	5.B Cropland, Living biomass	CO ₂		Level	Trend		Level	Trend
LULUCF	5.B Cropland, Mineral soils	CO ₂	Level	Level		Level	Level	
LULUCF	5.B Cropland, Organic soils	CO ₂	Level	Level		Level	Level	
LULUCF	5.C Grassland, Living biomass	CO ₂		Level	Trend		Level	Trend
LULUCF	5.D Wetlands, Soils	CO ₂						Trend
LULUCF	5(IV) Cropland Limestone	CO ₂	Level		Trend	Level		Trend

7.1.4 Methods

Approximately 2/3 of the total Danish land area is cultivated and 14.7 per cent forested. Together with high number of cattle and pigs there is a high (environmental) pressure on the landscape. To reduce the impact an active policy has been adopted to protect the environment. The adopted policy aims at doubling the forested area within the next 80-100 years, restoration of former wetlands and establishment of protected national parks. In Denmark almost all natural habitats and all forests are protected. Therefore only limited conversions from forest or wetlands into cropland or grassland are occurring.

Estimation of carbon stock changes in the Danish forests is based on a combination of surveys and the National Forest Inventory (NFI). Changes in carbon stock in mineral cropland soils are estimated with a nationally developed Tier 3 model, whereas the emission calculation from organic soils is based on nationally developed emission factors.

Since the last submission has a minor revision of the land use matrix for the whole period taken place. Primarily because of a misclassification of some urban parks as forest. The revision has only minor impact on the emission estimate.

No permanent snow cover exists in Denmark and only a very small insignificant area with rocks and cliffs. OL is restricted to beaches and sand dunes. The official land area is 43 098 km². The new land use matrix has estimated the total area to 43 056 km². This area includes rivers and lakes. The small discrepancy is due to differences in the definition of the 7000 km long coast-line. The land use matrix uses the latest official vector maps from Danish Geodata Agency.

The emission data are reported in the CRF format under IPCC categories 5A (Forestry), 5B (Cropland), 5C (Grassland), 5D (Wetlands) and 5E (Settlements) and 5F (Other Land). Denmark is free from ice and rocks and Other Land therefore represents beaches and sand dunes. Fertilisation of Forests and Other Land is negligible and all fertiliser consumption is therefore reported in the agricultural sector. All liming is reported under Cropland because only very limited amounts are used in forestry and on permanent grassland. Field burning of biomass is prohibited in Denmark. Wildfires in forest are reported. This is normally around 0-10 hectares per year. Con-

trolled burning of heathland is taking place of approximately 300-700 hectares to maintain the heath.

Savannas and rice cultivation do not occur in Denmark.

The Cropland and Grassland area are based on agricultural EU subsidiary systems and very detailed. A drawback is, however, that one field in one year can be classified as CL and the next year as GL and then again converted back to CL. This creates large conversion rates between cropland and grassland but mainly towards grassland as an extensification currently takes place in Denmark (Table 7.3). The switching between CL and GL will, however, have no effect on the emission estimates except for an estimated release of N₂O from mineralisation of organic matter. This issue is also recognized in the IPCC 2006 Guidelines. Table 7.3 shows the overall development from 1990 to 2012. Afforestation is mainly taking place on CL and GL not previously classified as forest. Areas, which are deforested are mainly converted to WE as a consequence of clearing of some areas in the State forests towards more open areas, to CL due to conversion of christmas trees agricultural cropland. Since 1990 more than 31 100 hectares have been changed into SE and other infrastructures. No land is converted into OL.

In the new land use matrix has a linear approach for all land use changes been adopted for the period 1990 to 2005 and from 2005 to 2011. From 2011 and onwards is used annually updated data from the different data suppliers. Some of these data are not updated annually and thus a time lag in the implementation of the land use changes may occur in some areas. Conversion to annual updates may create more fluctuating area changes than in the previous years.

Table 7.3 Land Use Change from 1990 to 2012 based on GIS vector layers and Earth Observations. The figures are given in hectares.

1990\2012	Forest	Cropland	Grassland	Wetlands	Lakes	Settlements	Other	Sum
Forest	538 189	1 270	1 713	2 145	346	310	0	543 973
Cropland	48 355	2 590 240	49 600	12 074	2 691	15 403	0	2 718 362
Grassland	45 985	74 740	260 279	3 294	2 691	15 384	0	402 374
Wetlands	18	560	0	69 853	6	71	0	70 509
Lakes	0	0	0	0	58 357	1	0	58 358
Settlements	0	0	0	0	0	485 543	0	485 543
Other	0	0	0	0	0	0	26 433	26 433
Sum	632 547	2 666 811	311 592	87 367	64 092	516 712	26 433	4 305 552
Percentage	14.7%	61.9%	7.2%	2.0%	1.5%	12.0%	0.6%	100.0%

Table 7.4 gives an overview of the emission from the LULUCF sector in Denmark. Forests have been sinks in Denmark for the last decade but due to the age distribution of the forests - containing a majority of mature forests - a slight decrease of the carbon stock is observed, as the old forests are regenerated with young trees and a net source were observed. The changes occur before the 2008-2012 period and the results can also partly be attributed to the recalculations - as described later. Currently the NFI indicates that forests are a sink. Cropland is ranging from being a net source from up to 4,642 Gg in 1990 to be a net source of 3,185 Gg in 2010. High fluctuations in the emission from CL between years are related to the actual crop yield that year and the climatic conditions. Low crop yields combined with high temperatures reduce the total amount of carbon in agricultural soils whereas a year with a high yield and low temperatures increase the carbon stock in soil.

From 1990 and onwards a general decrease in the emission from Cropland is estimated due to a higher incorporation of straw (ban of field burning), demands of growing of catch crops in the autumn, a change from low yielding spring barley to high yielding winter wheat, an increased carbon stock in hedgerows, a reduced consumption of lime and that organic agricultural soils are disappearing. The area with restored wetlands has increased and consequently the accumulation of organic matter has also increased here leading to a lower net source.

Table 7.4 Overall emission (Gg CO₂) from the LULUCF sector in Denmark, 1990- 2012.

Greenhouse gas source and sink categories	1990	1995	2000	2005	2008	2009	2010	2011	2012
5. Land Use, Land-Use Change and Forestry, CO ₂	5,265.22	3,655.80	3,219.87	4,461.90	-1,611.35	3,032.18	-337.70	-2,756.28	-851.70
A. Forest Land	126.2	-954.7	-764.1	615.4	-5716.4	31.1	-4046.5	-6311.3	-4453.54
B. Cropland	4855.0	4389.2	3760.5	3473.5	3730.7	2625.3	3349.4	3156.2	2955.24
C. Grassland	183.6	152.2	158.9	234.6	269.3	256.5	246.4	277.4	553.52
D. Wetlands	87.5	51.3	41.8	99.5	59.7	71.9	63.5	69.6	2.27
E. Settlements	12.9	17.8	22.7	39.0	45.3	47.4	49.5	51.7	90.81
F. Other Land	NA, NO	NA, NO	NA, NO	NA, NO	NA, NO	NA, NO	NA, NO	NA, NO	NA, NO
5. Land Use, Land-Use Change and Forestry, N ₂ O (CO ₂ eq)	16.5	15.2	14.2	13.2	12.7	12.5	12.5	12.4	13.1
A. Forest Land	16.1	14.7	13.8	12.9	12.3	12.1	12.1	12.1	12.1
B. Cropland	0.3	0.4	0.3	0.2	0.2	0.2	0.2	0.2	0.9
C. Grassland	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
D. Wetlands	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
E. Settlements	NO	NO	NO	NO	NO	NO	NO	NO	NO
F. Other Land	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO
5. Land Use, Land-Use Change and Forestry, total CO ₂ eq	5282.2	3671.0	3234.1	4475.2	-1598.7	3044.7	-325.2	-2743.8	-838.5

7.2 Forest remaining forest (5.A.1)

7.2.1 Forest census

From 1881 to 2000, a National Forest Census has been carried out roughly every 10 years based on questionnaires sent to forest owners (Larsen and Johannsen, 2002). Since the data was based on questionnaires and not field observations, the actual forest definition may have varied. The basic definition was that the tree covered area should be minimum 0.5 ha to be a forest. There were no specific guidelines as to crown cover or the height of the trees. Open woodlands and open areas within the forest were generally not included. All values for growing stock, biomass or carbon pools based on data from the National Forest Census were estimated from the reported data on forest area and its distribution to main species, age class and site productivity classes. The two last censuses were carried out in 1990 and 2000.

The 1990 National Forest Census was based on reported forest statistics from 22,300 respondents, resulting in information on area, main species, age class distribution and productive indicators. The estimated forest area was 445 000 ha or 10.3 % of the land. Of the total forest area 64 % was coniferous forest and 34 % was deciduous forest (the remainder was temporarily unstocked). The total volume was estimated at 55.2 million cubic metres of which 57 % was coniferous.

The number of respondents in the 2000 National Forest Census was 32,300, which is considerably higher than in the 1990 survey. The change in the number of respondents probably contributed to the observed increase in forest area and growing stock between the 1990 and 2000 census. The estimated forest area was 486 000 ha or 11.3 % of the land. Of the total forest area 60 % was coniferous forest and 36 % was deciduous forest (the remainder was temporarily unstocked). The total volume was estimated at 77.9 million cubic metres of which 63 % was coniferous.

7.2.2 National forest inventory

In 2002, a new sample-based National Forest Inventory (NFI) was initiated (Nord-Larsen et al., 2008). This type of forest inventory is very similar to inventories used in other countries, e.g. Sweden or Norway. The NFI has replaced the National Forest Census.

The NFI is a continuous sample-based inventory with partial replacement of sample plots based on a 2 x 2 km grid covering the Danish land surface. At each grid intersection, a cluster of four circular plots (primary sampling unit, PSU) for measuring forest factors (e.g. wood volume) are placed in a 200 x 200 m grid. Each circular plot (secondary sampling unit, SSU) has a radius of 15 meters. When plots are intersected by different land-use classes or different forest stands, the individual plot is divided into tertiary sampling units (TSU).

About one third of the plots is assigned as permanent and is re-measured in subsequent inventories every five years. Two thirds are temporary and are moved randomly within the particular 2x2 km grid cell in subsequent inventories. The sample of permanent and temporary field plots has been systematically divided into five non-overlapping, interpenetrating panels that are each measured in one year and constitute a systematic sample of the entire country. Hence all the plots are measured in a 5-year cycle.

Based on analysis of aerial photos, each sample plot (SSU) is allocated to one of three basic categories, reflecting the likelihood of forest or other wooded land (OWL) cover in the plot: (0) Unlikely to contain forest or other wooded land cover, (1) Likely to contain forest, and (2) Likely to contain other wooded land. All plots in the last two categories are inventoried in the field.

In the most recent five-year rotation of the NFI (2008-2012) the number of clusters (PSU) and sample plots (SSU) were 4.138 and 9.425, respectively. On average 1,885 yr⁻¹ plots (SSU) were identified as having forest or other wooded land cover based on the aerial photos and were thus selected for inventory. Only 3 sample plots were missing in the 2008-2012 inventories, due to difficulties of access.

Table 7.5 Number of measured clusters and sample plots in the five year rotation 2008-2012. Forest covered sample plots not inventoried in the field are denoted "Missing".

Year	Clusters			Sample plots		
	Total	Forest	Missing	Total	Forest	Missing
2008	2 212	804	2	8 644	1 896	3
2009	2 195	783	0	8 604	1 800	0
2010	2 196	793	0	8 614	1 855	0
2011	2 173	850	0	8 520	1 896	0
2012	2 200	908	0	8 617	1 978	0
Total	10 976	4 138	2	42 999	9425	3

Each plot is divided into three concentric circles with radius 3.5, 10 and 15 m. A single caliper measurement of diameter is made at breast height for all trees in the 3.5 m circle. Trees with diameter larger than 10 cm are measured in the 10 m circle and only trees larger than 40 cm are measured in the 15 m circle. On a random sample of 2-6 trees further measurements of total height, crown height, age and diameter at stump height are made and the presence of defoliation, discoloration, mast, mosses and lichens are recorded. The presence of regeneration on the plots is registered and the species, age and height of the regeneration are recorded. Stumps from trees harvested within a year from the measurement are measured for diameter.

Deadwood is measured on the sample plots. Standing deadwood with a diameter at breast height diameter larger than 4 cm is measured according to the same principles as live trees. Lying deadwood with a diameter of more than 10 cm is measured within the 15 m radius sample plot. Length of the lying deadwood is measured as the length of the tree that exceeds 10 cm in diameter and is within the sample plot. The diameter is measured at the middle of the lying deadwood measured for length. In addition to the size measurements of deadwood the degree of decay is recorded on an ordinal scale.

On each plot the presence and state of ditches and drainage conditions are recorded. Further, the presence of peatland is recorded and the depth of the peat is measured. Finally, the depth of the humus layer is measured on all plots.

7.2.3 Forest area mapping

Due to differences in methodologies major inconsistencies in forest areas and other forest variables are observed between the different forest inventories (i.e. the 1990 and 2000 Forest Census and the 2006 National Forest Inventory). With the objective to obtain time consistent and precise estimates of forest areas to report to UNFCCC and under the Kyoto protocol, two projects have aimed at mapping the forest area in Denmark based on satellite images. Forest area and forest area change have been estimated for the years 1990, 2005 and 2011.

A land use/land cover map was produced for the base year 1990 and for the year 2005 based on EO data (23 August 1990) and other data collected from 1992-2005 and for 2005 using NFI in situ data. Forest maps are developed using Landsat imagery mainly Landsat 5 (TM) and 7 (ETM+) data to classify and estimate the area of forest cover types in Denmark. Portions of seven scenes covering the whole country were classified into forest and non-forest classes. The approach involved the integration of sampling, image processing, and estimation. A detailed QA/QC process was conducted in 2011/2012. Maps for 2011 were produced in 2012 (Huber & Tøttrup 2012). In order to map the forest cover, multi-spectral and multi-temporal Landsat data of June 2010 and April 2011 with a spatial pixel resolution of 30 m were used. Except of Bornholm, none of the scenes was cloud-free. To still obtain a national forest cover map without gaps, the forest cover map of some minor areas is solely based on one image.

The product is specified by a Minimum Mapping Unit (MMU) of 0.5 ha, a geometric accuracy of < 15 m RMS and a thematic accuracy of 90 % +/- 5 % for the land use class Forest.

7.2.4 Forest definition

The forest definition adopted in the NFI is identical to the FAO definition (TBFRA, 2000). It includes “wooded areas larger than 0.5 ha, that are able to form a forest with a height of at least 5 m and crown cover of at least 10 %. The minimum width is 20 m.” Temporarily non-wooded areas, fire breaks and other small open areas, that are an integrated part of the forest, are also included.

7.2.5 Estimation of forest carbon pools

In the following, procedures for estimating forest carbon pools are described. For the specific formulas used in the calculations, readers are referred to Annex 3F.

Estimation of forest area

Based on analysis of aerial photos, each sample plot (SSU) is allocated to one of three forest status categories (Z), reflecting the likelihood of forest or other wooded land (OWL) in the plot: (0) Unlikely to be covered by forest or other wooded land, (1) Likely to be covered by forest, and (2) Likely to be covered by other wooded land.

On individual sample plots (j) the forest cover percentage (X) is calculated as the proportion of the forest area (A) to the total plot area of the 15 m radius circle (A_{15}). The average forest percentage (\bar{X}) on plots with forest status $Z=1$ (and 2) is calculated as the sum of the forest percentages times an indicator variable (R) that is 1 if Z equals 1 (or 2) and 0 otherwise, divided by the number of plots with forest status $Z=1$ (or 2).

The overall average forest percentage (\bar{X}) is calculated as the sum of: (1) observed forest cover percentages of the individual sample plots, (2) the number of unobserved sample plots with forest status $Z=1$ times the average forest cover percentage of sample plots with forest status 1, and (3) the number of unobserved sample plots with forest status 2 times the average forest cover percentage of observed sample plots with forest status $Z=2$ divided by the number of observed and unobserved sample plots. In this context sample plots with forest status 0 are regarded as observed and assumed to have a forest cover percentage of 0. Finally, the overall forest area (A_{Forest}) is calculated as the overall average forest percentage times the total land area (A_{total}).

When estimating the forest area with a specific characteristic (k), such as forest established before or after 1990, the proportion of the plot area with the particular characteristic is found by summing the forested plot areas times an indicator variable (R) that is 1 if the plot has the k th characteristic and 0 otherwise. Subsequently the plot area with the k th characteristic is divided by the total forested plot area.

The total forest area with a particular characteristic (A_k) is found as the forest area percentage with the particular characteristic k times the total forest area.

Estimation of volume, biomass and carbon pools

For estimation of volume of individual trees, we use the volume functions developed for the most common Danish forest tree species (Madsen, 1985, Madsen 1987 and Madsen and Heusèrr 1993). The functions use individual

tree diameter and height as well as quadratic mean diameter of the forest stand as independent variables.

Based on the trees measured for both height and diameter, diameter-height regressions are developed for each species and growth region. The functions use the observed mean height and mean diameter on each sample plot for creating localized regressions using the regression form suggested by Sloboda et al. (1993). For plots where no height measurements are available, generalized regressions are developed based on the Näslund-equation modified by Johannsen (1992).

The next step is to estimate the quadratic mean diameter of the trees on the sample plot. As the trees are measured in different concentric circles depending on their diameter, the basal area on each sample plot is estimated by scaling the basal area of each tree (standing or felled) according to the circular area in which the tree has been measured. A similar calculation has been made for the number of stems. Finally, mean squared diameter is calculated from the basal area and stem numbers.

Based on the diameter, estimated or measured height of individual trees and the squared mean diameter before thinning, the volume of individual trees is estimated using the species specific volume functions by Madsen (1987) and Madsen & Heusèrr (1993). The volume of trees less than 3 meters tall is estimated using an alternative function. The calculated volumes are total stem volume over bark for conifers and total above ground volume over bark for deciduous species.

Based on the estimated individual tree volumes, above ground biomass of the individual tree (stem biomass for conifers and total above ground biomass for broadleaves) is subsequently calculated as the total volume times the basic density. Species specific basic densities are based on Moltesen (1988), Skovsgaard et al. (2011) and Skovsgaard & Nord-Larsen (2012). Finally, total biomass (below and above ground) is estimated using expansion factors. For coniferous species an expansion factor model developed for Norway spruce (Skovsgaard et al. 2011) is applied whereas for deciduous species an expansion factor model developed for beech (Skovsgaard & Nord-Larsen, 2012) is used.

Total or regional volume, biomass and pools of carbon are estimated based on the estimates of individual tree volumes, biomass and carbon. First, volume, biomass or carbon per hectare is estimated for each of the concentric circles ($c=3.5, 10$ or 15 m radius) on each plot as the plot area depends on the diameter of the tree. Using the estimates from individual plots, the area weighted mean volume, biomass or carbon per hectare for the three concentric circles is estimated. The overall mean volume, biomass or carbon is estimated as the sum of the average volumes for the three circles. Finally, the total or regional volume, biomass or carbon is estimated as the forest area times the overall mean volume.

The total volume, biomass or carbon pools with a given characteristic are estimated in a similar way as the total figures. First, volume, biomass or carbon per hectare with the given characteristic is estimated for each of the concentric circles ($c=3.5, 10$ or 15 m radius) on each plot. Using the estimates from individual plots, the area weighted mean volume, biomass or carbon per hectare with the given characteristic for the three concentric circles is es-

timated. The overall mean volume, biomass or carbon is estimated as the sum of the average volumes for the three circles. Finally, the total or regional volume, biomass or carbon with the given characteristic is estimated as the forest area times the overall mean volume.

Dead wood volume, biomass and carbon content

The volume of standing dead trees is calculated similarly to the calculations for live trees. The volume of lying dead trees within the sample plot is calculated as the length of the dead wood times the cross sectional area at the middle of the dead wood. Biomass of the dead wood is calculated as the volume times the species specific basic density and a reduction factor according to the structural decay of the wood. Finally, carbon content for each standing or lying dead tree is calculated by multiplying the dead wood biomass by 0.5.

Total or regional volume, biomass and carbon pools of deadwood are estimated based on the estimates of volumes, biomass and carbon for individual dead trees or pieces of dead wood. First, deadwood volume, biomass or carbon per hectare is estimated for each of the concentric circles ($c=3.5, 10$ or 15 m radius). Estimates for lying dead wood are made using the 15 m circle. Using the estimates from individual plots, the area weighted mean volume, biomass or carbon per hectare of deadwood for the three concentric circles is estimated. The overall mean deadwood volume, biomass or carbon is estimated as the sum of the average volumes for the three circles. Finally, the total or regional deadwood volume, biomass or carbon is estimated as the forest area times the overall mean volume.

Forest floor

On each NFI plot (SSU) the depth of the forest floor is measured. As peatlands are reported specifically, a maximum depth of 15 cm is used in the calculations. Forest floor carbon for individual species is estimated by multiplication of the forest floor depth with the plot area, a species specific density (Vesterdal & Raulund-Rasmussen, 1998) and the fraction of the individual species. The fractions are based on the proportion of basal area of the individual species and total forest floor carbon is estimated by summation of forest floor carbon of the different species.

Average forest floor carbon is estimated by summation of forest floor carbon on the individual plots and dividing the total by the total plot area. Total forest floor carbon is subsequently estimated by multiplication of the average forest floor carbon by the total forest area.

Forest floor carbon stocks were assessed in the Forest Soil Inventory described below. However, there was no good basis to estimate change over time for this C pool as historic data were very scarce (see below). Hence changes in this C pool were based on depth measurements performed on all NFI plots.

Forest mineral soil

The NFI monitoring was supplemented by an additional forest soil inventory in order to document that forest soils is not an overlooked source for CO_2 emissions. The monitoring of soil C-stocks concerns two of the five carbon pools identified by IPCC (2003), litter (forest floor) and mineral soil to a depth of minimum 30 cm.

There is relatively good information from various soil profile databases on mineral soil carbon stocks to 1 m depth for well-drained Danish forest soils (Vejre et al., 2003; Krogh et al., 2003). However, there has been no spatially systematic study performed on temporal change in forest soil carbon. This has limited the possibility to explore the development in forest soil carbon stocks over time. This need is most pronounced for the quickly changing litter carbon pool and previous C stock estimations (Vejre et al., 2003) did not include moist and wet forest soils.

According to decision 16/CMP "A Party may choose not to account for a given pool in a commitment period if transparent and verifiable information is provided that the pool is not a source." The forest soil inventory aims to document that forest soils are not a major source for emissions of CO₂, i.e. that there is no detectable depletion in soil carbon. This may be called the "no source principle" (Somogyi & Horvath, 2007). According to IPCC (2003) the necessary documentation may come from various sources such as:

- Representative and verifiable sampling and analysis to show that the pool has not decreased
- Reasoning based on sound knowledge of likely system responses
- Surveys of peer-reviewed literature for the activity, ecosystem type, region and pool in question
- Combined methods.

Based on literature and reasoning based on sound knowledge there is little evidence to support that the soil C pool in forest remaining forest would currently be changing to an extent that would be detectable by sampling with decadal frequency. For well-drained soils there may be temporal changes in soil carbon stocks at fine spatial resolution (ha-level) due to clear-cutting and replanting, but for the entire forest area with the whole range of age classes, the assumption is that soil carbon stocks are unchanged over time. In fact, the conversion toward close-to-nature forestry with continuous crown cover and abandonment of clearcutting suggests a future increase in soil carbon stocks rather than depletion (Brunner et al., 2005; Yanai et al., 2000). Areas with wet forest soils have probably been sources for increased CO₂ emissions in a period after ditching and drainage activities took place from the late 19th century. These activities led to increased mineralization of peaty soils. However, during the last 20 years, drainage activities have diminished strongly and has almost ceased in state forests as a direct effect of the Strategy for the State forests to convert to more Close to Nature Forest Management, including restoration of more natural hydrology (see more on <http://www.naturstyrelsen.dk/>). Here, the natural hydrological conditions are actively restored by filling up ditches in some areas. It is expected that this change in management will lead to sequestration of carbon as these forest soils gradually get wetter and the rate of decomposition decreases more than rates of organic matter inputs from litterfall. Exact information on the extent of restored natural hydrology is not available, but is being assessed based on expert judgement and information from forest managers.

Since the reporting in 2009 for 1990-2007, quantitative information has gradually become available; a project (SINKS) initiated in 2007 has delivered data on soil C change based on repeated sampling of soil C pools in forests remaining forests, and more data on soil C pools are being made available. The preliminary data suggest that forest soil C pools are not sources for CO₂

and thus support that more accurate estimates of litter and soil C pool removals/emissions do not need to be included in the reporting.

New data

The only existing systematic sampling of Danish forest soils has been conducted within the so-called Kvadratnet ("Agricultural Network", <http://www-landbrugsinfo.dk/Planteavl/Goedskning/Naeringsstoffer/Kvadratnet-for-nitratundersogelser/Sider/Startside.aspx>). It was established in the 1980's in order to optimize the applied amounts of fertilizer in agriculture by monitoring nitrate leaching to groundwater resources in the most common land uses. Soil samples from the years around 1990 exist in soil storages. Given the time constraints of the commitment period and reporting deadlines, changes in soil carbon stocks could only be assessed by repeated sampling of soils within this monitoring grid.

The Agricultural monitoring grid is 7x7 km and by 1990 it included 108 plots with forest cover (Østergaard & Mamsen, 1990). Soil sampling and analysis was conducted in 1986-90 in all 108 forest plots of Agricultural Network, and a subset of 25 plots was resampled in 1994 (Breuning-Madsen & Olsson, 1995) as a part of the Pan-European forest monitoring programme, which uses these 25 plots for assessment of the forest condition. The 25 plots resampled in 1994 have been resampled again in 2007 as a part of the demonstration project BioSoil (<http://forest.jrc.ec.europa.eu/contracts/-biosoil>), under the Pan-European forest monitoring programme Forest Focus and in 2008/2009 the other 83 plots were resampled, except for one plot for which the land owner did not grant access to re-sampling.

Mineral soil samples from 1990 are thus available from 108 forest plots. The sampling was complete for the period 2007-2010, while soil-archive samples from 1990 were missing for six plots. Soil samples from 1986-1987 were used for one of these plots while it was not possible to retrieve archived soil samples for the last five plots. The sampling of O-horizons was also complete for the 108 plots for the period 2007-2009, while O-horizon samples from 1990 were of a very poor quality and only available from 32 plots. Consequently, forest floor samples from 1994 were used to represent forest floors in 1990, while results based on soil samples 0-100 cm from 1994 were only used to check other data.

The plots were in all cases (with a few exceptions due to practical circumstances) designed as a 50 x 50 m square. In 2007-09 ten forest floor and mineral soil cores were collected along a transect determined as the diagonal from the south-west to north-east corner of the square. In the 1990 16 soil cores were taken randomly across the square plot, while forest floor samples were only collected occasionally in an unspecified manner.

The O-horizon samples from 2007-2009 were area-based samples (Vesterdal & Raulund-Rasmussen, 1998) removed from a 25 x 25 cm area, that were brought to the laboratory in separate bags.

The mineral soil samples from 2007-2009 were taken in the ten sampling points where O-horizons had been removed. A 2-3 cm thick soil corer was used. The mineral soil samples from around 1990 were taken in a similar manner for the 16 sampling points. Samples from 4-5 different horizons were pooled in the field. Only one joint sample pr. plot per depth were ana-

lysed for carbon content. Hence, information on within-plot variation in soil carbon contents is not available. The division into horizons differed slightly between the three sampling campaigns: 1986-1990, 2007 and 2008/2009. In 1986-1990 the division was 0-25, 25-50, 50-75 and 75-100 cm; in 2007 (the 25 BioSoil plots) it was 0-10, 10-20, 20-40, 40-80 and 80-100 cm; and in 2008/2009 0-10, 10-25m 25-50, 50-75 and 75-100 cm.

In the lab, all samples were dried at 40°C until constant weight. Before sieving through a 2 mm sieve, more clay-rich mineral soil samples were crushed in a mortar, while sandy soil samples were gently crushed or sieved directly. The stones (>2 mm) left after sieving were weighed (DW_{stone}), while the fine soil (<2 mm) was dried at 40°C for at least 48 h, and then weighed (DW_{soil}). A sub-sample of the fine soil, about 20 g, was removed after thorough mixing for finer grinding in an agate mortar.

The ten O-horizon samples from each plot were weighed separately, and then ground in Retsch grinder through a 2 mm net. From each of the ten samples, 10 % of the material was removed after thorough mixing to get a pooled sample for the plot. About 100 ml of the pooled sample was removed after thorough mixing and then ground more finely in a Tecator mill.

Mineral soil samples were analysed by dry combustion (Elementar Analyz-er) for total organic carbon (TOC) and O-horizon samples for total carbon by a laboratory certified according to ISO 10694. Analyses were done by Agro-lab/ Institut Koldingen, Sarstedt, Germany.

For each of the plots, the mineral soil carbon stocks in 2007-2009, C_{m-2009} (tonne C ha⁻¹), was calculated as

$$C_{m-2009} = \sum_{i=1}^{4(or5)} d_{m-2009} \cdot 10000 \cdot (1 - RV_{stone-2009}) \cdot \rho_{soil} \cdot c_{soil-2009}$$

where d_m is the depth of a given horizon (m), and ρ_{soil} is the bulk density of soils (g cm⁻³) assessed by use of published pedotransfer functions (Vejre et al., 2003). $c_{soil-2009}$ is the C concentration (mg g⁻¹). RV_{stone} is the relative volume of the stone (versus that of the fine soil):

$$RV_{stone-2009} = \frac{DW_{stone-2009} / \rho_{stone}}{DW_{stone-2009} / \rho_{stone} + DW_{soil-2009} / \rho_{soil}}$$

where $\rho_{stone}=2.65$ g cm⁻³, $DW_{soil-2009}$ (g) is the dry weight of the fine soil (<2 mm) in the soil samples from 2007-2009 and $DW_{stone-2009}$ (g) is correspondingly the weight of stones in the soil sample (>2 mm).

For each of the plots, the forest floor carbon stocks in 2007-2009, $C_{ff-2009}$ (t C ha⁻¹), was calculated as

$$C_{ff-2009} = \sum_{i=1}^{10} DW_{ff-2009,i} \cdot 0.0016 \cdot c_{ff-2009}$$

where $DW_{ff-2009,i}$ (g dry weight) is the dry weight of sample number i, i=1-10 and $c_{ff-2009}$ is the C concentration of the pooled sample per plot (mg g⁻¹)

The mineral soil dry weight in 1990 was calculated in the same manner as for 2007-2009, assuming that the relative stone volume was identical to that

of 2007-2009. The forest floor depth was, however, not measured in 1990, nor was an area-based forest floor weight recorded. Forest floor depth (d_{ff} , m) measured for profiles on 25 plots in 1994, was used instead, while forest floor densities for the individual plots were obtained from the new measurements performed in 2007. For these 25 plots, forest floor C-stocks in 1990, $C_{ff-1990}$ (tonne C ha⁻¹) were calculated as

$$C_{ff-1990} = d_{ff} \cdot 10000 \cdot \rho_{ff-2007} \cdot c_{ff-1990}$$

where $c_{ff-1990}$ (mg g⁻¹) is the carbon concentration of the forest floor samples from 1994 (measured in 2007), and $\rho_{ff-2007}$ (g m⁻³) is the average bulk density of the forest floor for the individual plot as measured in 2007:

$$\rho_{ff-2009} = \frac{\sum_{i=1}^{10} DW_{ff,i}}{0.25 \cdot 0.25}$$

Considering the forest structure in Denmark with many small forests (about 70 % of the forest estates are of less than 5 ha) the “Kvadrantet” is a very coarse grid. Even if the grid was fully sampled, it is therefore unlikely that the 108 plots represent the Danish forest area of approximately 500 000 ha. We thus evaluated based on power analyses that further sampling was necessary for future monitoring and chose to include a randomly selected subset of the permanent plots of the National Forest Inventory (NFI) for this purpose. A total of 277 plots were sampled.

It will not be possible, as with the Agricultural Network, to resample soils of the NFI plots for changes in soil C within the short time frame before Kyoto Protocol reporting. From 2012 and onward the NFI plots can be resampled to better support the work to demonstrate that soil carbon stocks are not a source for CO₂ emissions. As the Danish reporting of the three forest carbon pools aboveground biomass, belowground biomass and dead wood is based on the NFI, this will also ensure the consistency of monitoring of all five forest carbon pools defined by IPCC (2003). In the first reporting efforts, however, information on C-stocks and site properties from the NFI will enable better upscaling of results from Agricultural Network to the Danish forest area.

Changes in forest soil carbon stocks in forests planted before 1990

The preliminary results from the Agricultural Network showed that there is a large variation in soil C pools among sites for both forest floors (only 32 plots) and mineral soils. The mean C pool of forest floors was about 22 and 28 tonnes C ha⁻¹ in 1990 and 2007-09, respectively. The corresponding C pools for mineral soils were 156 in 1990 and 157 tonnes C ha⁻¹ in 2007-09 (Table 7.6). A simple t-test of the mean changes in forest floor and mineral soils pools between 1990 and 2007-2009 (5.6 and 1.5 tonnes C ha⁻¹ yr⁻¹ respectively) indicate that changes were not significant (Table 7.7, Figure 7.1a-b).

Table 7.6 Basic statistics on soil C pools measured in the Agricultural Network.

	Mean Pool	Standard deviation	Minimum	Maximum
	tonne C ha ⁻¹			
Forest floor 1990	22.12	19.12	0.76	80.34
Forest floor 2007-2009	27.68	30.05	3.94	164.48
Mineral soil 1990	155.78	115.91	29.31	848.14
Mineral soil 2007-2009	157.26	100.34	18.66	853.08

Table 7.7 Basic statistics on the differences in C soil pools between 1990 and 2007-2009 and statistics from a simple t-test (H_0 : change in soil C-stock = 0).

	Total number of sites	Number of sites in t-test	Mean change	Std	Minimum	Maximum	P-value
				(tonne C ha ⁻¹)			
Forest floor	108	31	5.56	24.78	-61.44	84.13	0.22
Mineral soil	108	104	1.48	47.56	-182.62	131.51	0.75




Figure 7.1 C pools in forest soils for forest before 1990. a) Forest floor C in 2007-2009 versus 1990, b) Mineral soil C in 2007-2009 versus 1990. Lines: $y=x$.

Some mineral soils had one or several horizons of organic origin and these soils had very high soil C-stocks to 1 m depth (>300 tonne C ha⁻¹), and these will probably be handled separately in further work with the data. Determination of true changes in organic soils requires that the total depth of the organic layer is known, while soils were only sampled until 1 m in SINKS.

Carbon pools 2008-2012

Carbon pools in live and dead biomass estimated for the most recent rotation of the NFI (2008-2012) is 41,0 million tonnes C. The live above ground biomass carbon makes up about 82 % of the total carbon in biomass and dead wood makes up only 1.7 % of the total. Carbon in biomass in forests established after 1990 make up 3.0 % of the total.

Table 7.8 Carbon in forest biomass for NFI rotations with reference years 2008-2012.

			2008	2009	2010	2011	2012
Forests established before 1990	Area		540.124	539.600	539.076	538.553	538.189
	Live biomass	Above ground	31.109	31.011	31.395	32.563	33.347
		Below ground	6.086	6.087	6.160	6.403	6.580
	Dead wood		545	566	628	653	673
	Forest soil	Litter	6.042	6.059	6.577	6.837	7.091
		Soil					
Forests established after 1990	Area		79.787	84.033	88.279	92.525	94.358
	Live biomass	Above ground	735	782	840	886	885
		Below ground	162	169	178	184	179
	Dead wood		23	24	23	20	16
	Forest soil	Litter	342	359	381	378	380
		Soil					

The amount of carbon in biomass in forests established before 1990 has been slowly increasing since 2006. Based on preliminary results of an evaluation of the subsequent measurement cycles 2002-2006 and 2007-2011, the increase is at least partly caused by an increased average biomass per hectare. However, part of the increase is also due to an increase in forest area, which is caused by improved detection of forest caused by improvements of aerial photos used for this.




Figure 7.2 Forest carbon in forests established before 1990 estimated from NFI data from 2002-2011. Note that estimates for 2002-2005 are based on only 1-4 years of measurements. Only from 2006 the estimates are based on a full five-year rotation of the NFI.

The amount of carbon in biomass in forests established after 1990 has been increasing rapidly during the time of NFI measurements. The very low estimates of forest carbon at the beginning of the NFI measurements may in part be due to a large number of plots not measured in the field as a result of start up problems which may have biased the results. Also, in the early measurements aerial photographs were of a poorer quality and recent afforestations may have been difficult to detect.




Figure 7.3 Forest carbon in forests established after 1990 estimated from NFI data from 2002-2011. Note that estimates for 2002-2005 are based on only 1-4 years of measurements. Only from 2006 the estimates are based on a full five-year rotation of the NFI.

7.2.6 Uncertainties and time series consistency

Danish national forest inventories have developed over the years from the earliest inventories more than a century ago. More recently the development has been quite rapid, thus influencing the estimation of forest carbon pools in relation to LULUCF.

In the 1990 forest census the number of questionnaires sent to respondents was 22,300. In the subsequent inventory the number of respondents increased to 32,300. Not unexpectedly this led to a substantial increase in estimated forest area, which is not possible to separate from the actual increase in forest area that occurred during that period of time. Also, it is not possible to single out the effect of the increased number of questionnaires on estimates of species distribution, carbon pools etc.

In 2002, the sample based forest inventory released the previous forest census for the first time enabling annual forest statistics. The NFI includes areas and forest owners that have not previously been included in the forest census. Firstly because not every forest owner was included in the previous surveys and secondly because not all forest areas according to the FAO definitions would be perceived as forest by the respondents. Consequently, the change from questionnaire based forest census to sample based forest inventory has led to an increase in forest area estimates that is not possible to separate from the actual increase in forest area that occurred during that period of time.

Specifically, in relation to the reporting of carbon pools in forest, the change from questionnaire based forest census to sample based forest inventory has changed the calculation of forest volume, biomass and carbon. In the forest census, forest carbon is estimated from the reported forest area within different species, age and site classes and a number of forest growth models. In the forest inventory, forest volume (and subsequently carbon) is measured on the plots. The observed forest area and carbon is subsequently expanded to regions or the entire country using statistical models. This has led to a substantial increase in forest volume, biomass and carbon estimates, mainly due to methodological differences.

In the estimation of carbon emissions from existing forests, the information collected in relation to different forest census and inventories is combined with the satellite based land use/land cover map for the base year 1990 and for the year 2005. Hereby, consistent estimates of emissions from existing forests are obtained utilising as much information from the data sources as possible and hereby providing best possible time series. For the period from 2006 and onwards - there is full consistency of the data.

The uncertainty of the estimates of the carbon pools have been analysed by the use of bootstrap analysis. For the total carbon pool of the living biomass standard error is estimated to be 0.6 tonne C pr. ha or equalling 0.9 per cent. Applying the stock change method the emission/sink estimates of the different parts of the carbon pools depend on the certainty of each pool at two consecutive times.

The uncertainty of the estimates for subsets of the full forest area is related to the sampling intensity. With more subdivisions the uncertainty increases as the sampling size is reduced. An initial bootstrap analysis of this has been performed.

Table 7.9 Tier 1 estimate of the uncertainty in the forest.

		1990	2012	Activity data, %	Emission factor, %	Combined uncertainty	Total uncertainty, %	Uncertainty, 95 %, Gg CO ₂ eqv.
		Emission/sink, Gg CO ₂ eqv.	Emission/-sink, Gg CO ₂ eqv.					
5. LULUCF		5282.55	-838.44				24.7	207.4
5.A Forests		141.9	-4441.5				15.0	664.8
Forest remaining forest	CO ₂	49.6	-4491.4	15	2.00	15.1	15.1	679.7
Land converted to forest	CO ₂	76.6	37.8	15	8.74	17.4	17.4	6.6
Drainage of forest soils	N ₂ O	15.6	12.1	30	10	31.6	31.6	3.8

7.2.7 QA/QC and verification

A continuous focus on the measurements of carbon pools in forest will contribute to QA/QC and verification in the following submissions. As we gain more data through resampling of permanent plots in the NFI this will further support the verification of the data reported. These will be available for the reporting performed in 2013.

On-going development of the NFI in terms of sampling procedures and estimation methods is essential for the continued QA/QC process of the NFI.

Integration with multi-phase and multi scale inventory - through e.g. other in-situ data like LiDAR scanning or remote sensing like satellite imagery will through research contribute to the continued QA/QC process of the NFI and the carbon stock estimates for forests.

7.2.8 Recalculations and changes made in response to the review process

The estimation methods are similar to the last reporting. There are sampling errors, but basically the continuous sampling, with partial replacement, provide stable estimates of the carbon pools in forests.

7.2.9 Planned improvements

Below is a list of planned improvements.

- A further QA/QC of the Land Use matrix will be performed.
- Further analysis of the changes in forest area - including afforestation and deforestation - and if possible more detailed information on these will be collected and analysed.
- Documentation for carbon pools in soil and litter is expected to be further improved in the next submission.
- Documentation for extent of restored natural hydrology will be included if available and will mainly refer to the period of the NFI since 2002, as no systematic information is available prior to this time and changes are first detectable based on the full analysis of the re-measurements performed in 2011. This is expected to confirm the reduced drainage of forest areas.
- Further analysis of uncertainty estimates for all the carbon pools in the forest areas based on the remeasurements and bootstrap analyses.

7.3 Land converted to forests

7.3.1 Forest area

See section 7.2.1 Information on approaches used for representing land areas and on land-use databases used for the inventory preparation

7.3.2 Forest definition

See section 7.2.1 Land-use definitions and the classification systems used and their correspondence to the LULUCF categories (e.g. land use and land-use change matrix)

7.3.3 Methodological issues for land converted to forest

See also section 7.2.1

Living biomass

With respect to the option for distinguishing forest with and without harvesting, it is not possible with the available data. Data from the NFI is utilised based on the land use mapping to identify sample plots on afforested/reforested (AR) areas. It is - however not possible to determine the amount of harvesting. Furthermore, Denmark applies an approach utilising total carbon stock change, both growth and harvesting is included in the overall estimation.

When converting land to forest land the standing living above- and below ground biomass are removed from the land. In Table 7.10 the default values for the amount of living biomass is shown.

For land converted from cropland a standard default loss value of 9 577 kg DM (dry matter) per hectare in above-ground biomass and 2 298 kg DM per hectare in below-ground biomass is used. This value is equivalent to the average harvest of living biomass for all cereals grown in Denmark from 2000 to 2010, including straw, stubble and glumes. For conversion from DM to carbon a default fraction of 0.5 kg C per kg DM is used.

Table 7.10 Default values for the amount of DM (dry matter, kg per hectare) used for estimating carbon stock changes where land use conversions take place.

		Dry matter, kg DM pr hectare	
		Above ground biomass	Below ground biomass
Cropland		9 577	2 298
Grassland	Improved Grassland	2 400	6 720
	Unmanaged Grassland	2 200	6 160
Wetlands	Peat extraction	0	0
	Other Wetland	3 600	10 080
Settlements		2 200	2 200
Other land		0	0

Soils

The included soil carbon pool changes concerned only carbon sequestration due to development of forest floors, i.e. the organic layer on top of the mineral soil. Carbon sequestration was included in this layer since national scientific projects had indicated that this was the soil compartment mainly prone to changes following land-use change. The previous NIR reports did not account for possible changes in carbon pools of the mineral soil; based on chronosequence studies of afforested stands, no consistent changes had been detected in mineral soil organic matter during the first 30 years following afforestation (Vesterdal et al., 2002a; Vesterdal et al., 2007). This is also supported by the finding that there was no significant difference in mineral soil carbon stocks in paired forest-cropland sites at 28 different sites in Denmark (Vesterdal et al., 2002b). These conclusions are supported by data from the new national forest soil inventory.

New data

New information on carbon pools in forest soils is available from the national project, SINKS. In this project forest soils are sampled in two grids, Agricultural Network and the National Forest Inventory (NFI), see Section 7.2.1 for a description.

Apart from 108 plots in forests planted before 1990, the Agricultural Network included 15 plots in afforestation since 1990. The sampling took place together with the sampling in forests planted before 1990, and was thus complete for the period 2007-2009. Archived soil samples from 1990, when the plots were arable land, were missing for 1 plot.

The sampling, the sample preparation, chemical analyses and calculations were similar to that performed from forests planted before 1990, see 7.2.1

A few of the 277 plots sampled in the NFI grid are probably also located in forests planted since 1990. The data calculations are currently being performed, and these data will be reported in the next NIR.

7.3.4 Changes in forest soil carbon stocks in forests planted on arable land since 1990

The average carbon sequestration rates for forest floors for broadleaves and conifers were estimated from the information from scientific projects in afforestation chronosequences; the average annual sequestration of carbon in forests floors was 0.09 and 0.31 tonne C ha⁻¹ yr⁻¹ under broadleaves and conifers, respectively (Table 7.11.). These rates of change have been used for calculation of forest floor carbon sequestration in afforested land, however, the

accumulation of conifer forest floors is assumed to start only after eight years based on observations from chronosequence and other field data.

Table 7.11 Forest floor carbon sequestration rates in afforestation areas for different species in national chronosequential studies.

Species in national chronosequential studies.						
Tree species category	Tree species	Study type	Age (yr)	Forest floor C sequestration (tonne C ha ⁻¹ yr ⁻¹)	Source*	
Broadleaves	Oak	Chronosequence	29	0.08	1	
	Oak	Stand	30	0.02	2	
	Oak	Stand	30	0.05	2	
	Oak	Stand	30	0.04	2	
	Oak	Stand	30	0.02	2	
	Oak	Stand	30	0.13	3	
	Oak	Stand	40	0.09	3	
	Beech	Stand	30	0.09	2	
	Beech	Stand	30	0.10	2	
	Beech	Stand	30	0.12	2	
	Beech	Stand	30	0.13	2	
	Beech	Stand	30	0.18	3	
	Beech	Stand	40	0.14	3	
	Average (SEM)				0.09 (0.01)	
Conifers	Norway	Chronosequence	30	0.35	1	
	Spruce	Chronosequence	41	0.43	1	
		Stand	30	0.21	2	
		Stand	30	0.15	2	
		Stand	30	0.20	2	
		Stand	30	0.30	2	
		Stand	30	0.30	3	
		Stand	40	0.65	3	
		Sitka spruce	Stand	30	0.43	2
		Stand	30	0.24	2	
		Stand	30	0.22	2	
		Stand	30	0.25	2	
	Average (SEM)				0.31 (0.04)	

* 1) Vesterdal et al. (2007), 2) Vesterdal & Raulund-Rasmussen (1998), 3) Vesterdal et al. (2008).

The results from scientific projects have lately been checked by analysis of preliminary results from the Agricultural Network. The afforested plots in the monitoring grid also revealed large variation in soil carbon pools among for both forest floors and mineral soils (Table 7.12). The mean carbon pool of the forest floor among the afforested sites was about 2.5 t C ha⁻¹ in 2007-2009 (and supposedly 0 t C ha⁻¹ at the time of the afforestation) while the mean carbon pools for mineral soils were 114 and 108 t C ha⁻¹ in 1990 and 2007-2009 respectively (Table 7.18). A simple t-test on the mean change in mineral soils pools between 1990 and 2007-2009 (-1.87 t C ha⁻¹ yr⁻¹) showed that the change was not significant (Table 7.12 and Figure 7.2) while there, as expected, was a significant sequestration of carbon in the forest floor due to litterfall inputs and subsequent buildup of the organic layer (Table 7.13, Figure 7.3). The age of the afforested stands ranged from 8-19 years, so only the establishment phase was covered.

Table 7.12 Basic statistics on soil carbon pools measured in the "Agricultural Network".

	Mean C pool	Std	Min	Max
	(tonne C ha ⁻¹)			
Forest floor at the time of the afforestation	-	-	-	-
Forest floor 2007-2009	2.53	1.79	0.25	5.56
Mineral soil 1990	113.63	35.37	68.00	186.06
Mineral soil 2007-2009	107.83	41.25	52.82	220.06

Table 7.13 Statistics from a simple t-test on the change in soil carbon between ca. 1990 and 2009 for forests after 1990.

	Total number of sites	Number of sites in t-test	Mean change	Std	Min	Max	P-value
			(tonne C ha ⁻¹)				
O-horizon	15	15	2.53	1.79	0.25	5.56	<.0001
Mineral soil	15	14	-1.87	17.59	-35.32	34.00	0.70




Figure 7.4 Carbon pools in mineral soils in 2009 versus 1990. Forests established on arable land since 1990. Line: $y=x$.

The amount of carbon in the forest floors increased with the age of the afforested stand (Figure 7.4), while this was not the case for the mineral soil (Figure 7.5).




Figure 7.5 Forest floor carbon pools in forests afforested since 1990 in the Agricultural Network. The regression was forced through (0,0) ($C \text{ acc.} = 0.2057 \times \text{age}$, $R^2=0.3124$, $p<0.0001$).




Figure 7.6 Change in mineral soil carbon stocks for forests since 1990. Line: $y=0$ (Regression line (not shown): $R^2=0.0005$, $p=0.9356$).

Average carbon sequestration rates for forest floors for broadleaves and conifers were also estimated from “Agricultural Network” in order to check the forest floor carbon sequestration rates used in reporting; in this case the average annual sequestration of carbon in forests floors was 0.16 and 0.20 t C ha⁻¹ yr⁻¹ under broadleaves and conifers, respectively (Table 7.14). These values are lower compared to the values obtained from 30-40 yr-old stands. (Table 7.11).

Table 7.14 Forest floor carbon sequestration rates in afforestation areas for different species. Data from the “Agricultural Network”.

Tree species category	Tree species	Study type	Age (year)	Forest floor C sequestration (tonne C ha ⁻¹ yr ⁻¹)	Site
Broadleaves	Oak	Monitoring plots	14	0,19	837
	Oak	Monitoring plots	17	0,22	301
	Maple	Monitoring plots	9	0,06	485
	Lime	Monitoring plots	10	0,20	571
	<i>Average (SEM)</i>			<i>0.16 (0.07)</i>	
Conifers	Norway spruce	Monitoring plots	19	0,21	479
	Sitka spruce	Monitoring plots	13	0,41	335
	Sitka spruce	Monitoring plots	10	0,03	340
	Normann fir	Monitoring plots	13	0,31	31
	Normann fir	Monitoring plots	16	0,04	171
	Normann fir	Monitoring plots	12	0,13	235
	Normann fir	Monitoring plots	8	0,08	292
	Normann fir	Monitoring plots	12	0,24	689
	Silver fir	Monitoring plots	19	0,29	66
	Larch	Monitoring plots	8	0,43	334
	Mixed conifers	Monitoring plots	10	0,08	509
	<i>Average (SEM)</i>			<i>0.20 (0.14)</i>	

Lastly we combined all data to explore the trends in forest floor carbon stocks among broadleaves and conifers (Figure 7.7). The rates used seem reasonable, even if the inclusion of new data indicate that it might be too high for conifers in the stand establishment phase. Thus, accumulation of conifer forest floors is assumed to start after 8 years of chronosequences. This is reasonable since observations in chronosequences indicate that there is little litterfall in conifer stands to build up forest floors during the first 10 years.




Figure 7.7 Forest floor carbon pools in afforested plots. All available data from chronosequential studies and the “Agricultural Network” are included. Lines show the carbon sequestration rates used in the reporting: 0.31 tonne carbon ha⁻¹ yr⁻¹ for conifers and 0.09 tonne carbon ha⁻¹ yr⁻¹ for broadleaves.

Several previous national field studies mentioned above (Vesterdal et al. 2002a, 2002b, 2007) did not suggest measurable decadal changes in mineral soil carbon following afforestation. In the Forest Soil Inventory (SINKS project), soil carbon content to 100 cm in forest land remaining forest land was compared with soil carbon in the same depth found in a parallel project for cropland soils (Table 7.15). These data also support that mineral soils are neither sinks nor sources for CO₂ following afforestation of former cropland. Using a transition time of 50 years, these soil carbon contents were used to calculate the small rates of soil carbon stock change for cropland to forest conversion.

Table 7.15 Mineral soil carbon content (Mg ha⁻¹) in cropland and forest land based on the Agricultural Network. N: number of plots, mean and standard deviation (std).

Land use	Sandy soils			Loamy soils		
	N	mean	std	N	mean	std
Cropland		137			158	
Forest land	261	154	79	116	164	103
Grassland and Other land	19	150	84		150a	84a
Settlements		120 b			120 b	

^a Same data as for sandy soils.

^b Agreed with the UFCCC-ERT during the 2011 review.

In conclusion, preliminary results from the Forest Soil Inventory project show no evidence that mineral soil carbon pools for forests on former arable land are neither sinks nor sources for CO₂. The data from the SINKS project support the conclusions drawn from Vesterdal et al. (2002, 2007), Vesterdal and Raulund-Rasmussen (1998), and Vesterdal et al. (2008) for forest floors. The comparison between Danish land-uses (Table 7.15) suggests that particularly sandy soils would sequester carbon following afforestation of cropland, whereas carbon stocks in loamy soils are quite similar between land uses. Thus, a no-source principle would be justified in case of land-use conversions to forest.

Until final results from the Forest Soil Inventory are available we continue to use the previously used average carbon sequestration rates: 0.09 tonne carbon ha⁻¹ yr⁻¹ for broadleaves and 0.31 for conifers.

The sequestration of CO₂ in forest floors in forests established since 1990 has gradually increased and the annual CO₂ sequestration will increase much more over the next decades when cohorts of afforestation areas enter the stage of maximum current increment.

The reporting of the forest floor in the afforestation in the 2008-2012 period is based on the NFI monitoring of forest floor depth as described above.

7.3.5 Uncertainties and time series consistency

See Section 7.2.1 and 7.2.2 for recalculation since 1990.

7.3.6 QA/QC and verification

A continuous focus on the measurements of carbon pools in land converted to forest will contribute to QA/QC and verification in the following submissions. See also Chapter 7.2.1

7.3.7 Recalculations, including changes made in response to the re-view process

In the updated land use matrix that now includes mapping of three years: 1990, 2005 and 2011, significant changes have been noted related to land use and land use changes. This includes increased afforestation in areas without support from public funds. This includes establishment of minor forests areas, to improve hunting options and to produce biomass. Some forest areas have been established through natural succession, a method now approved by the Forest Act (from 2005). In the previous reporting, mainly afforestation based on subsidies were expected and included in the reporting.

7.3.8 Planned improvements

A QA/QC of the Land Use matrix is a continuous process.

The basic information utilised to give the data for the emission estimates for units of land subjected to afforestation/reforestation is based on National Forest Inventory (NFI) observations of stock change, specific related to the afforested areas. This will include all changes in carbon pools - also if affected by harvest - including thinnings of young stands. Based on the NFI it will be possible - for the next reporting to provide some indications of the frequency of harvesting/thinning occurring on the afforested areas. Given the fact that the afforested area is still a relatively small part of the full forest area, there will be more uncertainty on the estimate related to afforested areas compared to the area of forest remaining forest.

Documentation for carbon pools in soil and litter is expected to be further improved in the next submission.

7.4 Cropland (5B)

7.4.1 Cropland and cropland management (5B1)

The total Danish cropped agricultural area of approximately 2.7 million hectare can relate to approximately 620 000 individual fields, which again is lo-

cated at 200 000 land parcels. This gives an average field size of less than four ha. The actual crop grown in each land parcel (LPIS) is known from 1998 and onwards. Since 1990 the agricultural area recorded by Statistics Denmark has decreased from 2.78 million hectare to 2.64 million hectare (Table 7.16). The total crop yield given as kernel, root fruits and grass as measured in dry matter (million kg dry matter per year) is, however, at the same level and increasing due to improved cropping techniques, Figure 7.8




Figure 7.8 Total crop yield given as kernel, root fruits and grass as measured in dry matter (Million kg dry matter per year)

The main reason for the loss of land for agricultural purposes is urbanisation and afforestation. The major part of the agricultural area is grown with annual crops: cereals, grass in rotation, oilseed, sugar beets, potatoes and temporarily set-a-side. Permanent grass outside rotation with none or very little fertiliser application rates (>63 kg N per ha per year) is reported under Grassland. All fertilisation with nitrogen is reported under Agriculture 4D2.

Table 7.16 shows the development in the agricultural area from 1990 to 2012 (Statistics Denmark). A general trend is a continuous decrease of 6 000 - 7 000 ha per year in the agricultural area.

Table 7.16 Cropland area in Denmark 1990- 2012 according to Statistics Denmark and the Land Use Matrix, hectares.

	1990	1995	2000	2005	2010	2011	2012
Annual crops (CL) 1	2 236 535	1 969 275	1 938 633	1 953 306	2 049 304	2 050 108	2 053 093
Grass in rotation (CL)	306 325	310 568	330 834	342 417	327 319	336 061	331 512
Permanent grass (CL and GL)	217 235	207 122	166 261	192 968	199 859	186 652	200 413
Horticulture – vegetables (CL)	16 428	12 915	10 803	9 557	10 812	11 215	10 305
Perennial fruit trees – perennial wood- en crops (CL)	10 267	10 669	9 892	9 464	8 181	7 477	7 570
Set-a-side and other land (CL)	3 861	217 801	192 441	200 751	51 309	48 273	41 800
Total agricultural land area reported by Statistics Denmark	2 788 276	2 726 048	2 646 982	2 707 236	2 646 400	2 639 905	2 644 631
Willow and other crops for energy purposes (CL)	588	588	695	1 320	4 049	4 795	5 410
Hedgerows (CL)	61 326	61 019	60 554	60 170	59 791	59 732	59 659
“Other agricultural land”	80 783	107 253	119 847	61 115	105 720	94 385	86 433

¹CL refers to that the area is treated under Cropland. GL refers to Grassland.

Cropland area

The Cropland area is defined as the agricultural area as given by Statistics Denmark, Perennial wooden crops (fruit trees, orchards and willow), hedgerows (perennial trees/bushes not meeting the forest definition) in the agricultural landscape and "Other agricultural land". The latter is defined as the difference in the area between the total Cropland area as defined by land use matrix (see Table 7.3) minus agricultural crops in rotation as given by statistics Denmark minus the area with fruit trees and the area with hedgerows. "Other agricultural land" is thus comparable small areas and probably without agricultural and wooden crops, which cannot be allocated to other land use categories. In the inventory carbon in living biomass for "Other agricultural land" is given the same value as for annual crops so that inter-annual changes in the cropland area from Statistics Denmark are eliminated.

The area with Perennial wooden crops are the area given by Statistics Denmark and for some categories it is split further down with data from the EU crop subsidiary system, which gives information on which crops are grown where on species level.

The main data for land use in Cropland (5.B.1) is the agricultural area given by Statistics Denmark. Both annual agricultural and wooden perennial crops are allocated into grids (climatic, soil type and municipality) with the help of the EU Land Parcel Information System (LPIS). LPIS contains information of the exact position of the field. The survey data from Statistics Denmark differs a little from the LPIS system ($<\pm 2\%$ for the major crops). Area and yield data from each region is used for the calculations as reported by Statistics Denmark.

The area with hedgerows is based on analysis of aerial photos from 1990 and 2005 combined with planting and removal statistics of hedges from the Ministry of Food, Agriculture and Fisheries. The major part of the hedge erection is subsidies in Denmark and therefore monitored.

Cropland definition

The land area under "CL" consists of: Cropland with annual crops, cropland with wooden perennial crops, area with hedgerows and "Other agricultural area". The latter consists of small undefined areas lying inside the area, which is allocated as cropland in the cropland area.

For purposes of the calculations for annual crops a division as follows is used: Winter and spring wheat, rye, triticale, winter and spring barley, oat, winter and spring rape, grass for grass seed production, grassland in rotation, potatoes, sugar beets, peas, maize for silage, cereals for silage, vegetables and miscanthus.

For purposes of perennial wooden crops a division as follows is used: Apple, Pears, Cherries, Plumes, Rosehips, Elderberries, Hazel and Walnuts, Grapes, Other fruit trees, Black current, Other fruit bushes, Hedgerows and Willows.

Biomass from Christmas trees in the agricultural area is reported under forests.

Cropland - Methodological issues

The following data sources are used for determination of cropland area, for determination of any land-use changes, for allocation of natural and administrative parameters, for development of emission factors for soils and biomass and for calculation of carbon stocks in soils and biomass at various times.

- Agricultural area data from Statistics Denmark, 1980 to 2012
- Area and harvest surveys from Statistics Denmark, 1980 to 2012.
- Area with willow from the agricultural subsidiary system.
- EUs Land Parcel Information System, 1998 and onwards (grown crops on field and soil level).
- Digital soil map, 1:25.000.
- Arial photos of hedgerows in 1990 and 2005.
- Hedgerow planting data 1977 to 2012.
- Lime consumption data 1990 to 2012.

The model for carbon stock changes in hedges is based on a growth model from the National Forest Inventory (NFI) classified into plant and soil type and height.

Emissions from living biomass

For annual agricultural crops on cropland remaining cropland (5B1) it is assumed that no changes in above-ground, below-ground, dead biomass and litter are occurring cf. IPPC 2003 (3.3.1.1.1). The variations in the actual agricultural area collected by Statistics Denmark may be up to 100,000 hectares per year. When estimating the carbon stock in living biomass such changes may create large variations between years, which may be artefacts. As the amount of living biomass is defined according to the time where the peak of living biomass is occurring the variation in the area from Statistics Denmark create large fluctuations in the carbon stock in living biomass compared to other sources. To counteract this problem the sub-division "Other agricultural land" has been created with a default carbon stock of living biomass as in the designated agricultural area. The default carbon stock in living biomass is equivalent to an average spring barley crop with aboveground biomass of 9 577 kg DM (dry matter) pr hectare and a below ground DM of 2 298 kg pr hectare. Default dry matter values for the different crop categories used in the inventory was given in Table 7.10.

Fruit trees and other perennial wooden plants

Fruit trees, other perennial commercial wooden plants and durable horticultural plantations are reported separately under Cropland (Table 5.B). These are only of minor importance in Denmark. The total area for different main classes and the used carbon stock in above-ground and below-ground biomass are given in Table 7.17. Due to the limited area and small changes between years the CO₂ removal/emission is calculated without a growth model for the different tree categories. Instead the average stock figures are used in Table 7.17 multiplied with changes in the area to estimate the annual emissions/removals. Perennial horticultural crops account for approximately 0.07 % of the standing carbon stock. Christmas trees are reported under forest (5.A).

The carbon fraction of dry matter (DM) is assumed to be 0.5 for all species. For parameter estimation of living biomass, see Gyldenkerne et al. 2005 for fruit trees, for willow and *Miscanthus*:

Table 7.17 Mg living biomass per hectare and area, ha, with perennial wooden trees and – bushes, 1990- 2012.

	Living biomass, Mg DM per ha	1990	1995	2000	2005	2007	2008	2009	2010	2011	2012
Black currant	5,20	1 269	1 828	1 492	2 001	1 856	2 071	1 848	1 935	2 041	1 855
Other berries	5,20	663	547	611	698	708	589	578	533	608	734
Rosehip	13,99	0	0	0	0	0	120	159	197	197	34
Cherries	25,45	1 787	2 653	2 804	2 131	2 169	1 951	1 864	1 743	1 466	1 401
Plumes	25,45	0	0	0	0	0	60	63	68	65	73
Hazelnut and Walnuts	25,45	0	0	0	15	13	45	14	14	23	28
Aples	33,76	2 726	1 658	1 678	1 751	1 812	1 797	1 730	1 684	1 550	1 703
Pears	13,99	351	546	441	413	466	442	372	357	336	344
Elderberry	25,45	0	0	0	9	15	10	12	9	16	14
Grapes	5,20	0	0	0	18	22	31	37	45	50	57
Other fruit trees	13,99	0	0	0	110	125	21	48	60	74	67
Rowan-berries	33,76	0	0	0	0	0	14	16	16	10	12
Willow	17,43	588	588	695	1 320	1 736	1 832	2 736	4 049	4 795	5 410
Miscanthus	17,43	1	1	6	33	88	80	80	156	774	70
Total		7 385	7 821	7 727	8 499	9 010	9 062	9 556	10 865	12 005	11 804

Hedgerows

Since the beginning of the early 1970s governmental subsidies have been given to increase the area with hedgerows to reduce soil erosion. Annually financial support is given to approximately 600-800 km of hedgerow. There are no figures on how many hedgerows have been removed in the same period as these to a large extend are not protected. Therefore 144 aerial photos on a 2x2 km² square for 1990 and 2005 have been analysed to monitor and detect changes in the landscape. The squares are distributed throughout Denmark in a stratified way according to primarily soil and wind conditions (Figure 7.9). A very large dynamic in the location of the hedges between 1990 and 2005 was observed (Figure 7.9). Only areas not meeting the definition of forests and areas not classified under Perennial Wooden crops (fruit trees, willows etc.) were included in the analysis. The hedges were further allocated into eight different regions, mainly according to soil type (e.g. growth pattern).




Figure 7.7 Designated areas with different types/classes of hedgerows.




Figure 7.9 The dynamics of hedgerows in the Danish Landscape 1990 to 2005. Blue colour indicates no changes, red colours are removed hedges and green colours are new hedges (Source: M. Fuglsang, DCE).

The overall results from the analysis of hedges are shown in Table 7.18. The total area with hedges has decreased with 2 % but the total volume and the carbon stock has increased due to changed sizes and composition.

Table 7.18 Hedges in the cropland 1990 and 2005.

	1990	2005	Change in percent 1990-2005
Area, ha	61 326	60 093	-2.0
Volume, million. m3	4 139	4 402	6.4
Carbon stock, Gg	939	1 072	14.2

In Table 7.19 the actual planting and removal rates for hedgerows is shown. The 1970s and 1980s have a high concern to protect and maintain the hedgerows and a substantial replacement took place. Currently is the governmental subsidiary targeted to broadleaved hedgerow replacing old single-rowed conifers (mainly white spruce (*Picea glauca*)). In 1990 75 % of the replaced conifers hedgerows were replaced with 3- to 6-rowed broad-leaved hedges. In 2005 only 20 % are replacements and the remaining is new hedges cf. Table 7.19. Over the years a decrease in the number of subsidized hedgerows has taken place. The Ministry of Food, Agriculture and Fisheries is responsible for all administration, registration and mapping of all subsidised hedgerow planting in Denmark.

Table 7.19 Hedges planted and removed under the governmental subsidiary system 1985 to 2012.

	1985	1990	1995	2000	2005	2008	2009	2010	2011	2012
Planted 3-rowed, km	1 082	928	560	852	390	141	128	109	96	107
Planted 6-rowed, km	0	0	252	250	115	41	38	29	37	33
Planted small biotopes, ha							96	64	52	33
Percentage removed, %	75	75	36	27	20	20	20	20	20	20
Percentage new, %	25	25	64	74	80	80	80	80	80	80
Hedges removed, ha	608	522	218	219	76	83	25	21	20	21

The biomass estimation of the hedges is based on measurements made in the Danish NFI where plots with similar height and plant species are used as transfer functions (Figure 7.10).




Figure 7.10 Biomass function estimated as m^3 biomass per m^3 versus tree height in NFI plots less than 15 meter (Courtesy Thomas Nord-Larsen, SL, LIFE, KU).

Emission from soils

Based on a GIS analysis of the data in the LPIS and a newly produced soil map of the organic soil the agricultural area is distributed between mineral soils and organic soils and subdivided into cropland and permanent grassland.

Mineral soils – 5B1

For carbon changes in for agricultural crops a 3-pooled dynamic soil model is used (Petersen, 2003; Petersen et al. 2002, 2005, 2010, Gyldenkerne et al. 2005) to calculate the soil carbon dynamics in relation to the Danish commitments to UNFCCC. C-TOOL is only used in CL. No change in the carbon stock in soils under perennial wooden plants, hedgerows and “Other agricultural cropland” is expected and reported as NA. These areas are also only a very minor part of the cropland area. For agricultural crops C-TOOL is run on a regional level.

C-TOOL

C-TOOL is a 3-pooled dynamic model, where the approximate average half-life times for the three different pools, Fresh organic matter (FOM), Humified organic matter (HUM) and ROM (Resilient Organic Matter) are 0.6-0.7 years, 50 years and 600-800 years, respectively. The main part of biomass returned to soil each year is in the first and easiest degradable FOM pool. This pool consists of mainly fresh straw, fresh manure, root residues, fungi and small animals and fluctuates very much between years depending on the harvest yield and climatic conditions. A simple diagram of C-TOOL is shown in Figure 7.11.

C-TOOL is parameterised and validated against long-term field experiments (100-150 years) conducted in Denmark, UK (Rothamsted) and Sweden and is “State-of-the-art”. A detailed description of C-TOOL can be found at www.agrsci.dk/c-tool/index.htmls. More recent investigations have shown that C-TOOL is not properly parameterised on soils having more than 6 % organic carbon. Soils having 6-12 % organic carbon is therefore treated as or-

ganic soils with an emission factor of 50 % of organic soils > 12 % organic carbon.




Figure 7.11 A simple diagram of C-TOOL. Please refer to www.agrsci.dk/c-tool for more information.

Input data to C-TOOL and out put

As carbon input to each region for each year is taken the actual crop area and crop yield from Statistics Denmark for that particular region and crop species as given by Statistics Denmark (www.dst.dk Table AFG, AFG07, HST7 and HST77). The dry matter content depends on the actual crop. For cereals it is 15 %.

The amount of agricultural residues returned to soil is the amount estimated by Statistics Denmark (www.dst.dk Table HALM and HALM1). The dry matter content depends on the actual crop. For cereals it is 16 %.

More detailed figures on the distribution as an example of the crop yield and areas are given in Annex 3F, Table 3.F10-12 for a specific region (Eastern Jutland) in year 2010.

The overall input to C-TOOL varies between years (Figure 7.8) due to the actual growing conditions in that year. 2012 was a fairly good year in terms of harvested crops. The cereal yield was the third highest since 1990 and on a general decreasing area. 2010 and 2011 were medium years, whereas 2009 were the best cereal year ever. The variation in the input to C-TOOL gives a large inter-annual variation in the carbon input to the soil for all years of the time series. Combined with large inter-annual differences in the temperature this creates large inter-annual differences in the net carbon stock change in

mineral soils, where low yields combined with high temperatures reduce the total amount of carbon in agricultural soils, whereas in years with a high yield and low temperatures the carbon stock in soils is increased.

The amount of animal manure produced and applied to soil is estimated with the same methodology as in the Agricultural sector for estimating CH₄ and N₂O emission where annually updated feeding and excreting data are provided for the regulation of the animal production in Denmark. Here detailed data on the number of animal, housing and manure type are available on farm level. This also includes data whether the manure has been bio-gassed or not. The manure data are used as input to C-TOOL.

In Figure 7.12 is shown the overall input of C to the agricultural soils. Due to a ban of field burning in 1990, increased management and demand on catch crops an increase in the C input to soils can be seen.




Figure 7.12 Calculated C input to the soil (red squares) and modelled (blue curve) development of the FOM pools, million tonnes C.

Since 1997 there has been a demand for growing N catch crops in Denmark in order to reduce N leaching. Besides reducing the N leaching these crops increase the carbon stock in the soil. Between 120 000 and 200 000 hectares of the agricultural area has this additional crop every year. The demand for catch crops has altered the way of farming in two main ways. For farmers with cattle the farmers are sowing grass seed in their normal cereal fields. This grass sword must not be ploughed into the soil before winter/next spring. For farmers growing grass seed, which is common in Denmark, old grass seed fields are not ploughed before next spring in contradiction to the current situation where it would be ploughed early autumn. It has been estimated that the obligatory catch crops are increasing the amount of C returned to soil with 0.36 to 2.14 tonnes carbon per hectare per year (Olesen et al. 2004). The area with catch crops in each region is estimated from each farms obligatory N accounting, in which the area of catch crops area given on farm level (www.pdir.dk).

C-TOOL is initiated with data from 1980 and run multipliable times until stability, before the emissions from 1980 and onwards was calculated. Actual monthly average temperatures are used as temperature driver. The main drivers in the degradation of soil biomass are temperature and humidity. The Danish climate is quite humid with winter temperatures around zero degrees Celsius and hence the importance of soil humidity on the model outcome is low in contradiction to temperature, which has a high effect on the emission. As mentioned, when biomass is returned to the soil the major

part of it is quite easily degradable. Warm winters with unfrozen soils in connection with high inputs of biomass will therefore, as a result, yield high emissions from the soil compared to more cold years, which will yield low emissions.

In recent years (1999- 2012) Denmark has experienced very warm winters although 2010 was very cold and below the average from 1961 to 1990. In 18 out of the last 20 years the annual average temperature has been above the average temperature from 1961 to 1990. Year 2012 had an average temperature of 8.3 °C or 0.6 °C above the average from 1961 to 1990.

Year 2006 resulted in a high loss due to the warmest year up to now combined with a harvest yield 5 % below the average for 1997 to 2009 (measured as kernel yield from cereals) (Figure 7.13). In this year the organic matter input from crop residues and animal manure were not able to compensate for the loss (Figure 7.13). 2007 was not as warm, which led to an increase in the carbon stock. 2009 was cooler than 2008 but 2009 gave the highest cereal yield ever monitored in Denmark despite the fact that the agricultural area has decreased since 1980. This led to a very high input of organic matter into the soil, which again increased the soil carbon stock.

2010 were very cold with low harvest yields and 2011 were moderate too due to draught. An overall decreasing C stock in mineral soils is therefore estimated.

The combination of a higher yield in 2012 than in 2011 combined with a lower temperature in 2012 than in 2011 has resulted in a lower C loss from soils in 2012 than in 2011.

The FOM-pool (Fresh Organic Matter), which has a very fast turnover rate, consists of approx. 1.0 % of the total carbon content in the agricultural soil. Because of its large fluctuation between individual years and its small impact on the overall trend in the long-term development of the carbon stock in the soil, it has been agreed with the previous ERT during the in-country review in 2010, that all input sources are included in the modelling but in the reporting on the development an instant turnover of the FOM pool is used. The reported development is thus the two pools, HUM (Humified Organic Matter) and ROM (Resilient Organic Matter) which account for 99 % of the total amount of carbon in the soil. Figure 7.13 shows the development in the two pools. As can be seen there is a small increase in the total modelled carbon stock from 2008 to 2010 but a decrease in HUM and ROM. A new warm year with normal harvest yields will speed up the degradation of the FOM pool and as a consequence the two lines will get closer again.




Figure 7.13 The development in the C-stock in agricultural soils, Tg C (million tonne C).

As a whole the modelled emissions are found to be the most realistic emissions estimates for Denmark. As described in the agricultural sector the Danish farmers have faced increased demands for lower environmental impact since the mid-1980s. The general effect on the carbon stock in soil is that the 1980s showed a decrease in the carbon stock. In the 1990s the carbon stock seemed to stabilise due to the higher input of organic matter. Due to the increased global warming a declining carbon stock was modelled between 2000 and 2011. Since 1990 C-TOOL has estimated a loss of 1.5 % of the total carbon stock in the mineral agricultural soils. No precise uncertainty calculation has been made. However, it must be assumed that uncertainty in the estimate in the annual loss/gain is around 25 %. As Denmark has very good data on harvest yields and area data the uncertainty in the trend is very low. The estimated annual amounts of carbon in the agricultural soils are given in Table 7.20.




Figure 7.14 Estimated annual emissions from mineral soils 1990 to 2012 (million tonne CO₂ yr⁻¹).

Table 7.20 Modelled carbon stock (0-100 cm) in mineral soils from 1980 to 2011, Tg C.

Year	All pools	Fast FOM model
1980	433,795	428,295
1981	431,976	427,201
1982	430,412	426,068
1983	428,834	424,763
1984	428,764	423,954
1985	428,208	423,168
1986	427,508	422,382
1987	426,710	421,654
1988	425,496	420,625
1989	424,394	419,568
1990	425,419	419,182
1991	425,356	418,920
1992	423,752	418,029
1993	425,330	417,983
1994	423,994	417,545
1995	423,646	417,254
1996	424,436	417,329
1997	423,524	417,032
1998	424,346	416,949
1999	422,883	416,516
2000	423,050	416,266
2001	422,716	416,055
2002	421,398	415,406
2003	421,410	415,075
2004	421,323	414,798
2005	421,127	414,534
2006	419,695	413,975
2007	419,908	413,481
2008	419,755	413,113
2009	419,969	413,014
2010	419,354	412,700
2011	419,058	412,432
2012	418.897	412.274

Independent verification of C-TOOL

An independent validation of C-TOOL has been performed by soil sampling in the Danish Agricultural grid. The grid was established in 1987 and in a 7 x 7 km² grid square. In 1987 > 600 agricultural plots were sampled and analysed for carbon. Half of them were resampled in 1998 and a full resampling of 464 plots was made in 2008/2009. Figure 7.15 shows the development in the carbon stock in 0-100 cm depth in the paired plots. It can be seen that there has been an increase in the soil C stock in the sandy soils (Coarse Sand, Fine Sand and Loamy Sand). This is mainly due to that the Danish cattle herd is located on these soils combined with large areas with grass in rotation. This favour the soil C stock. Contrary to this is observed a loss in the C stock on the loamy soils (Sandy Loam and Loam). On these soils are annual crops the most common cultivars combined with a limited number of cattle and pigs. On these soils it seems difficult to maintain the soil C stock. Although there is some variability the overall conclusion is that there is a small loss from the Danish agricultural soils.

C-TOOL has estimated an overall loss from 1987 to 2009 of 7-10 million tonnes C and in the soil sampling grind is found an average loss of approx. 5 tonnes C per ha. With approx. 2 million hectares in rotation this gives a total

loss of 10 million tonnes C from 1987 to 2009. The conclusion is therefore that the modelled outcome from C-TOOL represents a proper value for the development of the carbon stock in the Danish agricultural soils.




Figure 7.15 The change in carbon stock in soil (0 - 100 cm) in >460 paired agricultural plots from 1987 to 2009 (Olesen et al. in prep.)

Organic soils - 5B1

A complete new soil map of the organic soils was made in 2010 for the inventory (Figure 7.16). The new soil map is a statistical map based on >10 000 soil samples down to the mineral soil in 30 cm intervals combined with a very detailed digital elevation map (DEM) for each 1.6 × 1.6 m² covering the entire Denmark, water table maps and old maps with organic soils. The definition of an organic soil in the new map is 20 % organic matter with a depth of minimum 30 cm (Greve et al., 2013, submitted). The total area with organic soils has been estimated to approx. 106 642 ha.



Figure 7.16 The new organic soil map for Denmark for year 2010, > 12 % OC (Greve et al. 2012, submitted).

On top of the organic soil map digital maps has been laid a map where 99 % of all Danish farmed fields (>636 000 fields) from the EU subsidiary system are precisely mapped with an uncertainty down to ± 0.5 meter. The actual grown crop is known for each field. In total more than 270 different crop types or combination of crop and crop management are recorded. In 2012 40 706 hectares with annual crops were located to be grown on the organic soil area and 10 180 hectares with grass in rotation.

The previous Danish soil classification was carried out in 1975. In 1975 it was estimated that there were 178 000 hectares of organic soils (>12 % C). Of this were 118 000 ha in the Cropland and the Grassland area and the remaining 60 000 ha were located in the Forests, Wetlands, Settlements and Other land. Overlay between the field map and the soil map has shown that only around 51 000 hectare in 2012 is farmed in the Cropland area and 17 200 hectare in Grassland and that the depth of the organic layer has become very shallow. The major reason for the drastic reduction is that Denmark is quite flat with shallow organic layers, which combined with intensive agricultural utilisation with high drainage rates has oxidized a major part of the organic matter.

The outcome is that during recent years more and more previously organic soils do not qualify to be organic by definition and that the area will decrease rapidly in future.

Emission factors for organic soils

An intensive research programme has been carried out to monitor the CO₂ emission from three organic soils in Denmark with annual crops in rotation and permanent fertilized grassland (Elsgaard et al. 2012, in prep). The overall result is shown in Table 7.21 compared with the IPCC default values. Maljanen et al. (2010) recently reviewed the GHG balance of managed organic peatlands in the Nordic countries. For areas with agricultural grasslands, the available studies suggested a net CO₂ emission of 4.9 ± 3.2 t C m⁻² yr⁻¹ (mean \pm standard deviation, n = 4). The available studies (n = 4) represented three Finnish and one Norwegian site (Lohila et al., 2004; Maljanen et

al., 2001, 2004; Grønlund et al., 2008). The upscaled annual emission from the Danish declining carbon stock is in line with these figures when taking into account the differences in temperatures. Considering that the IPCC temperate cold zone covers the major part of Europe the measured Danish values also seems to be in line with the IPCC guidelines. Emissions from organic soils on permanent grassland are reported under Grassland (CRF Table 5.C.1).

The dominating use of the organic soils is fertilised annual crops and grass in rotation. As C-TOOL has shown not to be able to simulate the emissions from soils having >6 % organic carbon fixed emission factor have been used for this area. No data has been found in the literature as they in the scientific world do not qualify as organic and hence little attention has been paid to these soils. Normally mineral soils in equilibrium will have an organic matter of 1-4 % organic carbon. Soils having higher contents are most likely developed under humid conditions with low degradation rates. Drained and managed soils having > 6 % organic carbon can therefore not be seen as being in their equilibrium state and will evidently lose carbon. We have therefore decided to allocate an emission of 50 % of what we have measured for soils > 12 % organic carbon in an attempt to account for these losses. These emissions are reported under 5B organic soils.

Table 7.21 Emission factors from organic soils, tonne C per ha per year.

	Cropland		Grassland	Uncertainty
	Annual crops and grass in rotation	Fertilised permanent grass	Permanent grass	Percent
Soils > 12 % OC	8.7	5.17	1.25	90
Soils 6-12 % OC	4.36	2.59	-	90
IPCC, Cold temperate	5.0		1.25 ^a	90
IPCC, Warm temperate	10.0		2.5	90

^a There seems to be an error in the guidelines on the emission from grassland. It is assumed that the figure should be one fourth of the emission from annual crops (5 t C per ha).

As emission factor for N₂O the IPCC 2003 default value of 8 kg N₂O-N per ha per year is used for the area with > 12 % organic carbon. This emission is reported in the agricultural sector, 4D2.

To estimate the emission from the organic soils a linear decrease in the area with organic soils between 1975 and 2010 has been assumed. All CO₂ emissions from organic soils converted from other Land Use categories to Cropland are reported under 5.B.1 and not under the respective land use conversion classes 5.B.2.1 to 5.B.2.5. The related N₂O emission is reported in the agricultural sector in CRF Table 4.Ds1.

The total emissions from the organic soils are given in Table 7.22.

Table 7.22 Emissions from cropland organic soils 1990 to 2012.

	1990	1995	2000	2005	2008	2009	2010	2011	2012
Cropland, 6-12 % OC, ha	46 270	43 045	39 820	36 595	34 660	34 015	33 370	32 734	32734
Cropland, > 12 % OC, ha	74 473	69 282	64 092	58 901	55 786	54 748	53 710	52 687	50886
Cropland, total, ha	120 743	112 327	103 912	95 496	90 446	88 763	87 080	85 420	83620
Emission, total, Gg C	-787	-733	-678	-623	-590	-579	-568	-558	-540
Emission, total, Gg CO ₂	-2 887	-2 686	-2 485	-2 284	-2 163	-2 123	-2 082	-2 045	-1981

Uncertainties and time series consistency

A Tier 1 uncertainty analysis has been made for part of the LULUCF sector cf. Table 7.21. The uncertainty in the activity data for the agricultural sector is very low. The highest uncertainty is associated with the emission factors. Especially the emission/sink from mineral soils and organic soils has a high influence on the overall uncertainty.

The LULUCF sector contributes to a large extend to the total estimated uncertainty. In recognition of the difficulties in analyses of uncertainty, the estimated uptake of CO₂ in the forestry sector must be treated with caution.

Table 7.21 Tier 1 uncertainty analysis for Cropland for 2012.

		1990	2012					
		Emission/sink, Gg CO ₂ eqv.	Emission/sink, Gg CO ₂ eqv.	Activity data, %	Emission factor, %	Combined uncertainty	Total uncertainty, %	Uncertainty 95 %, Gg CO ₂ eqv.
5.B Cropland		4232.3	2763.9				67.2	1856.8
Living biomass	CO ₂	-73.5	193.0	10	50	51.0	51.0	98.4
Dead organic matter	CO ₂	2.9	12.2			51.0	51.0	
Mineral soils	CO ₂	1415.3	577.1	10	75	75.7	75.7	436.6
Organic soils	CO ₂	2887.3	1980.7	10	90	90.6	90.6	1793.6

The time series are complete.

QA/QC and verification

A general QA/QC plan is developed for cropland. The following Points of Measures (PM) are taken into account.

- Collection and error check on in-data.
- Control of sums.
- Comparison with other data.

The area estimates for cropland and grassland in 2012 are very precise due to unrestricted access to detailed data from EUs Integrated Administration and Control System (IACS) on agricultural crops on field level and the use of the vector based Land Parcel Information System (LPIS). This access includes both Statistics Denmark and DCE. The total uncertainty in the major crop data is estimated by Statistics Denmark to be <2 %. Together with detailed soil maps this gives a unique possibility to estimate the agricultural crops on different soil types and hence track changes in land use. However, IACS and LPIS are only available from 1998 and onwards, and estimates for 1990 are therefore more uncertain. The QA of crop data is made by Statistics Denmark.

Data on newly planted and removed hedgerows are based on subsidised hedgerows and QA is carried out by the Ministry of Food, Agriculture and Fisheries, who is responsible for the administration of the subsidy scheme. The uncertainty in the number of plants used for the hedgerows is not estimated but is assumed to be very low because of the subsidy system.

There is an unknown uncertainty in the number of un-registered removal of hedgerows. A linear approach has therefore been made for "missing" hedgerows over the years. Establishment of wetlands is based on vector maps received from every county in Denmark. The uncertainty is not estimated but assumed to be very low due to the subsidised system.

As shown in Figure 7.12 and 7.13 the loss estimated by C-TOOL seems very close to the results from 464 paired soil samples.

A range of experts from the Faculty of Agricultural Sciences, Aarhus University, are repeatedly involved in discussions and report writings on topics related to the inventory.

Recalculations, including changes made in response to the review process

Recalculations have been made for emission estimates which are affected of the land use matrix due to the small changes. This has only a small influence on the emission estimate as the major sources are the mineral and the organic soils in cropland remaining cropland.

N₂O emissions from Grassland and Wetlands converted to Cropland have been included. These emissions account for 0.034 % of the total emission from cropland.

All recalculations have reduced the overall emission level from cropland with 190 Gg CO₂ in 1990 and with 180 Gg CO₂ in 2011 equivalent to app. 0.3 % of the total emission from cropland

All changes have been implemented for all years.

Planned improvements

A new version of C-TOOL has been developed. It is expected that the new version will be used in the next submission. Verification and investigation of the hedgerows will take place in 2016. A new soil sampling in the agricultural network is planned in 2018/2019.

7.4.2 Land converted to cropland (5B2)

Agriculture covers more than 63 % of the total area giving a large impact on the environment. As a consequence there are many initiatives to transfer agricultural land into natural habitats and forest, and the continuous development of infrastructure demands more land. Land converted to cropland is therefore not an issue. The largest challenge is that the farmers in one year may report that a certain field is cropland and the next year is permanent grassland where it could stay for several years before it again is ploughed and turned into annual cropland for one year. Despite or rather because of the detailed information which is available, is it impossible to have a conservative land use transition between these two land use categories. The new land use matrix showed that 49 600 hectares were converted from GL to CL from 1990 to 2012 and that 74 740 hectares were in a transition stage from CL to GL. The difference between these two figures also indicate a conversion to less intensive agriculture. No conversion from the other land use categories to CL has been found.

Approaches used for representing land

The area converted from other land use to Cropland is based on remote sensing of the Danish area in 1990, 2005, 2011 and 2012 combined with data in LPIS on which crops are grown in each field.

Methodological issues

Change in carbon stock in living biomass

For land converted to cropland a standard default gain value of 9 577 kg DM (dry matter) per hectare in above-ground biomass and 2 298 kg DM per hectare

ture in below-ground biomass is used. This value is equivalent to the average harvest of living biomass for all cereals grown in Denmark from 2000 to 2010, including straw, stubble and glumes. For conversion from DM to carbon a default fraction of 0.5 kg C per kg DM is used.

For conversion from cropland to other land use categories the same value is used but recorded as a loss of carbon in the respective category (5A2, 5C2, 5D2 and 5E2).

The loss in living biomass for conversion from another land use category into CL is estimated as the default value for DM in that particular land use category. I.e. for deforested areas the average carbon stock per hectare for all deforested areas is used.

Change in carbon stock in dead organic matter

When forest land is converted to cropland it is assumed that all dead organic matter will have an instant oxidation. The actual amount depends on which type of forest is converted. Based on the measured Danish data on the nitrogen content in the litter and by using an emission factor of 1.25 % is the N₂O emission from the litter layer 4.5 kg N₂O-N per ha from broadleaves and 9.1 kg N₂O-N from conifers. This emission is reported under Cropland 5(III) for all land use conversions cases despite the land use conversion are to other land uses, see Section 7.11.

Conversion from other categories is assumed as NO as no dead organic matter is reported for these categories.

Change in carbon stock in soils

The actual amount depends on which type of land it is converted from, see Table 7.22. To reach the new equilibrium state is used a default transition period of 50 years. The default IPCC-value of 20 years seems according to Danish investigations not to be applicable for Danish conditions.

N₂O emissions for land converted to Cropland is based on the Tier 2 methodology with the default C stock as given in Table 7.15 and using a C:N value of 15 and an emission factor of 0.0125 kg N₂O-N kg N⁻¹ released.

Uncertainties and time series consistency

The time series are complete.

See uncertainties and time series consistency in Section 7.4.1.

QA/QC and verification

See QA/QC and verification in Section 7.4.

Recalculation

See recalculation in Section 7.4.

Planned improvements

See planned improvements in Section 7.4.

7.5 Grassland (5C)

7.5.1 Grassland remaining grassland (5C1)

Denmark is an intensive agricultural country with many small holders and small fields where CL and GL are mixed together making it difficult to distinguish between dedicated CL and dedicated GL. According to the Danish Land Parcel Information System (LPIS) there are approx. 100 000 fields of total 200 000 ha with permanent GL in 2012 giving an average size of two ha. Some of them cannot be regarded as permanent GL and are therefore included in CL.

Grassland area

The total area with grassland has been estimated in the Land Use matrix. In 1990 the grazing grassland were 217 235 hectares and Other grassland were 183 611 hectares. In 2012 the reported area with grazing grassland were 200 413 hectares and 111 179 hectare with other grassland. The main reason for the decrease in grassland is afforestation and conversion to SE but a better classification of the area which has allowed classifying some of the areas as cropland is also important. The fluctuation is mainly due to difficulties to estimate the land use change between CL and GL. Especially from 2011 to 2012 a large reclassification from GL to CL combined with an increase in the area with grazed grassland from Statistics Denmark is observed.

Grassland definition

Grassland is split into Grazing grassland and Other grassland. Grazing grassland is the area with permanent grassland as recorded by Statistics Denmark. Other Grassland is the difference between the grassland area in the Land Use matrix and the area reported by Statistics Denmark.

Other grassland includes heath land and other areas, e.g. scrub land, which may be grazed by cattle and sheep or land which is kept open for recreational purposes. "Other Grassland" may contain bushes and other wooden plants, which do not meet the thresholds for forest. This is land where the crown cover is below 10 % and where the height at maturity do not reach 5 meter. It includes also nature protection sites, military training sites, electricity network lines etc.

Methodological issues for grassland

The area for grazing grassland is the area reported by statistics Denmark and the rest of the Grassland is the residual part of the grassland area. The area with organic soils in Grassland is estimated from the new organic soil map with an overlay of the fields where the farmers are reporting agricultural crops. Permanent grass fields receiving >63 kg N per ha per year is reported under Grassland. If the farmers are reporting permanent grassland but are using >63 kg N per ha per year it is assumed that this field is grass in rotation because of the fertilization level.

Change in carbon stock in living biomass

No changes in living biomass are assumed for GL remaining GL except for a minor conversion between "Grazing land" and "Other grassland". Because of an increased area reported as grazed grassland in 2012 compared to previous years a quite large emission from GL has been estimated in 2012 compared to previous years.

Change in carbon stock in dead organic matter

No changes in dead organic matter are estimated as this is not occurring for this category.

Change in carbon stock in soils

No changes in the carbon stock in mineral soils are assumed. For organic soils the default IPCC 2003 EF of 1 250 kg C per ha per year is used (there is a likely error in the guidelines as the value is given as 0.25 kg C per ha per year).

Uncertainties and time series consistency

Table 7.22 Tier 1 uncertainty analysis for Grassland for 2012.

		1990	2012						
		Emission/- sink, Gg CO ₂ eqv.	Emission/- sink, Gg CO ₂ eqv.	Activity data, %	Emission factor, %	Combined uncertainty	Total uncertainty, %	Uncertainty 95 %, Gg CO ₂ eqv.	
5.C.Grassland		183.6	553.5				44.6	246.8	
Living biomass	CO ₂	75.3	463.2	10	50	51.0	51.0	236.2	
Dead organic matter	CO ₂	1.6	7.0	10	50	51.0	51.0	3.6	
Mineral soils	CO ₂	0.2	4.5	10	75	75.7	75.7	3.4	
Organic soils	CO ₂	106.6	78.7	10	90	90.6	90.6	71.3	

The time series are complete.

QA/QC and verification

See QA/QC and verification in Section 7.3.1.

Recalculations

The area has been recalculated because of the updated land use matrix.

Planned improvements

None.

7.5.2 Land converted to grassland (5C2)

As agriculture covers more than 63 % of the land area and in order to reduce the environmental impact there is a strategy for turning CL into GL or FL and where deforestation takes place it is often turned into GL or WE.

Approaches used for representing land

The area converted from other land use to GL is based on use of Land Parcel Information data, Natura 2000 vector layers, other vector maps and remote sensing of the Danish area in 1990, 2005, 2011 and 2012. Areas used for gravel digging are normally converted to GL because the normal procedure is removal of the topsoil, and then gravel digging. After having finished the gravel digging the topsoil is reversed to the land and the area turned into marginal grassland/recreational area. To avoid too many land conversions are gravel digging converted directly from CL to GL instead of CL-SE-GL. For an example with an open gravel pit and a restored area please see: [Hedeland resort](#)

Methodological issues

Change in carbon stock in living biomass

For land converted to “grazing land” a standard default gain value of 2 400 kg DM (dry matter) per hectare in above-ground biomass (IPCC 2006, Table 6.4) and 6 720 kg DM per hectare in below-ground biomass (IPCC 2006, Table 6.1) is used. For “Other grassland” not purely free of wooden trees/bushes it is assumed that there is a living biomass of 2 200 kg DM per ha in above ground biomass and 6 160 kg DM per ha in below ground biomass (R:S-factor of 2.8, IPCC 2003 default guideline). For conversion from DM to C a default fraction of fraction of 0.5 kg C per kg DM is used (Table 7.10).

For conversion from GL to other land use categories the same value is used, but recorded as a loss of carbon in the respective category (5A2, 5B2, 5D2 and 5E2).

Change in carbon stock in dead organic matter

When forest land is converted to GL it is assumed that all dead organic matter will be cleared and instant oxidation is taking place.

Conversion from other categories is assumed as NA as no dead organic matter is reported for this category.

Change in carbon stock in soils

The actual amount depends on which type of land it is converted from, see Table 7.15. To reach the new equilibrium state a default transition period of 50 years is used. The default IPCC-value of 20 years seems according to Danish investigations not to be applicable for Danish conditions.

Uncertainties and time series consistency

See Section 7.5.1.

7.6 Wetlands (5D)

Wetland includes:

- unmanaged fully water covered wetlands (lakes and rivers)
- unmanaged partly water covered wetlands (fens and bogs)
- managed water reservoirs (currently not occurring in Denmark)
- managed drained land for peat extraction
- managed partly water covered wetlands (re-established wetlands on primarily former cropland and grassland).

In the beginning of 1990 the total area with wetland has been estimated to 128 867 hectares. By end of 2012 this area has increased to 151 458 hectares. Of this was 58 538 ha lakes and rivers in 1990 increasing to 64 092 ha by end of 2012 inside the > 7000 km long coastline.

7.6.1 Wetlands remaining wetlands – peat extraction (5D1)

The new land use matrix has provided updated figures on the area with partly water covered and fully water covered wetland areas. Partly water covered areas are moors and other areas with raised water table. Fully water covered areas are lakes and rivers.

Wetland area

In the beginning of 1990 the total area with partly covered WE remaining WE has been estimated to 70 509 hectares and the area with lakes to 58 358 hectares. By end of 2012 the area with partly water covered WE remaining WE has decreased slightly to 69 853 hectares. The estimated area with lakes has been reduced with one hectar The total area with peat extraction is about 300 hectares open surface (Lykke Larsen, Pindstrup Mosebrug, personal comm.). Based on aerial photos it is estimated that 1 596 hectares are land connected to the peat extraction areas.

Approaches used for representing land areas

The area for wetlands remaining wetlands is primarily based on data from Danish Geodata Agency and Natura 2000 maps (moors and other natural habitats). The area with peat excavation is a vector map layer made by DCE based on aerial photos of the four excavation sites (Figure 7.17). The actual three locations are Fuglsø mose on Djursland, Lille Vildmose and Store Vildmose – both in Northern Jutland. All locations are nutrient poor raised bogs.




Figure 7.17 Areas with established wetlands, increased water tables and peat extraction in 2008.

Methodological issues for partly water covered wetlands

No changes in the carbon stocks and emissions are reported.

Methodological issues for peat extraction areas

Change in carbon stock in living biomass

No changes in living biomass occurring on the area are reported.

Change in carbon stock in dead organic matter

Dead organic matter is not occurring.

Change in carbon stock in soils

The surface emission from the open peat extraction area is calculated according to Tier 1 (IPCC, 2003) for nutrient poor areas with an emission factor of 0.5 tonnes C per hectare land with peat extraction per year.

The amount of excavated peat (m³ per year) is for each individual extraction site reported to and published by Statistics Denmark (www.dst.dk, Table RST). The total amount of peat excavated has since 1990 been reduced from 399 000 m³ to 152 000 m³ in 2012. This is a >50 % reduction compared to the 10 years ago. For conversion to carbon a density factor of 200 kg per m³ is used (personal comm. with Pindstrup Mosebrug, www.pindstrup.dk who is responsible for the majority of the extraction sites). Furthermore, a DM content of 0.5, an ash content of 0.02 (www.pdir.dk) and a carbon content of 0.58 kg C per kg OM are applied.

For other areas in WE remaining WE is no changes reported.

Nitrous oxide emission

The nitrous oxide emission from peat land extraction areas is estimated from the total N-turnover multiplied with the default IPCC emission factor of 1.25 %. The C:N-ratio in the peat is estimated to 36 in an analysis from the Danish Plant Directorate (www.pdir.dk). Hence the N₂O emission is estimated to 0.546 kg N₂O per tonnes C. Only nitrogen in the degradation of the surface is accounted for in the inventory. N₂O from N in the excavated peat is not estimated.

Uncertainties and time series consistency

Table 7.23 Tier 1 uncertainty analysis for WE remaining WEs and re-established WE for 2012.

		1990	2012					
		Emission/-	Emission/-	Activity	Emission	Combined	Total	Uncertainty 95 %
		sink,	sink,	data, %	factor, %	uncertainty	uncertainty, %	Gg CO ₂ eqv.
		Gg CO ₂ eqv.	Gg CO ₂ eqv.					
5.D Wetlands		87.9	2.4				88.1	2.1
Living biomass	CO ₂	2.6	-0.2	10	50	51.0	51.0	0.1
Dead organic matter	CO ₂	0.5	0.0	10	100	100.5	0	0
Soils	CO ₂	84.7	2.5	10	100	100.5	100.5	2.5
Land for peat extraction	N ₂ O	0.1	0.1	10	100	100.5	100.5	0.1

The time series are complete.

QA/QC and verification

The peat excavation area has been verified with aerial photos and the amount of excavated peat is made by Statistics Denmark.

Recalculation

None.

Category-specific planned improvements

No improvements are planned.

7.6.2 Land converted to wetland (5D2)

In order to restore nature and reduce the environmental impact Denmark has actively re-established WE (Figure 7.17). The size of each restoration project range from less than 1 ha up to 2 500 ha. The benefit of the restoration

programme is more nature but also a reduction in leaching of nitrogen into lakes, rivers and coastal water. The establishment of WE takes place either as large areas turned into lakes or low laying fens.

Since 1990 23 248 ha have been established. These are primarily on CL and GL. Of this is 5 734 hectares converted into new lakes. A major part is restored as a part of the Danish Action Plan for the Aquatic Environment part two (VMP II, running from 1997 to 2006) where land was bought for this purpose but also 2 145 hectares of forest has been converted to partly water covered wetlands. This has primarily taken place in the state owned forest. It is accounted for that the establishment often takes place in connection to existing wetlands.

Water reservoirs for human purposes have not been established for the past 100 years and therefore currently reported as NO.

Approaches used for representing land areas

Geographical vector layers are available for almost all established WE.

Methodological issues

Change in carbon stock in living biomass

For land converted to partly covered wetland a standard default gain value of 4 000 kg DM (dry matter) per hectare in above-ground biomass and 1 200 kg DM per hectare in below-ground biomass is used. For conversion from DM to carbon a default fraction of 0.5 kg C per kg DM is used.

For conversion from wetland to other land use categories the same value but recorded as a loss of carbon in the respective category (5A2, 5B2, 5C2 and 5E2) are used.

Change in carbon stock in dead organic matter

When forest land is converted to wetland it is assumed that all dead organic matter will be cleared with instant oxidation.

Conversion from other categories is assumed as NA as no dead organic matter is reported for these categories.

Change in carbon stock in soils

A default carbon sequestration of 0.5 tonnes C per hectare is assumed for land converted to partly water covered WE.

Nitrous oxide emission

No estimates for the N₂O emission from re-established wetlands have been made.

Methane emission

CH₄ emissions are not estimated due to lack of methodology.

Uncertainties and time series consistency

The time series are complete.

QA/QC and verification

No verification has been made yet.

Recalculation

A recalculation has been made due to new area data on the established WE has been obtained.

Planned improvements

None.

7.7 Settlements (5E)

The annual changes in carbon stock in settlements are assumed to be negligible, and because no estimates have been made, most changes are reported as NA in the CRF Table 5.E. For reporting purposes for land use conversions a default biomass in low buildings, grave yards is established.

The total area with SE has been estimated to 485 543 hectares in 1990 increasing to 516 712 hectares by end of 2012 or to approx. 11 % of the total Danish area.

7.7.1 Settlements remaining settlement (5E1)

Settlement area

The total area with SE has been estimated to 465 779 hectares in 1990 increasing to 491 286 hectares in 2012 or approx. 12 % of the total Danish area. The area is estimated from the cadastral maps and the date where the land parcel was included in the cadastral map, e.g. a change from agriculture to a permanent residence or a road.

Settlement definition

Settlements are defined as all areas with infrastructures, roads, grave yards, sport facilities etc.

Methodological issues

Change in carbon stock in living biomass

No changes in carbon stocks are reported for SE remaining SE.

Change in carbon stock in dead organic matter

No changes in carbon stocks are reported for SE remaining SE.

Change in carbon stock in soils

No changes in carbon stock in soils are assumed.

Uncertainties and time series consistency

Table 7.24 Tier 1 uncertainty analysis for Settlements for 2012.

		1990	2012					Uncertainty 95
		Emission/-	Emission/-	Activity	Emission	Combined	Total	
		sink,	sink,	data, %	factor, %	uncertainty	uncertainty, %	%, Gg CO ₂
		Gg CO ₂ eqv.	Gg CO ₂ eqv.					eqv.
5.E Settlements		12.9	90.8				37.0	33.6
Living biomass	CO ₂	11.4	56.0	10	50	51.0	51.0	28.5
Dead organic matter	CO ₂	0.6	0.0	10	50	51.0	51.0	0.0
Soils	CO ₂	1.0	34.8	10	50	51.0	51.0	17.7

The time series are complete.

QA/QC and verification

No QA/QC has been performed.

Recalculations

A recalculation of the area has been made due to new land use matrix, which slightly has increased the SE area.

Planned improvements

No improvements are planned.

7.7.2 Land converted to settlement (5E2)

Land converted to SE is mostly taking place around the big cities and primarily on cropland and grassland.

Settlement area

The area converted to SE is based on cadastral maps and other digital maps. For simplicity for the years 1990 to 2012 is only used three occasions 1990, 2005 and 2012 with a linear increase in the area in the years between. In future will annual updates take place so that the increase from 2012 to 2013 will be all new houses and roads included in the cadastral map from 31.12.2012 to 31.12.2013

Methodological issues*Change in carbon stock in living biomass*

For land converted to single-family houses a standard default gain value of 2 200 kg DM (dry matter) per hectare in above-ground biomass and 2 200 kg DM per hectare in below-ground biomass is used. For conversion from DM to carbon a default fraction of 0.5 kg carbon per kg DM is used.

For conversion from settlements to other land use categories the same value is used, but recorded as a loss of carbon in the respective category (5A2, 5B2, 5C2 and 5D2).

Change in carbon stock in dead organic matter

When forest land is converted to settlements it is assumed that all dead organic matter will be cleared. Conversion from other categories is assumed as NA as no dead organic matter is reported for these categories.

The N₂O emission is estimated from an instant oxidation of the litter layer.

Change in carbon stock in soils

A default value of 120 tonnes carbon per ha is assumed to be areas Settlements (Table 7.15). For all areas converted from other land use to Settlement is assumed that equilibrium state will be reached after 100 years from the carbon stock in the previous land use category. This is agreed with the UN-FCCCs review team during the review in 2012.

Uncertainties and time series consistency

See uncertainties and time series consistency in Section 7.7.1

The time series are complete.

QA/QC and verification

No QA/QC has been performed.

Category-specific recalculations

The review team in 2011 argued that carbon losses due to deforestation into settlements should be included in the inventory despite there is no IPCC guidelines for this. During the review it was agreed on that an appropriate equilibrium carbon stock in Settlements could be 120 tonnes of carbon per ha (0-100 cm) which is approximately 20-25 % lower than found in FL, CL and GL.

Planned improvements

No improvements are planned.

7.8 Other Land

No permanent snow cover exists in Denmark and only a very small insignificant area with rocks and cliffs. OL is restricted to beaches and sand dunes and estimated to 26 433 hectares.

No land use changes from 5A, 5B, 5C, 5D and 5E is reported.

7.9 Direct N₂O emissions from N fertilization of Forest Land and Other land use – 5(I)

Only a very small amount of nitrogen fertilisers are used in the Danish forests and primarily to Christmas trees. All emissions are reported under Agriculture CRF Table 4.Ds1 since there is only one common national statistics for N fertilization in agriculture and forestry.

7.10 Non-CO₂ emissions from drainage of forest soils and wetlands – 5(II)

A large proportion of the Danish forest area may be considered as drained in the sense that the natural hydrology has been modified by establishment of ditches. Large forest areas have been drained in order to enable establishment of Norway spruce in depressions, fens and pond areas. As an example, a major state forest Gribskov in Northern Zealand by 1850 had an estimated wetland area 400 % larger than that of 1988

(<http://www.skovognatur.dk/Ud/Beskrivelser/Hovedstaden/Gribskov/VandetTilbage.htm>). During the recent years, there has been an effort to restore wetland habitat in the state forests and several drained areas have been restored by filling up ditches, and in many areas of the state forests ditches are no longer maintained and will be gradually more and more ineffective over time. This is a direct consequence of the strategic plan for the state forests to convert to more Close to Nature Forest Management with a specific aim to restore natural hydrology in as many places as possible.

Very few data exist for N₂O emissions in Danish forests. A national project and EU projects have provided data for hydrological gradients in mini-catchments in an old-growth forest and an afforestation area (Christiansen et al., in prep.) and data for one intensively studied beech forest plot (Skiba et al., 2009). For general application at the national level, tier 1 methods will be applied based on default emission factors (IPCC GPG). Emission factors will be compared with the few examples of emission factors data from national projects.

7.10.1 Methodological issues

Equation 3a.2.1 of IPCC GPG was used for estimation of direct N₂O emissions from drained forest soils (tier 1).

Default emission factors (IPCC GPG, Table 3a.2.1) were used for calculation of N₂O emissions. Danish organic soils were considered to be “Nutrient Rich” based on the general presence of fens (minerotrophic peat) and the relatively high N deposition to Danish forests. Rewetted forest soils were assumed to have an N₂O emission corresponding to the natural level and emissions were therefore by default set to zero in accordance with IPCC GPG.

7.10.2 Areas of drained forest soils

Based on expert judgment, the area of drained forest soils were 65 % of mineral forest soils and 75 % of organic forest soils in 1990. It is further judged that the amount of drained forest soils have decreased in the period until 2008 resulting in an area of drained forest soils with 55 % of mineral forest soils and 50 % of organic forest soils. Organic soils constituted 5 % of the forest area based on information on presence of peat from the NFI. A more detailed analysis of forest soils including a mapping is under preparation. A detailed analysis of the re-measurements of the NFI since 2002 will give some indications of the changes in that period, but no data exists prior to 2002. The combined analysis will be included in the next reporting.

7.10.3 Emissions of N₂O from drained forest soils

Estimates of N₂O emissions (Gg N₂O per year) from drained forest soils are based on the IPCC 2003 values. This means that for mineral soils is 0.06 kg N₂O-N per ha per year and for organic soils 0.6 kg N₂O-N per ha per year.

Emission factors are generally in reasonable accordance with those obtained in national projects. In mini-catchments Christiansen et al. (in prep.) found average annual emissions of 0.56±1.1 kg N₂O-N per ha per year for an afforested stand (30 years) and of 0.78±4.2 kg N₂O-N per ha per year for an old-growth forest. Both sites included hydrological gradients from wet/moist to well-drained conditions. For a well-drained Danish beech forest site, Skiba et al. (2009) reported average annual emissions of 0.45±0.48 kg N₂O-N per ha per year.

7.11 N₂O emissions from disturbance associated with land-use conversion to cropland – 5(III)

The main land-use conversion involving deforestation is the conversion from forest to cropland and grassland and a minor deforestation to SE. A major part of the deforestation in 2012 is conversion of christmas trees plantations to cropland and grassland.

This land-use change is expected to be a source for N₂O emissions due to the decomposition of forest floors and corresponding increased mineralization of N. It is assumed that forest floors are completely decomposed during the conversion. Emissions of N₂O are based on default emission factors (IPCC, 2003).

7.11.1 Methodological issues

For all deforested areas it is assumed that the forest floor disappears regardless if the land use conversion is into CL, GL, WE or SE. This is in con-

tradition to the guidelines and CRF Table 5(III), which is only related to disturbance associated with land-use conversion to CL.

The average nitrogen content of forest floors based on the repeated soil inventory was used to estimate the N mineralized for conifers and broadleaves, respectively. A proportion of 1.25 % of the N stock mineralized is assumed to be emitted as N₂O-N.

7.11.2 Emissions of N₂O from deforestation and land-use conversion

The average N content of broadleaf and conifer forest floors for Danish forest plots are given in Table 7.25 together with the estimated N fraction emitted as N₂O. According to IPCC (2003), a default fraction of 1.25 % is assumed emitted as N₂O-N during mineralization of the total N content following conversion.

Table 7.25 Total N content of forest floors in Denmark from the systematic grid "Agricultural Network". The total N content is used for estimation of the amount of N (1.25%) emitted as N₂O during mineralization of the total forest floor N content following land-use change from forest to grassland.

Tree species	Number of plots	Mean N content (kg ha ⁻¹)	Standard dev. (kg ha ⁻¹)	Min N content (kg ha ⁻¹)	Max N content (kg ha ⁻¹)	N ₂ O-N, (kg ha ⁻¹)
Broadleaves	48	359	310	42	1472	4.5
Conifers	60	728	637	20	3447	9.1

In 1990, emissions of N₂O from deforestation were estimated at 0.0098 Gg N₂O for mineral soils and 0.0005 Gg N₂O for organic soils. In 2012 the figures were 0.0019 and 0.0001 Gg N₂O for mineral and organic soils, respectively.

For land use conversion from GL and WE to CL is used the default methodology from IPCC (IPCC 2003) (Eq. 3.3.13 – 3.3.15). The used average carbon stocks are given in Table 7.15. The default methodology assume that an N₂O emission only occur if there is a decrease in the carbon stock the methodology will only estimate a N₂O emission if the land converted from has a higher carbon stock than the land converted to. As the carbon stock in Danish GL soil has been estimated to a lower value than in cropland soils the default methodology will not estimate a N₂O emission for these occasions but estimate a N₂O emission when converted from CL to GL.

7.12 CO₂ emissions from agricultural lime application – 5(IV)

Liming of agricultural soils has taken place for many years. Only a very little amount of lime is applied in forests (<0.5 %) and on permanent grassland. Therefore all liming is included in the inventory under cropland (CRF Table 5(IV)).

The Danish Agricultural Advisory Centre (DAAC) has published the lime consumption for agricultural purposes annually since 1960 (Table 7.26). DAAC are collecting data from all producers and importers. By legislation all producers and importers are obligated to have their products analysed for acid neutralisation content. The analysis is carried out by the Danish AgriFish Agency and published annually (PDIR 2004). The published data from DAAC are corrected for acid neutralisation contents for each product and thus given in pure CaCO₃. For that reason there is no need to correct for inert materials and differ between lime and dolomite as made in the guide-

lines, as this has already been included in the background data. The data from DAAC includes all different products used in agriculture, including e.g. CaCO_3 from the sugar refineries.

The amount of lime used in private gardens has been estimated from the main supplier to private gardens. According to the company (Kongerslev Havekalk A/S, pers. comm.) they are responsible for 80 % of the sale to private gardens. Their sales figures have been used to estimate the total consumption in private gardens. Furthermore, the figures are corrected for acid neutralisation capacity according to the data from the Danish AgriFish Agency. This gives an approximate amount of 2 300 tonnes CaCO_3 per year in private gardens. This figure has been used for all years.

A very small consumption of CAN (Calcium Ammonium Nitrate) and Urea is taking place in Denmark. The amount of CO_2 included in these two fertilisers is included in the lime consumption. Data is taken from the annual fertiliser statistics published by Statistics Denmark.

The amount of lime used for agricultural purposes has declined with 70 % since 1990. From 2000 to 2012 the consumption has been very stable around 400-500 Gg CaCO_3 . The sold amount in 2012 was 426 Gg CaCO_3 . 500 Gg is expected to be the lowest consumption needed to maintain appropriate pH values in the Danish agricultural soils at the moment. The main reason for the reduced lime consumption is a decreased need for acid neutralisation due to less SO_x deposition in Denmark (which also can be seen in the Norwegian inventory) and a reduced consumption of fertilisers containing ammonium. The inter-annual variation is primarily due to weather conditions (if it is possible to drive in the fields) and the economy in agriculture.

The amount of carbon is calculated according to the guidelines where the carbon content is 12/100 of the CaCO_3 . It is assumed that all carbon disappear as CO_2 the same year as the lime is applied.

Table 7.26 Lime and CAN application to cropland and grassland and in forests, 1990-2012.

Greenhouse gas source and sink categories	1990	1995	2000	2005	2008	2009	2010	2011	2012
Agriculture, Gg CaCO_3	1283	1125	590	497	518	410	345	365	426
Private gardens, Gg CaCO_3	2,3	2,3	2,3	2,3	2,3	2,3	2,3	2,3	2,3
CAN and Urea	131,4	149,9	89,9	4,3	2,9	8,3	9,0	9,0	9,0
Total, Gg CaCO_3	1417	1277	682	504	523	421	356	376	437
Total, Gg C pr yr	623	562	300	221	230	185	157	165	192

Table 7.27 Tier 1 uncertainty analysis for Liming for 2012.

	1990	2012					
	Emission/sink, Gg CO_2 eqv.	Emission/sink, Gg CO_2 eqv.	Activity data, %	Emission factor, %	Combined uncertainty	Total uncertainty, %	Uncertainty 95 %, Gg CO_2 eqv.
5(IV) Liming CO_2	622.9	192.3	5	50	50.2	50.2	96.6

The time series is complete.

The collected data is assumed to be very reliable. It is assumed that the uncertainty is in the range of 5 %. The emission factor may be overestimated due to expected leaching of CO_3^{--} , however no data is available on this issue.

7.13 Biomass burning – 5(V)

Burning of forest is prohibited as well as burning of wooden debris from hedgerows are very seldom. Only controlled burning of heathland of approximately 300 hectares per year is taking place. However in 2012 the area with controlled burning of heathland increased to 709 hectares. Due to the humid climate wildfires in the forest are very seldom and normally 0-10 hectares per year.

Data on wild and controlled fires has been collected by the Danish Nature Agency from the forest departments for the period 1990 to 2012. The emission factors are taken from the IPCC 2006 guidelines. As the burned forest is located on poor sandy soils the default standing carbon stock assumed to be 150 Cubic meter per hectare, which is slightly lower than the average standing carbon stock in the Danish forests. The fraction burned for forest is taken from the guidelines whereas for heat land a factor of 0.33 is used. It is based on expert judgment made by the Danish Nature Agency who is responsible for the controlled burning.

Table 7.27 Burned areas 1990 –2012, ha per year.

	1990	1995	2000	2005	2006	2007	2008	2009	2010	2011	2012
Forestland area burned	150	1	0	0	6	0	0	0	0	0	0
Heathland area burned	47	53	122	638	202	270	282	296	359	377	709
Total burned area	197	54	122	638	208	270	282	296	359	377	709

Table 7.28 Tier 1 uncertainty analysis for Biomass burning for 2012.

		1990	2012	Activity data, %	Emission factor, %	Combined uncertainty	Total uncertainty, %	Uncertainty 95 %, Gg CO ₂ eqv.
		Emission/sink, Gg CO ₂ eqv.	Emission/sink, Gg CO ₂ eqv.					
5(V) Biomass Burning	CH ₄	0.5	0.0	50	30	58.3	58.3	0.014
5(V) Biomass Burning	N ₂ O	0.4	0.0	50	30	58.3	58.3	0.020

7.14 Uncertainty, Tier 2 analysis

A first approach has been done to run a Tier 2 Monte Carlo uncertainty simulation for the LULUCF sector. The results are given in Table 7.29 and Table 7.30.

Table 7.29 shows the estimated median emission for the major sources in the LULUCF sector. The organic agricultural soil shows the highest uncertainty in 2012 and Forest remaining Forest the 2nd highest.

In the overall uncertainty estimates for Denmark is only included Tier 1 uncertainty estimates. The Tier 2 uncertainty estimates shown below are only for the LULUCF sector in 2012.

Table 7.29 Detailed uncertainty tier 2 estimates for the LULUCF sector in 2012, Gg CO₂-eqv.

Category	Emissions, Median	Below 2.5%	Above 97.5%	Plus 95%	Minus 95%
5.	-739.30	-2227.59	2002.49	-371%	201%
5.B.1 Cropland, Organic soils	1973.7	848.6	4643.3	-361%	152%
5.A.1 Forest remaining forest	-4489.2	-5216.1	-3867.5	-84%	98%
5.B.1 Cropland, Mineral soils	578.8	283.8	1186.7	-82%	40%
5.C.1 Grassland, Living biomass	346.8	211.0	563.2	-29%	18%
5.B.1 Cropland, Living biomass	271.7	167.8	443.8	-23%	14%
5(IV) Cropland Limestone	192.3	119.4	314.1	-16%	10%
5.C.2 Land converted to grassland	127.4	77.8	207.4	-11%	7%
5.C.1 Grassland, Organic soils	78.8	38.5	160.8	-11%	5%
5.D.1 Wetlands, Soils	34.4	13.6	88.9	-7%	3%
5.B.2 Land converted to cropland	-70.8	-115.4	-42.6	-4%	6%
5.D.2 Land converted to Wetlands	-32.6	-81.5	-12.8	-3%	7%
5.E.2 Settlements, Living biomass	55.8	34.2	91.6	-5%	3%
5.E.2 Settlements, Soils	34.8	21.6	56.5	-3%	2%
5.A.2 Land converted to forest	37.9	31.9	44.9	-1%	1%
5(II) Forest Land.	12.0	8.9	16.3	-1%	0%
5(III) Disturbance, Land converted to cropland	1.0	0.5	2.1	0%	0%
5(II) Wetlands	0.1	0.1	0.3	0%	0%
5.E.2 Settlements, Dead organic matter	0.0	0.0	0.1	0%	0%
5(V) Biomass Burning	0.1	0.0	0.1	0%	0%

The estimated trend uncertainty is shown in Table 7.30.

Table 7.30 Danish uncertainty estimates for the LULUCF sector, tier 2 approach, 2012.

	Uncertainty		Trend	Uncertainty of trend,	
	of total emission, %		1990-2012, %	%age points	
CO ₂	201.3%	-370.9%	114.2%	28.5%	-20.8%

In the next submission an update will take place of the uncertainty factors and the Tier 2 uncertainty estimates in the LULUCF sector will be included in the overall Danish inventory.

7.15 References

Breuning-Madsen, H. & Olsson, M. 1995: Jordbundsundersøgelser i EU's Kvadratnet for monitoring af skovsundhed, Danmark. Rapport, Geografisk Institut, Københavns Universitet, 15 s + 8 bilag.

Grønlund, A., Hauge, A., Hovde, A. & Rasse, D.P. 2008: Carbon loss estimates from cultivated peat soils in Norway: a comparison of three methods, *Nutr. Cy. Agroecos.*, 81, 157-167.

Gundersen, P. 1998: Effects of enhanced nitrogen deposition in a spruce forest at Klosterhede, Denmark, examined by moderate NH₄NO₃ addition. *For. Ecol. Manage.* 101, 251-268.

Johannsen, V.K. 2002: Selection of diameter-height curves for even-aged oak stands in Denmark. Dynamic growth models for Danish forest tree species, Working paper 16, Danish Forest and Landscape Research Institute, Hørsholm, Denmark. 70 p.

Larsen, P.H. & Johannsen, V.K. (eds.) 2002: Skove og plantager 2000. Danmarks Statistik, Skov & Landskab og Skov- og Naturstyrelsen. 171 p. ISBN: 87-501-1287-2.

Lohila, A., Aurela, M., Tuovinen, J.-P. & Laurila, T. 2004: Annual CO₂ exchange of a peat field growing spring barley or perennial forage, J. Geophys. Res., 109, D18116, doi:10.1029/2004JD004715.

Madsen, S.F. 1987: Vedmassefunktioner for nogle vigtige danske skovtræarter. Det Forstlige Forsøgsvæsen 40, 47-242.

Madsen, S.F. 1987: Vedmassefunktioner for nogle vigtige danske skovtræarter. Det Forstlige Forsøgsvæsen 40, 47-242.

Madsen, S.F. & Heusèrr, M. 1993: Volume and stem taper functions for Norway spruce. Forest and Landscape Research 1, 51-78.

Maljanen, M., Martikainen, P.J., Walden, J. & Silvola, J. 2001: CO₂ exchange in an organic field growing barley or grass in eastern Finland, Global Change Biol., 7, 679-692.

Maljanen, M., Komulainen, V.-M., Hytönen, J., Martikainen, P.J. & Laine, J. 2004: Carbon dioxide, nitrous oxide and methane dynamics in boreal organic agricultural soils with different soil management, Soil Biol. Biochem., 36, 1801-1808.

Maljanen, M., Sigurdsson, B.D., Guðmundsson, J., Óskarsson, H., Huttunen, J.T. & Martikainen, P.J. 2010: Greenhouse gas balances of managed peatlands in the Nordic countries – present knowledge and gaps. Biogeosciences 7, 2711-2738.

Moltesen, P. 1985: Skovtræernes ved og anvendelse. Skovteknisk Institut, Frederiksberg. 132 p.

Nord-Larsen, T., Johannsen, V.K., Bastrup-Birk, A & Jørgensen, B.B. (eds.) 2008: Skove og plantager 2006. Skov og Landskab and Skov- og Naturstyrelsen, Hørsholm. 185 p. ISBN: 978-87-7903-368-9.

Petersen, B.M. 2010: Available at: www.agrsci.dk/c-tool

Prins, E. 2009: Prins Engineering: Available at: <http://www.prinsengineering.com>

Skovsgaard, JP, Bald, C & Nord-Larsen, T. 2011: Functions for biomass and basic density of stem, crown and root system of Norway spruce (*Picea abies* (L.) Karst.) in Denmark, Scandinavian Journal of Forest Research, vol. 26 (Suppl 11), pp. 3-20.

Skovsgaard, J.P. & Nord-Larsen, T. 2011: Biomass, basic density and biomass expansion factor functions for European beech (*Fagus sylvatica* L.) in Denmark. European Journal of Forest Research. DOI: 10.1007/s10342-011-0575-4, 1-19.

Sloboda, B., Gaffrey, D. & Matsumura, N. 1993: Regionale und lokale Systeme von Höhenkurven für gleichaltrige Waldbestände. Allg. Forst- u. J. Ztg. 164, 225-228.

Vejre, H., Callesen, I., Vesterdal, L. & Raulund-Rasmussen, K. 2003: Carbon and Nitrogen in Danish forest soils – Contents and distribution determined by soil order. Soil Science Society of America Journal 67: 335-343.

Vesterdal, L. & Raulund-Rasmussen, K. 1998: Forest floor chemistry under seven tree species along a soil fertility gradient. Canadian Journal of Forest Research 28: 1636-1647.

Vesterdal, L., Dalsgaard, M., Felby, C., Raulund-Rasmussen, K. & Jørgensen, B.B. 1995: Effects of thinning and soil properties on accumulation of carbon, nitrogen and phosphorus in the forest floor of Norway spruce stands. Forest Ecology and Management 77: 1-10.

Vesterdal, L., Ritter, E. & Gundersen, P. 2002a: Change in soil organic carbon following afforestation of former arable land. Forest Ecology and Management 169: 141-151.

Vesterdal, L., Jørgensen, F.V., Callesen, I., Raulund-Rasmussen, K. 2002b: Skovjordes kulstoflager - sammenligning med agerjorde og indflydelse af intensiveret biomasseudnyttelse. In: Christensen, B.T. (ed.), Biomasse til energiformål - konsekvenser for jordens kulstofbalance i land- og skovbrug. DJF rapport Markbrug nr. 72.

Vesterdal L., Rosenqvist, L., van der Salm, C., Hansen, K., Groenenberg B.-J. & Johansson, M.-B. 2007: Carbon sequestration in soil and biomass following afforestation: experiences from oak and Norway spruce chronosequences in Denmark, Sweden and the Netherlands. In: Heil G., B. Muys and K. Hansen. Environmental Effects of Afforestation in North-Western Europe - From Field Observations to Decision Support. Springer, Plant and Vegetation 1: 19-52.

Vesterdal, L., Schmidt, I.K., Callesen, I., Nilsson, L.O. & Gundersen, P. 2008: Carbon and nitrogen in forest floor and mineral soil under six common European tree species. For. Ecol. Manage. 255: 35-48.

8 Waste (CRF sector 6)

8.1 Overview of the sector

The waste sector consists of the CRF source categories: 6.A Solid Waste Disposal on Land, 6.B. Wastewater Handling, 6.C. Waste Incineration and 6.D. Waste Other. The data presented in Chapter 8 relate to Denmark only, whereas information for Greenland is included in Chapter 16 and for the Faroe Islands in Annex 8.

For the CRF category 6.A. Solid Waste Disposal on Land, the CH₄ emissions reported in this chapter are a result of calculations in continuation of previously used and reported methodology. Changes in the time trend for this year's submission are due to re-allocation and verification of reported waste types according to the harmonised European waste codes (Statutory Order no. 1309, 18/12/2012). The new harmonised European waste code data collection system and the old ISAG system result in 18 new waste types and associated model input parameters as described in Chapter 8.2 and Thomsen & Hjelgaard (2014).

For the CRF category 6.B. Wastewater Handling, the emissions reported in this chapter are a result of calculations in continuation of previously used and reported methodology. Changes in the time trend for this year's submission are below 1% and due to updated activity data on the biological oxygen demand and a change in the methane correction factor, MCF, from 1 to 0.8 according to the 2006 IPCC guidelines (IPCC, 2009) and in agreement with available plant level data on methane production, flaring and venting as presented in Chapter 8.5.5. It has not been possible to obtain completeness in plant specific data regarding the methane production, flaring and venting. Status of such review are presented in Chapter 8.3 and in Thomsen & Hoffmann, 2014 (see Chapter 8.9).

For the CRF source category 6.C. Waste Incineration, the main emissions are included in the energy sector since all incineration of municipal, industrial, medical and hazardous waste in Denmark is done with energy recovery. The Waste Incineration category includes CH₄ and N₂O emissions from the minor sources of cremation of corpses and carcasses.

The source sector 6.D. Waste Other covers emissions from combustion of biogas in biogas production plants (mentioned as Gasification of biogas in the CRF tables) for the years 1994-2005 where these emissions existed. This activity is not occurring in 2006 - 2012. The Waste Other category also includes CO₂, CH₄ and N₂O emissions from the sources: accidental building fires, accidental vehicle fires and compost production.

Chapter 8.7 and 8.8 presents improved QA/QC procedures and recalculations reflecting the recommended improvements of the 2011 centralised review.

In Table 8.1.1, an overview of all emissions from the waste sector is presented. The emissions are taken from the CRF tables and are presented as rounded figures. The full time series is presented in Annex 3G, Table 3G-1.1.

Table 8.1.1 Emissions for the waste sector, Gg CO₂ equivalents.

		1990	1995	2000	2005	2010	2011	2012
6 A. Solid Waste Disposal on Land	CH ₄	1366	1209	1047	862	720	731	698
6 B. Wastewater Handling	CH ₄	65	68	73	72	74	75	74
6 B. Wastewater Handling	N ₂ O	105	108	90	84	76	79	73
6 C. Waste Incineration	CH ₄	0.01	0.01	0.01	0.01	0.02	0.01	0.01
6 C. Waste Incineration	N ₂ O	0.20	0.21	0.22	0.24	0.29	0.27	0.27
6 D. Waste Other	CO ₂	18	20	18	18	18	18	16
6 D. Waste Other	CH ₄	31	41	70	73	88	90	92
6 D. Waste Other	N ₂ O	13	23	160	62	110	118	127
6. Waste	Total	1598	1469	1459	1173	1087	1113	1080

6.A. *Solid Waste Disposal on Land* is the dominant source in the waste sector with contributions in the time series varying from 85 % (1990) to 65 % (2012) of the total emission, given in CO₂ equivalents. Throughout the time series, the emissions are decreasing due to a reduction in the amount of waste deposited. Comparing 2012 with 1990, the emissions from Solid Waste Disposal Sites have decreased with 49%.

6.B. *Wastewater Handling*. For this source, N₂O contributes the most to the sectorial total, varying between contributions of 5 % (2003) and 8 % (2008). In 2012 the contribution is 7 %. CH₄ from this source contributes with between 4 % (1990) and 7 % (2012) of the sectorial total. The CH₄ emissions increase steadily over the time series. Comparing 2012 with 1990, the emissions from Wastewater Handling have decreased with 14 %.

6.C. *Waste Incineration*. This source contributes with CH₄ and N₂O emissions from human and animal cremations. The contribution to CO₂ equivalent emissions from the sum of CH₄ and N₂O is for the time series 1990-2012 between 0.01 % (1991) and 0.03 % (2010). The trend for the total emissions 1990 - 2012 from this source is increasing; compared to 1990 the 2012 and 2010 emissions have increased with 38 % and 49 % respectively. This increase is almost entirely caused by the increase in animal cremation as this activity has risen with 825 % from 1990 to 2012.

6.D. *Waste Other*. This source contributes with CO₂, CH₄ and N₂O emissions from accidental fires and compost production. The contribution to the total emissions from the waste sector varies from 4 % (1990) to 22 % (2002). Throughout the time series, emissions from Waste Other are increasing; from 1990 to 2012 this category increases with 285 %. The great increase in emissions from 6.D is almost entirely caused by the increasing use of composting at municipal treatment sites.

As a result for the entire waste sector, the sectorial total emission in units of CO₂ equivalents (provided in Table 8.1.1) is decreasing throughout the time series; the emission in 2012 has decreased with 32 % compared to 1990.

Table 8.1.2 specifies the origin and type of the methods and emission factors applied in the Danish inventory.

Table 8.1.2 Reported emissions, calculated methods and type of emissions factors for the subcategory waste handling in the Danish inventory. (CS=country specific, D=default, OTH=other).

CR F	Source	Emissions reported	Method	Emission factor
6 A.	Solid Waste Disposal on land	CH ₄	Tier 2,CS	CS,D
6 B.	Wastewater Handling	CH ₄	CS	CS
		N ₂ O	CS	CS
6 C.	Waste Incineration	CH ₄	Tier 1	Tier 1/CS
		N ₂ O	Tier 1	Tier 1/CS
6 D.	Waste Other	CO ₂	Tier 1, CS	CS, OTH
		CH ₄	Tier 1, CS	CS, OTH
		N ₂ O	Tier 1, CS	CS, OTH

8.1.1 Key category identification

In the key category analysis (KCA) the waste emissions are divided into fourteen categories. In the Tier 1 and Tier 2 KCA, three of the fourteen source categories are identified as key categories in 2012 (Table 8.1.3). The Tier 1 key source identification is based on ranking of absolute quantitative emission, while the Tier 2 KCA takes into account the uncertainties in the calculated emissions, cf. Chapter 1.5).

Off the fourteen categories, Solid Waste Disposal on Land and Composting are the only categories identified as key sources for level. According to the level assessment for both Tier 1 and Tier 2 KCAs Solid Waste Disposal on Land is a key source for level for both year 1990 and 2012. Solid Waste Disposal on Land and Composting are also key category contributions to the trend of the national total of greenhouse gases, calculated in CO₂ equivalents, from 1990 to 2012 for both Tier 1 and Tier 2 KCAs.

Identified key categories within the waste sector are presented in Table 8.1.3. For further information on the KCA level and trend assessments please refer to Chapter 1.5 and Annex 1.

Table 8.1.3 Key category identification Tier1 and Tier 2 from the waste sector 1990 and 2012.

			Tier 1			Tier 2		
			1990	2012	1990-2012	1990	2012	1990-2012
6 A.	Solid Waste Disposal on Land	CH ₄	Level	Level	Trend	Level	Level	Trend
6 B.	Wastewater Handling, direct	CH ₄	-	-	-	-	-	-
		N ₂ O	-	-	-	-	-	-
6 B.	Wastewater Handling, indirect	N ₂ O	-	-	-	-	-	-
6 C.	Incineration of corpses	CH ₄	-	-	-	-	-	-
		N ₂ O	-	-	-	-	-	-
6 C.	Incineration of carcasses	CH ₄	-	-	-	-	-	-
		N ₂ O	-	-	-	-	-	-
6 D.	Accidental fires, buildings	CO ₂	-	-	-	-	-	-
		CH ₄	-	-	-	-	-	-
6 D.	Accidental fires, vehicles	CO ₂	-	-	-	-	-	-
		CH ₄	-	-	-	-	-	-
6 D.	Compost production	CH ₄	-	-	-	-	Level	Trend
		N ₂ O	-	-	Trend	-	Level	Trend

As may be observed from Table 8.1.3 category 6B. and 6C. are divided into two sub-categories, whereas category 6D is divided into three sub-categories. Sub-categories are defined according to inventory reporting and emission models used and the outcome of the KCA influenced by the level of disaggregation into sub-categories.

8.2 Solid waste disposal on land (CRF source category 6A)

For many years, only managed waste disposal sites have existed in Denmark. Unmanaged and illegal disposal of waste is considered to play a negligible role in the context of this category. The amount of deposited waste has decreased markedly throughout the time series.

In 2010, the Danish EPA implemented the new Waste Data System to collect waste statistics. The design of the Waste Data System is considerably different from the ISAG Waste Information System it succeeds. Unlike the previous ISAG system, all waste operators, and not only the plants receiving waste, must now report to the Waste Data System. The fact that waste operators must report to the system makes it possible to collect more accurate data about what industry from which the waste originates. However, the 2010 and 2011 datasets are characterised by the fact that waste operators and waste reception facilities have had to get used to reporting to the system, and even though the Danish EPA has conducted quality assurance of the figures, there is still some uncertainty in the data (The Danish Government, 2013).

The general development for solid waste at disposal sites is a result of action plans by the Danish government called the "Action plan for Waste and Recycling 1993-1997" and "Waste 21 1998-2004" (The Danish Government, 1999). The latter plan had, inter alia, the goal to recycle 64 %, incinerate 24 % and deposit 12 % of all waste. The goal for deposited waste was met in 2000. Further, in 1996 a municipal obligation to assign combustible waste to incineration was introduced. In 2003, the Danish Government set up targets for the year 2008 for waste handling in a "Waste Strategy 2005-2008" report (The Danish Government, 2003). According to this strategy, the target for 2008 is a maximum of 9 % of the total waste to be deposited. In the waste statistics report for the year 2004, data shows that this target was met, since 7.7 % of total waste was deposited in 2004 (DEPA, 2006a). Waste Strategy 2009-12, part I (The Danish Government, 2009) was the sixth waste management plan or strategy adopted by the successive governments dating back to 1986. Waste Strategy 2009-12 set up targets for 2012 according to which a maximum of 6 % of the total waste produced is to be deposited (The Danish Government, 2009). In 2009, it appears that this target has already been met as only 5.6 % of all produced waste was deposited. Data on this level of information from the ISAG database/waste statistics (1994-2009) is presented in Annex 3G, Table 3G-2.1.

Waste Strategy 2009-2012, Part II includes goals of continued decrease in the amount of waste being deposited in Denmark and an increase in reuse, recycling and recovery (Danish Ministry of Environment, 2010). This report includes an evaluation of the capacity of Danish solid waste disposal sites divided into waste classes: inert, mineral, mixed and hazardous waste (DEPA, 2010c). The same waste classes are defined in the new Statutory Order for Landfill (Statutory Order no. 719, 24/06/2011), which refers to the Statutory Order for Waste (Statutory Order no. 1309, 18/12/2012) regarding characterisation of the waste according to the European waste code system; the EWC-code list included in Annex 2 of the statutory Order no. 1319. The New Danish Waste Reporting System (www.mst.dk) is based on the EWC-code system, which forms the basis for the estimation of yearly deposited 18 waste types presented in this year's NIR; details are further described in this chapter, in Annex 3G and in Thomsen and Hjelgaard, 2014).

8.2.1 Source category description

From 1994 to 2005, the number of registered solid waste disposal sites (SWDSs) landfill sites in Denmark has decreased from 176 to 134 (DEPA, 2006b, 2013). Of the closed and still active solid waste disposal sites (SWDS) existing today, 81 of the 134 was closed in 2003, leaving 53 still active SWDS reported to the new waste data system in 2012. Methane collections from 26 of these SWDS are reported to be used at energy-producing installations and 29 are included in the Energy statistics (DEPA, 2003a; Interministerial report, 2007; DEA, 2013a and b) (cf. chapter 8.9).

A quantitative overview of the source category are provided in Table 8.2.1 presenting the amounts of landfilled waste, the annual gross emissions of CH₄, the recovered CH₄ in terms of collected biogas at the landfill sites used for energy production, the amount of CH₄ oxidised in the top layers and the resulting net CH₄ emissions. The CH₄ emission estimate has decreased with 49 % from 1990 to 2012.

A full time series (1990-2012) of these data are shown in Annex 3G, Table 3G-2.2. The amount of waste and the resulting CH₄ emission can also be found in the CRF tables submitted

(http://unfccc.int/national_reports/annex_i_ghg_inventories/national_inventories_submissions/items/6598.php).

Table 8.2.1 Annual amounts of deposited waste, generated methane, recovered methane collected for biogas production, oxidised methane in the top layer and resulting net emission for the Danish SWDS.

Year	Landfilled waste	Gross methane emission	Recovered methane	Methane oxidised in the top layers	Net methane emission	
	Gg	Gg CH ₄	Gg CH ₄	Gg CH ₄	Gg CH ₄	Gg CO ₂ eq
1990	3190	72.8	0.5	7.2	65.0	1366
1995	1969	71.6	7.6	5.8	57.6	1209
2000	1489	66.7	11.3	5.5	49.9	1047
2005	983	55.6	9.9	4.6	41.1	862
2006	1786	44.5	6.4	3.8	34.3	720
2007	2439	42.7	4.0	3.9	34.8	731
2008	1107	41.2	4.3	3.7	33.2	698
2009	3190	72.8	0.5	7.2	65.0	1366
2010	1969	71.6	7.6	5.8	57.6	1209
2011	1489	66.7	11.3	5.5	49.9	1047
2012	983	55.6	9.9	4.6	41.1	862

The decrease in the emission throughout the time series is not quite as steep as the general decrease in the amount of waste deposited. This is partly due to the lag time involved in the exponential degradation processes generating the CH₄ (cf. eq. 8.2.4) and partly due to a significant decrease in the amount of degradable organic waste deposited at landfills in Denmark (cf. Annex 3G, Table 3G-2.3).

Methodological issues

The estimation of CH₄ emission from Danish SWDSs is based on a First Order Decay (FOD) model equivalent to the IPCC Tier 2 methodology (IPCC 1997, 2000 and 2006). The model calculations are performed using national statistics on landfill waste categories reported in the national waste statistics.

This year's submission is based on allocation of the old ISAG and the new waste data for which amount are reported according to the European waste codes into 18 defined waste types with individual content of degradable organic matter and half-life's as provided in table 8.2.2.

The degradation of a deposited waste type of quantity N is modelled according to first order kinetics. The mathematical formulation of this type of exponential decay is

$$\frac{dN}{dt} = -k \cdot N \quad \text{Eq. 8.2.1}$$

where k is the decay constant. Equation 8.2.1 can be solved for the simple case of a momentarily single deposition at time t (W_i) yielding:

$$N(t) = W_i \cdot e^{-k \cdot t} \quad \text{Eq. 8.2.2}$$

where k relates to the half-life time for the content of degradable organic carbon (DOC) in the bulk waste, as:

$$t_{1/2} = \frac{\ln 2}{k} \Rightarrow k = \frac{\ln 2}{t_{1/2}} \quad \text{Eq. 8.2.3}$$

The content of degradable organic carbon (DOC_i), half-life times ($t_{1/2}$) and the corresponding methane generation constants (k) are presented in Table 8.2.2.

Table 8.2.2 Half-life times ($t_{1/2}$), degradation rates constants (k) and content of degradable organic matter (DOC_i) according to 18 waste type, of which 11 are characterised as inert*.

Waste type ¹	DOC_i , [% ww] ²	$t_{1/2}$, [yr, ww] ³	k , [yr ⁻¹ , ww]
Food	15	4	0.173
Paper and cardboard	40	12	0.058
Wood	43	23	0.0
Plastic*	0		
Textile, fur and leather	24	12	0.058
Biodegradable garden waste	20	7	0.099
Chemicals, inert*	0		
Electric & Hazardous*	0		
Glass*	0		
Metal*	0		
Scrap vehicles*	0		
Demolition	4	23	0.030
Soil & Stone*	0		
Particulate matter and dust*	0		
Sludge, inert*	0		
Sludge, degradable	57	12	0.058
Ash & Slag*	0		
Other not combustible waste*	0		

¹Waste types marked "*" are characterised as being inert, meaning that these fraction do not decompose, i.e. $DOC_i = 0$.

²Default IPCC, 2006, Vol. 5, Chapter 2, Table 2.4.

³Default IPCC, 2006, Vol. 5, Chapter 3, Table 3.4.

⁴For demolition waste, the degradable fraction is assumed to be wood and the half-life for wood is therefore used.

The amount of generated methane decreases exponentially over time according to first order degradation kinetics of the content of degradable organic carbon in the deposited waste.

At a given year (t) the amount of degradable organic carbon ($DDOCm(t)$) which decomposes is a result of accumulated contributions from all former years deposit of waste ($W(x)$), where x is year since depositing. The residue of organic matter, i.e. decomposable DOC, left from waste deposited at land-fill sites x years ago, is calculated using the exponential decomposition rule (Eq. 8.2.4).

$$DDOCm(t) = W_i \cdot DOC_i \cdot DOC_f \cdot MCF + DDOCm(t-1) \cdot e^{-k} \quad \text{Eq. 8.2.4}$$

where the methane conversion factor, MCF, is set to the default value of 1 for managed SWDS corresponding to the situation in Denmark (page 3.14, IPCC 2006). DOC_i is the mass fraction of degradable organic carbon in the deposited waste types (Table 8.2.2), and DOC_f represents the fraction of the degradable organic carbon that decompose as function of e.g. pH, temperature and waste composition at the SDWS. For Denmark the default DOC_f value is set to 0.5 (IPCC 2006, page 3.13).

Eq. 8.2.4 assumes that the deposition of degradable organic carbon takes place momentarily once a year and just after the time t , where t is defined as whole years (integer: $t=1,2,..$), so Eq. 8.2.4 consists of two overall contributions that may be expressed as

$$DDOCm(t) = \text{New deposit} + \text{Remaining part of former years deposit}$$

The total amount of degraded organic matter during year t ($DDOCm_{decomp_T}$) is assumed to be equal to the degradation during year t of the organic matter that was deposited at the beginning of the year ($DDOCm(t-1)$):

$$DDOCm_{decomp_T} = DDOCm(t-1) \cdot (1 - e^{-k}) \quad \text{Eq. 8.2.5}$$

Based on Equation 8.2.4 and 8.2.5 it is possible to calculate the degraded amount of organic matter in a step wise manner based on last year result. The degraded amount of organic matter is assumed to generate the CH_4 as described by

$$CH_4 \text{ generated}_T = DDOCm_{decomp_T} \cdot F \cdot 16/12 \quad \text{Eq. 8.2.6}$$

where F , which is the fraction of methane in the gas from landfills, is set equal to 0.41 (DGC, 2009) and 16/12 is the conversion factor from units of C to CH_4

For deriving the net emissions, the amount of recovered or collected methane as well as the amount of oxidised methane in the SWDS top layers needs to be subtracted from the generated methane:

$$CH_4 \text{ Emissions} = \left(\sum_x CH_4 \text{ generated}_{x,T} - R_T \right) \cdot (1 - OX_T) \quad \text{Eq. 8.2.7}$$

where $CH_4 \text{ Emissions}$ is the methane emitted in year T , in units of Gg. T is the inventory year, x is the waste category or type.

R_T is the amount of recovered CH_4 at the Danish disposal sites which are used for energy production. Energy producing installations at 16 sites are registered. The Danish Energy Agency registers the biogas amounts recovered at disposal sites in energy units (TJ) (DEA, 2012). The amount of gas in energy unit is converted to volume of gas using the net calorific value of 15.19 MJ per Nm^3 (DGC, 2009; Vattenfall, 2010; Verdo, 2011). As for the FOD model, the content of CH_4 in the gas recovered is estimated to 41 % and the density of CH_4 is 0.678 kg per m^3 .

OX_T is the assumed oxidation of CH_4 in the top layer. The amount oxidised is uncertain and varies according to SWDS characteristics and management practices. For the Danish model an oxidation factor (OX) of 0.1 used; i.e. the default value for industrialised countries with well-managed disposal sites (IPCC, 2000 and 2006).

The amount of CH_4 recovered, $R(t)$, is calculated as:

$$R_T = \frac{B \cdot 0.41 \cdot 0.678 \text{ kg/m}^3}{15.19 \text{ MJ/m}^3} \quad \text{Eq. 8.2.8}$$

where B is the collected amount of biogas as reported by the DEA in units of MJ. The CH_4 recovered is reported in Table 8.2.1 and 8.2.9 in units of Gg.

Model results and activity data

The amounts of waste deposited are registered and published in the national ISAG and new waste system (www.mst.dk) databases and have been allocated into 18 waste types as presented in Table 8.2.3 and in Annex 3G-2-3

Table 8.2.3 Waste amounts divided between eighteen waste types of which eleven* have been identified as inert waste fractions (cf. Table 8.2.2), Gg.

Waste types	1990	1995	2000	2005	2010	2011	2012
Food	111.7	52.0	26.5	4.6	1.0	0.7	0.9
Paper and cardboard	180.2	84.1	43.0	7.5	1.9	2.6	1.9
Wood	201.5	260.9	254.8	2.6	70.7	96.6	54.7
Plastic	27.0	14.2	8.8	4.6	5.5	7.8	4.3
Textile, fur and leather	5.0	3.1	2.3	0.8	2.2	3.1	2.4
Biodegradable garden waste	136.0	65.2	35.2	7.0	0.0	22.9	5.2
Chemicals, inert	7.7	4.7	3.6	1.4	1.0	0.6	0.1
Electric & Hazardous*	0.5	0.3	0.7	83.7	0.0	0.1	1.4
Glass*	37.3	18.5	10.6	4.8	5.4	5.3	2.9
Metal*	184.3	127.8	107.4	77.9	134.9	156.0	132.6
Scrap vehicles	104.5	64.5	48.8	48.7	16.0	17.2	1.5
Demolition, inert*	282.8	174.5	132.0	87.1	94.8	176.7	155.4
Soil & Stone*	466.1	308.6	271.3	174.0	1321.0	1774.0	624.1
Particulate matter and dust*	32.1	0.0	0.3	0.1	2.5	5.4	25.0
Sludge, inert	90.7	44.5	25.0	10.7	2.2	9.8	11.1
Sludge, degradable	210.7	135.7	107.1	37.7	18.5	22.8	13.4
Ash & Slag	465.8	145.0	8.5	33.8	48.4	54.2	23.3
Other not combustible waste	645.9	464.8	402.9	395.9	59.9	82.8	46.6
Total inert, [Gg]	2062.0	1193.0	887.9	835.6	1596.8	2113.2	872.9
Total, [Gg]	3189.8	1968.5	1488.8	983.0	1786.0	2438.6	1106.8

Data on the amounts of municipal solid waste deposited at managed solid waste disposal sites are reported by the Danish Environmental Protection Agency (DEPA) in old database ISAG database for the years 1994-2009 and the new waste data system (2010-2012). The ISAG data system provides

landfill data for the years 1994-2009 (DEPA, 1996a, 1998a, 1999a, 2001a, 2001b, 2002a, 2004a, 2004b, 2005a, 2006a, 2006b, 2008, 2010a, 2011a) and the new waste data system provides data for 2011 (DEPA, 2013). Data for 2010 and 2012 have not yet been published by the Danish EPA and was obtained by direct contact in the DEPA (cf. chapter 8.7). The reporting year's 2010 (and to a lesser extent 2011-2012) are associated with increased uncertainty (DEPA, 2013).

For the years 2010-2012 allocations has been performed according to the reported European waste codes (Statutory Order no. 1309, 18/12/2012) in the new waste data system (cf. Annex 3G, Table 3G-2.4).

For the old ISAG database, 1994-2009, have been analysed in depth and specific waste fractions have been allocated according to the 18 defined waste types as provided in Table 8.2.2.

Waste characterization data for the year 1985 and information on the total amount of waste deposited at SWDSs in 1970 reported by the Danish EPA in 1993 (DEPA, 1993) was used in the back calculation of the time series from 1994-1985.

Data for 1971-1984 have been determined by assuming a linear development between 1970 and 1985. 1960-1969 data are assumed constant at the 1970 level.

Waste amounts for the whole time series, i.e. 1960- 2012, categorised, allocated and divided into 18 waste types as described above, are provided in Annex 3G, Table 3G-2.3. Corresponding annual fractional distributions of the total amount of deposited waste according to type, respecting mass conservation, is presented in units of mass fractions in Table 8.2.4 (for the whole time series the reader is referred to Annex 3G, Table 3G-2.5).

Table 8.2.4. Fractional distribution of reported waste, according to the old ISAG and the new waste data system (EWC), allocated according to the 18 waste types.

Waste types	1990	1995	2000	2005	2010	2011	2012
Food	3.50	2.64	1.78	0.46	0.06	0.03	0.09
Paper and cardboard	5.65	4.27	2.89	0.76	0.11	0.11	0.17
Wood	6.32	13.26	17.11	0.27	3.96	3.96	4.94
Plastic*	0.85	0.72	0.59	0.47	0.31	0.32	0.38
Textile, fur and leather	0.16	0.16	0.16	0.08	0.12	0.13	0.22
Biodegradable garden waste	4.26	3.31	2.37	0.72	0.00	0.94	0.47
Chemicals, inert*	0.24	0.24	0.24	0.14	0.05	0.02	0.01
Electric & Hazardous*	0.02	0.02	0.05	8.51	0.00	0.00	0.13
Glass*	1.17	0.94	0.71	0.49	0.30	0.22	0.26
Metal*	5.78	6.49	7.21	7.93	7.55	6.40	11.98
Scrap vehicles*	3.28	3.28	3.28	4.96	0.90	0.71	0.14
Demolition	8.87	8.87	8.87	8.87	5.31	7.25	14.04
- of which are concrete, bricks, tiles and ceramics	0.00	0.00	0.00	0.00	0.28	0.21	0.29
- of which are insulation & asbestos-containing materials	0.00	0.37	0.59	2.24	2.60	3.37	7.17
- of which are other demolition	0.00	0.00	0.00	0.00	2.43	3.67	6.58
Soil & Stone*	14.61	15.68	18.22	17.70	73.96	72.75	56.39
Particulate matter and dust*	1.01	0.00	0.02	0.01	0.14	0.22	2.26
Sludge, inert*	2.84	2.26	1.68	1.09	0.12	0.40	1.00
- of which originates from desanding and screenings at WWTPs*	0.00	0.00	0.00	0.00	0.12	0.22	0.75
- of which originates from physico-chemical treatments of water(*)	0.00	0.00	0.00	0.00	0.00	0.00	0.00
- of which originates from anaerobic treatment of waste*	0.00	0.00	0.00	0.00	0.00	0.04	0.06
- of which are sludge and filter cakes primarily from gas treatment*	0.00	0.00	0.00	0.00	0.00	0.05	0.08
- of which originates from primary filtration & clarification of water*	0.00	0.00	0.00	0.00	0.00	0.09	0.10
Sludge, degradable	6.60	6.90	7.19	3.83	1.04	0.93	1.21
- of which originates from treatment of urban WW	0.00	0.00	0.00	0.00	0.41	0.44	0.65
- of which originates from biological treatment of industrial WW	0.00	2.77	3.35	3.07	0.01	0.02	0.02
- of which originates from other treatment of industrial WW	0.00	0.00	0.00	0.00	0.02	0.02	0.03
- of which originates from physico-chemical treatments of water	0.00	0.07	0.04	0.06	0.00	0.17	0.01
- of which originates from oil/water separators and oil regeneration	0.00	0.00	0.00	0.00	0.00	0.01	0.02
- of which originates from sewage cleaning	0.00	0.00	0.00	0.00	0.23	0.04	0.12
- of which is categorised as manure	0.00	0.00	0.00	0.00	0.02	0.00	0.01
- of which is categorised as organic chemicals	0.00	0.00	0.00	0.03	0.35	0.24	0.34
- of which is categorised as other	0.00	0.87	0.74	0.47	NR	NR	NR
- of which 10-30% is dry matter	0.00	2.80	3.43	0.78	NR	NR	NR
- of which >30% dry is matter	0.00	2.65	1.31	0.52	NR	NR	NR
Ash & Slag*	14.60	7.36	0.57	3.44	2.71	2.22	2.10
- of which is categorised as bottom ash, slag and boiler dust*	0.00	7.36	0.57	3.24	1.93	1.92	1.47
- of which is categorised as which fly ash*	0.00	0.01	0.00	0.20	0.30	0.18	0.45
- of which is categorised as which other*	0.00	0.00	0.00	0.00	0.48	0.12	0.19
Other waste, inert*	20.25	23.61	27.06	40.27	3.35	3.39	4.21
- of which is categorised not combustible municipal waste	0.00	11.44	14.11	30.59	3.35	3.39	4.21
- of which is not combustible garden waste	0.00	0.75	0.23	0.05	0.00	0.00	0.00
- of which is not combustible industrial waste	0.00	8.08	6.32	0.63	0.00	0.00	0.00
- of which is not combustible building & construction waste	0.00	3.35	6.40	8.99	0.00	0.00	0.00

While Table 8.2.4 presents the fractional distribution of 18 identified waste types of known DOC_i values, corresponding methane generation potentials are presented in Table 8.2.5.

Table 8.2.5 Methane generation potential for each of the 18 waste types, Gg CH₄ per Gg waste

Waste types	$L_{o,i}/W_i$
Food	0.041
Paper and cardboard	0.109
Wood	0.118
Plastic*	0
Textile, fur and leather	0.066
Biodegradable garden waste	0.055
Chemicals, inert*	0
Electric & Hazardous*	0
Glass*	0
Metal*	0
Scrap vehicles*	0
Demolition	0.011
Soil & Stone*	0
Particulate matter and dust*	0
Sludge, inert*	0
Sludge, degradable	0.156
Ash & Slag*	0
Other waste, inert*	0

The content of degradable organic matter, DOC_i values, in each waste type is shown separately in Table 8.2.2 and has been kept constant for the whole time series. The methane generation potential per unit waste type i is obtained from the equation:

$$\frac{L_{o,i}}{W_i} = DOC_f \cdot MCF \cdot F \cdot 16/12 \cdot DOC_i$$

$$\Rightarrow \frac{L_{o,i}}{W_i} = 0.27 \cdot DOC_i$$

Eq. 8.2.9

where the yearly decomposable fraction of the organic carbon content, DOC_f , are set equal to 0.5, the methane conversion factor, MCF are set equal to 1 and the volume fraction of CH₄ in generated landfill gas, F , are 0.41 (DGC, 2009). The methane generation potential according to waste types are reported in Table 8.2.5. A detailed description of the reallocation of waste statistics according to the 18 waste types is presented in Thomsen and Hjelgaard, 2014.

The annual amounts of the waste types (Table 8.2.3) and their emission generation potentials per mass unit (Eq. 8.2.9 and Table 8.2.6) are used to calculate the deposited CH₄ generation potential and the actual generated CH₄ emission from the annually amount of deposited waste (Eq. 8.2.6).

Figure 8.2.1 shows the time trend in annual amounts of deposited methane generation potential for each of the deposited waste type per year.




Figure 8.2.1 Annual amounts of deposited methane generation potential per waste type.

As shown from Figure 8.2.1, there is a general tendency for decreasing solid waste deposition in Denmark. Also, significant fluctuations are observed; fluctuations that are even bigger for the inert waste types; i.e. waste fraction with a methane generation potential of zero (cf. the introduction section of chapter 8.2). The same fluctuations may be observed from the amount of deposited waste types, meaning that the amount of deposited waste types influences the yearly deposited methane generation potential more than the variation in degradable organic carbon for the individual waste types, DOC_i values.

However, only a fraction of the methane generation potential is release per year; i.e. a function of the waste type specific degradation rate constants, the content of degradable organic carbon and according to first order degradation kinetics for each waste type (Eq. 8.2.1 to 8.2.6 and Table 8.2.2). These seemingly significant fluctuations in the yearly amounts of deposited waste and methane generation potentials becomes insignificant when looking at the annual methane emission factors as illustrated in Figure 8.2.2.




Figure 8.2.2 Annual gross emission factors for each waste type.

Figure 8.2.2 shows the time trend in the gross methane emission factor calculated as the gross methane emission divided by the remaining amount of degradable organic carbon within each waste type. As may be observed from comparing Figure 8.2.2 with 8.2.1, food waste has the highest gross methane emission factor and one of the lowest yearly methane generation potentials. The highest methane emission factor (Figure 8.2.2) for food waste throughout the time series may be explained by the lowest half-life (high CH₄ release rate) and content of degradable organic carbon for food waste compared to other waste types. Still, the yearly amounts of deposited food waste is low and so is the yearly methane generation potential (Eq. 8.2.9).

The net CH₄ emission (Eq. 8.2.7) is obtained upon subtraction of the recovered CH₄, utilised for energy production by biogas combustion installations at some of the sites, and the amount of oxidised methane in the SWDS top layers from the gross methane emission. The annual total amounts of deposited waste, accumulated degradable organic waste, degraded organic matter and the calculated CH₄ emissions are presented in Table 8.2.6.

Table 8.2.6 Waste deposited, total organic degradable matter, amounts of annual degraded organic matter and resulting CH₄ emissions for 1990-2012.

Year	Total Deposited Waste	Accumulated amount of decomposable DDOCm Eq. 8.2.4	Annual amount of degraded DDOCm, Eq. 8.2.5	Annual deposited CH ₄ potential	Annual Gross CH ₄ emission, Eq. 8.2.6	Recovered methane	Annual net emission before oxidation	Annual net emission after oxidation, Eq. 8.2.7	Implied emissions factor	
									Gg CH ₄ /Gg waste	Gg CH ₄ /Gg DDOCm
	[Gg]			[Gg CH ₄]						
1990	3190	942	123	98.2	72.8	0.5	72.3	65.0	0.020	0.026
1995	1969	900	123	62.0	71.6	7.6	64.0	57.6	0.029	0.023
2000	1489	847	117	64.8	66.7	11.3	55.4	49.9	0.033	0.020
2005	983	710	99	10.3	55.6	9.9	45.6	41.1	0.042	0.020
2006	1002	682	95	8.6	53.0	5.6	47.4	42.7	0.043	0.021
2007	984	658	91	10.2	50.8	5.5	45.2	40.7	0.041	0.021
2008	1072	632	87	13.2	48.7	5.0	43.8	39.4	0.037	0.021
2009	779	606	84	8.9	46.6	4.7	41.9	37.8	0.048	0.021
2010	1786	577	80	6.4	44.5	6.4	38.1	34.3	0.019	0.020
2011	2439	551	77	12.6	42.7	4.0	38.7	34.8	0.014	0.021
2012	1107	524	74	18.6	41.2	4.3	36.9	33.2	0.030	0.020

The total waste amount in the second column of Table 8.2.6 is the sum of the amounts of the 18 different waste types (Table 8.2.3). The total waste amount is reported as the activity data for the Annual Municipal Solid Waste (MSW) at SWDSs in the CRF Table 6.A.

The implied emission factor (IEF) in the CRF tables reflects an aggregated emission factor for the model calculated as the net methane emission divided by respectively the total amount of waste deposited in the current year (tenth column in Table 8.2.6) and the total amount of decomposable degradable organic matter, DDOCm, (eleventh column in Table 8.2.6).

The time trend for the total decomposable DOC and annual degraded organic matter are provided in the third and fourth column in Table 8.2.6 and visualised in Figure 8.2.3.




Figure 8.2.3 Time trend in gross methane emission, collected methane, annual net methane emission, annual DDOCm (decomposable degradable organic matter) and remaining DDOCm.

In total a reduction in the net methane emission from 1990 to 2012 of 49 % is observed. This reduction in the methane emission is accompanied by an increase in the degraded amount of decomposable degradable organic matter, DDOCm of 39 % and a 92 % decrease in the remaining amount of DDOCm from 1990 to 2012. The fluctuation in the net methane emission is explained by the fluctuations in the amount of recovered methane.

8.3 Wastewater handling (CRF source category 6B)

The Danish wastewater treatment system is characterised by few big and advanced wastewater treatment plants (WWTPs) and many smaller WWTPs. From 1993 to 2012 the amount of wastewater treated at the most advanced technological WWTPs in Denmark has increased from 53 % to more than 90 %. Improvements of the decentralised wastewater treatment system as well as the sewer system are on-going in Denmark (DEPA, 2010b). For this part of the population, i.e. scattered houses, sludge from septic tanks are collected once per year or as appropriate by judgement of the local authorities (DEPA, 1999b). Municipal collection and transportation of sludge from septic tanks for treatment at the centralised WWTPs occurs to some extent, the frequency set by the authorities and in general septic tanks are emptied one time each year.

A presentation of methodological approach, emission factors, activity data and recalculations are presented in the following sub-chapters.

8.3.1 Source category description

This source category includes an estimation of the emission of CH₄ and N₂O from wastewater handling; i.e. wastewater collection and treatment. CH₄ is produced during anaerobic conditions and treatment processes, while N₂O may be emitted as a by-product from nitrification and denitrification processes under anaerobic as well as aerobic conditions (e.g. Adouani et al., 2010; Kampschreur et al., 2009).

No distinction between emissions from industrial and municipal WWTPs is made, as Danish industries to a great extent are connected to the municipal

sewer system. Wastewater streams from households and industries are therefore mixed in the sewer system prior to further treatment at centralised WWTPs. The contribution from the industry to the influent wastewater at the centralised WWTPs has increased from zero to around 40% from 1987 to 2010 with the highest influent contribution occurring at the biggest and most advanced technological WWTPs in Denmark (Thomsen & Lyck, 2005; ASEP 2010). Monitoring data on the biological oxygen demand (BOD) for the mixed household and industrial influent are available for all WWTPs with a capacity above 30 PE treating more than 90 % of the Danish wastewater (DEPA, 1989, 1999c, 2001d, 2003c, 2004d, 2009; Niero et al., 2014).

Documentation for a decreased fraction of the population not connected sewer system is still missing, and therefore the fraction of the population not connected to the collective sewer system is kept at 10%.

Regarding diffuse emissions from the sewer system, very little data are available (e.g. Lyngby-Taarbæk Kommune, 2013). It is known that centralized wastewater treatment plants are associated with increased residence times, which increases the risk of the occurrence of bottom sediments and thus biological decomposition of organic matter in the sewage system. The sewer system is, however, hydraulically designed to prevent the accumulation of bottom sediments and under such conditions, temporary anaerobic processes will be dominated by fermentation and sulfate reduction, which means that the possibility of methane formation may be ignored (Danva, 2008; Hvitved-Jacobsen, 2001).

It should be mentioned that no activity data are available for industrial WWTPs. Therefore, the emissions from industries having separate wastewater treatment is unknown and i.e. not included in the Danish inventory for category 6.B. Wastewater handling. Only the indirect N₂O emissions are included as effluent N data are available from DEPA reports (DEPA, 1994, 1996b, 1997, 1998b, 1999b, 2000, 2001c, 2002b, 2003b, 2004c, 2005b, 2005c and ASEP 2007, 2010, 2011, 2012, 2013).

Methane emission

The unspecified fugitive methane emission has this year been specified according to the identified systems and processes contribution to the fugitive methane emission from wastewater handling in Denmark. Fugitive methane releases from the municipal and private WWTPs have been divided into contributions from 1) the sewer system, primary settling tank and biological N and P removal processes, 2) from anaerobic treatment processes in closed systems with biogas generation and combustion for energy production and 3) septic tanks. The individual contribution to the net methane emission is given in Table 8.3.1 and Annex 3G, Table 3G-3.1.

Table 8.3.1 Produced, recovered and emitted CH₄ from wastewater treatment, Gg.

	1990	1995	2000	2005	2006	2007	2008	2009	2010	2011	2012
CH ₄ emitted from sewer system and WWTP	0.17	0.21	0.27	0.25	0.26	0.27	0.22	0.25	0.26	0.27	0.24
CH ₄ emission from anaerobic treatment	0.13	0.16	0.29	0.23	0.23	0.24	0.20	0.23	0.24	0.25	0.23
CH ₄ emitted from septic tanks	2.81	2.86	2.92	2.96	2.97	2.98	3.00	3.02	3.03	3.04	3.06
Net CH ₄ emission	3.12	3.23	3.48	3.45	3.46	3.49	3.41	3.50	3.53	3.56	3.52
CH ₄ gross anaerobic processes	13.30	16.03	29.04	22.94	23.17	24.23	19.69	22.81	23.63	24.63	22.62
CH ₄ recovered	13.17	15.87	28.75	22.71	22.94	23.99	19.49	22.58	23.40	24.38	22.39

The emission from the sewer system and WWTP processes (including anaerobic treatment) has increased by 53 % from 1990 to 2012. Regarding the

fraction of the population not connected to the collective sewer system (CH₄ emitted from septic tanks) an increase of 9 % is observed from 1990 to 2012.

Lastly, the amount of recovered methane for energy production has increased 70 % from 1990 to 2012 (cf. chapter 8.5.5 for further documentation).

Nitrous oxide emission

N₂O formation and releases both during the treatment processes at the WWTPs and also from discharged effluent wastewater are included.

The emission of N₂O from wastewater handling is calculated as the sum of contributions from wastewater treatment processes at the WWTPs and from sewage effluents. The emission from effluent wastewater, i.e. indirect emissions, includes separate industrial discharges, rainwater-conditioned effluents, effluents from scattered houses, from mariculture and fish farming. In Table 8.3.2, emission of N₂O from effluent and the contribution from direct N₂O emissions to the total N₂O emission, i.e. the sum of indirect and direct N₂O emissions, is presented.

Table 8.3.2 shows the total N₂O emission originating from treatment processes at the Danish WWTPs (direct emissions) and effluents to the Danish surface waters (indirect emissions). The full time series 1990-2011 is shown in Annex 3G, Table G-3.2.

Table 8.3.2 N₂O emissions from wastewater, Mg.

	1990	1995	2000	2005	2006	2007	2008	2009	2010	2011	2012
N ₂ O, indirect	265	238	157	111	109	116	103	108	109	106	104
N ₂ O, direct	73	111	134	161	127	154	214	127	136	150	131
N ₂ O, total	339	350	292	272	236	270	317	235	246	256	235

Regarding the time trend, the indirect N₂O emission has decreased 61 % N₂O from 1990 to 2012, while the direct N₂O emission has increased 79 %. In absolute figures the indirect emission is a major contributor and the resulting total N₂O emission has decreased 31 % from 1990 to 2012.

8.3.2 Methodology and data

The methodology developed for this submission for estimating emission of methane and nitrous oxide from wastewater handling follows the IPCC Guidelines (IPCC, 1997) and the IPCC Good Practice Guidance (IPCC, 2000). No methodological changes have occurred compared to 2010 (Nielsen et al., 2011a). For the methane emissions from anaerobic digestion, the methane correction factor was decreased from 1 to 0.8, which is in accordance to the IPCC guidelines 2006, and which have been further justified by plant specific data as described in Thomsen and Hoffmann (2014 and chapter 8.5.5). Besides the correction of MCF, smaller corrections in plant inlet TOW data have occurred corresponding to a change below 2 % throughout the time series (cf. Table 8.8.1).

This section is divided into methodological issues related to the CH₄ and N₂O emission calculations, respectively.

Methane emissions from private and municipal WWTPs

The methane emissions from WWTP are divided into a contribution from the sewer system, primary settling tank and biological N and P removal processes, $CH_{4, \text{sewer+MB}}$ and from anaerobic treatment processes in closed systems with biogas extraction for energy production, $CH_{4,AD}$:

$$CH_{4,WWTP} = CH_{4, \text{sewer+MB}} + CH_{4,AD} \quad \text{Eq. 8.3.1}$$

The fugitive emissions from the sewer system, primary settling tank and biological N and P removal processes, $CH_{\text{sewer+MB}}$, are estimated as:

$$\begin{aligned} CH_{4, \text{sewer+MB}} &= EF_{\text{sewer+MB}} \cdot TOW_{\text{inlet}} \\ \Downarrow \\ CH_{4, \text{sewer+MB}} &= B_o \cdot MCF_{\text{sewer+MB}} \cdot TOW_{\text{inlet}} \end{aligned} \quad \text{Eq. 8.3.2}$$

where

TOW_{inlet} equals the influent organic degradable matter measured as biological oxygen demand (BOD) in the influent wastewater flow,

B_o is the default maximum CH_4 producing capacity, i.e. 0.6 kg CH_4 per kg BOD (IPCC, 1997),

$MCF_{\text{sewer+MB}}$ is the fraction of DOC that is anaerobically converted in sewers and WWTPs. MCF_{WWTP} equals 0.003 based on an expert judgement (personal communication: Professor Jes Vollertsen) of a conservative estimate of the fugitive methane emission from the primary settling tanks and biological treatment processes is well below 0.1 % of influent BOD, while the fugitive emission from the sewer system is judged to be negligible or zero (DANVA, 2008).

The emission factor, $EF_{\text{sewer+MB}}$, for these three processes and systems equals 0.0018 kg CH_4 per kg BOD.

The methane emission from anaerobic digestion is calculated as:

$$\begin{aligned} CH_{4,AD} &= EF_{AD} \cdot TOW_{\text{inlet}} \cdot (1 - MR_{AD}) \\ \Downarrow \\ CH_{4,AD} &= B_o \cdot MCF_{AD} \cdot f_{AD} \cdot TOW_{\text{inlet}} \cdot (1 - MR_{AD}) \end{aligned} \quad \text{Eq. 8.3.3}$$

where

B_o is the default maximum CH_4 producing capacity, i.e. 0.6 kg CH_4 per kg BOD (IPCC, 1996; 2006),

MCF_{AD} is extent to which degradation occurs under anaerobic conditions is set equal to 0.8 (IPCC, 2006),

TOW_{inlet} equals the influent organic degradable matter measured as biological oxygen demand (BOD) in the influent wastewater flow,

f_{AD} is the fraction of sludge treated in anaerobic closed systems.

MR_{AD} is the methane generation and combustion efficiency = 99 %

The methane recovery, MR_{AD} , for the anaerobic wastewater treatment with biogas production has been set to 99 % according to expert knowledge (personal communication, Professor Jes Vollertsen; ASEP, 2010) which is further verified in Chapter 8.5.5.

Methane emissions from septic tanks

For the part of the population not connected to the collective sewer system, simple decentralised wastewater handling is assumed and modelled as septic tanks. Only little knowledge is available about the frequency of collection and no measurements of the methane emissions from septic tanks and the pumping and management of septage, including its transportation to a wastewater treatment facility exist. Methane emission from septic tanks is calculated as:

$$\begin{aligned} CH_{4,st} &= EF_{st} \cdot f_{nc} \cdot P \cdot DOC_{st} \\ \Downarrow \\ CH_{4,st} &= B_o \cdot MCF_{st} \cdot f_{nc} \cdot P \cdot DOC_{st} \end{aligned} \quad \text{Eq. 8.3.4}$$

where

B_o is the default maximum CH_4 producing capacity, i.e. **0.6 kg CH_4 per kg BOD** (IPCC, 1996).

MCF_{st} is the methane conversion factor. It depends on the extent to which BOD settles in the septic tanks. MCF_{st} has been set **equal to 0.5** (IPCC, 2006) assuming that degradation for the settled DOC occurs under 100 % anaerobic conditions.

f_{nc} is the fraction of the population that are not connected to the sewer system, i.e. scattered houses which is set equal to **10 %**.

DOC_{st} is the per capita produced degradable organic matter (DOC) which equals **18.250 kg BOD per 1000 persons per year** (IPCC, 2000).

P is the population number.

Using the default maximum methane producing capacity and a methane conversion factor of 0.5 (IPCC guidelines, 2006, Table 6.3) results in an emission factor, EF_{st} , **equal to 0.03**.

Annual activity data and emission factors used for calculation the net methane emission

Monitoring data on the influent biological oxygen demand (BOD) are available for mixed industrial and household wastewater, which are used for calculating the total organic waste (TOW) in the influent wastewater. From 1990 to 1998, the IPCC default methodology for household wastewater has been applied by accounting and correcting for the industrial influent load (Thomsen & Lyck, 2005). For the years 1999 to 2012 monitoring data from the national monitoring program exists. The time series for activity data on TOW are presented in Table 8.3.3.

Table 8.3.3 presents the total degradable organic waste (TOW) calculated by use of the default IPCC method corrected for contribution from industry to

the influent TOW (1990-1998) (Thomsen and Lyck, 2005) and country-specific data (1999-2012). The full time series is presented in Annex 3G, Table 3G-3.3.

Table 8.3.3 Calculated total degradable organic waste in the influent wastewater (TOW).

	1990	1995	2000	2005	2006	2007	2008	2009	2010	2011	2012
Contribution from industrial inlet BOD	2.5	22.2	42.0	40.5	40.5	40.5	40.5	40.5	40.5	40.5	40.5
Population (1000)	5135	5216	5330	5411	5427	5447	5476	5511	5535	5561	5581
TOW [Gg] corrected IPCC method	96.5	116.3									
TOW [Gg]; country-specific data			148.5	140.9	142.3	148.8	120.9	140.1	144.5	150.9	134.6

*TOW = $(1+I/100) \times (P \times D_{\text{dom}})$, where P is the Population number, $D_{\text{dom}} = 18\,250$ kg BOD per 1000

persons per year and I is the per cent contribution from industry to the influent wastewater TOW content.

A country-specific emission factor for calculating the amount of methane produced during anaerobic treatment processes, the gross methane emission (cf. Table 8.3.4), at the Danish WWTPs has been derived. From this emission factor the fugitive emissions from anaerobic treatment processes has been calculated according to Equation 8.3.3. The emission factor varies according to the national statistics on the fraction of wet weight sludge treated anaerobic as reported in the Danish sludge database and presented in Table 8.3.4.

Table 8.3.4 shows the country-specific emission factor for estimating the methane generated during anaerobic treatment processes. The full time series is presented in Annex 3G, Table 3G-3.4.

Table 8.3.4 Emission factor for estimating the methane generation, kg CH₄ per kg BOD.

	1990	1995	2000	2005	2006	2007	2008	2009	2010	2011	2012
MCF _{AD} * f _{AD} *	0.23	0.23	0.33	0.27	0.27	0.27	0.27	0.27	0.27	0.27	0.28
EF _{AD} = B ₀ * MCF _{AD} * f _{AD} [kg CH ₄ /kg BOD]	0.14	0.14	0.20	0.16	0.16	0.16	0.16	0.16	0.16	0.16	0.17

* (Eq.8.3.3).

The Danish sludge database have incomplete statistics due to lack of facility reporting for the years 2006-2011 why the fraction of sludge treated anaerobic have been set equal to the reported fraction in 2005.

Overall methane emission time trends

The trends in the CH₄ emission from the Danish WWTPs, as summarised in Table 8.3.1, are presented graphically in Figure 8.3.1.




Figure 8.3.1 Time trends for net methane emission, methane emission from sewer systems, from septic tanks and from anaerobic treatment processes.

The net methane emissions has increased from 3.2 Gg in 1990 to 3.52 Gg methane in 2012 corresponding to an increase in net methane emissions from wastewater handling of 13 %.

N₂O emissions from WWTPs

N₂O may be generated by nitrification (aerobic processes) and denitrification (anaerobic processes) during biological treatment. Starting material in the influent may be urea, ammonia and proteins, which are converted to nitrate by nitrification. Denitrification is an anaerobic biological conversion of nitrate into dinitrogen. N₂O is an intermediate of both processes. A Danish investigation indicates that N₂O is formed during aeration steps in the sludge treatment processes as well as during anaerobic treatments; the former contributing most to the N₂O emissions during sludge treatment (Gejlsberg et al., 1999). A review by Kampschreur et al. (2009) documents that around 90% of the emitted N₂O originates from activated sludge processes. Based on this review an average of two highest EF values, i.e. 0.6 % N₂O (Wicht et al., 1995) and 0.035 % (Czepiel et al., 1995), both reported in units of per cent N load in the influent wastewater, was used to derive a national EF for the direct emission of nitrous oxide.

The direct N₂O emission from wastewater treatment processes is calculated according to Equation 8.3.5:

$$E_{N_2O} = EF_{N_2O,direct} \cdot m_{N,inf luent} \cdot \frac{M_{N_2O}}{2 \cdot M_N} \quad \text{Eq. 8.3.5}$$

where

$EF_{N_2O,direct}$ is set equal to a fraction of 0.0032 of the N load in the influent wastewater.

$m_{N,inf luent}$ is the annually reported N load in the Danish Water Quality Parameter Database provided in Table 8.3.5.

M_{N_2O}/M_N is the mass ratio i.e. 44/28 to convert the fraction of discharged N emitted as nitrous oxide from total N.

The country-specific EF value of 0.0032 may be expressed as $EF_{N_2O,direct} = 4.99 \text{ g N}_2\text{O per kg N load in the influent wastewater}$ by reducing eq. 8.3.5 to:

$$E_{N_2O} = EF_{N_2O,direct} \cdot m_{N,inf luent} \quad \text{Eq. 8.3.6}$$

The methodology here adopted for estimating the direct N₂O emission only relies on the influent N load as activity data.

The indirect N₂O emission from WWTPs is calculated according to Equation 8.3.7:

$$E_{N_2O,WWTP,effluent} = D_{N,WWTP} \cdot EF_{N_2O,WWTP,effluent} \cdot \frac{M_{N_2O}}{2 \cdot M_N} \quad \text{Eq. 8.3.7}$$

where

$D_{N,WWTP}$ is the effluent discharged sewage nitrogen load consisting of contributions from municipal wastewater treatment plants, the separate industry,

effluent from mariculture and fish farming, rainwater conditioned effluents and scattered houses not connected to the sewage system (cf. Table 8.3.5).

$EF_{N_2O,WWTP,effluent}$ is the IPCC default emission factor of **0.01 kg N₂O-N per kg sewage-N** produced (IPCC, 1997, p 6.28)

M_{N_2O}/M_{N_2} is the mass ratio i.e. 44/28 to convert the fraction of discharged N emitted as nitrous oxide from total N.

Annual activity data and emission factors for calculating the nitrous oxide emission

Data on the N content in the influent and effluent wastewater flows are provided in Table 8.3.5. The effluent data provided in the table constitute a sum of the N content in effluent wastewater from municipal wastewater treatment plants, the separate industry, effluent from mariculture and fish farming, rainwater conditioned effluents and scattered houses. For the entire time series 1990-2011 cf. Annex 3G, Table 3G-3.5.

Table 8.3.5 Nitrogen content in the influent and effluent wastewater, Mg.

	1990	1995	2000	2005	2006	2007	2008	2009	2010	2011	2012
Influent wastewater to WWTPs*	14679 ***	22340	26952	32288	25401	30899	42808	25519	27357	30049	26316
Effluent wastewater from WWTP**	10268	8938	4653	3831	3634	4358	3575	4025	4025	3916	3849
Effluent wastewater, total**	16884	15152	10005	7038	6935	7381	6557	6878	6960	6770	6597

*Data on the influent wastewater N load from municipal WWTPs are available from the Danish Water Quality Parameter Database held by the Agency for Spatial and Environmental Planning ** Effluent wastewater, total includes separate industrial discharges, rainwater conditioned effluent, scattered houses, mariculture and fish farming and effluents from WWTPs (DEPA. 1994, 1996b, 1997, 1998b, 1999b, 2000, 2001c, 2002b, 2003b, 2004c, 2005b, 2005c and ASEP 2007, 2010, 2011, 2012, 2013).*** The significant lower number in 1990 compared to 1995 is due to step increase in the number of WWTPs above 30 PE after implementation of the first Water Action Plan in 1987 (Thomsen and Lyck, 2005; Annex 3G, Table 3G-3.5).

The reduction of N in the effluent wastewater from municipal WWTP compared to in influent wastewater has increased from a reduction efficiency of 30% in 1990 to a reduction efficiency of around 85% in 2012. The significant reduction in the effluent wastewater content of nitrogen has been a driver for the increasing direct N₂O emission from WWTPs. However, emerging wastewater treatment technologies may cause an increased N capture in the sludge (Kristensen og Jørgensen, 2008).

Overall nitrous oxide emission trends

The trends in the direct N₂O emission from WWTPs, the indirect emission from wastewater effluent and the total, as summarised in Table 8.3.2, are presented graphically in Figure 8.3.2.




Figure 8.3.2 Time trends for direct emission of N₂O, indirect emission, i.e. from wastewater effluents, and total N₂O emission.

The annual fluctuations may be caused by several factors such as e.g. climatic condition such as variations in precipitation and as a result varying contributions to the influent N and varying characteristics of especially the industrial contributions to the influent. Furthermore, infiltration of groundwater, as well as exfiltration of overload rainwater and wastewater (DEPA. 1994, 1996b, 1997, 1998b, 1999b, 2000, 2001c, 2002b, 2003b, 2004c, 2005b, 2005c; ASEP 2007, 2010, 2011; Vollertsen et al., 2002), may contribute to the “noise” or fluctuation in the trend of the calculated indirect N₂O emission.

The direct emission shows an increasing trend increases from 1900 to 2002, where a stable but fluctuating level is reached. Comparing 2012 with the base year 1990 an increase of 79 % is observed.

The decrease in the emission from effluent wastewater is due to the technical upgrade and centralisation of the Danish WWTPs following the adoption of the Action Plan on the Aquatic Environment in 1987. The indirect emission from wastewater effluent has decreased from 265 tonnes N₂O in 1990 to 104 tonnes N₂O in 2012 corresponding to a reduction of 61%.

The indirect emission is the major contributor to the emission of nitrous oxide in the period 1990-1997. However, from around year 2000, the direct N₂O emission is the major contributor to the total N₂O emission. Overall, a net reduction of 31 % is observed for the total N₂O emission from wastewater handling.

8.4 Waste incineration (CRF source category 6C)

The CRF source category 6.C. Waste Incineration, includes cremation of human bodies and cremation of animal carcasses.

Incineration of municipal, industrial, clinical and hazardous waste takes place with energy recovery, therefore the emissions are included in the relevant subsectors under CRF sector 1A. For documentation please refer to Chapter 3.2. Flaring off-shore and in refineries are included under CRF sector 1B2c, for documentation please refer to Chapter 3.5. No flaring in chemical industry occurs in Denmark.

Table 8.4.1 gives an overview of the Danish greenhouse gas emission from the CRF source category 6.C waste incineration.

CO₂ emissions from cremations of human bodies and animal carcasses are biogenic.

While emissions from human cremations have been steady over the last two decades, emissions from animal cremations have increased. In 1990, incineration of animal carcasses stood for 5 % of the total emission of CO₂ eqv. from cremations. In 2012 this number has increased to 32 %. GHG emissions from cremations are very small; 0.21 Gg in 1990 and 0.29 Gg in 2012. GHG emissions are shown in Table 8.4.1, for the full time series, see Annex 3G, Table 3G-4.1.

Table 8.4.1 Overall emission of greenhouse gases from the incineration of human bodies and animal carcasses.

Year		199	199	2000	2005	2010	2011	2012
CH ₄ emission from								
Human cremation	Mg	0.48	0.52	0.49	0.48	0.49	0.49	0.48
Animal cremation	Mg	0.03	0.04	0.08	0.14	0.26	0.22	0.22
Total	Mg	0.51	0.55	0.57	0.62	0.76	0.71	0.70
N ₂ O emission from								
Human cremation	Mg	0.60	0.64	0.61	0.60	0.62	0.61	0.60
Animal cremation	Mg	0.03	0.05	0.10	0.17	0.33	0.28	0.28
Total	Mg	0.64	0.69	0.71	0.77	0.95	0.88	0.88
6C. Waste incineration								
CO ₂ equivalents	Gg	0.21	0.23	0.23	0.25	0.31	0.29	0.29

8.4.1 Human cremation

The incineration of human corpses is a common practice that is performed on an increasing part of the deceased. All Danish crematoria use optimised and controlled cremation facilities, with temperatures reaching 800-850 °C, secondary combustion chambers, controlled combustion air flow and regulations for coffin materials.

Emissions from human cremation are calculated for greenhouse gases CH₄ and N₂O.

Methodological issues

During the 1990es all Danish crematoria were rebuilt to meet new standards. This included installation of secondary combustion chambers and in most cases, replacement of old primary combustion chambers (Schleicher et al., 2001). All Danish crematoria are therefore performing controlled incinerations with a good burn-out of the gases, and a low emission of pollutants.

Following the development of new technology, the emission limit values for crematoria were lowered again in January 2011. These new standards were originally expected from January 2009 but were postponed two years for existing crematoria. Table 8.4.2 shows a comparison of the emission limit values from February 1993 and the new standard limits.

Table 8.4.2 Emission limit values mg per Nm³ at 11 % O₂ (Schleicher et al., 2008).

Component	Report 2/1993	Standard terms (1/2011)
	Emission limit value mg per normal m ³ at 11 % O ₂	
CO ₂	500	500
Other demands:		
Stack height	3 m above rooftop	3 m above rooftop
Temperature in stack	Minimum 150 °C	Minimum 110 °C
Flue gas flow in stack	8 – 20 m/s	No demands
Temperature in after burner	850 °C	800 °C
Residence time in after burner	2 seconds	2 seconds

To meet the new standards, some crematoria have been rebuilt to larger capacity while others are closed (MILIKI, 2006). In 2012, there were 26 operating crematoria in Denmark, some with multiple furnaces. In 2010 there were 31 operating crematoria (DKL, 2013).

Crematoria that are not closed are equipped with flue gas cleaning (bag filters with activated carbon). The use of air pollution control devices. The use of air pollution control devices, will however not affect the greenhouse gas emissions.

Around half of the Danish crematoria are currently connected to the district heating system and in addition, a few crematoria produce heat for use in their own buildings. The bag filter cleaning system requires that the flue gas is cooled down to 125-150 °C, and the cheapest way to do so is to use the surplus heat in the district heating system (DKL, 2009). The heat contribution from crematoria is negligible compared to the total district heat production and is not part of the Danish energy statistics, therefore, it is not included in the Energy sector.

Activity data

Table 8.4.3 shows the time series of total number of nationally deceased persons (Statistics Denmark, 2013), number of cremations and the fraction of cremated corpses in relation to the total number of deceased (DKL, 2013). Annex 3G, Table 3G-4.2 presents data for the entire time series 1990-2012.

Table 8.4.3 Data human cremations, DKL (2013), Statistics Denmark (2013)

Year	1990	1995	2000	2005	2010	2011	2012
Nationally deceased	60926	63127	57998	54962	54368	52516	52325
Cremations	40991	43847	41651	40758	42050	41248	40909
Cremation fraction, %	67.3	69.5	71.8	74.2	77.3	78.6	79.6




Figure 8.4.1 Illustration of the development in cremations (DKL 2013), where the number of cremations, $N_{\text{cremations}}$, is shown at the left Y-axis. The cremation percentage, $F_{\text{cremations}}$, shows the percentage of cremated deceased of the total number of deceased for the years 1990-2012.

Even though the total number of annual cremations is fluctuating, the cremation percentage has been steadily increasing since 1990, and is likely to continue to increase.

The average body weight is assumed to be 65 kg (EEA, 2009).

Figure 8.4.2 presents the trend of the number of deceased persons together with the activity data for human cremation. The figure shows a direct connection between the number of deceased and the activity of human cremation as the two trends are quite similar. Figure 8.4.2 also shows the effect of the increasing fraction of cremations per deceased, as the number of cremations is not decreasing along with the number of deceased. The cremation fraction has increased from 67 % in 1990 to 80 % in 2012; the trend of this fraction is shown in Figure 8.4.1.




Figure 8.4.2 Trends of the activity data for cremation of human corpses and the national number of deceased persons.

Emission factors

For human cremation, emissions are calculated by multiplying the total number of human cremations by the emission factors. Since there are no continuous measurements available of the annual emission from Danish crematoria, the estimation of emissions is based on emission factors from literature.

A literature search has provided the emission factors shown in Table 8.4.4. It has not been possible to find any additional data to validate the emission factors.

Table 8.4.4 Emission factors for human cremation with references.

Pollutant name	Unit	Emission factor	Reference
CH ₄	g/body	11.8	Aasestad, 2008
N ₂ O	g/body	14.7	Aasestad, 2008

8.4.2 Animal cremation

The incineration of animal carcasses in animal crematoria follows much the same procedure as human cremation. Animal crematoria use similar two chambered furnaces and controlled incineration. However, animal carcasses are incinerated in special designed plastic (PE) bags rather than coffins. Emissions from animal cremation are similar to those from human cremation.

Animal cremations are performed in two ways, individually where the owner often pays for receiving the ashes in an urn or collectively which is most often the case with animal carcasses that are left at the veterinarian.

Methodological issues

Open burning of animal carcasses is illegal in Denmark and is not occurring, and small-scale incinerators are not known to be used at Danish farms. Livestock that is diseased or in other ways unfit for consumption is disposed of through rendering plants. Incineration of livestock carcasses is illegal and these carcasses are therefore commonly used in the production of fat and soap at Daka Bio-industries.

The only animal carcasses that are approved for cremation in Denmark are deceased pets and animals used for experimental purposes, where the incineration must take place at a specialised animal crematorium. There are four animal crematoria in Denmark but one of these is situated at a waste incineration company in northern Jutland called AVV. The specially designed cremation furnaces are at this location connected to the flue gas cleaning equipment of the municipal waste incineration plant with energy recovery and the emission from the cremations are therefore included in the annual inventory from AVV and consequently included under the energy sector in this report. Therefore only three animal crematoria are included in this section.

Animal by-products are regulated under the EU commission regulation no. 142/2011. This states that animal crematoria must be approved by the authority and comply either with the EU directive (2000/76/EC) on waste incineration or with Regulation (EC) No. 1069/2009 (EC, 2011).

The incineration of animal carcasses is, as the incineration of human corpses, performed in special incineration chambers. All Danish animal crematoria have primary combustion chambers with temperatures around 850 °C and secondary combustion chambers with temperatures around 1100 °C. The support fuel used at the Danish facilities is natural gas.

Emissions from pet cremations are calculated for CH₄ and N₂O.

Activity data

Activity data for animal cremation are gathered directly from the animal crematoria. There is no national statistics available on the activity from these facilities. The precision of activity data therefore depends on the information provided by the crematoria.

Table 8.4.5 lists the four Danish animal crematoria, their foundation year and provides each crematorium with an id letter.

Table 8.4.5 Animal crematoria I Denmark.

Id	Name of crematorium	Founded in
A	Dansk Dyrekremering ApS	May 2006
B	Ada's Kæledyrskrematorium ApS	Unknown, existed in more than 30 years
C	Kæledyrskrematoriet	2006
D	Kæledyrskrematoriet v. Modtage-station - Vendsyssel I/S	

Crematoria D is situated at the AVV municipal waste incineration site and the emissions from this site are, as previously mentioned, included in the

annual emission reporting from AVV and consequently included in the energy sector in this report as waste incineration with energy recovery. Therefore, only crematoria A-C are considered in this chapter.

Table 8.4.6 lists the activity data for animal crematoria A-C. The entire dataset for 1990-2012 is available in Annex 3G, Table 3G-4.3.

Table 8.4.6 Activity data. Source: direct contact with all Danish crematoria.

	1990	1995	2000	2005	2010	2011	2012
Total, Mg	150	200	443	762	1449	1219	1238

Crematorium B delivered exact annual activity data for the years 1998-2011. They were not certain about the founding year but believe to have existed since the early 1980es. Activity data for 1990-1997 and 2012 has therefor been estimated by the authors expert judgement. It is not possible to extrapolate data back to 1990 because the activity, due to the steep trend line, in this case would become negative.




Figure 8.4.3 The amount of animal carcasses cremated, in Mg. Data from 1998-2012 are delivered by the crematoria and is considered to be exact; these data are marked as points. Data from 1990-1997 are estimated and are shown as the thick line in the figure.

It is not possible to extrapolate data linearly back to 1980 because the activity, due to the steep increase, in this case would become negative from 1993 and back in time.

Emission factors

Concerning the incineration of animal carcasses in animal crematoria there is not much literature to be found.

Emission factors for CH₄ and N₂O are collected from the literature search on human cremation, and it is assumed that humans and animals are similar in composition for this purpose. Emission factors from human cremation are recalculated to match the activity data for animal cremation, emission per Mg.

Table 8.4.7 lists the emission factors and their respective references.

Table 8.4.7 Emission factors for animal cremation.

Pollutant name	Unit	Emission factor	Reference
CH ₄	g/Mg	182	Aasestad, 2008
N ₂ O	g/Mg	226	Aasestad, 2008

8.5 Waste Other (CRF Source Category 6D)

This category is a catch all for the waste sector. Emissions in this category could stem from compost production, accidental fires, sludge spreading, biogas production and other combustion without energy recovery. In the Danish inventory emissions from accidental fires, compost production and biogas production are included in this category.

Table 8.5.1 gives an overview of the Danish greenhouse gas emission from the CRF source category 6.D waste other.

CO₂ emissions from compost production are biogenic. For accidental fires in buildings, there is a high content of wood both in the structure and in the interior; this leads to 82 % of the CO₂ emission from accidental building fires to be biogenic, see Table 8.5.1.

Emissions from accidental fires are quite constant, with a peak in 2008 of 23 Gg CO₂ equivalents. Compost production is the largest source of greenhouse gas emissions in this sector. In 1990 composting stood for 69 % (42 Gg CO₂ equivalents) of the total greenhouse gas emission in CO₂ equivalents from the other waste category, in 2012 this number has increased to 92 % (217 Gg CO₂ equivalents). The full time series is shown in Annex 3G, Table 3G-5.1a-c.

Table 8.5.1 Overall emission of greenhouse gasses from accidental fires and composting.

		1990	1995	2000	2005	2010	2011	2012
CO ₂ emission from								
Accidental building fires	Gg	63.1	72.2	63.8	62.4	61.7	67.6	60.5
- of which non-biogenic	Gg	11.4	13.1	11.5	11.3	11.1	12.2	10.8
Accidental vehicle fires	Gg	6.1	6.5	6.9	6.9	7.3	6.3	5.6
Total, non-biogenic	Gg	17.5	19.6	18.4	18.1	18.3	18.4	16.4
CH ₄ emission from								
Compost production	Mg	1386.1	1860.2	3240.0	3419.9	4094.6	4204.3	4306.3
Accidental building fires	Mg	64.1	73.4	64.9	63.8	64.6	68.5	61.7
Accidental vehicle fires	Mg	12.8	13.6	14.3	14.3	15.1	13.1	11.6
Total	Mg	1463.0	1947.2	3319.2	3497.9	4174.3	4285.9	4379.6
N ₂ O emission from								
Compost production	Mg	41.5	72.8	515.7	200.2	355.2	381.4	409.1
Accidental building fires	Mg	NAV	NAV	NAV	NAV	NAV	NAV	NAV
Accidental vehicle fires	Mg	NAV	NAV	NAV	NAV	NAV	NAV	NAV
Total	Mg	41.5	72.8	515.7	200.2	355.2	381.4	409.1
6D. Waste other								
CO ₂ -equivalents	Gg	61.1	83.0	248.0	153.6	216.1	226.7	235.2

8.5.1 Compost production

This section covers the biological treatment of solid wastes called composting. Greenhouse gasses that are emitted from this process are CH₄, N₂O and CO₂.

Methodological issues

Emissions from composting have been calculated according to a country specific Tier 1 method. However, a Tier 1 default methodological guidance is available in the 2006 IPCC Guidelines (IPCC, 2006).

In Denmark, composting of solid biological waste includes composting of:

- garden and park waste (GPW),
- organic waste from households and other sources,
- sludge,
- home composting of garden and vegetable food waste.

In 2001, 123 composting facilities treated only garden and park waste (type 2 facilities), nine facilities treated organic waste mixed with GPW or other organic waste (type 1 facilities) and 10 facilities treated GPW mixed with sludge and/or “other organic waste” (type 3 facilities). 92 % of these facilities consisted entirely of windrow composting, which is a simple technology composting method with access to only natural air. It is assumed that all facilities can be considered as using windrow composting. (Petersen & Hansen, 2003)

Composting is performed with simple technology in Denmark; this implies that temperature, moisture and aeration are not consistently controlled or regulated. Temperature is measured but not controlled, moisture is regulated by watering the windrows in respect to weather conditions and aeration is assisted by turning the windrows. (Petersen & Hansen, 2003)

During composting a large fraction of the degradable organic carbon (DOC) in the waste material is converted into CO₂. Even though the windrows are occasionally turned to support aeration, anaerobic sections are inevitable and will cause emissions of CH₄. In the same manner, aerobic biological digestion of N leads to emission of N₂O (IPCC, 2006).

Activity data

All Danish waste treatment plants are obligated to statutory registration and reporting of all waste entering and leaving the plants. All waste streams are weighed, categorised with a waste type and a type of treatment and registered to the ISAG waste information system, which contain data for 1995-2009 (ISAG, 2010). The new waste data system that was supposed to replace ISAG in 2010 is not yet functioning; activity data for 2010-2012 has therefore been estimated by extrapolation.

Figure 8.5.1 illustrates the composted amount of waste divided in the four categories mentioned earlier.




Figure 8.5.1 National amount of composted waste, these data are also shown in Table 8.5.3.

Activity data for the years 1995-2009 are collected from the ISAG database for the categories: “sludge”, “organic waste from households and other sources” and “garden and park waste”. Activities for 2010-2012 are calculated by using the trend from earlier years.

The Danish legislation on sludge (DEPA, 2006c) was implemented in the summer of 2003. This stated that composted sludge may only be used as a fertilizer on areas not intended for growing foods of any kind for at least 2-3 years. This restriction caused the amount of composted sludge to drop drastically from 2003 to 2004.

The trend in composting of sludge does not demonstrate a convincing trend that can be used for estimation of activity data for previous years. Since this activity is insignificant for 1995-1997 (1-2 %) it is assumed to be “not occurring” for 1990-1994.

The amount of organic waste from households composted in the years 1990-1994 is estimated by multiplying the number of facilities treating this type of waste with the average amount composted per facility in the years 1995-2001 (2.6-3.8 Gg per facility per year). The following Table 8.5.2 shows the number of composting sites divided in the three types described in “Methodological issues” (Petersen, 2001 and Petersen & Hansen, 2003).

Table 8.5.2 Number of composting facilities in the years 1990-2001.

Facility type	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001
Type 1	5	6	7	8	9	13	14	13	14	13	11	9
Type 2	38	54	70	86	102	113	108	99	102	111	115	123
Type 3	1	2	2	3	4	9	9	11	10	10	7	10
Total	44	62	79	97	115	136	133	126	130	139	138	149

Type 1 waste treatment sites normally includes biogas producing facilities, but these have been excluded in Table 8.5.2.

The ISAG activity data for composting of garden and park waste (GPW) include wood chipping. Compost data for GPW provided by Petersen (2001) and Petersen & Hansen (2003) show that for 1997-2001, wood chipping accounts for about 3 % of the total chosen ISAG activity data for GPW. Activity data for GPW for the years 1985-1994 and 2010-2011 are estimated by extrapolating the trend.

The last waste category involved in composting is home composting of garden waste and vegetable waste. The activity data for this category are known from Petersen & Kielland (2003) to be 21.4 Gg in 2001. It is assumed that the following estimates made by Petersen & Kielland (2003) are valid for all years 1990-2012.

- 28 % of all residential buildings with private gardens (including summer cottages) are actively contributing to home composting.
- 14 % of all multi-dwelling houses are actively contributing to home composting.
- 50 kg waste per year will in average be composted at every contributing residential building.
- 10 kg waste per year will in average be composted at every contributing multi-dwelling house.

Multi-dwelling houses include apartment buildings, it is very un-common for people in these types of buildings to compost their bio waste and the average amount of composted waste is therefore lower in spite of the higher number of residents. The total number of occupied residential buildings, summer cottages and multi-dwelling houses are found at the Statistics Denmark's website.

The calculated activity data for composting are shown in Table 8.5.3 and in Annex 3G, Table 3G-5.2.

Table 8.5.3 Activity data composting, Gg.

	1990	1995	2000	2005	2010	2011	2012
Composting of garden and park waste	288	376	677	737	877	901	924
Composting of organic waste from house_holds and other sources	16	40	47	45	58	59	59
Composting of sludge	NO	7	218	50	120	132	145
Home composting of garden and vegetable food waste	20	21	21	22	23	23	23
Total	324	444	963	854	1078	1115	1151

NO = Not occurring

Emission factors

The emissions from composting strongly depend on both the composition of the treated waste and on process conditions such as aeration, mechanical agitation, moisture control and temperature pattern (Amlinger et al., 2008).

The emission factors stated in Table 8.5.4 are considered the best available for the calculation of Danish emissions from composting.

Table 8.5.4 Composting emission factors.

	Garden and park waste (GPW)	Organic waste from households and other sources	Sludge	Home composting of garden and vegetable food waste
Unit	kg per Mg	kg per Mg	kg per Mg	kg per Mg
CH ₄	4.20	4.00	0.41	5.63
N ₂ O	0.12	0.30	1.92	0.11
Source	Boldrin et al., 2009	EEA, 2009	MST, 2013	Boldrin et al., 2009

Emission factors for composting of GPW and for home composting of garden and vegetable food waste are derived from Boldrin et al. (2009). No other sources were found that describe the emission from home composting.

Boldrin et al. (2009) and MST (2013) do not directly provide any emission factors, the following assumptions were made to derive the factors shown in Table 8.5.4;

- 0.5 % N per dry matter waste water sludge
- 25 % moisture in waste water sludge.
- 2 % N per dry matter garden waste (incl. home composting)
- 25-50 % DOC per dry matter garden waste (incl. home composting)
- 50 % moisture in garden waste (incl. home composting)

The CO₂ produced and emitted during composting is short-cycled C and is therefore regarded as CO₂ neutral. (Boldrin et al., 2009).

8.5.2 Accidental building fires

Emissions from accidental fires are categorised under CRF category 6D Other Waste. Emissions that escape from building fires are CO₂ and CH₄.

Methodological issues

Emissions from building fires are calculated by multiplying the number of building fires with selected emission factors. Six types of buildings are distinguished with different emission factors: detached houses, undetached houses, apartment buildings, industrial buildings, additional buildings and containers.

Activity data

In January 2005 it became mandatory for the local authorities to register every rescue assignment in the *online data registration- and reporting system called ODIN*, *ODIN is developed and run by the Danish Emergency Management Agency (DEMA, 2007)*.

Activity data for accidental building fires are given by ODIN (DEMA, 2013). Fires are classified in four categories: full, large, medium and small. The emission factors comply for full scale fires and the activity data are therefore recalculated as a full scale equivalent where it is assumed that a full, large, medium and a small scale fire leads to 100 %, 75 %, 30 % and 5 % of a full scale fire respectively.

In practice, a full scale fire is defined as a fire where more than three fire hoses were needed for extinguishing the fire, a full scale fire is considered as a complete burnout. A large fire is in this context defined as a fire that involves the use of two or three fire hoses for fire extinguishing and is assumed to typically involve the majority of a house, an apartment, or at least part of an industrial complex. A medium size fire is in this context defined as a fire involving the use of only one fire hose for fire-fighting and will typically involve a part of a single room in an apartment or house. And a small size fire is in this context defined as a fire that was extinguished before the arrival of the fire service, extinguished by small tools or a chimney fire.

The total number of registered fires is known for the years 1990-2012. For the years 2007-2012 the total number of registered building fires is known with a very high degree of detail.

Table 8.5.5 shows the occurrence of all types of fires (registered for 1990-2012) and the occurrence of building fires (2007-2012) registered at DEMA. In 2007-2010 the average per cent of building fires, in relation to all fires, was 60 %. The total numbers of building fires 1990-2006 are calculated using this percentage. The full time series is presented in Annex 3G, Table 3G-5.3.

Table 8.5.5 Occurrence of all fires and building fires.

	1990	1995	2000	2005	2010	2011	2012
All fires	17025	19543	17174	16551	16728	16157	14084
Building fires	10187	11694	10276	9903	9325	11447	9932

The building fires that occurred in the years 2007-2012 are subcategorised into six building types; detached houses, undetached houses, apartment buildings, industrial buildings, additional buildings and container fires.

Table 8.5.6 presents the calculated averages of the registered activity data for building fires for the years 2007-2010, divided in both damage size and building type. These data describe the average share of building fires from 2007-2010 of a certain type and size, in relation to all building fires in the same four years period.

Table 8.5.6 Average registered occurrence of building fires for 2007-2010. (DEMA).

	Size	Detached	Undetached	Apartment	Industry	Additional	Container	All building fires
Average, %	full	2.46	0.50	0.31	0.73	0.44	0.17	4.61
	large	4.01	1.14	1.09	1.69	3.08	1.92	12.93
	medium	5.24	2.33	6.15	2.92	4.30	18.46	39.40
	small	11.77	4.24	12.64	5.36	4.79	4.27	43.06
	all	23.47	8.21	20.19	10.70	12.61	24.82	100.00

It is assumed that the average percentages provided by the years 2007-2010 shown in Table 8.5.6 are compliant for the years 1990-2006. Hereby, similar activity data for building fires can be estimated back to 1990.

By applying the damage rates of 100 %, 75 %, 30 % and 5 % corresponding to the damage sizes full, large, medium and small, a full scale equivalent can be determined. Table 8.5.7 shows the calculated full scale equivalents (FSE). The full time series is shown in Annex 3G, Table 3G-5.4.

Table 8.5.7 Accidental building fires full scale equivalent activity data.

	1990	1995	2000	2005	2010	2011	2012
Container fires	750	861	756	729	594	729	584
Detached house fires	777	892	784	755	833	818	742
Undetached house fires	231	265	233	224	194	206	181
Apartment building fires	367	421	370	357	348	362	327
Industry building fire	320	368	323	311	281	334	298
Additional building fires	437	501	440	424	429	740	610

Emission factors

For building fires, emissions are calculated by multiplying the number of full scale equivalent fires with the emission factors. The emission factors are produced from different measurements and assumptions from literature and expert judgements. When possible, emission factors are chosen that represent conditions that are comparable to Denmark. By comparable is meant countries that have similar building traditions, with respect to the materials used in building structure and interior.

In the process of selecting the best available emission factors for the calculation of the emissions from Danish accidental building fires, a range of different sources has been studied. Unfortunately it is difficult to do an interrelated comparison of the different sources because they all establish emission factors on different assumptions and many of these assumptions are not fully accounted for.

Table 8.5.8 lists the emission factors that were chosen for 2012 as the best reliable and their respective references.

Table 8.5.8 Emission factors building fires, per FSE fire, 2012.

Compound	Unit /fire	Detached house	Undetached house	Apartment building	Industrial building	Additional building	Container	Source
CO ₂ - total	Mg	32.4	26.2	15.2	78.1	3.9	1.8	Blomqvist et al., 2002
CO ₂ - biogenic	Mg	26.4	21.4	12.4	67.6	3.2	0.2	Blomqvist et al., 2002
CO ₂ - non-biogenic	Mg	6.0	4.9	2.8	10.5	0.7	1.7	Blomqvist et al., 2002
CH ₄	kg	43.0	34.7	20.2	52.0	2.1	0.3*	NAEI, 2009

*Container fires have a different source of CH₄ emission factor than the other five categories; Blomqvist et al. 2002.

Emission factors for detached, undetached and apartment fires depend on the annual average floor space; see Table 8.5.9. Industrial, additional and container fires on the other hand are assumed to have a constant size/volume throughout the time series. Emission factors for detached, undetached and apartment fires for 1990-2012 are shown in Annex 3G, Table 3G-5.5a-c.

Emission factors from Aasestad (2008) are already specified for four of the six building types; detached houses, undetached houses, apartment buildings and industrial buildings. Aasestad (2008) and all other sources considered were altered to match the six building types. This alternation was performed simply by adjusting the average floor space for each of the building types respectively, whereas factors like loss rate and mass of combustible contents per area are not altered.

The average floor space in Danish buildings is stated in Table 8.5.9. The data are collected from Statistics Denmark and takes into account possible multiple building floors but not attics and basements. For the full time series see Annex 3G, Table 3G-5.6. The average floor space in industrial buildings, schools etc. is estimated to 500 square meters for all years and the average floor space for additional buildings, sheds etc. is estimated to 20 square meters for all years.

Table 8.5.9 Average floor space in building types (Statistics Denmark, 2013).

	1990	1995	2000	2005	2010	2011	2012
Detached houses	156	155	156	162	163	164	165
Undetached houses	129	129	131	131	134	132	134
Apartment buildings	75	75	75	76	77	78	78

Some emission factors are delivered in mass emission per mass burned. In order to connect these emission factors to the activity data, the total combustible building masses are estimated using the data from Table 8.5.10.

Table 8.5.10 Building mass per building type.

	Unit	Detached house	Un-detached house	Apartment building	Industry building	Additional building	Container
Average floor area*	m ²	165	134	78	500	20	-
Building mass per floor area	kg per m ²	40	40	35	30	30	-
Total building mass	Mg per fire	6.6	5.4	2.7	15.0	0.6	1

* 2012 numbers

Emission factors for container fires cannot be calculated based on an average floor space but on an average mass. The average mass of a container is set to 1 Mg and covers all types of containers, from small residential garbage containers to large shipping containers and waste/goods in storage piles.

No data was available for N₂O.

For more information on the emission factors, please refer to Hjelgaard (2013).

8.5.3 Accidental vehicle fires

Emissions that escape from vehicle fires are CO₂ and CH₄.

Methodological issues

Emissions from vehicle fires are calculated by multiplying the mass of vehicle fires with selected emission factors. Emission factors are not available for different vehicle types, whereas it is assumed that all the different vehicle types leads to similar emissions. The activity data are calculated as an annual combusted mass by multiplying the number of different full scale vehicle fires with the Danish registered average weight of the given vehicle type.

Activity data

As with accidental building fires, data for accidental vehicle fires are available through the Danish Emergency Management Agency (DEMA). DEMA provides very detailed data for 2007-2012; the remaining years back to 1990 are estimated by using surrogate data.

Table 8.5.11 shows the occurrence of fires in general and vehicle fires registered at DEMA. In 2007-2010 the average per cent of vehicle fires, in relation to all fires, was 20 %. The total numbers of vehicle fires in 1990-2006 are calculated using this percentage. The full time series is presented in Annex 3G, Table 3G-5.3.

Table 8.5.11 Occurrence of all fires and vehicle fires.

	1990	1995	2000	2005	2010	2011	2012
All fires	17025	19543	17174	16551	16728	16157	14084
Vehicle fires	3354	3850	3383	3260	3459	3255	2889

There are fourteen different vehicle categories. The activity data are categorised in passenger cars (lighter than 3500 kg), buses, light duty vehicles (vans and motor homes), heavy duty vehicles (trucks and tankers), motorcycles/mopeds, other transport, caravans, trains, boats, airplanes, bicycles, tractors, combine harvesters and machines.

In the same manner as accidental building fires, the 2007-2012 data from DEMA can be divided in four categories according to damage size. It is assumed that a full scale fire is a complete burnout of the given vehicle, and that a large, medium and small scale fire corresponds to 75 %, 30 % and 5 % of a full scale fire respectively. The total number of full scale equivalent (FSE) fires can be calculated for each of the fourteen vehicle categories for 2007-2012.

The total number of registered vehicles is known from Jensen et al. (2013) and Statistics Denmark (2013). By assuming that the share of vehicle fires in relation to the total number of registered vehicles, of every category respectively, can be counted as constant, the number of vehicle fires is estimated for the years 1990-2006.

Table 8.5.12 states the total number of national registered vehicles and the number of full scale equivalent vehicle fires. The full time series 1990-2012 is shown in Annex 3G, Table 3G-5.7a-c.

Table 8.5.12 Number of nationally registered vehicles and full scale equivalent vehicle fires.

	Passenger Cars		Buses		Light Duty Vehicles		Heavy Duty Vehicles	
	Registered	FSE fires	Registered	FSE fires	Registered	FSE fires	Registered	FSE fires
1990	1645454	479	8109	12	192317	19	45664	58
1995	1733242	504	14371	21	228074	22	48077	61
2000	1916364	557	15051	22	272386	27	50227	64
2005	2012216	585	15131	22	372674	36	49311	63
2010	2246675	646	14577	23	362385	38	44813	60
2011	2281539	584	13915	13	343355	43	43640	54
2012	2326778	514	13177	11	318668	32	42326	53
<i>Continued</i>								
	Motorcycles/Mopeds		Caravans		Train		Ship	
	Registered	FSE fires	Registered	FSE fires	Registered	FSE fires	Registered	FSE fires
1990	163133	58	86257	24	7156	9	2324	26
1995	165272	58	95831	26	6854	8	1911	21
2000	233309	82	106935	29	4907	6	1759	19
2005	273904	97	121350	33	3195	4	1792	20
2010	301562	83	142354	37	2740	2	1773	16
2011	295488	91	142764	34	2943	3	1768	21
2012	295798	82	142654	33	3055	2	1772	14
<i>Continued</i>								
	Airplane		Tractor		Combined Harvester		Bicycle	Other transport Machine
	Registered	FSE fires	Registered	FSE fires	Registered	FSE fires	FSE fires	FSE fires
1990	1055	1	131880	82	33594	56		
1995	1058	1	130028	81	27986	46		
2000	1070	1	111736	69	23272	39		
2005	1073	1	104551	65	20965	35		
2010	1152	1	102619	77	18889	32	4	58
2011	1132	0	102619	59	18889	21	3	50
2012	1111	0	102619	68	18889	18	2	50

The average weights of a passenger car, bus, light commercial vehicle, truck and motorcycle/moped are known for every year back to 1993 (Statistics Denmark, 2012). The corresponding weights from 1990 to 1992 and the average weight of the units from the remaining categories are estimated by an expert judgment, see Table 8.5.13 and Annex 3G, Table 3G-5.8.

Table 8.5.13 Average weight of different vehicle categories, kg.

Year	Cars	Buses	Vans	Trucks	Motorcycles/ Mopeds
1990	850	10000	2000	15000	86
1995	923	10807	2492	14801	97
2000	999	11195	3103	15214	103
2005	1068	11560	3793	13258	116
2010	1144	11804	4498	11883	133
2011	1154	11907	4296	11291	135
2012	1160	11625	4150	10844	136

It is assumed that the average weight of a boat equals that of a bus. That tractors and vans weigh the same and that trains, airplanes and combine harvesters have the same average weight as trucks.

Bicycles, machines and other transport can only be calculated for the years 2007-2012 due to the lack of surrogate data (number of nationally registered vehicles). The average weight of a bicycle, caravan, machine and other

transport is estimated as 12 kg, 90 % of a car, 50 % of a car and 40 % of a car respectively.

By multiplying the number of full scale fires with the average weight of the vehicles respectively, the total amount of combusted vehicle mass can be calculated. The result is shown in Table 8.5.14 and in Annex 3G, Table 3G-5.9a-c.

Table 8.5.14 Burnt mass of different vehicle categories, Mg.

Vehicle category	1990	1995	2000	2005	2010	2011	2012
Passenger cars	407	466	557	625	739	674	592
Buses	116	223	242	251	266	160	130
Light duty vehicles	37	55	82	138	171	185	133
Heavy duty vehicles	869	902	969	829	715	606	579
Motorcycle, moped	5	6	8	11	11	12	11
Other transport	-	-	-	-	33	29	29
Caravan	30	36	44	53	63	59	57
Train	128	121	89	51	24	28	23
Ship	257	228	218	229	189	249	160
Airplane	12	11	12	10	7	3	5
Bicycle	-	-	-	-	0	0	0
Tractor	164	202	216	247	347	254	283
Combine harvester	530	476	425	409	398	271	236
Machine	-	-	-	-	43	51	53
Total	2555	2727	2863	2858	3025	2624	2319

Emission factors

In the process of selecting the most reliable emission factors for the calculation of the emissions from Danish vehicle fires, a range of different sources have been studied. Unfortunately it is difficult to make an interrelated comparison of the different sources because they all establish emission factors on different assumptions and many of these assumptions are not fully accounted for. Table 8.5.15 lists the accepted emission factors and their respective references.

Table 8.5.15 Emission factors for vehicle fires, per Mg.

	Unit	Emission factor	Source
CO ₂	Mg	2.4	Lönnermark et al., 2006
CH ₄	kg	5	NAEI, 2009
N ₂ O	-	NAV	-

NAV = not available

8.5.4 Sludge spreading

Sludge from wastewater treatment plants is only spread out in the open with the purpose of fertilising crop fields. Greenhouse gas emissions from this activity are included in the agricultural sector, see Chapter 6.

8.5.5 Biogas production

Emissions from biogas production are divided and reported in different sections of this inventory according to the waste type and method.

Emissions from the combustion of biogas regardless of the origin are included in the energy sector and are allocated to the appropriate subsector in the Danish energy statistics. For the biogas production from organic waste with

the purpose of energy production, the fuel consumption rate of the biogas production plants refers to the Danish energy statistics. The applied emission factors are the same as for biogas boilers (see Chapter 3, Energy).

Biogas production from manure and the reduced emissions of CH₄ and N₂O from gasification of manure is included in Chapter 6, Agriculture.

Fugitive emissions of CH₄ from anaerobic digestion of sludge have been kept equal to last year's NIR submission, i.e. 1% of the biogas production, and are included in Chapter 8.3. In the below section a presentation of status for available plant level data supporting a methane generation and combustion efficiency of 99% (cf. Eq. 8.3.3) is presented.

Flaring and venting from biogas production at WWTPs

Flaring and venting may occur in different degrees at WWTPs which have implemented anaerobic treatment of sludge for biogas generation. Venting may occur intentionally or unintentionally if there are technical problems at the plant. Flaring is intentional combustion of biogas and occurs for regulation of the gas pressure.

Table 8.5.16 presents available information on the amount of flared and vented biogas in absolute numbers as well as in per cent of the recovered biogas at three of the biggest wastewater treatment plants in Denmark.

Table 8.5.16 Biogas production data for the WWTPs Lynetten, Avedøre and Damhusåen.

WWTP		2007	2008	2009	2010	2011	2012
Lynetten¹							
Biogas produced	Nm ³ /year		6330381	5942571	5792838	6695142	7154932
Flaring	Nm ³ /year		284615	659576	494972	946468	903613
	%		4,50%	11,10%	8,54%	14,14%	12,63%
Venting	Nm ³ /year		NR	NR	NR	NR	NR
	%		NR	NR	NR	NR	NR
Biogas consumed at plant	Nm ³ /year		6045766	5282995	5297866	5748674	6251319
Biogas reported to DEA ³	Nm ³ /year	4417670	4953913	4650708	4533525	3969338	6251318
	%		82%	88%	86%	69%	100%
Avedøre³							
Biogas produced	Nm ³ /year	3300000	3400000	3100000	3300000	3100000	3300000
Flaring	Nm ³ /year	140000	140000	54000	170000	36000	10000
	%	4,24%	4,12%	1,74%	5,15%	1,16%	0,30%
Venting	Nm ³ /year	0	2661	9179	54400	130063	50246
	%	0%	0,08%	0,30%	1,65%	4,20%	1,52%
Biogas consumed at plant	Nm ³ /year	3200000	3300000	3000000	3200000	2900000	3300000
Biogas reported to DEA ³	Nm ³ /year	2874932	3161242	2813589	2769597	2581438	2966742
	%	90%	96%	94%	87%	89%	90%
Damhusåen²							
Biogas produced	Nm ³ /year		2690037	1665416	2123357	1997333	1918325
Flaring	Nm ³ /year		57750	57750	307335	94150	236950
	%		2,15%	3,47%	14,47%	4,71%	12,35%
Venting	Nm ³ /year		NR	NR	NR	NR	NR
	%		NR	NR	NR	NR	NR
Biogas consumed at plant	Nm ³ /year		2632287	1607666	1816022	1903183	1681375
Biogas reported to DEA ³	Nm ³ /year		NR	NR	NR	NR	NR
	%		NR	NR	NR	NR	NR

¹Lynettefællesskabet (2009, 2010, 2011, 2012, 2013); ²Spildevandscenter Avedøre (2012, 2013); ³DEA (2013); ⁴NR: Not Reported.

As may be observed from Table 8.5.16, the amount of flaring is varying from year to year for the same plant as well as between WWTPs. The average flaring is 10 % at Lynetten (data for five years), 2.8 % at Avedøre (data for six years), and 7.4 % at Damhusåen (data for five years). Venting is only reported for Avedøre and constitute in average 1.3 % of the produced amount of biogas. Work is ongoing to extent the documentation for flaring and venting at biogas producing WWTPs (cf. Chapter XX).

The methodology used for estimating the CH₄ and N₂O emissions from wastewater handling are described in Chapter 8.3, Wastewater Handling.

Methodological issues

Biogas production in this sector covers fugitive emissions from the handling of biological waste, sludge and manure. This includes activities like storage, pre- and post-treatment during which anaerobic conditions may occur, and fugitive emissions from the anaerobic digestion that is the actual production. However, emissions from these activities are not currently included in the inventory.

This source also covers emissions from combustion of biogas in biogas production plants (mentioned as Gasification of biogas in the CRF tables) for the years 1994-2005 where these emissions existed. This activity is not occurring in 2006 - 2012. Pollutants from this activity are CH₄ and N₂O.

Activity data

Activity data for this source category are collected from the energy statistics (DEA, 2012). Combustion of biogas in biogas production plants occurred for the years 1994-2005¹.

Table 8.5.16 Combusted biogas.

	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
GJ	857	4711	4503	4447	35416	43688	40990	27270	30567	35544	28744	22034




Figure 8.5.2 Combusted biogas at biogas production plants.

¹ Activity data for combustion of biogas is in need of an updating, this will be done as soon as the energy statistics for 2013 is available.

Emission factors

Emission factors for combustion of biogas in biogas production plants are presented in Table 8.5.17. These are outdated emission factors for biogas boilers and will be updated in the next sector report.

Table 8.5.17 Emission factors for combustion of biogas, per GJ.

Pollutant	Unit	Emission factor
CH ₄	g	4
N ₂ O	g	2

8.5.6 Other

Other combustion sources included under Waste Other are the open burning of yard waste and bonfires.

Due to the cold and wet climatic conditions in Denmark wild fires very seldom occur. Controlled field burnings and the occasional wild fires are categorised under the Chapters on 6 Agriculture and 7 Land Use, Land Use Change and Forestry (LULUCF) respectively.

In Denmark, the open burning of private yard waste is under different restrictions according to the respective municipality. These restrictions involve what can be burned but also the quantity, how, when and where, or in some cases a complete ban is imposed. The burning of yard waste is not allowed within urban areas (DEPA, 2011b). There is no registration of private waste burning and the activity data on this subject are very difficult to estimate. Citizens are generally encouraged to compost their yard waste or to dispose of it through one of the many waste disposal/recycling sites.

The occurrence of bonfires at Midsummer's Eve and in general are likewise not registered, therefore it has not been possible to obtain activity data and consequently, bonfires are not included in this inventory.

8.6 Uncertainties and time series consistency

Two set of uncertainty estimates are made for the Danish emission inventory for greenhouse gases based on Tier 1 and Tier 2 methodology, respectively. The uncertainty models follow the methodology in the IPCC Good Practise Guidance (IPCC, 2000). Tier 1 is based on the simplified uncertainty analysis and Tier 2 is based on Monte Carlo simulations.

8.6.1 Input data

Solid Waste Disposal on Land

The waste amounts for solid waste disposal on land are registered in a national database held by the Danish EPA and assessed to be of high quality resulting in the adoption of an uncertainty for reported waste amounts of 10 %.

Input parameter uncertainties for SWDS considered in the Tier 1 uncertainty analysis are based on the IPCC (IPCC 2000, page 5.12, Table 5.2) default values and provided in Table 8.6.1.

Table 8.6.1 Tier 1 input parameter uncertainty, %.

Parameter	Parameter Uncertainty		Note
	ID	%	
The Waste amount sent to SWDS	<i>W</i>	10	Since the amounts are based on weighing at the SWDS the lower value in IPCC (2000), is used
Degradable Organic Carbon	<i>DOC_i</i>	50	Highest value, IPCC 2000, page 5.12, Table 5.2
Fraction of DOC dissimilated	<i>DOC_f</i>	30	Highest value, IPCC 2000, page 5.12, Table 5.2
Methane Correction Factor	<i>MCF</i>	10	IPCC, 2006
Fraction of CH ₄ in landfill gas		10	Medium value, IPCC 2000, page 5.12, Table 5.2
Methane Generation Rate Constant	<i>k</i>	100	IPCC 2000, page 5.12, Table 5.2

The waste amounts for solid waste disposal on land are registered in a national database held by the Danish EPA and assessed to be of high quality resulting in the adoption of an uncertainty for reported waste amounts of 10 %. The default uncertainty range for the methane generation constant, *k*, is: -40 % to +300 %.; for the Tier 1 uncertainty calculation it has been set to 100 % (Limpert et al., 2001). For the remaining parameters default uncertainties are used until country-specific parameters becomes available.

The uncertainty on the implied emission factor, *U_{ief}*, is based on uncertainty estimates in Table 8.6.1 and is approximated with IPCC (2000) Equation 6.4 equals

$$U_{ief} \% = \text{SQRT}(50^2 + 30^2 + 10^2 + 10^2 + 100^2) = 117.9 \%$$

These uncertainties give the combined Tier 1 uncertainty on the emission from SWDS of: $\text{SQRT}(10^2 + 117.9^2) = 118.3 \%$.

Wastewater Handling

The uncertainty levels used in the Tier 1 and 2 uncertainty models are shown in Table 8.6.2.

Table 8.6.2 Estimated uncertainty rates for activity data and emission factors, %.

95 % confidence interval uncertainties	Activity data	Emission factor
N ₂ O, WWT, direct	20	53
N ₂ O, WWT, indirect	42	42
CH ₄ , Sewer system and WWTP processes	24	32
CH ₄ , Anaerobic digestion	24	39
CH ₄ , Septic tanks (scattered houses)	31	32

Default IPCC values are assumed to be given at 95 % confidence level. For the country-specific activity data, the standard deviation of different data sources has been used for deriving per cent uncertainty estimates. Annex 3G, Table 3G-3.6 elaborates on the different values and their references.

Uncertainties have been derived from IPCC default values and uncertainties in country-specific parameters, respectively (cf. Annex 3G, Table 3G-3.6).

Waste Incineration

The uncertainty of the number of human cremations is miniscule, however for the purpose of uncertainty calculation it has been set to 1 %. The uncertainty of the activity data from animal cremations is also minimal for the most recent years (1998-2012) but is increasing back in time (to 67 % in 1990). The uncertainty is set to 67 % in 1990 and 5 % for 2012 (Authors expert judgement).

Table 8.6.4 lists the 95 % confidence interval uncertainties for activity data and emission factors used in this inventory and at the present level of available information.

Table 8.6.4 Estimated uncertainty rates for activity data and emission factors, %.

95 % confidence interval uncertainties	CO ₂	CH ₄	N ₂ O
Human cremation			
Activity data	-	1	1
Emission factor	-	150	150
Animal cremation			
Activity data	-	5/67	5/67
Emission factor	-	150	150

Waste Other

Activity data for composting are estimated for the years 1990-1994 and 2010-2012 resulting in a higher level of uncertainty these years; this is set at 40 %.

The uncertainty of the total number of accidental fires is very small, but the division into building and transportation types and also the calculation of full scale equivalents will lead to some uncertainty, partly caused by the category "other". The uncertainty for both building and vehicle activity data is therefore set to 10 % for all years. The uncertainty is however lowest for the most recent years (2007-2012) (Authors expert judgement).

Activity data for combustion of biogas at biogas production facilities are available from the national energy statistics; the uncertainty for this activity is set to 5 %. However, this activity only exists in 1994-2005 and is therefore not part of the uncertainty calculations.

Table 8.6.5 lists the 95 % confidence interval uncertainties for activity data and emission factors used in this inventory and at the present level of available information. The uncertainties are assumed valid for all years 1990-2011.

Table 8.6.5 Estimated uncertainty rates for activity data and emission factors, %.

95 % confidence interval uncertainties	CO ₂	CH ₄	N ₂ O
Compost production			
Activity data	-	40	40
Emission factor	-	100	100
Accidental building fires			
Activity data	10	10	-
Emission factor	300	500	-
Accidental vehicle fires			
Activity data	10	10	-
Emission factor	500	700	-

The input parameter uncertainties are at the same aggregation level as reported in last year's NIR (Nielsen et al., 2013).

8.6.2 Tier 1 uncertainty results

The Tier 1 uncertainty estimates for the waste sector are calculated from 95 % confidence interval uncertainties, results are shown in Table 8.6.6.

The overall uncertainty interval for greenhouse gases (GHG) is estimated to be ± 86.8 % and the trend in GHG emission, calculated as the per cent change in GHG emissions in 2011 compared to 1990, is -37.8 % ± 13.6 %.

Table 8.6.6 National Tier 1 uncertainty estimates for the waste sector.

Pollutant	National emission, 2012, Gg	Total emission uncertainty, %	Trend* 1990-2011, %	Trend uncertainty, %
GHG**	1063.97	± 78.3	-32.4	± 20.7
CO ₂	16.36	± 261.1	-6.7	± 11.0
CH ₄	41.14	± 89.1	-38.7	± 14.0
N ₂ O	0.64	± 73.9	69.4	± 175.4

*Per cent change in emission in 2011 with respect to the base year 1990.

**GHG emissions are calculated in units of CO₂ equivalents.

8.6.3 Tier 2 uncertainty results

The Tier 2 uncertainty estimates for the waste sector are calculated from the input data presented in Section 8.6.1; results are shown in Table 8.6.7. The calculations are based on a Monte Carlo approach as described in Chapter 1.7.

Table 8.6.7 National tier 2 uncertainty estimates for the waste sector, [Gg]

	1990 National emission, [Gg]			2012 National emission, [Gg]			1990-2012 Trend, [Gg]		
	Uncertainty interval, Mean [%]			Uncertainty interval, Mean [%]			Uncertainty, Mean [%]		
	lower (-)	upper (+)		lower (-)	upper (+)		lower (-)	upper (+)	
GHG	1629	-56	162	1138	-45	116	-454	-97	159
CO ₂	27	-66	355	25	-67	344	-2	-6	32
CH ₄	88	-62	179	31	-51	137	-54	-76	180
N ₂ O	0.04	-53	139	0.43	-51	118	0.38	-1134	457

Greenhouse gas (GHG) emissions are calculated in CO₂ equivalents.

8.6.4 Time series consistency and completeness

Solid Waste Disposal on Land

Registration of the amount of waste has been carried out since the beginning of the 1990s in order to measure the effects of action plans. The activity data are, therefore, considered to be consistent through the time series to make the activity data input to the FOD model reliable.

The consistency of the emissions and the emission factor is a result of the same methodology and the same model used for the whole time series. The parameters in the FOD model are the same for the whole time series. The use of a model of this type is recommended in IPCC (1997) and IPCC (2000).

As regards completeness, the waste amounts used, as registered in the ISAG system, do not only include traditional Municipal Solid Waste (MSW), but

also non-MSW such as Industrial Waste, Building and Construction Waste and Sludge. The composition of these waste types is, according to Danish data, used to estimate DOC values for the waste types (refer IPCC 2000, page 5.10). Improvement are planned to increase completeness and appropriateness of waste types as described in Chapter 8.9

Wastewater Handling

Consistency and completeness have been improved by access to the Danish Water Quality Parameter Database (www.miljoportal.dk) and the Danish Sludge Database held by DEPA.

Data regarding industrial on-site wastewater treatment processes is not available at a level that allows for calculation of the on-site industrial contribution to CH₄ or N₂O emissions. The degree to which industry is covered by the estimated emission is, therefore, dependent on the amount of industrial wastewater connected to the municipal sewer system. Any direct emissions from pre-treatment on-site are not covered in this inventory.

Waste Incineration

Activity data for human cremation is considered to be consistent as these data have been collected by DKL throughout the time series. Activity data for animal cremation on the other hand is not fully consistent. Data for 1998-2012 are gathered directly from the crematoria and data for 1990-1997 are estimated by the author's expert judgement, no surrogate data or data regression is possible.

Emission factors and calculation method are consistent throughout the time series for both human and animal cremation.

Cremation of both corpses and carcasses is considered to be complete. Open burning of carcasses is illegal and therefore not occurring in Denmark, and small-scale incinerators are not known to be used at Danish farms.

Waste Other

For compost production, activity data are not consistent as data are only available for 1995-2009. Data for 1990-1994 and 2010-2012 along with data for home composting are estimated through linear regression and with surrogate data respectively. Emission factors and calculation method are consistent throughout the time series.

Emissions from compost production are believed to be complete; calculations include composting at all nationally registered sites and best available estimated data for home composting.

For accidental fires, DEMA provides detailed data for 2007-2012 and the total number of nationally registered fires for 1990-2012. Activity data for accidental fires are there for believed to be consistent. Both emission factors and calculation method are also consistent throughout the time series.

Emissions from accidental fires are believed to be complete. Field burning of agricultural residue is included in Chapter 6 Agriculture.

8.7 QA/QC and verification

In general terms, for this part of the inventory, the Data Storage (DS) Level 1, 2 and 4 and the Data Processing (DP) Level 1 can be described as follows.

8.7.1 Data Storage Level 1

The external data level refers to the placement of the original input data used for estimating annual activity and emission factors in the waste sector. Data references in terms of reports and databases used for deriving input for the emission calculations. Reports and a list of links to external data sources are stored in a common data storage system including all sectors of the annual NIR.

Table 8.7.1a Overview of annually stored external data sources at DS level1.

http, file or folder name	Description	AD or EF	Reference	Contact	Data agreement/ Comment
DCE data-exchange folder U:\ST_ENVS-Luft-Emi\Inventory\2012\6_Waste\Level_1a_Storage\ge*	Inventory data storage system	AD and EF	DCE		
Report series published and available from the Danish Environmental Protection Agency www.mst.dk	Reported sludge and water quality parameters	AD	Report series from DEPA: "Wastewater sewage sludge from municipal and private wastewater treatment plants" (1997-2005) "Point sources" (1993-2005)	Marianne Thomsen (mth@dmu.dk)	Public available reports
Report series published by the Agency for Spatial and Environmental Planning (ASEP) and available from the Danish Nature Agency (DNA): www.nst.dk			Report series: "Point sources" (2006-2012)	Naturstyrelsen Vestjylland Anna Gade Holm (angho@nst.dk) Marianne Thomsen (mth@dmu.dk)	Public available reports
Danish Water Quality parameter Database	Annually reported wastewater characteristics at plant level which includes all years 1990- 2012	AD	www.miljoeportalen.dk	Naturstyrelsen Vestjylland Anna Gade Holm (angho@nst.dk) Marianne Thomsen (mth@dmu.dk)	Authorised access
Danish Sludge Database	Annually reported sludge characteristics at plant level	AD	DEPA	Linda Bagge (bagge@mst.dk) Marianne Thomsen (mth@dmu.dk)	none
I:\ROSPROJ\LUFT_EMI\Inventory\2011\6_WaU:\ST_ENVS-Luft-Emi\Inventory\2012\data 6_Waste\Level_1a_Storage\6A Solid Waste Disposal	Emi\Inventory\2012\data 6_Waste\Level_1a_Storage\6A SWDS	Activity data	The Danish Environmental Protection Agency, database on all registered Danish waste. Available at: http://www2.mst.dk	Unit for Soil and Waste Eik Kristensen (eikri@mst.dk)	The amounts are registered due to statutory requirements

Continued

I:\Rosproj\LUFT_EMI\Energy\2011	Basic data DS1 Dataset for energy- producing SWDS	CH ₄ recovery (DEA) data	The Danish Energy Agency	Peter Dal (pd@ens.dk)	Prepared due to the obligation of DEA
I:\ROSPROJ\LUFT_EMI\Inventory\2011\6_WaExcel file with the ste\Level_1b_Processes\6A Solid Waste Disposal	FOD model swds_fod_model 2011.xls DP1	Parameters of the FOD model	IPCC 1997 and 2000	Marianne Thomsen (mth@dmu.dk)	
http://www.dkl.dk	Number for crema- tions	AD	Association of Danish Crematories	Hanne Ring hr@dkl.dk	Public access
http://www.statistikbanken.dk	Statistics for popula- tion, buildings and vehicles	AD	Statistics Denmark		Public access

*The data storage level 1 consists of DEPA reports and data extracted from other sources listed in the Table.

Table 8.7.1b Overview of annually stored external data sources at DS level1 (Continued).

http, file or folder name	Description	AD or EF	Reference	Contact	Data agree- ment/ Comment
U:\ST_ENVS-Luft- Emi\Inventory\2012\6_Waste\Level_1a_Stocarcasses rage\6C Waste Incineration	Cremated animal	AD	Dansk Dyrekremering ApS	Knud Ribergaard in-fo@danskdyrekremering.dk	Personal contact
U:\ST_ENVS-Luft- Emi\Inventory\2012\6_Waste\Level_1a_Stocarcasses rage\6C Waste Incineration	Cremated animal	AD	Ada's Kæledyrskrematorium ApS	Anders Oxholm anders@adakrem.dk	Personal contact
U:\ST_ENVS-Luft- Emi\Inventory\2012\6_Waste\Level_1a_Stocarcasses rage\6C Waste Incineration	Cremated animal	AD	Kæledyrskrematoriet	Annette Laursen dyrepension@skylinemail.dk	Personal contact
https://statistikbank.brs.dk	Categorized fires	AD	The Danish Emergency Management Agency	Steen Hjere Nonnemann shn@beredska.bs styrelsen.dk	Public access
http://www2.mst.dk/udgiv/publikationer/2010/978-87-92668-21-9/pdf/978-87-92668-22-6.pdf	Waste categories for composting	AD	Danish Environmental Protection Agency (DEPA), Waste Statistics		Public access

8.7.2 Data Processing Level 1

This level comprises a stage where the external data extracted from the waste data system (DEPA, 2013) are processed internally. For SWDS data are prepared for the DCE First Order of Decay model by allocation of the reported waste amounts according to the European Waste Codes (EWC) as presented in Thomsen et al., 2014 and in Annex 3G, Table 3G-2.4. The model runs in excel and the output are stored inside the excel file. For Wastewater Handling data are prepared for the input to the country-specific models. Programming as to automatically calculations based on activity data and

emission factors are not yet fully operational. Calculations are carried out and the output stored in a not editable format each year. The DP at level 1 has been improved to fit into a more uniform and easily accessible data reporting format. Regarding the derivation of activity data and emission factors used in the model calculations, this year's improvements are documented in Chapter 8.3.1.

For the CRF categories 6C and 6D, the activity data and emission factors are recalculated to match each other by using national average data like the average floor space in houses etc.

8.7.3 Data Storage Level 2

Data Storage Level 2 is the placement of selected output data from the calculation of emissions as inventory data on SNAP levels in the Access (CollectER) database.

8.7.4 Data Storage Level 4

Data Storage Level 4 is the placement of the calculated output data from the calculation of emissions as data on SNAP levels in the CRFs.

8.7.5 Points of measurement

The present stage of QA/QC for the Danish emission inventories for the waste sector is described below for DS level 1, 2 and 4 and DP level 1 Points of Measurement (PMs). This is to be seen in connection with the general QA/QC description in Section 1.6 and, especially, 1.6.10 on specific description of PMs common to all sectors, general to QA/QC.

Data Storage level 1	1. Accuracy	DS.1.1.1	General level of uncertainty for every dataset including the reasoning for the specific values
----------------------	-------------	----------	--

The sources of data described in the methodology sections and in DS.1.2.1 and DS.1.3.1 are used in this inventory. It is the accuracy of these data that define the uncertainty of the inventory calculations.

With regard to the general level of uncertainty for SWDS, the amounts in waste fractions/categories are reasonably certain (per cent uncertainty set equal to 10 %, cf. Table 8.6.1. Due to the statutory environment for these data, while the distribution of waste fractions according to waste type and their content of DOC is more uncertain (per cent uncertainty set equal to 50 %, cf. Table 8.6.1). It is generally accepted that FOD models for CH₄ emission estimates offer the best and the most certain way of estimation. The half-life in the FOD models is an important parameter with some uncertainty (cf. Table 8.6.1).

The input parameter uncertainties for Wastewater Handling have been derived from standard deviations between activity data extracted from national databases and reported national statistics as shown in Table 8.6.2. Uncertainties on defaults numbers are taken from the IPCC (1997 and 2000). Uncertainty of activity data are based on simple standard deviations accompanying the annual reported monitoring data.

For Waste Incineration and Waste Other the level of uncertainty is generally low for activity data but higher for emission factors, cf. Table 8.6.4 and 8.6.5. Expert judgments are used whenever default uncertainties are not available.

Data Storage level 1	2.Comparability	DS.1.2.1	Comparability of the emission factors/calculation parameters with data from international guidelines, and evaluation of major discrepancies.
----------------------	-----------------	----------	--

Comparison of Danish data values from external data sources with corresponding data from other countries has been carried out in order to evaluate discrepancies.

Comparison of Danish data values with data sources from other countries has been carried out as presented in the national verification report by Fauser et al., 2007 and 2014 and in the methodology report by Thomsen & Lyck, 2005 and Thomsen and Hjelgaard (2014).

Data Storage level 1	3.Completeness	DS.1.3.1	Ensuring that the best possible national data for all sources are included, by setting down the reasoning behind the selection of datasets.
----------------------	----------------	----------	---

SWDS

- Danish Environmental Protection Agency (DEPA), ISAG database and the new waste data system: amounts of the various waste fractions deposited (refer to Section 8.2.1).
- A Danish investigation and verification of the overall mass balance upon allocating waste fractions within the old ISAG and the new waste data system into 18 defined waste types (Thomsen and Hjelgaard, 2014).
- Danish Energy Agency (DEA): Official Danish energy statistics: CH₄ recovery data.

The selection of sources is obvious. The ISAG database is based on statutory registrations and reporting from all Danish waste treatment plants for all waste entering or leaving the plants. Information concerning waste in the previous year must be reported to the DEPA each year, no later than January 31. Registration is made by mass for the individual waste categories. A New waste reporting system has been implemented and the nine waste types reported in the last NIR extended to 18 waste fractions of which 11 are characterised as inert. The individual waste type characteristics have been documented in Table 8.2.2 and Table 8.2.3 as well as in Annex 3G, Table G3-2.3 and G3-2.4.

For recovery data, the DEA registers the energy produced from plants where installations recover CH₄ for the energy statistics (cf. chapter 8.9 on planned improvements for the waste category 6.A..

For the parameters of the FOD model, references are made to IPCC (1997, 2000 and 2006).

WWT

- The Danish sludge database (Table 8.3.4)
- The Danish Water Quality Parameter Database (www.miljoportal.dk)

Waste Incineration

- Tables from Association of Danish Crematories available online
- Direct contact with the Danish animal crematories
- Emission factors from literature

Data from the Association of Danish Crematories is based on annual reporting from all Danish crematories. Specific reported data are available for the complete time series.

Waste Other

- Waste Statistics (DEPA, 1996, 1998, 1999, 2001a, 2001b, 2002, 2004a, 2004b, 2005, 2006a, 2006b, 2008, 2010a, 2011a)
- Danish Emergency Management Agency (DEMA) database
- Emission factors from literature

The waste statistics are based on data from the ISAG database, which is the only Danish registration of waste amounts. Also the DEMA database is the only provider of data on accidental fires; data for newer years (2007-2011) are extremely detailed.

Data Storage level 1	4.Consistency	DS.1.4.1	The original external data has to be archived with proper reference.
----------------------	---------------	----------	--

Data are predominantly extracted from the internet (ISAG, Water Quality Parameter, sludge, Statistics Denmark, DEMA database, human cremation). The origin of external activity data has been preserved as much as possible by saving them as original copies in their original form. Files are saved for each year of reporting, in this way changes to previously received data and calculations is reflected and explanations are given. Specific information from reports, industries and experts are saved as e-mails and pdf files.

Data Storage level 1	6.Robustness	DS.1.6.1	Explicit agreements between the external institution holding the data and DCE about the conditions of delivery.
----------------------	--------------	----------	---

As stated in DS.1.4.1 most data are obtained from the internet. It is a statutory requirement that amounts of waste are reported annually to DEPA, no later than January 31 for the previous year. No explicit agreements have been made with external institutions.

However for Wastewater Handling, this point may still be critical due to the missing timing full reporting and completeness of the databases held by the ASEP and DEPA respectively with respect to the submission date of the annual NIR.

Data Storage level 1	7.Transparency	DS.1.7.1	Listing of all archived datasets and external contacts.
----------------------	----------------	----------	---

Contact persons related to the delivery of specific data are provided in Table 8.7.1a and b.

For a listing of all archived external datasets, see DS 1.3.1.

Data Processing level 1	1. Accuracy	DP.1.1.1	Uncertainty assessment for every data source not part of DS.1.1.1 as input to Data Storage level 2 in relation to type and scale of variability.
-------------------------	-------------	----------	--

No data are used in addition to those included in DS.1.1.1. Uncertainties are reported in Section 8.6.

Data Processing level 1	2.Comparability	DP.1.2.1	The methodologies have to follow the international guidelines suggested by UNFCCC and IPCC.
-------------------------	-----------------	----------	---

The methodological approach is based on the detailed methodology as outlined in the Emission Inventory Guidebook. The calculation used for SWDS is a Tier 2 methodology from IPCC (1997, 2000 and 2006). For WWT the calculations follow the IPCC (1997 and 2000). Exemptions have been documented whenever occurring. The inventory calculations for Waste Incineration and Waste Other are a simple multiplication of activity data and emission factors. See also DS.1.3.1.

Data Processing level 1	3.Completeness	DP.1.3.1	Identification of data gaps with regard to data sources that could improve quantitative knowledge.
-------------------------	----------------	----------	--

For SWDS there is no quantitative knowledge in the methodology on either (1) the shift in waste fractions within waste categories for 1960-1984 and 1986-1993, (2) the development over time of the DOC content in individual waste fractions or (3) possible individual conditions relating to the SWD sites. On-going research might change this lack.

Data on separate industrial WWTPs. Information on methane emissions for separate industries may be of importance.

Emission factors for cremation and accidental fires are gathered from literature studies. There is no Danish literature or measurements available on greenhouse gas emissions from these categories.

Activity data for accidental fires for the years 1990-2006 are not subcategorised into vehicles, buildings or sizes.

Data Processing level 1	4.Consistency	DP.1.4.1	Documentation and reasoning of methodological changes during the time series and the qualitative assessment of the impact on time series consistency.
-------------------------	---------------	----------	---

There is no change in calculation procedure during the time series and the activity data are, as far as possible, kept consistent for the calculation of the time series. Any changes in calculation procedures are noted for each year's inventory, cf. Section 8.6.4.

Data Processing level 1	5.Correctness	DP.1.5.1	Verification of calculation results using time series
-------------------------	---------------	----------	---

The time series of activities and emissions in the model output, in the SNAP source categories and in the CRF format have been prepared. The time series are examined and significant changes are checked and explained. Comparison is made with the previous year's estimate and any major changes are verified.

Data Processing level 1	5. Correctness	DP.1.5.2	Verification of calculation results using other measures
-------------------------	----------------	----------	--

The correct interpretation in the model/calculation of the methodology and the parameterisation has been checked as far as possible.

Data Processing level 1	7. Transparency	DP.1.7.1	The calculation principle, the equations used and the assumptions made must be described.
-------------------------	-----------------	----------	---

The calculation principle and equations are described in “Methodological issues”, Section 8.2.1, 8.3.2, 8.4.1, 8.4.2, 8.5.1, 8.5.2 and 8.5.3.

Data Processing level 1	7. Transparency	DP.1.7.2	Clear reference to dataset at Data Storage level 1
-------------------------	-----------------	----------	--

Refer to the table at the start of this Section and DS.1.1.1 (8.7.1a and b).

The calculation principle and equations are described in “Methodological issues”, Section 8.2.1, 8.3.2, 8.4.1, 8.4.2, 8.5.1, 8.5.2 and 8.5.3.

Data Processing level 1	7. Transparency	DP.1.7.3	A manual log to collect information about recalculations.
-------------------------	-----------------	----------	---

Recalculation and changes in the emission inventories are described in the NIR whenever occurring. The logging of the changes takes place in the annual model file.

Data Storage level 2	5. Correctness	DS.2.5.1	Check if a correct data import to level 2 has been made
----------------------	----------------	----------	---

The transfer of emission data from level 1, storage and processing, to data storage level 2 is manually checked. This check is performed, comparing model output and report files made by the CollectER database system.

Data Storage level 4	4. Consistency	DS.4.4.3	The IEFs from the CRF are checked both regarding level and trend. The level is compared to relevant emission factors to ensure correctness. Large dips/jumps in the time series are explained.
----------------------	----------------	----------	--

See DP.1.5.1 and DP.1.5.2.

8.8 Source specific recalculations

Table 8.8.1 presents the recalculations to the waste sector for this year's inventory. Tables with the full time series 1990-2011 are shown in Annex 3G, Table 3G-6.1, 3G-6.2, 3G-6.3, 3G-6.4 and 3G-6.5.

Table 8.8.1. Changes in emissions from the waste sector compared with last year's submission.

	Unit	1990	1995	2000	2005	2010	2011	2012
Solid Waste Disposal on Land								
CH ₄ , previous inventory	Gg	70.4	60.5	0.57	40.8	34.3	33.3	
CH ₄ , recalculated	Gg	65.0	57.6	0.57	41.1	34.3	34.8	33.2
Change, CO ₂ -equivalents	Gg	-112.0	-60.1	0.71	5.9	0.2	32.3	
Change	%	-7.6	-4.7	0.71	0.7	0.0	4.6	
Wastewater Handling								
CH ₄ , previous inventory	Gg	3.15	3.27	3.52	3.54	3.59	3.62	
CH ₄ , recalculated	Gg	3.12	3.23	3.48	3.45	3.53	3.56	3.52
N ₂ O, previous inventory	Gg	0.34	0.35	0.29	0.27	0.25	0.26	
N ₂ O, recalculated	Gg	0.34	0.35	0.29	0.27	0.25	0.26	0.23
Change, CO ₂ -equivalents	Gg	-0.70	-0.84	-0.90	-1.93	-1.24	-1.29	
Change	%	-0.41	-0.48	-0.55	-1.21	-0.82	-0.83	
Waste Incineration								
CH ₄ , previous inventory	Mg	0.51	0.55	0.57	0.62	0.76	0.71	
CH ₄ , recalculated	Mg	0.51	0.55	0.57	0.62	0.76	0.71	0.70
N ₂ O, previous inventory	Mg	0.64	0.69	0.71	0.77	0.95	0.88	
N ₂ O, recalculated	Mg	0.64	0.69	0.71	0.77	0.95	0.88	0.88
Change, CO ₂ -equivalents	Gg	0.00	0.00	0.00	0.00	0.00	0.00	
Change	%	0.00	0.00	0.00	0.00	0.00	0.00	
Waste Other								
CO ₂ , previous inventory	Gg	18.28	20.11	18.77	18.20	18.19	18.21	
CO ₂ , recalculated	Gg	17.54	19.60	18.40	18.14	18.35	18.45	16.36
CH ₄ , previous inventory	Gg	1.40	1.80	3.06	3.31	3.91	4.02	
CH ₄ , recalculated	Gg	1.46	1.95	3.32	3.50	4.17	4.29	4.38
N ₂ O, previous inventory	Gg	0.04	0.05	0.13	0.10	0.14	0.14	
N ₂ O, recalculated	Gg	0.04	0.07	0.52	0.20	0.36	0.38	0.41
Change, CO ₂ -equivalents	Gg	1.61	9.18	123.38	33.42	73.10	79.76	
Change	%	2.70	12.43	99.03	27.80	51.10	54.28	
Total Waste								
CO ₂ , previous inventory	Gg	18.3	20.1	18.8	18.2	18.2	18.2	
CO ₂ , recalculated	Gg	17.5	19.6	18.4	18.1	18.3	18.4	16.4
CH ₄ , previous inventory	Gg	74.9	65.5	57.2	47.6	41.8	40.9	
CH ₄ , recalculated	Gg	69.6	62.8	56.7	48.0	42.0	42.7	41.1
N ₂ O, previous inventory	Gg	0.4	0.4	0.4	0.4	0.4	0.4	
N ₂ O, recalculated	Gg	0.4	0.4	0.8	0.5	0.6	0.6	0.6
Change, CO ₂ -equivalents	Gg	-111.1	-51.7	107.5	37.4	72.0	110.8	
Change	%	-6.5	-3.4	8.0	3.3	7.1	11.1	

8.8.1 Solid waste disposal on land recalculations

The recalculation of emissions from Solid Waste Disposal on Land is caused by an in depth 1) disaggregation and re-allocation of the old ISAG waste categories according to 18 characterised waste types and 2) allocation of waste amounts reported in the new waste data system into the same 18 defined waste types within this years NIR. The 18 waste types are characterised in terms of their content of degradable organic matter, DOC_i, and half-life times, t_{1/2}, as provided in Table X. A detailed description of the waste characterisation of ISAG and EWC waste types and the weights used for allocation into the new 18 waste types have been according to mass conservation. The impacts of all changes made to the input parameters and waste amounts (according to types) on the calculated emissions from SWDS are provided in Table 8.8.1.

8.8.2 Wastewater Handling recalculations

For Wastewater Handling recalculations have been made to the N₂O emission. Minor changes in the effluent tonnes N for the years 2007-2010 have been made due to updated information from the Danish EPA and updated data on amount of biogas produced from the anaerobic digestion of sludge. The major reason for the observed reduction of the total emission from Sector 6.B is due to the elimination of a correction factor that was not justified after verification of nitrogen effluent data with the newest reporting of effluent data in the report series “point sources” published by the Danish EPA (DEPA, 2012). As mentioned in Section 8.9 a verification of completeness in the activity data throughout the time series has been initiated and will be published in a sector report (Chapter 8.9).

The effect of the correction is shown in Table 8.8.1.

8.8.3 Waste Incineration recalculations

No recalculations were made for Waste Incineration.

8.8.4 Waste Other recalculations

For sector 6.D. Waste Other several recalculations were made. Changes were made in the vehicle fires and composting source categories.

For vehicle fires; the time-series for vehicle population delivered from an external source has been updated along with the estimated average weights of some vehicle types, the result is a decrease of CH₄ and CO₂ emissions from 1990-2006 and an increase for 2007-2011.

Emission factors have been updated for composting of sludge and organic municipal waste which have resulted in increased emissions of CH₄ and N₂O.

The joint effect of these recalculations is an increase in CH₄ emissions between 2 % (1990) and 12 % (2002), for CO₂ emissions have decreased from 1990 (4 %) to 1997 (0.4 %) and increased from 1998 (14 %) to 2011 (1 %), and finally the N₂O emissions have increased for all years with between 10 % (1990) and 370 % (2002).

8.9 Source specific planned improvements

For the category 6a.SDWS, plant specific data was made available from the new waste reporting system in November 2013, which forms the basis for exploring the possibility of obtaining model input parameter data at plant level. The first step towards a plant level emission model for SWDS in Denmark is to review available data on methane emissions and recovery data as well as plant specific input parameters for the FOD model (e.g. [DEPA, 2011](#)). Special focus before the next reporting year, will be to verify plant level emission data, using the 18 defined waste types, against emission data reported to the National PRTR reporting system, as well as to verify plant level recovered emissions against the Energy statistics and Energy Producer Accounting system produced by the Danish Energy Agency.

Regarding 6B. Wastewater Handling, new data on extension of the collective sewerage system in Denmark have not yet become available. We plan still to document reduced emissions from scattered settlements and to include such

documentation in the next inventory in terms of a reduction in the fraction of the population not connected to the municipal sewer system.

Likewise, alternative solutions to the treatment of wastewater from scattered houses as well as development in aquaculture and marine fish farming activities in Denmark will influence indirect N₂O emissions, why improvements are expected. However, these improvements are long term aspects realised as the necessary documentation becomes available.

There are no other planned improvements for the waste sector.

8.10 References

Aasestad, K. 2008: The Norwegian Emission Inventory 2008, Documentation of methodologies for estimating emissions of greenhouse gases and long-range transboundary air pollutants. Available at: http://www.ssb.no/english/subjects/01/04/10/rapp_emissions_en/rapp_200848_en/rapp_200848_en.pdf (05/02-2014).

Adouania, N., Lendormia, T., Limousya, L. & Sirea, O. 2010: Effect of the carbon source on N₂O emissions during biological denitrification. *Resources, Conservation and Recycling*, 54, 299-302.

Agency for Spatial and Environmental Planning (ASEP), 2007: Point sources 2006. In Danish: Punktkilder 2006. Det nationale program for overvågning af vandmiljøet; Fagdatacenterrapport. Available at: http://www.naturstyrelsen.dk/Udgivelser/Aarstal/2007/Punktkilder_2006_Det_nationale_program_for_overvaagning_af_vandmiljoet_Fagdatacenterrapport.htm (26/02-2014).

Agency for Spatial and Environmental Planning (ASEP), 2009: Point sources 2007. In Danish: Punktkilder 2007. Det nationale program for overvågning af vandmiljøet; Fagdatacenterrapport. Available at: http://www.naturstyrelsen.dk/NR/rdonlyres/CBCB6E2F-D00E-418F-805C-3F6E54BA90DD/0/Punktkilderrapport_.pdf (26/02-2014).

Agency for Spatial and Environmental Planning (ASEP), 2010: Point sources 2008. In Danish: Punktkilder 2008. Det nationale program for overvågning af vandmiljøet; Fagdatacenterrapport. Available at: <http://www.naturstyrelsen.dk/NR/rdonlyres/40130016-7F05-47F9-BF2E-A9756D83EDE6/0/PunktkildeRapport200815022010.pdf> (26/02-2014).

Agency for Spatial and Environmental Planning (ASEP), 2010: Point sources 2009. In Danish: Punktkilder 2009. Det nationale program for overvågning af vandmiljøet; Fagdatacenterrapport. Available at: <http://www.naturstyrelsen.dk/NR/rdonlyres/10B91A8A-3937-4AB8-A246-12F0DB5C2126/119653/punktkilder20091.pdf> (26/02-2014).

Agency for Spatial and Environmental Planning (ASEP), 2011: Point sources 2010. In Danish: Punktkilder 2010. Det nationale program for overvågning af vandmiljøet; Fagdatacenterrapport. Available at: <http://www.naturstyrelsen.dk/Udgivelser/Aarstal/2011/Punktkilderapport2010.htm> (26/02-2014).

Agency for Spatial and Environmental Planning (ASEP), 2012: Point sources 2011. In Danish: Punktkilder 2011. Det nationale program for overvågning af vandmiljøet; Fagdatacenterrapport. Available at: http://www.naturstyrelsen.dk/Udgivelser/Aarstal/2012/punktkilder_2011.htm (26/02-2014).

Agency for Spatial and Environmental Planning (ASEP), 2013: Point sources 2012. In Danish: Punktkilder 2012. Det nationale program for overvågning af vandmiljøet; Fagdatacenterrapport. Available at: <http://www.naturstyrelsen.dk/NR/rdonlyres/B76CDB71-702A-440C-87FF-7837E9E84AFB/0/Punktkilderrapport2012.pdf> (26/02-2014).

Amlinger, F., Peyr, S. & Cuhls, C. 2008: Greenhouse gas emissions from composting and mechanical biological treatment. Florian Amlinger, Stefan Peyr and Carsten Cuhls. Waste Management & Research 2008; 26; 47. Sage publications, International Solid Waste Association. Available at: <http://wmr.sagepub.com/cgi/reprint/26/1/47.pdf> (05/02-2014).

Blomqvist, P., Persson, B. & Simonson, M. 2002: Utsläpp från bränder till miljön, Utsläpp av dioxin, PAH och VOC till luften, Räddnindsverket, Karlstad, FoU rapport, SP Brandteknik, Borås. Available at: <http://rib.msb.se/Filer/pdf%5C18193.pdf> (05/02-2014) (Swedish).

Boldrin, A., Andersen, J.K. & Christensen, T.H. 2009: Environmental assessment of garden waste management in Århus Kommune (Miljøvurdering af haveaffald i Århus kommune), Technical University of Denmark - Miljø, 3R. Available at: <http://www.aarhus.dk/~media/Subsites/AVA/Om-AVA/Bibliotek/Rapporter-og-undersogelser/Aarhus-haveaffald-engelsk-rapport.pdf> (05/02-2014)

Czepiel, P., Crill, P. & Harriss, R. 1995: Nitrous oxide emissions from municipal wastewater treatment, Environmental Science and Technology, 29, 2352-2356.

Danish Emergency Management Agency (DEMA), 2007: Beredskabsstyrelsen, Redningsberedskabets Statistiske Beretning. Available at: http://brs.dk/viden/publikationer/Documents/RBS_Statistisk_Beretning_2007.pdf (05/02-2014) (Danish).

Danish Emergency Management Agency (DEMA), 2013: Beredskabsstyrelsen, statistikbanken. Available at: <https://statistikbank.brs.dk/sb/main/p/a0109> (05/02-2014) (Danish).

Danish Energy Agency (DEA), 2013a: The Danish energy statistics, Available at: <http://www.ens.dk/en/info/facts-figures/energy-statistics-indicators-energy-efficiency/annual-energy-statistics> (2014-02-14)

Danish Energy Agency (DEA), 2013b: The Danish energy statistics, Energiproducenttællingen 1990-2012. Unpublished.

Danish Environmental Protection Agency (DEPA), 1989: Wastewater from municipal and private wastewater treatment plants in 1987. In Danish: Spil-

devandsslam fra kommunale og private renseanlæg i 1987, Orientering fra Miljøstyrelsen, nr. 10. Available at:

<http://www2.mst.dk/Udgiv/publikationer/1989/87-503-7944-5/pdf/87-503-7944-5.pdf> (28/02-2014).

Danish Environmental Protection Agency (DEPA), 1993: Affaldsstrømmene i Danmark, Arbejdsrapport fra Miljøstyrelsen Nr. 15 1993. Available at:

<http://www.mst.dk/Publikationer/Publikationer/1993/01/87-7909-716-2.htm> (28/02-2014).

Danish Environmental Protection Agency (DEPA), 1994: Point Sources 1993. In Danish: Punktkilder 1993, Orientering fra Miljøstyrelsen, nr. 8. Available at:

<http://www2.mst.dk/Udgiv/publikationer/1994/87-7810-279-0/pdf/87-7810-279-0.pdf> (28/02-2014).

Danish Environmental Protection Agency (DEPA), 1996a: Affaldsstatistik 1995 (Waste Statistics 1995. Orientering fra Miljøstyrelsen nr 14 1997) Available at:

<http://www2.mst.dk/Udgiv/publikationer/1996/87-7810-704-0/pdf/87-7810-704-0.pdf> (28/02-2014).

Danish Environmental Protection Agency (DEPA), 1996b: Point Sources 1995. In Danish: Punktkilder 1995, Orientering fra Miljøstyrelsen, nr. 16. Available at:

<http://www2.mst.dk/Udgiv/publikationer/1996/87-7810-709-1/pdf/87-7810-709-1.pdf> (28/02-2014).

Danish Environmental Protection Agency (DEPA), 1997: Point Sources 1996. In Danish: Punktkilder 1996, Orientering fra Miljøstyrelsen, nr. 16. Available at:

<http://www2.mst.dk/common/Udgivramme/Frame.asp?http://www2.mst.dk/Udgiv/publikationer/1998/87-7909-162-8/html/default.htm> (28/02-2014).

Danish Environmental Protection Agency (DEPA), 1998a: Waste Statistics 1996. Available at:

http://www2.mst.dk/common/Udgivramme/Frame.asp?pg=http://www2.mst.dk/Udgiv/publications/1998/87-7909-106-7/html/default_eng.htm (28/02-2014).

Danish Environmental Protection Agency (DEPA), 1998b: Point Sources 1997. In Danish: Punktkilder 1997, Orientering fra Miljøstyrelsen, nr. 9. Available at:

<http://www2.mst.dk/common/Udgivramme/Frame.asp?http://www2.mst.dk/Udgiv/publikationer/1998/87-7909-162-8/html/default.htm> (28/02-2014).

Danish Environmental Protection Agency (DEPA), 1999a: Waste Statistics 1997. Available at:

http://www2.mst.dk/common/Udgivramme/Frame.asp?pg=http://www2.mst.dk/Udgiv/publications/2000/87-7909-433-3/html/default_eng.htm (28/02-2014).

Danish Environmental Protection Agency (DEPA), 1999b: Point Sources 1998. In Danish: Punktkilder 1998, Orientering fra Miljøstyrelsen, nr. 6. Available at:

<http://www2.mst.dk/Udgiv/publikationer/1999/87-7909-554-2/pdf/87-7909-554-2.pdf> (28/02-2014).

Danish Environmental Protection Agency (DEPA), 1999c: Wastewater from municipal and private wastewater treatment plants in 1997. In Danish: Spildevandsslam fra kommunale og private renseanlæg i 1997, Miljøprojekt, nr. 473. Available at:

<http://www2.mst.dk/Udgiv/publikationer/1999/87-7909-330-2/pdf/87-7909-330-2.pdf> (28/02-2014).

Danish Environmental Protection Agency (DEPA), 2000: Point Sources 1999. In Danish: Punktkilder 1999, Orientering fra Miljøstyrelsen, nr. 16. Available at:

<http://www2.mst.dk/Udgiv/publikationer/2000/87-7944-298-6/pdf/87-7944-299-4.pdf> (28/02-2014).

Danish Environmental Protection Agency (DEPA), 2001a: Waste Statistics 1998. Available at:

http://www2.mst.dk/common/Udgivramme/Frame.asp?pg=http://www2.mst.dk/Udgiv/publications/2001/87-7944-351-6/html/default_eng.htm (28/02-2014).

Danish Environmental Protection Agency (DEPA), 2001b: Waste Statistics 1999. Available at:

http://www2.mst.dk/common/Udgivramme/Frame.asp?pg=http://www2.mst.dk/Udgiv/publications/2001/87-7944-381-8/html/default_eng.htm (28/02-2014).

Danish Environmental Protection Agency (DEPA), 2001c: Point Sources 2000. In Danish: Punktkilder 2000, Orientering fra Miljøstyrelsen, nr. 13. Available at: <http://www2.mst.dk/Udgiv/publikationer/2001/87-7944-943-3/pdf/87-7944-944-1.pdf> (28/02-2014).

Danish Environmental Protection Agency (DEPA), 2001d: Wastewater from municipal and private wastewater treatment plants in 1999. In Danish: Spildevandsslam fra kommunale og private renseanlæg i 1999, Orientering fra Miljøstyrelsen, nr. 3. Available at:

<http://www2.mst.dk/Udgiv/publikationer/2001/87-7944-356-7/pdf/87-7944-357-5.pdf> (28/02-2014).

Danish Environmental Protection Agency (DEPA), 2002a: Waste Statistics 2000. Available at:

http://www2.mst.dk/common/Udgivramme/Frame.asp?pg=http://www2.mst.dk/Udgiv/publications/2002/87-7972-027-7/html/default_eng.htm (28/02-2014).

Danish Environmental Protection Agency (DEPA), 2002b: Point Sources 2001. In Danish: Punktkilder 2001, Orientering fra Miljøstyrelsen, nr. 7. Available at: <http://www2.mst.dk/Udgiv/publikationer/2002/87-7972-384-5/pdf/87-7972-385-3.pdf> (28/02-2014).

Danish Environmental Protection Agency (DEPA), 2003a: Hvad koster det at reducere CO₂-mankoen? Reduktionspotentiale og omkostninger i udvalgte sektorer, Arbejdsrapport fra Miljøstyrelsen Nr. 7 2003. Available at:

<http://www2.mst.dk/udgiv/publikationer/2003/87-7972-464-7/pdf/87-7972-465-5.pdf> (28/02-2014).

Danish Environmental Protection Agency (DEPA), 2003b: Point Sources 2002. In Danish: Punktkilder 2002, Orientering fra Miljøstyrelsen, nr. 10. Available at:
<http://www2.mst.dk/Udgiv/publikationer/2003/87-7614-074-1/pdf/87-7614-076-8.pdf> (28/02-2014).

Danish Environmental Protection Agency (DEPA), 2003c: Wastewater from municipal and private wastewater treatment plants in 2000 and 2001. In Danish: Spildevandsslam fra kommunale og private renselanlæg i 2000 og 2001, Orientering fra Miljøstyrelsen, nr. 9. Available at:
<http://www2.mst.dk/Udgiv/publikationer/2003/87-7614-005-9/pdf/87-7614-006-7.pdf> (28/02-2014).

Danish Environmental Protection Agency (DEPA), 2004a: Waste Statistics 2001. Available at:
http://www2.mst.dk/common/Udgivramme/Frame.asp?pg=http://www2.mst.dk/Udgiv/publications/2004/87-7614-105-5/html/default_eng.htm (28/02-2014).

Danish Environmental Protection Agency (DEPA), 2004b: Waste Statistics 2002. Available at:
http://www2.mst.dk/common/Udgivramme/Frame.asp?pg=http://www2.mst.dk/Udgiv/publications/2004/87-7614-107-1/html/default_eng.htm (28/02-2014).

Danish Environmental Protection Agency (DEPA), 2004c: Point Sources 2003. In Danish: Punktkilder 2003, Orientering fra Miljøstyrelsen, nr. 16. Available at:
<http://www2.mst.dk/Udgiv/publikationer/2004/87-7614-482-8/pdf/87-7614-483-6.pdf> (28/02-2014).

Danish Environmental Protection Agency (DEPA), 2004d: Wastewater from municipal and private wastewater treatment plants in 2002. In Danish: Spildevandsslam fra kommunale og private renselanlæg i 2002, Orientering fra Miljøstyrelsen, nr. 5. Available at:
<http://www2.mst.dk/Udgiv/publikationer/2004/87-7614-195-0/pdf/87-7614-196-9.pdf> (28/02-2014).

Danish Environmental Protection Agency (DEPA), 2005a: Waste Statistics 2003. Available at:
http://www2.mst.dk/common/Udgivramme/Frame.asp?pg=http://www2.mst.dk/Udgiv/publications/2005/87-7614-585-9/html/default_eng.htm (28/02-2014).

Danish Environmental Protection Agency (DEPA), 2005b: Point Sources 2004. In Danish: Punktkilder 2004, Orientering fra Miljøstyrelsen, nr. 9. Available at:
<http://www2.mst.dk/Udgiv/publikationer/2005/87-7614-865-3/pdf/87-7614-866-1.pdf> (28/02-2014).

Danish Environmental Protection Agency (DEPA), 2005c: Point Sources 2003. Revision. In Danish: Punktkilder 2003 – revideret udgave, Orientering fra Miljøstyrelsen, nr. 1. Available at:
<http://www2.mst.dk/Udgiv/publikationer/2005/87-7614-512-3/pdf/87-7614-513-1.pdf> (28/02-2014).

Danish Environmental Protection Agency (DEPA), 2006a: Waste Statistics 2004. Available at:

http://www2.mst.dk/common/Udgivramme/Frame.asp?pg=http://www2.mst.dk/Udgiv/publications/2006/87-7614-962-5/html/default_eng.htm
(28/02-2014)

Danish Environmental Protection Agency (DEPA), 2006b: Affaldsstatistik 2005 (Waste Statistics 2005). Orientering fra Miljøstyrelsen nr. 6. Available at:

<http://www2.mst.dk/common/Udgivramme/Frame.asp?pg=http://www2.mst.dk/Udgiv/publikationer/2006/87-7052-284-7/html/default.htm>
(28/02-2014) (Danish).

Danish Environmental Protection Agency (DEPA), 2006c: Legislation on sludge. Bekendtgørelse om anvendelse af affald til jordbrugsformål (Slambekendtgørelsen), BEK nr. 1650 af 13/12/2006. Available at:

<https://www.retsinformation.dk/Forms/R0710.aspx?id=13056> (05/02-2014) (Danish).

Danish Environmental Protection Agency (DEPA), 2008: Affaldsstatistik 2006. (Waste Statistics 2006) Orientering fra Miljøstyrelsen nr 2 2008. Available at:

<http://www2.mst.dk/udgiv/publikationer/2008/978-87-7052-753-8/pdf/978-87-7052-754-5.pdf> (28/02-2014) (Danish).

Danish Environmental Protection Agency (DEPA), 2009: Wastewater from municipal and private wastewater treatment plants in 2005. In Danish: Spildevandsslam fra kommunale og private renseanlæg i 2005, Orientering fra Miljøstyrelsen, nr. 3. Available at:

<http://www2.mst.dk/udgiv/publikationer/2009/978-87-7052-993-8/pdf/978-87-7052-994-5.pdf> (28-02-2014).

Danish Environmental Protection Agency (DEPA), 2010a: Affaldsstatistik 2007 og 2008. (Waste Statistics 2007 and 2008). Available at:

<http://www.mst.dk/NR/rdonlyres/204B82C6-C3BE-43A4-824A-C30EED54CF9F/0/Affaldsstatistik2007og2008.pdf> (28/02-2014) (Danish).

Danish Environmental Protection Agency (DEPA), 2010b: Anvendelse af affald til jordbrugsformål. Vejledning fra Miljøstyrelsen Nr. 1 2010. Available at:

http://www.landbrugsinfo.dk/Regelinfo/Kommenteringer/Filer/pl_ri_11_054_b1.pdf (28/02-2014).

Danish Environmental Protection Agency (DEPA), 2010c: Deponeringskapacitet i Danmark - i perioderne 2009-12 og 2013-20 (English title: Depositing capacity in Denmark - in periods 2009-12 and 2013-20). Environment project no.1318. Available at:

<http://www2.mst.dk/udgiv/publikationer/2010/978-87-92617-55-2/pdf/978-87-92617-56-9.pdf> (28/02-2014).

Danish Environmental Protection Agency (DEPA), 2011a: Affaldsstatistik 2009 og fremskrivning af affaldsmængder 2011-2050. (Waste Statistics 2009). Available at:

<http://www2.mst.dk/udgiv/publikationer/2011/10/978-87-92779-44-1/978-87-92779-44-1.pdf> (28-02-2014) (Danish).

Danish Environmental Protection Agency (DEPA), 2011b: Legislation on waste (Affaldsbekendtgørelsen), BEK nr. 1415 af 12/12/2011 Gældende, Chapter 7, §46 and §47. Available at: <https://www.retsinformation.dk/forms/r0710.aspx?id=139654#Kap7> (In Danish) (05/02-2014).

Danish Gas Technology Center (DGC), 2009: DGC målerapport 731-28-5 November 2009, Gasmotoranlæg, Jenbacher J 316, deponigas, Steen D. Andersen. Available at: [http://www.energinet.dk/SiteCollectionDocuments/Danske%20dokumenter/Klimaogmiljo/Emissionsprojektet%20-%20Anl%C3%A6g%2005%20-%20Motor%20\(biogas%20deponigas\).pdf](http://www.energinet.dk/SiteCollectionDocuments/Danske%20dokumenter/Klimaogmiljo/Emissionsprojektet%20-%20Anl%C3%A6g%2005%20-%20Motor%20(biogas%20deponigas).pdf) (05/02-2014).

Danish Ministry of Environment, 2010. Waste strategy '10 (Waste Strategy 2009-2012, Part II) (<http://www.mst.dk/NR/rdonlyres/5944D6BF-C455-4A94-A7D1-9062ED0173A0/0/Affaldsstrategi10.pdf>, (23/02-2014)

Danske Krematoriernes Landsforening (DKL), 2013: Association of Danish crematories, statistisk materiale. Available at: <http://dkl.dk/Info/statistik.php> (05/02-2014) (Danish).

DANVA, 2008. Eksplosiv atmosfære i kloakker (English title: Explosive atmospheres in sewers). Report no. 77. Available at: [file:///C:/Users/mth/Downloads/Eksplosiv atmosf%C3%A6re i kloakke_r_samlet_rapport%20\(2\).pdf](file:///C:/Users/mth/Downloads/Eksplosiv%20atmosf%C3%A6re%20i%20kloakke%20samlet%20rapport%20(2).pdf) (26/02-2014) (Danish).

European Environment Agency (EEA), 2009: The EMEP/EEA air pollutant emission inventory guidebook, Technical report No 9/2009. Available at: <http://www.eea.europa.eu/publications/emep-eea-emission-inventory-guidebook-2009> (05-02-2014).

Fausser, P., Thomsen, M., Nielsen, O., Winther, M., Gyldenkerne, S., Hoffmann, L., Lyck, E. & Illerup, J.B. 2007: Verification of the Danish emission inventory data by national and international data comparisons. Danmarks Miljøundersøgelser, Aarhus Universitet. 53 pp - NERI Technical Report No. 627. Available at: http://www2.dmu.dk/Pub/FR627_Final.pdf (01/03-2014).

Fausser, P., Nielsen, M., Winther, M., Plejdrup, M. S., Gyldenkerne, S., Mikkelsen, M. H., Albrektsen, R., Hoffmann, L., Thomsen, M., Hjelgaard, K. & Nielsen, O-K., 2013. Verification of the Danish GHG Key Categories and Annex II Indicators. Greenhouse Gas Measurement & Management 3(3-4), 107-127.

Fausser, P., Sørensen, P. B., Nielsen, M., Winther, M., Plejdrup, M. S., Hoffmann, L., Gyldenkerne, S., Mikkelsen, M. H., Albrektsen, R., Lyck, E., Thomsen, M., Hjelgaard, K. & Nielsen, O-K., 2011. Monte Carlo (Tier 2) Uncertainty Analysis of Danish Greenhouse Gas Emission Inventory. Greenhouse Gas Measurement & Management, 1(3/4), 145-161.

Fontelle, J.-P., Allemand, N., Andre, J.-M., Beguier, S., Chang, J.-P., Gueguen, C., Jacquier, G., Martinet, Y., Mathias, E., Nguyen, V., Oudart, B., Serveau, L. & Vincent, J., 2008: Organisation et méthodes des inventaires nationaux des émissions atmosphériques en France, 5^{ème} édition.

Hjelgaard, 2013: Hjelgaard, K. 2013. Danish Emission Inventory for Waste Incineration and Other Waste. Inventories until year 2011. Aarhus University, DCE – Danish Centre for Environment and Energy, 96 pp. Scientific Report from DCE – Danish Centre for Environment and Energy No. 70. Available at: <http://www.dmu.dk/Pub/SR70.pdf> (05/02-2014).

Hvitved-Jacobsen, T., 2001. Sewer Processes – Microbial and Chemical Process Engineering of Sewer Networks, CRC Press.

Intergovernmental Panel on Climate Change (IPCC), 1997: Revised 1996 IPCC Guidelines for National Greenhouse Gas Inventories. Available at: <http://www.ipcc-nggip.iges.or.jp/public/gl/invs1.html> (05/02-2014).

Intergovernmental Panel on Climate Change (IPCC), 2000: Good Practice Guidance and Uncertainty Management in National Greenhouse Gas Inventories. Available at: <http://www.ipcc-nggip.iges.or.jp/public/gp/english/index.html> (05/02-2014).

Intergovernmental Panel on Climate Change (IPCC), 2006: 2006 IPCC Guidelines for National Greenhouse Gas Inventories. Available at: <http://www.ipcc-nggip.iges.or.jp/public/2006gl/index.html> (05/02-2014).

Inter-ministerial group on organizational change and regulation of waste incineration and landfill area, 2007. Deponering af affald – baggrundsrapport (English title: Landfilling of waste - Background Report). Available at: (<http://www.mst.dk/NR/rdonlyres/A340FD39-C091-482A-9A94-DB21627A8C1C/0/Baggrundsrapportomdeponering.pdf>) (28/02-2014).

ISAG, 2011: The Danish Environmental Protection Agency, database on all registered Danish waste. Available at: <http://www2.mst.dk/databaser/isag/Default.asp?advanced=Yes> (28/02-2014) (Danish).

Kampschreur, M.J., Temmink, H., Kleerebezem, R., Jetten, M.S.M. & van Loosdrecht, M.C.M., 2009: Nitrous oxide emission during wastewater treatment. Water Research 43, 4093-4103.

Kristensen, G.H., Jørgensen, P.E., 2008. Renseteknik – oversigt over ny miljøteknologi indenfor spildevandshåndtering (English title: Wastewater treatment Technologies - Overview of new environmental technology in waste water management). ASEP report. Available at: <http://www2.blst.dk/udgiv/Publikationer/2008/978-87-92256-39-3/pdf/978-87-92256-40-9.pdf> (26/02-2014)

Limpert, E., Stahel, W.A. & Abbt, M. 2001: Log-normal Distributions across the Sciences: Keys and Clues. BioScience Vol. 51 No. 5.

Lyngby-Taarbæk commune, 2013. Midlertidig tilslutningstilladelse til udledning af byggegrubevand fra bygge- og anlægsarbejdet på Kanalvej 7 (English title: Temporary connection permit for the discharge of water in relation to the construction work on Kanalvej 7). Available at: http://www.ltk.dk/sites/default/files/uploads/public/user_uploads/CM P/Hoeringer/tilslutningstilladelse.pdf (26/02-2014).

Lönnermark, A. & Blomqvist, P., 2006: Emissions from an automobile fire. *Chemosphere* 62 (2006) 1043–1056. Available at: <http://www.sciencedirect.com/science/article/pii/S0045653505006351> (05/02-2014).

MILIKI, 2006: Ministeriet for Ligestilling og Kirke (Ministry for Gender Equality and Ecclesiastical Affairs), Arbejdsgruppe foreslår, at der skal være færre krematorier, december 2006. Available at: http://miliki.dk/fileadmin/share/publikationer/Krematorier/Rapport_om_krematorier.pdf (05/02-2014) (Danish).

MST, 2013: Livscyklusvurdering og samfundsøkonomisk analyse for anvendelse af spildevandsslam, Miljøministeriet, Miljøstyrelsen (MST), Miljøprojekt nr. 1459, 2013, ISBN nr. 978-87-92903-81-5. Available at: <http://www.mst.dk/Publikationer/Publikationer/2013/Maj/978-87-92903-81-5.htm> (28/02-2014) (Danish).

National Atmospheric Emission Inventory of UK (NAEI), 2009. Available at: <http://www.naei.org.uk/emissions/selection.php> (05/02-2014).

Nielsen, O.-K., Winther, M., Nielsen, M., Mikkelsen, M.H., Albrektsen, R., Gyldenkerne, S., Plejdrup, M.S., Hoffmann, L., Thomsen, M., Hjelgaard, K. & Fauser, P. 2011b: Projection of Greenhouse Gas Emissions 2010 to 2030. National Environmental Research Institute, Aarhus University. 176 s. (NERI Technical Report; 841). Available at: <http://www.dmu.dk/Pub/FR841.pdf> (28/02-2012).

Nielsen, O.-K., Plejdrup, M.S., Winther, M., Nielsen, M., Gyldenkerne, S., Mikkelsen, M.H., Albrektsen, R., Thomsen, M., Hjelgaard, K., Hoffmann, L., Fauser, P., Bruun, H.G., Johannsen, V.K., Nord-Larsen, T., Vesterdal, L., Møller, I.S., Caspersen, O.H., Rasmussen, E., Petersen, S.B., Baunbæk, L. & Hansen, M.G. 2013. Denmark's National Inventory Report 2013. Emission Inventories 1990-2011 - Submitted under the United Nations Framework Convention on Climate Change and the Kyoto Protocol. Aarhus University, DCE - Danish Centre for Environment and Energy, 1202pp. Scientific Report from DCE - Danish Centre for Environment and Energy. Available at: <http://www.dmu.dk/Pub/SR56.pdf> (05/02-2014).

Persson, B. & Simonson, M. 1998: Fire emissions into the atmosphere, Swedish National Testing and Research Institute, Birås, Sweden, Fire Technology, Vol. 34, No. 3, 1998. Available at: http://download.springer.com/static/pdf/16/art%253A10.1023%252FA%253A1015350024118.pdf?auth66=1391779932_7fe1a282ee6536d6cc46aefb69742e23&ext=.pdf (05/02-2014).

Petersen, C. 2001: Statistik for behandling af organisk affald fra husholdninger - Revideret udgave (Statistics for treatment of organic waste from households - revised version). Miljøprojekt Nr. 654 2001, Miljøstyrelsen, Miljøministeriet. Available at: <http://www2.mst.dk/Udgiv/publikationer/2001/87-7944-932-8/pdf/87-7944-933-6.pdf> (05/02-2014) (Danish).

Petersen, C. & Hansen, V.L. 2003: Statistik for behandling af organisk affald fra husholdninger 2001 (Statistics for treatment of organic waste from households). Miljøprojekt Nr. 856 2003, Miljøstyrelsen, Miljøministeriet.

Available at:

<http://www2.mst.dk/Udgiv/publikationer/2003/87-7972-962-2/pdf/87-7972-963-0.pdf> (05/02-2014) (Danish).

Petersen, C. & Kielland, M. 2003: Statistik for hjemmekompostering (Statistics on home composting), Econet A/S. Miljøprojekt Nr. 855 2003, Miljøstyrelsen, Miljøministeriet. Available at:

<http://www2.mst.dk/udgiv/publikationer/2003/87-7972-960-6/pdf/87-7972-961-4.pdf> (05/02-2014) (Danish).

Schleicher, O. & Gram, L.K. 2008: Miljøprojekt Nr. 1191, Analyse af omkostningerne for rensning for kviksølv på krematorier i Danmark, Miljøministeriet, Miljøstyrelsen. Available at:

<http://www2.mst.dk/common/Udgivramme/Frame.asp?http://www2.mst.dk/udgiv/publikationer/2008/978-87-7052-594-7/html/helepubl.htm> (05/02-2014) (Danish).

Schleicher, O., Jensen, A.A. & Blinksbjerg, P. 2001: Miljøprojekt Nr. 649, Måling af dioxinemissionen fra udvalgte sekundære kilder, Miljøministeriet, Miljøstyrelsen. Available at:

<http://www2.mst.dk/Udgiv/publikationer/2001/87-7944-868-2/pdf/87-7944-869-0.pdf> (05/02-2014) (Danish).

Statistics Denmark, 2013: StatBank Denmark. Available at:

<http://www.statistikbanken.dk/statbank5a/default.asp?w=1024> (05/02-2014) (Danish/English).

Statutory order for Landfilling, no. 719, 24/06/2011 (Deponeringsbekendtgørelsen) (<https://www.retsinformation.dk/forms/R0710.aspx?id=137791>) (28/02-2014).

Statutory order for Waste, no 1309 af 18/12/2012 (Affaldsbekendtgørelsen). Available at:

<https://www.retsinformation.dk/Forms/R0710.aspx?id=144826> (05/02-2014).

The Danish Government, 1999: Waste 21. Waste Management plan 1998-2004. Available at

<http://www2.mst.dk/Udgiv/publications/1999/87-7909-571-2/pdf/87-7909-570-4.pdf> (04-02-2014).

The Danish Government, 2003: Affaldsstrategi 2005-2008. (In Danish: Waste Strategy 2005-2008. Available at:

<http://www2.mst.dk/Udgiv/publikationer/2003/87-7972-971-1/pdf/87-7972-973-8.pdf> (04-02-2014).

The Danish Government, 2009: Affaldsstrategi 2009-2012 1. delstrategi. (In Danish: Waste Strategy 2009-2012. Available at:

<http://www.mst.dk/NR/rdonlyres/747FBCE2-A3D4-444F-BF60-D1747C36516D/0/Endelig1delafAffaldsstrategi200912.pdf> (04-02-2014).

The Danish Government, 2013. Denmark without waste. Recycle more – incinerate less. http://www.mim.dk/NR/rdonlyres/F58E8C8B-3EB9-47DF-8DF3-4BF9207C9DFE/0/Ressourcestrategi_UK_web.pdf

Thomsen, M. & Lyck, E. 2005: Emission of CH₄ and N₂O from wastewater treatment plants (6B). NERI Research Note No. 208. Available at: http://www2.dmu.dk/1_viden/2_Publikationer/3_arbrapporter/rapporter/AR208.pdf (04-02-2014).

Vattenfall, 2010: Fynsværket CO₂-udledningsrapport 2010.

Verdo, 2011: Verdo (Randers) CO₂-kvote indberetning 2011.

Vollertsen, J., Vorkamp, K., Hvitved-Jacobsen, T. & Jensen, N.A., 2002: Udsivning af spildevand fra afløbssystemer. Miljøprojekt Nr. 685.

Wicht, H. & Beier, M. 1995: N₂O emission aus nitrifizierenden und denitrifizierenden Kläranlagen. Korrespondenz Abwasser 42, 404–406.

9 Other (CRF sector 7)

In CRF Sector 7, there are no activities and emissions for the inventories of Denmark. Until the 2009 submission in the inventories of the Kingdom of Denmark (Denmark, Faroe Islands and Greenland) emissions from Faroe Islands and Greenland were reported in Sector 7. This has been changed so that Greenland and Faroe Islands are included in full CRF's.

For further detail on the emissions from Greenland and the Faroe Islands please see Chapter 16 and Annex 9.

10 Recalculations and improvements

Previously the recalculation tables in the CRF have been incorrect due to the different geographical scopes of the inventory and technical limitations in the CRF Reporter software. However, by running five different installations of the CRF Reporter software, the data presented in Table 8 of the CRF are accurate. Explanations for the recalculations of the Danish inventory are included in Chapter 10.1.1.

The overall impact of recalculations is shown in Table 10.1. A more detailed overview is provided in Tables 10.2 – 10.5.

Information on recalculations for the aggregated submission of Denmark and Greenland under the Kyoto Protocol are included in Chapter 17.

10.1 Explanations and justifications for recalculations

Explanations and justifications for the recalculations performed in this submission, since submission of data to the UNFCCC due April 15, 2013 for Denmark, are given in the following sector chapters:

Energy:

- Stationary Combustion Chapter 3.2.8
- Transport Chapter 3.3.7
- Fugitive emissions Chapter 3.5.8

Industry:

- Mineral products Chapter 4.2.5
- Chemical industry Chapter 4.3.4
- Metal production Chapter 4.4.4
- Food and drink Chapter 4.5.4
- Consumption of f-gases Chapter 4.7

Solvents and Other Product Use Chapter 5.6

Agriculture Chapter 6.10

LULUCF

- Forest Land Chapter 7.2.1, 7.2.2
- Cropland Chapter 7.3
- Grassland Chapter 7.4
- Wetlands Chapter 7.5
- Settlements Chapter 7.6

Waste

- Solid Waste Disposal on Land Chapter 8.8.1
- Wastewater Chapter 8.8.2
- Waste incineration Chapter 8.8.3
- Waste, Other Chapter 8.8.4

KP-LULUCF

- ARD Chapter 11.3.5

- FM Chapter 11.4.5
- CM Chapter 11.5.5
- GM Chapter 11.6.4

The main recalculations since the 2013 submission are:

10.1.1 Energy

Stationary Combustion

For stationary combustion plants, the emission estimates for the years 1990-2011 have been updated according to the latest energy statistics published by the Danish Energy Agency. The update included both end use and transformation sectors as well as a source category update. The changes in the energy statistics are largest for the years 2009, 2010 and 2011.

For CO₂ the largest recalculation is in source category Manufacturing industries and constructions. The recalculation is related to liquid fuels and is a result of correction of an error. The consumption of residual oil was underestimated in the former inventories. The CO₂ emission from liquid fuels applied in manufacturing industries and construction for 2011 is 7% higher in the 2014 reporting than in the 2013 reporting.

The CH₄ emission from residential wood combustion has been recalculated based on improved emission factors for stoves. This has caused a 16 % increase of the CH₄ emission reported for biomass fuels in residential plants for 2011.

For N₂O the largest recalculation is in source category Manufacturing industries and constructions. This recalculation is also related to the former underestimate for residual oil. The N₂O emission from liquid fuels applied in manufacturing industries and constructions for 2011 is 13 % higher in the 2014 reporting than in the 2013 reporting.

Mobile sources

The following recalculations and improvements of the emission inventories have been made since the emission reporting in 2013.

Road transport

Based on the updated version of COPERT IV launched in 2013, new vehicle sub categories have been introduced in the emission inventories for mopeds and passenger cars. For mopeds a division is now made between 2-stroke and 4-stroke engine technologies and for passenger cars small engine sizes below 0.8 l. for gasoline and below 1.4 l. for diesel have been included. Also NO_x emission factors for euro 5 diesel passenger cars have been updated in the model based on the new COPERT IV version.

Small errors in input gasoline fuel consumption for the years 2009-2011 and for input diesel fuel consumption in the years 2010-2011 have been corrected.

The percentage emission change interval and year of largest percentage differences (low %; high %, year) for the different emission components are: CO₂ (-0.5 %; -0.05 %, 2008), CH₄ (-0.2 %; 2.4 %, 2011) and N₂O (-0.4 %; 0.3 %, 2008).

Navigation

Minor changes in ferry input data has been made for the years 2008-2011 causing minor emission changes for domestic navigation. The following largest percentage differences (in brackets) for domestic navigation are noted for: CO₂ (-1.0 %), CH₄ (-0.8 %) and N₂O (-1.1 %).

Agriculture/forestry/fisheries

The number and engine size of machine pool tractors has been updated for the years 2007-2011. The number of ATV's has been changed for the years 2009-2011. Errors in the fuel consumption for fisheries in 2000, 2010 and 2011 have been corrected.

In 2000 the following percentage differences (in brackets) for agriculture/forestry/fisheries are noted for: CO₂ (9.1 %), CH₄ (3.7 %) and N₂O (11.9 %) due to fuel consumption changes in fisheries.

For other years than 2000, the following largest percentage differences (in brackets) are noted for: CO₂ (-3.7 %), CH₄ (5 %) and N₂O (-4.9 %).

Industry

The number of mini loaders has been updated for the years 2004-2011.

The following largest percentage differences (in brackets) for industrial non road machinery are noted for: CO₂ (1.0 %), CH₄ (1.6 %) and N₂O (1.6 %).

Civil aviation

An error in the CH₄ emission factor has been corrected for the years 1985-2000. The emission factors are now in line with the factors proposed by the EMEP/EEA emission inventory guidebook. The CH₄ emission percentage differences are between -31 % and -42 %.

Military

Emission factors derived from the new road transport simulations have caused some emission changes from 1985-2010. The following largest percentage differences (in brackets) for military are noted for: CO₂ (0 %), CH₄ (0.6 %) and N₂O (0.2 %).

Fugitive emissions

In the emission inventory reported in 2014 for the years 1990-2012 the following recalculations regarding fugitive emissions from fuels have been applied:

Natural gas transmission and distribution

Activity data and IEF for the time series 1990-2011 has been updated for transmission and distribution according to annual environmental reports and the latest national energy statistics, respectively. Further, the CH₄ EFs are updated for one town gas distribution company following an update of the estimated fugitive losses per distribution. The recalculation has changed the CO₂ emission by 0.01 ktonnes and CH₄ emission by (-0.09) - 0.07 ktonnes, corresponding to < 0.003 % and (-2) % - 2% of the total fugitive CO₂ and CH₄ emission.

Venting

EFs for CH₄ have been added for the years 1990-1993 for one gas storage plant. In these years the plant is treated as an area source in the national system, while it is treated separately as a point source in the following years.

EFs are based on data from annual reports for 1995-1999, as no data are available for the years 1990-1994. Further, a minor error has been applied for venting in 2011, according to the annual report from one of the natural gas storage facilities. The recalculation has changed the CH₄ emission by 0.06 ktonnes, corresponding to 2% of the total fugitive CH₄ emission.

Flaring in gas storage and treatment plants

EFs for CO₂ and CH₄ have been added for the years 1990-1993 for the gas treatment plant. In these years the plant is treated as an area source in the national system, while it is treated separately as a LPS in the following years. EFs are based on data from annual reports for 1995-1999, as no data are available for the years 1990-1994. The recalculation has changed the CO₂ emission by 2.2 ktonnes and the CH₄ emission by 0.01 ktonnes, corresponding to 0.4 % and 0.4 % of the total fugitive CO₂ and CH₄ emission.

Flaring in refineries

The CO₂ EF for one refinery has been updated for the years 1994-2006 and now reflects the average EF from the first five EU-ETS reports (2007-2011). The NO_x EF for two refineries has been changed to the standard EF in the EMEP/EEA Guidebook, as references to the previously applied EFs are outdated or not existing.

The CO₂ EF for a refinery that was shut down in 1996 has been changed for the years 1990-1996 to match the existing refineries, as no better data is available. CO₂ EF for 2010-2011 has been updated and now corresponds to the EU-ETS reporting. The recalculation has changed the CO₂ emission by (-0.05) - 0.49 ktonnes and CH₄ emission by (-0.06) ktonnes, corresponding to (-0.01) - 0.1% and (-2) % of the total fugitive CO₂ and CH₄ emission.

10.1.2 Industrial Processes

Lime production

EU-ETS data have been implemented for one lime production plant for the years 2011 and 2012 leading to a minor decrease in the overall emission of CO₂ emission by 5.51 ktonnes. This change in methodology will be implemented for the previous years in the next submission.

Limestone and Dolomite Use

Activity data for flue gas cleaning have been changed for three power and waste incineration plants in 2011. Consumption of CaCO₃ has increased with 236 tonnes and CaCO₃ containing residues have increased with 669 tonnes resulting in an increased CO₂ emission of 162 tonnes.

An error in transferring data from the emission database to CRF Reporter has been corrected for 1995-2011 for two individual plants.

Chemical industry

The process emission has been adjusted to reflect the total emission reported in environmental reports minus the energy related emissions reported to EU-ETS.

Consumption of Halocarbons and SF₆ – SF₆ – Other

A minor correction has been made to emission of SF₆ from double glazed windows for 2010.

Potential emissions revised for 2005-7 to reflect the fact that potential emission of SF₆ is the same as yearly consumption. Consumption of SF₆ for double glaze windows stopped in 2001.

10.1.3 Solvents and Other Product Use

Recalculations have been made for Other Product Use, where changes were made for the activity data of all four emission sources; candles (2009-2011), fireworks (2009-2011), tobacco (1980-1999, 2011) and charcoal used for barbeques (1980-1987, 2009-2011). These changes have caused recalculations for NMVOC, N₂O and CO₂. NMVOC emissions for the years 1980-1999 have increased between 6.0 % (1997) and 9.1 % (1982) and decreased for 2009-2011 with 2.1 % (2009) to 5.7 % (2011). CO₂ emissions have decreased for the years 2009 (13.4 %) and 2010 (6.1 %) and increased for 2011 (0.7 %), no or minor recalculations were performed for 1980-2008. N₂O emissions have increased for 1980-1999 with between 0.5 % (1999) and 2.8 % (1982) and decreased for the years 2009-2011 with between 0.3 % (2011) and 1.0 % (2009).

10.1.4 Agriculture

Some changes of emissions from the agricultural sector have taken place. These changes reflect decreased emissions in the years 1990-2011 up to 0.2 % compared to the total CO₂-equivalent emission from the agricultural sector. The decrease in 1990-2011 is due to a decrease in the emissions of both N₂O and CH₄.

The CH₄ emission decreases both for emission from enteric fermentation and manure management. The number of geese has been changed for all years and the number of weaners and fattening pigs has been changed for 2011, this affects both emissions from enteric fermentation and manure management. The amount of straw used for bedding for heifers has been changed for 1990-2002 and the amount of biogas treated manure in 2010 has been changed. This affects the emission of CH₄ from manure management.

For the N₂O emission a range of changes have been made, which have both increasing and decreasing effect. EF for NH₃ from synthetic fertiliser has been changed for all years and this affects the emission of N₂O. The emission of N₂O from atmospheric deposition increases due to the change for NH₃ from synthetic fertiliser while the emission of N₂O from synthetic fertiliser decreases. Change in the number of geese decrease the emission of N₂O from grazing for all years. The change of amount of biogas treated manure decreases the N₂O emission in 2010. The emission from manure, manure on soil and leaching is decreased in 2011 due to updated numbers of weaners and fattening pigs.

10.1.5 LULUCF

An update of the LULUCF matrix has taken place for all years. The update was necessary because there were errors in the maps received last year from the Danish Geodata Agency. The problems were especially allocated to a misclassification of recreational areas and a misclassification of parks inside some cities as being forests.

The forest area has been decreased slightly in 2011 (app. 0.3%) compared to the submission last year. This has only had a very limited effect on the total carbon stock in the Danish forest as the carbon stock is estimated in the Na-

tional Forest Inventory, which is independent of the LULUCF matrix. Furthermore has some unclassified areas inside and around the cities which previous was classified as grassland now been removed to Settlement.

Forestry

Since the NFI was initiated in 2002 and have a 5-year rotation, a full measurement is available from 2006. Calculation of carbon stock in the period 2000-2005 is based on interpolation between the carbon stock observed in the NFI in 2006 and the carbon stock as calculated for 2000. For 2006-2012 carbon stock is calculated solely on the basis of the NFI - with additional information about the total forest area from satellite image mapping. Reported values from the NFI correspond to the last year of a five year measurement cycle (i.e. reported values for 2010 rely on data from 2006-2010). Same approach was applied in the 2014 submission.

Cropland, grassland, wetlands and settlements

As the land use matrix is slightly changed the emissions from land use conversion for all sectors are changed slightly for the whole time series. These changes have no effect on the emissions from agricultural soils as these are based information from the EU Land Parcel Information System, i.e. the actual land use. Two minor technical errors have been found in the accounting estimate and corrected: living biomass in Settlements and the area accounted for in Cropland Management and Grassland Management under article 3.4. These errors have only a small impact on the inventory.

10.1.6 Waste

For the category 6A SWDS, an in depth disaggregation of deposited waste for the years 2010, 2011 and 2012 have been performed based on the new waste reporting system in Denmark. 18 categories have been identified of which eleven have been evaluated as inert waste. A detailed description of the waste characterisation, allocation into 18 categories and back casting from 2010 to 1990 are described in Nielsen et al. and Thomsen et al., (2014a and b). The overall results of this detailed characterisation and reallocation of the deposited waste results in a decrease in the CH₄ emission in 1990 of 7.6% and an increase in the CH₄ emission in 2011 of 4.6%.

For the category 6B wastewater handling, no recalculations were made for N₂O. For the methane emissions, the methane correction factor was decreased from 1 to 0.8, which is in accordance to the IPCC guidelines 2006, and which have been further justified by plant specific data as described in Nielsen et al. and Thomsen et al. (2014a, b). Besides the correction of MCF, smaller corrections in plant inlet TOW data have occurred corresponding to a change below 1% throughout the time series.

There are no recalculations in the waste incineration category, however; a correction in the rounding of decimals has caused an increase of 0.07 % of the N₂O emissions from animal cremation for all years 1990-2011.

For the category waste other; changes were made in the vehicle fires and composting source categories. For vehicle fires; the time-series for vehicle population has been updated along with the estimated average weights of some vehicle types, the result is a decrease of CH₄ and CO₂ emissions from 1990-2006 and an increase for 2007-2011. Emission factors have been updated for composting of sludge and organic municipal waste which have resulted in increased emissions of CH₄ and N₂O. The joint effect of these recal-

culations is a decrease in CH₄ emissions from 1980 (2.4 %) to 1984 (2.0 %) and an increase from 1985 (16.7 %) to 2011 (275 %), for CO₂ emissions have decreased from 1980 (1.1 %) to 2006 (0.004 %) and increased from 2007 (0.1 %) to 2011 (0.2 %), finally the N₂O emissions have increased for all years between 1985 (1.1 %) and 2011 (238 %).

10.1.7 KP-LULUCF

A recalculation for KP-LULUCF has been performed for all areas as a consequence of the new land area matrix, see the section on LULUCF.

10.2 Implications for emission levels

For the national total CO₂ equivalent emissions without Land-Use, Land-Use Change and Forestry, the general impact of the improvements and recalculations performed is small and the changes for the whole time-series are between -0.09 % (1990) and 0.48 % (2011). Therefore, the implications of the recalculations on the level and on the trend, 1990-2011, of the national total are small, see Table 10.1.

For the national total CO₂ equivalent emissions with Land-Use, Land-Use Change and Forestry, the general impact of the recalculations is larger due to recalculations in the LULUCF sector. The changes vary between -0.34 % (1990) and +0.55 % (2010), see Table 10.1.

Table 10.1 Recalculation performed in the 2014 submission for 1990-2011. Differences in pct. of CO₂ equivalents between this submission and the May 2013 submission for DK, excluding Greenland and Faroe Islands.

Year	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
Total CO ₂ eqv. Emissions with Land-Use Change and Forestry	-0.34	-0.03	-0.02	-0.01	0.01	0.03	0.03	0.06	0.10	0.17	0.46
Total CO ₂ eqv. Emissions without Land-Use Change and Forestry	-0.09	-0.07	-0.06	-0.04	-0.02	0.00	0.01	0.04	0.08	0.16	0.46
<i>Continued</i>	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011
Total CO ₂ eqv. Emissions with Land-Use Change and Forestry	0.18	0.29	0.35	0.17	-0.11	-0.05	0.27	-0.26	0.54	0.55	0.36
Total CO ₂ eqv. Emissions without Land-Use Change and Forestry	0.23	0.35	0.40	0.21	0.23	0.24	0.29	0.22	0.37	0.30	0.48

10.3 Implications for emission trends, including time series consistency

It is a high general priority in the considerations leading to recalculations back to 1990 to have and preserve the consistency of the activity data and emissions time-series. As a consequence activity data, emission factors and methodologies are carefully chosen to represent the emissions for the time-series correctly. Often considerations regarding the consistency of the time-series have led to recalculations for single years when activity data and/or emission factors have been changed or corrected. Furthermore, when new sources are considered, activity data and emissions are as far as possible introduced to the inventories for the whole time-series based on preferably the same methodology.

The implication of the recalculations is further shown in Tables 10.2-10.5.

Table 10.2 Recalculation for CO₂ performed in the 2014 submission for 1990-2011. Differences in Gg CO₂ eqv. between this and the May 2013 submission for DK. Excluding Greenland and Faroe Islands.

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
Total National Emissions and Removals	-128	89	88	88	86	84	82	86	84	84	232
1. Energy	63	63	63	65	64	62	62	64	63	64	211
1.A. Fuel Combustion Activities	61	61	61	62	64	62	62	64	63	64	211
1.A.1. Energy Industries	-	-	-	1	68	5	12	11	11	12	12
1.A.2. Manufacturing Industries and Construction	59	59	60	60	48	56	50	52	51	52	52
1.A.3. Transport	0	0	0	0	0	0	0	0	0	0	0
1.A.4. Other Sectors	2	1	1	1	-52	0	1	1	1	0	147
1.A.5. Other	-	-	-	-	-	-	-	-	-	-	-
1.B. Fugitive Emissions from Fuels	2	3	2	3	0	0	0	0	0	0	0
2. Industrial Processes	-	-	-	-	-	1	0	3	2	3	5
2.A. Mineral Products	-	-	-	-	-	1	1	2	2	3	5
2.B. Chemical Industry	-	-	-	-	-	-	-1	1	-	-	0
2.C. Metal Production	-	-	-	-	-	-	-	-	-	-	-
2.D. Other Production	-	-	-	-	-	-	-	-	-	-	-
2.G. Other	-	-	-	-	-	-	-	-	-	-	-
3. Solvent and Other Product Use	-	-	-	-	-	-	-	-	-	-	-
4. Agriculture	-	-	-	-	-	-	-	-	-	-	-
5. Land Use, Land-Use Change and Forestry (net)	-190	26	25	24	23	22	21	20	19	18	17
5.A. Forest Land	7	7	7	7	7	7	7	7	7	7	7
5.B. Cropland	-192	26	26	27	27	27	27	27	27	27	28
5.C. Grassland	0	0	0	0	0	0	0	0	0	0	0
5.D. Wetlands	-3	-5	-6	-7	-8	-9	-10	-11	-12	-13	-14
5.E. Settlements	-3	-3	-3	-3	-4	-4	-4	-4	-4	-4	-4
5.F. Other Land	-	-	-	-	-	-	-	-	-	-	-
5.G. Other	-	-	-	-	-	-	-	-	-	-	-
6. Waste	-1	-1	-1	-1	-1	-1	-1	0	0	0	0
6.C. Waste Incineration	-	-	-	-	-	-	-	-	-	-	-
6.D. Other	-1	-1	-1	-1	-1	-1	-1	0	0	0	0
<i>Continued</i>	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011
Total National Emissions and Removals	41	42	82	80	-121	-106	99	-247	239	248	66
1. Energy	65	64	102	100	99	108	108	50	136	100	145
1.A. Fuel Combustion Activities	65	64	102	100	99	108	108	50	136	100	145
1.A.1. Energy Industries	11	11	34	12	11	7	6	13	34	29	8
1.A.2. Manufacturing Industries and Construction	54	53	84	88	90	86	125	110	108	153	133
1.A.3. Transport	0	0	0	0	0	0	0	-69	0	-4	-13
1.A.4. Other Sectors	0	0	-16	0	-2	15	-23	-3	-5	-78	18
1.A.5. Other	-	-	-	-	-	-	-	-	-	-	-
1.B. Fugitive Emissions from Fuels	0	0	0	0	0	0	0	0	0	0	0
2. Industrial Processes	2	1	1	-1	-1	3	1	1	1	4	-2
2.A. Mineral Products	2	1	1	1	1	3	2	3	2	4	-2
2.B. Chemical Industry	0	0	-1	-2	-2	-1	-1	-1	-1	-	-1
2.C. Metal Production	-	-	-	-	-	-	-	-	-	-	-
2.D. Other Production	-	-	-	-	-	-	-	-	-	-	-
2.G. Other	-	-	-	-	-	-	-	-	-	-	-
3. Solvent and Other Product Use	-	-	-	-	-	-	-	-	-12	-7	1
4. Agriculture	-	-	-	-	-	-	-	-	-	-	-
5. Land Use, Land-Use Change and Forestry (net)	-25	-23	-21	-19	-219	-216	-10	-299	114	150	-78
5.A. Forest Land	-34	-30	-27	-23	-51	-49	157	-132	280	316	88
5.B. Cropland	28	27	27	27	-181	-181	-181	-181	-181	-181	-181
5.C. Grassland	0	0	0	0	32	31	31	31	30	30	29
5.D. Wetlands	-15	-16	-17	-18	-14	-13	-13	-12	-11	-11	-10
5.E. Settlements	-4	-4	-4	-4	-5	-5	-5	-4	-4	-4	-4

<i>Continued</i>											
5.F. Other Land	-	-	-	-	-	-	-	-	-	-	-
5.G. Other	-	-	-	-	-	-	-	-	-	-	-
6. Waste	0	0	0	0	0	0	0	0	0	0	0
6.C. Waste Incineration	-	-	-	-	-	-	-	-	-	-	-
6.D. Other	0	0	0	0	0	0	0	0	0	0	0

Table 10.3 Recalculation for CH₄ performed in the 2014 submission for 1990-2011. Differences in Gg CO₂ eqv. between this and the May 2013 submission for DK. Excluding Greenland and Faroe Islands.

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
Total National Emissions and Removals	-113.9	-108.2	-101.0	-89.3	-74.7	-60.7	-50.6	-37.5	-28.3	-20.5	-17.2
1. Energy	5.6	6.8	6.4	7.3	7.3	8.6	7.7	7.0	6.1	4.7	5.7
1.A. Fuel Combustion Activities	4.8	5.6	5.8	6.4	6.8	7.1	7.5	6.6	5.7	6.2	7.5
1.A.1. Energy Industries	0.0	0.0	0.0	0.0	0.6	1.9	3.4	2.2	2.0	2.2	2.3
1.A.2. Manufacturing Industries and Construction	0.0	0.0	0.0	0.0	0.0	-1.2	-2.8	-2.5	-2.3	-2.4	-2.8
1.A.3. Transport	-0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
1.A.4. Other Sectors	4.9	5.7	5.9	6.5	6.2	6.4	6.9	6.9	6.1	6.3	8.1
1.A.5. Other	-	-	-	-	-	-	-	-	-	0.0	0.0
1.B. Fugitive Emissions from Fuels	0.8	1.2	0.6	0.9	0.5	1.5	0.2	0.4	0.4	-1.5	-1.8
2. Industrial Processes	-	-	-	-	-	-	-	-	-	-	-
3. Solvent and Other Product Use	-	-	-	-	-	-	-	-	-	-	-
4. Agriculture	-8.0	-8.9	-9.6	-10.4	-10.8	-11.6	-12.7	-12.7	-12.6	-12.3	-12.4
4.A. Enteric Fermentation	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
4.B. Manure Management	-8.0	-8.9	-9.6	-10.4	-10.8	-11.6	-12.7	-12.7	-12.6	-12.3	-12.4
4.F. Field Burning of Agricultural Residues	-	-	-	-	-	-	-	-	-	-	-
5. Land Use, Land-Use Change and Forestry (net)	-	-	-	-	-	-	-	-	-	-	-
6. Waste	-111.5	-106.2	-97.8	-86.2	-71.3	-57.7	-45.6	-31.8	-21.8	-12.9	-10.5
6.A. Solid Waste Disposal on Land	-112.0	-106.9	-98.9	-87.5	-72.7	-60.1	-47.6	-34.3	-24.5	-16.4	-14.9
6.B. Waste-water Handling	-0.7	-0.7	-0.7	-0.7	-0.8	-0.8	-1.0	-1.2	-1.1	-1.3	-0.9
6.C. Waste Incineration	-	-	-	-	-	-	-	-	-	-	-
6.D. Other	1.2	1.5	1.8	2.0	2.2	3.2	3.0	3.7	3.8	4.9	5.4
<i>Continued</i>	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011
Total National Emissions and Removals	-15.2	-6.8	13.8	15.5	21.3	28.5	37.4	36.7	42.3	22.2	53.4
1. Energy	7.8	7.9	9.9	11.1	13.4	16.2	20.6	18.3	16.7	18.7	16.9
1.A. Fuel Combustion Activities	7.9	7.9	9.7	11.0	13.4	16.1	20.3	18.5	16.9	19.1	16.9
1.A.1. Energy Industries	1.9	2.1	1.9	2.2	2.0	2.0	1.8	1.7	2.0	2.0	2.1
1.A.2. Manufacturing Industries and Construction	-3.0	-3.5	-3.0	-3.5	-3.3	-2.3	-1.8	-2.0	-2.2	-0.9	-0.8
1.A.3. Transport	0.0	0.0	0.0	0.0	0.1	0.0	0.1	0.0	0.3	0.3	0.3
1.A.4. Other Sectors	9.0	9.3	10.8	12.3	14.6	16.4	20.2	18.7	16.9	17.8	15.2
1.A.5. Other	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
1.B. Fugitive Emissions from Fuels	0.0	0.0	0.1	0.1	0.1	0.1	0.3	-0.2	-0.2	-0.4	0.0
2. Industrial Processes	-	-	-	-	-	-	-	-	-	-	-
3. Solvent and Other Product Use	-	-	-	-	-	-	-	-	-	-	-
4. Agriculture	-13.0	-12.4	0.0	0.0	0.0	0.0	0.0	0.0	0.0	-1.0	-0.2
4.A. Enteric Fermentation	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	-0.1
4.B. Manure Management	-13.0	-12.4	0.0	0.0	0.0	0.0	0.0	0.0	0.0	-1.0	-0.2
4.F. Field Burning of Agricultural Residues	-	-	-	-	-	-	-	-	-	-	-
5. Land Use, Land-Use Change and Forestry (net)	-	-	-	-	-	-	-	-	-	-	-
6. Waste	-10.0	-2.4	4.0	4.5	7.9	12.3	16.8	18.5	25.6	4.4	36.7
6.A. Solid Waste Disposal on Land	-14.4	-8.2	-1.9	2.0	5.9	9.6	13.9	17.1	21.2	0.2	32.3
6.B. Waste-water Handling	-1.4	-1.8	-1.9	-2.1	-1.9	-1.6	-1.3	-3.0	-1.9	-1.2	-1.3
6.C. Waste Incineration	-	-	-	-	-	-	-	-	-	-	-
6.D. Other	5.7	7.7	7.8	4.6	3.9	4.3	4.2	4.3	6.3	5.5	5.7

Table 10.4 Recalculation for N₂O performed in the 2014 submission for 1990-2011. Differences in Gg CO₂ eqv. between this and the May 2013 submission for DK. Excluding Greenland and Faroe Islands.

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
Total National Emissions and Removals	-7.6	-6.9	-6.1	-7.4	-6.8	-2.3	-1.2	0.7	26.1	69.6	-7.6
1. Energy	1.7	2.3	1.9	1.4	1.4	0.9	1.6	1.8	0.6	2.2	1.7
1.A. Fuel Combustion Activities	1.8	2.4	1.9	1.5	1.4	0.9	1.6	1.8	0.6	2.2	1.8
1.A.1. Energy Industries	0.7	1.2	0.8	0.4	0.4	0.3	0.9	1.2	0.0	1.5	0.7
1.A.2. Manufacturing Industries and Construction	1.1	1.1	1.1	1.1	1.1	0.6	0.7	0.6	0.6	0.6	1.1
1.A.3. Transport	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
1.A.4. Other Sectors	-	0.0	0.0	0.0	-0.1	0.0	0.0	0.1	0.0	0.0	-
1.A.5. Other	-	-	-	-	-	-	-	-	-	0.0	-
1.B. Fugitive Emissions from Fuels	-0.1	-0.1	-0.1	-0.1	-	-	-	-	-	-	-0.1
2. Industrial Processes	-	-	-	-	-	-	-	-	-	-	-
3. Solvent and Other Product Use	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
4. Agriculture	-10.5	-10.5	-9.7	-10.7	-10.3	-9.8	-8.7	-8.2	-7.6	-6.8	-10.5
4.A. Enteric Fermentation	-	-	-	-	-	-	-	-	-	-	-
4.B. Manure Management	-	-	-	-	-	-	-	-	-	-	-
4.D. Agricultural Soils	-10.5	-10.5	-9.7	-10.7	-10.3	-9.8	-8.7	-8.2	-7.6	-6.8	-10.5
4.F. Field Burning of Agricultural Residues	-	-	-	-	-	-	-	-	-	-	-
5. Land Use, Land-Use Change and Forestry (net)	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.1	0.0
5.A. Forest Land	-0.2	-0.2	-0.2	-0.2	-0.2	-0.2	-0.2	-0.2	-0.2	-0.2	-0.2
5.B. Cropland	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
5.C. Grassland	-	-	-	-	-	-	-	-	-	-	-
5.D. Wetlands	-	-	-	-	-	-	-	-	-	-	-
5.E. Settlements	-	-	-	-	-	-	-	-	-	-	-
5.F. Other Land	-	-	-	-	-	-	-	-	-	-	-
5.G. Other	-	-	-	-	-	-	-	-	-	-	-
6. Waste	1.1	1.3	1.6	1.8	2.1	6.5	5.9	7.0	33.1	74.2	1.1
6.B. Waste-water Handling	-	-	-	-	-	-	-	-	-	-	-
6.C. Waste Incineration	-	-	-	-	-	-	-	-	-	-	-
6.D. Other	1.1	1.3	1.6	1.8	2.1	6.5	5.9	7.0	33.1	74.2	1.1
<i>Continued</i>	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011
Total National Emissions and Removals	115.2	108.6	181.7	176.5	26.4	24.9	36.5	48.7	49.7	59.7	65.2
1. Energy	2.9	-1.0	-1.5	-0.6	-0.2	0.2	2.0	1.9	1.5	1.9	2.1
1.A. Fuel Combustion Activities	2.9	-1.0	-1.5	-0.6	-0.2	0.2	2.0	1.9	1.5	1.9	2.1
1.A.1. Energy Industries	0.0	0.1	0.0	0.4	0.7	0.8	0.6	0.0	0.1	0.3	0.0
1.A.2. Manufacturing Industries and Construction	0.0	-1.1	-1.6	-0.5	-1.0	-1.0	1.4	2.2	1.7	1.9	3.8
1.A.3. Transport	0.0	0.0	0.1	0.1	0.2	0.4	-0.2	-0.1	-0.6	0.0	0.0
1.A.4. Other Sectors	2.9	0.0	0.0	-0.6	0.0	0.0	0.2	-0.4	0.2	-0.3	-1.8
1.A.5. Other	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
1.B. Fugitive Emissions from Fuels	-	-	-	-	-	-	-	-	-	-	-
2. Industrial Processes	-	-	-	-	-	-	-	-	-	-	-
3. Solvent and Other Product Use	-	-	-	-	-	-	-	-	-	0.0	0.0
4. Agriculture	-6.1	-5.6	-5.0	-5.0	-5.2	-4.8	-4.2	-4.2	-4.6	-3.5	-4.3
4.A. Enteric Fermentation	-	-	-	-	-	-	-	-	-	-	-
4.B. Manure Management	-	-	-	-	-	-	-	-	-	-	-0.6
4.D. Agricultural Soils	-6.1	-5.6	-5.0	-5.0	-5.2	-4.8	-4.2	-4.2	-4.6	-3.5	-3.7
4.F. Field Burning of Agricultural Residues	-	-	-	-	-	-	-	-	-	-	-
5. Land Use, Land-Use Change and Forestry (net)	0.1	0.1	0.1	0.1	0.1	-0.1	-0.1	-0.1	0.0	0.0	0.0
5.A. Forest Land	-0.2	-0.2	-0.2	-0.2	-0.2	-0.2	-0.2	-0.2	-0.2	-0.2	-0.2
5.B. Cropland	0.2	0.2	0.2	0.3	0.3	0.1	0.1	0.1	0.1	0.1	0.2
5.C. Grassland	-	-	-	-	-	-	-	-	-	-	-
5.D. Wetlands	-	-	-	-	-	-	-	-	-	-	-
5.E. Settlements	-	-	-	-	-	-	-	-	-	-	-

<i>Continued</i>												
5.F. Other Land	-	-	-	-	-	-	-	-	-	-	-	-
5.G. Other	-	-	-	-	-	-	-	-	-	-	-	-
6. Waste	118.4	115.0	188.1	182.0	31.7	29.6	38.8	51.1	52.9	61.4	67.4	
6.B. Waste-water Handling	-	-	-	-	-	-	-	-	-	-	-	-
6.C. Waste Incineration	-	-	-	-	-	-	-	-	-	-	-	-
6.D. Other	118.4	115.0	188.1	182.0	31.7	29.6	38.8	51.1	52.9	61.4	67.4	

Table 10.5 Recalculation for HFCs, PFCs and SF₆ performed in the 2014 submission for 1990-2011. Differences in Gg CO₂ eqv. between this and the May 2013 submission for DK. Excluding Greenland and Faroe Islands.

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	
HFC	-	-	-	-	-	-	-	-	-	-	-	
PFC	-	-	-	-	-	-	-	-	-	-	-	
SF ₆	-	-	-	-	-	-	-	-	-	-	-	
<i>Continued</i>	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	
HFC	-	-	-	-	-	-	-	-	0,0	0,0	-	
PFC	-	-	-	-	-	-	-	-	-	-	-	
SF ₆	-	-	-	-	-	-	-	-	-	-0,4	-	

10.4 Recalculations, including those in response to the review process, and planned improvements to the inventory (e.g. institutional arrangements. inventory preparations)

The review on the submissions in 2007 and 2008 was finalised and the report was published April 15, 2009. For the 2009 submission the review report was finalised and published April 15 2010. The review report of the in-country review of the 2010 submission was published March 3 2011. The draft review report for the review of the 2011 submission was available February 9, 2012. The final review report was published April 30 2012. The draft review report of the 2012 submission was made available April 30 2013 and the final review report was dated August 2 2013. The significant delay made it impossible to ensure that all findings by the ERT were addressed in the 2013 submission and with the late publication even the inclusion in the 2014 submission was challenging.

As of April 7 2014 Denmark has not received a draft of the review report from the centralised review carried out in September 2013. Therefore, it has not been possible to address the outcome of the 2013 review in this report. The main recommendations from the reviews of the 2008, 2009, 2010, 2011 and 2012 submissions are listed in Table 10.6.

To keep the table transparent the recommendations that have been completed from the review of the 2008, 2009, 2010 and 2011 submissions have been deleted.

Table 10.6 Main recommendations from the reviews of the 2008, 2009, 2010, 2011 and 2012 submissions.

CRF	ERT Comment	Denmark's response	Reference
2008 submission (Review report: http://unfccc.int/resource/docs/2009/arr/dnk.pdf)			
Energy, road transport – Paragraph 41	The change of non-CO ₂ EFs associated with the use of bioethanol in gasoline blends has not been taken into account when estimating the corresponding emissions. The ERT suggests that Denmark assess probable changes to these EFs in its next annual submission.	No data has previously been available indicating different CH ₄ and N ₂ O emission factors for blends of fossil and biogenic fuels. This issue is being followed in case new research indicates otherwise.	Chapter 3.3.2.
2009 submission (Review report: http://unfccc.int/resource/docs/2010/arr/dnk.pdf)			
CRF	ERT Comment	Denmark's response	Reference
Industrial Processes, limestone and dolomite use – Paragraph 62	The ERT noted that estimates for more recent years could be underestimated in comparison to previous years in the time-series, and the full time-series may not be consistent. Besides, the ERT also found some potential inconsistencies in the time-series of AD: an increase of 28.1 per cent in 2005–2006 and a 28.8 per cent decrease in 2006–2007. The ERT recommends that Denmark provide in the NIR of its next annual submission information on the specific procedures and verifications the Party used.	The sector (2A3) comprises a number of different processes: consumption of CaCO ₃ to flue gas cleaning at power plants and waste incineration plants, production of mineral wool and refining of sugar. The activity data are not comparable or even confidential or lacking for part of the period. The inventory team are working on improvement of the documentation of AD. <u>2014 update:</u> The limestone use in sugar production was reallocated so that the category now only comprises flue gas desulphurisation and production of stone wool. When implementing the 2006 IPCC Guidelines for the 2015 submission the allocation of emissions from the use of limestone and dolomite will be further analysed.	
Solvent and other product use, use of N ₂ O – Paragraph 64	The ERT encourages the Party to provide estimates of emissions of N ₂ O from use as anaesthesia for the period 1990–2004 in order to complete the time-series.	The producers and distributors of N ₂ O will be contacted again and if data cannot be given for 1990-2004 this will be clearly explained in the report. <u>2013 update:</u> Based on contact with the industry, data for 2000-2004 have been obtained and included in the inventory.	
Agriculture, manure management – Paragraph 74	Denmark treats some of its animal slurries in biogas plants, capturing the CH ₄ generated and using it for electricity and CHP production. In response to a question from the ERT, Denmark stated that some of the information in table 6.12 of the NIR might be misleading with regard to the energy production values expressed in TJ, as they are not directly related to the estimation of CH ₄ captured, but rather	DK agree that the information on the energy production can be misleading. The calculation of the lower CH ₄ emission as a consequence of biogas treatment is based on the amount of biogas treated slurry, which is received from the Danish Energy Agency. Table 6.1 includes data concerning the amount of slurry, the VS content in the treated slurry and the reduced emission. DK has planned to improve the possibilities to verify the calculation of the reduced emission from biogas treated slurry. This could be done by contacting a biogas plant in preparation for potential data based on	Chapter 6.4.2 Table 6.18

Table 10.6 Main recommendations from the reviews of the 2008, 2009, 2010, 2011 and 2012 submissions.

CRF	ERT Comment	Denmark's response	Reference
	<p>were obtained independently from the Danish Energy Agency. The ERT recommends that the Party report estimates of energy production and CH₄ recovery in a consistent way, and correct table 6.12 in the next annual submission.</p> <p>The ERT recommends that Denmark, for the sake of improving transparency, provide plant-specific data regarding energy output and quantities of slurry treated from one or more of the larger biogas plants. The ERT also recommends that Denmark use the energy output from plants to assess the validity of the CH₄ reduction potentials for cattle and swine slurry as an additional QC check.</p>	<p>measurement from slurry.</p> <p><u>2014 update:</u> The Danish Energy Agency (DEA) provides annually the Danish energy statistics and from 2012 we succeed to receive plant specific data for the energy production based in biogas. Next step is to investigate the possibilities to improve the data regarding the energy potential in order to improve the estimate of biogas treated manure.</p>	
2010 submission (Review report: http://unfccc.int/resource/docs/2011/arr/dnk.pdf)			
Energy, Country-specific issues – Paragraph 55	To improve accuracy, the ERT recommends that Denmark make efforts to estimate CO ₂ emissions from gas oil used in Greenland by using country-specific EFs that are already available.	<p>This issue will be investigated further with the aim of revising the CO₂ emission factor for gas oil combusted in Greenland.</p> <p><u>2014 update:</u> Based on fuel analysis the CO₂ emission factor for gas oil in Greenland has been updated.</p>	Chapter 16
LULUCF, Land converted to cropland – Paragraph 123	<p>For land converted to cropland, net carbon stock change of mineral and organic soils is reported as "IE" for many conversions. During the review, Denmark explained that these have been included in cropland remaining cropland.</p> <p>To improve transparency, the ERT recommends that Denmark report net carbon stock change of mineral and organic soils separately under cropland remaining cropland and land converted to cropland in the next annual submission.</p>	<p>The recommendation from the ERT is difficult to follow. Although we have very detailed information on the individual fields it will be a very time consuming task with little effect.</p> <p>The area with soil in agricultural use is based on the detailed information on the position of the field and the actual crop grown in that field. The minor areas which is converted to cropland and its use is included in the modelling with C-TOOL and for the organic soils we use an overlay of the current used fields to see their position in relation to the organic soil map.</p>	
2011 submission (Review report: http://unfccc.int/resource/docs/2012/arr/dnk.pdf)			
Industrial processes, cement produc-	The ERT also questioned the Party, during the	The work is on-going.	Chapter 4.2.2

Table 10.6 Main recommendations from the reviews of the 2008, 2009, 2010, 2011 and 2012 submissions.

CRF	ERT Comment	Denmark's response	Reference
tion – Paragraph 71	review, as to whether it accounts for imports and exports for the early years of the time series, which are required to be taken into account when using a tier 1 approach. The Party responded to the ERT that it believes that clinker production at that time was solely for the company's own use, but that it will research this further and confirm in its next annual submission. The ERT recommends that Denmark conduct this research to ensure that the tier 1 approach is being implemented in accordance with the IPCC good practice guidance for estimating emissions for the early years of the time series.	<u>2014 update:</u> Denmark has collected the data for import/export of clinker and documented it in the NIR.	
Industrial processes, cement production – Paragraph 72	The ERT further questioned Denmark on its consideration of cement kiln dust (CKD) in the time series of emission estimates, in particular for the earlier years. Denmark responded that, although it is known that the emission estimates are based on the different types of clinker used, there is no information to indicate whether CKD is included in the emission estimates. The ERT recommends that Denmark continue to pursue any information that could clarify whether CKD is included in the emission estimates for all years of the time series.	The ERT has been informed that no further information is available for the years 1990-1997. The work with including CKD in the emission estimates is on-going. <u>2014 update:</u> According to information from the plant CKD is included in the emission calculation for 1990-1997.	
Industrial processes, Consumption of halocarbons – Paragraph 74	The ERT concluded that Denmark has provided complete estimates for these disposal emissions. However, the ERT recommends that Denmark be more transparent and provide the rationale for this determination in the NIR of its next annual submission.	This work is on-going. <u>2014 update:</u> Denmark has expanded the discussion regarding de-commissioning of f-gas containing products.	Chapter 4.7.2
2012 submission (Review report: http://unfccc.int/resource/docs/2013/arr/dnk.pdf)			
General, completeness – Paragraph	Provide a complete set of CRF tables, includ-	By mistake the explanations for recalculation were not included in the	

Table 10.6 Main recommendations from the reviews of the 2008, 2009, 2010, 2011 and 2012 submissions.

CRF	ERT Comment	Denmark's response	Reference
10	ing complete information in CRF table 8(b), in accordance with the requirements of the "Guidelines for the preparation of national communications by Parties included in Annex I to the Convention, Part I: UNFCCC reporting guidelines on annual inventories"	aggregate CRF. However, all recalculations were thoroughly explained and documented in the NIR. It will be ensured that it is checked in the future that the table is completed.	
Energy, International bunker fuels – Paragraph 39	Provide explanations in the NIR on the large inter-annual variations in CO ₂ emissions from international bunker fuels for the years 2008–2010	Explanations on the fluctuations in the trend have been included in the NIR.	Chapter 3.3.1
Energy, International bunker fuels – Paragraph 40	Include data on international bunkers for lubricants in CRF table 1.C and improve the associated QC procedures	Data for lubricants used in international bunkers were included in the 2014 submission. The QC procedures have been updated.	CRF
Energy, Feedstocks and non-energy use of fuels – Paragraph 42	Include a reference in CRF table 3.A-D to clarify the reporting of CO ₂ emissions from white spirits	A reference was added in the CRF.	CRF
Energy, Feedstocks and non-energy use of fuels – Paragraph 43	Document and justify in the NIR why a carbon storage factor of 1.00 has been used	<p>The documentation in the NIR Chapter 3.4 has been improved as has the documentation in the CRF</p> <p>In CRF Table 1A.(c) a reference to NIR chapter 3.4 have been added.</p> <p>In CRF Table 1A (c) the following comment have been added: Non-energy use of fuels is not included in sector 1A in the Danish National Approach (IE). Fuel consumption for non-energy is subtracted in Reference Approach to make results comparable. This have been done by setting the fraction stored equal to 1.</p> <p>In CRF Table 1A(d) the following comments have been added:</p> <p>1.AD.2 Lubricants:All emissions from the non-energy use of the three fuels identified in the Danish energy statistics are included in the Industrial Processes and the Solvent and Other Product Use sectors. In order to make the reference approach comparable with the sectoral approach the fraction of carbon stored has been set to 1. This is to exclude it from the comparison with the sectoral approach. The relevant emissions are included in the sectors identified in the table.</p> <p>1.AD.3 Bitumen:All emissions from the non-energy use of the three fuels identified in the Danish energy statistics are included in the Indus-</p>	Chapter 3.4 CRF Table 1.A(d) and CRF Table 1.A(c)

Table 10.6 Main recommendations from the reviews of the 2008, 2009, 2010, 2011 and 2012 submissions.

CRF	ERT Comment	Denmark's response	Reference
		<p>trial Processes and the Solvent and Other Product Use sectors. In order to make the reference approach comparable with the sectoral approach the fraction of carbon stored has been set to 1. This is to exclude it from the comparison with the sectoral approach. The relevant emissions are included in the sectors identified in the table.</p> <p>1.AD.10 White Spirit: All emissions from the non-energy use of the three fuels identified in the Danish energy statistics are included in the Industrial Processes and the Solvent and Other Product Use sectors. In order to make the reference approach comparable with the sectoral approach the fraction of carbon stored has been set to 1. This is to exclude it from the comparison with the sectoral approach. The relevant emissions are included in the sectors identified in the table.</p>	
Energy, Stationary combustion: solid and liquid fuels – Paragraph 44	Reflect the results of the analysis of the CO ₂ EFs for fuel oil in the NIR	The analysis of the CO ₂ emission factors for fuel oil like in the case of coal did not yield any useable results.	Chapter 3.2.5
Energy, Road transportation: liquid fuels – Paragraph 45	Provide brief descriptions of the methods used to obtain the fleet and mileage data necessary for the COPERT IV model	The description has been improved in the NIR.	Chapter 3.3.2
Energy, Aviation: liquid fuels - Paragraph 48	Provide information on the number of domestic landings and take-offs (LTOs) per representative aircraft type for each of the Danish airports, including flights between Denmark and Greenland/the Faroe Islands, the average LTO fuel consumption and the EFs per representative aircraft type	Information has been included.	Annex 3B-10
Industrial processes, Cement production – Paragraph 55	Provide information on the imports and exports of cement for the years 1990–1997, in order to ensure that the tier 1 method is being implemented in accordance with the IPCC Good Practice Guidance and Uncertainty Management in National Greenhouse Gas Inventories (the IPCC good practice guidance)	The information has been included.	Chapter 4.2.2
Industrial processes, Cement production – Paragraph 56	Provide relevant information to clarify whether cement kiln dust is included in the emission estimates for the years prior to 1998	The work is ongoing.	
Agriculture, Sector overview - Paragraph 66	Provide a description of the use of the biogas and the associated energy output	More descriptive text has been implemented and furthermore annex 3E now includes two tables with detailed information on data used to calcu-	Annex 3E-12a and 3E-12b

Table 10.6 Main recommendations from the reviews of the 2008, 2009, 2010, 2011 and 2012 submissions.

CRF	ERT Comment	Denmark's response	Reference
		late the reduction from biogas treated manure.	
Agriculture, Manure management – Paragraph 70	Follow the methodology provided in the IPCC good practice guidance, make further efforts to obtain the necessary AD, as outlined in the planned inventory improvements in the 2012 NIR, and make recalculations for 2010	The Danish Energy Agency (DEA) provides annually the Danish energy statistics and from 2012 we succeed to receive plant specific data for the energy production based on biogas.	Chapter 6.4.2 Table 6.18
Agriculture, Manure management – Paragraph 71	Provide explanations for the inter-annual changes in the NH ₃ IEF	The main documentation for the Danish NH ₃ emission inventory is the Informative Inventory report. This is reported annually to the UNECE Convention on Long-Range Transboundary Air Pollution. This report is included with a direct link in the list of references.	
Agriculture, Manure management – Paragraph 72	Explain whether the reduction in the EF for ammonium sulphate was the result of using different methodologies and protocols for the national review of for other reasons	The main documentation for the Danish NH ₃ emission inventory is the Informative Inventory report. This is reported annually to the UNECE Convention on Long-Range Transboundary Air Pollution. This report is included with a direct link in the list of references.	
Agriculture, Direct emissions from soils – Paragraph 73	Improve transparency by providing disaggregated data on the amount of crop residues that are used for each purpose (i.e. feeding, bedding and energy production)	Additional description is included in the NIR and a new table in Annex 3E, which shows the amount of harvested straw used for fodder, fuel and bedding.	Annex 3E-18 Chapter 6.5.2
Agriculture, Direct emissions from soils – Paragraph 74	Include, in the NIR, an explanation to substantiate the reduction in N ₂ O emissions from nitrogen-fixing crops	Additional documentation was added in the NIR regarding the basis for estimating N ₂ O emissions from N-fixing crops.	Chapter 6.5.2 Annex 3E-15
LULUCF, Sector overview – Paragraph 78	Improve the QA/QC processes for the LULUCF sector and report on the improvements made, particularly those related to the NFI in terms of sampling procedures and estimation methods	The QC procedures have been further expanded for all parts of the LULUCF sector.	QC chapters within Chapter 7
LULUCF, Forest land remaining forest land – Paragraph 80	Make continuous efforts to ensure time-series consistency	The work to improve time-series consistency is continuously ongoing. Any changes will be reflected in the NIR.	
LULUCF, Forest land remaining forest land – Paragraph 81	Provide additional information to explain the large inter-annual variations in the carbon stock in forest land remaining forest land (i.e. information on changes in the composition of tree species and the age structure of forest stands; the area and volume of clear cutting; and the area subject to destructive disturb-	Additional information has been provided in the NIR.	Chapter 7.2

Table 10.6 Main recommendations from the reviews of the 2008, 2009, 2010, 2011 and 2012 submissions.

CRF	ERT Comment	Denmark's response	Reference
	ance)		
LULUCF, Cropland remaining cropland – Paragraph 83	Include the underlying data that support the explanations of the emissions from cropland remaining cropland, particularly those that demonstrate the link between temperature and yield	There is not clear relationship between temperature and crop yield. The crop yield depends on many different factors where precipitation and sun hours combined with decent temperatures are the most important. The temperature is only used in the degradation model for organic matter in the soil. Here are used monthly mean values for the different years. More detailed data e.g. on a daily basis has very little influence on the output from the degradation C-TOOL. Annual average temperature file has been included in Annex 3F table 14.	Annex 3F table 14
Waste, Sector overview – Paragraph 86	Provide more complete information on the assumptions made and methodologies used, and on the choice of EFs and AD	More information has been provided in the NIR.	Chapter 8
Waste, Sector overview – Paragraph 87	Verify all data used for the estimation of emissions in the waste sector in the next annual submission, including verifying that the provided expert judgement is conservative with reference to the default value for flowing sewers, which is zero according to the IPCC 2006 guidelines. If data are not available, to verify that the chosen MCF is conservative, perform direct source testing to verify existing relevant literature on fugitive emissions from sewer systems	A descriptive section on available data on VOC (e.g. HS) emissions from sewer systems has been included in chapter 8. The estimated fraction of the population not connected to the sewer system has been kept at 10% (unrealistic high) until the ongoing mapping of extension of the sewer system in Denmark I finalised. As the contribution to the methane emission is dominated by the contribution from septic tanks, the methane emissions from 6B is conservative and overestimated. However, updated information documenting any fugitive emissions from the Danish sewer system will be included as soon as data becomes available.	Chapter 8.3
Waste, Solid waste disposal on land – Paragraph 89	Analyse and report, in the next annual submission, updated information on the composition of the waste category “other combustibles”, divide it into different well characterised waste types, in order to document and assign each type a DOC value	The information regarding the waste characterisation in the new waste database has been improved.	Chapter 8.2
Waste, Wastewater handling - Paragraph 91	Perform a review of the biogas production data from plants for the entire time series, consistent with the requirements of the IPCC good practice guidance, and include a time trend for the amount of recovered methane, in its next	Data collection has been done for a number of plants. However, the work to collect the data is substantial and therefore ongoing. The results achieved so far are documented in the NIR.	Chapter 8.3.2 and 8.5.5

Table 10.6 Main recommendations from the reviews of the 2008, 2009, 2010, 2011 and 2012 submissions.

CRF	ERT Comment	Denmark's response	Reference
	annual submission		
Waste, Wastewater handling - Paragraph 91	Provide information on the physical characteristics of the sewer systems in the NIR	A descriptive section on structural characteristics and ongoing improvements and extensions of the sewer system have been included in chapter 8	Chapter 8.3

More information on the specific responses to the review has been given in the sectoral chapters of this report.

10.5 Explanations, justifications and implications of recalculations for KP-LULUCF inventory

10.5.1 Recalculations

Almost all sectors in the KP-LULUCF have been recalculated.

This is due to:

- A revision of the land use matrix for the entire period 1990 to 2012
- Updated data from the Danish National Forest Inventory (NFI) for carbon stock changes in above/below ground, dead wood and litter

For more information on KP-LULUCF recalculations please refer to Chapter 11.

Table 10.7 Effect of the recalculations in the KP-LULUCF sector for 1990 and 2011, Net CO₂ equivalents.

GREENHOUSE GAS SOURCE AND SINK ACTIVITIES	Net CO ₂ equivalent emissions/removals					
	1990			2011		
	2013 submission	2014 submission	Change, %	2013 submission	2014 submission	Change, %
A. Article 3.3 activities				9.7	-34.1	-450.4
A.1. Afforestation and Reforestation				-73.1	-119.2	63.1
A.2. Deforestation				82.8	85.1	2.7
B. Article 3.4 activities	5.237.8	5088.3	-2.9	-2711.1	-2739.4	1.0
B.1. Forest Management				-6313.6	-6180.0	-2.1
B.2. Cropland Management	5.053.9	4844.7	-4.1	3367.8	3196.1	-5.1
B.3. Grazing Land Management	184.0	177.4	-3.6	234.6	244.5	4.2

10.5.2 Review recommendations

The main recommendations for KP-LULUCF are included in Table 10.8.

Table 10.8 Recommendations from the UNFCCC review process concerning KP-LULUCF.			
CRF	ERT Comment	Denmark's response	Reference
2010 submission (Review report: http://unfccc.int/resource/docs/2011/arr/dnk.pdf)			
2011 submission (Review report: http://unfccc.int/resource/docs/2012/arr/dnk.pdf)			
2012 submission (Review report: http://unfccc.int/resource/docs/2013/arr/dnk.pdf)			
KP-LULUCF, Afforestation and reforestation – Paragraph 97	Provide further and verified information on the rationale for changing the method used to estimate the carbon pools	The methods for the estimations have not been changed since the NFI was started in 2002. Prior to this it was not possible to give precise estimates. The NFI and the data have been reviewed in 2012/2013. The updated land use matrix influences the overall estimates.	
KP-LULUCF, Afforestation and reforestation – Paragraph 98	Provide any available data on harvested areas and the associated estimation of emissions and removals	The method use to estimate forest carbon pools is based on stock change approach, and hence there is no need for direct estimation of harvested areas and related emissions/removals, as they are reflected directly in the measured carbon pools in the forests.	
KP-LULUCF, Forest management – Paragraph 100	Provide further and verified information on the rationale for changing the method used to estimate the carbon pools	The methods for the estimations have not been changed since the NFI was started in 2002. Prior to this it was not possible to give precise estimates. The NFI and the data have been reviewed in 2012/2013. The updated land use matrix influences the overall estimates.	
KP-LULUCF, Cropland management – Paragraph 102	Provide all relevant information on the selection of appropriate EFs	Further information has been implemented in the NIR. More detailed data with examples have been included in Annex3.F_LULUCF	Abnnex3.F_LULUCF Table 3.10-3.12

11 KP-LULUCF

11.1 General information

In the following text the abbreviations is used in accordance with definitions in the IPCC guidelines:

A:	Afforestation
R:	Reforestation
D:	Deforestation
FF:	Forest remaining Forest, areas remaining forest after 1990
FL:	Forest Land meeting the Danish definition of forests
CL:	Cropland
GL:	Grassland
SE:	Settlements
OL:	Other land, unclassified land
FM:	Forest Management, areas managed under article 3.4
CM:	Cropland Management, areas managed under article 3.4
GM:	Grazing land Management, areas managed under article 3.4

11.1.1 Definition of forest and any other criteria

For the estimation of anthropogenic emissions by sources and removals by sinks associated with afforestation (A), reforestation (R) and deforestation (D) since 1990 under Article 3.3 and forest management (FM) under Article 3.4 of the Kyoto Protocol, the following forest definition will be applied:

- Minimum values for tree crown cover: 10 % tree crown cover for forests.
- Minimum values for land area: 0.5 ha.
- Minimum value for tree height: trees must be able to reach a minimum height of 5 m in the site.

In addition, the forest area includes temporarily unstocked areas, smaller open areas in the forest needed for management purposes and fire breaks. Forests in national parks, reserves, or areas under special protection are included. Windbreaks and groves covering more than 0.5 ha and with a minimum width of 20 m are also considered as forests. Farmlands, fruit plantations for commercial purposes, orchards, gardens (houses and summer houses) are NOT included in the forest area. Willow plantations on agricultural soils for bioenergy purposes are included in Cropland (CL).

11.1.2 Elected activities under Article 3, paragraph 4, of the Kyoto Protocol

As regards the possibility of including in the first commitment period emissions and removals associated with land use, land-use change and forestry activities under Article 3.4 of the Kyoto Protocol, it has been decided to include emissions and removals from forest management (FM), cropland management (CM) and grazing land management (GM). The Danish territory covers mainland Denmark and Greenland and not the Faroe Islands.

The tables given below covers only the Danish territory and not data from Greenland and thus only data, which shall be included in the submission

to the European Union (EU). The Danish CRF and KP tables are named: DNM

For Greenland separate CRF and KP tables are produced, see Chapter 15. The Greenlandic tables are named: **GRL**.

The Greenlandic impact on the overall estimates is very low: <0,01 % and thus the figures given below can be regarded as very proximate values for both Denmark and Greenland.

The Danish and the Greenlandic CRF and KP tables are merged into one set of CRF and KP tables and named: **DKE**.

The Faroe Islands has not signed the Kyoto-Protocol and has therefore not submitted KP tables or been included in the Danish and the Greenlandic submission.

The national system has identified land areas associated with the activities under Article 3.4 of the Kyoto Protocol in accordance with definitions, modalities, rules and guidelines relating to land use, land-use change and forestry activities under the protocol by satellite monitoring, use of Land Parcel Information System (LPIS) from the EU subsidiary system as well as the Greenlandic subsidiary system, detailed crop information data on field level, soil mapping and sample plots from the national forest inventory (NFI).

Inventories of emissions and removals under Article 3.3 and Article 3.4 are prepared for 2011, and reported annually in 2013 together with the other greenhouse gas inventory information.

11.1.3 Description of how the definitions of each activity under Article 3.3 and each elected activity under Article 3.4 have been implemented and applied consistently over time

The definition of afforestation, reforestation and deforestation is in accordance with the GPG (IPCC 2003).

Afforestation or reforestation is identified when areas have wooded tree cover and fulfils the forest definition given above. The time of the A is given by the time of action - i.e. planting of trees. For R the time is given by the first spontaneous regeneration of tress, typically either by absence of management or by management inducing natural regeneration. All types of establishment of forest (A or R) is considered human induced, as all land area of Denmark is under management or as minimum specifically left for spontaneous revegetation. Regulations and support for A and R include natural revegetation as a specific method, often supplementing already existing forest areas. (Danish Forest and Nature Agency, Support for Sustainable Forestry - active until 2010.

<http://www.skovognatur.dk/Skov/Privat/Tilskud/Baeredygtig/>)

Deforestation is identified where areas in 1990 were covered by forest and where subsequent information (through remote sensing or NFI) is recorded to have another land use. Deforestation occurs for a number of reasons, e.g. nature restoration which in the period 1990 - 2011 have been the predominant reason. Other reasons can be urban or infrastructure development.

Temporarily unstocked areas - as integral part of forest management or as result of windthrow - which is expected to continue in forest management is not considered deforestation.

As for the forest management (Article 3.4) - the forest areas fulfilling the definition given above are included under this activity. All forest areas are considered managed due to the intense utilisation of the land area of Denmark. All inventories apply this approach. The Forest Act in Denmark gives the frame for most of the forest area ('Fredskov') - thereby ensuring continued forest cover - or by deforestation at least afforestation of a similar area or in most cases the double area. As described in Chapter 7 the changes in forest floor and mineral soils pools are not significant in the period observed (1990-2011) and are hence not considered being a source of emissions.

For Cropland and Grassland the area accounted for under Art. 3.4 has been estimated with the EO mapping combined with agricultural data from Statistics Denmark, Statistics Greenland and the EU agricultural subsidiary system. Only areas which are reported as CL and GL are included in the accounted area.

11.1.4 Description of precedence conditions and/or hierarchy among article 3.4 activities and how they have been consistently applied in determining how land was classified

All Forest activities have precedence, after this Cropland activities and then Grassland activities.

Afforestation has precedence. All land converted to forest are included as afforested area. Deforested areas are reported under D. The following categories in the Convention reporting are included under afforestation:

- 5A21 CL to A
- 5A22 GL to A
- 5A23 WE to A
- 5A24 SE to A
- 5A25 OL to A

Deforestation is estimated as:

- 5B21 to CL
- 5C21 to GL
- 5D21 to WE
- 5E21 to SE
- 5F21 to OL

FM activities are only related to:

- 5A1 Forest remaining Forest

CM activities are related to:

- 5B1 CL remaining CL
- 5B22 GL to CL
- 5B23 WE to CL (not occurring)
- 5B24 SE to CL
- 5B25 OL to CL
- 5D22 CL to WE
- 5E22 CL to SE
- 5F22 CL to OL (not occurring)

GM activities are related to:

- 5C1 GL remaining GL
- 5C22 CL to GL
- 5C23 WE to GL (not occurring)
- 5C24 SE to GL
- 5C25 OL to GL
- 5D23 GL to WE
- 5E23 GL to SE
- 5F23 GL to OL (not occurring)

No elected land has left land, which is accounted for. Land conversion between elected activities (FM, CM and GM) has been allowed. FL, CL and GM, which has been converted to WE and SE is still included in the accounted area. No land elected under 3.4 activities has been converted to Other Land. No Other land has been converted to land included in Art. 3.3 and 3.4 activities. As a consequence there has been a small increase in land, which is accounted for under Art. 3.3 and Art. 3.4 (Table 11.1) with two hectares from 1990 to 2012.

Table 11.1 The development in the different KP classes, which are included in the accounting (only Denmark) 1990 to 2012.

	1990	2008	2009	2010	2011	2012
AF	4 187	79 787	84 033	88 279	92 525	94 358
D	117	3 849	4 373	4 896	5 420	5 784
FM	543 856	540 124	539 600	539 076	538 553	538 189
CM	2 715 136	2 628 797	2 631 508	2 634 219	2 636 930	2 643 697
GM	401 414	315 123	308 248	301 373	294 498	285 900
Total area, Hectares	3 664 709	3 567 679	3 567 761	3 567 844	3 567 926	3 567 928

The Land Use matrix developed for the purpose of reporting Art. 3.3 and 3.4 activities for 2012 are shown in Table 11.2.

Table 11.2 Land Use matrix for art. 3.3 and 3.4 activities in 2012, in 1000 hectares.

To current inventory From previous inventory year		Article 3.3 activities		Article 3.4 activities				Other ⁽⁵⁾	Total area at the beginning of the current inventory year ⁽⁶⁾
		Afforestation and Reforestation	Deforestation	Forest Management (if elected)	Cropland Management (if elected)	Grazing Land Management (if elected)	Revegetation (if elected)		
		(kha)							
Article 3.3 activities	Afforestation and Reforestation	92.52	NO						92.52
	Deforestation		5.42						5.42
Article 3.4 activities	Forest Management (if elected)		0.36	538.19					538.55
	Cropland Management ⁽⁴⁾ (if elected)	1.73	NA		2,622.53	12.67	NA		2,636.93
	Grazing Land Management ⁽⁴⁾ (if elected)	0.11	NA		21.16	273.23	NA		294.50
	Revegetation ⁽⁴⁾ (if elected)	NA			NA	NA	NA		NA
Other ⁽⁵⁾		0.00	0.00	NO	0.00	NO	NA	737.62	737.63
Total area at the end of the current inventory year		94.36	5.78	538.19	2,643.70	285.90	NA	737.62	4,305.55

The above given information in the hierarchy between the Contention and the KP-LULUCF activities ensures that emission from activities under article 3.4 are not double counted under both article 3.3 and 3.4 activities.

11.2 Land-related information

11.2.1 Spatial assessment unit used for determining the areas of the units of land under Article 3.3

Afforestation and reforestation is identified where areas in 1990 were not covered by forest and where subsequent information (through remote sensing or NFI) is recorded to have forest cover fulfilling the forest definition.

Even though the definition for A and R refers to the time of establishment, there may be a slight time delay in the actual recording of the A/AR. This will be improved through more frequent land use mapping and improved methods for mapping in the coming years.

Deforestation is identified where areas at the beginning of the commitment period were covered by forest and where subsequent information (through remote sensing or NFI) is recorded to have another land use. The identification of the areas is in most cases supported by reports on e.g. nature restoration or establishment of settlements.

11.2.2 Methodology used to develop the land transition matrix

A land use/land cover map was produced for the Kyoto reference year 1990, 2005 and 2011 based on EO data for the forest land use. For mostly all other land uses the main data comes from detailed vector maps. These include data such as different vector layers from cadastral maps, road maps, wetland areas, agricultural land use data, vector layers of established wetlands, gravel maps etc. as well as aerial photos. The primary data used for the forest land use mapping is Landsat imagery mainly Landsat 5 (TM) and 7 (ETM+) data to classify and estimate the area and in combination with NFI data and other sources of data, including LiDAR data. The product is specified by a Minimum Mapping Unit (MMU) of 0.5 ha, a geometric accuracy of < 15 m RMS and a thematic accuracy of 90% +/- 5%.

The land use was allocated to the six major Kyoto classes: Forest, Cropland, Grassland, Wetland, Settlements, and Other. Highest priority was given to maps having the highest reliability in the production of the land use matrix. To avoid transition artefacts due minor updates in the precision of the vector maps a Minimum Mapping Unit (MMU) for land use change has been set to 0.5 ha which is the same as the elected Danish minimum MMU for forests in the Initial Report under the Kyoto protocol: [Initial Report](#)

In Chapter 7, Table 7.1 shows the overall development from 1990 to 2012. The preliminary result is an increase in the afforested area of 94 358 hectares, but also that deforestation has taken place of approximately 5 784 ha. Afforestation is mainly taking place on CL and GL. Areas, which are deforested, are mainly converted to CL and GL as the far major part of D is a conversion of existing Christmas tree fields to agricultural crops in rotation or permanent grass. Only to a little extend is forest converted to SE.

Since 1990 almost 31 169 hectares have been changed into SE and other infrastructures. No FF, CL and GL are converted into OL by definition.

Based upon the combination of the satellite image classified land use map and the combined vector layer of know information a full land use map for 1990, 2005 and 2011 was produced.

11.2.3 Maps and/or database to identify the geographical locations, and the system of identification codes for the geographical locations

The entire Danish territory except the Faroe Islands is included. This chapter includes only the territory of Denmark without Greenland. Denmark is reported as one unit and no sub-geographical locations are used.

Greenland is submitting a full separate NIR and CRF to be included in the submission to UNFCCC (Chapter 16).

11.3 Afforestation, Reforestation & Deforestation (ARD)

11.3.1 Methods for carbon stock change and GHG emission and removal estimates

For afforestation the carbon stock change in the period 1990 - 2011 is based both on the area of afforestation, the information on species composition from the Forest Census 2000 and from the NFI.

In the afforestation a steady increase in carbon stock is found. The species composition is based on the information from the 2000 Forest Census for the period 1990-2000. Subsequently the NFI provides information on the afforestation area and the carbon pools in these areas - up till 2007. The estimates for the carbon pools in the afforestation are similar to previous estimates, with a slight increase due to the new knowledge on species composition and average carbon stock in those areas based on the NFI data.

Carbon stock change caused by deforestation is estimated based on the deforested area and the mean values of carbon stock in the total forest area in the period 1990-2005. Based on analysis by aerial photographs and LiDAR data of the deforested areas in the period 2005-2011 is it estimated that 50 pct. of this deforestation is happening in very young forests or forests with low biomass (e.g. Christmas tree plantations or small open forests on the edge of agricultural land). This biomass carbon removed from these areas is estimated to be 15 t C/ha whereas the remaining deforested areas is assumed to have average carbon pools as the remaining forest area.

Where deforestation is taking place is the living and dead biomass removed and oxidized instantly. This includes also the litter layer in the forest. For the litter layer is further more included a N₂O-emission from nitrogen in the litter layer as well as changes in the C stock in mineral soils multiplied with a C:N ratio of 15 and a EF of 1.25 %. A large part of the deforestation is conversion of forest to create wetlands by removing the forest and closing the drainage system. For land converted to wetlands is assumed an average increase in the soil carbon stock of 0.5 ton C per ha per year which are and reported under mineral soils.

Further details are available in Johannsen et al. 2009.

11.3.2 Description of the methodologies and the underlying assumptions used

The climate in Denmark is cold and wet, which gives limitations to the growth of the forests and therefore afforestation in Denmark are on long rotations (>50 years) to give a reasonable amount of wood and wood products. Furthermore, the afforested areas are in many cases protected against deforestation. Therefore, afforested areas under article 3.3. will seldom be harvested during the commitment period. In the current submission is no estimates for "Units of land harvested since the beginning of the commitment period" in table 5(KP-I)A.1.2 given and stated as IE.

The basic information utilised to give the data for the emission estimates for units of land subjected to afforestation/reforestation is based on National

Forest Inventory (NFI) observations of stock change, specific related to the afforested areas. This will include all changes in carbon pools - also if affected by harvest - including thinnings of young stands.

Based on the NFI it will be possible - for the next reporting also to give some indications of the frequency of harvesting/thinning occurring on the afforested areas. Given the fact that the afforested area still is a relatively small part of the full forest area - there will be more uncertainty on the estimate related to afforested areas compared to the area of forest remaining forest.

11.3.3 Justification when omitting any carbon pool or GHG emissions/removals from ARD

When deforestation occurs it is assumed that all dead organic matter will be cleared. The actual amount depends on which type of forest is converted.

11.3.4 Information on whether or not indirect and natural GHG emissions and removals have been factored out

No factoring out has been performed in the emission and removal estimates.

11.3.5 Changes in data and methods since the previous submission (recalculations)

Minor recalculations have been made as updated values from the NFI have become available; also minor changes in the Land Use Matrix have occurred. See more in Chapter 7.3.7.

11.3.6 Uncertainty estimates

The uncertainty has been estimated based on the net changes in the different carbon stock with the IPCC Tier 1 methodology, Table 11.3 and Table 11.4.

The uncertainty in 2012 for Afforestation has been estimated to ± 9.7 Gg CO₂ equivalents and for Deforestation to ± 10.0 Gg CO₂ equivalents. The low uncertainty for afforestation is associated with that the figure is around zero.

Table 11.3 Uncertainty assessment for Afforestation for year 2012.

KP A.1.1 Afforestation and Reforestation		Emission	Activity	Emission	Combined	Total	Uncertainty
			data, %	factor, %	uncertainty	uncertainty, %	95 %, Gg CO ₂ eqv.
Area subject to the activity, ha		94 358					
Area of organic soils, ha		10 251					
Net CO2 emissions/ removals , Gg		37.8				18.8	7.1
Carbon stock change in above-ground biomass	Net change	12.4	15	8.5	17.2	17.2	2.1
	Net change	5.3	15	8.7	17.4	17.4	0.9
Carbon stock change in below-ground biomass		-1.4	15	1.0	15.0	15.0	0.2
Net carbon stock change in litter		3.2	15	3.3	15.4	15.4	0.5
Net carbon stock change in dead wood					52.2	52.2	
Net carbon stock change in soils	Mineral soils	-12.6	15	50.0			6.6
	Organic soils	3.5	15	50.0	52.2	52.2	1.8

Table 11.4 Uncertainty assessment for Deforestation. 1000 hectares and Gg CO₂ eqv.

KP A.2 Deforestation		Emission	Activity	Emis-	Combined	Total uncer-	Uncertainty 95
			data,	sion	uncertainty	tainty, %	%, Gg CO ₂ eqv.
			%	factor, %			
Area subject to the activity, ha		5 784					
Area of organic soils, ha		NO					
Net CO ₂ emissions/ removals, Gg CO ₂ (1)		109.3				12.1	13.2
Carbon stock change in above-ground biomass	Net change	20.6	15	8.5	17.2	17.2	3.5
	Net change				17.4	17.4	
Carbon stock change in below-ground biomass		4.4	15	8.7			0.8
Net carbon stock change in litter		4.8	15	1.0	15.0	15.0	0.7
Net carbon stock change in dead wood		0.5	15	3.3	15.4	15.4	0.1
Net carbon stock change in soils	Mine- ral soils	-0.5	15	50	52.2	52.2	0.3

1) The total emission of 109.3 in this table is slightly lower than the total emission from Deforestation this figure include only CO₂ emissions and not N₂O emissions from deforestation.

11.3.7 Information on other methodological issues

See Chapter 7.

11.3.8 The year of the onset of an activity, if after 2008

Not applicable.

11.4 Forest Management (FM)

11.4.1 Methods for carbon stock change and GHG emission and removal estimates

See Chapter 7 in LULUCF on "Forest remaining forest (5.A.1)".

11.4.2 Methodologies and the underlying assumptions

See Chapter 7 in LULUCF on "Forest remaining forest (5.A.1)".

11.4.3 Omission of pools from FM

No pools omitted.

11.4.4 Factoring out

No factoring out has been made.

11.4.5 Recalculations

Minor recalculations have been made due to updated values from the NFI on carbon stocks. Main change since last reporting is correction of years to which the data refer. This will cause a shift in reported changes. See more in Chapter 7.2.8.

11.4.6 Uncertainty estimates

The uncertainty has been estimated based on the net changes in the different carbon stock with the IPCC Tier 1 methodology. The uncertainty in 2012 for Forest Management has been estimated to ± 686.3 Gg CO₂ equivalents and ± 9.9 Gg CO₂ equivalents from drainage of organic soils in the forest.

Table 11.5 Uncertainty assessment for Forest Management.

KP B.1 Forest Management	Emission	Activity data, %	Emission factor, %	Combined uncertainty	Total uncertainty, %	Uncertainty 95 %, Gg CO ₂ eqv.	Emission
Area subject to the activity, ha		538 189					
Area of organic soils, ha		26 267					
Net CO ₂ emissions/ removals		-4491.4				10.3	461.1
Carbon stock change in above-ground biomass	Net change	-784.0	15	1.8	15.1	15.1	118.5
Carbon stock change in below-ground biomass	Net change	-176.4	15	2.0	15.1	15.1	26.7
Net carbon stock change in litter		-253.5	15	1.0	15.0	15.0	38.1
Net carbon stock change in dead wood		-20.0	15	3.3	15.4	15.4	3.1
Net carbon stock change in soils	Mineral soils	NO	15	50	52.2	-	-
	Organic soils	8.9	30	90	94.9	94.9	8.5

Table 11.6 Uncertainty assessment associated with drainage of forest soils.

KP-II 2 N ₂ O from drainage of soils	Emission	Activity data, %	Emission factor, %	Combined uncertainty	Total uncertainty, %	Uncertainty 95%, Gg CO ₂ eqv.
Area of drained soils, ha	294 691					
Emission	12.1				80.8	9.7
N ₂ O, Gg CO ₂ eqv.	12.1	30	75	80.8	80.8	9.7

11.4.7 Information on other methodological issues

See Chapter 7 in LULUCF on "Forest remaining forest (5.A.1)".

11.4.8 The year of the onset of an activity, if after 2008

Not applicable.

11.5 Cropland Management (CM)

11.5.1 Methods for carbon stock change and GHG emission and removal estimates

CL is subdivided in four classes: agricultural CL, wooded perennial fruit plantations, hedgerows and "other agricultural CL".

11.5.2 Methodologies and the underlying assumptions used

The area with agricultural CL are given as the agricultural area in Statistics Denmark for cereals, fodder crops, grass for seed, sugar beets, potatoes and other root crops.

Land converted from other Land use categories to CL is included under CL. Land converted to forest is reported under forest (AR). Land which according to the land use matrix is converted to WE and SE are still included in CM. Land conversion to OL is not allowed.

The same methodology as used in the Convention reporting, is used in the KP reporting.

11.5.3 Omission of pool from CM

Aboveground and belowground living biomass, litter and dead organic are only reported for perennial woody crops in accordance with IPCC GPG 2003. No litter and dead organic matter are reported under CL as this is seen as not occurring or as very insignificant as it is only related to the small area with fruit plantations and hedges. Only above- and belowground living biomasses for perennial fruit plantations, hedgerows and willow plantations for bioenergy purposes on agricultural land are therefore reported under cropland. Christmas trees are reported under FL.

11.5.4 Factoring out

The dramatic increase in the temperature in the latter years results in a higher turn-over rate of organic matter in soils leading to an increased emission from soils compared to pre 1990. For agricultural soils Denmark is using a dynamical temperature dependent model (Tier 3), which is expected to give the best estimate of the actual emission from soils compared to most other methods. If Denmark had used the default IPCC Tier 1 or 2 there would likely have been a *negative* factoring out, because the emission factor (EF) in these methods are based on long-term scientific data and thus not having the recent increase in temperatures included. Therefore by using the actual temperature in the Tier 3 no factoring out has been made.

11.5.5 Recalculations

Recalculations has been made due to an updated land use matrix and implementation of N₂O emissions from GL and WE converted to CL. The latter account for a very limited emission. No recalculations have been made for emissions from mineral soils, organic soils, hedgerows, perennial wooden crops and liming which are by far the major sources. The recalculations has reduced the overall emission from CL with 190 Gg CO₂ in the base year and with 180 Gg CO₂ in 2012.

11.5.6 Uncertainty estimates

The uncertainty has been estimated based on the net changes in the different carbon stock with the IPCC Tier 1 methodology. The uncertainty in 2012 for Cropland Management has been estimated to ± 2001.1 Gg CO₂ equivalents, ± 0.1 Gg CO₂ equivalents associated with disturbance from land use change and ± 83.2 Gg CO₂ equivalents from lime application, Table 11.7, 11.8 and 11.9, respectively.

Table 11.7 Uncertainty assessment for Cropland Management.

KP B.2 Cropland Management		Emission	Activity data, %	Emission factor, %	Combined uncertainty	Total uncertainty, %	Uncertainty 95 %, Gg CO ₂ eqv.
Area subject to the activity, ha		2 643 697					
Area of organic soils, ha		50 886					
Net CO ₂ emissions/ removals		2765.3				66.9	1850.6
Carbon stock change in above-ground biomass	Net change	53.8	10	50	51.0	51.0	27.5
	Net change				51.0	-	
Carbon stock change in below-ground biomass	Net change	IE	10	50			-
	Mineral soils				75.7	75.7	
Net carbon stock change in soils	Organic soils	160.1	10	75	90.6	90.6	121.2
	Organic soils	540.2	10	90			489.2

Table 11.8 Uncertainty assessment for N₂O associated with land use conversion.

KP-II 3 N ₂ O associated from disturbance of land use change	Emission	Activity data, %	Emission factor, %	Combined uncertainty	Total uncertainty, %	Uncertainty 95 % Gg CO ₂ eqv.
Land area converted, ha	5 438					
Emission						
N ₂ O Gg CO ₂ eqv	0.9				76.5	0.7

Table 11.9 Uncertainty assessment for lime consumption.

KP-II 4 Lime consumption	Emission	Activity data, %	Emission factor, %	Combined uncertainty	Total uncertainty, %	Uncertainty 95 % Gg CO ₂ eqv.
Total amount of lime applied	437 341					
Emission						
CO ₂	192.3				50.2	96.6

11.5.7 Information on other methodological issues

None.

11.5.8 The year of the onset of an activity. if after 2008

Not applicable.

11.6 Grazing land management (GM)

11.6.1 Methods for carbon stock change and GHG emission and removal estimates

Grazing land is defined as land used for permanent grazing as well as dry land not meeting the definitions for FF, CL, WE or SE. GL is subdivided into two types: Land strictly used for grazing and other grassland. Land used for grazing has no wooden vegetation whereas other grassland may have some wooden vegetation that does not meet the forest definition. The area with strict grazing land is remaining area between the grazing area and the grassland area in the land use matrix.

11.6.2 Description of the methodologies and the underlying assumptions used

As all the grazed grassland is more or less unimproved without fertiliser or limited fertilisation no changes in management practice has been applied. This is in accordance with IPCC GPG 2003 (3.4.1.2.1.2).

For land converted to GL and not purely free of wooden trees/bushes it is assumed that there is a living biomass of 2.200 kg DM per ha in above ground biomass and 6.160 kg DM per ha in below ground biomass (IPCC. 2003). In Grassland it is assumed that no changes in soil carbon stock in mineral soils are occurring. For organic soils is assumed an emission as reported in Section 7.

11.6.3 Factoring out

No factoring out has been made.

11.6.4 Recalculations

See section 10.5.5 as this also affect GM.

11.6.5 Uncertainty estimates

The uncertainty has been estimated based on the net changes in the different carbon stock with the IPCC Tier 1 methodology. The uncertainty in 2011 for Grassland Management has been estimated to ± 101.0 Gg CO₂ equivalents.

Table 11.10 Uncertainty assessment for Grassland Management.

KP B.3 Grassland Management		Emission	Activity data.	Emission factor.	Combined uncertainty	Total uncertainty.	Uncertainty 95 %
			%	%		%	Gg CO ₂ eqv.
Area subject to the activity		285 900					
Area of organic soils		17 171					
Net CO ₂ emissions/ removals							
Carbon stock change in above-ground biomass	Net change	523.2				44.7	233.7
Carbon stock change in below-ground biomass	Net change				51.0	51.0	
Net carbon stock change in soils	Mineral soils	119.0	10	50			60.7
	Organic soils	IE	10	50	51.0	-	-
		2.2	10	75	75.7	75.7	1.7

11.6.6 Information on other methodological issues

None.

11.6.7 The year of the onset of an activity, if after 2008

Not applicable.

11.7 Article 3.3

11.7.1 Information that demonstrates that activities under Article 3.3 began on or after 1 January 1990 and before 31 December 2012 and are direct human-induced

The land use mapping in 1990, 2005, 2011 and 2011 is the documentation that activities under Article 3.3 began after 1.1.1990. As all land area is under management all changes are evaluated as direct human induced. This also includes A and R, which are based on approved methods of establishing new forest - both planting and natural revegetation. In some cases the absence of removal of tree growth is an easy and cheap method for establishing new forest. Hence this method has also been supported through public support for establishment of new forest areas.

11.7.2 Information on how harvesting or forest disturbance that is followed by the re-establishment of forest is distinguished from deforestation

Deforestation is detected by analysis of satellite images. Furthermore deforestation of larger areas is confirmed by e.g. projects on nature restoration. Temporarily unstocked areas are typically located within larger forest areas and will in most cases be reforested within a period of 10 years as according to the Forest Act of Denmark, which applies to all Legal Forest Reserves (Fredsskov) and equals approximately 70 % of the total forest area. Clearcuts outside forests - e.g. small plantations of conifers on former cropland - is considered deforestation.

Most forest areas - including new forest areas - are subject to intermediate thinnings - harvesting of small trees. This is done with the purpose of reducing stem number and often to produce firewood or wood chips. Clearcuts of

new forest areas occurs in most cases first at maturity of the stand – after 50-100 years. A subset of the new forest area are managed as coppice like management. e.g. for production of Christmas trees.

11.7.3 Information on the size and geographical location of forest areas that have lost forest cover but which are not yet classified as deforested

This information will be available after the QA/QC analysis of the land use maps of 1990, 2005 and 2011, which will be performed during 2013.

11.7.4 Uncertainty on article 3.3 activities

A Tier 1 uncertainty analysis has been performed for Article 3.3 activities according to the GPG 2000. In total, the overall uncertainty in the year has been estimated to 50.94 %.

Table 11.11 Uncertainty assessment for Article 3.3. activities inclusive trend uncertainty.

	Year 2012 emission Gg CO ₂ eqv.
A. Article 3.3 activities	148.03
KP A.1.1 Afforestation and Reforestation	
KP A.2 Deforestation	37.83
Overall uncertainty in the year	
Total uncertainties (%)	109.26

11.8 Article 3.4

11.8.1 Information that demonstrates that activities under Article 3.4 have occurred since 1 January 1990 and are human-induced

Forest Management

In FM all forest area is under management and changes in carbon stock are hence seen as human induced. The baseline for 1990 is estimated as documented in Johannsen et al. 2009.

Cropland Management

Since 1990 major changes in Danish Agriculture has taken place. Due to environmental demands for “green crops during winter” the previous major crop, spring barley, has been replaced by primarily winter wheat. Furthermore, a ban on field burning was implemented in January 1990 (Executive order NO. 142 of 08/03/1989). This has reduced the burning of field residues, which were widely occurring until then. Furthermore, as part of reducing the leaching of nitrogen, executive order NO. 624 of 15/07/1997 demands of the farmers that a certain percentage of the area shall be grown with an extra crop after harvest of annual crops. Currently about eight per cent of the agricultural area is having an extra crop. From 2003 agricultural areas has been taken out of rotation due to demanded borders along watersheds to protect the watersheds.

Grassland Management

No specific activities have taken place in Grassland to increase or decrease the carbon stock. GM was elected so that all human induced activities affecting the carbon stock in the landscape are included in the Danish commitments under the Kyoto Protocol. Furthermore, it is very difficult to distinguish between activities in CM and GM in the heterogenic patchy Danish landscape.

11.8.2 Information relating to Cropland Management. Grazing Land Management and Revegetation, if elected, for the base year

No further information is available.

11.8.3 Information relating to Forest Management

No further information is available.

11.8.4 Uncertainty on article 3.4 activities

An Tier 1 uncertainty analysis has been performed for Article 3.3 activities according to the GPG 2000. In total the overall uncertainty in the year has been estimated to 170.50 % and the trend uncertainty to 117.52 %.

Table 11.12 Uncertainty assessment for Article 3.4. activities inclusive trend uncertainty.

	Base year 1990 emission	Year 2012 emission	Uncertainty in trend in national emissions intro- duced by emission factor uncertainty	Uncertainty in trend in national emissions intro- duced by activity data uncertainty	Uncertainty introduced into the trend in total national emissions
	Gg CO ₂ eqv.	Gg CO ₂ eqv.	%	%	%
B. Article 3.4 activities	5,088.29	-998.39			
KP B.1 Forest Management	NA	-4,491.37	NA	-18.72	NA
KP B.2 Cropland Management	4,221.74	2,765.31	52.53	7.69	53.09
KP B.3 Grassland Management	177.39	523.16	8.22	1.45	8.35
KP-II 2 N ₂ O from drainage of soils	NA	12.07	NA	0.10	NA
KP-II 3 N ₂ O associated from disturbance of land use change	0.26	0.94	0.01	0.00	0.02
KP-II 4 Lime consumption	622.92	192.30	3.09	0.27	3.10
Total uncertainties	Overall uncertainty in the year (%):				317.39

11.9 Other information

11.9.1 Key category analysis for Article 3.3 activities and any elected activities under Article 3.4

According to the IPCC Good Practice Guidance for LULUCF a category that is identified as key in the UNFCCC inventory should also be considered key under the Kyoto Protocol (IPCC GPG. 2003).

In 2012 the following LULUCF categories were identified as key categories at the level in the UNFCCC reporting:

- Forest land remaining forest land.
- Cropland remaining cropland – living biomass
- Cropland remaining cropland – organic soils
- Cropland remaining cropland – mineral soils
- Grassland remaining grassland – living biomass

According to Table 5.4.4 in the IPCC GPG for LULUCF this means that the following Kyoto Protocol activities are initially considered key.

Table 11.13 Relationship between activities in the UNFCCC LULUCF and the KP-LULUCF.

LULUCF activity	KP-LULUCF activities
Forest land remaining forest land	FM, GM, CM
Land converted to forest land	AR
Cropland remaining cropland	CM
Grassland remaining grassland	GM

For Denmark the relevant KP-LULUCF activity corresponding to forest land remaining forest land identified as being a key category in the UNFCCC reporting is FM. For land converted to forest afforestation/reforestation is a key category. For cropland remaining cropland the relevant KP-LULUCF activity is CM. For grassland remaining grassland the relevant KP-LULUCF activity is GM.

Therefore AR, FM, CM and GM are considered key categories in the Danish KP-LULUCF inventory.

For the full list of identified key categories please refer to Annex 1.

11.10 Information relating to Article 6

There are no Article 6 projects (Joint Implementation) on the Danish territory.

12 Information on accounting of Kyoto units

Referring to Decision 15/CMP.1 on Guidelines for the preparation of the information required under Articles 7 of the Kyoto Protocol (UNFCCC, 2006), this chapter and chapters 13, 14 and 15 include information and references to Denmark's and Greenland's annual non-inventory information under the Kyoto Protocol.

12.1 Background information

In accordance with paragraph 10 of the annex to Decision 15/CMP.1 information on emission reduction units, certified emission reductions, temporary certified emission reductions, long-term certified emission reductions, assigned amount units and removal units will be reported for the first calendar year in which these units will be transferred or acquired.

12.2 Summary of information reported in the SEF tables

The information required is contained in the UNFCCC Standard Electronic Format (SEF) application version 1.2.1.

12.3 Discrepancies and notifications

Annex 1 parties are also required to submit four reports according to paragraphs 12 to 16 of the annex to decision 15/CMP.1. These reports are:

- Paragraph 12 – List of discrepancies identified by the ITL.
- Paragraph 13/14 – List of notifications from the CDM Executive Board regarding ICERs.
- Paragraph 15 – List of non-replacement identified by the ITL.
- Paragraph 16 – List of invalid Kyoto units.

The list described in paragraph 12 is contained in Annex 6 as “Report – List of discrepancies identified by the ITL according to paragraph 12 of the annex to decision 15/CMP.1”.

The lists described in paragraph 13-15 are not included in this NIR, as there are no tCERs or ICERs in the Danish Registry. For paragraph 16, the Danish Registry has yet to receive invalid Kyoto units. This also renders this list unnecessary to submit. The discrepancies have been found in the daily reconciliation and have all been solved by manual intervention by either the Danish Registry or the CITL/ITL depending on which stage the transaction was in.

12.4 Publicly accessible information

Information to be publically available from the SEF will be included in SEF 2012 Denmark. The SEF report will also be publically available on the Danish Business Authority website (<http://www.dba.erhvervsstyrelsen.dk/eu-ets-registry-kyoto>).

Other information that is required to be publically available can be found on the EUTL website: <http://ec.europa.eu/environment/ets/>.

This information includes information on each account as required in paragraph 45 of the annex to Decision 13/CMP.1. Please note that the contact information (paragraph 45 (d) and (e)) requires the consent of the account holder according to EU law. Thus, all of this information is not publically available.

Information required in paragraph 45 (c) of the annex to Decision 13/CMP.1 can be found at the Danish Business Authority webpage:

<http://www.dba.erhvervsstyrelsen.dk/eu-ets-registry-kyoto>

Information on article 6 projects is not available as Denmark to this date has not approved any Joint Implementation projects in Denmark.

12.5 Calculation of the commitment period reserve

The calculation of the Commitment Period Reserve (CPR) is based on the assigned amount of 276 838 955 tonnes of CO₂ equivalents (UNFCCC, 2007). Subsequently, the CPR calculated as 90 % of the assigned amount is 249 155 060 tonnes CO₂ equivalent, during the commitment period and has not changed since the Report of the review of the initial report of Denmark published on 2 November 2007 (UNFCCC, 2007). The commitment period reserve has not changed since the previous submission, as 100 % times the most recent inventory times five would amount to a higher value.

12.6 KP-LULUCF accounting

At the time of preparation for this report Denmark has issued 6 942 723 RMUs and net-source cancelled 597 294 units (552 195 RMUs and 45 099 AAUs).

Referring to the KP-LULUCF inventory the accounting quantity is 8 613 701 tonnes CO₂ equivalent as RMUs on the basis of activities in 2008-2012 under Articles 3.3 and 3.4 of the Kyoto Protocol.

The accounting of RMUs based on the 2013 and 2014 submissions will not begin until after publication of the review report from the review of the submissions.

Table 12.1 Information on accounting for activities under articles 3.3 and 3.4 of the Kyoto Protocol.

Greenhouse gas source and sink activities	Net emissions/-removals							Accounting Parameters	Accounting Quantity
	Base year	2008	2009	2010	2011	2012	Total		
	(Gg CO ₂ equivalent)								
A. Article 3.3 activities									
A.1. Afforestation and Reforestation		393.4	-216.2	-279.8	-119.2	37.8	-183.9		-183.9
A.2. Deforestation		81.3	80.9	82.4	85.1	110.2	439.8		439.8
B. Article 3.4 activities									
B.1. Forest Management		-6097.6	259.4	-3754.7	-6180.0	-4479.3	-20252.2		-1172.6
3.3 offset								255.9	-255.9
FM cap								916.7	-916.7
B.2. Cropland Management		4844.7	3768.0	2663.5	3388.5	3196.2	2957.8	24223.3	-8249.3
B.3. Grazing Land Management		177.6	236.0	223.4	214.8	245.7	524.5	887.9	556.5
Total		-1460.1	-2270.7	-1616.3	-1614.5	-1391.9			-8609.4

Table 12.2 shows the average accounting quantity for 2008-2012.

Table 12.2 Annual average accounting quantity, Gg CO₂ equivalent.

Average for 2008-2012	
Afforestation and Reforestation	-36,78
Deforestation	87,97
Forest Management	-234,52
Cropland Management	-1649,86
Grazing Land Management	111,31
Total	-1721,88

¹ Calculated as the FM cap plus the Article 3.3 offset divided by five.

12.7 References

EC, 2004: COMMISSION REGULATION (EC) No 2216/2004 of 21 December 2004 for a standardised and secured system of registries pursuant to Directive 2003/87/EC of the European Parliament and of the Council and Decision No 280/2004/EC of the European Parliament and of the Council.

Available at:

<http://eur-lex.europa.eu/LexUriServ/LexUriServ.do?uri=OJ:L:2004:386:0001:0077:EN:PDF>

UNFCCC, 2006: Report of the Conference of the Parties serving as the meeting of the Parties to the Kyoto Protocol on its first session, held at Montreal from 28 November to 10 December 2005. Available at:

<http://unfccc.int/resource/docs/2005/cmp1/eng/08a02.pdf>

UNFCCC, 2007: Report of the review of the initial report of Denmark. Available at: <http://unfccc.int/resource/docs/2007/irr/dnk.pdf>

13 Information on changes in the national system

Since the 2013 submission no changes have been made to the national system.

14 Information on changes in the national registry

The ETS operates in 30 countries: the 27 EU Member States plus Iceland, Liechtenstein and Norway. It covers CO₂ emissions from installations such as power stations, combustion plants, oil refineries and iron and steel works, as well as factories making cement, glass, lime, bricks, ceramics, pulp, paper and board.

The following changes to the national registry of Denmark have occurred in 2013:

Reporting Item	Description
15/CMP.1 annex II.E paragraph 32.(a) Change of name or contact	<p>The Danish Business Authority The Danish Kyoto Registry Dahlerups Pakhus Langelinie Allé 17 DK-2100 København Ø Telephone 1: +45 3529 1000 Telephone 2: +45 7220 0030 E-mail: co2register@erst.dk www.erhvervsstyrelsen.dk/kvoteregisteret http://danishbusinessauthority.dk/eu-ets-registry-kyoto</p> <p>The registry Staff has changed to: Registry Manager Ms. Susanne Petersen Phone: +45 3529 1884 e-mail: susbod@erst.dk</p> <p>Registry staff: Mr. Peter W. Bentzen Phone: +45 3529 1883 e-mail: petben@erst.dk</p> <p>Ms. Hanne Paulli Phone: +45 3529 1881 e-mail: hanpau@erst.dk</p> <p>Ms. Anita Smed Phone: +45 3529 1622 e-mail: anisme@erst.dk</p> <p>Mr Ulrik Barkentin Overby Phone: +45 3529 1636 e-mail: Ulrove@erst.dk</p>
15/CMP.1 annex II.E paragraph 32.(b) Change regarding cooperation arrangement	No change of cooperation arrangement occurred during the reported period.
15/CMP.1 annex II.E paragraph 32.(c) Change to database structure or the capacity of national registry	<p>An updated diagram of the database structure is attached as Annex A.</p> <p>Iteration 5 of the national registry released in January 2013 and Iteration 6 of the national registry released in June 2013 introduces changes in the structure of the database.</p> <p>Changes introduced in release 5 and 6 of the national registry were limited and only affected EU ETS functionality. No change was required to the database and application backup plan or to the disaster recovery plan.</p> <p>No change to the capacity of the national registry occurred during the reported period.</p>

Reporting Item	Description
15/CMP.1 annex II.E paragraph 32.(d) Change regarding conformance to technical standards	Changes introduced in release 5 and 6 of the national registry were limited and only affected EU ETS functionality. However, each release of the registry is subject to both regression testing and tests related to new functionality. These tests also include thorough testing against the DES and were successfully carried out prior to the relevant major release of the version to Production (see Annex B). Annex H testing was carried out in February 2014 and the successful test report has been attached. No other change in the registry's conformance to the technical standards occurred for the reported period.
15/CMP.1 annex II.E paragraph 32.(e) Change to discrepancies procedures	No change of discrepancies procedures occurred during the reported period.
15/CMP.1 annex II.E paragraph 32.(f) Change regarding security	No change of security measures occurred during the reporting period The Danish Registry staff has made 3 changes to increase the security on the administration: <ol style="list-style-type: none"> 1. The staff members are not allowed to leave their cell phones un-attended. They are obliged to either have the phones with them, or the phone is locked in a cabinet. Only the staff member has the key to his cabinet. 2. All staff members have been undergoing an extended Police investigation. 3. The building hosting the registry administration has been secured. Only employees in The Danish Business Authority have access to the floor of the Danish Registry Staff.
15/CMP.1 annex II.E paragraph 32.(g) Change to list of publicly available information	EC: No change to the list of publicly available information occurred during the reporting period. DK: UK: http://danishbusinessauthority.dk/public_information http://danishbusinessauthority.dk/danish_emission_trading_registry DK: http://www.erhvervsstyrelsen.dk/offentlig_information http://www.erhvervsstyrelsen.dk/kyoto-registeret The publicly available information is updated on a monthly basis and confidential information is clearly marked as confidential. The information is available in English and Danish. Publicly available information concerning transactions, holdings and total volumes via the EUTL is concerned confidential, this information is not publicly available before year x+5, where "x" is the year of the transaction. Furthermore the following information is considered confidential: <ul style="list-style-type: none"> • Account identifier, • Representative's identifier, name, and contact information • Holdings of all accounts, • All transactions made, • The unique unit identification code of the allowances • The unique numeric value of the unit serial number of the Kyoto units held or affected by a transaction No public information is available concerning article-6 projects. Denmark has not approved any joint implementation projects in Denmark.
15/CMP.1 annex II.E paragraph 32.(h) Change of Internet address	No change of the registry internet address occurred during the reporting period. The internet address of the Danish registry is: https://ets-registry.webgate.ec.europa.eu/euregistry/DK/index.xhtml
15/CMP.1 annex II.E paragraph 32.(i) Change regarding data integrity measures	No change of data integrity measures occurred during the reporting period.
15/CMP.1 annex II.E paragraph 32.(j) Change regarding test results	Changes introduced in release 5 and 6 of the national registry were limited and only affected EU ETS functionality. Both regression testing and tests on the new functionality were successfully carried out prior to release of the version to Production. The site acceptance test was carried out by quality assurance consultants on behalf of and assisted by the European Commission; the report is attached as Annex B. Annex H testing was carried out in February 2014 and the successful test report has been attached.

Reporting Item	Description
The previous Annual Review recommendations	<p>In response to the previous Annual Review recommendations, the following document was submitted as a second addendum to Chapter 14: 'Information on changes in national registry' of the Annual Inventory Submission for the reporting year 2012.</p> <p>8. The assessor notes that Denmark is not fully reporting changes in the national registry related to change of test results. While the Party has resubmitted these items during the assessment cycle, the provided test report reveals a test plan which was of insufficient scope. This is evidenced by the limited number of Kyoto processes covered and absence of DES compliance demonstration through Annex H testing. Compliance with the DES requirements is essential to maintain confidence that national registry continues to perform the functions set out in the annex to decision 13/CMP.1 and the annex to decision 5/CMP.1. Therefore, the assessor strongly recommends that the Party test each release thoroughly against the DES as part of each major release cycle and provide the complete results in its annual NIR.</p> <p><u>Response:</u> The consolidated EU system of registries successfully completed a full certification procedure in June 2012. Notably, this procedure includes connectivity testing, connectivity reliability testing, distinctness testing and interoperability testing to demonstrate capacity and conformance to the Data Exchange Standard (DES). This included a full Annex H test. All tests were executed successfully and led to successful certification on 1 June 2012. The October 2012 release (version 4.0) was only a minor iteration and changes were limited to EU ETS functionality and had no impact on Kyoto Protocol functions in the registry. The test script previously provided reflects this.</p> <p>However, each major release of the registry is subject to both regression testing and tests related to new functionality. These tests include thorough testing against the DES and were successfully carried out prior to the relevant major release of the version to Production</p>
	<p>9. The assessor notes that Denmark is not fully reporting changes in the national registry related to the description of database structure. While the Party has resubmitted a simplified data model during the assessment cycle, the information contained within the model is not sufficient. This is evidenced by the lack of descriptions of each entity in the diagram and the omission of some diagram entities mandated in the Data Exchange Standard. The assessor recommends that following major changes, the party provide a data model which contains all DES required entities complete with descriptions in its annual NIR.</p> <p><u>Response:</u> The complete description of the consolidated registry was provided in the common readiness documentation and specific readiness documentation for the national registry of EU and all consolidating national registries. Since the successful certification of the registry on 1 June 2012, Iteration 4 of the registry, introduced in October 2012, added a limited number of new entities, none of them relating to DES entities.</p> <p>A data model was attached which more clearly shows the relevant entities "RECONCILIATIONS", "NOTIFICATIONS", "RESPONSES", "INTERNAL AUDIT LOG" and "MESSAGE LOG." As specified in the DES (Section VII. Data Logging Specifications/E. Message Archive), a copy of messages sent and received is stored in standalone files in one of two managed servers in the hosting environment. For that reason, the Message Archive is not shown in the model. The "MESSAGE LOG" object holds the location of the entire message, for each Message_ID.</p> <p>Since the successful certification of the registry on 1 June 2012, there has been no change in the capacity of the registry or change of its infrastructure.</p> <p>The recommendations is implemented and the requested reports will be provided by EC</p>

The mentioned Annex A and Annex B contains confidential information and is therefore not part of the NIR. The information has been submitted to the UNFCCC as confidential.

15 Information on the minimization of adverse impacts in accordance with Article 3, paragraph 14

No changes have occurred since the information reported in NIR 2011.

16 Methodology applied for the greenhouse gas inventory for Greenland

16.1 Introduction

The following sections contain a report of Greenland's part of the National Inventory Report (NIR) 2014. The structure of the report follows the UNFCCC guidelines on reporting and review (UNFCCC, 2002).

The report is to a far extent structured according to the recommended outline provided by the UNFCCC secretariat.

Previous to 2010 the greenhouse gas (GHG) inventory and this report were completed exclusively by The Danish Centre for Environment and Energy, Aarhus University (DCE), with input from the Environmental and Nature Protection Agency (APA), Ministry of Domestic Affairs, Nature and Environment (current Ministry of Environment and Nature).

In 2008 an energy statistic was officially initiated at Statistics Greenland with the intention to "... create an important tool, which in regard to political and economical priorities, can contribute to the identification of efforts on energy matters..." and which "... in regard to environmental aspects will create a basis for assessing the development in regard to Greenland's meetings of the Kyoto protocol ...". The first results on the new energy statistics, covering the period 2004-2007, were published in November 2008.

The GHG inventory submitted in April 2014 is completed by Statistics Greenland and the Ministry of Environment and Nature, Greenland Government, with technical support from DCE. This report on methodology is written by Statistics Greenland with assistance from the Ministry of Environment and Nature and documental support by DCE.

The annual emission inventories for Greenland for the years 1990-2012, are reported in the full CRF format.

The GHG's reported are:

- | | |
|------------------------|------------------|
| • Carbon dioxide | CO ₂ |
| • Methane | CH ₄ |
| • Nitrous Oxide | N ₂ O |
| • Hydrofluorocarbons | HFCs |
| • Perfluorocarbons | PFCs |
| • Sulphur hexafluoride | SF ₆ |

16.1.1 A description of the institutional arrangement for inventory preparation

Statistics Greenland and The Greenland Ministry of Environment and Nature are responsible for the annual preparation of the Greenlandic contribution to the National Inventory Report and the GHG inventories in the Common Reporting Format in accordance with the UNFCCC Guidelines. Statistics Greenland provides the data to DCE. DCE is responsible for ag-

gregating the Danish and Greenlandic CRF submissions and reporting the aggregated CRF and the National Inventory Report to the UNFCCC.

The inventory for LULUCF and KP-LULUCF is carried out by DCE and the documentation of the inventory (Sections 16.7 and 16.11) is completed by the Danish LULUCF experts.

Formerly, the provision of data was on a voluntary basis, but a formal contract between DCE and the Greenland Government came in place for the 2009 GHG inventory report.

The work concerning the annual GHG emission inventory is carried out in co-operation with other Greenlandic ministries, research institutes, organisations and companies:

Statistics Greenland (Ministry of Finance)

Annual energy statistics in a format suitable for the emission inventory work and fuel-use data for the large combustion plants. Since 2009 annual survey on emissions of F-gases.

Agricultural Advisory Service (Ministry of Fisheries, Hunting and Agriculture)

Background data on cropland and grassland, and statistics on livestock (sheep and reindeer).

Ministry of Domestic Affairs, Nature and Environment

Data on waste and emissions of F-gases. Annual Survey carried out by the Ministry of Domestic Affairs, Nature and Environment until 2008 and by Statistics Greenland from 2009 and onwards.

Ministry of Fisheries, Hunting and Agriculture and the Greenlandic Arboretum

Background data on forestry.

Greenland Airport Authority (Ministry of Health and Infrastructure)

Statistics on domestic flights and foreign flights to and from Greenland.

16.1.2 Brief description of the process of inventory preparation - data collection, data processing, data storage

The background data (activity data and emission factors) for estimation of the Greenlandic emission inventories is collected and stored in central databases at Statistics Greenland. The databases are in SAS format and handled with software from the SAS Institute Inc. The SAS programs are designed by Statistics Greenland. The methodologies and data sources used for the different sectors are described briefly in Section 16.1.4 and more in depth in Sections 16.3 to 16.8 and Section 16.11.

The material is placed on servers at Statistics Greenland. The servers are subject to routine backup services. Material, which have been backed up is archived safely.

16.1.3 General description of methodologies and data sources used

The GHG inventory for Greenland includes the following sectors:

- Energy sector
- Industrial processes
- Solvent and other product use

- Agriculture
- Land Use, Land-use Change and Forestry
- Waste
- KP-LULUCF

The applied methodologies follow the IPCC Guidelines and IPCC Good Practice Guidance. In some cases the methodology is identical to the methodology applied in the Danish inventory, however, the availability of data – especially site specific data – do not allow the same methodology to be used for all the sectors. The brief methodological description is included below for the different sectors. More thorough descriptions are included in Sections 16.3-16.8 and 16.11.

Energy sector

Fuel combustion

The Greenlandic emission inventory for fuel combustion has been performed according to the IPCC tier 1 methodology. The inventory is based on activity data from the Greenlandic energy statistics and on emission factors for different fuels, plants and sectors.

Total fuel combustion is based on data from Polaroil, Statoil and Malik Supply A/S. Polaroil imports fuel and distributes fuel in all parts of Greenland. Statoil imports and distributes fuel in Kangerlussuaq. Malik Supply A/S, a Danish company, re-distributes fuel bought from Polaroil to Greenlandic trawlers, ships etc. By using detailed data from Polaroil, Statoil and Malik Supply A/S it is possible to determine total import, total export, total international bunkers and total domestic fuel combustion.

Total domestic fuel combustion is then divided into sectors and private households by using data from a survey on energy consumption, company specific sales data from Polaroil and local fuel distributors, company tax accountings, municipality and the Government of Greenland accountings, and by estimation.

Fuel combustion in private households is estimated using detailed information from a number of local fuel distributors. Fuel deliveries are registered by buildings. In Greenland each building has a unique number registered in the Greenlandic Area Register (NIN). By combining the NIN-register and the GER-register (see above) with statistics on housing and population each building is labelled *private household* or located to a sector describing the main activity in the building. This new building-sector register, completed annually, is used extensively to determine the buyer of fuel delivered by Polaroil or local fuel distributors.

Fuel combustion in road traffic is based on a model designed by Statistics Greenland. The model contains data on the vehicle stock obtained from the Greenland Police Department's register on engine data. The vehicles are divided into broad categories of type i.e. personal car, lorry, taxi, truck, ambulance, motorbike etc. Each category is assigned with ratios on fuel type and mileage. Input data on mileage is derived from an annual survey among businesses and private road traffic since 2008. Each vehicle is divided in business categories or labelled *private vehicle* according to the owner. For each group the emissions are estimated by combining vehicle and annual mileage numbers with standard emission factors according to the type of fuel. The model does not take cold start or hot engines into account.

For air traffic annual emissions are based on activity data from Air Greenland A/S and sales data from the Greenland Airport Authority. For navigation, ferries and freight, annual emissions are based on activity data from Royal Arctic Line A/S (freight), Royal Arctic Tankers A/S (freight), Royal Arctic Bygdeservice A/S (freight/passengers), and Arctic Umiaq Line A/S (passengers) and the liquidated Assartuivik A/S (passengers).

For further information please refer to Section 16.3.

Fugitive emission

Greenland has no coal mines, no off-shore activities, no oil refineries, no natural gas transmission or distribution. For that reason there have been no fugitive emissions from such activities in 1990-2009. However in 2010 a scottish company initiated a search for oil along the westcoast of Greenland. Three wells were drilled and tested in 2010. Five wells in 2011. The oil company reported a total fugitive emission of 44 tonnes CO₂-eqv methane from all three wells in 2010. This information has been recognised as the total fugitive emission of CH₄ from fuels in 2010, whereas the calculation of fugitive emission of CO₂ and N₂O in 2010 and all three gases in 2011 has been based on IPCC Guideline emission factors, see Section 16.3. There were no oil exploration activities in 2012.

Furthermore, some fugitive emission occurs in the distribution of fuel e.g. when refuelling from ships to on-shore tanks, onshore loading of fuel to ships and offshore loading of ships. The emission would only be in the form of NMVOC. The fugitive emission from loading/unloading of ships is currently not estimated.

Industrial processes

Mineral products

CO₂ emissions occur from limestone and dolomite use, road paving with asphalt and asphalt roofing. Import statistics of asphalt and limestone are used as activity data for estimating the emissions.

Chemical industry

Greenland has no chemical industry.

Metal production

Greenland has no metal production.

Other production

There are several manufacturers of fish products and one tannery. Emissions of NMVOC are estimated, but there are no emissions of greenhouse gases occurring.

F-gases

Greenland has no production of halocarbons or SF₆. Data on consumption of F-gases (HFCs and SF₆) are obtained from an annual survey on consumption of halocarbons and SF₆ conducted by Statistics Greenland. Information on emission of industrial gases is available from 1995 onwards. Greenland has no consumption of PFCs.

For further information on the methodology for calculating emissions from industrial processes please refer to Section 16.4.

Solvent and other product use

The emission estimates for solvent and other product use are prepared by using import statistics of pure chemicals that fits the criteria for being considered a NMVOC compound. Additionally import statistics are used for products containing NMVOC's. The NMVOC emission is then calculated in to a CO₂ emission by using a standard value for carbon content in the NMVOC's. For further information see Section 16.5.

Agriculture

Enteric Fermentation Manure Management

Agriculture is sparse in Greenland due to climatic conditions. However sheep and reindeer are considered to contribute to emission of greenhouse gases. Enteric fermentation and manure management is assumed to contribute to emission of CH₄, and nitrogen excretion is assumed to contribute to emission of N₂O.

The emissions are given in CRF: Table 4 Sectoral Report for Agriculture and Table 4.A, 4.B(a), 4.B(b) and 4.D Sectoral Background Data for Agriculture. The calculation of emissions from the agricultural sector is based on methods described in the IPCC Guidelines (IPCC, 1996) and the Good Practice Guidance (IPCC, 2000). Activity data for livestock is on a one year average basis from the agriculture statistics published by Statistics Greenland. Data concerning the land use and crop yield is obtained from the Agricultural Advisory Service.

Data concerning the feed consumption and nitrogen excretion from sheep is based on information from the Agricultural Advisory Service supplemented by data on imported feed. Data concerning the feed consumption and nitrogen excretion from reindeer is based on information from the Agricultural Advisory Service and information from an article on reindeer management in Greenland.

Emission of N₂O is closely related to the nitrogen balance. Thus, quite a lot of the activity data is related to the calculation of ammonia emission. National standards are used to estimate the amount of ammonia emission. When estimating the N₂O emission the IPCC standard value is used for all emission sources. The emission of CO₂ from Agricultural Soils is included in the LULUCF sector.

For a more thorough description of the methodology for the agricultural sector please refer to Section 16.6.

Land use, land-use change and forestry

Greenland is the world's largest non-continental island on the northern American continent between the Arctic Ocean and the North Atlantic Ocean, northeast of Canada. The northernmost point of Greenland, Cape Morris Jesup, is only 740 km from then North Pole. The southernmost point is Cape Farewell, which lies at about the same latitude as Oslo in Norway. Greenland is covering approx. 2,166,086 km². It has been estimated that 81 % is covered permanently with ice leaving only 410,449 km² ice free. The climate is Arctic to sub arctic with cool winters and cold summers. The capitol Nuuk is having an average temperature of 1.4°C.

Due to its cold climate the LULUCF sector is of minor importance in relation to the emission of green house gases. Only a very minor area is covered by forest of which the major part has been planted within the last 40

years. Cropland was introduced in year 2000 and grassland management within the last 30 years. The cold climate slows down the biological processes making all growth rates very low.

In total has the emission from the LULUC sector in 2012 been estimated to a net source of 1.32 Gg CO₂ equivalent or 0.2 % of the total Greenlandic emission.

Forest land

Greenland has a few forests, which may qualify to the FAO criteria of forest definitions. The major forest areas are:

A natural forest in the Qinnngua valley of 45 ha consisting mainly of *Betula Pubescens ssp. czerepanovii* which in the period 1990 to 2012 has had an average height of six meters and approx. 100 trees per ha. It is thus assumed that it has had the same biomass for the whole period.

187 ha other planted forest. The largest of this is an arboretum (a research area) where different species and origins of trees are investigated which are adaptable to the harsh climate.

Cropland

In 1990 no annual crops were grown in Greenland. In 2012 10.5 ha of cropland was used for annual crops. The primary production is potatoes. Potato fields are mainly managed by hand and primarily fens with a high content of organic matter which is used for this purpose. It is thus assumed that the IPCC standard emission factor for boreal/cold areas of five tonnes C pr ha can be used although it is probably an overestimation due to the cold climate and the current management practice.

Grassland

In total is 242,000 hectare reported as grassland. The grassland is located in mountainous areas used for grazing of sheep. Due to the global warming are there some smaller areas which have become improved fertilised grassland. The total area with improved grassland has increased from 490 ha in 1990 to 1,060 ha in 2012.

Wetlands

Reported area with wetlands consists only of water-reservoirs. Due to lack of methodology for methane emissions under arctic conditions has no emission estimates has been made which is in accordance with the IPCC GPG 2003 guidelines.

Settlements

The few settlements are mainly built on cliffs with very sparse vegetation. Hence it is assumed that no changes in C stock occur.

Other land

No emission estimates has been made since no data is available which is in accordance with IPCC GPG 2003 guidelines.

For a more thorough description of the methodology applied for LULUCF and KP-LULUCF please refer to Section 16.7 and 16.11.

Waste

Solid waste management

The solid waste management in Greenland can be divided in the following processes:

- Managed waste disposal combined with open burning.
- Unmanaged waste disposal combined with open burning.
- Waste incineration with energy recovery.
- Waste incineration without energy recovery.

Waste incineration with energy recovery is according to IPCC Guidelines included under the energy sector.

Information on amount of waste produced pr year, amount of waste treated in the different processes, distribution between household and commercial waste, composition of the household waste and commercial waste, respectively, are provided by the Ministry of Environment and Nature.

Wastewater handling

N₂O emission from human sewage is estimated. The calculation of the N₂O emission uses population data from Statistics Greenland and an estimate for average protein consumption combined with default values from the IPCC Guidelines. No emissions of CH₄ are assumed to occur.

For more information please refer to Section 16.8.

Memo Items

International Aviation Bunkers

Emissions from international aviation bunkers are considered to be of negligible importance. The Greenland Airport Authority has reported the annual amount of jet fuel loaded into foreign aircrafts including Danish aircrafts. However, it is not possible to distinguish between Danish aircrafts and other aircrafts. Since most foreign aircrafts by far are Danish the annual amount of jet fuel loaded into foreign aircrafts are therefore included as part of the IPCC category 1A3a Civil aviation.

International Marine Bunkers

Emissions from international marine bunkers are included from 2004 and onwards. Before 2004 international marine bunkers are considered to be of negligible importance.

KP-LULUCF

Regarding the possibility of including in the first commitment period emissions and removals associated with land use, land-use change and forestry activities under Article 3.4 of the Kyoto Protocol, Greenland as part of the Kingdom of Denmark has included emissions and removals from forest management (FM), cropland management (CM) and grazing land management (GM).

The national system has identified land areas associated with the activities under Article 3.4 of the Kyoto Protocol in accordance with definitions, modalities, rules and guidelines relating to land use, land-use change and forestry activities under the protocol. All land converted from other activities into Cropland and Grassland is accounted for. No land has been allowed to leave elected areas under art. 3.4, see Section 16.11 for further details.

16.1.4 Brief description of key categories

A key category analysis (KCA) for year 1990 and 2012 has been carried out in accordance with the IPCC Good Practice Guidance.

The categorisation used results in a total of 32 categories. In the level KCA for the inventory for 1990, five key categories were identified. In the KCA for 2012, six categories were identified as key categories due to both level and trend. Further, three categories were key categories due to the trend.

Of the six key sources due to level for the reporting year 2012 four are in the energy sector, of which CO₂ from liquid fuels excluding transport in the analysis contributes most with 74.0 % of the national total (this contribution and the percentage contributions in the following are results from the level KCA based on the absolute values of the emissions; this contribution as percentages may differ somewhat from the percentage used in the sectoral chapters). The remaining level key categories in the energy sector are all CO₂ from the transport sector. Civil aviation, road transportation and domestic navigation comprise respectively 7.6 %, 5.4 % and 4.9 % of the national total. The last key categories are N₂O from wastewater handling and the consumption of HFCs.

The trend assessment shows that CO₂ from from Grassland remaining grassland, direct N₂O emissions from agricultural soils and CO₂ emission from waste incineration are key categories to the trend.

The categorisation used, results, etc. are included in Section 16.12 (Annex 1).

16.1.5 Information on QA/QC plan including verification

A number of measures are in place to ensure the quality of the Greenlandic greenhouse gas inventory.

The general QC activities include:

- Check that data are correctly moved between data processing steps, e.g. it is ensured that the data are imported correctly from the emission spreadsheets/databases to the CRF Reporter.
- The time series are analysed. Any large fluctuations are investigated and explained/corrected.
- The recalculations are analysed and the consistency of the emission estimates are verified.
- The completeness of the inventory is checked utilising the completeness checker incorporated in the CRF Reporter as well as expert knowledge from the inventory compilers.
- All references are checked and it is ensured that the citations are correct.

These types of QC checks are recommended as tier 1 QC checks in the IPCC Good Practice Guidance (IPCC, 2000).

The Greenlandic emission inventory is reviewed by Danish emission experts, who provide input to the Greenlandic inventory compilers on necessary improvements etc. This is done as a QA procedure. When the emission estimates are transferred to DCE, the quality control system of the Danish emission inventory is applied to the Greenlandic data.

All information related to the Greenlandic emission estimates are documented and archived securely annually. This is done in order to ensure that any part of the inventory can be reproduced at a later stage if necessary.

In addition source specific QA/QC activities are conducted; please see the associated paragraphs in the sectoral chapters.

16.1.6 General uncertainty evaluation

The uncertainty estimates are based on the Tier 1 methodology in the IPCC Good Practice Guidance (GPG) (IPCC, 2000). Uncertainty estimates for the following sectors are included in the current year: fuel combustion, fugitive emissions, industrial processes, solid waste, wastewater treatment and waste incineration, solvents and other product use, agriculture and LU-LUCF.

The uncertainties for the activity rates and emission factors are shown in Table 16.1.4. The estimated uncertainties for total GHG and for CO₂, CH₄, N₂O and F-gases are shown in Table 16.1.3. The base year for F-gases is 1995 and for all other sources the base year is 1990. The total Greenlandic GHG emission is estimated with an uncertainty of $\pm 3.8\%$ and the trend in GHG emission since 1990 has been estimated to be $-6.1\% \pm 2.7\%$ -age points. The GHG uncertainty estimates do not take into account the uncertainty of the GWP factors.

The uncertainty on CO₂ and N₂O from liquid fuels in fuel combustion, N₂O emission from waste water treatment and CH₄ emission from enteric fermentation are the largest sources of uncertainty for the Greenlandic GHG inventory. The result is skewed by the fact that more than 90 % of the Greenlandic Greenhouse gas emission is from fuel combustion of liquid fuels.

Table 16.1.3 Uncertainties 1990-2012.

	Uncertainty [%]	Trend [%]	Uncertainty in trend [%-age points]
GHG	± 3.8	-6.1	± 2.7
CO ₂	± 2.8	-7.1	± 2.6
CH ₄	± 56	-7.8	± 8.8
N ₂ O	± 82	-10.1	± 26
F-gases	± 51	+11 808	$\pm 5\ 250$

Table 16.1.4 Uncertainty rates for each emission source.

IPCC Source category	Gas	Base year	Year t	Activity data	Emission factor
		emission	emission	uncertainty	uncertainty
		Gg CO ₂ eqv	Gg CO ₂ eqv	%	%
1A Liquid fuels	CO ₂	613	562	2	2
1A Municipal waste	CO ₂	2	7	2	25
1A Liquid fuels	CH ₄	1	1	2	100
1A Municipal waste	CH ₄	0	0	2	100
1A Biomass	CH ₄	0	0	2	100
1A Liquid fuels	N ₂ O	2	2	2	500
1A Municipal waste	N ₂ O	0	0	2	500
1A Biomass	N ₂ O	0	0	2	200
1B2 Oil Exploration	CO ₂	0	0	2	1 000
1B2 Oil Exploration	CH ₄	0	0	2	1 000
1B2 Oil Exploration	N ₂ O	0	0	2	1 000
2A3 Limestone and dolomite use	CO ₂	0	0	5	5
2A5 Asphalt roofing	CO ₂	0	0	5	25
2A6 Road paving with asphalt	CO ₂	0	0	5	25
2F Consumption of HFC	HFC	0	7	10	50
2F Consumption of SF ₆	SF ₆	0	0	10	50
3A Paint application	CO ₂	0	0	10	15
3B Degreasing and dry cleaning	CO ₂	0	0	10	15
3C Chemical products, manufacturing and processing	CO ₂	0	0	10	15
3D5 Other	CO ₂	0	0	10	20
4A Enteric Fermentation	CH ₄	6	5	10	100
4B Manure Management	CH ₄	0	0	10	100
4B Manure Management	N ₂ O	1	1	10	100
4D1 Direct N ₂ O emissions from agricultural soils	N ₂ O	0	1	20	50
4D2 Pasture range and paddock	N ₂ O	1	1	20	25
4D3 Indirect N ₂ O emissions from agricultural soils	N ₂ O	1	1	20	50
5A Forest	CO ₂	0	0	5	50
5B Cropland	CO ₂	0	0	5	50
5C Grassland	CO ₂	0	1	5	50
6A Solid Waste Disposal on Land	CH ₄	4	4	10	100
6B Wastewater Handling	N ₂ O	15	12	30	100
6C Waste incineration	CO ₂	3	3	10	25
6C Waste incineration	CH ₄	2	2	10	50
6C Waste incineration	N ₂ O	1	1	10	100

16.1.7 General assessment of completeness

The present Greenlandic greenhouse gas emission inventory includes all major sources identified by the Revised IPCC Guidelines.

16.1.8 References

Ministry of Environment and Nature: Data on waste and ozone depleting substances and greenhouse gases HFCs, PFCs and SF₆.

Agricultural Advisory Service: Statistics on livestock (sheep and reindeer) and background data on land use (cropland and grassland).

Ministry of Fisheries, Hunting and Agriculture: Background data for Forestry.

IPCC, 1997: Revised 1996 IPCC Guidelines for National Greenhouse Gas inventories. Available at:

<http://www.ipcc-nggip.iges.or.jp/public/gl/invs6.htm> (15-04-2007).

IPCC, 2000: Good Practice Guidance and Uncertainty Management in National Greenhouse Gas Inventories. Available at:

<http://www.ipcc-nggip.iges.or.jp/public/gp/english/> (15-04-2007).

IPCC, 2006: 2006 IPCC Guidelines for National Greenhouse Gas Inventories, Prepared by the National Greenhouse Gas Inventories Programme, Eggleston H.S., Buendia L., Miwa K., Ngara T. and Tanabe K. (eds). Published: IGES, Japan. Available at:

<http://www.ipcc-nggip.iges.or.jp/public/2006gl/index.html>

Rasmussen, H.E., 1992: Reindeer Management in Southern Greenland (1992). Published in Danish as "Den sydgrønlandske rendrift" (1992).

Statistics Greenland: Energy Consumption in Greenland 2004-2012 published in Danish. Available at:

<http://www.stat.gl/publ/da/EN/201303/pdf/Grønlands%20energiforbrug%202012.pdf> as "Grønlands energiforbrug 2012" (01-11-2013).

Statistics Greenland, 2013: The Greenlandic energy statistics aggregated to SINK categories. Not published.

16.2 Trends in Greenhouse Gas Emissions

16.2.1 Description and interpretation of emission trends for aggregated greenhouse gas emission

The GHG emissions are estimated according to the IPCC guidelines and are aggregated into seven main sectors; Energy incl. Transport, Industrial Processes, Solvent and Other Product Use, Agriculture, LULUCF, and Waste. In Figure 16.2.3 and Figure 16.2.4 CO₂ emissions from fuel combustion in the Energy Sector is split into several sub-categories i.e. Energy Industries, Manufacturing Industries and Construction, Commercial and Institutional, Transport, Residential, Agriculture and Fishing.

The greenhouse gases include CO₂, CH₄, N₂O, HFCs, PFCs and SF₆. However, Greenland has no consumption of PFC. In 2012 total emission of greenhouse gases excluding LULUCF was 610.38 Gg CO₂ equivalent, and 611.70 Gg CO₂ equivalent including LULUCF.

Figure 16.2.1 shows total greenhouse gas emission in CO₂ equivalents from 1990 to 2012. The emissions are not corrected for temperature variations. CO₂ is the most important greenhouse gas. In 2012 CO₂ contributed to the total emission in CO₂ equivalent excluding LULUCF (Land Use and Land-Use Change and Forestry) with 93.8 %, followed by N₂O with 3.0 %, CH₄ 2.0 % and F-gases (HFCs and SF₆) with 1.2 %. Since 1990 these percentages have been increasing for F-gases, almost constant for CO₂ and falling for N₂O and CH₄.




Figure 16.2.1 Greenhouse gas emission in CO₂ equivalents, time series 1990-2012.

Stationary combustion plants and transport represent the largest categories. Energy excluding transport contributed to the total emission in CO₂ equivalents excluding LULUCF with 75.7 % in 2012; see Figure 16.2.2. Transport contributed with 18.1 %. Industrial processes, solvent and other products use, agriculture and waste contributed to the total emission in CO₂ equivalents with 6.2 %.

The net CO₂ emission forestry etc. is 0.2 % of the total emission in CO₂ equivalents in 2012. The total GHG emission in CO₂ equivalents excluding LULUCF has decreased by 6.2 % from 1990 to 2012 and decreased 6.1% including LULUCF. Comments on the overall trends etc. seen in Figure 16.2.1 and Figure 16.2.2 are given in the sections below on the individual greenhouse gases.




Figure 16.2.2 Greenhouse gas emission in CO₂ equivalents distributed on main sectors for 2012.

16.2.2 Description and interpretation of emission trends by gas

Carbon Dioxide

Emission of CO₂ accounted for 93.8 % of the total GHG emission in 2012. The largest source to the emission of CO₂ is the energy sector comprising Fuel Combustion (Sectoral Approach) and Fugitive Emissions from Fuels. In 2012 the energy sector contributed to 99.4 % of the total CO₂ emission.

In Figure 16.2.3 and Figure 16.2.4 CO₂ emissions are split into several subcategories i.e. Energy Industries, Manufacturing Industries and Construction, Transport, Commercial and Institutional, Residential, Agriculture and Fishing all subcategories from the energy sector. All remaining sectors are included in the subcategory Other including Waste incineration, Solvents and Other Product Use and Industrial Processes.

The largest source to the emission of CO₂; the energy sector includes combustion of fossil fuels like gas oil, gasoline, jet kerosene etc. From this sector Agriculture and Fishing (AFF) contributes with 23.2 % making AFF the largest contributor in 2012 followed by Residential with 19.6 % and Energy Industries and Transport both sharing 19.1 % of the total CO₂ emission in 2012.

Emissions from Energy Industries have been reduced a great deal in later years due to massive investments in hydro power plants. However, in 2010 and 2011 oil explorations were initiated along the west coast increasing fuel combustion and thus emissions in the Energy Industries to rise to the highest point ever. In 2012 oil exploration activities came to a standstill, which send energy combustion in Energy Industries to the lowest level ever in the time series since 1990.

Commercial and Institutions contributes with 9.4 % of the total CO₂ emission and Manufacturing Industries and Construction with 6.3 %. The category *Other* contributed with 3.3 % of the CO₂ emissions in 2012.

CO₂ emissions excluding LULUCF decreased by 20.2 % from 2011 to 2012. The main reason for this decrease was a standstill in the oil explorations in 2012. In 2012, the actual CO₂ emission was 7.2 % lower than the emission in 1990 excluding LULUCF.




Figure 16.2.3 CO₂ emissions, time series for 1990-2012.




Figure 16.2.4 CO₂ emissions, distribution according to the main sectors for 2012.

Nitrous oxide

Waste, particularly waste water handling is the most important N₂O emission source in 2012 contributing 68.5 % to the total N₂O emissions, see Figure 16.2.6. Agricultural activities contributed 20.8 % to the total N₂O emissions in 2012. Fuel combustion including transport contributed 10.7 %. Since 1990 total emission of N₂O has decreased by 10.1 %.

The N_2O emission from agriculture decreased during the early nineties due to a decrease in reindeer livestock from 1990 to 1994. Since 1995 the emission of N_2O has increased and decreased for shorter periods depending on changes in the livestock and the use of fertiliser. In 2008, the actual N_2O emission was double the emission in 1990, see Figure 16.2.5. The cause of this was a significant increase in the use of fertilisers in 2008.

Besides from a temporary increase in 2011 total N_2O emission was reduced in 2009-2010 and 2012 by 18.4 %, 2.1 %, and 7.6 % due to a fall in the amount of waste water handling from industrial fishing plants and reduced use of synthetic fertilisers in agricultural activities, see Figure 16.2.5.




Figure 16.2.5 N_2O emissions, time series for 1990-2012.




Figure 16.2.6 N_2O emissions, distribution according to the main sectors in 2012.

Methane

The largest sources of anthropogenic CH_4 emissions are agricultural activities contributing with 45.6 % of total CH_4 emission in 2012; see Figure 16.2.8. Waste handling contributes to 45.5 % of total emission and the energy sector to 9.0 % of total CH_4 emission in 2012. The emission from agriculture derives from enteric fermentation (97.7 %) and management of animal manure (2.3 %).

Since 1990 the overall number of sheep has increased, while the overall number of reindeer has decreased. From 1990 to 2012 the emission of CH_4 from agricultural activities has decreased by 10.5 %.

The emission of CH_4 from waste derives from solid waste disposal (70.9 %) and waste incineration (29.1 %). From 1990 to 2012 the emission of CH_4 from solid waste disposal has increased by 7.2 %, while emissions from waste incineration have decreased by 29.4 %. Overall emission of CH_4 from waste handling has decreased by 6.8 % from 1990 to 2012.




Figure 16.2.7 CH₄ emissions, time series for 1990-2012.




Figure 16.2.8 CH₄ emissions, distribution according to the main sectors in 2012.

HFCs, PFCs and SF₆

This part of the Greenlandic inventory only comprises a full data set for HFCs and SF₆ from 1995. Greenland has no consumption that leads to emission of PFCs. From 1995 to 2012 there has been a continuous and substantial increase in the contribution from F-gases calculated as the sum of emissions in CO₂ equivalents, see Figure 16.2.9. This increase is caused by and simultaneous with an increase in the emission of HFCs. For the time series 2004-2012 the increase is lower than for the years 1995 to 2004. The increase from 1995 to 2004 is 8,892 %. From 2004 to 2012 total emission increased by 32.4 %. SF₆ contributed to the F-gas sum in 1995 with 59.4 %. Environmental awareness and regulation of this gas under Danish law has reduced its use considerably since 1995. In 2012 the contribution from SF₆ to the emission of F-gases was only 0.04 %.

The use of HFCs has increased to a great extent. Today HFCs are by far the dominant F-gas, comprising 40.6 % in 1995, but 99.96 % in 2012. HFCs are mainly used as a refrigerant.




Figure 16.2.9 F-gas emissions, time series for 1990-2012.

16.2.3 Description and interpretation of emission trends by category

Energy

The emission of CO₂ from energy has decreased by 7.4 % from 1990 to 2012. Emissions decreased from 1990 until 1994 due to the implementation of the first hydro power plant. However, since 1994 combustion of fuel increased continuously causing emissions to increase as well. The reason for this increase was primarily higher demand for transportation and heating. Combustion of fuel may decrease in certain years due to milder temperatures. In 2010 and 2011 emissions increased significantly due to a significant increase in fuel combustion due to the initiation of oil exploration, which caused CO₂ emission from energy to rise by 14.6 % in 2010 and by 6.9 % in 2011. However, in 2012 oil exploration activities came to a standstill, while Greenland's fifth hydro power plant went into operation. This caused CO₂ emissions from energy to decrease by 20.3 % in 2012.

Emission of CH₄ has increased by 3.0 % from 1990 to 2012 primarily due to an increase in the use of fuel for transportation as well as in the energy recovery from waste. The CH₄ emission from the transport sector has increased by 54.4 % from 1990 to 2012, mainly due to increasing domestic aviation.

Emission of N₂O has increased by 5.3 % from 1990 to 2012.

Industrial processes

Emissions from industrial processes (consumption of halocarbons and SF₆) other than fuel combustion amount to 1.2 % of the total emission in CO₂ equivalents excluding LULUCF in 2012. The main source is consumptions of HFCs. Emission of F-gases have increased considerably since 1990.

Agriculture

The agricultural sector contributes with 1.6 % of the total GHG emissions excluding LULUCF in 2012, 45.8 % of the total CH₄ emission and 22.1 % of the total N₂O emission. The total emission from the sector has increased by 6.4 % from 1990 to 2012. This increase is due to a rise in the use of synthetic fertilisers. The number of reindeer has decreased from 6,000 heads in 1990 to 3,000 heads in 2012. The number of sheep has increased from 19,929 heads in 1990 to 20,107 heads in 2012. The CH₄ emission has decreased by 10.5 % from 1990 to 2012 primarily due to the fall in the number of reindeers. In the same period N₂O emission has increased by 46.2 % due to a significant increase in the use of fertilisers.

LULUCF

Emissions from the LULUCF sector amount to just 0.2 % of the total emission in CO₂ equivalents in 2012. Forests are assumed to be a sink for the whole period increasing from approximately zero in 1990 to 42.3 tonnes CO₂ in 2012. The emission from cropland is estimated to zero in 1990 as there were no cropland in Greenland in 1990 and a net source in 2012 of 48.1 tonnes CO₂. The emission from grassland has been estimated to 214 tonnes CO₂ in 1990 increasing to 1,316 tonnes CO₂ in 2012.

Waste

The waste sector contributes with 3.5 % of the total greenhouse gas emissions in 2012, 45.5 % of the total CH₄ emission and 73.2 % of the total N₂O emission. The total emission from the sector has decreased by 13.1 % from 1990 to 2012. This decrease is caused by a drop in the CH₄ emission from

waste incineration by 29.4 %, and a decrease in N₂O emission from waste water handling by 20.8 %.

Total GHG emission from waste incineration without energy recovery has decreased by 6.3 % from 1990 to 2012 due to an increasing amount of waste incineration with energy recovery and a decrease in waste water handling from industrial fishing plants in 2012. Emission from incinerated waste used for heat production is included in the 1A1 IPCC category Energy Industries.

16.2.4 Description and interpretation of emission trends for indirect greenhouse gases and SO₂

NO_x

The largest sources to emission of NO_x are AFF (Agriculture, Forestry and Fisheries) followed by Transport and combustion in Energy Industries (public power and district heating plants). The AFF-sector is the most contributing sector to the emission of NO_x. In 2012, 50.8 % of the Greenlandic emission of NO_x came from AFF-related activities.

The emission of NO_x from AFF varies from year to year. In recent years emission of NO_x from AFF has been relatively stable with a slightly decreasing tendency since 2000.

The emissions from transport obtain 26.5 % of total emissions in 2012. From 1990 to 2012 emission of NO_x from AFF has increased by 28.9 %, while emissions from transport have increased by 17.3 %. In the same period total emission of NO_x has increased by 15.6 %.

The emissions from energy industries obtain 6.9 % of total emission in 2012. The emission from energy industries have decreased by 39.3 % from 1990 to 2012. The decrease is due to a continuous substitution of fossil fuels with hydro power.

Emission of NO_x from waste handling obtains 0.9 % of total emission, see Figure 16.2.10.




Figure 16.2.10 NO_x emissions. Distribution according to the main sectors (2012), and time series (1990-2012).

CO

Mobile sources like transport and AFF (agriculture, forestry and fisheries) contribute significantly to the total emission of this pollutant. Transport is the largest contributor to the total CO emission, see Figure 16.2.11.

Total CO emission has increased by 43.8 % from 1990 to 2012, largely due to increasing emissions from road transportation and civil aviation. Emission

sions from energy industries have more than doubled from 1990 to 2012, while emissions from transport have nearly doubled since 1990.




Figure 16.2.11 CO emissions. Distribution according to the main sectors (2012), and time series (1990-2012).

NMVOC

The emissions of NMVOC originate from many different sources and can be divided into two main groups: incomplete combustion and evaporation. Road vehicles and other mobile sources such as national navigation vessels fishing vessels and off-road machinery are the main sources of NMVOC emissions from incomplete combustion processes. Road transportation and fishing vessels are the main contributors to this pollutant. Road transportation is included under transportation, which obtain 37.4 % of the total NMVOC emission in 2012. Fishing vessels are included under AFF (agriculture, forestry and fisheries), which obtain 31.8 % of total NMVOC emission in 2012, see Figure 16.2.12.

The evaporative emissions mainly originate from the use of solvents and the extraction, handling and storage of oil. Emissions from solvents and other product use have decreased by 12.5 % from 1990 to 2012.

The total anthropogenic emissions have increased by 24.6 % from 1990 to 2012, largely due to the increase in road transportation and AFF activities.




Figure 16.2.12 NMVOC emissions. Distribution according to the main sectors (2012), and time series (1990-2012).

SO₂

The main part of the SO₂ emission originates from the combustion of fossil fuels mainly gas oil in public power and district heating plants. From 1990 to 2012, total emission of SO₂ decreased by 10.7 %.

Emissions from AFF (Agriculture, Forestry and Fisheries) obtain 34.5 % of total SO₂ emission in 2012 followed by Energy Industries obtaining 23.6 % in 2012. Also emissions from other industrial combustion plants, non-

industrial combustion plants and mobile sources are important. Transport contributed with 13.6 % of total SO₂ emission in 2012.




Figure 16.2.13 SO₂ emissions. Distribution according to the main sectors (2012), and time series (1990-2012).

16.3 Energy (CRF sector 1)

16.3.1 Overview of sector

The emission of greenhouse gases from energy activities includes CO₂, CH₄ and N₂O emission from fuel combustion. In 2010 fugitive emission of CO₂, CH₄ and N₂O occurred for the first time due to the initiation of well drilling and testing for oil and gas. The emissions are reported in CRF Tables 1.A(a), 1.A(b), 1.A(c), 1.A(d) and 1.B. Furthermore, the emission of non-methane volatile organic compounds (NMVOC), NO_x, CO and SO₂ from fuel combustion is given in CRF Table 1.

Summary tables for the energy sector are shown below.

Table 16.3.1 CO₂ emission from the energy sector.

Greenhouse gas source and sink categories	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
	Gg										
1. Energy	614.6	600.0	586.0	536.6	487.1	525.1	587.0	607.4	586.4	584.4	657.0
A. Fuel Combustion (Sectoral Approach)	614.6	600.0	586.0	536.6	487.1	525.1	587.0	607.4	586.4	584.4	657.0
1. Energy Industries	180.4	175.2	171.1	154.8	138.5	119.6	120.4	127.3	125.3	127.4	130.8
2. Manufacturing Industries and Construction	26.0	25.2	24.6	22.2	19.8	43.2	43.8	45.5	39.4	45.2	47.5
3. Transport	95.1	94.6	92.6	86.3	80.0	87.9	91.8	95.7	100.1	103.5	104.8
4. Other Sectors	305.0	297.1	290.0	266.3	242.6	267.8	324.3	332.3	315.0	301.8	367.2
5. Other	8.1	7.9	7.7	7.0	6.2	6.5	6.5	6.5	6.6	6.6	6.6
B. Fugitive Emissions from Fuels	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
<i>continued</i>	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011
1. Energy	608.0	570.1	639.4	630.1	634.2	652.3	643.3	667.6	583.6	668.7	714.7
A. Fuel Combustion (Sectoral Approach)	608.0	570.1	639.4	630.1	634.2	652.3	643.3	667.6	583.6	668.7	714.7
1. Energy Industries	131.9	132.6	133.2	137.1	135.8	141.0	133.8	142.6	124.8	224.3	249.3
2. Manufacturing Industries and Construction	45.1	42.6	49.2	50.2	54.5	55.1	56.8	58.8	42.7	38.3	46.8
3. Transport	95.1	91.5	100.3	112.4	110.8	120.0	109.3	116.0	104.9	107.4	114.4
4. Other Sectors	329.2	296.8	350.2	322.9	325.8	326.7	335.8	340.4	295.3	274.6	283.1
5. Other	6.6	6.6	6.6	7.4	7.2	9.6	7.6	9.9	15.8	24.1	21.1
B. Fugitive Emissions from Fuels	NO	NO	NO	NO	NO	NO	NO	NO	NO	0.02	0.03
<i>continued</i>	2012										
1. Energy	569.3										
A. Fuel Combustion (Sectoral Approach)	569.3										
1. Energy Industries	109.6										
2. Manufacturing Industries and Construction	36.2										
3. Transport	109.6										
4. Other Sectors	298.4										
5. Other	15.5										
B. Fugitive Emissions from Fuels	NO										

Table 16.3.2 CH₄ emission from the energy sector.

Greenhouse gas source and sink categories	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
	Gg										
1. Energy	0.05	0.05	0.05	0.04	0.04	0.04	0.05	0.05	0.05	0.05	0.05
A. Fuel Combustion (Sectoral Approach)	0.05	0.05	0.05	0.04	0.04	0.04	0.05	0.05	0.05	0.05	0.05
1. Energy Industries	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
2. Manufacturing Industries and Construction	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
3. Transport	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.01	0.01	0.01	0.00
4. Other Sectors	0.04	0.03	0.03	0.03	0.03	0.03	0.03	0.04	0.03	0.03	0.04
5. Other	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
B. Fugitive Emissions from Fuels	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
<i>continued</i>	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011
1. Energy	0.05	0.05	0.05	0.06	0.06	0.06	0.06	0.06	0.06	0.05	0.06
A. Fuel Combustion (Sectoral Approach)	0.05	0.05	0.05	0.06	0.06	0.06	0.06	0.06	0.06	0.05	0.05
1. Energy Industries	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
2. Manufacturing Industries and Construction	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
3. Transport	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
4. Other Sectors	0.04	0.04	0.03	0.04	0.04	0.04	0.04	0.04	0.04	0.03	0.03
5. Other	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
B. Fugitive Emissions from Fuels	NO	NO	NO	NO	NO	NO	NO	NO	NO	0.002	0.001
<i>continued</i>	2012										
1. Energy	0.05										
A. Fuel Combustion (Sectoral Approach)	0.05										
1. Energy Industries	0.01										
2. Manufacturing Industries and Construction	0.00										
3. Transport	0.01										
4. Other Sectors	0.03										
5. Other	0.00										
B. Fugitive Emissions from Fuels	NO										

Table 16.3.3 N₂O emission from the energy sector.

Greenhouse gas source and sink categories	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
	Gg										
1. Energy	0.01	0.01	0.01	0.01	0.00	0.01	0.01	0.01	0.01	0.01	0.01
A. Fuel Combustion (Sectoral Approach)	0.01	0.01	0.01	0.01	0.00	0.01	0.01	0.01	0.01	0.01	0.01
1. Energy Industries	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
2. Manufacturing Industries and Construction	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
3. Transport	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
4. Other Sectors	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
5. Other	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
B. Fugitive Emissions from Fuels	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
<i>continued</i>	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011
1. Energy	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
A. Fuel Combustion (Sectoral Approach)	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
1. Energy Industries	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
2. Manufacturing Industries and Construction	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
3. Transport	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
4. Other Sectors	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
5. Other	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
B. Fugitive Emissions from Fuels	NO	NO	NO	NO	NO	NO	NO	NO	NO	0.00	0.00
<i>continued</i>	2012										
1. Energy	0.01										
A. Fuel Combustion (Sectoral Approach)	0.01										
1. Energy Industries	0.00										
2. Manufacturing Industries and Construction	0.00										
3. Transport	0.00										
4. Other Sectors	0.00										
5. Other	0.00										
B. Fugitive Emissions from Fuels	NO										

16.3.2 Source category description

In this section emission source categories, fuel consumption data and emission data are presented.

Activity data on fuel consumption is based on annual statistics on energy published by Statistics Greenland and information on waste incineration with energy recovery. The annual statistics on energy is divided into sectors according to the Greenlandic Business Register (GB2000). The register comprises 577 business categories. The official statistics on energy is published by aggregation into 34 categories.

In the Greenlandic emission database, all activity rates and emissions are based on the official statistics on energy. However, in order to fit the CRF format fuel consumption from the official statistics on energy is further aggregated into 15 sectors increased to 16 sectors in 2010 with the energy sub-sector "Manufacture of Solid Fuels and Other Energy Industries" that contains emissions from oil exploration.

Fuel combustion

In 2012, total fuel combustion was 8,014 TJ of which 7.826 TJ was liquid fossil fuels.




Figure 16.3.1 Fuel combustion rates, fossil fuels 2012 (Statistics Greenland).

In Greenland gas oil, kerosene and gasoline are used in fuel combustion. From 2010 fuel oil is also being imported and combusted in ships. Gas oil and kerosene are the most utilised fuels. Gas oil is used in power plants to produce electricity and heat, as well as in district heating, private households, industries and for transportation. In 2010 and 2011 the combustion of gas oil increased significantly due to oil explorations. Due to a standstill in oil explorations total fuel combustion decreased significantly in 2012.

Kerosene is primarily used in aviation, but also for heating in smaller settlements.

A time series on the consumption of Liquid Petrol Gas (LPG) was introduced for the first time in the 2013 inventory submission. However, the consumption of LPG amount to less than 1 % of the total fuel combustion, see Figure 16.3.2. It has been possible to construct a time series on LPG consumption running from 2004 and onwards.




Figure 16.3.2 Fuel combustion, 2012 (Statistics Greenland).

Time series on fuel consumption are presented in Figure 16.3.3. Total fuel consumption has decreased by 6.4 % from 1990 to 2012. This overall decrease in fuel consumption is entirely caused by a drop in the consumption

of liquid fossil by 8.1 %. Consumption of renewable waste-energy has increased continuously with a total increase of more than 300 % from 1990 to 2012.




Figure 16.3.3 Fuel consumption time series 1990-2012 (Statistics Greenland).

Fuel consumption is dominated by liquid fuels e.g. gas oil, kerosene and gasoline. In 2012 total fuel consumption consists of 98 % liquid fuels, 1 % solid fuels and 1 % biomass.

In 2012 Energy Industries accounted for 20 % of total fuel consumption. From 1990 to 1995 fuel consumption in Energy Industries decreased significantly due to the introduction of the first hydro power plant in 1993, and the introduction of burning waste to produce heat for district heating networks in 1989. Dependence on gas oil conversion decreased immediately. Nevertheless, from 1995 onwards consumption of gas oil once again increased due to the general economic development. In 2007 fuel consumption in Energy Industries decreased due to a relatively warm winter. Contrary to this, the winter in 2008 was relatively colder, which increased fuel consumption to produce heat. In 2009 hydro power productions increased further when a fourth plant was opened. Together with a relatively warm 2009 winter fuel consumption in Energy Industries decreased additionally. In 2010 and 2011 fuel consumption increased significantly due to oil explorations along the westcoast of Greenland. In 2012, most recently fuel consumption decreased significantly due to a standstill in the oil exploration and the opening of the fifth hydro power plant.

Fuel consumption in Agriculture, Forestry and Fisheries accounted for 23 % of total fuel consumption in 2012. Fuel consumption in this sector has increased since 2010. Before 2004, annual fuel combustion in this sector varied a great deal due to fluctuations in fishing activities from year to year. However, some uncertainty is expected in the 1990-2003 time series on fuel consumption in Agriculture, Forestry and Fisheries.

Residential fuel consumption also accounted for 19 % of total fuel consumption in 2012. Fluctuations in fuel consumption are largely a result of variation in outdoor temperatures from year to year, which also causes fluctuations in fuel consumption in Energy Industries.

For 2004-2012 Statistics Greenland has conducted statistics on energy including detailed information on fuel consumption divided into 33 business categories and private households; see Section 16.3.3.1. Compared to the new statistics on energy the historic construction of time series on fuel consumption in 1990-2003 was based on a much simpler method. Some uncertainty is therefore to be expected in the 1990-2003 time series on sector-divided fuel consumption.

Fugitive emissions from fuels

Greenland has no coal mines, no off-shore activities, no oil refineries, no natural gas transmission or distribution. For that reason there have been no fugitive emissions from such activities in 1990-2009. However in 2010 a scotish company initiated a search for oil along the westcoast of Greenland. Three wells were drilled and tested in 2010. Five wells in 2011. The oil company reported a total fugitive emission of 44 tonnes CO₂-eqv methan from all three wells in 2010. This information has been recognised as the total fugitive emission of CH₄ from fuels in 2010. The calculation of fugitive emission of CO₂ and N₂O, and total fugitive emissions in 2011 has been based on IPCC Guideline emission factors (IPCC 2000 GPG, Table 2.16).

All oil exploration activities ere put on halt in 2012. No wells were drilled and tested in 2012. Hence, no fugitive emissions occurred in 2012.

Furthermore, some fugitive emission occurs in the distribution of fuel e.g. when refuelling from ships to on-shore tanks, onshore loading of fuel to ships and offshore loading of ships. The emission would only be in the form of NMVOC. The fugitive emission from loading/unloading of ships is currently not estimated.

International bunker fuels

International Aviation Bunkers

Emissions from international aviation bunkers are considered to be of negligible importance. The Greenland Airport Authority has reported the annual amount of jet fuel loaded into foreign aircrafts including Danish aircrafts. However, it is not possible to distinguish between Danish aircrafts and other aircrafts. Since most foreign aircrafts by far are Danish the annual amount of jet fuel loaded into foreign aircrafts are therefore included as part of the IPCC category 1A3a Civil aviation.

International Marine Bunkers

Emission from international marine bunkers is included from 2004 and onwards. Before 2004 international marine bunkers are considered to be of negligible importance.

Feedstocks and non-energy use of fuels

At the moment Greenland has no production or use of feedstocks. Emissions from non-energy use of fuels (e.g. bitumen and solvents) are included in other sectors of the Greenlandic inventory (Industrial Processes (CRF sector 2) and Solvent and Other Product Use (CRF sector 3)).

16.3.3 Methodological issues

Activity data

The Greenlandic emission inventory for fuel combustion has been performed according to the IPCC tier 1 methodology. The inventory is based on activity data from the Greenlandic energy statistics and on emission factors for different fuels, plants and sectors.

Total fuel combustion is based on data from Polaroil, Statoil and Malik Supply A/S. Polaroil imports and distributes fuel in all parts of Greenland. Statoil imports and distributes fuel in Kangerlussuaq. Malik Supply A/S, a Danish company, re-distributes fuel bought from Polaroil to Greenlandic trawlers, ships etc. By using detailed data from Polaroil, Statoil and Malik Supply A/S it is possible to determine total import, total export, total international bunkers and total domestic fuel combustion.

Total domestic fuel combustion is then divided into sectors and private households by using data from a survey on energy consumption, company specific sales data from Polaroil and local fuel distributors, company tax accountings, municipal accountings and Greenland Government accountings, and by estimation.

Since 2008 Statistics Greenland has conducted an annual survey among larger companies. By completing a questionnaire each company returned detailed information on the consumption of specific types of fuel in 2004-2012. The survey covered 53.2 % of total GHG emission from energy combustion in 2012, see Table 16.3.4. The decreased coverage by the survey is due to a drop in fuel combustion in companies that are covered by the survey primarily companies in Energy Industries.

By using detailed information on sales from Polaroil and local fuel distributors it is possible to determine fuel combustion in private companies and public offices with an automatic deal on supply. The sales data covered 11.8 % of total GHG emission from energy combustion in 2012, see Table 16.3.4.

Tax accountings in DKK are used to determine annual consumption of fuel in private companies, in municipalities, and within the Greenland Government. At the moment tax accountings are primarily used for determining fuel combustion in municipalities and public offices in settlements. Accountings cover 13.1 % of total GHG emission from energy combustion in 2012, see Table 16.3.4.

The remaining amount of total inland fuel combustion is divided into sectors and private households by estimation. This work is carried out by involving statistical material on population, housing, public finances, fisheries and hunting, and national accountings. The Greenlandic Business Register (GER) is used to divide remaining companies into sectors. Information on employees, operating units, vehicles etc. is used to determine the activity in each company.

Fuel combustion in private households is estimated using detailed information from a number of local fuel distributors. Fuel deliveries are registered by buildings. In Greenland each building has a unique number registered in the Greenlandic Area Register (NIN). By combining the NIN-register and the GER-register (see above) with statistics on housing and

population each building is labelled *private household* or located to a sector describing the main activity in the building. This new building-sector register, completed annually, is used extensively to determine the buyer of fuel delivered by Polaroil or local fuel distributors.

Fuel combustion in road traffic is based on a model designed by Statistics Greenland. The model contains data on the vehicle stock obtained from the Greenland Police Department's register on engine data. The vehicles are divided into broad categories of type i.e. personal car, lorry, taxi, truck, ambulance, motorbike etc. Each category is assigned with ratios on fuel type and mileage. Input data on mileage is derived from an annual survey among businesses and private road traffic in 2008-2013. Each vehicle is divided in business categories or labelled *private vehicle* according to the owner. For each group the emissions are estimated by combining vehicle and annual mileage numbers with standard emission factors according to the type of fuel. The model does not take cold start or hot engines into account.

For air traffic annual emissions are based on activity data from Air Greenland A/S and sales data from the Greenland Airport Authority. For navigation, ferries and freight, annual emissions are based on activity data from Royal Arctic Line A/S (freight), Royal Arctic Tankers A/S (freight), Royal Arctic Bygdeservice A/S (freight/passengers), and Arctic Umiaq Line A/S (passengers) and the liquidated Assartuivik A/S (passengers).

Table 16.3.4 shows the part of total CO₂ emission divided into sources - survey, specific sales data, tax accountings, and estimation.

Table 16.3.4 CO₂ emission from fuel combustion by sources to sectoral division (2004-2012).

	2004	2005	2006	2007	2008	2009	2010	2011	2012
	pct.								
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Survey	48.8	48.4	47.9	49.6	50.3	52.8	63.0	61.3	53.2
Sales data from Polaroil	2.7	3.4	3.7	3.6	3.4	3.0	4.2	5.0	5.7
Sales data from local fuel distributors	0.0	0.0	3.2	5.1	6.6	6.5	5.0	5.6	6.1
Accountings	12.7	12.1	12.9	12.8	12.2	12.7	10.8	11.0	13.1
Estimation	35.8	36.1	32.3	29.0	27.5	25.0	17.0	17.0	21.8

The procedure described above is used to divide total fuel combustion into sectors and private households during the period 2004-2012. Formerly, the period 1990-2003, activity data on sectors and private households were estimated using aggregated statistics on population, housing, companies, data on sales from Polaroil, and data on energy consumption in larger companies.

An increasing part of municipal waste incineration is utilised for heat and power production. Thus, incineration with energy-recovery is included in the Energy sector.

Table 16.3.5 shows the activity data on fuel combustion for the period 1990-2012.

Table 16.3.5 Activity data on fuel combustion (SINK categories).

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
	TJ										
Total	8 559	8 358	8 166	7 484	6 801	7 331	8 190	8 475	8 189	8 172	8 559
Energy industries	2 519	2 447	2 393	2 169	1 944	1 685	1 698	1 794	1 766	1 805	2 519
Manufacturing and construction	360	349	340	307	274	598	607	630	546	626	360
Domestic aviation	541	556	547	524	500	581	636	660	775	748	541
Road transport	501	488	476	437	397	370	369	387	361	401	501
National navigation	288	280	273	248	224	285	285	299	275	308	288
Commercial/Institutional	682	662	645	583	520	724	733	757	667	753	682
Residential	2 120	2 062	2 014	1 832	1 651	1 710	1 731	1 787	1 576	1 777	2 120
AFF	1 436	1 405	1 372	1 288	1 205	1 287	2 039	2 070	2 134	1 663	1 436
Other	113	110	107	97	86	91	91	91	91	91	113
<i>Continued</i>	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011
Total	8 514	7 995	8 964	8 840	8 898	9 153	9 031	9 371	8 207	9 387	10 026
Energy industries	1 885	1 900	1 915	1 976	1 959	2 032	1 934	2 057	1 813	3 192	3 542
Manufacturing and construction	624	590	680	696	755	763	787	814	592	531	649
Domestic aviation	632	603	646	608	633	691	701	753	635	654	723
Road transport	399	388	433	508	505	575	505	535	493	478	479
National navigation	308	297	334	464	420	421	334	347	350	378	405
Commercial/Institutional	725	699	796	1 014	979	1 107	939	969	784	641	694
Residential	1 748	1 670	1 895	2 155	2 032	2 271	1 804	1 888	1 628	1 577	1 615
AFF	2 101	1 755	2 174	1 317	1 516	1 161	1 921	1 871	1 691	1 600	1 628
Other	91	91	91	103	100	132	105	138	219	335	292
<i>Continued</i>	2012										
Total	8 014										
Energy industries	1 609										
Manufacturing and construction	501										
Domestic aviation	660										
Road transport	469										
National navigation	413										
Commercial/Institutional	742										
Residential	1 554										
AFF	1 851										
Other	215										

Sources: Statistics Greenland. Notes: Data on fuel combustion in 1993 are interpolated from 1992 and 1994, since no data is available for 1993.

Emission factors

For each fuel and source category a set of general area source emission factors has been determined. The emission factors are either nationally referenced or based on the IPCC Reference Manual (IPCC, 1997).

CO₂

The CO₂ emission factors applied are presented in Table 16.3.6. For municipal waste and all other fuels the same emission factor is applied for 1990-2012.

In 2013 a technical analysis has been conducted on the arctic gas oil that is by far the most dominant type of fuel in Greenland. The analysis was conducted by the Danish Technological Institute in order to gain a country specific emission factor on the Greenlandic gas oil, see Table 16.3.6 and Section 16.3.7 for further details.

In reporting to the Climate Convention, the CO₂ emission is aggregated to three fuel types: Liquid fuel, Biomass and Other fuel.

The CO₂ emission from incineration of municipal waste with energy-recovery (75.1 + 37.0 kg pr GJ) is divided into two parts: the emission from combustion of the plastic content of waste (which is included in the Greenlandic total) and the emission from combustion of the rest of the waste – the biomass part (which is reported as a memo item). In the IPCC reporting, the fossil part of the waste and the associated emissions from fuel combustion of the plastic content of the waste is reported in the fuel category, *Other fuels*. The emission factors on municipal waste were revised in the 2012 inventory submission. Greenland uses the Danish emission factors on municipal waste, which have been revised recently due to new information.

Table 16.3.6 CO₂ emission factors 1990-2012.

Fuel	Emission factor	Unit	Reference type	IPCC fuel Category
Gas oil	72.237	kg pr GJ	Country specific	Liquid
Kerosene	71.148	kg pr GJ	IPCC reference manual	Liquid
Jet-Kerosene	70.785	kg pr GJ	IPCC reference manual	Liquid
Gasoline	68.607	kg pr GJ	IPCC reference manual	Liquid
Fueloil	76.593	kg pr GJ	IPCC reference manual	Liquid
LPG	63.100	kg pr GJ	IPCC reference manual	Liquid
Waste oil	76.593	kg pr GJ	IPCC reference manual	Liquid
Municipal waste – biomass	75.100	kg pr GJ	Country specific	Biomass
Municipal waste – fossil fuel	37.000	kg pr GJ	Country specific	Other fuels

The CO₂ emission has been calculated by using the same methodology as described in the IPCC Guidelines (IPCC, 1997). This methodology implies use of C content per fuel type (default) and fraction of carbon oxidised (default); see the equation below.

$$E_{CO_2} = \sum Act_a \times EF_{C,a} \times Ox \times 44/12$$

where:

Act_a = activity; consumption of fuel a

EF_{C,a} = C emission factor for fuel a

Ox = oxidation factor

The emissions of CH₄, N₂O, NO_x, CO and NMVOC have been calculated at sector/fuel level by using IPCC default emission factors combined with measured/Danish EF waste incineration (with energy recovery), se Table 16.3.7 – Table 16.3.9 below.

The equation applied for each pollutant is:

$$E = \sum (EF_{ab} \times Act_{ab})$$

where:

EF	= emission factor
Act	= activity; fuel input
a	= fuel type
b	= sector activity

CH₄

The CH₄ emission factors applied for 1990-2012 are presented in Table 16.3.7. Emission factors for municipal waste refer to emission measurements carried out in Danish plants (Nielsen et al., 2010). Other emission factors refer to the IPCC Guidelines (IPCC, 1997).

Table 16.3.7 CH₄ emission factors 1990-2012.

Fuel group	Fuel	CRF sector	Emission factor, g pr GJ	Reference
Liquid	Gas oil	1A1 Energy Industries	3	IPCC, 1997
		1A2 Manufacturing Industries and Constructions	2	IPCC, 1997
		1A3a Transport – Civil aviation	0.5	IPCC, 1997
		1A3b Transport – Road transportation	5	IPCC, 1997
		1A3d Transport – Navigation	5	IPCC, 1997
		1A4a Other sectors – Commercial / Institutional	10	IPCC, 1997
		1A4b Other sectors – Residential	10	IPCC, 1997
		1A4c Other sectors – AFF stationary	10	IPCC, 1997
		1A4c Other sectors – AFF mobile	5	IPCC, 1997
		1A5b Other – Military mobile	5	IPCC, 1997
	Kerosene	1A1 Energy Industries	3	IPCC, 1997
		1A2 Manufacturing Industries and Constructions	2	IPCC, 1997
		1A3a Transport – Civil aviation	0.5	IPCC, 1997
		1A3b Transport – Road transportation	20	IPCC, 1997
		1A3d Transport – Navigation	5	IPCC, 1997
		1A4a Other sectors – Commercial / Institutional	10	IPCC, 1997
		1A4b Other sectors – Residential	10	IPCC, 1997
		1A4c Other sectors – AFF stationary	10	IPCC, 1997
		1A4c Other sectors – AFF mobile	5	IPCC, 1997
		1A5b Other – Military mobile	5	IPCC, 1997
	Gasoline	1A1 Energy Industries	3	IPCC, 1997
		1A2 Manufacturing Industries and Constructions	2	IPCC, 1997
		1A3a Transport – Civil aviation	0.5	IPCC, 1997
		1A3b Transport – Road transportation	20	IPCC, 1997
		1A3d Transport – Navigation	5	IPCC, 1997
		1A4a Other sectors – Commercial / Institutional	10	IPCC, 1997
		1A4b Other sectors – Residential	10	IPCC, 1997
		1A4c Other sectors – AFF stationary	10	IPCC, 1997
		1A4c Other sectors – AFF mobile	5	IPCC, 1997
		1A5b Other – Military mobile	5	IPCC, 1997
	Fueloil	1A1 Energy Industries	3	IPCC, 1997
		1A2 Manufacturing Industries and Constructions	2	IPCC, 1997
		1A3a Transport – Civil aviation	0.5	IPCC, 1997
		1A3b Transport – Road transportation	5	IPCC, 1997
		1A3d Transport – Navigation	5	IPCC, 1997
		1A4a Other sectors – Commercial / Institutional	10	IPCC, 1997
		1A4b Other sectors – Residential	10	IPCC, 1997
		1A4c Other sectors – AFF stationary	10	IPCC, 1997
		1A4c Other sectors – AFF mobile	5	IPCC, 1997
		1A5b Other – Military mobile	5	IPCC, 1997
	LPG	1A1 Energy Industries	1	IPCC, 1997
		1A2 Manufacturing Industries and Constructions	5	IPCC, 1997
		1A3a Transport – Civil aviation	-	IPCC, 1997
		1A3b Transport – Road transportation	50	IPCC, 1997
		1A3d Transport – Navigation	-	IPCC, 1997
		1A4a Other sectors – Commercial / Institutional	5	IPCC, 1997
		1A4b Other sectors – Residential	5	IPCC, 1997
		1A4c Other sectors – AFF stationary	5	IPCC, 1997
		1A4c Other sectors – AFF mobile	5	IPCC, 1997
		1A5b Other – Military mobile	-	IPCC, 1997
	Waste oil	1A1 Energy Industries	3	IPCC, 1997
Biomass	Municipal waste	1A1 Energy Industries	30Nielsen et al., 2010	
Other fuel	Municipal waste	1A1 Energy Industries	30Nielsen et al., 2010	

N₂O

The N₂O emission factors applied for 1990-2012 are presented in Table 16.3.8. Emission factors for municipal waste refer to emission measure-

ments carried out in Danish plants (Nielsen et al., 2010). Other emission factors refer to the IPCC Guidelines (IPCC, 1997).

Table 16.3.8 N₂O emission factors 1990-2012.

Fuel group	Fuel	CRF sector	Emission factor g pr GJ	Reference
Liquid	Gas oil	1A1 Energy Industries	0.6	IPCC, 1997
		1A2 Manufacturing Industries and Constructions	0.6	IPCC, 1997
		1A3a Transport – Civil aviation	2	IPCC, 1997
		1A3b Transport – Road transportation	0.6	IPCC, 1997
		1A3d Transport – Navigation	0.6	IPCC, 1997
		1A4 Other sectors	0.6	IPCC, 1997
		1A5b Other – Military mobile	0.6	IPCC, 1997
	Kerosene	1A1 Energy Industries	0.6	IPCC, 1997
		1A2 Manufacturing Industries and Constructions	0.6	IPCC, 1997
		1A3a Transport – Civil aviation	2	IPCC, 1997
		1A3b Transport – Road transportation	0.6	IPCC, 1997
		1A3d Transport – Navigation	0.6	IPCC, 1997
		1A4 Other sectors	0.6	IPCC, 1997
		1A5b Other – Military mobile	0.6	IPCC, 1997
	Gasoline	1A1 Energy Industries	0.6	IPCC, 1997
		1A2 Manufacturing Industries and Constructions	0.6	IPCC, 1997
		1A3a Transport – Civil aviation	2	IPCC, 1997
		1A3b Transport – Road transportation	0.6	IPCC, 1997
		1A3d Transport – Navigation	0.6	IPCC, 1997
		1A4 Other sectors	0.6	IPCC, 1997
		1A5b Other – Military mobile	0.6	IPCC, 1997
	Fueloil	1A1 Energy Industries	0.6	IPCC, 1997
		1A2 Manufacturing Industries and Constructions	0.6	IPCC, 1997
		1A3a Transport – Civil aviation	2	IPCC, 1997
		1A3b Transport – Road transportation	0.6	IPCC, 1997
		1A3d Transport – Navigation	0.6	IPCC, 1997
		1A4 Other sectors	0.6	IPCC, 1997
		1A5b Other – Military mobile	0.6	IPCC, 1997
	LPG	1A1 Energy Industries	0.1	IPCC, 1997
		1A2 Manufacturing Industries and Constructions	0.1	IPCC, 1997
		1A3a Transport – Civil aviation	-	IPCC, 1997
		1A3b Transport – Road transportation	0.1	IPCC, 1997
		1A3d Transport – Navigation	-	IPCC, 1997
		1A4 Other sectors	0.1	IPCC, 1997
		1A5b Other – Military mobile	0.1	IPCC, 1997
	Waste oil	1A1 Energy Industries	0.6	IPCC, 1997
Biomass	Municipal waste	1A1 Energy Industries	4	Nielsen et al., 2010
Other fuel	Municipal waste	1A1 Energy Industries	4	Nielsen et al., 2010

SO₂, NO_x, NMVOC and CO

Emission factors for SO₂, NO_x, NMVOC and CO are listed in Table 16.3.9. The same emission factors have been applied in the period 1990-2012.

Table 16.3.9 SO₂, NO_x, NMVOC and CO emission factors 1990-2012 (g pr GJ).

Fuel group	Fuel	CRF sector	NO _x	CO	NMVOC	SO ₂	Ref
Liquid	Gas oil	1A1 Energy Industries	200	15	5	141	1
		1A2 Manufacturing Industries and Constructions	200	10	5	141	1
		1A3a Transport – Civil aviation	300	100	50	141	1
		1A3b Transport – Road transportation	800	1 000	200	141	1
		1A3d Transport – Navigation	1 500	1 000	200	141	1
		1A4a,b Other sectors	100	20	5	141	1
		1A4c Other sectors – AFF stationary	100	20	5	141	1
		1A4c Other sectors – AFF mobile	1 200	1 000	200	141	1
		1A5b Other – Military mobile	1 500	1 000	200	141	1
	Kerosene	1A1 Energy Industries	200	15	5	23	1
		1A2 Manufacturing Industries and Constructions	200	10	5	23	1
		1A3a Transport – Civil aviation	300	100	50	23	1
		1A3b Transport – Road transportation	600	8 000	1 500	23	1
		1A3d Transport – Navigation	1 500	1 000	200	23	1
		1A4a,b Other sectors	100	20	5	23	1
		1A4c Other sectors – AFF stationary	100	20	5	23	1
		1A4c Other sectors – AFF mobile	1 200	1 000	200	23	1
		1A5b Other – Military mobile	1 500	1 000	200	23	1
	Gasoline	1A1 Energy Industries	200	15	5	46	1
		1A2 Manufacturing Industries and Constructions	200	10	5	46	1
		1A3a Transport – Civil aviation	300	100	50	46	1
		1A3b Transport – Road transportation	600	8 000	1 500	46	1
		1A3d Transport – Navigation	1 500	1 000	200	46	1
		1A4a,b Other sectors	100	20	5	46	1
		1A4c Other sectors – AFF stationary	100	20	5	46	1
		1A4c Other sectors – AFF mobile	1 200	1 000	200	46	1
		1A5b Other – Military mobile	1 500	1 000	200	46	1
	Fueloil	1A1 Energy Industries	200	15	5	492	1
		1A2 Manufacturing Industries and Constructions	200	10	5	492	1
		1A3a Transport – Civil aviation	300	100	50	492	1
		1A3b Transport – Road transportation	600	8 000	1 500	492	1
		1A3d Transport – Navigation	1 500	1 000	200	492	1
		1A4a,b Other sectors	100	20	5	492	1
		1A4c Other sectors – AFF stationary	100	20	5	492	1
		1A4c Other sectors – AFF mobile	1 200	1 000	200	492	1
		1A5b Other – Military mobile	1 500	1 000	200	492	1
	LPG	1A1 Energy Industries	150	20	5	0.13	1
		1A2 Manufacturing Industries and Constructions	150	30	5	0.13	1
		1A3a Transport – Civil aviation	-	-	-	-	1
		1A3b Transport – Road transportation	600	400	5	0.13	1
		1A3d Transport – Navigation	-	-	-	-	1
		1A4a,b Other sectors	50	50	5	0.13	1
		1A4c Other sectors – AFF stationary	50	50	5	0.13	1
		1A4c Other sectors – AFF mobile	1 000	400	5	0.13	1
		1A5b Other – Military mobile	-	-	-	-	1
	Waste oil	1A1 Energy Industries	200	15	5	477	1
Biomass	Municipal waste	1A1 Energy Industries	100	1 000	50	6	2
Other fuel	Municipal waste	1A1 Energy Industries	100	1 000	50	6	2

Sources: 1) IPCC Guidelines (IPCC, 1997). 2) Nielsen et al., 2010.

16.3.4 Emissions

The greenhouse gas (GHG) emissions are listed in Table 16.3.10. The total emission of greenhouse gases from energy (fuel combustion and fugitive emissions) accounts for 93.6 % of total Greenlandic GHG emission in 2012.

The CO₂ emission from energy accounts for 99.4 % of the Greenlandic CO₂ emission (excluding net CO₂ emission from Land Use, Land Use Change and Forestry (LULUCF)). The CH₄ emission from fuel combustion (Sectoral Approach) accounts for 9.0 % of the Greenlandic emission and the N₂O emission from fuel combustion accounts for 10.7 % of the Greenlandic N₂O emission, see Table 16.3.10.

Table 16.3.10 Greenhouse gas emission for the year 2012.

	CO ₂	CH ₄	N ₂ O
	Gg CO ₂ equivalent		
1A1 Fuel consumption, Energy Industries	109.6	0.2	0.5
1A2 Fuel consumption, Manufacturing Industries and Construction	36.2	0.0	0.1
1A3 Fuel consumption, Transport	109.6	0.2	0.6
1A4 Fuel consumption, Other sectors	313.9	0.7	0.8
1B2 Fugitive emissions from fuel, Oil and natural gas	NO	NO	NO
Total emission from energy	569.3	1.1	2.0
Greenlandic emission (excluding net emission from LULUCF)	572.7	12.1	18.4
	%		
Emission share for energy	99.4	9.0	10.7

CO₂ is the most important GHG pollutant and accounts for 99.5 % of the GHG emission in CO₂ equivalents from energy, see Figure 16.3.4.




Figure 16.3.4 GHG emissions (CO₂ equivalent) from stationary combustion plants.

Figure 16.3.5 depicts the time series of GHG emission in CO₂ equivalents from energy. As shown by the blue curve the development in total GHG emission follows the CO₂ emission development very closely. Emission of CO₂ and total GHG emission are respectively 7.3 % and 7.4 % lower in 2012 compared to 1990.




Figure 16.3.5 GHG emission time series for energy.

From 1990 to 1994 total GHG emission was reduced by 21 %. This was primarily due to the introduction of the first hydropower plant in 1993 but also to the introduction of burning waste to produce heat for district heating network in 1989. Dependence on gas oil conversion decreased immediately. Nevertheless, from 1995 an onwards consumption of gas oil once again increased due to the general economic development

In 2001-2002 total GHG emission decreased due to a minor recession in the economy. However since 1994 GHG emissions have increased in general with some fluctuations from year to year. The fluctuations are largely a result of outdoor temperature variations from year to year i.e. in 2008 the winter was relatively colder than in 2007. As a result fuel consumption increased in 2008 increasing GHG emission from fuel combustion. In 2009 GHG emission decreased by 13 % due to a significantly substitution in Energy Industries from fuel consumption to hydro power production together with a relatively warmer winter. However, in 2010 and 2011 GHG emission increased by 15 % and 7 % due to the initiation of oil exploration. In the most recent year, 2012 GHG emission decreased by 7.3 % due to the standstill in the oil exploration activities and a drop in fuel combustion in Energy Industries due to the opening of Greenlands fifth hydro power plant.

CO₂

CO₂ emission from energy accounts for 99.4 % of the total Greenlandic CO₂ emission. Table 16.3.11 lists the CO₂ emission inventory for the energy sector in 2012 as well as the relative percentage for each category under the sectoral approach.


The table reveals that Agriculture, Forestry and Fisheries (AFF) accounts for 23.3 % of the CO₂ emission. Other large CO₂ emission sources are Residential, Transportation and Energy Industries. These are sectors, which also account for a considerable share of fuel consumption.

Table 16.3.11 CO₂ emission from energy 2012.

		2012	
		Gg	%
1A1	Energy Industries	109.6	19.3
1A2	Manufacturing Industries	36.2	6.4
1A3	Transport	109.6	19.3
1A4a	Commercial / Institutional	53.6	9.4
1A4b	Residential	112.2	19.7
1A4c	Agriculture / Forestry / Fisheries	132.6	23.3
1A5	Other	15.5	2.7
1B2	Fugitive emissions from fuel, oil and natural gas	NO	NO
Total		569.3	100.0

The CO₂ emission from combustion of biomass fuels is not included in the total CO₂ emission data, since biomass fuels are considered CO₂ neutral. The CO₂ emission from biomass combustion is reported as a memo item in the Climate Convention reporting. In 2012, the CO₂ emission from biomass combustion was 14.1 Gg.

Time series for CO₂ emissions are provided in Figure 16.3.6. Fluctuations in CO₂ emission from AFF primarily regard fluctuations in fishing activities from year to year. Fluctuations in CO₂ emission from residential plants are largely a result of outdoor temperature variations from year to year. This also causes fluctuations in CO₂ emission from Energy Industries which cover electricity and heat production. However, the significant increase in emission from Energy Industries in 2010 continuing in 2011 is caused by the initiation of oil exploration in 2010, which is reported in the subsector "Manufacture of Solid Fuels and Other Energy Industries. Due to a stand-still in oil exploration fuel combustion in (and emissions from) Energy Industries dropped in 2012.

Figure 16.3.6 CO₂ emission time series for fuel combustion (Sectoral Approach).

Detailed trend discussion on CRF category level is available in Section 16.2.

CH₄


CH₄ emission from energy accounts for 9.0 % of the Greenlandic CH₄ emission. Table 16.3.12 lists the CH₄ emission inventory for energy in 2012. The table reveals that Residential plants accounted for 30.1 % of the CH₄ emission from energy in 2012. Energy Industries accounted for 19.27 % of the emission in 2012.

Table 16.3.12 CH₄ emission from fuel combustion 2012.

	2012	
	Mg	%
1A1 Energy Industries	9.9	19.2
1A2 Industry	1.0	2.0
1A3 Transport	7.4	14.3
1A4a Commercial / Institutional	7.4	14.4
1A4b Residential	15.5	30.1
1A4c Agriculture / Forestry / Fisheries	9.3	18.0
1A5 Other	1.1	2.1
1B2 Fugitive emissions from fuel, Oil and natural gas	NO	NO
Total	51.5	100.0

The CH₄ emission from energy has increased by 3.0 % since 1990. You may notice that CH₄ emission has decreased from 1990 to 2012 while CO₂ emission from energy has fallen in the same period. The reason for this is that the amount of recovered energy from waste has increased, while the consumption of liquid fossil fuel has decreased from 1990 to 2012. And in view of the fact that CH₄ emission from energy recovered waste by far exceeds the overall CH₄ emission from liquid fossil fuel, total CH₄ emission from energy has from 1990 to 2012, while CO₂ emissions dropped.

Time series for CH₄ emissions are provided in Figure 16.3.7. Fluctuations in CH₄ emission from AFF primarily regard fluctuations in fishing activities from year to year. Fluctuations in CH₄ emission from residential plants are largely a result of outdoor temperature variations from year to year. This also causes fluctuations in CH₄ emission from Energy Industries, which cover electricity and heat production and manufacture of solid fuels and other Energy Industries. The increase of CH₄ emission in 2010 and 2011 was caused by the initiation of activities concerning oil exploration, while the decrease of CH₄ emission in 2012 is due to a standstill in oil explorations in 2012.

Figure 16.3.7 CH₄ emission time series for energy.

Detailed trend discussion on CRF category level is available in Section 16.2.

N₂O


The N₂O emission from energy accounts for 10.7 % of the Greenlandic N₂O emission. Table 16.3.13 lists the N₂O emission inventory for energy in 2012. The table reveals that Transportations accounted for 29.0 % of the N₂O emission from energy. Energy Industries accounted for 25.2 % of the emissions in 2012.

Table 16.3.13 N₂O emission from energy 2012.

		2012	
		Mg	%
1A1	Energy Industries	1.6	25.2
1A2	Industry	0.3	4.7
1A3	Transport	1.8	29.0
1A4a	Commercial / Institutional	0.4	7.0
1A4b	Residential	0.9	14.6
1A4c	Agriculture / Forestry / Fisheries	1.1	17.4
1A5	Other	0.1	2.0
1B2	Fugitive emissions from fuel, oil and natural gas	NO	NO
Total		6.4	100.0

Figure 16.3.8 shows the time series for the N₂O emission from energy. The N₂O emission has increased by 5.3 % from 1990 to 2012. Similar to the increase in CH₄ emissions, N₂O emissions have increased from 1990 to 2012 due to an increase in the use of recovered energy from waste simultaneously to a decrease in the consumption of liquid fuels.

Once again, the 2010 and 2011 increases in N₂O emission from Energy Industries are predominantly caused by the startup of oil explorative activities, while the decrease of N₂O emission in 2012 is due to a standstill in oil explorations in 2012.

Figure 16.3.8 N₂O emission time series for energy.

Detailed trend discussion on CRF category level is available in Section 16.2.

SO₂, NO_x, NMVOC and CO

The emissions of SO₂, NO_x, NMVOC and CO from energy in 2012 are presented in Table 16.3.14. SO₂ from energy accounts for 99.4 % of the Greenlandic SO₂ emission. NO_x, CO and NMVOC account for 99.1, 89.6 % and 75.5 % respectively, of the Greenlandic emissions for these substances.

Table 16.3.14 SO₂, NO_x, NMVOC and CO emission from energy 2012.

	NO _x	CO	NMVOC	SO ₂
	Gg	Gg	Gg	Gg
1A1 Fuel consumption, Energy Industries	0.3	0.2	0.0	0.2
1A2 Fuel consumption, Manuf. Industries and Constr.	0.1	0.0	0.0	0.1
1A3 Fuel consumption, Transport	1.2	2.2	0.4	0.1
1A4 Fuel consumption, Other sectors	2.8	2.1	0.4	0.6
1B2 Fugitive emissions from fuel, Oil and natura gas	NO	NO	NE	NO
Total emission from fuel consumption and fugitive emissions from fuel	4.3	4.5	0.9	1.0
Greenlandic emission	4.4	5.0	1.2	1.0
	%			
Emission share for fuel consumption	99.1	89.6	75.5	99.4

16.3.5 Uncertainties

A tier 1 uncertainty assessment has been carried out in accordance with the IPCC GPG (IPCC, 2000). The uncertainty has been estimated for all sources included in the reporting for the energy sector. The uncertainties for the activity data and emission factors are shown in Table 16.3.15.

Table 16.3.15 Uncertainties for activity data and emission factors for the energy sector.

Subsector	Pollutant	Activity data uncertainty	Emission factor uncertainty
1A Liquid fuels	CO ₂	2	2
1A Municipal waste	CO ₂	2	25
1B2 Oil exploration	CO ₂	2	1 000
1A Liquid fuels	CH ₄	2	100
1A Municipal waste	CH ₄	2	100
1A Biomass	CH ₄	2	100
1B2 Oil exploration	CH ₄	2	1 000
1A Liquid fuels	N ₂ O	2	500
1A Municipal waste	N ₂ O	2	500
1A Biomass	N ₂ O	2	200
1B2 Oil exploration	N ₂ O	2	1 000

The activity data comes from the official Greenlandic energy statistics, which is considered to be of high quality, therefore the uncertainty of the activity data have been set to 2 %.

Regarding the emission factor uncertainty, the CO₂ emission factors are considered the most certain. In previous submissions the emission factor uncertainty for liquid fuels was set to 5 %. Due to a technical analysis a country specific emission factor is now available on the Greenlandic gas oil; the dominating liquid fuel comprising 79 % in 2012. Consequently, in this 2014 submission the CO₂ emission factor uncertainty has been revised from 5 % to 2 % for liquid fuels.

To account for the more inhomogeneous nature of municipal waste the emission factor uncertainty has been set to 25 %. For CH₄ the emission factor uncertainty has been set to 100 % in accordance with the IPCC GPG (IPCC, 2000). For N₂O the emission factor uncertainties have been estimated to between 200 % and 500 %. This is based on a first estimate and can be improved upon in the future.

Oil exploration has occurred in 2010 and 2011, but not in 2012. Regarding fugitive emissions from oil exploration emission factor uncertainty has been set to 1,000 % for CO₂, CH₄ and N₂O. In 2011 fugitive emission of CO₂, CH₄ and N₂O is calculated based on standard IPCC emission factors for drilling and testing. In 2010 the emission factor uncertainty for CH₄ was set to 200 % due to the fact that the amount of fugitive emission of CH₄ in 2010 was obtained directly from the scotish oil company and the uncertainty concerning CH₄ was therefore considered to be much lower than for CO₂ and N₂O.

The resulting uncertainties for the individual greenhouse gases and the total uncertainty on the greenhouse gas emission are shown in Table 16.3.16.

Table 16.3.16 Uncertainties for the emission estimates.

	Uncertainty %	Trend 1990-2012 %	Trend uncertainty %
GHG	± 3.2	-7.3	± 2.6
CO ₂	± 2.8	-7.4	± 2.6
CH ₄	± 89	3.0	± 10.6
N ₂ O	± 442	5.3	± 51

16.3.6 Source specific QA/QC

The elaboration of a formal QA/QC plan is to be completed.

However, the official Greenland energy statistics is continuously going through a great deal of quality work with regard to accuracy, comparability and completeness. Statistics Greenland is responsible for the official Greenlandic energy statistics, and as such responsible for the completeness of data. The uncertainties connected with estimating fuel consumption do not influence the coherence between the energy statistics and the datasets used in the emission inventory submission. For the remainder of the datasets, it is assumed that the level of uncertainty is relatively small. See chapter regarding uncertainties for further comments.

Statistics on fuel consumption is reported by Statistics Greenland in form of a spreadsheet. Annual consumption of gas oil, kerosene, gasoline and LPG are divided into business categories and private households. To ensure consistency data are compared with those from previous years and large discrepancies are checked.

All external data used for the emission inventory submission are archived in spreadsheets. Data are archived annually in order to ensure that the basic data for a given report are always available in their original form.

Safely stored and quality checked activity data are then processed by using a methodological approach consistent with international guidelines.

Calculated emission factors are compared with guideline emission factors to ensure that they are reasonable. The calculations follow the principle in international guidelines.

During data processing, it is checked that calculations are being carried out correctly. However, a documentation plan for this is to be elaborated.

Time series for activity data, emission factors and calculated emissions are used to identify possible errors in the calculation procedure. In fact, during the calculation, numerous controls take place to ensure correctness. Sums are checked of the various stages in the calculation procedure. Implied emission factors are compared to emission factors.

Every single time series imported to the CRF Reporter is checked for fuel rate, units for fuel rate, emission factor and plant-specific emissions. Additional checks are performed on the database. The database encloses every single activity data, emission factors, emission, notation key and comment imported to the CRF Reporter. In other words, no information is typed manually into the CRF Reporter. Instead, all information is imported to the CRF Reporter through XML-files to ensure maximum accuracy and completeness.

Reference approach

In addition to the sector-specific CO₂ emission inventories (the Greenlandic approach), the CO₂ emission is also estimated using the reference approach described in the IPCC Reference manual (IPCC, 1997). The reference approach is based on data for fuel production, import, export and stock change. The CO₂ emission inventory based on the reference approach is reported to the Climate Convention and used for verification of the official data in the Greenlandic approach.

Data for import, export and stock change used in the reference approach originate from the annual “basic data” table prepared by Statistics Greenland. The fraction of carbon oxidised has been assumed to be 1.00. The carbon emission factors are default factors originating from the IPCC Reference Manual (IPCC, 1997). The country-specific emission factors are not used in the reference approach, the approach being for the purposes of verification.

The Climate Convention reporting tables include a comparison of the Greenlandic approach and the reference approach estimates. To make results comparable, the CO₂ emission from incineration of the plastic content of municipal waste is added in the reference approach while the fuel consumption is subtracted.

In 2012 the fuel consumption rates in the two approaches differ by 0.0 % and the CO₂ emission differs by 0.9 %. In the period 1990-2012 both the fuel consumption and the CO₂ emission differ by 1 % or less at all times. The differences in energy consumption are below 1 % for all years. The difference in CO₂ emission is 1 % from 1990 to 1994, and below 1 % since 1995. According to IPCC Good Practice Guidance (IPCC, 2000) the difference should be within 2 %. A comparison of the Greenlandic approach and the reference approach is illustrated in Figure 16.3.9.




Figure 16.3.9 Comparison of the reference approach and the national approach.

16.3.7 Source specific recalculations and improvements

Improvements and recalculations since the 2013 emission inventory submission include:

- A technical analysis has been conducted on a sample of the Greenlandic gas oil. Due to the technical analysis, conducted by the Danish Technical Institute, a country specific emission factor is now available on the Greenlandic gas oil; the most dominant liquid fuel in Greenland with a percentage of 79 % of all liquid fuels used in 2012. The result of the technical analysis is found in Annex 8 (Section 16.19).
- Update of fuel rates according to the latest energy statistics. The update includes the years 2004-2011.
- And a further update of gas oil rates according to a new (country specific) calorific value on gas oil, see Annex 8 (Section 16.19). The update includes the years 2004-2011.
- Adjustment of municipal waste with energy recovery according to improvements in population statistics, which is used in the estimation of municipal waste.

Table 6.3.17 shows recalculations in the energy sector compared with the 2013 submission.

Table 16.3.17 Changes in GHG emission in the energy sector compared with the 2013 submission.

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
Previous inventory, Gg CO ₂ eqv.	625.7	610.8	596.6	546.2	495.7	534.4	597.5	618.2	596.7	594.7	668.7
Recalculated, Gg CO ₂ eqv.	617.5	602.9	588.9	539.2	489.5	527.7	589.9	610.4	589.3	587.4	660.3
Change in Gg CO ₂ eqv.	-8.2	-7.9	-7.7	-7.0	-6.2	-6.7	-7.6	-7.8	-7.4	-7.3	-8.4
Change in pct.	-1.3	-1.3	-1.3	-1.3	-1.3	-1.3	-1.3	-1.3	-1.2	-1.2	-1.3
<i>continued</i>	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011
Previous inventory, Gg CO ₂ eqv.	618.8	580.2	650.9	639.0	643.1	661.4	652.2	676.9	591.7	677.9	724.4
Recalculated, Gg CO ₂ eqv.	611.0	573.0	642.7	633.4	637.5	655.7	646.6	671.1	586.7	672.2	718.3
Change in Gg CO ₂ eqv.	-7.8	-7.2	-8.2	-5.6	-5.6	-5.7	-5.6	-5.8	-5.0	-5.8	-6.1
Change in pct.	-1.3	-1.2	-1.3	-0.9	-0.9	-0.9	-0.9	-0.9	-0.8	-0.8	-0.8
<i>continued</i>	2012										
Previous inventory, Gg CO ₂ eqv.	-										
Recalculated, Gg CO ₂ eqv.	572.4										
Change in Gg CO ₂ eqv.	-										
Change in pct.	-										

16.3.8 Source specific planned improvements

Some planned improvements to the emission inventories are discussed below.

1) Improved documentation for emission factors

The reporting of, and references for, the applied emission factors have been improved in the current year and will be further developed in future inventories. This will happen on the advice from the Danish National Environmental Research Institute.

2) Improvements in plant specific fuel combustion

Plant specific fuel combustion will be further improved according to the developments made by Statistics Greenland in the energy statistics.

3) Uncertainty estimates

Uncertainty estimates are largely based on the default uncertainty levels for activity rates and emission factors. More country-specific uncertainty estimates will be incorporated in future inventories.

4) Country specific emission factors

Statistics Greenland has acquired a technical analysis on the gas oil that is imported to and used in Greenland. The technical analysis conducted by the Danish Technical Institute has provided a country specific emission factor on the Greenlandic gas oil. The goal was to implement the new country specific emission factors in the 2015 submission, but with the technical results all ready in hand it was managed to implement the new country specific emission factor on gas oil in this 2014 submission. The arctic gas oil stands for 79 % of all liquid fuels in 2012.

The plan is to obtain additional country specific emission factors on other liquid fuels, but only if the UNFCCC recommend it as in the case of the Greenlandic gas oil.

16.3.9 References

Astrup, T., Larsen, A. W., Fuglsang, K. & Pedersen, N. H., 2012: PSO-0213, Biogenic Carbon in Danish Combustible Waste. DTU 2012. In press.

Consulting Company Carl Bro, 1996: Survey on waste from private households and companies in Greenland.

Consulting Company Carl Bro, 2001: Survey on waste from private households and companies in Greenland.

IPCC, 1997: Revised 1996 IPCC Guidelines for National Greenhouse Gas inventories. Available at:

<http://www.ipcc-nggip.iges.or.jp/public/gl/invs6.htm> (15-04-2007).

IPCC, 2000: Penman, J., Kruger, D., Galbally, I., Hiraishi, T., Nyenzi, B., Emmanuel, S., Buendia, L., Hoppaus, R., Martinsen, T., Meijer, J., Miwa, K. & Tanabe, K. (Eds). Good Practice Guidance and Uncertainty Management in National Greenhouse Gas Inventories. Published: IPCC/OECD/-IEA/IGES, Hayama, Japan. Available at:

<http://www.ipcc-nggip.iges.or.jp/public/gp/english/> (15-04-2007).

European Environmental Agency (EEA), 2007: EMEP/CORINAIR Emission Inventory Guidebook 2007, Technical Report No 16/2007. Prepared by the UNECE/EMEP Task Force on Emissions Inventories and Projections, 2007 update. Available at:

<http://www.eea.europa.eu/publications/EMEPCORINAIR5> (2009-02-27).

Previous version of the Guidebook can be linked from this page.

KEMIN, 2008: Ministry of Climate and Energy. Law on CO₂ quotas, nr. 348 of 09/05/2008. (Danish legislation). Available (in Danish) at:

<https://www.retsinformation.dk/Forms/R0710.aspx?id=117147>

(2009-03-01).

Greenland Government, Ministry of Environment and Nature: Data on municipal waste incineration with energy-recovery.

Nielsen M., Nielsen, O.-K. & Thomsen, M. 2010: Emissions from decentralised CHP plants 2007 - Energinet.dk Environmental project no. 07/1882. Project report 5 - Emission factors and emission inventory for decentralised CHP production. National Environmental Research Institute, Aarhus University. 113 pp. - NERI Technical report No. 786. Available at:

<http://www.dmu.dk/Pub/FR786.pdf> (2011-02-28).

Nielsen, O.-K., Plejdrup, M.S., Winther, M., Nielsen, M., Gyldenkerne, S., Mikkelsen, M.H., Albrektsen, R., Thomsen, M., Hjelgaard, K., Hoffmann, L., Fauser, P., Bruun, H.G., Johannsen, V.K., Nord-Larsen, T., Vesterdal, L., Møller, I.S., Caspersen, O.H., Rasmussen, E., Petersen, S.B., Baunbæk, L., Hansen, M.G., 2013: Denmark's National Inventory Report 2013 - Emission Inventories 1990-2012 - Submitted under the United Nations Framework Convention on Climate Change and the Kyoto Protocol. National Environmental Research Institute, University of Aarhus. 1,203 pp. - Scientific Report from DCE - Danish Center for Environment and Energy no. 56. Available at:

<http://www2.dmu.dk/Pub/SR56.pdf>

Pulles, T. & Aardenne, J.v. 2004: Good Practice for CLRTAP Emission Inventories, 24. June 2004. Available at:

<http://www.eea.europa.eu/publications/EMEPCORINAIR4/BGPG.pdf>

(2009-02-27).

Statistics Greenland: Energy Consumption in Greenland 2004-2012 published in Danish. Available at:

<http://www.stat.gl/publ/da/EN/201303/pdf/Grønlands%20energiforbrug%202012.pdf> as “Grønlands energiforbrug 2012” (01-11-2013).

Statistics Greenland, 2013: The Greenlandic energy statistics aggregated to SINK categories. Not published.

16.4 Industrial processes (CRF sector 2)

16.4.1 Overview of sector

In this chapter industrial emissions of greenhouse gases, not related to generation of energy, are presented.

The emission of greenhouse gases from industrial processes includes CO₂, HFCs and SF₆. The emissions are reported in CRF Tables 2(I), 2(I).A, 2(II), 2(II).C, 2(II).E and 2(II).F. Furthermore, the emission of non-methane volatile organic compounds (NMVOC) and CO from industrial processes related to asphalt roofing, road paving with asphalt and production of food and drink are given in CRF Table 2(I).

An overview of sources identified is presented in Table 16.4.1 with an indication of the contribution to the industrial part of the emission of greenhouse gases in 2012. Emissions are extracted from the CRF tables.

Table 16.4.1 Overview of greenhouse gas sources 2012.

Process	IPCC Code	Substance	Emission tonnes CO ₂ eqv.	%
Mineral Products				
Limestone and Dolomite Use	2A	CO ₂	19.57	0.272
Asphalt Roofing	2A	CO ₂	0.23	0.003
Road Paving with Asphalt	2A	CO ₂	0.10	0.001
Consumption of Halocarbons and SF₆				
Refrigeration and Air Conditioning Equipment	2F	HFCs	7 185	99.684
Electrical Equipment	2F	SF ₆	2.90	0.040
Total emission			7 208	100

The subsectors *Mineral Products* (2A) constitutes 0.276 % and *Consumption of Halocarbons and SF₆* (2F) constitutes 99.724 % of the industrial emission of greenhouse gases. The total emission of greenhouse gases (excl. LULUCF) in Greenland is estimated to 610.4 Gg CO₂ equivalent, of which industrial processes contribute with 7.208 Gg CO₂ equivalent (1.2 %). The emission of greenhouse gases from industrial processes from 1990-2012 are presented in Figure 16.4.1.

Greenland has no chemical industry, metal production or production of halocarbons or SF₆. Greenland has no consumption of PFCs.




Figure 16.4.1 Emission of greenhouse gases from industrial processes 1990-2012.

The key category in the industrial sector *Consumption of Halocarbons and SF₆* constitutes 1.2 % of the total emission of greenhouse gases. The trends in greenhouse gases from the industrial sector/subsectors are presented in Table 16.4.2. The emissions are extracted from the CRF tables.

Table 16.4.2 Emission of greenhouse gases from industrial processes in different subsectors from 1990-2012.

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
CO ₂ (tonnes CO ₂)											
A. Mineral Products	0.11	0.11	0.11	0.11	0.10	0.11	0.10	0.13	0.12	0.13	4.09
CH ₄	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
N ₂ O	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
HFCs (tonnes CO ₂ eqv.)											
F. Consumption of Halocarbons and SF ₆	NE	NE	NE	NE	16	25	77	390	713	1 279	1 871
PFCs (tonnes CO ₂ eqv.)											
F. Consumption of Halocarbons and SF ₆	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
SF ₆ (tonnes CO ₂ eqv.)											
F. Consumption of Halocarbons and SF ₆	NE	NE	NE	NE	NE	35.9	3.4	3.4	3.3	3.3	3.3
<i>continued</i>	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011
CO ₂ (tonnes CO ₂)											
A. Mineral Products	2.94	1.46	3.05	2.06	0.52	0.20	1.71	3.24	0.20	5.08	0.30
CH ₄	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
N ₂ O	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
HFCs (tonnes CO ₂ eqv.)											
F. Consumption of Halocarbons and SF ₆	2 964	3 898	4 750	5 425	5 499	5 558	6 065	6 527	6 568	6 771	7 144
PFCs (tonnes CO ₂ eqv.)											
F. Consumption of Halocarbons and SF ₆	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
SF ₆ (tonnes CO ₂ eqv.)											
F. Consumption of Halocarbons and SF ₆	3.2	3.2	3.2	3.1	3.1	3.1	3.0	3.0	3.0	3.0	2.9
<i>continued</i>	2012										
CO ₂ (tonnes CO ₂)											
A. Mineral Products	19.90										
CH ₄	NO										
N ₂ O	NO										
HFCs (tonnes CO ₂ eqv.)											
F. Consumption of Halocarbons and SF ₆	7 185										
PFCs (tonnes CO ₂ eqv.)											
F. Consumption of Halocarbons and SF ₆	NO										
SF ₆ (tonnes CO ₂ eqv.)											
F. Consumption of Halocarbons and SF ₆	2.9										

Greenland has no production of halocarbons or SF₆. Data on consumption of F-gases (HFCs and SF₆) are obtained from the Statistics Greenland (imports) and by an annual survey on consumption halocarbons and SF₆. Information on consumption of F-gases is available from 1995 onwards. Greenland has no consumption of PFCs.

One single plant in Greenland has reported use of SF₆ in 1995. The emission of SF₆ was 35.9 tonnes CO₂ equivalents in 1995. The annual emission from 1996 and onwards is assumed to be 0.5 % of the amount filled into the plant in 1995. This causes a relative high emission of SF₆ in 1995 and a much lower emission in the period 1996-2012.

Energy consumption associated with industrial processes and emissions thereof are included in the Energy sector of the inventory.

16.4.2 Source category description

Mineral products

The subsector *Mineral products* (2A) cover the following processes:

- Limestone and dolomite use.
- Roof covering with asphalt materials.
- Road paving with asphalt.

The time series for the emission of CO₂ from Mineral products (2A) are presented in Table 16.4.3. The emissions are extracted from the CRF tables and the values are rounded.

Table 16.4.3 Time series for emission of CO₂ (tonnes) from Mineral products (2A).

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
3. Limestone and dolomite use	-	-	-	-	-	-	-	-	-	-	3.96
5. Asphalt roofing	0.01	0.01	0.01	0.01	0.00	0.01	0.01	0.01	0.01	0.00	0.01
6. Road paving	0.10	0.10	0.10	0.10	0.10	0.10	0.09	0.11	0.11	0.13	0.12
Total	0.11	0.11	0.11	0.11	0.10	0.11	0.10	0.13	0.12	0.13	4.09
<i>continued</i>	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011
3. Limestone and dolomite use	2.77	1.32	2.64	1.80	0.11	0.03	1.51	2.96	0.03	4.94	0.00
5. Asphalt roofing	0.00	0.02	0.04	0.07	0.03	0.05	0.05	0.08	0.06	0.06	0.04
6. Road paving	0.17	0.12	0.37	0.19	0.38	0.12	0.15	0.20	0.11	0.07	0.26
Total	2.94	1.46	3.05	2.06	0.52	0.20	1.71	3.24	0.20	5.08	0.30
<i>continued</i>	2012										
3. Limestone and dolomite use	19.57										
5. Asphalt roofing	0.23										
6. Road paving	0.10										
Total	19.90										

The use of limestone and dolomite started in 2000. Hence there is no emission from limestone and dolomite use before 2000. The use of limestone and dolomite has been estimated from the annual import of these products to Greenland. Imports seem to vary a great deal from year to year, which causes the estimated use to vary as well.

In 2012 the most significant CO₂ emission came from the use of limestone and dolomite, which constituted 98.3 % of the total CO₂ emission from mineral products in 2012. According to Table 16.4.3 the emission of CO₂ from the use of limestone and dolomite exploded from nearly nothing in 2011 to 19.57 tonnes CO₂ in 2012. However, this increase in CO₂ emission

from limestone and dolomite is merely an expression of the fact that imports of limestone and dolomite increased substantial in 2012.

The relative increase in CO₂ emission is most significant for the use of asphalt roofing. From 1990 to 2012, CO₂ emissions increased from 0.01 to 0.23 tonnes CO₂; an increase of 2 441 %. The increase in CO₂ from asphalt roofing has primarily taken place from 2002 and onwards.

The CO₂ emission from subsectors under mineral products fluctuates a great deal from year to year. This is caused by fluctuations in building activities and road paving. However fluctuations in CO₂ are also caused by the fact that activity data for mineral products are based on import data, which do not allow distinction of imported amount into consumption and stockpiling.




Figure 16.4.2 Emission of CO₂ from mineral products.

Consumption of Halocarbons and SF₆

The subsector *Consumption of Halocarbons and SF₆* (2F) includes the following source categories and the following F-gases of relevance for Greenlandic emissions:

- 2F1: Refrigeration: HFC32, 125, 134a, 143a, unspecified HFCs.
- 2F8: Electrical equipment: SF₆.

A quantitative overview is given below for each of these source categories and each F-gas, showing their emissions in tonnes through the time series. The data is extracted from the CRF tables that form part of this submission and the data presented is rounded values. It must be noticed that the inventories for the years 1990-1993 (1994) might not cover emissions of these gases in full. The chosen base-year for these gases is 1995 for Greenland.

Table 16.4.4 Emission of HFCs from refrigeration (t).

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
HFC32	NE	NE	NE	NE	NE	NA	0.00	0.00	0.00	0.00	0.00
HFC125	NE	NE	NE	NE	NE	NA	0.01	0.04	0.08	0.15	0.22
HFC134a	NE	NE	NE	NE	0.01	0.02	0.03	0.06	0.10	0.17	0.24
HFC143a	NE	NE	NE	NE	NE	NA	0.01	0.05	0.09	0.16	0.24
Unspecified HFCs	NE	NE	NE	NE	NE	NA	0.00	0.00	0.00	0.00	0.00
<i>continued</i>	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011
HFC32	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
HFC125	0.35	0.46	0.56	0.64	0.64	0.65	0.71	0.76	0.77	0.80	0.84
HFC134a	0.35	0.45	0.55	0.63	0.65	0.65	0.68	0.67	0.64	0.62	0.63
HFC143a	0.39	0.51	0.63	0.71	0.72	0.72	0.79	0.86	0.88	0.91	0.97
Unspecified HFCs	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<i>continued</i>	2012										
HFC32	0.01										
HFC125	0.85										
HFC134a	0.59										
HFC143a	0.98										
Unspecified HFCs	0.00										

Table 16.4.5 Emission of SF₆ from electrical equipment (kg).

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
SF ₆	NE	NE	NE	NE	NE	1.50	0.14	0.14	0.14	0.14	0.14
<i>continued</i>	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011
SF ₆	0.14	0.13	0.13	0.13	0.13	0.13	0.13	0.13	0.13	0.12	0.12
<i>continued</i>	2012										
SF ₆	0.12										

The emission of SF₆ was highest in 1995, when one single plant in Greenland reported use of SF₆. The emission of SF₆ was 1.5 kg in 1995. Since 1995 the annual emission is assumed to be 0.5 % of the amount filled into the plant in 1995. This causes a relative high emission of SF₆ in 1995 and a much lower emission in the following years. In 2012 the emission of SF₆ was 0.12 kg.

HFCs are used in various types of refrigeration in industry, retail, buildings and onboard ships. In 1994 and 1995 consumption of HFC134a was the only reported HFC used for refrigeration. Since 1996 consumption of HFC32, 125, 134A, 143A has been reported continuously. The emission of HFCs has increased rapidly since 1995.

Table 16.4.6 and Figure 16.4.3 and Figure 16.4.4 quantify an overview of the emissions of the gases in CO₂ eqv. The reference is the trend table as included in the CRF table for year 2012.

Table 16.4.6 Time series for emission of HFCs and SF₆ (tonnes CO₂ eqv.).

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
HFCs	NE	NE	NE	NE	16	25	77	390	713	1 279	1 871
SF ₆	NE	NE	NE	NE	NE	35.9	3.4	3.4	3.3	3.3	3.3
<i>continued</i>	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011
HFCs	2 964	3 898	4 750	5 425	5 499	5 558	6 065	6 527	6 568	6 771	7 144
SF ₆	3.2	3.2	3.2	3.1	3.1	3.1	3.0	3.0	3.0	3.0	2.9
<i>continued</i>	2012										
HFCs	7 185										
SF ₆	2.9										





Figure 16.4.3 Emission of HFCs (from refrigeration).

Figure 16.4.4 Emission of SF₆ (from electrical equipment).

HFCs is by far the most dominant group. HFCs constitute a key category both with regard to the key category level and the trend analysis.

16.4.3 Methodological issues

General

The CO₂ emission from the use of limestone and dolomite, asphalt materials used for roof covering and road paving has been estimated from the annual import of these products to Greenland.

The emissions of HFCs and SF₆ have been estimated from data on consumption of F-gases. Activity data includes annual imports and data on consumption of halocarbons and SF₆ obtained from an annual survey among importers and consumers of F-gases.

The following sections contain a description of activity data and emission factors used for the subsectors under industrial processes. The section is

concluded by a description of the emissions of greenhouse gases from industrial processes.

Activity data

Activity data for subsectors *Mineral Products* (2A) and *Other Production* (2D) are presented in Table 16.4.7. Activity data under subsector *Other Production* (2D) are used for calculation of emission of non-methane volatile organic compounds (NMVOC).

The activity data are rounded. Notice that production of beer is given in hectolitre (hl). All other activity data are given in tonnes (t).

Statistics on imports are used to estimate annual consumption of mineral products. Statistics on imports of whole coffee beans and yeast for baking are used to estimate annual production of coffee and bread. Statistics on landings of fish and seafood to domestic plants are used to determine domestic processing of fish and seafood. Statistics on imports are produced by Statistics Greenland (2012b).

Production of beer including a fermentation process has taken place at the brewery “Godthåb Bryghus” since 2005 (Godthåb Bryghus, 2013). The brewery has reported annual production in rounded hectolitre. The much larger company “Nuuk Imeq” has no production of beer including a fermentation process. As a bottling company the activity at “Nuuk Imeq” only includes diluting of the concentrated quantities imported to Greenland and afterwards bottling of the beer.

Table 16.4.7 Time series for activity data for Mineral Products and Other Production (Godthåb Bryghus, 2013, Statistics Greenland, 2013b).

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001
Mineral Products												
2A3 Limestone and dolomite use (t)	-	-	-	-	-	-	-	-	-	-	9	6
2A5 Asphalt materials used for roofing (t)	37	35	39	39	13	56	29	59	39	7	26	11
2A6 Asphalt used for road paving (t)	591	581	595	604	597	577	532	664	649	752	694	988
Other Production												
Food and Drink -												
2D2 Beans roasted to produce coffee (t)	0	0	0	0	-	0	-	-	0	0	0	1
Food and Drink -												
2D2 Production of bread (t)	356	346	339	358	501	244	415	500	847	689	687	566
Food and Drink -												
2D2 Landings of fish and seafood (t)	76872	39565	55359	423	47967	78660	66262	24467	24763	75074	10566	929
Food and Drink -												
2D2 Production of beer (hl)	-	-	-	-	-	-	-	-	-	-	-	-
<i>continued</i>	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	Source	Source
Mineral Products												
2A3 Limestone and dolomite use (t)	3	6	4	0	0	3	7	0	11	0	45	1
2A5 Asphalt materials used for roofing (t)	81	149	263	114	193	209	321	241	256	173	929	1
2A6 Asphalt used for road paving (t)	705	2 218	1 127	2 258	698	912	1 206	629	443	1 529	583	1
Other Production												
Food and Drink -												
2D2 Beans roasted to produce coffee (t)	-	0	0	0	0	1	0	0	0	0	1	2
Food and Drink -												
2D2 Production of bread (t)	1 020	1 048	1 338	1 014	1 134	859	931	587	790	584	563	2
Food and Drink -												
2D2 Landings of fish and seafood (t)	85		102	103	111	118	109	102		104		
2D2 Landings of fish and seafood (t)	97080	667	570	642	351	260	420	393	99 829	020 97	532	3
Food and Drink -												
2D2 Production of beer (hl)	-	-	-	1 000	2 000	2 000	1 850	1 650	2 010	2 115	2 080	4

Sources:

- 1) Statistics on imports are used to estimate annual consumption of mineral products.
- 2) Statistics on imports of whole coffee beans and yeast for baking are used to estimate annual production of coffee and bread.
- 3) Statistics on landings of fish and seafood to domestic plants are used to determine domestic processing of fish and seafood.
- 4) Data from the brewery "Godthåb Bryghus" are used to determine annual production of beer.

The data for emission of HFCs and SF₆ has been obtained in continuation on the work on inventories for previous years. The determination includes the quantification and determination of any import and export of HFCs and SF₆ contained products and substances in stock form. This is in accordance with IPCC guidelines (IPCC (1997), vol. 3, p. 2.43ff), as well as the relevant decision trees from the IPCC Good Practice Guidance (IPCC (2000) p. 3.53ff).

The following sources of information have been used (Statistics Greenland, 2013a):

- Importers, wholesaler and suppliers.
- Statistics Greenland.
- Consuming enterprises.

Importers and suppliers provide consumption data of F-gases. Emission factors are defaults from the GPG. Import/export data for sub-source categories where import/export is relevant are quantified on estimates from

import/export statistics of products + default values of the amount of gas in the product.

The determination of emissions of F-gases is based on a calculation of the actual emission. The actual emission is the emission in the evaluation year, accounting for the time lapse between consumption and emission. The actual emission includes Greenlandic emissions from production and from products during their lifetimes. Consumption and emissions of F-gases are, whenever possible for individual substances, even though the consumption of certain HFCs has been limited. This has been varied out to ensure transparency of evaluation in the determination of GWP values. However, the continued use for Other HFCs has been necessary since not all importers and suppliers have specified records of sales for individual substances.

Only the actual emission has been calculated. Thus, the potential emission is assumed to be the same as the actual emission in the CRF tables.

Table 16.4.8 Content (w/w%) of “pure” HFC in HFC-mixtures, used as trade names.

HFC mixtures	HFC32	HFC125	HFC134a	HFC143a	Unspecified HFCs
	%	%	%	%	%
HFC-134, total			100		
HFC-404, total		44	4	52	
HFC-407c, total	23	25	52		
HFC-507a, total		50		50	
Unspecified HFCs					100

The substances have been accounted for in the survey according to their trade names, which are mixtures of HFCs used in the CRF. In the transfer to the “pure” substances used in the CRF reporting schemes, the ratios shown in Table 16.4.8 have been used.

Activity data for the consumption of F-gases is shown in Table 16.4.9. The activity data are rounded and given in kg.

Table 16.4.9 Time series for activity data for the consumption of F-gases by trade-names (Statistics Greenland, 2013a).

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
	Kg										
HFC-134											
Domestic	NE	NE	NE	264	139	91	187	134	453	319	289
Commercial and Industry	NE	NE	NE	-	-	-	123	123	247	247	493
Transport	NE	NE	NE	-	-	-	64	64	128	128	256
HFC-404a											
Commercial and Industry	NE	NE	NE	-	-	-	488	488	976	976	1 952
Transport	NE	NE	NE	-	-	-	82	82	164	164	328
HFC-407c											
Commercial and Industry	NE	NE	NE	-	-	-	34	34	68	68	135
HFC-507a											
Transport	NE	NE	NE	-	-	-	113	113	225	225	450
Unspecified HFCs											
Commercial and Industry	NE	NE	NE	-	-	-	45	45	90	90	180
SF ₆											
Electrical Equipment	NE	NE	NE	-	-	30	-	-	-	-	-
<i>continued</i>	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011
HFC-134											
Domestic	492	774	635	635	-	-	-	-	-	-	-
Commercial and Industry	493	493	493	260	208	680	329	312	195	484	340
Transport	256	256	256	120	120	30	30	-	-	-	-
HFC-404a											
Commercial and Industry	1 952	1 952	1 952	1 324	1 041	2 033	2 069	1 950	2 089	2 993	2 687
Transport	328	328	328	154	222	369	413	384	241	205	205
HFC-407c											
Commercial and Industry	135	135	135	68	83	31	4	112	90	-	90
HFC-507a											
Transport	450	450	450	-	-	120	180	-	120	-	180
Unspecified HFCs											
Commercial and Industry	180	180	180	326	314	556	698	309	400	576	600
SF ₆											
Electrical Equipment	-	-	-	-	-	-	-	-	-	-	-
<i>continued</i>	2012										
HFC-134											
Domestic	-										
Commercial and Industry	225										
Transport	-										
HFC-404a											
Commercial and Industry	2 760										
Transport	205										
HFC-407c											
Commercial and Industry	45										
HFC-507a											
Transport	180										
Unspecified HFCs											
Commercial and Industry	35										
SF ₆											
Electrical Equipment	-										

Emission factors

The CO₂ emission factors applied for mineral products in 2012 are presented in Table 16.4.10. The same emission factor has been applied for 1990-2012.

Table 16.4.10 CO₂ emission factors 2012.

Product	Emission factor	Unit	Reference	IPCC Category
Limestone and dolomite use	440	kg pr tonne	IPCC, 1997	2A3
Asphalt materials used for roofing	0.25	kg pr tonne	Nielsen et al., 2011	2A5
Asphalt used for road paving	0.168	kg pr tonne	Nielsen et al., 2011	2A6

The CO emission factors applied for the consumption of asphalt products under mineral products in 2012 are presented in Table 16.4.11. The same emission factor has been applied for 1990-2012.

Table 16.4.11 CO emission factors 2012.

Product	Emission factor	Unit	Reference	IPCC Category
Asphalt materials used for roofing	0.01	kg pr tonnes	Nielsen et al., 2011	2A5
Asphalt used for road paving	0.075	kg pr tonnes	Nielsen et al., 2011	2A6

The NMVOC emission factors applied for the consumption of asphalt products under mineral products and products used in the production of food and drink in 2012 are presented in Table 16.4.12. The same emission factor has been applied for 1990-2012.

Table 16.4.12 NMVOC emission factors 2012.

Product	Emission factor	Unit	Reference	IPCC Category
Asphalt materials used for roofing	0.08	kg pr tonnes	Nielsen et al., 2011	2A5
Asphalt used for road paving	0.015	kg pr tonnes	Nielsen et al., 2011	2A6
Food and Drink - Beans roasted to produce coffee	0.55	kg pr tonnes	IPCC, 1997	2D2
Food and Drink - Production of bread	8	kg pr tonnes	IPCC, 1997	2D2
Food and Drink - Landings of fish and seafood	0.3	kg pr tonnes	IPCC, 1997	2D2
Food and Drink - Production of beer	0.0625	kg pr hl	Nielsen et al., 2011	2D2

16.4.4 Emissions

The greenhouse gas (GHG) emissions are listed in Table 16.4.13. The emission from industrial processes accounts for 1.2 % of the Greenlandic GHG emission.

The CO₂ emission from industrial processes accounts for just 0.00348 % of the Greenlandic CO₂ emission (excluding net CO₂ emission from Land Use, Land Use Change and Forestry (LULUCF)). The HFC emission from industrial processes accounts for 100 % of the Greenlandic emission and the SF₆ emission accounts for 100 % of the Greenlandic SF₆ emission.


Table 16.4.13 Greenhouse gas emission for the year 2012.

	CO ₂	HFC	SF ₆
	Tonne CO ₂ equivalent		
2A3 Limestone and Dolomite Use	19.57	NA	NA
2A5 Asphalt Roofing	0.23	NA	NA
2A6 Road Paving with Asphalt	0.10	NA	NA
2F1 Refrigeration	NA	7 185	NA
2F8 Electrical Equipment	NA	NA	2.9
Total emission from industrial processes	19.90	7 185	2.9
Greenlandic emission (excluding net emission from LULUCF)	572 699	7 185	2.9
	%		
Emission share for industrial processes	0.00348	100	100

HFC is the most important GHG pollutant and accounts for 99.7 % of the GHG emission in CO₂ equivalents from industrial processes. Illustration of the percentage of share in a figure is omitted due to the large share of HFC, which completely dominates as the most significant GHG pollutant from industrial processes.

CO₂

Figure 16.4.5 depicts the time series of CO₂ emission from industrial processes. As shown by the blue curve total CO₂ emission follows the CO₂ emission from use of limestone and dolomite closely. Limestone and dolomite was not imported to Greenland before 2000. Thus emission of CO₂ from the use of mineral products increased significantly in 2000. The emission of CO₂ has increased by a factor 184 from 1990 to 2012 primarily due to the introduction of limestone and dolomite import in 2000. In 2012 limestone and dolomite imports increased significantly causing emissions from mineral products to increase as well. Data on imports are used to estimate the annual use of limestone and dolomite. This causes a great deal of fluctuations from year to year. Hence, in years with none or low import of limestone and dolomite, i.e. 2009 and 2011, CO₂ emission from the use of limestone and dolomite are also low.

Figure 16.4.5 Emission of CO₂ from industrial processes.

Emission of HFCs and SF₆ are illustrated in Figure 16.4.3 and Figure 16.4.4.

NMVOC and CO

The emissions of NMVOC and CO from industrial processes in 2012 are presented in Table 16.4.14. NMVOC and CO account for 2.92 % and 0.001 % respectively, of the Greenlandic emissions for these substances.

Table 16.4.14 NMVOC and CO emission from industrial processes 2012.

		NMVOC	CO
		Tonnes	
2A5	Asphalt Roofing	0.07	0.01
2A6	Road Paving with Asphalt	0.01	0.04
2D2	Food and Drink	33.90	NA
Total emission from industrial processes		33.98	0.05
Greenlandic emission		1 163.8	5 011.3
		%	
Emission share for industrial processes		2.92	0.001

16.4.5 Uncertainties

A tier 1 uncertainty assessment has been carried out in accordance with the IPCC GPG (IPCC, 2000). The uncertainty has been estimated for all sources included in the reporting for industrial processes. The uncertainties for the activity data and emission factors are shown in Table 16.4.15.

Table 16.4.15 Uncertainties for activity data and emission factors for industrial processes.

Subsector	Pollutant	Activity data uncertainty	Emission factor uncertainty
2A3 Limestone and dolomite use	CO ₂	5	5
2A5 Asphalt roofing	CO ₂	5	25
2A6 Road paving with asphalt	CO ₂	10	50
2F Consumption of HFC	HFC	10	50
2F Consumption of SF ₆	SF ₆	10	50

The activity data comes from the import statistics, which is considered to be of high quality, therefore the uncertainty of the activity data has been set to 5 % for limestone and dolomite use and asphalt roofing, while it is assumed to be 10 % for road paving and consumption of HFCs and SF₆.

Regarding the emission factor uncertainty, the CO₂ emission factor for limestone and dolomite use is considered very certain. It is derived from stoichiometric calculations. Thus an emission factor of 5 % has been assumed. The uncertainty levels for asphalt roofing and road paving are expert judgements. The emission of F-gases is dominated by emissions from refrigeration equipment and, therefore, the uncertainties assumed for this sector will be used for all the F-gases. The IPCC propose an uncertainty at 30-40 % for regional estimates. However, Greenlandic statistics have been developed over a number of years and, therefore the uncertainty on activity data is assumed to be 10 %. The uncertainty on the emission factor is, on the other hand, assumed to be 50 %. The base year for F-gases for Greenland is 1995.

The resulting uncertainties for the individual greenhouse gases and the total uncertainty on the greenhouse gas emission are shown in Table 16.4.16.

Table 16.4.16 Uncertainties for the emission estimates.

	Uncertainty %	Trend 1990-2012 ¹ %	Trend uncertainty %
GHG	± 36	16 856	± 4 679
CO ₂	± 7	18 258	± 4 440
HFC	± 51	29 210	± 4 145
SF ₆	± 51	-92	± 1.1

¹ For f-gases the base year of 1995 is used.

16.4.6 Source specific QA/QC

The elaboration of a formal QA/QC plan is to be completed.

However, the official Greenland import statistics has gone through a great deal of quality work with regard to accuracy, comparability and completeness. Statistics Greenland is responsible for the official Greenlandic import statistics, and as such responsible for the completeness of data.

Statistics on imports is reported by Statistics Greenland in form of a spreadsheet. Annual import of limestone and dolomite, asphalt materials used for roof covering and road paving, whole coffee beans and yeast for baking are compared with imports in previous years and large discrepancies are checked. The same procedure is used to ensure accuracy in annual use of F-gases and statistics on landings of fish and seafood to domestic plants.

All external data used for the emission inventory submission are archived in spreadsheets. Data are archived annually in order to ensure that the basic data for a given report are always available in their original form.

Safely stored and quality checked activity data are then processed by using a methodological approach consistent with international guidelines.

Calculated emission factors are compared with guideline emission factors to ensure that they are reasonable. The calculations follow the principle in international guidelines.

During data processing, it is checked that calculations are being carried out correctly. However, a documentation plan for this needs to be elaborated.

Time series for activity data, emission factors and calculated emissions are used to identify possible errors in the calculation procedure. In fact, during the calculation, numerous controls take place to ensure correctness. Sums are checked in the various stages in the calculation procedure. Implied emission factors are compared to emission factors.

Every single time series imported to the CRF Reporter is checked for annual activity, units for activity, emission factor and emissions. Additional checks are performed on the database. The database encloses every single activity data, emission factors, emission, notation key and comment imported to the CRF Reporter. In other words, no information is typed manually into the CRF Reporter. Instead, all information is imported to the CRF Reporter through XML-files to ensure maximum accuracy and completeness.

16.4.7 Source specific recalculations and improvements

The sectors *Mineral Products* (2A) and *Other Production* (2D) were included in the inventory for the first time in the 2010 submission. During implementation the following improvements were made:

Introduction of new activity data on non-energy use of limestone and dolomite, products containing bitumen used for asphalt roofing, and road paving with asphalt.

Introduction of new activity data on consumption of products used in the production of food and drink i.e. raw coffee beans, yeast used for baking, landings of fish, shellfish, seals and whales, and production of beer. Use of these products caused no CO₂ emission only non-methane volatile organic compounds (NMVOC).

Improved data on use of F-gases. Activity data on F-gases are now divided into domestic, commercial and industry, transport, and electrical equipment. Further more the substances, which are accounted according to their trade names, are now transferred into "pure" substances.

In the 2014 emission inventory submission there have been no further improvements or recalculations compared to the 2013 submission. Therefore Table 16.4.17 shows no changes in recalculations relating to industrial processes compared with the 2013 submission.

Table 16.4.17 Changes in GHG emission in the industrial processes sector compared with the 2013 submission.

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
Previous inventory, Gg CO ₂ eqv.	0.0	0.0	0.0	0.0	0.0	0.1	0.1	0.4	0.7	1.3	1.9
Recalculated, Gg CO ₂ eqv.	0.0	0.0	0.0	0.0	0.0	0.1	0.1	0.4	0.7	1.3	1.9
Change in Gg CO ₂ eqv.	-	-	-	-	-	-	-	-	-	-	-
Change in pct.	-	-	-	-	-	-	-	-	-	-	-
<i>continued</i>	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011
Previous inventory, Gg CO ₂ eqv.	3.0	3.9	4.8	5.4	5.5	5.6	6.1	6.5	6.6	6.8	7.1
Recalculated, Gg CO ₂ eqv.	3.0	3.9	4.8	5.4	5.5	5.6	6.1	6.5	6.6	6.8	7.1
Change in Gg CO ₂ eqv.	-	-	-	-	-	-	-	-	-	-	-
Change in pct.	-	-	-	-	-	-	-	-	-	-	-
<i>continued</i>	2012										
Previous inventory, Gg CO ₂ eqv.	-										
Recalculated, Gg CO ₂ eqv.	7.2										
Change in Gg CO ₂ eqv.	-										
Change in pct.	-										

16.4.8 Source specific planned improvements

Some planned improvements to the emission inventories are discussed below.

1) Distribution of unspecified mix of HFCs into single HFCs

An unspecified mix of HFCs is used in commercials and industries. In future inventories attempts will be made in order to distribute the unspecified mix of HFCs into single substances.

16.4.9 References

Godthåb Bryghus (Brewery in Nuuk), 2013: Data on production of beer 2006-2013. Not published.

IPCC, 1997: Revised 1996 IPCC Guidelines for National Greenhouse Gas inventories. Available at:

<http://www.ipcc-nggip.iges.or.jp/public/gl/invs6.htm> (15-04-2007).

IPCC, 2000: Good Practice Guidance and Uncertainty Management in National Greenhouse Gas Inventories. Available at:

<http://www.ipcc-nggip.iges.or.jp/public/gp/english/> (15-04-2007).

Nielsen, O.-K., Plejdrup, M.S., Winther, M., Nielsen, M., Gyldenkerne, S., Mikkelsen, M.H., Albrektsen, R., Thomsen, M., Hjelgaard, K., Hoffmann, L., Fauser, P., Bruun, H.G., Johannsen, V.K., Nord-Larsen, T., Vesterdal, L., Møller, I.S., Caspersen, O.H., Rasmussen, E., Petersen, S.B., Baunbæk, L., Hansen, M.G., 2013: Denmark's National Inventory Report 2013 – Emission Inventories 1990-2012 - Submitted under the United Nations Framework Convention on Climate Change and the Kyoto Protocol. National Environmental Research Institute, University of Aarhus. 1,203 pp. – Scientific Report from DCE – Danish Center for Environment and Energy no. 56. Available at:

<http://www2.dmu.dk/Pub/SR56.pdf>

Statistics Greenland, 2013a: Annual survey among importers, suppliers and consumers of F-gases in Greenland in 2012. Not published.

Statistics Greenland, 2013b: Foreign Trade, Import and Export. Available at: <http://www.stat.gl/dialog/main.asp?lang=da&sc=IE&version=201301> as “Grønlands udenrigshandel 2012” (27-03-2013). Data more detailed than the published version of the foreign trade statistics are used in order to access imports at the most detailed level.

16.5 Solvent and other product use (CRF sector 3)

16.5.1 Overview of sector

This section presents the methodology used for calculating CO₂ and NMVOC emissions from use of solvents in industrial processes and households that are related to the source categories Paint application (CRF sector 3A), Degreasing and dry cleaning (CRF sector 3B), Chemical products, manufacture and processing (CRF sector 3C) and Other (CRF sector 3D).

Solvents are chemical compounds that are used on a global scale in industrial processes and as constituents in final products to dissolve e.g. paint, cosmetics, adhesives, ink, rubber, plastic, pesticides, aerosols or are used for cleaning purposes, i.e. degreasing. NMVOCs are main components in solvents - and solvent use in industries and households is typically the dominant source of anthropogenic NMVOC emissions. In industrial processes where solvents are produced or used NMVOC emissions to air and as liquid can be recaptured and either used or destroyed. Solvent containing products are used indoor and outdoor and the majority of solvent sooner or later evaporate. A small fraction of the solvent ends up in waste or as emissions to water and may finally also contribute to air pollution by evaporation from these compartments.

In this section the methodology for the Greenland NMVOC emission inventory for solvent use is presented and the results for the period 1990 – 2012 are summarised. The method is based on the detailed approach described in EMEP/CORINAIR (2004) and emissions are calculated for the CRF sectors mentioned above.

16.5.2 Source category description

Table 16.5.1 and Figure 16.5.1 show the emissions of chemicals from 1990 to 2012, where the used amounts of single chemicals have been assigned to specific products and CRF categories.

Table 16.5.2 shows the used amounts of chemicals for the same period. Table 16.5.1 is derived from Table 16.5.2 by applying emission factors relevant to individual chemicals and production or use activities. Table 16.5.3 shows the used amounts of products from 1990 to 2012.

The default NMVOC-CO₂ conversion factor of $0.85 * 3.667 = 3.11$ is used.




Figure 16.5.1 Emission of NMVOC from solvent and other product use. The methodological approach for finding emissions is described in the text. Figures can be seen in Table 16.5.1.

Table 16.5.1 Emission of chemicals in tonnes per year.

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
Paint application (3A)	3.1	3.0	2.9	2.8	2.5	3.4	2.1	5.2	3.8	2.5	3.1
Degreasing and dry cleaning (3B)	0.1	0.1	0.1	0.1	0.4	NO	NO	0.1	0.2	NO	NO
Chemical products, manufacturing and processing (3C)	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.0	0.1	0.0
Other (3D)	81.2	80.1	79.4	81.2	85.4	85.2	64.9	79.2	82.8	96.8	76.4
Total NMVOC	84.4	83.3	82.5	84.1	88.3	88.7	67.1	84.4	86.9	99.4	79.5
Total CO ₂	263.4	259.7	257.4	262.5	275.6	276.7	209.3	263.4	271.0	310.1	247.9
<i>continued</i>	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011
Paint application (3A)	1.9	2.1	1.3	4.4	3.9	2.2	4.7	1.8	2.0	2.5	1.6
Degreasing and dry cleaning (3B)	NO	NO	NO	NO	NO	NO	NO	0.0	0.0	NO	NO
Chemical products, manufacturing and processing (3C)	0.0	0.1	0.0	0.2	0.1	1.2	1.3	0.4	0.6	0.7	0.0
Other (3D)	69.8	72.7	99.3	84.3	100.5	71.1	65.8	68.3	105.5	65.1	69.9
Total NMVOC	71.7	74.8	100.7	88.9	104.5	74.5	71.8	70.5	108.2	68.3	71.5
Total CO ₂	223.6	233.5	314.0	277.5	326.1	232.5	224.0	219.9	337.5	213.0	223.1
<i>continued</i>	2012										
Paint application (3A)	0.6										
Degreasing and dry cleaning (3B)	0.0										
Chemical products, manufacturing and processing (3C)	0.2										
Other (3D)	73.0										
Total NMVOC	73.9										
Total CO ₂	230.5										

Table 16.5.2 Used amounts of chemicals in tonnes per year.

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
Paint application (3A)	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
Degreasing and dry cleaning (3B)	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
Chemical products, manufacturing and processing (3C)	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
Other (3D)	37.0	36.6	35.1	34.8	59.6	43.5	45.4	32.8	27.1	36.5	18.6
Total NMVOC	37.0	36.6	35.1	34.8	59.6	43.5	45.4	32.8	27.1	36.5	18.6
<i>continued</i>	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011
Paint application (3A)	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
Degreasing and dry cleaning (3B)	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
Chemical products, manufacturing and processing (3C)	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
Other (3D)	33.0	20.0	31.9	27.5	27.4	30.4	24.2	26.2	68.2	36.7	23.9
Total NMVOC	33.0	20.0	31.9	27.5	27.4	30.4	24.2	26.2	68.2	36.7	23.9
<i>continued</i>	2012										
Paint application (3A)	NO										
Degreasing and dry cleaning (3B)	NO										
Chemical products, manufacturing and processing (3C)	NO										
Other (3D)	25.3										
Total NMVOC	25.3										

Table 16.5.3 Used amounts of products in tonnes per year.

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
Paint application (3A)	3.9	3.8	3.7	3.5	3.1	4.3	2.7	6.5	4.8	3.1	3.8
Degreasing and dry cleaning (3B)	0.2	0.2	0.1	0.1	0.8	NO	NO	0.1	0.4	NO	NO
Chemical products, manufacturing and processing (3C)	0.3	0.2	0.2	0.2	0.5	0.1	0.1	0.1	0.1	0.8	0.0
Other (3D)	84.6	83.5	83.5	85.8	84.9	84.5	61.8	81.8	90.9	105.7	83.8
Total products	89.0	87.7	87.5	89.7	89.4	89.0	64.6	88.6	96.1	109.5	87.6
<i>continued</i>	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011
Paint application (3A)	2.4	2.6	1.6	5.5	4.8	2.8	5.8	2.3	2.6	3.1	2.0
Degreasing and dry cleaning (3B)	NO	NO	NO	NO	NO	NO	NO	0.0	0.0	NO	NO
Chemical products, manufacturing and processing (3C)	0.1	0.4	0.2	0.5	0.3	11.5	13.9	4.4	6.1	5.7	0.3
Other (3D)	72.2	83.3	109.5	96.2	107.2	77.8	69.8	71.6	97.5	66.9	75.9
Total products	74.6	86.2	111.4	102.2	112.3	92.1	89.5	78.2	106.2	75.6	78.2
<i>continued</i>	2012										
Paint application (3A)	0.8										
Degreasing and dry cleaning (3B)	0.0										
Chemical products, manufacturing and processing (3C)	2.7										
Other (3D)	88.4										
Total products	91.9										

16.5.3 Methodological issues

Emission modelling of solvents can basically be done in two ways: 1) By estimating the amount of (pure) solvents consumed, or 2) By estimating the amount of solvent containing products consumed, taking account of their solvent content (EMEP/CORINAIR, 2004).

In 1) all relevant solvents must be estimated, or at least those together representing more than 90 % of the total NMVOC emission, and in 2) all relevant source categories must be inventoried or at least those together contributing more than 90 % of the total NMVOC emission. A simple approach is to use a pr capita emission for each category, whereas a detailed approach is to get all relevant consumption data (EMEP/CO-RINAIR, 2004).

The detailed method 1) is used in the emission inventory for solvent use, thus representing a chemicals approach, where each chemical (NMVOC) and chemical containing product (group) is estimated separately. The sum of emissions of all estimated NMVOCs used as solvents equals the NMVOC emission from solvent use.

Activity data

The definitions of solvents and VOC that are used are as defined in the solvent directive (Directive 1999/13/EC) of the EU legislation: "Organic solvent shall mean any VOC which is used alone or in combination with other agents, and without undergoing a chemical change, to dissolve raw materials, products or waste materials, or is used as a cleaning agent to dissolve contaminants, or as a dissolver, or as a dispersion medium, or as a viscosity adjuster, or as a surface tension adjuster, or a plasticiser, or as a preservative". VOCs are defined as follows: "Volatile organic compound shall mean any organic compound having at 293.15 K a vapour pressure of 0.01 kPa or more, or having a corresponding volatility under the particular condition of use".

Import figures of chemicals and chemical containing products are obtained from Statistics Greenland. There is no production or export of chemicals and chemical containing products, therefore the import amount is assumed to be equivalent to the used amount.

Emission factors

For some chemicals the emission factors are precise but for others they are rough estimates. In the Danish inventory emission factors are divided into four categories: 1) chemical industry (lowest EF), 2) other industry, 3) non-industrial activities, 4) domestic and other diffuse use (highest EF). This implies that high emission factors are applicable for use of solvent containing products and lower emission factors are applicable for use in industrial processes

The emission factors used in the Greenlandic inventory are the same as developed for the Danish inventory (please refer to Chapter 5). For the chemicals assumed to be used for industrial purposes the mean value of category 1 and 2 above is used.

16.5.4 Emissions

Table 16.5.1 and Figure 16.5.1 show the emissions of chemicals from 1994 to 2012, where the used amounts of single chemicals have been assigned to specific products and CRF categories. Table 16.5.2 shows the used amounts of chemicals for the same period. Table 16.5.1 is derived from Table 16.5.2 by applying emission factors relevant to individual chemicals and production or use activities. Table 16.5.3 showing the used amount of products is derived from Table 16.5.2, by assessing the amount of chemicals that is comprised within products belonging to each of the four source categories. The default NMVOC-CO₂ conversion factor of $0.85 * 3.667 = 3.11$ is used.

16.5.5 Uncertainties

A tier 1 uncertainty assessment has been carried out in accordance with the IPCC GPG (IPCC, 2000). The uncertainty has been estimated for all sources included in the reporting for solvent and other product use. The uncertainties for the activity data and emission factors are shown in Table 16.5.4.

Table 16.5.4 Uncertainties for activity data and emission factors for solvents.

Subsector	Pollutant	Activity data uncertainty	Emission factor uncertainty
3A Paint application	CO ₂	10	15
3B Degreasing and dry cleaning	CO ₂	10	15
3C Chemical products, manufacturing and processing	CO ₂	10	15
3D5 Other	CO ₂	10	20

The activity data comes from the import statistics, which is considered to be of high quality, therefore the uncertainty of the activity data has been set to 10 %.

Regarding the emission factor uncertainties, the uncertainty comprises of both the uncertainty of the NMVOC emission factor, and the uncertainty of the conversion factor of NMVOC to CO₂.

The resulting uncertainty for CO₂ is shown in Table 16.5.5.

Table 16.5.5 Uncertainties for the emission estimates.

	Uncertainty %	Trend 1990-2012 %	Trend uncertainty %
CO ₂	± 22.1	-12.5	± 12.2

16.5.6 Source specific QA/QC

Time series of activity data and emissions are analysed. Large inter annual variations is investigated further to ensure the accuracy of the estimates.

16.5.7 Source specific recalculations and improvements

Emissions from solvent and other product use were included in the Greenlandic emission inventory for the first time in the 2010 submission.

There have been no improvements in the 2014 emission inventory submission.

Priorily the notation key NE has been used regarding N₂O from fire extinguishers. However, a Danish research on the matter has showed that N₂O is not used in fire extinguishers. Since Greenland imports all fireextinguishers from Denmark, the notation key on N₂O in fire extinguishers has been changed from NE to NO concerning every year in the time series 1990-2011. The changed happened in the 2013 submission. With regard to aerosol cans, we are aware that N₂O is found in the products. Since we can not find any activity data on aerosol cans, we continue to report the notation key NE for N₂O in aerosol cans.

Table 16.5.6 Changes in GHG emission in the industrial processes sector compared with the 2013 submission.

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
Previous inventory, Gg CO ₂ eqv.	0.3	0.3	0.3	0.3	0.3	0.3	0.2	0.3	0.3	0.3	0.2
Recalculated, Gg CO ₂ eqv.	0.3	0.3	0.3	0.3	0.3	0.3	0.2	0.3	0.3	0.3	0.2
Change in Gg CO ₂ eqv.	-	-	-	-	-	-	-	-	-	-	-
Change in pct.	-	-	-	-	-	-	-	-	-	-	-
<i>continued</i>	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011
Previous inventory, Gg CO ₂ eqv.	0.2	0.2	0.3	0.3	0.3	0.2	0.2	0.2	0.3	0.2	0.2
Recalculated, Gg CO ₂ eqv.	0.2	0.2	0.3	0.3	0.3	0.2	0.2	0.2	0.3	0.2	0.2
Change in Gg CO ₂ eqv.	-	-	-	-	-	-	-	-	-	-	-
Change in pct.	-	-	-	-	-	-	-	-	-	-	-
<i>continued</i>	2012										
Previous inventory, Gg CO ₂ eqv.	-										
Recalculated, Gg CO ₂ eqv.	0.2										
Change in Gg CO ₂ eqv.	-										
Change in pct.	-										

16.5.8 Source specific planned improvements

It will be investigated whether use of N₂O is occurring in Greenland.

16.5.9 References

Nielsen, O.-K., Plejdrup, M.S., Winther, M., Nielsen, M., Gyldenkerne, S., Mikkelsen, M.H., Albrektsen, R., Thomsen, M., Hjelgaard, K., Hoffmann, L., Fauser, P., Bruun, H.G., Johannsen, V.K., Nord-Larsen, T., Vesterdal, L., Møller, I.S., Caspersen, O.H., Rasmussen, E., Petersen, S.B., Baunbæk, L., Hansen, M.G., 2013: Denmark's National Inventory Report 2013 - Emission Inventories 1990-2012 - Submitted under the United Nations Framework

Convention on Climate Change and the Kyoto Protocol. National Environmental Research Institute, University of Aarhus. 1,203 pp. – Scientific Report from DCE – Danish Center for Environment and Energy no. 56. Available at:
<http://www2.dmu.dk/Pub/SR56.pdf>

Emission Inventory Guidebook 3rd edition, prepared by the UNECE/EMEP Task Force on Emissions Inventories and Projections, 2002update. Available at: <http://reports.eea.eu.int/EMEP-CORINAIR3/en> (07-11-2003).

Directive 1999/13/EC of 11 March 1999 on the limitation of emissions of volatile organic compounds due to the use of organic solvents in certain activities and installations, Brüssel, 1999.

Statistics Greenland, 2013b: Foreign Trade, Import and Export. Available at: <http://www.stat.gl/dialog/main.asp?lang=da&sc=IE&version=201301> as “Grønlands udenrigshandel 2012” (27-03-2013). Data more detailed than the published version of the foreign trade statistics are used in order to access imports at the most detailed level.

16.6 Agriculture (CRF sector 4)

The emission of greenhouse gases from agricultural activities includes CH₄ emission from enteric fermentation, CH₄ and N₂O emission from manure management and N₂O emission from agricultural soils. The emissions are reported in CRF Tables 4.A, 4.B and 4.D.

Emission from rice production, burning of agricultural crop residue and burning of savannas does not occur in Greenland and the CRF Tables 4.F, 4.C and 4.E have, consequently, not been completed.

Emission of non-methane volatile organic compounds (NMVOC) from agricultural activities has not been estimated.

16.6.1 Overview of sector

In CO₂ equivalents, the agricultural sector (without LULUCF) contributes with 1.5 % of the overall greenhouse gas emission (GHG) in 2012. From 1990 to 2012 emissions increased from 8.78 Gg CO₂ equivalents to 9.34 Gg CO₂ equivalents, which correspond to an increase of 6.4 %, see Table 16.6.1. This emission increase is primarily caused by an growing use of synthetic fertiliser.

Table 16.6.1 Emission of GHG in the agricultural sector 1990-2012 in Gg CO₂ equivalents

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
CH ₄	6.16	6.21	5.58	4.89	5.33	5.72	5.89	6.42	6.16	5.52	5.36
N ₂ O	2.62	2.64	2.40	2.15	2.33	2.47	3.48	2.99	3.89	3.94	3.48
Total	8.78	8.85	7.98	7.03	7.66	8.19	9.37	9.41	10.05	9.46	8.84
<i>continued</i>	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011
CH ₄	5.46	5.26	5.33	5.60	5.82	5.64	5.77	5.63	5.52	5.66	5.54
N ₂ O	3.60	3.41	3.47	3.70	3.87	3.90	3.39	5.16	3.77	3.69	4.06
Total	9.06	8.67	8.80	9.30	9.70	9.54	9.16	10.79	9.29	9.35	9.60
<i>continued</i>	2012										
CH ₄	5.51										
N ₂ O	3.83										
Total	9.34										

As showed in Figure 16.6.1, CH₄ emission contributed with 59.0 % of the total GHG emission from the agricultural sector in 2012 and N₂O contributed with the remaining 41.0 % given in CO₂ equivalents. The major part of the emission is related to livestock production, which in Greenland particularly means the production of sheep. A smaller part is related to the reindeer production. Concerning the emission from agricultural soils, the main sources are use of synthetic fertiliser, nitrogen leaching from leaching and run-off and emission from grassing animals.




Figure 16.6.1 Emission of greenhouse gases from agriculture in 2012.

16.6.2 Source category description

The calculations of the emissions are based on methods described in the IPCC Reference Manual (IPCC, 1997) and the Good Practice Guidance (IPCC, 2000).

Statistics Greenland is responsible for collecting of data, preparation of emission inventory and reporting. Inputs of data are basically obtained from Statistics Greenland and the Greenland Agricultural Consulting Services (ACS). Data on climate are supplied by the Danish Meteorological Institute (DMI) and Greenland Survey (ASIAQ), and published by Statistics Greenland.

Table 16.6.2 List of institutes involved in the emission inventory for the agricultural sector.

References	Link	Abbrevia- tion	Data/information
Statistics Greenland	www.stat.gl	GST	<ul style="list-style-type: none"> - reporting - data collecting - no. of animal - feed import - use of synthetic fertiliser - spring temperature
The Agricultural Consulting Services	http://nunalerineq.org/	ACS	<ul style="list-style-type: none"> - N-excretion - milk yield - feed consumption and composition - stable- and grassing situation - animal growth and weight - land use - crop production
The Danish Plant Directorate	www.pdir.dk	PD	- N content in different fertiliser types
The Danish Agricultural Advisory Centre, Aarhus University	www.lr.dk	DAAC	- N content in crop residue

16.6.3 CH₄ emission from Enteric Fermentation (CRF sector 4A)

Description

The major part of the agricultural CH₄ emission originates from digestive processes. In 2012, this source accounts for 57.7 % of the total GHG emission from agricultural activities. The emission is primarily related to ruminants, which in Greenland is sheep. In 2012 sheep contributed with 87 % and the remaining 13 % from reindeer.

Methodological issues

The implied emission factors for all animal categories are based on the Tier 2/Country Specific (CS) approach. Feed consumption and composition for sheep and reindeer is based on data from Statistics Greenland and the Agricultural Consulting Services (ACS), which has information concerning the agricultural conditions in practice. Default values for the methane conversion rate (Y_m) for sheep given by the IPCC are used, as an average of mature sheep and lambs, which mean an Y_m value of 6 %.

Gross energy intake (GE)

The gross energy intake for sheep and reindeer is based on feeding plans for sheep from the Greenland Agricultural Consulting Services supplemented by data on imported feed. For reindeer information on gross energy intake is based on an article on reindeer management in Greenland.

Table 16.6.3 Parameters for calculation of emission from enteric fermentation.

Animal Category	Gross Energy (GE)	Methane conversion factor (Y_m)	Emission factor
	MJ pr head pr day		Kg CH ₄ pr head pr yr
Sheep	28.4	0.06	11.2
Reindeer	27.5	0.06	10.7

The default CH₄ emission factor for sheep Tier 1 methodology is estimated to 8 kg CH₄ per animal per year. The default GE is given as 20 MJ/head/yr, which is lower than the calculated GE for Greenland, and can explain the lower emission factor. Another reason could be the fact that the

national value for feed intake includes lambs. After lambing, ewes and lambs are put out to pasture. Thus lambs only feed through their mother and grass. Lambs are not fed separately before slaughter.

There is no default GE for reindeer. However, Norway, Sweden and Finland have estimated gross energy intake for reindeer to 29.6 – 31.6 MJ/head/day. Based on an article on reindeer management in southern Greenland by H.E. Rasmussen in 1992, the Greenlandic gross energy intake for reindeer has been estimated to 27.5 MJ pr head pr day, which is lower than Norway, Sweden and Finland. However, holding in mind that food conditions for reindeer is more scarcely in Greenland compared to conditions in Norway, Sweden and Finland, which have more forest, and that reindeer in Greenland are not fed separately, the estimated of gross energy intake for reindeer in Greenland seems acceptable.

Activity data

Table 16.6.4 shows the development in livestock. The number of sheep is varying slightly. The number of reindeer has decreased considerably since 1990. The reindeer livestock decreased significantly in 1999, when one of two reindeer stations closed. Since 1999 there has been only one reindeer station in Greenland.

Table 16.6.4 Number of animals from 1990-2012 (CRF Table 4.A. 4.B (a) and 4.B (b)).

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
Sheep	19 929	20 134	17 900	16 256	17 818	19 464	20 163	23 134	19 929	21 007	20 444
Reindeer	6 000	6 000	5 600	4 300	4 600	4 600	4 600	3 800	6 000	2 106	2 000
<i>continued</i>	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011
Sheep	20 394	18 967	19 259	20 383	21 317	21 289	21 704	21 080	20 139	20 729	20 232
Reindeer	2 480	3 100	3 100	3 100	3 100	2 318	2 441	2 500	3 000	3 000	3 000
<i>continued</i>	2012										
Sheep	20 107										
Reindeer	3 000										

Implied emission factor

The implied emission factor (IEF) could vary across years for sheep and reindeer due to changes in feed consumption. However, no existing data can document a change in feed intake. Therefore the same IEF is used for all years.

Time series consistency

The emission from enteric fermentation is given in Table 16.6.5. From 1990 to 2012, the emission has decreased by 10.5 % due to a fall in number of reindeer.

Table 16.6.5 Emission of CH₄ from Enteric Fermentation 1990–2012, tonnes CH₄.

CRF 4.A	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
Sheep	222	225	200	181	199	217	225	258	222	234	228
Reindeer	64	64	60	46	49	49	49	41	64	23	21
Total, tonnes CH ₄	287	289	260	227	248	266	274	299	287	257	250
Total, tonnes CO ₂ eqv.	6 018	6 066	5 452	4 775	5 208	5 594	5 758	6 275	6 018	5 396	5 240
<i>continued</i>	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011
Sheep	228	212	215	227	238	238	242	235	225	231	226
Reindeer	27	33	33	33	33	25	26	27	32	32	32
Total, tonnes CH ₄	254	245	248	261	271	262	268	262	257	263	258
Total, tonnes CO ₂ eqv.	5 336	5 141	5 209	5 473	5 692	5 510	5 635	5 502	5 393	5 532	5 415
<i>continued</i>	2012										
Sheep	224										
Reindeer	32										
Total, tonnes CH ₄	256										
Total, tonnes CO ₂ eqv.	5 386										

16.6.4 CH₄ and N₂O emission from Manure Management (CRF sector 4B)

Description

The emissions of CH₄ and N₂O from manure management are given in CRF Table 4.B (a) and 4.B (b). This source contributes with 8.3 % of the total emission from the agricultural sector in 2012. The major part of the emission originates from the production of sheep.

Methodological issues

CH₄ emission

The IPCC Tier 2/CS methodology has been used for the estimation of the CH₄ emission from manure management. Calculation of volatile solids, VS is based on national value of gross energy intake (GE). Default values is used for the maximum methane producing capacity (B₀), digestibility (DE), the ash content and the methane conversion factor (MCF).

For reindeer no default values exists. Thus DE, ASH and B₀ estimates for sheep are used. Sheep and reindeer are similar creatures, both ruminants. Greenlandic reindeer weigh an average of 70 kg. Greenlandic sheep weight approximately 50 kg. However, while sheep are fed relative more intensively, reindeer only feed on what they find in nature all year around. On these arguments the best estimate is to use DE, ASH and B₀ estimates for sheep on reindeer as well.

Table 16.6.6 CH₄ – Manure management – use of national parameters and IPCC default values.

Parameter	Unit	Sheep	Reindeer	Default or national value
Gross energy intake (GE)	MJ pr head pr day	28.4	27.2	National
Digestibility (DE)	Percent	60	60	IPCC default
Ash content (ASH)	Percent	8	8	IPCC default
Volatile solids (VS)	Kg VS pr head pr day	0.57	0.54	National
Max. methane producing capacity (B ₀)	M ³ pr kg VS	0.19	0.19	IPCC default
CH ₄ conversion factor (MCF), solid storage and pasture	Percent	1	1	IPCC default
Emission factor	Kg CH ₄ pr head pr yr	0.26	0.25	Tier 2

There are no changes in stable conditions or feed intake during the years 1990 to 2012. The implied emission factor is therefore the same for all years.

The default emission factor for sheep is 0.19 kg CH₄ per head per year. The higher national value is due to a higher estimate for gross energy intake.

Table 16.6.7 shows a decrease in the CH₄ emission from manure management from 1990 to 2012 by 11.2 %, which primarily is related to the fall in the production of reindeer.

Table 16.6.7 Emission of CH₄ from Manure Management 1990-2012, tonnes CH₄.

CRF 4.A	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
Sheep	5.2	5.2	4.7	4.2	4.6	5.1	5.2	6.0	5.2	5.5	5.3
Reindeer	1.5	1.5	1.4	1.1	1.2	1.2	1.2	1.0	1.5	0.5	0.5
Total, tonnes CH ₄	6.7	6.7	6.1	5.3	5.8	6.2	6.4	7.0	6.7	6.0	5.8
<i>continued</i>	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011
Sheep	5.3	4.9	5.0	5.3	5.5	5.5	5.6	5.5	5.2	5.4	5.3
Reindeer	0.6	0.8	0.8	0.8	0.8	0.6	0.6	0.6	0.8	0.8	0.8
Total, tonnes CH ₄	5.9	5.7	5.8	6.1	6.3	6.1	6.3	6.1	6.0	6.1	6.0
<i>continued</i>	2012										
Sheep	5.2										
Reindeer	0.8										
Total, tonnes CH ₄	6.0										

N₂O emission

Based on information from the Greenland Agricultural Consulting Services it is estimated that for sheep 55 % of the N-excretion is taken place in stable and all manure is handled as solid manure. The IPCC default emission value is applied, which means 2.0 % of the N-excretion for solid manure.

Reindeer is grassing all year. The emission from manure deposits on grass is included in "Pasture, Range and Paddock".

The total nitrogen excretion for sheep has increased by 0.9 % from 1990 to 2012 (Table 16.6.8) due to an increase in the number of sheep.

Table 16.6.8 Total nitrogen excretion for sheep, 1990-2012, tonnes N.

CRF table 4.B(b)	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
Livestock category											
N-excreted, tonnes in total	120	121	107	98	107	117	121	139	120	126	123
N-excretion, tonnes in stable	66	66	59	54	59	64	67	76	66	69	67
<i>continued</i>	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011
Livestock category											
N-excreted, tonnes in total	122	114	116	122	128	128	130	126	121	124	121
N-excretion, tonnes in stable	67	63	64	67	70	70	72	70	66	68	67
<i>continued</i>	2012										
Livestock category											
N-excreted, tonnes in total	121										
N-excretion, tonnes in stable	66										

Time series consistency

As shown in Table 16.6.9 total emission from manure management from 1990 to 2012 in CO₂ equivalents has decreased by 1.2 % due to a decrease in the number of reindeer.

Table 16.6.9 Emissions of N₂O and CH₄ from Manure Management 1990-2012.


	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
N ₂ O emission, tonnes CO ₂ eqv.	641	647	576	523	573	626	648	744	641	675	657
CH ₄ emission, tonnes CO ₂ eqv.	140	141	127	111	121	130	134	146	140	126	122
Total, tonnes CO ₂ eqv.	781	789	703	634	694	756	783	890	781	801	779
<i>continued</i>	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011
N ₂ O emission, tonnes CO ₂ eqv.	656	610	619	655	685	684	698	678	647	666	650
CH ₄ emission, tonnes CO ₂ eqv.	124	120	121	128	133	128	131	128	126	129	126
Total, tonnes CO ₂ eqv.	780	730	741	783	818	813	829	806	773	795	777
<i>continued</i>	2012										
N ₂ O emission, tonnes CO ₂ eqv.	646										
CH ₄ emission, tonnes CO ₂ eqv.	126										
Total, tonnes CO ₂ eqv.	772										

16.6.5 N₂O emission from Agricultural Soils (CRF sector 4D)

Description

The N₂O emissions from agricultural soils CRF Table 4.D contributed in 2012 with 34.0 % of the total emission from the agricultural sector. Figure 16.6.2 shows the overall development from 1990 to 2012 and the distribution on different sources. Since 1990 N₂O emissions increased suddenly in 1996, when farmers increased their use of synthetic fertiliser significantly. From 1997 to 2007 the emission of N₂O varied with an increasing trend. In 2008 the emission of N₂O increased considerably due to a considerable increase in the use of synthetic fertiliser caused by a periodical drought in the agricultural part of Greenland. In 2009 the use of synthetic fertiliser returned back to a more normal level, thus the emission of N₂O dropped as well. In 2012 the use of synthetic fertiliser decreased by of 13.7 % compared to 2011.

Emission from synthetic fertiliser and nitrogen leaching is an essential part of the total emission from agricultural soils and contributes totally with 58.7 %. Of the remaining sources the greatest part of the emission, by 20.4 %, origins from pasture, range and paddocks. Emissions from all sources have increased from 1990 to 2012 except from grassing animal where a fall in number of reindeer has taken place.

Figure 16.6.2 N₂O emissions from agricultural soils 1990-2012.

Methodological issues

To calculate the N₂O emission a combination of IPCC Tier 1a and Tier 1b is used. Tier 1b is used in calculation of emission from crops residue. Emissions of N₂O are closely related to the nitrogen balance. Data concerning

the N-excretion, evaporation of ammonia from synthetic fertiliser and grassing animal are based on national values.

The NH₃ and N₂O emission factor survey is presented in Table 16.6.10 and shows that except from histosols all N₂O emission factor is based on IPCC default values. The estimated emissions from the different sub-sources are described in the text which follows.

Table 16.6.10 Emissions factor - N₂O emission from the Agricultural Soils 1990-2012.

Agricultural soils – emission sources CRF Table 4.D	Ammonia emission factor	N ₂ O emission factor (country specific value)	N ₂ O emission factor (IPCC default value)
	Kg NH ₃ -N pr kg N	kg N ₂ O-N pr ha	kg N ₂ O -N pr kg N
1. Direct Soil Emissions			
Synthetic Fertiliser Applied to Soils	0.01 (CS)		0.0125
Animal Wastes Applied to Soils	0.20 (IPCC default)		0.0125
N-fixing Crops			0.0125
Crop Residue			0.0125
Cultivation of Histosols		1.06*	
2. Animal Production	0.07 (CS)		0.02
3. Indirect Soil Emissions			
Atmospheric Deposition			0.01
Nitrogen Leaching and Runoff			0.025

CS = country specific value.

* Include both emission from cropland and improved grassland. For further details see Section 16.7.

Direct emissions

Synthetic fertiliser

The calculation of nitrogen (N) applied to soil from use of synthetic fertiliser is based on data on imports from the Statistics Greenland. No data is available before 1994. The consumption for 1990 to 1993 is assumed to be on the same level as 1994. The nitrogen content for each fertiliser type is estimated based on expert judgement from the Danish Plant Directorate (Troels Knudsen, pers. comm.).

Table 16.6.11 shows the consumption of each type of fertiliser. Furthermore, the ammonia emission factor for each fertiliser is given, based on the values given in EMEP/EEA emission inventory guide book 2009 (Table 3-2). The emission factors are depending on the mean spring temperature estimated to seven degrees in Greenland. The spring temperature has to reflect the time where the fertilisers are applied, which in Greenland normally is June.

Table 16.6.11 Synthetic fertiliser consumption 2012 and the NH₃ emission factors.

Synthetic fertiliser	Calculation of ammonia emission factor ¹	NH ₃ emission factor ¹ kg NH ₃ -N pr kg N	Consumption ² t N
Fertiliser type			
Ammonium sulphate	=0.0107+0.0006*ts	1.49	NO
Ammonium nitrate	=0.008+0.0001*ts	0.87	7
Calcium ammonium nitrate	=0.008+0.0001*ts	0.87	0
Anhydrous ammonia	=0.0127+0.0012*ts	2.11	NO
Urea	=0.1067+0.0035*ts	13.12	0
Nitrogen solutions	=0.0481+0.0025*ts	6.56	NO
Ammonium phosphates	=0.0107+0.0006*ts	1.49	NO
Other NK and NPK	=0.008+0.0001*ts	0.87	133
Total consumption of N in synthetic fertiliser			141
National emission of NH ₃ -N, tonnes	1.2		
Average NH ₃ -N emission (FracGASF)	0.01		

*ts= means spring temperature=7 degree.

¹) EMEP/EEA (2009).

²) Statistics Greenland and the Danish Plant Directorate.

The Greenlandic value for the FracGASF is estimated to less than 0.01 in 2012, which is considerably lower than the recommended default value in IPCC, i.e. 0.10. The major part of the fertiliser types used in Greenland is related to ammonia nitrate and NPK fertiliser where the emission factor is quite low, i.e. 0.0087 kg NH₃-N pr kg N. Before 1995 urea accounted for a higher fraction. The value of FracGASF for these years is estimated to 0.10-0.13.

Table 16.6.12 FracGASF, 1990-2012.

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
FracGASM	0.13	0.13	0.13	0.13	0.13	0.10	0.02	0.03	0.01	0.01	0.01
<i>Continued</i>	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011
FracGASM	0.01	0.01	0.01	0.01	0.01	0.01	0.02	0.01	0.01	0.01	0.01
<i>Continued</i>	2012										
FracGASM	0.01										

Table 16.6.13 shows an increase in use of fertiliser and a particularly high increase in 2008. Due to a relatively small number of farms the individual handling of one farmer has a high effect on the total consumptions. With consumption of fertilisers being based on imports of fertilisers it is not possible to account for fertilisers bought for stockpiling. Thus it is possible that the relative high increase in use of fertilisers in 2008 is due to stockpiling. Another explanation could be that both 2007 and 2008 were relative dry years leading to a considerable decrease in amount of hay harvested. Hence, it is possible that farmers have tended to increase the use of fertilisers in 2008 to produce more feed. The use of fertiliser returned to a more normal level in 2009 and 2010. In 2011 the use of synthetic fertilisers increased 35.6 %. Most recently the use of synthetic fertilisers decreased 13.7 % in 2012.

Table 16.6.13 Nitrogen applied as fertiliser to agricultural soils 1990-2012.

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
N content in synthetic fertiliser, tonnes N	9	9	9	9	9	6	102	28	135	158	117
NH ₃ -N emission, tonnes	1	1	1	1	1	1	2	1	1	1	1
N in fertiliser applied on soil, tonnes N	8	8	8	8	8	6	100	27	134	157	116
N ₂ O emission, tonnes	0.16	0.16	0.16	0.16	0.16	0.11	1.97	0.53	2.63	3.08	2.28
<i>Continued</i>	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011
N content in synthetic fertiliser, tonnes N	126	114	117	128	136	144	86	273	134	120	163
NH ₃ -N emission, tonnes	1	1	1	1	1	1	1	2	1	1	1
N in fertiliser applied on soil, tonnes N	125	113	116	127	135	142	85	271	133	119	162
N ₂ O emission, tonnes	2.45	2.22	2.27	2.49	2.65	2.80	1.67	5.32	2.61	2.34	3.17
<i>Continued</i>	2012										
N content in synthetic fertiliser, tonnes N	141										
NH ₃ -N emission, tonnes	1										
N in fertiliser applied on soil, tonnes N	139										
N ₂ O emission, tonnes	2.74										

Manure applied to soil

The amount of nitrogen applied to soil from sheep on stables is estimated as the N-excretion in stables minus the ammonia emission, which occur in stables, under storage and in relation to the application of manure. There are no measurements of ammonia emission from stables in Greenland. Thus IPCC default is used. However, the FracGASM default at 0.20 (IPCC 1997, Table 4-19) match the Danish emission ammonia from sheep, which are estimated to 24 % in 1990 reduced to 19 % in 2008. A lower ammonia emission in Greenland is expected due to the cold climate, but on the other hand no ammonia reducing measures are implemented as in Denmark. The FracGASM at 0.20 are therefore considered as reliable.

Table 16.6.14 shows the development in nitrogen excretion in stables, the estimated amount of N applied on soil and the N₂O emission.

Table 16.6.14 Nitrogen applied as manure to agricultural soils 1990-2012.

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
N-excretion in stable, tonnes N	66	66	59	54	59	64	67	76	66	69	67
NH ₃ -N emission, tonnes N	13	13	12	11	12	13	13	15	13	14	13
N in manure applied on soil, tonnes N	53	53	47	43	47	51	53	61	53	55	54
N ₂ O emission, tonnes N ₂ O	1.03	1.04	0.93	0.84	0.92	1.01	1.05	1.20	1.03	1.09	1.06
<i>Continued</i>	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011
N-excretion in stable, tonnes N	67	63	64	67	70	70	72	70	66	68	67
NH ₃ -N emission, tonnes N	13	13	13	13	14	14	14	14	13	14	13
N in manure applied on soil, tonnes N	54	50	51	54	56	56	57	56	53	55	53
N ₂ O emission, tonnes N ₂ O	1.06	0.98	1.00	1.06	1.11	1.10	1.13	1.09	1.04	1.07	1.05
<i>Continued</i>	2012										
N-excretion in stable, tonnes N	66										
NH ₃ -N emission, tonnes N	13										
N in manure applied on soil, tonnes N	53										
N ₂ O emission, tonnes N ₂ O	1.04										

Crop residue

The cultivated area is approximately 1,070 ha with the main part as grass fields, only 10.5 ha are used for potato production. The cultivated area has

decreased compared to 2009 due to the shutdown of four farms. To estimate the emission from crop residue, IPCC Tier 1b has been applied. N₂O emissions from crop residues are calculated based on the total above-ground N-content in crop residue returned to soil, which in Greenland includes residue of leafs from grass fields and the top from potatoes.

National values for nitrogen content used are provided by the Faculty of Agricultural Sciences, Aarhus University (Djurhuus and Hansen 2003). Values are calculated based on relatively few observations related to Danish conditions, but are at present the best available data.

Table 16.6.15 N-content in crops residue 2012.

Crop type	Stubble	Husks	Top	Leafs	Frequency of	Nitrogen content	
					ploughing	in crop residue	
	kg N pr ha	kg N pr ha	kg N pr ha	kg N pr ha	No. of year before ploughing	kg N pr ha pr yr	kg N pr yr
Potatoes (top), non-harvest	-	-	48.7	-	1	48.7	17 441
Grass- and clover field in rotation	32.3		-	10.0	5	16.5	511
Total N from crop residue – 2012, kg							17 952

Reference: Djurhuus and Hansen 2003

To calculate the N₂O emission the IPCC standard emission factor 1.25 % is used. The national emission from crop residues has more than doubled from 1990 to 2012 (Table 16.6.16) as a result of increasing agricultural area.

Table 16.6.16 Emissions from crop residue 1990-2012.

Crop residue	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
Grass stub/leaves, kg N	8 071	8 498	8 925	9 352	9 778	10 205	10 632	11 059	11 486	11 912	12 339
Potato tops, kg N	-	-	-	-	-	-	-	-	-	-	-
Crop residue total, kg N	8 071	8 498	8 925	9 352	9 778	10 205	10 632	11 059	11 486	11 912	12 339
N ₂ O emission, kg	159	167	175	184	192	200	209	217	226	234	242
<i>Continued</i>	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011
Grass stub/leaves, kg N	12 766	14 005	14 384	14 614	15 176	15 823	16 018	16 378	17 925	17 546	17 624
Potato tops, kg N	244	244	244	244	244	244	244	244	317	317	511
Crop residue total, kg N	13 010	14 249	14 628	14 857	15 420	16 066	16 262	16 621	18 241	17 863	18 135
N ₂ O emission, kg	256	280	287	292	303	316	319	326	358	351	356
<i>Continued</i>	2012										
Grass stub/leaves, kg N	17 441										
Potato tops, kg N	511										
Crop residue total, kg N	17 952										
N ₂ O emission, kg	353										

Frac vaules

There is no cultivation of nitrogen fixing crops, why the Fraction value $Frac_{NCRBF}$ is not relevant. Until national data is available, the default value of $Frac_{NCRO}$ by 0.015 is used. The default value of $Frac_R$ is not current for the Greenlandic conditions, where the main part of the aboveground biomass is harvest and used for ensilage. Until national data is available, the $Frac_R$ is registered as “Not Estimated”.

Cultivation of histosols

N₂O emissions from histosols are based on the area with organic soils multiplied by the emission factor of 1.06 kg N₂O-N pr. kg N in 2012. See Section 16.7 on LULUCF for further description on cultivation of histosols.

Table 16.6.17 shows an increase in the N₂O emission from 1990 to 2012 due to extend of the agricultural area.

Table 16.6.17 Activity data and emission from cultivation of histosols 1990-2012.

CRF – Table 4.D	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
Cultivated histosols, ha	123	129	136	142	149	155	161	168	174	181	187
N ₂ O emission, kg	201	211	222	232	243	254	264	275	285	296	307
<i>Continued</i>	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011
Cultivated histosols, ha	195	214	220	223	232	242	245	250	274	268	270
N ₂ O emission, kg	325	356	366	371	385	401	406	415	456	447	455
<i>Continued</i>	2012										
Cultivated histosols, ha	268										
N ₂ O emission, kg	451										

Pasture, Range and Paddock

The amount of nitrogen deposited on grass includes grassing from reindeer 365 days a year and from sheep 164 days a year. An ammonia emission factor of 7 % is used for all animal categories based on investigations from the Netherlands and the United Kingdom (Jarvis et al., 1989a. Jarvis et al., 1989b and Bussink, 1994). EMEP/EEA emission inventory guidebook 2009 use a similar emission factor at 6 % for grassing dairy cattle (calculated from 4B, Appendix B).

Table 16.6.18 shows the estimated values of N-excretion from grassing animals, ammonia emission, the N₂O emission and the FracGRAZ value. As a consequence of an overall drop in number of reindeer, both the N₂O emission and the FracGRAZ value have decreased from 1990 to 2012.

Table 16.6.18 Emission from grassing animals 1990-2012.

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
N-excretion on grass, tonnes N	88	89	81	69	75	79	81	84	88	69	67
NH ₃ -N emission, tonnes	6	6	6	5	5	6	6	6	6	5	5
N deposited on grass, tonnes N	82	83	75	64	69	73	75	78	82	64	62
N ₂ O emission, tonnes	2.58	2.60	2.35	2.01	2.18	2.31	2.36	2.46	2.58	2.01	1.95
FracGRAZ	0.57	0.57	0.58	0.56	0.56	0.55	0.55	0.52	0.57	0.50	0.50
<i>Continued</i>	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011
N-excretion on grass, tonnes N	69	69	70	73	75	71	73	71	72	73	72
NH ₃ -N emission, tonnes	5	5	5	5	5	5	5	5	5	5	5
N deposited on grass, tonnes N	64	64	65	68	70	66	68	66	67	68	67
N ₂ O emission, tonnes	2.03	2.02	2.04	2.13	2.20	2.07	2.12	2.08	2.09	2.14	2.10
FracGRAZ	0.51	0.52	0.52	0.52	0.52	0.50	0.50	0.51	0.52	0.52	0.52
<i>Continued</i>	2012										
N-excretion on grass, tonnes N	72										
NH ₃ -N emission, tonnes	5										
N deposited on grass, tonnes N	67										
N ₂ O emission, tonnes	2.09										
FracGRAZ	0.52										

Indirect emissions

Atmospheric deposition

Atmospheric deposition includes ammonia emission from manure management, use of synthetic fertiliser and from grassing animals.

The N₂O emission from atmospheric deposition is nearly unaltered from 1990 to 2012. The fall in the reindeer production compensate for an increase in the number of sheep and a rise in use of synthetic fertiliser.

Table 16.6.19 Emission from atmospheric deposition 1990-2012.

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
NH ₃ -N manure management, tonnes	13	13	12	11	12	13	13	15	13	14	13
NH ₃ -N synthetic fertiliser, tonnes	1	1	1	1	1	1	2	1	1	1	1
NH ₃ -N pasture, tonnes	6	6	6	5	5	6	6	6	6	5	5
NH ₃ -N total, tonnes	21	21	19	17	18	19	21	22	21	20	19
N ₂ O emission, tonnes	0.32	0.33	0.29	0.26	0.29	0.30	0.33	0.34	0.32	0.32	0.30
<i>Continued</i>	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011
NH ₃ -N manure management, tonnes	13	13	13	13	14	14	14	14	13	14	13
NH ₃ -N synthetic fertiliser, tonnes	1	1	1	1	1	1	1	2	1	1	1
NH ₃ -N pasture, tonnes	5	5	5	5	5	5	5	5	5	5	5
NH ₃ -N total, tonnes	19	18	19	20	21	20	21	21	19	20	20
N ₂ O emission, tonnes	0.30	0.29	0.29	0.31	0.32	0.32	0.33	0.33	0.31	0.31	0.31
<i>Continued</i>	2012										
NH ₃ -N manure management, tonnes	13										
NH ₃ -N synthetic fertiliser, tonnes	1										
NH ₃ -N pasture, tonnes	5										
NH ₃ -N total, tonnes	20										
N ₂ O emission, tonnes	0.31										

Nitrogen leaching and Run-off

The amount of nitrogen lost by leaching and run-off is calculated by using the IPCC default FracLEACH at 0.3 (IPCC 1997, Table 4-24).

The N₂O emission from N-leaching and runoff more than doubled from 1990 to 2008. However, lately in 2009-2010 and 2012 total N₂O emission has decreased each year. In 2012 the N₂O emission from N-leaching and runoff amounted to 3.28 tonnes, which is 70.5 % more than in 1990.

From 1990 to 2012 total nitrogen content in manure has decreased due to a fall in the reindeer production. However, in the same period the use of synthetic fertilisers has increased significantly causing the overall N₂O emission from N-leaching and runoff to increase.

Table 16.6.20 Emission from N-leaching and runoff 1990-2012.

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
N-excretion total, tonnes N	154	155	140	122	133	143	147	161	154	138	134
N in synthetic fertiliser, tonnes	9	9	9	9	9	6	102	28	135	158	117
N ₂ O emission, tonnes	1.92	1.94	1.75	1.55	1.68	1.76	2.94	2.22	3.41	3.50	2.96
<i>Continued</i>	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011
N-excretion total, tonnes N	137	132	133	140	146	141	144	141	138	142	139
N in synthetic fertiliser, tonnes	126	114	117	128	136	144	86	273	134	120	163
N ₂ O emission, tonnes	3.09	2.89	2.95	3.16	3.32	3.36	2.72	4.88	3.20	3.09	3.55
<i>Continued</i>	2012										
N-excretion total, tonnes N	138										
N in synthetic fertiliser, tonnes	141										
N ₂ O emission, tonnes	3.28										

Activity data

Table 16.6.21 provides an overview on activity data from 1990 to 2012 used to the estimation of N₂O emission from agricultural soils. For all emission

sources the unit tonnes of nitrogen are used except from cultivation of histosols, where the unit is given as hectare.

Table 16.6.21 Activity data - agricultural soils 1990-2012, tonnes N (cultivation of histosols = ha).

CRF – Table 4.D	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
1. Direct Emissions											
Synthetic Fertiliser	8	8	8	8	8	6	100	27	134	157	116
Animal Manure Applied to Soils	53	53	47	43	47	51	53	61	53	55	54
Crop Residue	8	8	9	9	10	10	11	11	11	12	12
Cultivation of histosols	123	129	136	142	149	155	161	168	174	181	187
2. Pasture, Range and Paddock Manure	82	83	75	64	69	73	75	78	82	64	62
3. Indirect Emissions											
Atmospheric Deposition	21	21	19	17	18	19	21	22	21	20	19
Nitrogen Leaching and Run-off	49	49	45	39	43	45	75	56	87	89	75
<i>Continued</i>	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011
1. Direct Emissions											
Synthetic Fertiliser	125	113	116	127	135	142	85	271	133	119	162
Animal Manure Applied to Soils	54	50	51	54	56	56	57	56	53	55	53
Crop Residue	13	14	15	15	15	16	16	17	18	18	18
Cultivation of histosols	195	214	220	223	232	242	245	250	274	268	270
2. Pasture, Range and Paddock Manure	64	64	65	68	70	66	68	66	67	68	67
3. Indirect Emissions											
Atmospheric Deposition	19	18	19	20	21	20	21	21	19	20	20
Nitrogen Leaching and Run-off	79	74	75	80	85	85	69	124	82	79	90
<i>Continued</i>	2012										
1. Direct Emissions											
Synthetic Fertiliser	139										
Animal Manure Applied to Soils	53										
Crop Residue	18										
Cultivation of histosols	268										
2. Pasture, Range and Paddock Manure	67										
3. Indirect Emissions											
Atmospheric Deposition	20										
Nitrogen Leaching and Run-off	84										

Time series consistency

The N₂O emissions from agricultural soils have increased from 6.4 tonnes N₂O in 1990 to 14.5 tonnes N₂O in 2008. The more than doubled emission is a consequence of a significant increase in use of nitrogen in synthetic fertiliser. In 2012 N₂O emissions from agricultural soils decreased primarily due to a fall in the use of synthetic fertiliser.

Table 16.6.22 Emissions of N₂O from Agricultural Soils 1990–2012, tonnes N₂O.

CRF – Table 4.D	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
Total N ₂ O emission	6.4	6.4	5.9	5.2	5.7	5.9	9.1	7.2	10.5	10.5	9.1
1. Direct Emissions											
Synthetic Fertiliser	0.2	0.2	0.2	0.2	0.2	0.1	2.0	0.5	2.6	3.1	2.3
Animal Manure Applied on Soil	1.0	1.0	0.9	0.8	0.9	1.0	1.0	1.2	1.0	1.1	1.1
Crop Residue	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
Cultivation of Histosols	0.2	0.2	0.2	0.2	0.2	0.3	0.3	0.3	0.3	0.3	0.3
2. Pasture, Range and Paddock Manure	2.6	2.6	2.4	2.0	2.2	2.3	2.4	2.5	2.6	2.0	1.9
3. Indirect Emissions											
Atmospheric Deposition	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3
Nitrogen Leaching and Run-off	1.9	1.9	1.8	1.5	1.7	1.8	2.9	2.2	3.4	3.5	3.0
<i>Continued</i>	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011
Total N ₂ O emission	9.5	9.0	9.2	9.8	10.3	10.4	8.7	14.5	10.1	9.7	11.0
1. Direct Emissions											
Synthetic Fertiliser	2.4	2.2	2.3	2.5	2.6	2.8	1.7	5.3	2.6	2.3	3.2
Animal Manure Applied on Soil	1.1	1.0	1.0	1.1	1.1	1.1	1.1	1.1	1.0	1.1	1.0
Crop Residue	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.4	0.4	0.4
Cultivation of Histosols	0.3	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.5	0.4	0.5
2. Pasture, Range and Paddock Manure	2.0	2.0	2.0	2.1	2.2	2.1	2.1	2.1	2.1	2.1	2.1
3. Indirect Emissions											
Atmospheric Deposition	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3
Nitrogen Leaching and Run-off	3.1	2.9	2.9	3.2	3.3	3.4	2.7	4.9	3.2	3.1	3.6
<i>Continued</i>	2012										
Total N ₂ O emission	10.3										
1. Direct Emissions											
Synthetic Fertiliser	2.7										
Animal Manure Applied on Soil	1.0										
Crop Residue	0.4										
Cultivation of Histosols	0.5										
2. Pasture, Range and Paddock Manure	2.1										
3. Indirect Emissions											
Atmospheric Deposition	0.3										
Nitrogen Leaching and Run-off	3.3										

16.6.6 Uncertainties

A tier 1 uncertainty assessment has been carried out in accordance with the IPCC GPG (IPCC, 2000). The uncertainty has been estimated for all sources included in the reporting for agricultural sector. The uncertainties for the activity data and emission factors are shown in Table 16.6.23.

Table 16.6.23 Uncertainties for activity data and emission factors for agriculture.

Subsector	Pollutant	Activity data uncertainty	Emission factor uncertainty
4A Enteric Fermentation	CH ₄	10	100
4B Manure Management	CH ₄	10	100
4B Manure Management	N ₂ O	10	100
4D1 Direct N ₂ O emissions from agricultural soils	N ₂ O	20	50
4D2 Pasture range and paddock	N ₂ O	20	25
4D3 Indirect N ₂ O emissions from agricultural soils	N ₂ O	20	50

The resulting uncertainties for the individual greenhouse gases and the total uncertainty on the greenhouse gas emission are shown in Table 16.6.24.

Table 16.6.24 Uncertainties for the emission estimates.

	Uncertainty %	Trend 1990-2012 %	Trend uncertainty %
GHG	± 59	6.4	± 17
CH ₄	± 98	-10.5	± 12
N ₂ O	± 31	46.2	± 28

16.6.7 Source specific QA/QC

The elaboration of a formal QA/QC plan is to be completed.

However, data on livestock, land-use categories, synthetic fertilisers and cultivation of histosols has gone through a great deal of quality work with regard to accuracy, comparability and completeness.

All external data used for the emission inventory submission are archived in spreadsheets. Data are archived annually in order to ensure that the basic data for a given report are always available in their original form.

Annual data on livestock, land-use categories, synthetic fertilisers and cultivation of histosols are compared with previous years and large discrepancies are checked.

Safely stored and quality checked activity data are then processed by using a methodological approach consistent with international guidelines.

Calculated emission factors are compared with guideline emission factors to ensure that they are reasonable. The calculations follow the principle in international guidelines.

During data processing, it is checked that calculations are being carried out correctly. However, a documentation plan for this needs to be elaborated.

Time series for activity data, emission factors and calculated emissions are used to identify possible errors in the calculation procedure. In fact, during the calculation, numerous controls take place to ensure correctness. Sums are checked of the various stages in the calculation procedure. Implied emission factors are compared to emission factors.

Every single time series imported to the CRF Reporter is checked for annual activity, units for activity, emission factor and emissions. Additional checks are performed on the database. The database encloses every single activity data, emission factors, emission, notation key and comment imported to the CRF Reporter. In other words, no information is typed manually into the CRF Reporter. Instead, all information is imported to the CRF Reporter through XML-files to ensure maximum accuracy and completeness.

16.6.8 Source specific recalculations and improvements

Table 16.6.25 shows recalculations in the agricultural sector compared with the 2013 submission. In this 2012 submission the use of synthetic fertilisers in 2011 has been changed compared to the 2013 submission. Therefore Table 16.6.25 shows a change in the recalculations with regard to the emissions in 2011.

Table 16.6.25 Changes in GHG emission in the agricultural sector compared with the 2013 submission.

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
Previous inventory, Gg CO ₂ eqv.	8.8	8.9	8.0	7.0	7.7	8.2	9.4	9.4	10.1	9.5	8.8
Recalculated, Gg CO ₂ eqv.	8.8	8.9	8.0	7.0	7.7	8.2	9.4	9.4	10.1	9.5	8.8
Change in Gg CO ₂ eqv.	-	-	-	-	-	-	-	-	-	-	-
Change in pct.	-	-	-	-	-	-	-	-	-	-	-
<i>Continued</i>	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011
Previous inventory, Gg CO ₂ eqv.	9.1	8.7	8.8	9.3	9.7	9.5	9.2	10.8	9.3	9.3	8.7
Recalculated, Gg CO ₂ eqv.	9.1	8.7	8.8	9.3	9.7	9.5	9.2	10.8	9.3	9.3	9.6
Change in Gg CO ₂ eqv.	-	-	-	-	-	-	-	-	-	-	0.9
Change in pct.	-	-	-	-	-	-	-	-	-	-	10.3
<i>Continued</i>	2012										
Previous inventory, Gg CO ₂ eqv.	-										
Recalculated, Gg CO ₂ eqv.	9.3										
Change in Gg CO ₂ eqv.	-										
Change in pct.	-										

16.6.9 Source specific planned improvements

The Greenlandic emission inventory for the agricultural sector largely meets the request as set down in the IPCC Good Practice Guidance. Thus for the moment improvements especially concern the QA/QC practice.

16.6.10 References

Greenland Agricultural Consulting Service: Statistics on livestock (sheep and reindeer) and background data on land use (cropland and grassland).

Bussink, D.W. 1994: Relationship between ammonia volatilisation and nitrogen fertilizer application rate, intake and excretion of herbage nitrogen by cattle on grazed swards. *Fertil. Res.* 38, 111-121.

CRF, Common Reporting Format. Available at http://cdr.eionet.europa.eu/dk/Air_Emission_Inventories

Djurhuus, J. & Hansen, E.M. 2003: Notat vedr. tørstof og kvælstof i efterladte planterester for landbrugsjord – af 21. maj 2003. Forskningscenter Foulum, Tjele. (In Danish).

EMEP/EEA Guidebook, 2009: Draft version of the new EMEP/EEA

IPCC, 1997: Revised 1996 IPCC Guidelines for National Greenhouse Gas Inventories. Available at <http://www.ipcc-ggip.iges.or.jp/public/gl/invs1.html> (08-02-2010).

IPCC, 2000: Good Practice Guidance and Uncertainty Management in National Greenhouse Gas Inventories. Available at <http://www.ipcc-nggip.iges.or.jp/public/gp/english/> (08-02-2010).

Jarvis, S.C., Hatch, D.J. & Roberts, D.H., 1989a: The effects of grassland management on nitrogen losses from grazed swards through ammonia volatilization; the relationship to extral N returns from cattle. *J. Agric. Sci. Camb.* 112,205-216.

Jarvis, S.C., Hatch, D.J. & Lockyer, D.R., 1989b: Ammonia fluxes from grazed grassland annual losses from cattle production systems and their relation to nitrogen inputs. *J. Agric. Camp.* 113, 99-108.

Nielsen, O.-K., Plejdrup, M.S., Winther, M., Nielsen, M., Gyldenkerne, S., Mikkelsen, M.H., Albrektsen, R., Thomsen, M., Hjelgaard, K., Hoffmann, L., Fauser, P., Bruun, H.G., Johannsen, V.K., Nord-Larsen, T., Vesterdal, L., Møller, I.S., Caspersen, O.H., Rasmussen, E., Petersen, S.B., Baunbæk, L., Hansen, M.G., 2013: Denmark's National Inventory Report 2013 – Emission Inventories 1990-2012 - Submitted under the United Nations Framework Convention on Climate Change and the Kyoto Protocol. National Environmental Research Institute, University of Aarhus. 1,203 pp. – Scientific Report from DCE – Danish Center for Environment and Energy no. 56. Available at:

<http://www2.dmu.dk/Pub/SR56.pdf>

Rasmussen, H.E., 1992: Reindeer Management in Southern Greenland (1992). Published in Danish as “Den sydgrønlandske rendrift” (1992).

Statistics Greenland, Stat Bank: Data on average temperature in June in Southern Greenland. Available at:

<http://bank.stat.gl/database/Greenland/Energy/Climate/Climate.asp>

Statistics Greenland, Statistical Yearbook 1990-2013: Data on livestock (sheep and reindeer).

16.7 LULUCF (CRF sector 5)

16.7.1 Overview of LULUCF

This LULUCF chapter covers only the territory of Greenland. Greenland is part of the Danish Kingdom.



Figure 16.7.1 Municipalities and major cities in Greenland.

Greenland is the world's largest non-continental island located on the northern American continent between the Arctic Ocean and the North Atlantic Ocean, northeast of Canada. The northernmost point of Greenland, Cape Morris Jesup, is only 740 km from the North Pole. The southernmost point is Cape Farewell, which lies at about the same latitude as Oslo in Norway. Geographical coordinates are 72 00 N, 40 00 W.

Greenland is covering approximately 2 166 086 km². It has been estimated that 81 % is covered permanently with ice leaving only 410 449 km² ice free. The distance from the South to the North is 2 670 km, and from East to West 1 050 km.

The terrain is flat to gradually sloping ice cap, which covers all but a narrow, mountainous, barren, rocky coast. The ice cap is up to 3 km thick, and contains 10 per cent of the world's resources of freshwater.

The climate is arctic to sub-arctic with cool winters and cold summers in which the mean temperature does not exceed 10° C.

The mean temperature in January is for Nuuk, -8.6°, Kangerlussuaq, -17.0° and Ilulissat -9.6° (2007) and for July: Nuuk 7.7°, Kangerlussuaq 11.5° and Ilulissat 9.6° (2007).

Greenland is normally defined as having three different climatic zones. For the purpose of reporting is used the definition “Polar and Moist” according to IPCC 2006 Guidelines although some areas may qualify as arctic deserts.

The sparse population is confined to small settlements along the coast, but close to one-quarter of the population lives in the capital, Nuuk. The total population in January 2012 was 56 749 inhabitants.

Due to the cold climate and the small constant population there is almost no land use change occurring. The total area with Forests has been estimated to 218.5 hectares and 10.5 hectares with Cropland. Grassland is divided into improved Grassland covering 1060 hectares and unimproved Grassland covering 241 000 hectares. Wetlands consist of man made water reservoirs - in total 1076 hectares. Settlements cover 5655 hectares. Land classified as “Other Land” is then 99.9 % of the total area.

In the following text the abbreviations are used in accordance with definitions in the IPCC guidelines:

A:	Afforestation, areas with forest established after 1990 under Article 3.3.
R:	Reforestation, areas which have temporarily been unstocked for less than 10 years - included under Article 3.4.
D:	Deforestation, areas where forests are permanently removed to allow for other land use, included under Article 3.3.
FF:	Forest remaining Forest, areas remaining forest after 1990.
FL:	Forest Land meeting the definition of forests.
CL:	Cropland.
GL:	Grassland.
SE:	Settlements.
OL:	Other land, unclassified land.
FM:	Forest Management, areas managed under Article 3.4.
CM:	Cropland Management, areas managed under Article 3.4.
GM:	Grazing land Management, areas managed under Article 3.4.

The LULUCF sector differs from the other sectors in that it contains both sources and sinks of carbon dioxide. LULUCF are reported in the CRF format. Removals are given as negative figures and emissions are reported as positive figures in accordance with the guidelines.

In total the LULUCF sector has been estimated as a net source of 1.32 Gg CO₂ equivalents in 2012 equivalent to 0.2 % of the total Greenlandic emission.

The overall land use change from 1990 to 2012 is very small. Afforestation has been made on 14 hectares. No deforestation has occurred and the Cropland area has increased from none to 10.5 hectares.

The emission data are reported in the new CRF format under IPCC categories 5A (Forestry), 5B (Cropland), 5C (Grassland), 5D (Wetlands) and 5E (Settlements) and 5F (Other Land).

Fertilisation of forests and other land is not occurring and all fertiliser consumption is therefore reported in the agricultural sector. No drainage of forest soils is made. All liming is reported under Grassland because liming is not occurring in the forests and the very small area with Cropland. Field burning of wooden biomass is not occurring. Wildfires may occur sporadic in the mountains and these are reported as "Other land". Hence wildfires are reported as NO.

Table 16.7.1 gives an overview of the emission from the LULUCF sector in Greenland. The Forests are a net sink. Cropland is ranging from being zero in 1990 (no Cropland was occurring in 1990) to being a net source in 2010. GL has been estimated to be a small net source in 2010 due a decrease in the improved area with grassland compared to 2009, which has decreased the amount of living biomass in GL.

Table 16.7.1 Overall emission (Gg CO₂) from the LULUCF sector in Greenland, 1990-2010.

Greenhouse gas source and sink categories	1990	1995	2000	2005	2007	2008	2009	2010	2011	2012
5. Land Use, Land-Use Change and Forestry, CO ₂	0,21	0,39	0,53	0,64	0,96	0,86	0,15	1,42	1,21	1,32
A. Forest Land	NA	-0,02	-0,03	-0,05	-0,05	-0,06	-0,03	-0,04	-0,04	-0,04
B. Cropland	IE,NA,NIE,NA,NO O	IE,NA,N O	0,02	0,02	0,02	0,03	0,03	0,03	0,05	0,05
C. Grassland	0,21	0,41	0,56	0,66	0,98	0,89	0,16	1,43	1,20	1,32
D. Wetlands	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO
E. Settlements	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO
F. Other Land	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA

16.7.2 Forest remaining forest (5.A.1)

Forests and forest management

Greenland has virtually no forests and therefore there exist no official forest statistics. All forests are situated in the most southern part of Greenland. In an attempt to introduce trees to Greenland research were carried out to find species adaptable to the Greenlandic climate. This resulted in establishment of the Greenlandic Arboretum, which covers 150 hectares out of the total area of 218.5 hectares, Figure 16.7.2 and Table 16.7.2. Information about the Greenlandic Arboretum can be found at:

<http://ign.ku.dk/english/about/arboreta/arboretum-greenland/>



Figure 16.7.2 The position of the Greenlandic forests (Courtesy to Rasmus Enoksen Christensen).

Table 16.7.2 Forests in Greenland 1990 and 2012.

Location	Established	Dominant tree	Area, ha	1990 average tree height (m)	2012 average tree height	Density 1990 (trees pr ha)	Density 2009
Qinnua Valley	Natural	Birch and mountain ash	45	n.a	6	100	100
Qanassiassat Forest	1953-63	Conifer	1	5	11.5	1500	1000
Kuussuaq Forest	1962-64 -1982	Conifer	5	3	10.8	1300	900
Kuussuaq Forest	2008	Conifer	3	***	< 1	***	3500
Greenland Arboretum	(1976-1980)	Conifer	3	4	7	300	300
Greenland Arboretum	1980 -	Conifer	150	2	3	1500	1700
Itilleq	2004-2005	Conifer	6	***	< 1	***	3500
Upernaviarsuk	1954	Conifer	0,5	1,5	3	200	200
Lejrskolen	1999-2005	Conifer	4	***	1	***	2500
Klosterdalen	2000	Conifer	1	***	1	***	2000
Total			232.5				

Forest definition

The forest definition adopted in Greenland is almost identical to the FAO definition (TBFRA, 2000). It includes “wooded areas larger than 0.5 ha, that are able to form a forest with a height of at least 5 m and crown cover of at least 10 %. The minimum width is 20 m.” Temporarily non wooded areas, fire breaks, and other small open areas, that are an integrated part of the forest, are also included. However, due to extreme slow growing rates many of the forests are currently below 5 meters height.

Figure 16.7.3 shows a picture of the best developed forest in Greenland.




Figure 16.7.3 The forest in Kuusuaq. Photo: Rasmus E. Christensen, 2005.

Of special interest is the forest in Qinngua Valley. The Qinngua Valley is situated in a remote area. It consists of natural birch (*Betula pubescens* spp. *czerepanovii* and *B. glandulosa*.) which develops to forest like trees probably due to an introgressiv hybridisation (Pers. Comm. Rasmus Enoksen Christensen). This forest will probably not follow the FAO forest definition but are included in the inventory as a sub-division under forests. The Qinngua-valley is not included in the FAO forest statistics.



Figure 16.7.4 Kuusuaq, Tasermiut fjor. Photo: Rasmus Christensen, Juni 2004.

Methodological issues for forests

Estimation of volume, biomass and carbon pools

Due to lack of precise data and slow growth rates, simple functions are used that only include the height of the trees and the number per hectare.

The height of the trees has been estimated by Rasmus Enoksen Christensen based on data from the Aboretum. It is assumed that the trees are conical and the stem diameter at ground level is based on the general formula for even-aged forests (Vanclay, 2009).

$$D = \beta(H - 1.3) / \ln(N) \quad (\text{eq.1})$$

Where:

D = diameter at breast height, cm

β = slope, species dependent

H = Height of the trees (meters)

N = Number of trees per hectare

Eq. 1 has been simplified by omitting the breast height (1.3 meters) to

$$D = \beta(H) / \ln(N) \quad (\text{eq.2})$$

so that D is representing the diameter at ground level. The β -value used is given in Table 16.7.2.

Table 16.7.2 β -values for estimating the diameter of trees (from Vanclay, 2009).

	Betula, spp	Conifers
β -values	6.54	7.51

In order to estimate the C stock and C stock change is used the average default values from the IPCC 2006 guidelines for BCEF, density, C-content and Root-Shoot ratio for Boreal stands with a growing stock level of 21-50 m³, IPCC table 4.5, pp 4.50. The values are given in Table 16.7.3.

Table 16.7.3 Biomass expansion factors used for Greenland.

		Qinngua Walley (Betula, spp.) Birch	Conifers	Orpiuteqarfia (Larix sibirica) Siberian Larch
BCEF	Dimensionless	0.7	0.66	0.78
Density	kg dry matter per litre	0.51	0.4	0.46
C-content	kg C per kg dry matter	0.48	0.51	0.51
Root-shoot-ratio	Dimensionless	0.39	0.39	0.39
Dead Organic Matter	kg per kg aboveground biomass	0.1	0.2	0.1

Source: IPCC 2006 guidelines.

Dead wood volume, biomass and carbon

The volume of dead organic matter (DOM) is estimated as a fraction of the aboveground biomass (Table 16.7.3). It is assumed that litter is included in DOM.

Forest soils: forest floors and mineral soil

Following the cold climate and the slow growing rate it is assumed that no changes takes place in C-stock in the soil and hereby following the IPCC 2006 guidelines at Tier 1 level.

Uncertainties and time series consistency

The uncertainty in estimation of the C stock changes in the Greenlandic forests is very high. As there are very limited resources to visit and monitor in the remote areas there are very few data available. The current inventory is therefore based on the best knowledge available. It should also be taken into consideration that the importance of the forest sector in Greenland is marginal as only very little thinning is taking place as well as no deforestation and that the effect on the inventory is almost not measurable.

In the overall uncertainty section for the LULUCF is made a Tier 1 uncertainty analysis.

QA/QC and verification

Focus on the measurements of carbon pools in forest in Greenland will contribute to QA/QC and verification, but at the moment there are no plans to a further monitoring of the Greenlandic forests.

Recalculations and changes made in response to the review process

No recalculations have been made.

Planned improvements

No improvements are planned.

16.7.3 Land converted to forests

Forest area

See Section 16.2.1 Information on approaches used for representing land areas and on land-use databases used for the inventory preparation.

Forest definition

See Section 16.2.1 Land-use definitions and the classification systems used and their correspondence to the LULUCF categories (e.g. land use and land-use change matrix).

Methodological issues for land converted to forest

See also Section 16.2.1.

Since 1990 there has been a slight increase in the forest area of 14 hectares. This has taken place on land converted from "OL".

Uncertainties and time series consistency

See Section 16.2.1. For uncertainties, please see Chapter 16.7.13.

QA/QC and verification

No QA/QC plan has been made yet. The afforested area is known.

Recalculations, including changes made in response to the review process

None

Planned improvements

No improvements are planned.

16.7.4 Cropland – 5B

Cropland and cropland management – 5B1

In 1990 there were no cropland occurring in Greenland. Due to the global warming it is now possible to have a few crops which may mature. In 2001 the first five hectares with annual crops were established. These are reported under 5.B.2. A more intensive description of the agriculture in Greenland can be found at

<http://nunalerineq.gl/english/landbrug/jord/index-jord.htm>

Land converted to cropland – 5B2

In 2001 the first annual crops were grown in Greenland. Approximately five hectares with garden crops were grown. Of this it is assumed that 25 % of the area is on organic soils (pers. comm. with Kenneth Høeg, former

chief agricultural advisor in Greenland). The area converted to cropland was improved grassland.



Figure 16.7.5 Cropland and Grassland in Greenland.
(Photos from: <http://nunalerineq.gl/english/landbrug/landbrug/index-landbrug.htm>).

The region is generally characterized by a slightly podsol type of soil with a low pH value and small amounts of accessible plant nutrients. Larger concentrations of clay rarely occur, but considerable quantities of silt are often observable on the surface. Also, a certain amount of brown earth occurs in inland areas.

Methodological issues

Change in carbon stock in living biomass

For land converted to cropland is used a standard default value of 5 000 kg DM (dry matter) per hectare in above- and below-ground (IPCC 2006).

Change in carbon stock in dead organic matter

No organic matter is reported under CL.

Change in carbon stock in soils

No C stock changes in mineral soils are assumed. The emission in the 25 % organic soils is estimated by using the IPCC 2006 default value for cropland, Table 5.6 pp 5.19 of 5 000 kg C per ha per year.

Uncertainties and time series consistency

The time series are complete. For uncertainties, please see Chapter 16.7.13.

Category-specific QA/QC and verification

The number of hectares is provided by the Greenlandic Agricultural Consulting Services. As agricultural activities are economically subsidised in Greenland the figures are very accurate.

Category-specific recalculation

No recalculations have been made.

Category-specific planned improvements

No improvements are planned.

16.7.5 Grassland – 5C**Grassland remaining grassland – 5C1**

Grassland in Greenland is dominated by unimproved grassland where the sheep is grazing. The total area with GL has been estimated to 242 000 hectares. Of these only approximately 1 000 hectare is improved where stones have been removed combined with sowing of more high yielding species, see Figure 16.7.5.

Since 1990 the area with improved grassland has been extended from 460 hectares to 1060 hectares.

Methodological issues for grassland

Grassland is divided into improved and unmanaged Grassland.

Change in carbon stock in living biomass

As more GL becomes improved the amount of living biomass at peak is increased. To estimate the amount of living biomass in improved GL is using the same default value as for Cropland, e.g. 5 000 kg DM per hectare, IPCC 2006 default value for cropland, Table 5.9 pp 5.28. For unmanaged Grassland is used a default value of 1 700 kg DM per hectare according to IPCC 2006 default, Table 6.4 pp 6.27. No estimates for below-ground biomass are given. For conversion from DM to C is used a default value of 0.5 kg C per kg DM.

Change in carbon stock in dead organic matter

No changes in dead organic matter are estimated as this is not occurring for this category.

Change in carbon stock in soils

No changes in the carbon stock in mineral soils are assumed. For organic soils on improved grassland is used a default EF of 1 250 kg C per ha per year (IPCC, 2006) default value for grassland, Table 6.3 pp 6.17. For unmanaged grassland no carbon stock change is expected.

Uncertainties and time series consistency

The time series is complete. For uncertainties, please see Chapter 16.7.13.

Category-specific QA/QC and verification

The number of hectares is provided by the Greenlandic Agricultural Consulting Services. As the agriculture is subsidised in Greenland the figures are very accurate.

Recalculations

No recalculation has been made.

Planned improvements

No improvements are planned.

16.7.6 Wetlands – 5D

Wetland in Greenland includes only human made water reservoirs and not naturally occurring wetlands. In total 1 076 hectares with ponds and water reservoirs distributed on 48 locations are reported.

No emission estimates from these reservoirs has been made yet.

Uncertainties and time series consistency

Not estimated.

QA/QC and verification

QA and QC have been made by DCE and Statistics Greenland.

Recalculations

No recalculations have been made.

Category-specific planned improvements

No improvements are planned.

16.7.7 Settlements – 5E

In total there are approximately 56 000 inhabitants in Greenland with about one quarter of the population in the capital, Nuuk.

Table 7.4 Inhabitants and the area occupied with houses, hectares.

	1990	2000	2010	2012
Cities, inhabitants	44,427	45,734	47,641	48,232
Small villages, inhabitants	11,131	10,373	8,811	8,517
City area, ha	2,964	3,051	3,355	3,830
Villages, ha	1,825	1,825	1,825	1,825
Settlements, total, ha	4,789	4,876	5,1805	5,655

The cities are build on the rocky coastline where almost none vegetation occurs. As a consequence estimates for C stock in living biomass and in soil have been made.

The small increase in the area with Settlements since 1990 has taken place on “Other land”.

Currently, no official data or measurements of the area of villages and settlements are available. Alternatively, land utilized for villages and settlements have been measured by the use of NunaGIS, which is a digital internet atlas displaying maps over villages and settlements in Greenland. NunaGIS is available at www.nunagis.gl.

16.7.8 Other land

The far major part of Greenland is covered with snow or rocks. Thus Other Land consists of 99.9 % of the total area.

No emission estimates have been made for this area.

The global warming can be seen in Greenland with longer and warmer summers, which again increase the amount of living biomass. Especially

since the early 1990's there has been changes observed in the environment, e.g. as given in the area with Cropland and Grassland has increased. However, no methodology exists currently to estimate a proper estimate of the amount of living biomass in the large area classified as "Other land".

16.7.9 Direct N₂O emissions from N fertilization of Forest Land and Other land use – 5(I)

Not occurring.

16.7.10 Non-CO₂ emissions from drainage of forest soils and wetlands – 5(II)

Not occurring.

16.7.11 N₂O emissions from disturbance associated with land-use conversion to cropland – 5(III)

Not occurring.

16.7.12 CO₂ emissions from agricultural lime application – 5(IV)

As part of the agricultural practice liming is taking place on acidic agricultural soils (Kenneth Høeg, personal communication). The total amount of lime consumed in 2009, based on import statistics, is 5 tonnes lime and 5 tonnes dolomite. This figure is used for the period 2009-2012.

The amount of C is calculated according to the guidelines with a 90 % purity of lime and 95 % purity for dolomite. It is assumed that all C disappear as CO₂ the same year as the lime is applied.

Planned Improvements

None.

16.7.13 Biomass burning – 5(V)

No biomass burning takes place in Greenland, and wildfires rarely occur due to the moist climate.

16.7.14 Uncertainties

A tier 1 uncertainty assessment has been carried out in accordance with the IPCC GPG (IPCC, 2000). The uncertainty has been estimated for all sources included in the reporting for LULUCF. The uncertainties for the activity data and emission factors are shown in Table 16.7.4.

Table 16.7.4 Uncertainties for activity data and emission factors for LULUCF.

Subsector	Pollutant	Activity data uncertainty	Emission factor uncertainty
5A Forest	CO ₂	5	50
5B Cropland	CO ₂	5	50
5C Grassland	CO ₂	5	50

The assumed uncertainties represent expert judgement.

The resulting uncertainties for the individual greenhouse gases and the total uncertainty on the greenhouse gas emission are shown in Table 16.7.5.

Table 16.7.5 Uncertainties for the emission estimates.

	1990	2012					2012	
	Emission/sink, Gg CO ₂ eqv.	Emission/sink, Gg CO ₂ eqv.	Activity data, %	Emission factor, %	Combined uncertainty		Total	Uncertainty 95 %, Gg CO ₂ eqv.
5. LULUCF	-0,7	-1.2					50,0	
5.A Forests	-0,7	1.32					46.9	
Forests CO ₂	0	-0.04					50.2	0.620
5.B Cropland	NA	-0.04					50.2	
Cropland CO ₂	0,0	0.05	5	50	50.2		50.2	0.021
5.C.Grassland	NA	0.05					50.2	
Grassland CO ₂	-0,8	1.32	5	50	50.2		50.2	0.024
5(IV) Liming CO ₂	0.21	1.32					50.2	
CO ₂	0,0	0.004	5	50	50.2		50.2	0.661

16.7.15 References

Christensen, R.E. 2010: Information on Greenlandic forests. Not published.

IPCC, 2006: 2006 IPCC Guidelines for National Greenhouse Gas Inventories, Prepared by the National Greenhouse Gas Inventories Programme, Eggleston, H.S., Buendia, L., Miwa, K., Ngara, T. & Tanabe, K. (eds). Published: IGES, Japan. Available at:

<http://www.ipcc-nggip.iges.or.jp/public/2006gl/index.html>

Vanclay, J.K. 2009: Tree diameter, height and stocking in even-aged forests, Ann. For. Sci. 66. 702 Available online at: EDP Sciences, 2009. Available at: www.afs-journal.org DOI: 10.1051/forest/2009063.

16.8 Waste (CRF sector 6)

16.8.1 Overview of sector

The waste sector consists of the CRF source category 6.A. Solid Waste Disposal on Land, 6.B. Wastewater Handling, 6.C. Waste Incineration and 6.D. Other.

In CO₂ equivalents, the waste sector (without LULUCF) contributes with 3.5 % of the overall greenhouse gas emission (GHG) in 2012. This corresponds to an emission of 21.2 Gg CO₂ equivalents.

The Greenlandic inventory includes CH₄ emissions from solid waste disposal on land, CH₄ and N₂O from wastewater handling and CO₂, CH₄, N₂O, NO_x, CO, NMVOC and SO₂ from waste incineration. Only emissions from waste incineration without energy recovery are included in the waste sector. Emissions from waste incineration with energy recovery are included in the energy sector.

Table 16.8.1 shows the greenhouse gas emissions from the waste sector. The emissions are taken from the CRF tables and are presented as rounded figures.

Table 16.8.1 Emissions for the waste sector, Gg CO₂ equivalents.

		1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
6 A. Solid Waste Disposal on Land	CH ₄	3.6	3.7	3.8	3.8	3.9	3.9	4.0	4.1	4.1	4.1	4.2
6 B. Wastewater Handling	N ₂ O	14.9	14.9	14.9	14.9	14.9	14.9	14.9	14.9	14.9	14.9	14.9
6 C. Waste incineration	CO ₂	2.6	2.6	2.6	2.6	2.7	2.7	2.9	3.1	3.5	3.4	3.2
6 C. Waste incineration	CH ₄	2.3	2.3	2.3	2.3	2.3	2.4	2.3	2.3	2.2	2.1	1.8
6 C. Waste incineration	N ₂ O	1.1	1.1	1.1	1.1	1.1	1.1	1.1	1.1	1.1	1.0	0.9
6. Waste total		24.4	24.5	24.6	24.7	24.9	25.1	25.3	25.5	25.8	25.6	25.0
<i>Continued</i>		2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011
6 A. Solid Waste Disposal on Land	CH ₄	4.1	4.1	4.1	4.1	4.1	4.0	4.0	4.0	3.9	3.9	3.9
6 B. Wastewater Handling	N ₂ O	14.9	14.9	14.9	15.0	14.9	15.0	15.2	15.8	13.0	12.4	12.7
6 C. Waste incineration	CO ₂	3.3	3.2	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1
6 C. Waste incineration	CH ₄	1.8	1.7	1.6	1.6	1.6	1.6	1.6	1.6	1.6	1.6	1.6
6 C. Waste incineration	N ₂ O	0.9	0.9	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.8
6. Waste total		25.1	24.9	24.7	24.5	24.5	24.5	24.6	25.2	22.4	21.9	22.1
<i>Continued</i>		2012										
6 A. Solid Waste Disposal on Land	CH ₄	3.9										
6 B. Wastewater Handling	N ₂ O	11.8										
6 C. Waste incineration	CO ₂	3.1										
6 C. Waste incineration	CH ₄	1.6										
6 C. Waste incineration	N ₂ O	0.8										
6. Waste total		21.2										

The largest source of greenhouse gas emission in 2012 from the waste sector is N₂O emission from wastewater handling (56 %), more specifically from industrial effluents. Other large sources are CH₄ from solid waste disposal on land (18 %) and CO₂ from waste incineration (15 %).

The total greenhouse gas emission from the waste sector has decreased by 13.1 % from 1990 to 2012. In 2012 emissions from all sources except wastewater handling were more or less unchanged. However, N₂O from wastewater handling decreased by 6.9 %.

16.8.2 Solid waste management

Activity data for waste amounts for solid waste management are shown in Table 16.8.2.

Table 16.8.2 Waste amounts for solid waste management, tonnes.

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
6 A1. Managed Waste Disposal	6 056	6 124	6 168	6 232	6 334	6 428	6 410	6 416	6 145	5 697	4 876
6 A2. Unmanaged Waste Disposal	1 362	1 359	1 358	1 360	1 341	1 289	1 217	1 160	1 060	988	910
6 C. Waste incin., energy recovery	5 519	5 578	5 618	5 733	5 918	6 072	6 178	6 275	6 398	8 200	11 279
6 C. Waste incin., without energy rec.	16 566	16 713	16 808	16 955	17 195	17 460	17 828	18 162	18 756	17 827	16 068
6. Waste total	29 503	29 775	29 952	30 280	30 788	31 249	31 633	32 014	32 360	32 712	33 132
<i>continued</i>	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011
6 A1. Managed Waste Disposal	4 943	4 746	4 451	4 215	4 246	4 264	4 293	4 312	4 346	4 413	4 476
6 A2. Unmanaged Waste Disposal	868	843	835	828	826	818	791	763	746	722	692
6 C. Waste incin., energy recovery	11 526	12 658	14 084	15 312	15 572	15 788	16 056	16 366	16 686	17 077	17 500
6 C. Waste incin., without energy rec.	16 285	15 874	15 220	14 700	14 790	14 836	14 823	14 778	14 837	14 955	15 028
6. Waste total	33 623	34 121	34 589	35 055	35 435	35 705	35 964	36 220	36 614	37 168	37 695
<i>continued</i>	2012										
6 A1. Managed Waste Disposal	4 503										
6 A2. Unmanaged Waste Disposal	658										
6 C. Waste incin., energy recovery	17 854										
6 C. Waste incin., without energy rec.	15 027										
6. Waste total	38 043										

The waste amounts are based on municipal data on waste and waste incineration with energy recovery on local incinerator plants in 2004, and a survey by Consulting Company Carl Bro in 1996 and 2001, where waste amounts per person per year was identified as 650 kg and 455 kg for Greenlandic towns and villages, respectively. For the time series these amounts were regulated by 1 % per year upwards for years after 2004 and by 1 % per year downwards for years before 2004. Further, to construct the time series statistical data from Statistics Greenland on population in towns and villages were used. Other results of the survey used for the time series are that it was estimated that (1) 70 % of waste amounts is incinerated and 30 % deposited and (2) 80 % of combustible waste amounts deposited is burned in open burning.

Solid waste disposal

Source Category Description

The category consists of managed and unmanaged disposal of waste on land.

Methodological issues, activity data, emission factors and emissions

In Table 16.8.3 the composition of the waste according to the survey mentioned is shown.

Table 16.8.3 Composition of household and commercial waste before and after open burning.

Fraction	Household waste ²	Commercial waste ²	Household / Commercial Weighted %	After open burning	Weighted (after open burning)
Paper/cardboard, dry	8.00 ¹	20.00	11.84	2.37	7.66
Paper/cardboard, wet	10.00 ¹	7.00	9.04	1.81	5.85
Plastics	7.00 ¹	9.00	7.64	1.53	4.94
Organic waste	44.00 ¹	34.00	40.80	8.16	26.40
Other combustible	17.50 ¹	16.00	17.02	3.40	11.00
Glass	7.50 ¹	3.00 ¹	6.06	6.06	19.60
Metal	3.50 ¹	3.00 ¹	3.34	3.34	10.80
Other, non combustible	1.00 ¹	5.00	2.28	2.28	7.37
Hazardous waste	1.50 ¹	3.00 ¹	1.98	1.98	6.40
Total	100.00	100.00	100.00	30.93	100.00
Pct (%)	68 ³	32 ³		80 ⁴	

Notes:

¹ Measured values.

² Source: Former Environmental and Nature Agency, Ministry of Infrastructure and Environment. Survey from 2004.

³ Distribution of household and commercial waste.

⁴ Share of combustible waste burned at waste disposal sites.

A Tier 2 approach with a first order decay model is used for estimation of emissions of CH₄ from the solid waste disposals. For this purpose the activity data in Table 16.8.2 are estimated back to 1960 (not shown) based on the methodology described in connection to Table 16.8.2. Combining these activity data and the composition data in Table 16.8.3 time series for 1960-2012 with amounts of waste in waste fractions is calculated.

For these time series the waste fractions are associated to (1) Dissolved Organic Carbon (DOC) values according to Section 16.8.2 of this NIR and (2) emission factors based on DOC values and values of methane correction factors, fraction of DOC dissimilated and fraction of CH₄ in gas emitted according to the IPCC GL and GPG for managed disposals, Table 16.8.4 and unmanaged disposal, Table 16.8.5.

Table 16.8.4 DOC values and emission factors for CH₄ for managed disposals.

	Paper / cardboard, dry	Paper / cardboard, wet	Plastics	Organic waste	Other combustible	Glass	Metal	Other, non combustible	Hazardous waste
DOC weighted (after open burning) fraction	0.40	0.20	0.00	0.20	0.20	0.00	0.00	0.00	0.00
Emission factor kg CH ₄ /tonnes ¹	133.3	66.7	0.0	66.7	66.7	0.0	0.0	0.0	0.0
¹) based on:									
Methane correction factor	1								
Fraction of DOC dissimilated and emitted	0.5								
Fraction of CH ₄ in gas emitted	0.5								

Table 16.8.5 DOC values and emission factors for CH₄ for unmanaged disposals.

	Paper/ cardboard dry	Paper/ cardboard wet	Plastics	Organic waste	Other combustible	Glass	Metal	Other, non- combustible	Hazardous waste
DOC weighted (after open burn- ing) fraction	0.40	0.20	0.00	0.20	0.20	0.00	0.00	0.00	0.00
Emission factor kg CH ₄ /tonnes ¹	53.3	26.7	0.0	26.7	26.7	0.0	0.0	0.0	0.0
¹) based on:									
Methane correction factor				0.4					
Fraction of DOC dissimilated and emitted				0.5					
Fraction of CH ₄ in gas emitted				0.5					

For managed and unmanaged disposals the default half life time of 14 years and a time lag of 0.5 years are used. For the oxidation factor and according to the GPG for managed disposal 0.1 and for unmanaged 0.0 are used.

In Tables 16.8.6 and 16.8.7 selected data and results are shown for 1990-2012 for managed and unmanaged disposal, respectively. The data in the tables are as follows. The AD for the FOD model as amounts of waste in fractions, the potential emission of CH₄ calculated with emission factors on waste amounts in fractions, the annual generated emission of CH₄ calculated with the FOD model using the potential emissions, the oxidized CH₄ and the actual annual CH₄ emission calculated as the annual generated emission minus the CH₄ oxidized. Calculations are performed since 1960 and are not shown.

Table 16.8.6 Managed disposal. AD for the FOD model (amounts of waste in fractions), the potential emission of CH₄, the oxidized CH₄ and the annual CH₄ emission for 1990-2012.

	Paper /cardboard dry	Paper /cardboard wet	Plastics	Organic waste	Other combustible	Glass	Metal	Other, non combustible	Hazardous waste	Waste total	Potential emission	Annual generated emission	Annual oxidized emission	Annual emission
Unit	Tonnes	Tonnes	Tonnes	Tonnes	Tonnes	Tonnes	Tonnes	Tonnes	Tonnes	Tonnes	Tonnes CH ₄	Tonnes CH ₄	Tonnes CH ₄	Tonnes CH ₄
1990	464	354	299	1 598	667	1 187	654	446	388	6 056	232.7	174.8	17.5	157.3
1991	469	358	303	1 616	674	1 200	661	451	392	6 124	236.4	177.8	17.8	160.0
1992	472	361	305	1 627	679	1 209	666	455	395	6 168	239.0	180.7	18.1	162.6
1993	477	364	308	1 644	686	1 221	673	459	399	6 232	240.8	183.6	18.4	165.3
1994	485	370	313	1 671	697	1 241	684	467	405	6 334	243.3	186.5	18.6	167.8
1995	492	376	318	1 696	708	1 260	694	474	412	6 428	247.2	189.4	18.9	170.5
1996	491	375	317	1 691	705	1 256	692	473	410	6 410	250.9	192.4	19.2	173.2
1997	491	375	317	1 693	706	1 257	693	473	411	6 416	250.2	195.2	19.5	175.7
1998	471	359	304	1 621	676	1 204	664	453	393	6 145	250.5	197.9	19.8	178.1
1999	436	333	281	1 503	627	1 116	615	420	365	5 697	239.9	199.9	20.0	179.9
2000	373	285	241	1 286	537	955	527	359	312	4 876	222.4	201.0	20.1	180.9
2001	378	289	244	1 304	544	969	534	364	316	4 943	190.3	200.5	20.0	180.4
2002	363	277	234	1 252	522	930	513	350	304	4 746	193.0	200.1	20.0	180.1
2003	341	260	220	1 174	490	872	481	328	285	4 451	185.3	199.4	19.9	179.4
2004	323	246	208	1 112	464	826	455	311	270	4 215	173.7	198.1	19.8	178.3
2005	325	248	210	1 120	467	832	459	313	272	4 246	164.5	196.5	19.7	176.9
2006	326	249	211	1 125	469	836	460	314	273	4 264	165.7	195.0	19.5	175.5
2007	329	251	212	1 133	473	841	464	316	275	4 293	166.4	193.6	19.4	174.3
2008	330	252	213	1 138	475	845	466	318	276	4 312	167.6	192.4	19.2	173.2
2009	333	254	215	1 147	478	852	469	320	278	4 346	168.3	191.2	19.1	172.1
2010	338	258	218	1 164	486	865	477	325	283	4 413	169.6	190.2	19.0	171.2
2011	343	262	221	1 181	493	877	483	330	287	4 476	172.3	189.3	18.9	170.4
2012	345	263	222	1 188	496	882	486	332	288	4 503	174.7	188.6	18.9	169.8

Table 16.8.7 Unmanaged disposal. AD for the FOD model (amounts of waste in fractions), the potential emission of CH₄, the oxidized CH₄ and the annual CH₄ emission for 1990-2012.

	Paper /cardboard dry	Paper /cardboard wet	Plastics	Organic waste	Other combustible	Glass	Metal	Other, non combustible	Hazardous waste	Waste total	Potential emission	Annual generated emission	Annual oxidized emission	Annual emission
Unit	Tonnes	Tonnes	Tonnes	Tonnes	Tonnes	Tonnes	Tonnes	Tonnes	Tonnes	Tonnes	Tonnes CH ₄	Tonnes CH ₄	Tonnes CH ₄	Tonnes CH ₄
1990	104	80	67	359	150	267	147	100	87	1 362	21	15.8	0.0	15.8
1991	104	79	67	359	150	266	147	100	87	1 359	21	16.1	0.0	16.1
1992	104	79	67	358	149	266	147	100	87	1 358	21	16.3	0.0	16.3
1993	104	79	67	359	150	266	147	100	87	1 360	21	16.6	0.0	16.6
1994	103	78	66	354	148	263	145	99	86	1 341	21	16.8	0.0	16.8
1995	99	75	64	340	142	253	139	95	83	1 289	21	17.0	0.0	17.0
1996	93	71	60	321	134	238	131	90	78	1 217	20	17.1	0.0	17.1
1997	89	68	57	306	128	227	125	86	74	1 160	19	17.2	0.0	17.2
1998	81	62	52	280	117	208	115	78	68	1 060	18	17.3	0.0	17.3
1999	76	58	49	261	109	194	107	73	63	988	17	17.2	0.0	17.2
2000	70	53	45	240	100	178	98	67	58	910	15	17.2	0.0	17.2
2001	66	51	43	229	96	170	94	64	56	868	14	17.0	0.0	17.0
2002	65	49	42	222	93	165	91	62	54	843	14	16.8	0.0	16.8
2003	64	49	41	220	92	164	90	62	53	835	13	16.7	0.0	16.7
2004	63	48	41	218	91	162	89	61	53	828	13	16.5	0.0	16.5
2005	63	48	41	218	91	162	89	61	53	826	13	16.3	0.0	16.3
2006	63	48	40	216	90	160	88	60	52	818	13	16.2	0.0	16.2
2007	61	46	39	209	87	155	85	58	51	791	13	16.0	0.0	16.0
2008	58	45	38	201	84	150	82	56	49	763	12	15.8	0.0	15.8
2009	57	44	37	197	82	146	81	55	48	746	12	15.6	0.0	15.6
2010	55	42	36	191	80	142	78	53	46	722	12	15.4	0.0	15.4
2011	53	40	34	183	76	136	75	51	44	692	11	15.2	0.0	15.2
2012	50	38	32	174	72	129	71	48	42	658	11	15.0	0.0	15.0

16.8.3 Wastewater handling

Source category description

In Greenland no wastewater treatment occurs; although it should be mentioned some filtering of solid residues from industry may occur and likewise there are ongoing projects focussing on septic tanks at household levels. N₂O emission from human sewage is estimated. It is assumed that no methane emission occurs.

Methodological issues

According to the IPCC Guidelines (IPCC, 1997) the important factors for CH₄ production from handling of wastewater are: wastewater characteristics, handling systems, temperature and BOD vs. COD.

The Guidelines state that production of CH₄ generally requires temperatures above 15°C, and at temperatures below this the lagoon is principally a sedimentation tank (IPCC, 1997). Temperatures in Greenland rarely exceed 15°C, and the monthly average temperature has not exceeded 12°C during the period 1993-2011. Therefore CH₄ is reported as Not Applicable in the CRF.

N₂O emission from wastewater handling

The IPCC default methodology only includes N₂O emissions from human sewage based on annual per capita protein intake. The methodology account for nitrogen intake ("outcome"), i.e. faeces and urine, only and neither the industrial nitrogen input nor non-consumption protein from kitchen, bath and laundry discharges are included.

The formula used for calculation of the emission from effluent WWTP discharges is:

$$E_{\text{effluents}} = P \cdot F_N \cdot N_{\text{pop}} \cdot F_{\text{nc}} \cdot F \cdot EF \cdot \text{effluent} \cdot \frac{M_{\text{N}_2\text{O}}}{M_{\text{N}_2}}$$

where P is the annual protein per capita consumption per person per year set constant to 171.5 g/day (see below text).

F_N is the fraction of nitrogen in protein, i.e. 0.16 (IPCC, 1997).

N_{pop} is the Greenlandic population (source: Statistics Greenland).

F_{nc} is the fraction of the population not connected to the municipal sewer system, i.e. set to 1 as no wastewater treatment plants exists in Greenland at this point.

F is the fraction of non-consumption protein in domestic wastewater. i.e. 1.1 (IPCC, 2006).

$EF_{\text{N}_2\text{O.WWTP.effluent}}$ is the IPCC GL default emission factor of 0.01 kg N₂O-N/kg sewage-N produced (IPCC, 1996)¹.

$M_{\text{N}_2\text{O}}$ and M_{N_2} are the mass ratio. i.e. 44/28 to convert the discharged units in mass of total N to emissions in mass N₂O.

¹ The IPCC (2006) gives a default value for the N₂O emissions from domestic wastewater nitrogen effluent of 0.005 (0.0005 - 0.25) kg N₂O-N/kg N. However, the IPCC EF from the 1996 guidelines has been used.

For households

A large part of the diet originates from seafood, fish or sea mammals, but imported fabricated foods are expected to continue to take over an increasing part of human energy consumption. Due to weather conditions most of fresh food comes from wild animals or fish. Greenland has a production of lamb and a limited supply of vegetables; still most of the produced foods are imported from outside (Mulvad et al., 2007).

In Greenland, the traditional diet based on meat and fish has undergone diversification towards more carbohydrates with the development of a monetary economy; in 1855 the protein content of a mean diet was 377 g protein, whereas 80 years later, in 1935 – 43, the protein content of a mean diet was 257 g protein (Périssé and François, 1981). Today, the majority of young urbanised Greenlandic Inuit have Western dietary habits and consume less meat from marine mammals, terrestrial mammals and birds than Inuit from the hunting districts; Dietary profiles of Canadian Baffin Island Inuit with a high consumption of traditional foods have shown a mean daily protein intake of 144-199 g/day in 41- to 61-year-old (Laursen et al, 2001).

As no data on the protein intake are available a protein intake of 171.5 g/day, i.e. the average of the Canadian Inuit were adopted, as it is assumed that the protein intake has declined even more since 1935 due to increased number of urbanised Greenlandic Inuit. For comparison the Danish yearly protein consumption according to FAOSTAT has increased from 98 g/day in 1990 to 112 g/day in 2005. Using this number, the yearly protein intakes may be derived by multiplying with the population number and days in a year. Based on the above it was decided to set the protein intake to the average value of the Canadian Inuit data, 171.5 g/day. The N-content in effluent wastewater in Greenland was calculated the equation shown above.

From industries

The production of residue products from the fish industry in Greenland amounts to around 14,000 tons per year (Nielsen et al, 2005). Overall the waste amount from the Greenland halibut production is around 40 %, while the waste amount from codfish production is 50 %; this governs only the fish production including pre-processing.

According to IPCC, the fraction of nitrogen in protein is 0.16 (IPCC, 1996). The IPCC reports a range of 0.3 to 3.1 kg total N/ton fish referring to effluent loads from cod filleting; i.e. 0.0031. The report also presents values of the total N content of untreated wastewater from the fish industry in the range of 400-1000 mg/l corresponding to a fraction of corresponding. However, as it was not possible to find data for all fish groups, and as it was not possible to determine that fraction of fish, which was pre-processed and how big a fraction that was sold without pre-processing, the below approach was adopted.

From the EC BAT note (EC, 2003) the total N-content of untreated wastewater from the fishing industry was reported to be between 400 and 1000 mg/L with an average value of 700 mg/L. The number was multiplied by the water used within the fishing industry reported for 2004 to 2011 by Statistics Greenland. The effluent N-content for 1990 to 2002 was set equal to the estimated value for 2003.

Emissions

Emission of N₂O from wastewater handling is shown in Table 16.8.8.

Table 16.8.8 N₂O emissions from households and industries 1990-2012.

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
N ₂ O emission, effluents households, Gg	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010
N ₂ O emission, effluents industries, Gg	0.038	0.038	0.038	0.038	0.038	0.038	0.038	0.038	0.038	0.038	0.038
N ₂ O emission, effluents sum, Gg	0.048	0.048	0.048	0.048	0.048	0.048	0.048	0.048	0.048	0.048	0.048
<i>Continued</i>	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011
N ₂ O emission, effluents households, Gg	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010
N ₂ O emission, effluents industries, Gg	0.038	0.038	0.038	0.038	0.038	0.039	0.039	0.041	0.032	0.030	0.031
N ₂ O emission, effluents sum, Gg	0.048	0.048	0.048	0.048	0.048	0.049	0.049	0.051	0.042	0.040	0.041
<i>Continued</i>	2012										
N ₂ O emission, effluents households, Gg	0.010										
N ₂ O emission, effluents industries, Gg	0.028										
N ₂ O emission, effluents sum, Gg	0.038										

Total emission of N₂O increased until 2008 due to an increase in the emission from industrial effluents. However, in 2009-2012 total emission of N₂O decreased to a total level of 0.038-0.042 Gg (which is lower than 1990) due to a temporarily decrease in industrial effluents primarily caused by a decrease in the catches of shrimps.

16.8.4 Waste incineration

Source category description

In Greenland waste incineration is carried out both with and without energy recovery. According to IPCC Guidelines the emissions associated with waste incineration for energy production is included in the energy sector more specifically in the source category 1.A1a Public Electricity and Heat Production. The emissions from waste incineration without energy recovery is reported in source category 6.C. Waste Incineration. Additionally in Greenland open burning of waste occurs at landfill sites. Emissions associated with this are also reported under sector 6.C. Waste Incineration.

Methodological issues

The methodology used follows the IPCC Guidelines. For waste incineration the Danish emission factors are used, as it is trusted that they are also a good representation of Greenlandic conditions.

Neither the revised 1996 IPCC Guidelines (IPCC, 1997) nor the Good Practice Guidance (IPCC, 2000) contains a methodology for estimating emissions from open burning, therefore the methodology provided in the 2006 IPCC Guidelines (IPCC, 2006) is used.

The emission factors used for both waste incineration and open burning are included in Section 16.8.4.4.

Activity data

The amount of waste incinerated without energy recovery is presented in Table 16.8.9. The activity data is provided by the method described in Section 16.8.2.

Table 16.8.9 Activity data for waste incineration without energy recovery, Mg.

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
Waste incinerated without energy recovery, Mg	NO	NO	NO	NO	56	225	795	1 240	2 663	2 896	3 148
<i>continued</i>	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011
Waste incinerated without energy recovery, Mg	3 306	3 391	3 415	3 437	3 461	3 485	3 468	3 444	3 466	3 486	3 488
<i>continued</i>	2012										
Waste incinerated without energy recovery, Mg	3 501										

The open burning of waste is assumed to be 80 % of the waste deposited to landfills (Survey on waste by Carl Bro, 1996 and 2001). The activity data for open burning is presented in Table 8.10. The activity data for open burning is provided by the method described in Section 16.8.2.

Table 16.8.10 Activity data for open burning of waste, Mg.

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
Open burning of waste, Mg	16 566	16 713	16 808	16 955	17 140	17 235	17 033	16 922	16 093	14 930	12 920
<i>continued</i>	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011
Open burning of waste, Mg	12 979	12 483	11 804	11 263	11 329	11 350	11 355	11 335	11 371	11 470	11 540
<i>continued</i>	2012										
Open burning of waste, Mg	11 526										

Emission factors

Waste incineration

For waste incineration without energy recovery the same emission factors have been assumed as for waste incineration with energy recovery. The emission factors refer to the Danish emission factors (Nielsen et al., 2010). The greenhouse gas emission factors are shown in Table 16.8.11.

Table 16.8.11 Emission factors for greenhouse gases from waste incineration.

	Emission factor	Unit
CO ₂	32.5	Kg pr GJ
CH ₄	30	g pr GJ
N ₂ O	4	g pr GJ

The emission factors used for the indirect greenhouse gases are shown in Table 16.8.12.

Table 16.8.12 Emission factors for indirect greenhouse gases from waste incineration.

	NO _x	SO ₂	NM VOC	CO	Unit
Waste incineration	100	6	50	1 000	g pr GJ

Open burning

For open burning emissions are calculated using the methodology, standard parameters and emission factors provided by the 2006 IPCC Guidelines.

The CH₄ emission factor used is the recommended and default is 6,500 g per tonne MSW wet weight. This factor refers to US EPA (2001).

For N₂O a default emission factor of 150 g/t MSW dry weight is recommended (IPCC, 2006) this is corrected for the dry matter content to acquire an N₂O emission factor of 214 g per tonne MSW wet weight.

For calculating the CO₂ emission the dry matter content, carbon content and the fossil carbon content of the waste fractions are used. The parameters are included in Table 16.8.13.

Table 16.8.13 Parameter used in calculating CO₂ emissions from open burning.

	Dry matter content	Total carbon content, %	Fossil carbon content as percent of total carbon
Paper	0,9	46	1
Cardboard	0,9	46	1
Plastics	1,0	75	100
Organic waste	0,4	38	0
Other	0,9	3	100

Source: 2006 IPCC Guidelines, Volume 5, Chapter 2, Table 2.4

An oxidation factor of 58 % is assumed for open burning (IPCC, 2006).

The emission factors for NO_x, SO₂, NMVOC and CO are presented in Table 16.8.14. The emission factors are from the US EPA (1992).

Table 16.8.14 Emission factors for indirect greenhouse gases from open burning of waste.

	NO _x	SO ₂	NMVOC	CO	Unit
Open burning of municipal refuse	3	0.5	15	42	Kg pr Mg

Emissions

Total emission of greenhouse gases from sector 6.C. Waste Incineration is shown in Table 16.8.15. Figure 16.8.2 shows total emission of greenhouse gases from sector 6.C. Waste incineration is shown in Figure 16.8.1.

Table 16.8.15 Greenhouse gas emissions from waste incineration.

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
CO ₂ , Gg	2.6	2.6	2.6	2.6	2.7	2.7	2.9	3.1	3.5	3.4	3.2
CH ₄ , Mg	107.7	108.6	109.2	110.2	111.4	112.1	111.0	110.4	105.4	98.0	85.0
N ₂ O, Mg	3.5	3.6	3.6	3.6	3.7	3.7	3.7	3.7	3.6	3.3	2.9
CO ₂ eqv., Gg	5.9	6.0	6.0	6.0	6.1	6.2	6.4	6.5	6.8	6.5	5.9
<i>Continued</i>	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011
CO ₂ , Gg	3.3	3.2	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1	3.1
CH ₄ , Mg	85.4	82.2	77.8	74.3	74.7	74.9	74.9	74.8	75.0	75.7	76.1
N ₂ O, Mg	2.9	2.8	2.7	2.6	2.6	2.6	2.6	2.6	2.6	2.6	2.6
CO ₂ eqv., Gg	6.0	5.8	5.6	5.4	5.5	5.5	5.5	5.4	5.5	5.5	5.5
<i>Continued</i>	2012										
CO ₂ , Gg	3.1										
CH ₄ , Mg	76.0										
N ₂ O, Mg	2.6										
CO ₂ eqv., Gg	5.5										




Figure 16.8.1 Emission of greenhouse gases from waste incineration.

The emissions of indirect greenhouse gases from waste incineration are shown in Table 16.8.16.

Table 16.8.16 Emissions of indirect greenhouse gases from waste incineration, Mg.

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
NO _x	49.7	50.1	50.4	50.9	51.5	51.9	51.9	52.1	51.1	47.8	42.1
SO ₂	8.3	8.4	8.4	8.5	8.6	8.6	8.6	8.5	8.2	7.6	6.6
NMVOC	248.5	250.7	252.1	254.3	257.1	258.6	255.9	254.5	242.8	225.5	195.5
CO	695.8	701.9	705.9	712.1	720.4	726.1	723.7	723.7	703.9	657.5	575.7
<i>Continued</i>	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011
NO _x	42.4	41.0	39.0	37.4	37.6	37.7	37.7	37.6	37.8	38.1	38.3
SO ₂	6.7	6.4	6.1	5.8	5.9	5.9	5.9	5.9	5.9	5.9	6.0
NMVOC	196.4	189.0	178.9	170.7	171.7	172.1	172.1	171.8	172.4	173.9	174.9
CO	579.8	559.9	531.6	509.1	512.1	513.3	513.3	512.2	514.0	518.3	521.3
<i>Continued</i>	2012										
NO _x	38.3										
SO ₂	6.0										
NMVOC	174.7										
CO	520.9										

16.8.5 Uncertainties

A tier 1 uncertainty assessment has been carried out in accordance with the IPCC GPG (IPCC, 2000). The uncertainty has been estimated for all sources included in the reporting for the waste sector. The uncertainties for the activity data and emission factors are shown in Table 16.8.17.

Table 16.8.17 Uncertainties for activity data and emission factors for the waste sector.

Subsector	Pollutant	Activity data uncertainty	Emission factor uncertainty
6C Waste incineration	CO ₂	10	25
6A Solid Waste Disposal on Land	CH ₄	10	100
6C Waste incineration	CH ₄	10	50
6B Wastewater Handling	N ₂ O	30	100
6C Waste incineration	N ₂ O	10	100

The amount of waste incinerated and burned is relatively well known and the uncertainty is set to 10 %. The same is the case for the waste deposited to landfills. For waste water handling an uncertainty of 30 % on the activity data has been assumed.

Regarding the emission factor uncertainty, a value of 100 % has been used for CH₄ from solid waste disposal, N₂O from wastewater treatment and N₂O from waste incineration. This is in the same range as recommended by the IPCC GPG. For CO₂ and CH₄ from waste incineration emission factor uncertainties of 25 % and 50 % respectively have been chosen.

The resulting uncertainties for the individual greenhouse gases and the total uncertainty on the greenhouse gas emission are shown in Table 16.8.18.

Table 16.8.18 Uncertainties for the emission estimates.

	Uncertainty %	Trend 1990-2012 %	Trend uncertainty %
GHG	± 61	-13.1	± 21.5
CO ₂	± 27	22.9	± 17.4
CH ₄	± 73	-6.8	± 14.0
N ₂ O	± 98	-21.1	± 31.3

16.8.6 Source specific QA/QC

The elaboration of a formal QA/QC plan is to be completed.

However, data on solid waste disposal, waste water handling and waste incineration has gone through a great deal of quality work with regard to accuracy, comparability and completeness.

All external data used for the emission inventory submission are archived in spreadsheets. Data are archived annually in order to ensure that the basic data for a given report are always available in their original form.

Annual data on solid waste disposal, waste water handling and waste incineration are compared with previous years and large discrepancies are checked.

Safely stored and quality checked activity data are then processed by using a methodological approach consistent with international guidelines.

Calculated emission factors are compared with guideline emission factors to ensure that they are reasonable. The calculations follow the principle in international guidelines.

During data processing, it is checked that calculations are being carried out correctly.

Time series for activity data, emission factors and calculated emissions are used to identify possible errors in the calculation procedure. In fact, during the calculation, numerous controls take place to ensure correctness. Sums are checked in the various stages in the calculation procedure. Implied emission factors are compared to emission factors.

Every single time series imported to the CRF Reporter is checked for annual activity, units for activity, emission factor and emissions. Additional checks are performed on the database. The database encloses every single activity data, emission factors, emission, notation key and comment imported to the CRF Reporter. In other words, no information is typed manually into the CRF Reporter. Instead, all information is imported to the CRF Reporter through XML-files to ensure maximum accuracy and completeness.

16.8.7 Source specific recalculations and improvements

The sector *Waste Water Handling* (6B) was included in the inventory for the first time in the 2010 emission inventory submission. No improvements have been carried out regarding waste water handling in the 2014 submission.

Table 16.8.19 shows recalculations in the waste sector compared to the 2013 submission. No changes occurs.

Table 16.8.19 Changes in GHG emission in the waste sector compared with the 2013 submission.

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
Previous inventory, Gg CO ₂ eqv.	24.4	24.5	24.6	24.7	24.9	25.1	25.3	25.5	25.8	25.6	25.0
Recalculated, Gg CO ₂ eqv.	24.4	24.5	24.6	24.7	24.9	25.1	25.3	25.5	25.8	25.6	25.0
Change in Gg CO ₂ eqv.	-	-	-	-	-	-	-	-	-	-	-
Change in pct.	-	-	-	-	-	-	-	-	-	-	-
<i>continued</i>	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011
Previous inventory, Gg CO ₂ eqv.	25.1	24.9	24.7	24.5	24.5	24.5	24.6	25.2	22.4	21.9	22.1
Recalculated, Gg CO ₂ eqv.	25.1	24.9	24.7	24.5	24.5	24.5	24.6	25.2	22.4	21.9	22.1
Change in Gg CO ₂ eqv.	-	-	-	-	-	-	-	-	-	-	-
Change in pct.	-	-	-	-	-	-	-	-	-	-	-
<i>continued</i>	2012										
Previous inventory, Gg CO ₂ eqv.	-										
Recalculated, Gg CO ₂ eqv.	21.2										
Change in Gg CO ₂ eqv.	-										
Change in pct.	-										

16.8.8 Source specific planned improvements

Some planned improvements to the emission inventories are discussed below.

1) Improved data on solid waste disposals

In future inventories attempts will be made in order to improve data on solid waste disposals in general. Statistics Greenland has encouraged the mu-

municipal technical departments with responsibility for waste handling to start gathering data on the yearly amounts of waste handled.

2) Improved data on waste water handling

In future inventories attempts will be made in order to improve data on waste water handling in general. However, at the moment the municipal technical departments seem to have no data on waste water handling at all.

16.8.9 References

EC, 2003: Reference Document on BAT in the food, drink and milk industry. Available at:

<http://gasunie.eldoc.ub.rug.nl/FILES/root/2003/2665985/2665985.pdf>

IPCC, 1997: Revised 1996 IPCC Guidelines for National Greenhouse Gas Inventories. Available at:

<http://www.ipcc-nggip.iges.or.jp/public/gl/invs1.html>

IPCC, 2000: Good Practice Guidance and Uncertainty Management in National Greenhouse Gas Inventories. Penman, J., Kruger, D., Galbally, I., Hiraishi, T., Nyenzi, B., Emmanuel, S., Buendia, L., Hoppaus, R., Martinsen, T., Meijer, J., Miwa, K. & Tanabe, K. (Eds). Published: IPCC/OECD/IEA/IGES, Hayama, Japan. Available at:

<http://www.ipcc-nggip.iges.or.jp/public/gp/english/index.html>

IPCC, 2006: 2006 IPCC Guidelines for National Greenhouse Gas Inventories, Prepared by the National Greenhouse Gas Inventories Programme, Eggleston H.S., Buendia L., Miwa K., Ngara T. and Tanabe K. (eds). Published: IGES, Japan. Available at:

<http://www.ipcc-nggip.iges.or.jp/public/2006gl/index.html>

Laursen, J., Milman, N., Pedersen, H.S., Mulvad, G., Saaby, H. and Byg, K.-E., 2001: Elements in autopsy liver tissue samples from Greenlandic Inuit and Danes. III. zinc measured by X-ray fluorescence spectrometry. *Journal of Trace Elements in Medicine and Biology*, 15:209-214.

Mulvad, G., Petersen, H.S. and Olsen, J., 2007: Arctic health problems and environmental challenges in Greenland. In: Ørbæk, J.B., Kallenborn, R., Tombre, I., Hegseth, E.N., Falk-Petersen, S. and Hoel, A.H. *Arctic Alpine Ecosystems and People in a Changing Environment*. Available at:

<http://www.springerlink.com/content/tu377x6715200027/fulltext.pdf>

Nielsen M., Nielsen, O.-K. & Thomsen, M. 2010: Emissions from decentralised CHP plants 2007 - Energinet.dk Environmental project no. 07/1882. Project report 5 - Emission factors and emission inventory for decentralised CHP production. National Environmental Research Institute, Aarhus University. 113 pp. - NERI Technical report No. 786. Available at:

<http://www.dmu.dk/Pub/FR786.pdf> (2011-02-28).

Nielsen, O.-K., Plejdrup, M.S., Winther, M., Nielsen, M., Gyldenkerne, S., Mikkelsen, M.H., Albrechtsen, R., Thomsen, M., Hjelgaard, K., Hoffmann, L., Fauser, P., Bruun, H.G., Johannsen, V.K., Nord-Larsen, T., Vesterdal, L., Møller, I.S., Caspersen, O.H., Rasmussen, E., Petersen, S.B., Baunbæk, L., Hansen, M.G., 2013: Denmark's National Inventory Report 2013 - Emission Inventories 1990-2012 - Submitted under the United Nations Framework Convention on Climate Change and the Kyoto Protocol. National Environ-

mental Research Institute, University of Aarhus. 1,203 pp. – Scientific Report from DCE – Danish Center for Environment and Energy no. 56. Available at: <http://www2.dmu.dk/Pub/SR56.pdf>

Nielsen, U., Nielsen, K., Mai, P., Frederiksen, O., 2006: Organisk industriaffald i Grønland – Værktøjer til fremme af bedste tilgængelige teknik og nyttiggørelse af restprodukter nr. M. 127/001-0164. Realistiske muligheder for nyttiggørelse/udnyttelse af organisk industriaffald i Grønland. (In Danish)

Périssé, J. and François, P., 1981: Variability of the protein-calorie ratio in diets. FAO/WHO/UNUEPR/81/40. Available at: <http://www.fao.org/docrep/MEETING/004/AA054E/AA054E00.htm>

Statistics Greenland, 2013: Statistics on Population 1960-2013. Available at: <http://bank.stat.gl/database/Greenland/Population/Population%20in%20Greenland/Population%20in%20Greenland.asp>

US EPA, 1992: AP 42, Fifth Edition, Volume I Chapter 2.5: Open burning. Available at: <http://www.epa.gov/ttn/chief/ap42/ch02/final/c02s05.pdf>

US EPA, 2001: US-EPA Emission Inventory Improvement Program. Volume III Chapter 16 Open Burning. United States Environmental Protection Agency (USEPA). Available at: http://www.epa.gov/ttn/chief/eiip/techreport/volume03/iii16_apr2001.pdf

16.9 Other

In CRF Sector 7, there are no activities and emissions or removals for the inventory of Greenland.

16.10 Recalculations and improvements

The 2014 submission is the fifth year where Greenland on the request of the ERT submits a full CRF.

For recalculations and improvements please refer to Sections 16.3 - 16.8 and Section 16.11.

16.11 KP-LULUCF

16.11.1 General information

In the following text, the abbreviations used are in accordance with definitions in the IPCC guidelines:

A:	Afforestation
R:	Reforestation
D:	Deforestation
FF:	Forest remaining Forest, areas remaining forest after 1990
FL:	Forest Land meeting the Danish definition of forests
CL:	Cropland
GL:	Grassland
SE:	Settlements
OL:	Other land, unclassified land
FM:	Forest Management, areas managed under article 3.4

CM: Cropland Management, areas managed under article 3.4
GM: Grazing land Management, areas managed under article 3.4

Definition of forest and any other criteria

For the estimation of anthropogenic emissions by sources and removals by sinks associated with afforestation (A), reforestation (R) and deforestation (D) since 1990 under Article 3.3 and forest management (FM) under Article 3.4 of the Kyoto Protocol, the following forest definition will be applied:

- Minimum values for tree crown cover: 10 % tree crown cover for forests.
- Minimum values for land area: 0.5 ha.
- Minimum value for tree height: trees must be able to reach a minimum height of 5 m in the site.

In addition, the forest area includes temporarily unstocked areas, smaller open areas in the forest needed for management purposes and fire breaks. Forests in national parks, reserves or areas under special protection are included. Windbreaks and groves covering more than 0.5 ha and with a minimum width of 20 m are also considered as forests.

Woody biomass does not exist outside the forest and hence not reported under Cropland and Grassland.

Elected activities under Article 3, paragraph 4, of the Kyoto Protocol

As regards the possibility of including in the first commitment period emissions and removals associated with land use, land-use change and forestry activities under Article 3.4 of the Kyoto Protocol, it has been decided to include emissions and removals from forest management (FM), cropland management (CM) and grazing land management (GM).

The national system has identified land areas associated with the activities under Article 3.4 of the Kyoto Protocol in accordance with definitions, modalities, rules and guidelines relating to land use, land-use change and forestry activities under the protocol by Land Parcel Information System (LPIS), detailed crop information data on field level, soil mapping and forest maps.

Inventories of emissions and removals under Article 3.3 and Article 3.4 are prepared and reported annually together with the other greenhouse gas inventory information.

Description of how the definitions of each activity under Article 3.3 and each elected activity under Article 3.4 have been implemented and applied consistently over time

The definition of afforestation, reforestation and deforestation is in accordance with the GPG (IPCC 2003).

Afforestation or reforestation is identified when areas have wooded tree cover and fulfils the forest definition given above. The time of the AF is given by the time of action, i.e. planting of trees. No deforestation and reforestation is reported for Greenland as this is not occurring. All types of establishment of forest (AF or RF) are considered human induced.

As no reforestation has taken place Table 5(KP-I)A.1.2 "Units of land harvested since the beginning of the commitment period" is filled in as included elsewhere although it is not occurring.

As for the forest management (Article 3.4), the forest areas fulfilling the definition given above are included under this activity. All forest areas are considered managed except for the remote Qinnngua-valley.

For Cropland and Grassland the area accounted for under Art. 3.4 has been estimated with the best knowledge from the Greenlandic Agricultural Consulting Services. As the agriculture in Greenland is economically subsidized the area is estimated with a high accuracy. Only areas that are reported as CL and GL are included in the accounted area.

Description of precedence conditions and/or hierarchy among article 3.4 activities and how they have been consistently applied in determining how land was classified
All Forest activities have precedence, after this Cropland activities and then Grassland activities.

Afforestation has precedence. All land converted to forest are included as afforested area. Deforested areas are not reported as this is not occurring. The following categories in the Convention reporting are included under afforestation:

- 5A25 OL to A

FM activities are only related to:

- 5A1 Forest remaining Forest

CM activities are related to:

- 5B22 GL to CL

GM activities area related to:

- 5C1 GL remaining GL

No elected land has left land that is not accounted for. Land conversion between elected activities (FM, CM and GM) has been allowed but is currently not occurring. No land elected under article 3.4 activities has been converted to Other Land. Other land converted to elected activities is included in the respective category. As a consequence there has been a steady increase in the land which is accounted for under article 3.3 and 3.4 with 111 hectares from 1990 to 2011.

The Land Use matrix developed for the purpose of reporting article 3.3 and 3.4 activities for 2008 are shown in Table 16.11.1.

Table 16.11.1 Land Use matrix for art. 3.3 and 3.4 activities in 2012.

To current inventory From previous inventory year		Article 3.3 activities		Article 3.4 activities				Other ⁽⁵⁾	Total area at the beginning of the current inventory year ⁽⁶⁾
		Afforestation and Reforestation	Deforestation	Forest Management (if elected)	Cropland Management (if elected)	Grazing Land Management (if elected)	Revegetation (if elected)		
		(kha)							
Article 3.3 activities	Afforestation and Reforestation	0.01	NO						0.01
	Deforestation		NA						NA
Article 3.4 activities	Forest Management (if elected)		NO	0.20					0.20
	Cropland Management ⁽⁴⁾ (if elected)	NO	NO		0.01	NO	NA		0.01
	Grazing Land Management ⁽⁴⁾ (if elected)	NO	NO		0.00	241.99	NA		241.99
	Revegetation ⁽⁴⁾ (if elected)	NA			NA	NA	NA		NA
Other ⁽⁵⁾		NO	NO	NO	NO	NO	NA	216,366.38	216,366.38
Total area at the end of the current inventory year		0.01	NA,NO	0.20	0.01	241.99	NA	216,366.38	216,608.60

16.11.2 Spatial assessment unit used for determining the areas of the units of land under Article 3.3

Afforestation and reforestation are identified as areas which not were covered by forest in 1990. The increase in the forest area is planted.

Methodology used to develop the land transition matrix

The land use matrix is based on the best available data. No vector maps exist of the individual forests, cropland and grassland.

Maps and/or database to identify the geographical locations, and the system of identification codes for the geographical locations

The forests have been given individual names. For the Cropland and Grassland area no identification has been made.

16.11.3 Afforestation, Reforestation & Deforestation (ARD)

Methods for carbon stock change and GHG emission and removal estimates

For afforestation the carbon stock change in the period 1990 - 2012 is based both on the area of afforestation and the information on species composition.

In the afforestation a steady increase in carbon stock is found.

Description of the methodologies and the underlying assumptions used

See Chapter 16.7.

Justification when omitting any carbon pool or GHG emissions/removals from ARD

C stock changes in the soil is not expected due to the cold climate to occur and hence following the guidelines for a Tier 1 approach. As the afforestation is made by hand planting no damages of the existing soil C is expected to take place.

Information on whether or not indirect and natural GHG emissions and removals have been factored out

No factoring out has been performed in the emission and removal estimates.

Changes in data and methods since the previous submission (recalculations)

No recalculation has been performed.

Uncertainty estimates

Uncertainty estimates are given in Table 16.11.2.

Table 16.11.2 Uncertain estimates for 1990 and 2012 for the KP sector.

	1990			2012		
	Emission, Gg CO ₂ yr-1	Total uncertainty, %	Uncertainty 95 %, Gg CO ₂ eqv.	Emission, Gg CO ₂ yr-1	Total uncertainty, %	Uncertainty 95 %, Gg CO ₂ eqv.
KP total	0.179	55.1	0.1	1.322	78.3	1.035
A. Article 3.3 activities	IE,NA,NO,NR	-	-	IE,NA,NO,NR	-	-
B. Article 3.4 activities	0.179	57.8	0.1	1.322	78.6	1.038
KP A.1.1 Afforestation and Reforestation						
Area subject to the activity	NO	-	-	0.014	-	0.000
Area of organic soils(8)	NO	-	-	0.001	-	-
Net CO ₂ emissions/ removals (9)	NO	-	-	IE,NO,NR	-	-
KP A.2 Deforestation						
Area subject to the activity	NO			NO		
Area of organic soils	NO			NO		
Net CO ₂ emissions/ removals	NO			NO		
KP B.1 Forest Management						
Area subject to the activity	-	-	-	0.205		
Area of organic soils(8)	-	-	-	0.025		
Net CO ₂ emissions/ removals (9)	-	-	-	-0.042	36.0	0.0
KP B.2 Cropland Management						
Area subject to the activity	NO	-	-	0.011		
Area of organic soils(8)	NO	-	-	0.003		
Net CO ₂ emissions/ removals (9)	IE,NA	-	-	0.048	90.6	0.0
KP B.3 Grassland Management						
Area subject to the activity	242.0			242.0		
Area of organic soils(8)	7.368			7.493		
Net CO ₂ emissions/ removals (9)	0.171	57.8	0.1	1.312	83.9	1.1
KP-II 4 Lime consumption						
Total amount of lime applied	18.5			9.3		
Emission	0.008	50.2	0.0	0.004	50.2	0.0
Carbon	0.008	50.2	0.0	0.000	0.0	0.0

Information on other methodological issues

See Chapter 16.7.

The year of the onset of an activity, if after 2008

Not applicable.

16.11.4 Forest Management (FM)**Methods for carbon stock change and GHG emission and removal estimates**

See Chapter 16.7 in LULUCF on "Forest remaining forest (5.A.1)".

Methodologies and the underlying assumptions

See Chapter 16.7 in LULUCF on "Forest remaining forest (5.A.1)".

Omission of pools from FM

C changes in forest soils are omitted and hereby following GPG 2003 guidelines at a Tier 1 level.

Factoring out

No factoring out has been performed.

Recalculations

No recalculation has been performed.

Uncertainty estimates

See Table 16.11.2

Information on other methodological issues

See Chapter 16.7 in LULUCF on "Forest remaining forest (5.A.1)".

The year of the onset of an activity, if after 2008

Not applicable.

16.11.5 Cropland Management (CM)**Methods for carbon stock change and GHG emission and removal estimates**

Methodologies and the underlying assumptions used

The area with agricultural Cropland is reported as the area given in Statistics Greenland.

The same methodology as used in the Convention reporting is used in the KP reporting.

Omission of pool from CM

Aboveground and belowground living biomass, litter and dead organic are only reported for perennial woody crops in accordance with IPCC GPG 2003. No litter and dead organic matter are reported under Cropland as these are not occurring. Therefore only aboveground living biomasses are reported under Cropland. Below-ground biomass is included in above-ground biomass.

Factoring out

No factoring out has been made.

Recalculations

None.

Uncertainty estimates

See Table 16.11.2.

Information on other methodological issues

None.

The year of the onset of an activity, if after 2008

Not applicable.

16.11.6 Grazing land management (GM)**Methods for carbon stock change and GHG emission and removal estimates**

Grazing land is defined as land improved grassland and unmanaged grassland.

Description of the methodologies and the underlying assumptions used

The major part of the grassland is unmanaged (241 000 hectare). Only 1 060 hectares is improved grassland with occasional reseeding and fertiliser application. The methodology used is the default Tier 1. This is in accordance with IPCC GPG 2003 (3.4.1.2.1.2) as the total emission from LULUCF consists of less than 0.12 % of the total emission from Greenland.

Omission of pools from GM

Aboveground and belowground living biomass, litter and dead organic are only reported for perennial woody crops in accordance with IPCC GPG 2003. No litter and dead organic matter are reported under Cropland as these are not occurring. Therefore only aboveground living biomasses are reported under Cropland. Below-ground biomass is included in above-ground biomass.

Factoring out

No factoring out has been made.

Recalculations

No recalculation has been performed.

Uncertainty estimates

See Table 16.11.2.

Information on other methodological issues

None.

The year of the onset of an activity, if after 2008

Not applicable.

16.11.7 Article 3.3

Information that demonstrates that activities under Article 3.3 began on or after 1 January 1990 and before 31 December 2012 and are direct human-induced

All forests in Greenland are planted except for the Qinnngua valley, which is in a remote area.

Information on how harvesting or forest disturbance that is followed by the re-establishment of forest is distinguished from deforestation

No deforestation is occurring and therefore not applicable.

Information on the size and geographical location of forest areas that have lost forest cover but which are not yet classified as deforested

Not applicable.

16.11.8 Article 3.4

Information that demonstrates that activities under Article 3.4 have occurred since 1 January 1990 and are human-induced

Forest Management

In Forest Management all forest areas are under management and changes in carbon stock are hence seen as human induced.

Cropland Management

Due to the cold climate and the recent increase in temperature it has only very recently been possible to grow agricultural crops in Greenland with the first fields established around 2001. Today it is estimated that 10.5 hectares are regularly ploughed.

Grassland Management

Due to the cold climate in Greenland and the recent increase in temperature it has only recently been valuable to introduce management activities in the grassland to increase the crop yield. This is well documented in the Greenlandic subsidiary system to the farmers.

Information relating to Cropland Management, Grazing Land Management and Re-vegetation, if elected, for the base year

No further information is available.

Information relating to Forest Management

No further information is available.

16.11.9 Other information

Key category analysis for Article 3.3 activities and any elected activities under Article 3.4

According to the IPCC Good Practice Guidance for LULUCF a category that is identified as key in the UNFCCC inventory should also be considered key under the Kyoto Protocol (IPCC, 2003).

No LULUCF categories are reported as a key source. The total emission from the LULUCF sector is less than 0.2 % of the total emission from Greenland.

16.11.10 Information relating to Article 6

There are no Article 6 projects (Joint Implementation) on the Greenlandic territory.

16.12 Annex 1 Key categories

A Key Category Analysis (KCA) for year 1990 and 2012 for Greenland has been carried out in accordance with the IPCC Good Practice Guidance. For 1990 a level KCA has been carried out.

The base year in the analysis is the year 1990 for the greenhouse gases CO₂, CH₄, N₂O and 1995 for the greenhouse F-gases HFC, PFC and SF₆. The KCA approach is a Tier 1 quantitative analysis.

The level assessment of the Tier 1 KCA is a ranking of the source categories in accordance to their relative contribution to the national total of greenhouse gases calculated in CO₂ equivalents. The level key categories are found from the list of source categories ranked according to their contribution in descending order. Level key categories are those from the top of the list and of which the sum constitutes 95 % of the national total.

The trend assessment of the Tier 1 KCA is a ranking of the source categories according to their contribution to the trend of the national total of greenhouse gases, calculated in CO₂ equivalents, from the base year to the year under consideration. The trend of the source category is calculated relative to that of the national totals and the trend is then weighted with the contribution, according to the level assessment. The ranking is in descending order. As for the level assessment, the cut-off point for the sum of contribution to the trend is 95 % and the source categories from the top of the list to the cut-off line are trend key categories.

Result of the Key Category Analysis for Greenland for the year 1990 and 2012

The entries in the results of KCA in Tables 16.12.1 to 16.12.3 for the years 1990 and 2012 are composed from CRFs for those years in this report. Note that base-year estimates are not used in the level assessment analysis for year 2012, but are only included in Table 16.12.2 to make it more uniform with Tables 16.12.1 and 16.12.3.

The result of the Tier 1 KCA level assessment for Greenland for 1990 is shown in Table 16.12.1. For the assessment, 5 categories were identified as key categories and marked as shaded, refer Table 16.12.1.

The result of the Tier 1 KCA level assessment for Greenland for 2011 is shown in Table 16.12.2. For the assessment, 6 categories were identified as key categories, refer Table 16.12.2.

The result of the Tier 1 KCA trend assessment for Greenland for 1990/1995-2012 is shown in Table 16.12.3. For the trend assessment, 9 categories were identified as key categories, refer Table 16.12.3. Note that according to the GPG, the analysis implies that contributions to the trend are all calculated as mathematically positive to be able to perform the ranking. The LULUCF activities are in the table included with their sign, i.e. emissions: +, removals: -.

In Table 16.12.4 a summary of Key Category Analysis for Greenland is given for level assessment for year 1990/95 and 2012 and for trend for years 1990-2012. All the categories are listed by sector and key sources are shown with their ranking.

Table 16.12.1 Key Category Analysis base year 1990/1995, level assessment, Tier 1.

Table 7.A1 (of Good Practice Guidance)

Tier 1 Analysis - Level Assessment GRL – inventory

A			B	C	D	E
IPCC Source Categories (LULUCF included)			Direct GHG	Base Year Estimate Ex, o Assessment	Base Year Level Assessment	Base Year Cumulative total of Col. D
				Gg CO ₂ eqv.	Lx, o	
Energy	Combustion excluding transport	Liquid fuels	CO ₂	517.812	0.795	0.795
Energy	Civil aviation		CO ₂	38.321	0.059	0.854
Energy	Road transportation		CO ₂	36.058	0.055	0.909
Energy	Domestic navigation		CO ₂	20.732	0.032	0.941
Waste	Wastewater handling		N ₂ O	14.884	0.023	0.964
Agriculture	Enteric fermentation		CH ₄	6.018	0.009	0.973
Waste	Solid waste disposal on land		CH ₄	3.636	0.006	0.979
Waste	Waste incineration		CO ₂	2.550	0.004	0.983
Waste	Waste incineration		CH ₄	2.261	0.003	0.986
Energy	Combustion excluding transport	Other fuels	CO ₂	1.674	0.003	0.989
Energy	Combustion excluding transport		N ₂ O	1.392	0.002	0.991
Agriculture	Direct emissions from agricultural soils		N ₂ O	1.281	0.002	0.993
Waste	Waste incineration		N ₂ O	1.099	0.002	0.995
Energy	Combustion excluding transport		CH ₄	0.950	0.001	0.996
Agriculture	Indirect emissions from agricultural soils		N ₂ O	0.697	0.001	0.997
Agriculture	Manure management		N ₂ O	0.641	0.001	0.998
Energy	Civil aviation		N ₂ O	0.336	0.001	0.999
Solvents and other product use	Solvents		CO ₂	0.263	0.000	0.999
LULUCF	Grassland remaining grassland		CO ₂	0.214	0.000	0.999
Agriculture	Manure management		CH ₄	0.140	0.000	1.000
Energy	Road transportation		N ₂ O	0.093	0.000	1.000
Energy	Road transportation		CH ₄	0.064	0.000	1.000
Energy	Domestic navigation		N ₂ O	0.053	0.000	1.000
Industry	Consumption of SF ₆		SF ₆	0.036	0.000	1.000
Energy	Domestic navigation		CH ₄	0.030	0.000	1.000
Industry	Consumption of HFC's		HFCs	0.025	0.000	1.000
Energy	Civil aviation		CH ₄	0.006	0.000	1.000
Industry	Road Paving with asphalt		CO ₂	0.000	0.000	1.000
Industry	Asphalt roofing		CO ₂	0.000	0.000	1.000
Industry	Limestone and dolomite use		CO ₂	0.000	0.000	1.000
LULUCF	Forest land remaining forest		CO ₂	0.000	0.000	1.000
LULUCF	Conversion to cropland		CO ₂	0.000	0.000	1.000
Total				651.268	1.000	

Table 16.12.2 Key Category Analysis year 2012, level assessment, Tier 1.

Table 7.A1 (of Good Practice Guidance)

Tier 1 Analysis - Level Assessment GRL – inventory

A		B	C	D	E	F
IPCC Source Categories (LULUCF included)		Direct GHG	Base Year Estimate Ex,o Gg CO ₂ eqv	Year 2012 Estimate Ex,t Gg CO ₂ -eqv	Year 2012 Level Assessment Lx,t	Year 2012 Cumulative total of Col. E
Energy	Combustion excluding transport	Liquid fuels CO ₂	517.812	452.736	0.740	0.740
Energy	Civil aviation	CO ₂	38.321	46.669	0.076	0.816
Energy	Road transportation	CO ₂	36.058	33.231	0.054	0.871
Energy	Domestic navigation	CO ₂	20.732	29.741	0.049	0.919
Waste	Wastewater handling	N ₂ O	14.884	11.793	0.019	0.939
Industry	Consumption of HFC's	HFCs	0.025	7.185	0.012	0.950
Energy	Combustion excluding transport	Other fuels CO ₂	1.674	6.936	0.011	0.962
Agriculture	Enteric fermentation	CH ₄	6.018	5.386	0.009	0.970
Waste	Solid waste disposal on land	CH ₄	3.636	3.898	0.006	0.977
Waste	Waste incineration	CO ₂	2.550	3.135	0.005	0.982
Agriculture	Direct emissions from agricultural soils	N ₂ O	1.281	2.069	0.003	0.985
Waste	Waste incineration	CH ₄	2.261	1.597	0.003	0.988
Energy	Combustion excluding transport	N ₂ O	1.392	1.401	0.002	0.990
LULUCF	Grassland remaining grassland	CO ₂	0.214	1.316	0.002	0.992
Agriculture	Indirect emissions from agricultural soils	N ₂ O	0.697	1.113	0.002	0.994
Energy	Combustion excluding transport	CH ₄	0.950	0.928	0.002	0.996
Waste	Waste incineration	N ₂ O	1.099	0.810	0.001	0.997
Agriculture	Manure management	N ₂ O	0.641	0.646	0.001	0.998
Energy	Civil aviation	N ₂ O	0.336	0.409	0.001	0.999
Solvents and other product use	Solvents	CO ₂	0.263	0.230	0.000	0.999
Agriculture	Manure management	CH ₄	0.140	0.126	0.000	0.999
Energy	Road transportation	CH ₄	0.064	0.104	0.000	0.999
Energy	Road transportation	N ₂ O	0.093	0.087	0.000	1.000
Energy	Domestic navigation	N ₂ O	0.053	0.077	0.000	1.000
LULUCF	Conversion to cropland	CO ₂	0.000	0.048	0.000	1.000
Energy	Domestic navigation	CH ₄	0.030	0.043	0.000	1.000
LULUCF	Forest land remaining forest	CO ₂	0.000	-0.042	0.000	1.000
Industry	Limestone and dolomite use	CO ₂	0.000	0.020	0.000	1.000
Energy	Civil aviation	CH ₄	0.006	0.007	0.000	1.000
Industry	Consumption of SF ₆	SF ₆	0.036	0.003	0.000	1.000
Industry	Asphalt roofing	CO ₂	0.000	0.000	0.000	1.000
Industry	Road Paving with asphalt	CO ₂	0.000	0.000	0.000	1.000
Total			651.268	611.702	1.000	

Table 16.12.3 Key Category Analysis years 1990/1995-2012, trend assessment, Tier 1.

Table 7.A1 (of Good Practice Guidance)

Tier 1 Analysis - Trend Assessment GRL – inventory

A			B	C	D	E	F	G
IPCC Source Categories (LULUCF included)			Direct GHG	Base Year Estimate Ex,o Gg CO ₂ -eq	Year 2012 Estimate Ex,t Gg CO ₂ -eq	Trend Assessment Tx,t	Contribution to Trend	Cumulative total of col. F
Energy	Combustion excluding transport	Liquid fuels CO ₂		517.812	452.736	0.052	0.448	0.448
Energy	Civil aviation	CO ₂		38.321	46.669	0.016	0.142	0.590
Energy	Domestic navigation	CO ₂		20.732	29.741	0.016	0.137	0.726
Industry	Consumption of HFC's	HFCs		0.025	7.185	0.011	0.095	0.822
Energy	Combustion excluding transport	Other fuels CO ₂		1.674	6.936	0.008	0.071	0.893
Waste	Wastewater handling	N ₂ O		14.884	11.793	0.003	0.029	0.922
LULUCF	Grassland remaining grassland	CO ₂		0.214	1.316	0.002	0.015	0.937
Agriculture	Direct emissions from agricultural soils	N ₂ O		1.281	2.069	0.001	0.012	0.949
Waste	Waste incineration	CO ₂		2.550	3.135	0.001	0.010	0.959
Energy	Road transportation	CO ₂		36.058	33.231	0.001	0.008	0.967
Waste	Waste incineration	CH ₄		2.261	1.597	0.001	0.007	0.974
Waste	Solid waste disposal on land	CH ₄		3.636	3.898	0.001	0.006	0.981
Agriculture	Indirect emissions from agricultural soils	N ₂ O		0.697	1.113	0.001	0.006	0.987
Agriculture	Enteric fermentation	CH ₄		6.018	5.386	0.000	0.004	0.990
Waste	Waste incineration	N ₂ O		1.099	0.810	0.000	0.003	0.993
Energy	Civil aviation	N ₂ O		0.336	0.409	0.000	0.001	0.994
Energy	Combustion excluding transport	N ₂ O		1.392	1.401	0.000	0.001	0.996
LULUCF	Conversion to cropland	CO ₂		0.000	0.048	0.000	0.001	0.996
Agriculture	Manure management	N ₂ O		0.641	0.646	0.000	0.001	0.997
Energy	Road transportation	CH ₄		0.064	0.104	0.000	0.001	0.997
LULUCF	Forest land remaining forest	CO ₂		0.000	-0.042	0.000	0.001	0.998
Energy	Combustion excluding transport	CH ₄		0.950	0.928	0.000	0.000	0.998
Industry	Consumption of SF ₆	SF ₆		0.036	0.003	0.000	0.000	0.999
Energy	Domestic navigation	N ₂ O		0.053	0.077	0.000	0.000	0.999
Industry	Limestone and dolomite use	CO ₂		0.000	0.020	0.000	0.000	0.999
Solvents and other product use	Solvents	CO ₂		0.263	0.230	0.000	0.000	1.000
Energy	Domestic navigation	CH ₄		0.030	0.043	0.000	0.000	1.000
Agriculture	Manure management	CH ₄		0.140	0.126	0.000	0.000	1.000
Energy	Civil aviation	CH ₄		0.006	0.007	0.000	0.000	1.000
Energy	Road transportation	N ₂ O		0.093	0.087	0.000	0.000	1.000
Industry	Asphalt roofing	CO ₂		0.000	0.000	0.000	0.000	1.000
Industry	Road Paving with asphalt	CO ₂		0.000	0.000	0.000	0.000	1.000
Total				651.268	611.702		1.000	

Table 16.12.4 Summary of Key Category Analysis for Greenland for level assessment for year 1990/95 and 2012 and for trend for years 1990-2012.

Summary of Key Category analysis for Greenland			GHG	Key categories with number according to ranking in analysis		
IPCC Source Categories (LULUCF included)				Identification criteria		
				Level Tier1 1990	Level Tier1 2012	Trend Tier1 1990-2012
Energy	Combustion excluding transport	Liquid fuels	CO ₂	1	1	1
Energy	Combustion excluding transport	Other fuels	CO ₂			5
Energy	Combustion excluding transport		CH ₄			
Energy	Combustion excluding transport		N ₂ O			
Energy	Road transportation		CO ₂	3	3	
Energy	Road transportation		CH ₄			
Energy	Road transportation		N ₂ O			
Energy	Civil aviation		CO ₂	2	2	2
Energy	Civil aviation		CH ₄			
Energy	Civil aviation		N ₂ O			
Energy	Domestic navigation		CO ₂	4	4	3
Energy	Domestic navigation		CH ₄			
Energy	Domestic navigation		N ₂ O			
Industry	Limestone and dolomite use		CO ₂			
Industry	Asphalt roofing		CO ₂			
Industry	Road Paving with asphalt		CO ₂			
Industry	Consumption of HFC's		HFCs		6	4
Industry	Consumption of SF ₆		SF ₆			
Solvents and other product use	Solvents		CO ₂			
Agriculture	Enteric fermentation		CH ₄			
Agriculture	Manure management		CH ₄			
Agriculture	Manure management		N ₂ O			
Agriculture	Direct emissions from agricultural soils		N ₂ O			8
Agriculture	Indirect emissions from agricultural soils		N ₂ O			
Waste	Solid waste disposal on land		CH ₄			
Waste	Wastewater handling		N ₂ O	5	5	6
Waste	Waste incineration		CO ₂			9
Waste	Waste incineration		CH ₄			
Waste	Waste incineration		N ₂ O			
LULUCF	Forest land remaining forest		CO ₂			
LULUCF	Conversion to cropland		CO ₂			
LULUCF	Grassland remaining grassland		CO ₂			7

16.13 Annex 2 Detailed discussion of methodology and data for estimating CO₂ emission from fossil fuel combustion

Detailed information regarding the methodology and input data used to calculate CO₂ emissions from fossil fuel combustion is included in Section 16.3.

16.14 Annex 3 Other detailed methodological descriptions for individual source or sink categories

All methodological descriptions are included in Sections 16.3 – 16.8 and Section 16.11.

16.15 Annex 4 CO₂ reference approach and comparison with sectoral approach, and relevant information on the national energy balance

See Section 16.3.5.1 of this annex for the results of the comparison between the sectoral and reference approach.

16.16 Annex 5 Assessment of completeness and (potential) sources and sinks of greenhouse gas emissions and removals excluded

16.16.1 GHG inventory

The Greenlandic greenhouse gas emission inventories for 1990-2012 include all sources identified by the Revised 1996 IPCC Guidelines and the 2000 IPCC Good Practice Guidance except the following:

In the Solvent and other product use sector currently no N₂O emissions are included in CRF category 3D, Greenland will try to obtain activity data if they exist for uses of N₂O.

Direct and indirect CH₄ emissions from agricultural soils are not estimated. Direct and indirect soil emissions are considered of minor importance for CH₄. No methodology is recommended in IPCC-GPG.

In the LULUCF sector emissions/removals from wetlands, settlements and other land are currently not estimated due to the lack of available data. The lack of data availability is also an issue for other aspects of LULUCF, e.g. harvested wood products. For more detail please see Section 16. 7.

In the Waste sector CO₂ emissions from managed waste disposal on land are not estimated. According to the 1996 IPCC Guidelines: "Decomposition of organic material derived from biomass sources (e.g., crops, forests), which are re-grown on an annual basis is the primary source of CO₂ released from waste. Hence, these CO₂ emissions are not treated as net emissions from waste in the IPCC Methodology."

16.16.2 KP-LULUCF inventory

The KP-LULUCF inventory is considered complete. The carbon pools not estimated has been documented as not being sources, please see Section 16.11 for further documentation.

16.17 Annex 6 Additional information to be considered as part of the annual inventory submission and the supplementary information required under Article 7, paragraph 1, of the Kyoto Protocol or other useful reference information

No additional information for Greenland is deemed relevant.

16.18 Annex 7 Tables 6.1 and 6.2 of the IPCC good practice guidance

IPCC Source category	Gas	Base year emission	Year t emission	Activity data uncertainty	Emission factor uncertainty	Combined uncertainty	Combined uncertainty as % of total national emissions in year t	Type A sensitivity	Type B sensitivity	Uncertainty in trend in national emissions introduced by emission factor uncertainty	Uncertainty in trend in national emissions introduced by activity data uncertainty	Uncertainty introduced into the trend in total national emissions
		Input data Gg CO ₂ eq	Input data Gg CO ₂ eq	Input data %	Input data %	%	%	%	%	%	%	%
1A Liquid fuels	CO ₂	613	562	2	2	2.828	2.600	-0.020	0.864	-0.040	2.442	2.443
1A Municipal waste	CO ₂	2	7	2	25	25.080	0.284	0.008	0.011	0.206	0.030	0.208
1A Liquid fuels	CH ₄	1	1	2	100	100.020	0.158	0.000	0.001	0.001	0.004	0.004
1A Municipal waste	CH ₄	0	0	2	100	100.020	0.009	0.000	0.000	0.006	0.000	0.006
1A Biomass	CH ₄	0	0	2	100	100.020	0.011	0.000	0.000	0.008	0.000	0.008
1A Liquid fuels	N ₂ O	2	2	2	500	500.004	1.423	0.000	0.003	0.026	0.008	0.027
1A Municipal waste	N ₂ O	0	0	2	500	500.004	0.086	0.000	0.000	0.062	0.000	0.062
1A Biomass	N ₂ O	0	0	2	200	200.010	0.042	0.000	0.000	0.030	0.001	0.030
1B2 Oil exploration	CO ₂	0	0	2	1 000	1000.002	0.000	0.000	0.000	0.000	0.000	0.000
1B2 Oil exploration	CH ₄	0	0	2	1 000	1000.002	0.000	0.000	0.000	0.000	0.000	0.000
1B2 Oil exploration	N ₂ O	0	0	2	1 000	1000.002	0.000	0.000	0.000	0.000	0.000	0.000
2A3 Limestone and dolomite use	CO ₂	0	0	5	5	7.071	0.000	0.000	0.000	0.000	0.000	0.000
2A5 Asphalt roofing	CO ₂	0	0	5	25	25.495	0.000	0.000	0.000	0.000	0.000	0.000
2A6 Road paving with asphalt	CO ₂	0	0	5	25	25.495	0.000	0.000	0.000	0.000	0.000	0.000
2F Consumption of HFC	HFC	0	7	10	50	50.990	0.599	0.011	0.011	0.550	0.156	0.572
2F Consumption of SF ₆	SF ₆	0	0	10	50	50.990	0.000	0.000	0.000	-0.002	0.000	0.002
3A Paint application	CO ₂	0	0	10	15	18.028	0.000	0.000	0.000	0.000	0.000	0.000
3B Degreasing and dry cleaning	CO ₂	0	0	10	15	18.028	0.000	0.000	0.000	0.000	0.000	0.000
3C Chemical products, manufacturing and processing	CO ₂	0	0	10	15	18.028	0.000	0.000	0.000	0.000	0.000	0.000
3D5 Other	CO ₂	0	0	10	20	22.361	0.008	0.000	0.000	0.000	0.005	0.005
4A Enteric Fermentation	CH ₄	6	5	10	100	100.499	0.885	0.000	0.008	-0.041	0.117	0.124
4B Manure Management	CH ₄	0	0	10	100	100.499	0.021	0.000	0.000	-0.001	0.003	0.003
4B Manure Management	N ₂ O	1	1	10	100	100.499	0.106	0.000	0.001	0.007	0.014	0.016
4D1 Direct N ₂ O emissions from agricultural soils	N ₂ O	0	1	20	50	53.852	0.125	0.001	0.002	0.074	0.062	0.097

IPCC Source category	Gas	Base year emission	Year t emis- sion	Activity data uncertainty	Emission factor uncertainty	Combined uncertainty	Combined uncertainty as % of total national emissions in year t	Type A sensitivity	Type B sensitivity	Uncertainty in trend in national emissions introduced by emission factor uncer- tainty	Uncertainty in trend in national emissions introduced by activity data uncertainty	Uncertainty introduced into the total national emissions
		Input data Gg CO ₂ eq	Input data Gg CO ₂ eq	Input data %	Input data %	%	%	%	%	%	%	%
<i>Continued</i>												
4D2 Pasture range and paddock	N ₂ O	1	1	20	25	32.016	0.034	0.000	0.001	-0.004	0.028	0.028
4D3 Indirect N ₂ O emissions from agricultural soils (Atmospheric deposition)	N ₂ O	1	1	20	50	53.852	0.098	0.001	0.002	0.035	0.048	0.060
5A Forest	CO ₂	0	0	5	50	50.249	-0.003	0.000	0.000	-0.003	0.000	0.003
5B Cropland	CO ₂	0	0	5	50	50.249	0.004	0.000	0.000	0.004	0.001	0.004
5C Grassland	CO ₂	0	1	5	50	50.249	0.108	0.002	0.002	0.086	0.014	0.087
6A Solid Waste Disposal on Land	CH ₄	4	4	10	100	100.499	0.640	0.001	0.006	0.074	0.085	0.113
6B Wastewater Handling	N ₂ O	15	12	30	100	104.403	2.013	-0.003	0.018	-0.336	0.768	0.838
6C Waste incineration	CO ₂	3	3	10	25	26.926	0.138	0.001	0.005	0.028	0.068	0.074
6C Waste incineration	CH ₄	2	2	10	50	50.990	0.133	-0.001	0.002	-0.040	0.035	0.053
6C Waste incineration	N ₂ O	1	1	10	100	100.499	0.133	0.000	0.001	-0.034	0.018	0.038
Total		651	612				14,610					7,105
Total uncertainties				Overall uncertainty in the year (%):			3.822			Trend uncertainty (%):		2.665

16.19 Annex 8 Results of a technical analysis conducted on the Greenlandic gas oil

In 2013 a technical analysis has been conducted on the arctic gas oil that is by far the most dominant type of fuel in Greenland. The analysis was conducted by the Danish Technological Institute in order to gain a country specific emission factor on the Greenlandic gas oil.

Table 16.19.1 shows the results of the technological analysis on the Greenlandic gas oil.

Table 16.19.1 Results on the technical analysis on the Greenlandic gas oil.

	Test result	Method
C, %	85.4	Elementaranalyse
Upper calorific, J/g	45860	DS/CEN/TS 14918
Lower calorific, J/g	42900	Calculation
CO ₂ emission factor, kg CO ₂ /GJ	72.237	Calculation

17 Information regarding the aggregated submission for Denmark and Greenland

This chapter contains information on the aggregated submission for Denmark and Greenland submitted under the Kyoto Protocol. This chapter contains a trend discussion, a tier 1 uncertainty analysis, information on the aggregated reference approach, information relating to key categories and information on recalculations. Sector specific information is included for Denmark in Chapter 3-11 and for Greenland in Chapter 16.

The institutional arrangements and the overall QA/QC plan are described in Chapter 1. This description covers all the Danish submissions to the European Union, the UNFCCC and the Kyoto Protocol, and therefore information regarding the national system is not presented in this chapter. Information on the specific QA/QC activities concerning the aggregated submission is presented in Chapter 17.7.

In Chapter 17.6 a description of the aggregation process is provided. The chapter explains the technical issues in aggregating two CRF submissions, including the software used in the process and the handling of background data.

17.1 Trends in emissions

Due to the small emission originating from Greenland the trends for Denmark and Greenland are practically identical to the trends for Denmark presented in Chapter 2.

17.1.1 Greenhouse Gas Emissions

The greenhouse gas emissions are estimated according to the IPCC guidelines and are aggregated into seven main sectors. The greenhouse gases include CO₂, CH₄, N₂O, HFCs, PFCs and SF₆. Figure 17.1 shows the estimated total greenhouse gas emissions in CO₂ equivalents from 1990 to 2012. The emissions are not corrected for electricity trade or temperature variations. CO₂ is the most important greenhouse gas contributing in 2012 to the national total in CO₂ equivalents excluding LULUCF (Land Use and Land Use Change and Forestry) with 76.5 % followed by N₂O with 11.4 %, CH₄ 10.5 % and F-gases (HFCs, PFCs and SF₆) with 0.7 %. Seen over the time series from 1990 to 2012 these percentages have been increasing for F-gases and CH₄, almost constant for CO₂ and falling for N₂O. Stationary combustion plants, transport and agriculture represent the largest categories, followed by industrial processes, waste and solvents, see Figure 17.1. The net CO₂ uptake by LULUCF in 2012 is 1.6 % of the total emission in CO₂ equivalents excl. LULUCF. The national total greenhouse gas emission in CO₂ equivalents excluding LULUCF has decreased by 24.6 % from 1990 to 2012 and decreased 31.1 % including LULUCF. Comments on the overall trends seen in Figure 17.1 are given in the sections below on the individual greenhouse gases.




Figure 17.1 Greenhouse gas emissions in CO₂ equivalents distributed on main sectors for 2012 (excluding LULUCF) and time series for 1990 to 2012 (including LULUCF).

17.1.2 Carbon dioxide

The largest source to the emission of CO₂ is the energy sector, which includes combustion of fossil fuels like oil, coal and natural gas (Figure 17.2). Energy Industries contribute with 42 % of the emissions (excl. LULUCF). About 31 % come from the transport sector. The main reason for the fluctuations during the time series is the variations in electricity import/export. The CO₂ emission (excl. LULUCF) decreased by 10.7 % from 2010 to 2012. The main reasons for this decrease were increase in the share of renewable energy, decreasing fuel consumption, mainly for coal and natural gas, and increase in import of electricity. In 2012, the CO₂ emission (excl. LULUCF) was 25.3 % lower than the emission in 1990.




Figure 17.2 CO₂ emissions. Distribution according to the main sectors (2012) and time series for 1990 to 2012.

17.1.3 Nitrous oxide

Agriculture is the most important N₂O emission source in 2012 contributing 90 % (Figure 17.3) of which N₂O from agricultural soils accounts for 84 %. N₂O is emitted as a result of microbial processes in the soil. Substantial emissions also come from drainage water and coastal waters where nitrogen is converted to N₂O through bacterial processes. However, the nitrogen converted in these processes originates mainly from the agricultural use of manure and nitrogen fertilisers. The main reason for the drop in the emissions of N₂O in the agricultural sector of 35 % from 1990 to 2012 is legislation to improve the utilisation of nitrogen in manure. The legislation has resulted in less nitrogen excreted per unit of livestock produced and a considerable reduction in the use of nitrogen fertilisers. The basis for the N₂O emission is

then reduced. Combustion of fossil fuels in the energy sector, both stationary and mobile sources, contributes 6 %. The N_2O emission from transport contributes by 2 % in 2012. This emission has increased during the nineties because of the increase in the use of catalyst cars but is now decreasing due to improvements in technology. Production of nitric acid stopped in 2004 and the emissions from industrial processes is therefore not occurring from 2005 onwards. Other sources include e.g. use of N_2O for anaesthesia reported under Solvent and Other Product Use.




Figure 17.3 N_2O emissions. Distribution according to the main sectors (2012) and time series for 1990 to 2012.

17.1.4 Methane

The largest sources of anthropogenic CH_4 emissions are agricultural activities contributing in 2012 with 76 %, waste (16 %), public power and district heating plants (3 %), see Figure 17.4. The emission from agriculture derives from enteric fermentation and management of animal manure contributing with 53 % and 24 % of the national CH_4 emission in 2012. The CH_4 emission from public power and district heating plants increased in the nineties, mainly 1992-1996, due to the increasing use of gas engines in the decentralised co-generation plant sector. Up to 3 % of the natural gas in the gas engines is not combusted. The deregulation of the electricity market has made production of electricity in gas engines less favourable, therefore the fuel consumption has decreased and hence the CH_4 emission has decreased. Over the time series from 1990 to 2012, the emission of CH_4 from enteric fermentation has decreased 10.6 % due to the decrease in the number of cattle. However, the emission from manure management has in the same period increased 31.7 % due to a change in traditional animal housing systems towards an increase in slurry-based housing systems. Altogether, the emission of CH_4 from the agriculture sector has decreased by 0.7 % from 1990 to 2012. The emission of CH_4 from waste has decreased 40.8 % since 1990 due to an increase in the recycling, composting and incineration of waste and hence a steep drop in the amount of waste deposited in landfills.




Figure 17.4 CH₄ emissions. Distribution according to the main sectors (2012) and time series for 1990 to 2012.

17.1.5 HFCs, PFCs and SF₆

This part of the Danish KP inventory only comprises a full data set for all substances from 1995. From 1995 to 2000, there has been a continuous and substantial increase in the contribution from the range of F-gases as a whole, calculated as the sum of emissions in CO₂ equivalents, see Figure 17.5. This increase is simultaneous with the increase in the emission of HFCs. For the time series 2000-2012, the increase is lower than for the years 1995 to 2000 and since 2008 the emissions have been decreasing. The increase from 1995 to 2012 for the total F-gas emission is 143 %. SF₆ contributed considerably to the F-gas sum in earlier years, with 33 % in 1995. Environmental awareness and regulation of this gas under Danish law has reduced its use in industry, see Figure 17.5. As a result the contribution of SF₆ to F-gases in 2012 was only 14.9 %. The use of HFCs has increased several folds. HFCs have, therefore, become the even more dominant F-gases, comprising 67 % in 1995, but 84 % in 2012. HFCs are mainly used as a refrigerant. Danish legislation regulates the use of F-gases, e.g. since January 1, 2007 new HFC-based refrigerant stationary systems are forbidden. Refill of old systems are still allowed. The use of air conditioning in mobile systems and the amount of HFC for this purpose increases.




Figure 17.5 F-gas emissions. Time series for 1990 to 2012.

17.2 The reference approach

In addition to the sector-specific CO₂ emission inventories (the national approach), the CO₂ emission is also estimated using the reference approach described in the IPCC Reference Manual (IPCC, 1997). The reference approach is based on data for fuel production, import, export and stock change. The CO₂ emission inventory based on the reference approach is reported to the Climate Convention and used for verification of the official data in the national approach.

The reference approach for Denmark and Greenland is an aggregation of the individual reference approaches for the two. The reference approach for Denmark is described in Chapter 3.4 and the reference approach for Greenland is included in Chapter 16.

In 2012, the fuel consumption rates in the two approaches differ by -0.96 % and the CO₂ emission differs by -0.93 %. In the period 1990-2012, both the fuel consumption and the CO₂ emission differ by less than 2.0%. The differences are below 1 % for all years except 1998, 2009 and 2011. This is almost identical to the reference approach for Denmark, due to the very small emission from Greenland compared to Denmark. A comparison of the national approach and the reference approach is illustrated in Figure 17.6. The differences are related to the statistical differences in the Danish energy statistics, see Chapter 3.4.




Figure 17.6 Comparison of the reference approach and the national approach.

17.3 Uncertainties

An uncertainty estimate has been calculated for Denmark and Greenland. The uncertainty estimate for Denmark is included in Chapter 1.7 and for Greenland in Chapter 16.

The uncertainty estimates are based on the Tier 1 methodology in the IPCC Good Practice Guidance (GPG) (IPCC, 2000). Uncertainty estimates cover 100 % of the total net greenhouse gas emissions and removals. The emissions from Greenland have been treated separately due to the uncertainties being different than the uncertainties in the Danish inventory. The uncertainty of

the Greenlandic emissions has almost no effect on the overall uncertainty estimate, due to the low emissions originating from Greenland.

The estimated uncertainties for total GHG and for CO₂, CH₄, N₂O and F-gases are shown in Table 17.1. The base year for F-gases is 1995 and for all other sources the base year is 1990. The total net GHG emission from Denmark and Greenland is estimated with an uncertainty of ± 6.8 % and the trend in net GHG emission since 1990/1995 has been estimated to be -31.3 % ± 2.6 %-age points. The GHG uncertainty estimates do not take into account the uncertainty of the GWP factors.

The uncertainty on N₂O from leaching, CH₄ emission from solid waste disposal on land, N₂O from synthetic fertilizer, N₂O from animal waste applied to soils, N₂O from stationary combustion of biomass, CO₂ from forest remaining forest and CO₂ from cropland, organic soil are the largest sources of uncertainty for the aggregated greenhouse gas inventory for Greenland and Denmark.

Table 17.1 Uncertainties 1990-2012.

	Uncertainty [%]	Trend [%]	Uncertainty in trend [%-age points]
GHG incl. LULUCF	6.8	-31.3	2.6
GHG excl. LULUCF	5.5	-24.9	2.5
CO ₂	5.5	-33.4	2.4
CH ₄	19.3	-7.3	11.8
N ₂ O	42	-39	13
F-gases	43	143	41

The uncertainties for the activity rates and emission factors are shown in Table 17.2.

Table 17.2 Uncertainties for activity rates and emission factors.

IPCC Source category		Gas	Base year emission Input data Gg CO ₂ eqv.	2012 emission Input data Gg CO ₂ eqv.	Activity data uncertainty Input data %	Emission factor un- certainty Input data %
			23833.9			
Denmark	Stationary Combustion, Coal	CO ₂	1	10005.25	0.9	0.5
Denmark	Stationary Combustion, BKB	CO ₂	10.97	1.41	3.0	5.0
Denmark	Stationary Combustion, Coke	CO ₂	137.80	73.81	1.8	5.0
Denmark	Stationary Combustion, Fossil waste	CO ₂	573.46	1397.05	5.0	10.0
Denmark	Stationary Combustion, Petroleum coke	CO ₂	414.74	627.60	2.0	5.0
Denmark	Stationary Combustion, Residual oil	CO ₂	2495.97	570.79	1.2	2.0
Denmark	Stationary Combustion, Gas oil	CO ₂	4546.79	742.07	2.2	4.0
Denmark	Stationary Combustion, Kerosene	CO ₂	365.68	1.69	2.0	5.0
Denmark	Stationary Combustion, LPG	CO ₂	183.88	91.05	1.8	5.0
Denmark	Stationary Combustion, Refinery gas	CO ₂	816.13	905.83	1.0	2.0
Denmark	Stationary Combustion, Natural gas	CO ₂	4335.40	8293.11	1.0	0.4
Denmark	Stationary Combustion, SOLID	CH ₄	12.88	3.12	1.0	100.0
Denmark	Stationary Combustion, LIQUID	CH ₄	2.76	1.11	0.8	100.0
Denmark	Stationary Combustion, GAS	CH ₄	3.13	5.59	1.0	100.0
Denmark	Natural gas fuelled engines, GAS	CH ₄	4.64	120.43	1.0	2.0
Denmark	Stationary Combustion, WASTE	CH ₄	0.77	1.54	5.0	100.0
Denmark	Stationary Combustion, BIOMASS	CH ₄	101.83	134.83	15.7	100.0
Denmark	Biogas fuelled engines, BIOMASS	CH ₄	1.48	31.34	3.9	10.0
Denmark	Stationary Combustion, SOLID	N ₂ O	68.11	27.45	1.0	400.0
Denmark	Stationary Combustion, LIQUID	N ₂ O	44.02	11.08	0.8	1000.0
Denmark	Stationary Combustion, GAS	N ₂ O	16.38	24.45	1.0	750.0
Denmark	Stationary Combustion, WASTE	N ₂ O	6.58	15.72	5.0	400.0
Denmark	Stationary Combustion, BIOMASS	N ₂ O	37.92	90.89	2.2	1000.0
Denmark	Transport, Road transport	CO ₂	9283.52	11223.78	2.0	5.0
Denmark	Transport, Military	CO ₂	119.01	115.79	2.0	5.0
Denmark	Transport, Railways	CO ₂	296.75	249.33	2.0	5.0
Denmark	Transport, Navigation (small boats)	CO ₂	47.92	98.62	41.0	5.0
Denmark	Transport, Navigation (large vessels)	CO ₂	747.83	399.02	11.0	5.0
Denmark	Transport, Fisheries	CO ₂	590.60	479.10	2.0	5.0
Denmark	Transport, Agriculture	CO ₂	1272.47	1343.48	24.0	5.0
Denmark	Transport, Forestry	CO ₂	35.68	16.86	30.0	5.0
Denmark	Transport, Industry (mobile)	CO ₂	839.28	1021.40	41.0	5.0
Denmark	Transport, Residential	CO ₂	39.06	62.43	35.0	5.0
Denmark	Transport, Commercial/institutional	CO ₂	73.72	171.41	35.0	5.0
Denmark	Transport, Civil aviation	CO ₂	242.69	132.61	10.0	5.0
Denmark	Transport, Road transport	CH ₄	46.90	11.29	2.0	40.0
Denmark	Transport, Military	CH ₄	0.11	0.06	2.0	100.0
Denmark	Transport, Railways	CH ₄	0.26	0.15	2.0	100.0
Denmark	Transport, Navigation (small boats)	CH ₄	0.35	0.51	41.0	100.0
Denmark	Transport, Navigation (large vessels)	CH ₄	0.33	0.20	11.0	100.0
Denmark	Transport, Fisheries	CH ₄	0.27	0.25	2.0	100.0
Denmark	Transport, Agriculture	CH ₄	2.20	2.03	24.0	100.0
Denmark	Transport, Forestry	CH ₄	0.44	0.05	30.0	100.0
Denmark	Transport, Industry (mobile)	CH ₄	1.25	0.75	41.0	100.0
Denmark	Transport, Residential	CH ₄	1.07	1.36	35.0	100.0
Denmark	Transport, Commercial/institutional	CH ₄	2.08	3.18	35.0	100.0
Denmark	Transport, Civil aviation	CH ₄	0.09	0.05	10.0	100.0
Denmark	Transport, Road transport	N ₂ O	90.64	116.44	2.0	50.0
Denmark	Transport, Military	N ₂ O	1.15	1.27	2.0	1000.0

IPCC Source category		Gas	Base year emission Input data Gg CO ₂ eqv.	2012 emission Input data Gg CO ₂ eqv.	Activity data uncertainty Input data %	Emission factor un- certainty Input data %
<i>Continued</i>						
Denmark	Transport, Railways	N ₂ O	2.54	2.13	2.0	1000.0
Denmark	Transport, Navigation (small boats)	N ₂ O	0.39	1.08	41.0	1000.0
Denmark	Transport, Navigation (large vessels)	N ₂ O	14.60	7.78	11.0	1000.0
Denmark	Transport, Fisheries	N ₂ O	11.50	9.39	2.0	1000.0
Denmark	Transport, Agriculture	N ₂ O	15.27	17.70	24.0	1000.0
Denmark	Transport, Forestry	N ₂ O	0.17	0.17	30.0	1000.0
Denmark	Transport, Industry (mobile)	N ₂ O	10.62	13.44	41.0	1000.0
Denmark	Transport, Residential	N ₂ O	0.19	0.33	35.0	1000.0
Denmark	Transport, Commercial/institutional	N ₂ O	0.34	0.82	35.0	1000.0
Denmark	Transport, Civil aviation	N ₂ O	3.19	2.07	10.0	1000.0
Denmark	1.B.2 Flaring in refinery	CO ₂	22.95	22.33	11.0	2.0
Denmark	1.B.2 Flaring off-shore	CO ₂	301.85	194.57	7.5	2.0
Denmark	1.B.2 Land based activities	CO ₂	0.00	0.01	2.0	40.0
Denmark	1.B.2 Off-shore activities	CO ₂	2.38	3.70	2.0	30.0
Denmark	1.B.2 Transmission of natural gas	CO ₂	0.00	0.00	15.0	2.0
Denmark	1.B.2 Distribution of natural gas	CO ₂	0.00	0.00	25.0	10.0
Denmark	1.B.2 Venting in gas storage	CO ₂	0.00	0.00	15.0	2.0
Denmark	1.B.2 Flaring in refinery	CH ₄	0.16	0.16	11.0	15.0
Denmark	1.B.2 Flaring off-shore	CH ₄	0.73	0.32	7.5	125.0
Denmark	1.B.2 Refinery processes	CH ₄	0.78	46.76	1.0	125.0
Denmark	1.B.2 Land based activities	CH ₄	17.16	17.58	2.0	40.0
Denmark	1.B.2 Off-shore activities	CH ₄	14.87	36.62	2.0	30.0
Denmark	1.B.2 Transmission of natural gas	CH ₄	3.99	0.29	15.0	2.0
Denmark	1.B.2 Distribution of natural gas	CH ₄	5.42	3.08	25.0	10.0
Denmark	1.B.2 Venting in gas storage	CH ₄	1.23	1.21	15.0	2.0
Denmark	1.B.2 Flaring in refinery	N ₂ O	0.06	0.06	11.0	1000.0
Denmark	1.B.2 Flaring off-shore	N ₂ O	0.71	0.47	7.5	1000.0
Denmark	2A1 Cement production	CO ₂	882.40	871.08	1.0	2.0
Denmark	2A2 Lime production	CO ₂	115.53	40.17	5.0	5.0
Denmark	2A3 Limestone and dolomite use	CO ₂	13.69	25.62	5.0	5.0
Denmark	2A5 Asphalt roofing	CO ₂	0.02	0.02	5.0	25.0
Denmark	2A6 Road paving with asphalt	CO ₂	1.76	1.77	5.0	25.0
Denmark	2A7a Glass and Glass wool	CO ₂	17.41	9.69	5.0	2.0
Denmark	2A7b Yellow bricks	CO ₂	23.02	17.48	5.0	2.0
Denmark	2A7c Expanded clay	CO ₂	14.93	5.89	5.0	2.0
Denmark	2B5 Catalysts/Fertilizers, Pesticides and Sulphuric acid	CO ₂	0.80	1.41	5.0	5.0
Denmark	2C1 Iron and steel production	CO ₂	28.45	0.00	5.0	5.0
Denmark	2D2 Food and Drink	CO ₂	4.45	2.24	5.0	5.0
Denmark	2G Lubricants	CO ₂	49.71	31.70	2.0	5.0
Denmark	2B2 Nitric acid production	N ₂ O	1042.90	0.00	2.0	25.0
Denmark	2F Consumption of HFC	HFC	217.73	657.20	10.0	50.0
Denmark	2F Consumption of PFC	PFC	0.50	8.54	10.0	50.0
Denmark	2F Consumption of SF ₆	SF ₆	107.34	117.85	10.0	50.0
Denmark	3A Paint application	CO ₂	13.22	7.23	10.0	15.0
Denmark	3B Degreasing and dry cleaning	CO ₂	0.00	0.00	10.0	15.0
Denmark	3C Chemical products, manufacturing and processing	CO ₂	19.42	12.17	10.0	15.0
Denmark	3D5 Other	CO ₂	60.64	43.74	10.0	20.0
Denmark	3D5 Consumption of fireworks	CO ₂	0.06	0.15	8.0	100.0

IPCC Source category		Gas	Base year emission Input data Gg CO ₂ eqv.	2012 emission Input data Gg CO ₂ eqv.	Activity data uncertainty Input data %	Emission factor un- certainty Input data %
<i>Continued</i>						
Denmark	3D5 Use of candles	CO ₂	21.66	81.10	10.0	20.0
Denmark	3D1 Other - Use of N ₂ O for Anaesthesia	N ₂ O	0.00	9.34	5.0	5.0
Denmark	3D5 Use of tobacco	N ₂ O	0.25	0.16	20.0	30.0
Denmark	3D5 Use of charcoal for BBQ	N ₂ O	0.07	0.13	10.0	100.0
Denmark	3D5 Consumption of fireworks	N ₂ O	0.77	2.09	8.0	100.0
Denmark	3D5 Use of candles	N ₂ O	0.06	0.21	10.0	20.0
Denmark	4A Enteric Fermentation	CH ₄	3246.93	2903.79	2.0	20.0
Denmark	4B Manure Management	CH ₄	985.16	1297.17	5.0	20.0
Denmark	4F Field burning of agricultural residues	CH ₄	1.82	2.43	25.0	50.0
Denmark	4.B Manure Management	N ₂ O	599.78	390.77	22.4	100.0
Denmark	4.D1.1 Synthetic Fertilizer	N ₂ O	2353.53	1103.41	25.2	100.0
Denmark	4.D1.2 Animal waste applied to soils	N ₂ O	1111.74	1161.35	30.0	100.0
Denmark	4.D1.3 N-fixing crops	N ₂ O	269.47	256.21	20.0	100.0
Denmark	4.D1.4 Crop Residue	N ₂ O	361.22	311.19	20.0	100.0
Denmark	4.D1.5 Cultivation of histosols	N ₂ O	290.23	198.31	20.0	100.0
Denmark	4.D.2 Grassing animals	N ₂ O	334.25	210.88	25.5	100.0
Denmark	4.D3 Atmospheric deposition	N ₂ O	496.40	294.60	18.7	100.0
Denmark	4.D3 Leaching	N ₂ O	2447.13	1429.53	20.0	100.0
Denmark	4.D1.6 Sewage sludge and Industrial waste used as fertiliser	N ₂ O	27.92	38.76	20.0	100.0
Denmark	4.F Field Burning of Agricultural Residues	N ₂ O	0.70	0.93	25.0	50.0
Denmark	5.A.1 Forest remaining forest	CO ₂	49.60	-4491.37	15.0	2.0
Denmark	5.A.2 Land converted to forest	CO ₂	76.62	37.83	15.0	8.7
Denmark	5(II) Forest Land.	N ₂ O	15.64	12.07	30.0	10.0
Denmark	5.B Cropland, Living biomass	CO ₂	-73.52	192.99	10.0	50.0
Denmark	5.B Cropland, Dead organic matter	CO ₂	2.93	12.16	10.0	50.0
Denmark	5.B Cropland, Mineral soils	CO ₂	1415.31	577.08	10.0	75.0
Denmark	5.B Cropland, Organic soils	CO ₂	2887.33	1980.71	10.0	90.0
Denmark	5(III) Disturbance, Land converted to cropland	N ₂ O	0.25	1.01	50.0	75.0
Denmark	5.C Grassland, Living biomass	CO ₂	75.25	463.24	10.0	50.0
Denmark	5.C Grassland, Dead organic matter	CO ₂	1.63	7.04	10.0	50.0
Denmark	5.C Grassland, Mineral soils	CO ₂	0.16	4.54	10.0	75.0
Denmark	5.C Grassland, Organic soils	CO ₂	106.58	78.70	10.0	90.0
Denmark	5.D Wetlands, Living biomass	CO ₂	2.57	-0.22	10.0	50.0
Denmark	5.D Wetlands, Dead organic matter	CO ₂	0.54	0.00	10.0	100.0
Denmark	5.D Wetlands, Soils	CO ₂	84.67	2.50	10.0	100.0
Denmark	5(II) Wetlands	N ₂ O	0.13	0.13	10.0	100.0
Denmark	5.E Settlements, Living biomass	CO ₂	11.41	55.99	10.0	50.0
Denmark	5.E Settlements, Dead organic matter	CO ₂	0.55	0.03	10.0	50.0
Denmark	5.E Settlements, Soils	CO ₂	0.96	34.78	10.0	50.0
Denmark	5(IV) Cropland Limestone	CO ₂	622.92	192.30	5.0	50.0
Denmark	5(V) Biomass Burning	CH ₄	0.55	0.02	50.0	30.0
Denmark	5(V) Biomass Burning	N ₂ O	0.45	0.03	50.0	30.0
Denmark	6 A. Solid Waste Disposal on Land	CH ₄	1365.77	698.04	10.0	117.9
Denmark	6 B. Wastewater Handling	CH ₄	65.49	74.00	24.0	31.6
Denmark	6 B. Wastewater Handling - Direct	N ₂ O	22.70	40.70	21.7	49.6
Denmark	6 B. Wastewater Handling - Indirect	N ₂ O	82.25	32.14	59.2	42.0
Denmark	6.D Accidental fires, buildings	CO ₂	11.41	10.79	10.0	300.0
Denmark	6.D Accidental fires, vehicles	CO ₂	6.13	5.56	10.0	500.0

IPCC Source category		Gas	Base year emission Input data Gg CO ₂ eqv.	2012 emission Input data Gg CO ₂ eqv.	Activity data uncertainty Input data %	Emission factor un- certainty Input data %
<i>Continued</i>						
Denmark	6.C Incineration of corpses	CH ₄	0.01	0.01	1.0	150.0
Denmark	6.C Incineration of carcasses	CH ₄	0.00	0.00	40.0	150.0
Denmark	6.D Compost production	CH ₄	29.11	90.43	40.0	100.0
Denmark	6.D Accidental fires, buildings	CH ₄	1.35	1.30	10.0	500.0
Denmark	6.D Accidental fires, vehicles	CH ₄	0.27	0.24	10.00	700.00
Denmark	6.C Incineration of corpses	N ₂ O	0.19	0.19	1.00	150.00
Denmark	6.C Incineration of carcasses	N ₂ O	0.01	0.09	40.00	150.00
Denmark	6.D Compost production	N ₂ O	12.85	126.83	40.00	100.00
Greenland	1A Liquid fuels	CO ₂	612.92	562.38	2.0	2.0
Greenland	1A Municipal waste	CO ₂	1.67	6.94	2.0	25.0
Greenland	1A Liquid fuels	CH ₄	1.02	0.96	2.0	100.0
Greenland	1A Municipal waste	CH ₄	0.01	0.05	2.0	100.0
Greenland	1A Biomass	CH ₄	0.02	0.06	2.0	100.0
Greenland	1A Liquid fuels	N ₂ O	1.82	1.74	2.0	500.0
Greenland	1A Municipal waste	N ₂ O	0.03	0.10	2.0	500.0
Greenland	1A Biomass	N ₂ O	0.03	0.13	2.0	200.0
Greenland	1B2 Oil exploration	CO ₂	0.00	0.00	2.0	1000.0
Greenland	1B2 Oil exploration	CH ₄	0.00	0.00	2.0	1000.0
Greenland	1B2 Oil exploration	N ₂ O	0.00	0.00	2.0	1000.0
Greenland	2A3 Limestone and dolomite use	CO ₂	0.00	0.02	5.0	5.0
Greenland	2A5 Asphalt roofing	CO ₂	0.00	0.00	5.0	25.0
Greenland	2A6 Road paving with asphalt	CO ₂	0.00	0.00	5.0	25.0
Greenland	2F Consumption of HFC	HFC	0.02	7.19	10.0	50.0
Greenland	2F Consumption of SF ₆	SF ₆	0.04	0.00	10.0	50.0
Greenland	3A Paint application	CO ₂	0.01	0.00	10.0	15.0
Greenland	3B Degreasing and dry cleaning	CO ₂	0.00	0.00	10.0	15.0
Greenland	3C Chemical products, manufacturing and processing	CO ₂	0.00	0.00	10.0	15.0
Greenland	3D5 Other	CO ₂	0.25	0.23	10.0	20.0
Greenland	4A Enteric Fermentation	CH ₄	6.02	5.39	10.0	100.0
Greenland	4B Manure Management	CH ₄	0.14	0.13	10.0	100.0
Greenland	4.B Manure Management	N ₂ O	0.64	0.65	10.0	100.0
Greenland	4D1 Direct N ₂ O emissions from agricultural soils	N ₂ O	0.48	1.42	20.0	50.0
Greenland	4D2 Pasture range and paddock	N ₂ O	0.80	0.65	20.0	25.0
Greenland	4D3 Indirect N ₂ O emissions from agricultural soils	N ₂ O	0.70	1.11	20.0	50.0
Greenland	5A Forest	CO ₂	0.00	-0.04	5.0	50.0
Greenland	5B Cropland	CO ₂	0.00	0.05	5.0	50.0
Greenland	5.C Grassland	CO ₂	0.21	1.32	5.0	50.0
Greenland	6A Solid Waste Disposal on Land	CH ₄	3.64	3.90	10.0	100.0
Greenland	6B Wastewater Handling	N ₂ O	14.88	11.79	30.0	100.0
Greenland	6C Waste incineration	CO ₂	2.55	3.13	10.0	25.0
Greenland	6C Waste incineration	CH ₄	2.26	1.60	10.0	50.0
Greenland	6C Waste incineration	N ₂ O	1.10	0.81	10.0	100.0

17.4 Key category analysis

A tier 1 key category analysis (KCA) has been carried out on emissions from Denmark and Greenland. The key category analysis for Denmark is included

in Chapter 1.5 and Annex 1, and the key category analysis for Greenland is included in Chapter 16.

The KCA for 1990 and 2012 has been carried out in accordance with the IPCC Guidelines 2006 and IPCC Good Practice Guidance. The KCA has been carried out at CRF level, which is slightly more aggregated than the KCA carried out for Denmark. The categorisation used results in a total of 113 source categories of which 20 are LULUCF categories.

The KCA for Denmark and Greenland includes a total of six different analyses:

- base year, reporting year and trend,
- including and excluding LULUCF.

The six different KCA for Denmark and Greenland point out 16-24 key source categories each and a total of 27 different key source categories. The number of key categories in each of the main sectors is: Energy 7, Industrial Proc. 3, Solvents and other product use 0, Agriculture 10, LULUCF 6 and Waste 1.

An overview for all KCA is given in Table 17.3. The table includes ranking for each key category emission sources.

The six KCA for Denmark and Greenland are shown in Table 17.4-17.9.

Table 17.3 Key Category Analysis for Denmark and Greenland, overview.

IPCC Source Categories			GHG			Excluding LULUCF			Including LULUCF		
						Level Tier1	Level Tier1	Trend Tier1	Level Tier1	Level Tier1	Trend Tier1
						1990	2012	1990-2012	1990	2012	1990-2012
Energy	Combustion excluding transport, Liquid Fuels	CO ₂				2	4	2	2	4	2
Energy	Combustion excluding transport, Solid Fuels	CO ₂				1	2	1	1	2	1
Energy	Combustion excluding transport, Gaseous Fuels	CO ₂				4	3	3	4	3	4
Energy	Combustion excluding transport, Other Fuels	CO ₂				15	7	8	18	9	11
Energy	Combustion excluding transport, Liquid Fuels	CH ₄									
Energy	Combustion excluding transport, Solid Fuels	CH ₄									
Energy	Combustion excluding transport, Gaseous Fuels	CH ₄									
Energy	Combustion excluding transport, Biomass	CH ₄									
Energy	Combustion excluding transport, Other Fuels	CH ₄									
Energy	Combustion excluding transport, Liquid Fuels	N ₂ O									
Energy	Combustion excluding transport, Solid Fuels	N ₂ O									
Energy	Combustion excluding transport, Gaseous Fuels	N ₂ O									
Energy	Combustion excluding transport, Biomass	N ₂ O									
Energy	Combustion excluding transport, Other Fuels	N ₂ O									
Energy	Road transportation	CO ₂				3	1	4	3	1	5
Energy	Road transportation	CH ₄									
Energy	Road transportation	N ₂ O									
Energy	Civil aviation	CO ₂									
Energy	Civil aviation	CH ₄									
Energy	Civil aviation	N ₂ O									
Energy	Domestic navigation	CO ₂				13	14	13	15	17	18
Energy	Domestic navigation	CH ₄									
Energy	Domestic navigation	N ₂ O									
Energy	Railways	CO ₂					19			23	
Energy	Railways	CH ₄									
Energy	Railways	N ₂ O									
Energy	Fugitive emissions , 1B2ai Oil Exploration	CO ₂									
Energy	Fugitive emissions , 1B2ai Oil Exploration	CH ₄									
Energy	Fugitive emissions , 1B2ai Oil Exploration	N ₂ O									
Energy	Fugitive emissions , 1B2aii Oil Production	CO ₂									
Energy	Fugitive emissions , 1B2aii Oil Production	CH ₄									
Energy	Fugitive emissions , 1B2aii Oil Production	N ₂ O									

IPCC Source Categories			GHG			Excluding LULUCF			Including LULUCF		
						Level Tier1 1990	Level Tier1 2012	Trend Tier1 1990- 2012	Level Tier1 1990	Level Tier1 2012	Trend Tier1 1990- 2012
<i>Continued</i>											
Energy	Fugitive emissions , 1B2aiv, Oil refining and storage	CH ₄									
Energy	Fugitive emissions , 1B2b Natural gas transmission and distribution	CO ₂									
Energy	Fugitive emissions , 1B2biii, Gas transmission	CH ₄									
Energy	Fugitive emissions , 1B2biv, Gas distribution	CH ₄									
Energy	Fugitive emissions , 1B2c Venting	CO ₂									
Energy	Fugitive emissions , 1B2c Venting	CH ₄									
Energy	Fugitive emissions , 1B2c2ii, Flaring gas	CO ₂									
Energy	Fugitive emissions , 1B2c2ii, Flaring gas	CH ₄									
Energy	Fugitive emissions , 1B2c2ii, Flaring gas	N ₂ O									
Energy	Fugitive emissions , 1B2c2i, Flaring oil	CO ₂									
Energy	Fugitive emissions , 1B2c2i, Flaring oil	CH ₄									
Energy	Fugitive emissions , 1B2c2i, Flaring oil	N ₂ O									
Industrial processes	Cement production	CO ₂	12	11					14	13	
Industrial processes	Lime production	CO ₂									
Industrial processes	Limestone and dolomite use	CO ₂									
Industrial processes	Asphalt roofing	CO ₂									
Industrial processes	Road Paving with asphalt	CO ₂									
Industrial processes	Glass production	CO ₂									
Industrial processes	Yellow bricks	CO ₂									
Industrial processes	Expanded clay	CO ₂									
Industrial processes	Nitric acid production	N ₂ O	10		6				12		7
Industrial processes	Catalysts/Fertilizers, Pesticides and Sulphuric acid	CO ₂									
Industrial processes	Iron and steel production	CO ₂									
Industrial processes	Food and drink	CO ₂									
Industrial processes	Refrigeration and AC Equipment	HFC and PFC					13	10		15	13
Industrial processes	Foam Blowing	HFC									
Industrial processes	Aerosols	HFC									
Industrial processes	Electrical equipment	SF ₆									
Industrial processes	Other emissions of SF6 i.e. from double glaze windows and laboratories	SF ₆									
Industrial processes	Other i.e. Fibre Optics	HFC and PFC									
Industrial processes	Magnesium Production	SF ₆									
Industrial processes	Other, lubricants	CO ₂									
Solvents and other product use	Paint application	CO ₂									
Solvents and other product use	Degreasing and Dry Cleaning	CO ₂									
Solvents and other product use	Chemical Products, Manufacture and Processing	CO ₂									
Solvents and other product use	Other solvent	CO ₂									
Solvents and other product use	Other solvent	N ₂ O									
Agriculture	Enteric Fermentation	CH ₄	5	5	11	5	6	16			
Agriculture	Manure Management	CH ₄	11	8	12	13	10	17			
Agriculture	Manure management	N ₂ O	14	15	14	17	19	20			
Agriculture	Agriculture soils, direct emissions , Synthetic Fertilizers	N ₂ O	7	10	5	8	12	6			
Agriculture	Agriculture soils, direct emissions , Animal Manure Appl. to Soils	N ₂ O	9	9		11	11				
Agriculture	Agriculture soils, direct emissions , N-fixing Crops	N ₂ O		18			22				
Agriculture	Agriculture soils, direct emissions , Crop Residue	N ₂ O	17	16		20	20				
Agriculture	Agriculture soils, direct emissions , Cultivation of Histosols	N ₂ O									
Agriculture	Agriculture soils, direct emissions , Sludge	N ₂ O									
Agriculture	Agriculture soils, pasture, range and paddock	N ₂ O				21	24				
Agriculture	Agriculture soils, indirect, Atmospheric Deposition	N ₂ O	16	17	15	19	21	21			
Agriculture	Agriculture soils, indirect, Nitrogen Leaching and Run-off	N ₂ O	6	6	7	7	8	8			
Agriculture	Field Burning of Agricultural Residues	CH ₄									
Agriculture	Field Burning of Agricultural Residues	N ₂ O									
Waste	Solid Waste Disposal Sites	CH ₄	8	12	9	10	14	12			
Waste	N2O direct, Domestic and Commercial Wastewater	N2O									

IPCC Source Categories			GHG			Excluding LULUCF			Including LULUCF		
						Level Tier1 1990	Level Tier1 2012	Trend Tier1 1990- 2012	Level Tier1 1990	Level Tier1 2012	Trend Tier1 1990- 2012
<i>Continued</i>											
Waste	N ₂ O indirect from human sewage	N ₂ O									
Waste	Waste Water Handling	CH ₄									
Waste	Waste incineration	CO ₂									
Waste	Waste incineration	CH ₄									
Waste	Waste incineration	N ₂ O									
Waste	Waste, other	CO ₂									
Waste	Waste, other	CH ₄									
Waste	Waste, other	N ₂ O									
LULUCF	Forest Land remaining Forest L.	CO ₂							5	3	
LULUCF	Land converted to Forest L.	CO ₂									
LULUCF	Cropland, 5.B Cropland, Living biomass	CO ₂									19
LULUCF	Cropland, 5.B Cropland, Dead organic matter	CO ₂									
LULUCF	Cropland, 5.B Cropland, Mineral soils	CO ₂							9	16	10
LULUCF	Cropland, 5.B Cropland, Organic soils	CO ₂							6	7	9
LULUCF	Grassland, 5.C Grassland, Living biomass	CO ₂								18	15
LULUCF	Grassland, 5.C Grassland, Dead organic matter	CO ₂									
LULUCF	Grassland, 5.C Grassland, Mineral soils	CO ₂									
LULUCF	Grassland, 5.C Grassland, Organic soils	CO ₂									
LULUCF	Wetlands, 5D Wetlands Living biomass	CO ₂									
LULUCF	Wetlands, 5D Wetlands, dead org. matter	CO ₂									
LULUCF	Wetlands, 5D Wetlands, soils	CO ₂									
LULUCF	Settlements, 5E Total settlements	CO ₂									
LULUCF	Non-CO ₂ drainage of soils and wetlands, 5IID Forest Land.	N ₂ O									
LULUCF	Non-CO ₂ drainage of soils and wetlands, 5IID Wetlands. Peatland	N ₂ O									
LULUCF	N ₂ O Disturbance, Land converted to cropland, 5III Cropland	N ₂ O									
LULUCF	Agricultural lime application, 5IV Cropland Lime- stone	CO ₂							16		14
LULUCF	Biomass burning, 5V Biomass burning	CH ₄									
LULUCF	Biomass burning, 5V Biomass burning	N ₂ O									

Table 17.4 Key Category Analysis for Denmark and Greenland, level assessment for the base year, excl. LULUCF.

IPCC Source Categories (LULUCF excluded)		GHG	Base Year Estimate	Base Year Level	Base Year Cumulative total
			Ex,o	Assessment Lx,o	
			Gg CO ₂ eqv.		
Energy	Combustion excluding transport, Solid Fuels	CO ₂	23982.67	0.3448	0.3448
Energy	Combustion excluding transport, Liquid Fuels	CO ₂	12310.84	0.1770	0.5218
Energy	Road transportation	CO ₂	9319.58	0.1340	0.6558
Energy	Combustion excluding transport, Gaseous Fuels	CO ₂	4335.40	0.0623	0.7181
Agriculture	Enteric Fermentation	CH ₄	3252.95	0.0468	0.7649
Agriculture	Agriculture soils, indirect, Nitrogen Leaching and Run-off	N ₂ O	2447.73	0.0352	0.8000
Agriculture	Agriculture soils, direct emissions , Synthetic Fertilizers	N ₂ O	2353.58	0.0338	0.8339
Waste	Solid Waste Disposal Sites	CH ₄	1369.41	0.0197	0.8536
Agriculture	Agriculture soils, direct emissions , Animal Manure Appl. to Soils	N ₂ O	1112.06	0.0160	0.8696
Industrial processes	Nitric acid production	N ₂ O	1042.90	0.0150	0.8845
Agriculture	Manure Management	CH ₄	985.30	0.0142	0.8987
Industrial processes	Cement production	CO ₂	882.40	0.0127	0.9114
Energy	Domestic navigation	CO ₂	816.48	0.0117	0.9231
Agriculture	Manure Management	N ₂ O	600.42	0.0086	0.9318
Energy	Combustion excluding transport, Other Fuels	CO ₂	575.14	0.0083	0.9400
Agriculture	Agriculture soils, indirect, Atmospheric Deposition	N ₂ O	496.50	0.0071	0.9472
Agriculture	Agriculture soils, direct emissions , Crop Residue	N ₂ O	361.27	0.0052	0.9524
Agriculture	Agriculture soils, pasture, range and paddock	N ₂ O	335.05	0.0048	0.9572
Energy	Fugitive emissions , 1B2c2ii, Flaring gas	CO ₂	301.85	0.0043	0.9615
Energy	Railways	CO ₂	296.75	0.0043	0.9658
Agriculture	Agriculture soils, direct emissions , Cultivation of Histosols	N ₂ O	290.29	0.0042	0.9700
Energy	Civil aviation	CO ₂	281.01	0.0040	0.9740
Agriculture	Agriculture soils, direct emissions , N-fixing Crops	N ₂ O	269.47	0.0039	0.9779
Industrial processes	Foam Blowing	HFC	182.58	0.0026	0.9805
Industrial processes	Lime production	CO ₂	115.53	0.0017	0.9822
Energy	Combustion excluding transport, Biomass	CH ₄	103.75	0.0015	0.9837
Energy	Road transportation	N ₂ O	90.74	0.0013	0.9850
Waste	N ₂ O indirect from human sewage	N ₂ O	85.23	0.0012	0.9862
Energy	Combustion excluding transport, Liquid Fuels	N ₂ O	84.59	0.0012	0.9874
Solvents and other product use	Other solvent	CO ₂	82.61	0.0012	0.9886
Energy	Combustion excluding transport, Solid Fuels	N ₂ O	68.11	0.0010	0.9896
Industrial processes	Other emissions of SF ₆ i.e. from double glaze windows and laboratories	SF ₆	67.62	0.0010	0.9905
Waste	Waste Water Handling	CH ₄	65.49	0.0009	0.9915
Industrial processes	Other, lubricants	CO ₂	49.71	0.0007	0.9922
Energy	Road transportation	CH ₄	46.97	0.0007	0.9929
Energy	Combustion excluding transport, Biomass	N ₂ O	41.58	0.0006	0.9935
Industrial processes	Refrigeration and AC Equipment	HFC and PFC	35.68	0.0005	0.9940
Waste	N ₂ O direct, Domestic and Commercial Wastewater	N ₂ O	34.60	0.0005	0.9945
Energy	Fugitive emissions , 1B2aii Oil Production	CH ₄	32.03	0.0005	0.9949
Waste	Waste, other	CH ₄	30.72	0.0004	0.9954
Industrial processes	Iron and steel production	CO ₂	28.45	0.0004	0.9958
Agriculture	Agriculture soils, direct emissions , Sludge	N ₂ O	27.92	0.0004	0.9962
Industrial processes	Yellow bricks	CO ₂	23.02	0.0003	0.9965
Energy	Fugitive emissions , 1B2c2i, Flaring oil	CO ₂	22.95	0.0003	0.9969
Solvents and other product use	Chemical Products, Manufacture and Processing	CO ₂	19.42	0.0003	0.9971
Waste	Waste, other	CO ₂	17.54	0.0003	0.9974
Industrial processes	Glass production	CO ₂	17.41	0.0003	0.9976
Energy	Combustion excluding transport, Gaseous Fuels	N ₂ O	16.38	0.0002	0.9979
Energy	Domestic navigation	N ₂ O	15.05	0.0002	0.9981
Industrial processes	Expanded clay	CO ₂	14.93	0.0002	0.9983
Industrial processes	Limestone and dolomite use	CO ₂	13.69	0.0002	0.9985
Solvents and other product use	Paint application	CO ₂	13.23	0.0002	0.9987
Energy	Combustion excluding transport, Solid Fuels	CH ₄	12.88	0.0002	0.9989
Waste	Waste, other	N ₂ O	12.85	0.0002	0.9991
Energy	Combustion excluding transport, Liquid Fuels	CH ₄	11.10	0.0002	0.9992
Energy	Combustion excluding transport, Gaseous Fuels	CH ₄	7.77	0.0001	0.9993
Energy	Fugitive emissions , 1B2biv, Gas distribution	CH ₄	5.42	0.0001	0.9994
Industrial processes	Food and drink	CO ₂	4.45	0.0001	0.9995

IPCC Source Categories (LULUCF excluded)		GHG	Base Year Estimate	Base Year Level	Base Year Cumulative total
			Ex,o Gg CO ₂ eqv.	Assess- ment Lx,o	
<i>Continued</i>					
Energy	Fugitive emissions , 1B2biii, Gas transmission	CH ₄	3.99	0.0001	0.9995
Industrial processes	Electrical equipment	SF ₆	3.91	0.0001	0.9996
Energy	Civil aviation	N ₂ O	3.53	0.0001	0.9996
Energy	Combustion excluding transport, Other Fuels	N ₂ O	2.99	0.0000	0.9997
Waste	Waste incineration	CO ₂	2.55	0.0000	0.9997
Energy	Railways	N ₂ O	2.54	0.0000	0.9998
Energy	Fugitive emissions , 1B2aai Oil Production	CO ₂	2.38	0.0000	0.9998
Waste	Waste incineration	CH ₄	2.27	0.0000	0.9998
Agriculture	Field Burning of Agricultural Residues	CH ₄	1.82	0.0000	0.9998
Industrial processes	Road Paving with asphalt	CO ₂	1.76	0.0000	0.9999
Waste	Waste incineration	N ₂ O	1.30	0.0000	0.9999
Energy	Fugitive emissions , 1B2c Venting	CH ₄	1.23	0.0000	0.9999
Solvents and other product use	Other solvent	N ₂ O	1.14	0.0000	0.9999
Industrial processes	Catalysts/Fertilizers, Pesticides and Sulphuric acid	CO ₂	0.80	0.0000	0.9999
Energy	Fugitive emissions , 1B2aiv, Oil refining and storage	CH ₄	0.78	0.0000	0.9999
Energy	Fugitive emissions , 1B2c2ii, Flaring gas	CH ₄	0.73	0.0000	1.0000
Energy	Domestic navigation	CH ₄	0.71	0.0000	1.0000
Energy	Fugitive emissions , 1B2c2ii, Flaring gas	N ₂ O	0.71	0.0000	1.0000
Agriculture	Field Burning of Agricultural Residues	N ₂ O	0.70	0.0000	1.0000
Energy	Combustion excluding transport, Other Fuels	CH ₄	0.36	0.0000	1.0000
Energy	Railways	CH ₄	0.26	0.0000	1.0000
Energy	Fugitive emissions , 1B2c2i, Flaring oil	CH ₄	0.16	0.0000	1.0000
Energy	Civil aviation	CH ₄	0.10	0.0000	1.0000
Energy	Fugitive emissions , 1B2c2i, Flaring oil	N ₂ O	0.06	0.0000	1.0000
Industrial processes	Asphalt roofing	CO ₂	0.02	0.0000	1.0000
Energy	Fugitive emissions , 1B2b Natural gas transmission and distribution	CO ₂	0.01	0.0000	1.0000
Energy	Fugitive emissions , 1B2c Venting	CO ₂	0.00	0.0000	1.0000
Solvents and other product use	Degreasing and Dry Cleaning	CO ₂	0.00	0.0000	1.0000
Energy	Fugitive emissions , 1B2ai Oil Exploration	CH ₄	0.00	0.0000	1.0000
Energy	Fugitive emissions , 1B2ai Oil Exploration	CO ₂	0.00	0.0000	1.0000
Energy	Fugitive emissions , 1B2ai Oil Exploration	N ₂ O	0.00	0.0000	1.0000
Energy	Fugitive emissions , 1B2aai Oil Production	N ₂ O	0.00	0.0000	1.0000
Industrial processes	Aerosols	HFC	0.00	0.0000	1.0000
Industrial processes	Magnesium Production	SF ₆	0.00	0.0000	1.0000
Industrial processes	Other i.e. Fibre Optics	HFC and PFC	0.00	0.0000	1.0000
Total			69557.81	1.000	

Table 17.5 Key Category Analysis for Denmark and Greenland, level assessment for the base year, incl. LULUCF.

IPCC Source Categories (LULUCF included)		GHG	Base Year Estimate Ex,o Gg CO ₂ eqv	Base Year Level Assessment Lx,o	Base Year Cumulative total
Energy	Combustion excluding transport, Solid Fuels	CO ₂	23982.67	0.3198	0.3198
Energy	Combustion excluding transport, Liquid Fuels	CO ₂	12310.84	0.1642	0.4840
Energy	Road transportation	CO ₂	9319.58	0.1243	0.6083
Energy	Combustion excluding transport, Gaseous Fuels	CO ₂	4335.40	0.0578	0.6661
Agriculture	Enteric Fermentation	CH ₄	3252.95	0.0434	0.7095
LULUCF	Cropland, 5.B Cropland, Organic soils	CO ₂	2887.33	0.0385	0.7480
Agriculture	Agriculture soils, indirect, Nitrogen Leaching and Run-off	N ₂ O	2447.73	0.0326	0.7806
Agriculture	Agriculture soils, direct emissions , Synthetic Fertilizers	N ₂ O	2353.58	0.0314	0.8120
LULUCF	Cropland, 5.B Cropland, Mineral soils	CO ₂	1415.31	0.0189	0.8309
Waste	Solid Waste Disposal Sites	CH ₄	1369.41	0.0183	0.8491
Agriculture	Agriculture soils, direct emissions , Animal Manure Appl. to Soils	N ₂ O	1112.06	0.0148	0.8640
Industrial processes	Nitric acid production	N ₂ O	1042.90	0.0139	0.8779
Agriculture	Manure Management	CH ₄	985.30	0.0131	0.8910
Industrial processes	Cement production	CO ₂	882.40	0.0118	0.9028
Energy	Domestic navigation	CO ₂	816.48	0.0109	0.9137
LULUCF	Agricultural lime application, 5IV Cropland Limestone	CO ₂	622.93	0.0083	0.9220
Agriculture	Manure Management	N ₂ O	600.42	0.0080	0.9300
Energy	Combustion excluding transport, Other Fuels	CO ₂	575.14	0.0077	0.9377
Agriculture	Agriculture soils, indirect, Atmospheric Deposition	N ₂ O	496.50	0.0066	0.9443
Agriculture	Agriculture soils, direct emissions , Crop Residue	N ₂ O	361.27	0.0048	0.9491
Agriculture	Agriculture soils, pasture, range and paddock	N ₂ O	335.05	0.0045	0.9536
Energy	Fugitive emissions , 1B2c2ii, Flaring gas	CO ₂	301.85	0.0040	0.9576
Energy	Railways	CO ₂	296.75	0.0040	0.9615
Agriculture	Agriculture soils, direct emissions , Cultivation of Histosols	N ₂ O	290.29	0.0039	0.9654
Energy	Civil aviation	CO ₂	281.01	0.0037	0.9692
Agriculture	Agriculture soils, direct emissions , N-fixing Crops	N ₂ O	269.47	0.0036	0.9728
Industrial processes	Foam Blowing	HFC	182.58	0.0024	0.9752
Industrial processes	Lime production	CO ₂	115.53	0.0015	0.9767
LULUCF	Grassland, 5.C Grassland, Organic soils	CO ₂	107.14	0.0014	0.9782
Energy	Combustion excluding transport, Biomass	CH ₄	103.75	0.0014	0.9795
Energy	Road transportation	N ₂ O	90.74	0.0012	0.9808
Waste	N ₂ O indirect from human sewage	N ₂ O	85.23	0.0011	0.9819
LULUCF	Wetlands, 5D Wetlands, soils	CO ₂	84.67	0.0011	0.9830
Energy	Combustion excluding transport, Liquid Fuels	N ₂ O	84.59	0.0011	0.9841
Solvents and other product use	Other solvent	CO ₂	82.61	0.0011	0.9852
LULUCF	Land converted to Forest L., (blank)	CO ₂	76.62	0.0010	0.9863
LULUCF	Grassland, 5.C Grassland, Living biomass	CO ₂	74.90	0.0010	0.9873
LULUCF	Cropland, 5.B Cropland, Living biomass	CO ₂	73.52	0.0010	0.9882
Energy	Combustion excluding transport, Solid Fuels	N ₂ O	68.11	0.0009	0.9892
Industrial processes	Other emissions of SF ₆ i.e. from double glaze windows and laboratories	SF ₆	67.62	0.0009	0.9901
Waste	Waste Water Handling	CH ₄	65.49	0.0009	0.9909
Industrial processes	Other, lubricants	CO ₂	49.71	0.0007	0.9916
LULUCF	Forest Land remaining Forest L., (blank)	CO ₂	49.60	0.0007	0.9923
Energy	Road transportation	CH ₄	46.97	0.0006	0.9929
Energy	Combustion excluding transport, Biomass	N ₂ O	41.58	0.0006	0.9934
Industrial processes	Refrigeration and AC Equipment	HFC and PFC	35.68	0.0005	0.9939
Waste	N ₂ O direct, Domestic and Commercial Wastewater	N ₂ O	34.60	0.0005	0.9944
Energy	Fugitive emissions , 1B2aii Oil Production	CH ₄	32.03	0.0004	0.9948
Waste	Waste, other	CH ₄	30.72	0.0004	0.9952
Industrial processes	Iron and steel production	CO ₂	28.45	0.0004	0.9956
Agriculture	Agriculture soils, direct emissions , Sludge	N ₂ O	27.92	0.0004	0.9960
Industrial processes	Yellow bricks	CO ₂	23.02	0.0003	0.9963
Energy	Fugitive emissions , 1B2c2i, Flaring oil	CO ₂	22.95	0.0003	0.9966
Solvents and other product use	Chemical Products, Manufacture and Processing	CO ₂	19.42	0.0003	0.9968
Waste	Waste, other	CO ₂	17.54	0.0002	0.9971
Industrial processes	Glass production	CO ₂	17.41	0.0002	0.9973
Energy	Combustion excluding transport, Gaseous Fuels	N ₂ O	16.38	0.0002	0.9975
LULUCF	Non-CO ₂ drainage of soils and wetlands, 5IID Forest Land.	N ₂ O	15.64	0.0002	0.9977
Energy	Domestic navigation	N ₂ O	15.05	0.0002	0.9979
Industrial processes	Expanded clay	CO ₂	14.93	0.0002	0.9981
Industrial processes	Limestone and dolomite use	CO ₂	13.69	0.0002	0.9983
Solvents and other product use	Paint application	CO ₂	13.23	0.0002	0.9985
LULUCF	Settlements, 5E Total settlements	CO ₂	12.92	0.0002	0.9987
Energy	Combustion excluding transport, Solid Fuels	CH ₄	12.88	0.0002	0.9988

IPCC Source Categories (LULUCF included)		GHG	Base Year Estimate Ex,o Gg CO ₂ eqv	Base Year Level Assessment Lx,o	Base Year Cumulative total
<i>Continued</i>					
Waste	Waste, other	N ₂ O	12.85	0.0002	0.9990
Energy	Combustion excluding transport, Liquid Fuels	CH ₄	11.10	0.0001	0.9992
Energy	Combustion excluding transport, Gaseous Fuels	CH ₄	7.77	0.0001	0.9993
Energy	Fugitive emissions , 1B2biv, Gas distribution	CH ₄	5.42	0.0001	0.9993
Industrial processes	Food and drink	CO ₂	4.45	0.0001	0.9994
Energy	Fugitive emissions , 1B2biii, Gas transmission	CH ₄	3.99	0.0001	0.9994
Industrial processes	Electrical equipment	SF ₆	3.91	0.0001	0.9995
Energy	Civil aviation	N ₂ O	3.53	0.0000	0.9995
Energy	Combustion excluding transport, Other Fuels	N ₂ O	2.99	0.0000	0.9996
LULUCF	Cropland, 5.B Cropland, Dead organic matter	CO ₂	2.93	0.0000	0.9996
LULUCF	Wetlands, 5D Wetlands Living biomass	CO ₂	2.57	0.0000	0.9997
Waste	Waste incineration	CO ₂	2.55	0.0000	0.9997
Energy	Railways	N ₂ O	2.54	0.0000	0.9997
Energy	Fugitive emissions , 1B2aii Oil Production	CO ₂	2.38	0.0000	0.9998
Waste	Waste incineration	CH ₄	2.27	0.0000	0.9998
Agriculture	Field Burning of Agricultural Residues	CH ₄	1.82	0.0000	0.9998
Industrial processes	Road Paving with asphalt	CO ₂	1.76	0.0000	0.9998
LULUCF	Grassland, 5.C Grassland, Dead organic matter	CO ₂	1.63	0.0000	0.9999
Waste	Waste incineration	N ₂ O	1.30	0.0000	0.9999
Energy	Fugitive emissions , 1B2c Venting	CH ₄	1.23	0.0000	0.9999
Solvents and other product use	Other solvent	N ₂ O	1.14	0.0000	0.9999
Industrial processes	Catalysts/Fertilizers, Pesticides and Sulphuric acid	CO ₂	0.80	0.0000	0.9999
Energy	Fugitive emissions , 1B2aiv, Oil refining and storage	CH ₄	0.78	0.0000	0.9999
Energy	Fugitive emissions , 1B2c2ii, Flaring gas	CH ₄	0.73	0.0000	0.9999
Energy	Domestic navigation	CH ₄	0.71	0.0000	0.9999
Energy	Fugitive emissions , 1B2c2ii, Flaring gas	N ₂ O	0.71	0.0000	1.0000
Agriculture	Field Burning of Agricultural Residues	N ₂ O	0.70	0.0000	1.0000
LULUCF	Biomass burning, 5V Biomass burning	CH ₄	0.55	0.0000	1.0000
LULUCF	Wetlands, 5D Wetlands, dead org. matter	CO ₂	0.54	0.0000	1.0000
LULUCF	Biomass burning, 5V Biomass burning	N ₂ O	0.45	0.0000	1.0000
Energy	Combustion excluding transport, Other Fuels	CH ₄	0.36	0.0000	1.0000
Energy	Railways	CH ₄	0.26	0.0000	1.0000
LULUCF	N2O Disturbance, Land converted to cropland, 5III Cropland	N ₂ O	0.25	0.0000	1.0000
Energy	Fugitive emissions , 1B2c2i, Flaring oil	CH ₄	0.16	0.0000	1.0000
LULUCF	Grassland, 5.C Grassland, Mineral soils	CO ₂	0.16	0.0000	1.0000
LULUCF	Non-CO2 drainage of soils and wetlands, 5IID Wetlands. Peat- land	N ₂ O	0.13	0.0000	1.0000
Energy	Civil aviation	CH ₄	0.10	0.0000	1.0000
Energy	Fugitive emissions , 1B2c2i, Flaring oil	N ₂ O	0.06	0.0000	1.0000
Industrial processes	Asphalt roofing	CO ₂	0.02	0.0000	1.0000
Energy	Fugitive emissions , 1B2b Natural gas transmission and distribu- tion	CO ₂	0.01	0.0000	1.0000
Energy	Fugitive emissions , 1B2c Venting	CO ₂	0.00	0.0000	1.0000
Solvents and other product use	Degreasing and Dry Cleaning	CO ₂	0.00	0.0000	1.0000
Energy	Fugitive emissions , 1B2ai Oil Exploration	CH ₄	0.00	0.0000	1.0000
Energy	Fugitive emissions , 1B2ai Oil Exploration	CO ₂	0.00	0.0000	1.0000
Energy	Fugitive emissions , 1B2ai Oil Exploration	N ₂ O	0.00	0.0000	1.0000
Energy	Fugitive emissions , 1B2aii Oil Production	N ₂ O	0.00	0.0000	1.0000
Industrial processes	Aerosols	HFC	0.00	0.0000	1.0000
Industrial processes	Magnesium Production	SF ₆	0.00	0.0000	1.0000
Industrial processes	Other i.e. Fibre Optics	HFC and PFC	0.00	0.0000	1.0000
Total			74987.62	1.000	

Table 17.6 Key Category Analysis for Denmark and Greenland, level assessment for the reporting year, excl. LULUCF.

IPCC Source Categories (LULUCF excluded)		GHG	Reporting Year Estimate Ex,t Gg CO ₂ eqv	Reporting Year Level Assessment Lx,t	Reporting Year Cumulative total
Energy	Road transportation	CO ₂	11257.01	0.2155	0.2155
Energy	Combustion excluding transport, Solid Fuels	CO ₂	10080.46	0.1929	0.4084
Energy	Combustion excluding transport, Gaseous Fuels	CO ₂	8293.11	0.1587	0.5671
Energy	Combustion excluding transport, Liquid Fuels	CO ₂	6602.22	0.1264	0.6935
Agriculture	Enteric Fermentation	CH ₄	2909.18	0.0557	0.7492
Agriculture	Agriculture soils, indirect, Nitrogen Leaching and Run-off	N ₂ O	1430.55	0.0274	0.7765
Energy	Combustion excluding transport, Other Fuels	CO ₂	1403.99	0.0269	0.8034
Agriculture	Manure Management	CH ₄	1297.30	0.0248	0.8282
Agriculture	Agriculture soils, direct emissions , Animal Manure Appl. to Soils	N ₂ O	1161.67	0.0222	0.8505
Agriculture	Agriculture soils, direct emissions , Synthetic Fertilizers	N ₂ O	1104.26	0.0211	0.8716
Industrial processes	Cement production	CO ₂	871.08	0.0167	0.8883
Waste	Solid Waste Disposal Sites	CH ₄	701.94	0.0134	0.9017
Industrial processes	Refrigeration and AC Equipment	HFC and PFC	586.60	0.0112	0.9129
Energy	Domestic navigation	CO ₂	527.38	0.0101	0.9230
Agriculture	Manure Management	N ₂ O	391.42	0.0075	0.9305
Agriculture	Agriculture soils, direct emissions , Crop Residue	N ₂ O	311.30	0.0060	0.9365
Agriculture	Agriculture soils, indirect, Atmospheric Deposition	N ₂ O	294.70	0.0056	0.9421
Agriculture	Agriculture soils, direct emissions , N-fixing Crops	N ₂ O	256.21	0.0049	0.9470
Energy	Railways	CO ₂	249.33	0.0048	0.9518
Agriculture	Agriculture soils, pasture, range and paddock	N ₂ O	211.53	0.0040	0.9559
Agriculture	Agriculture soils, direct emissions , Cultivation of Histosols	N ₂ O	198.45	0.0038	0.9597
Energy	Fugitive emissions , 1B2c2ii, Flaring gas	CO ₂	194.57	0.0037	0.9634
Energy	Civil aviation	CO ₂	179.28	0.0034	0.9668
Energy	Combustion excluding transport, Biomass	CH ₄	167.06	0.0032	0.9700
Waste	Waste, other	N ₂ O	126.83	0.0024	0.9724
Energy	Combustion excluding transport, Gaseous Fuels	CH ₄	126.02	0.0024	0.9748
Solvents and other product use	Other solvent	CO ₂	125.22	0.0024	0.9772
Energy	Road transportation	N ₂ O	116.53	0.0022	0.9795
Industrial processes	Other emissions of SF6 i.e. from double glaze windows and laboratories	SF6	104.59	0.0020	0.9815
Energy	Combustion excluding transport, Biomass	N ₂ O	99.62	0.0019	0.9834
Waste	Waste, other	CH ₄	91.97	0.0018	0.9851
Waste	Waste Water Handling	CH ₄	74.00	0.0014	0.9866
Industrial processes	Foam Blowing	HFC	66.18	0.0013	0.9878
Energy	Combustion excluding transport, Liquid Fuels	N ₂ O	55.36	0.0011	0.9889
Energy	Fugitive emissions , 1B2aii Oil Production	CH ₄	54.20	0.0010	0.9899
Waste	N2O direct, Domestic and Commercial Wastewater	N ₂ O	49.46	0.0009	0.9909
Energy	Fugitive emissions , 1B2aiv, Oil refining and storage	CH ₄	46.76	0.0009	0.9918
Industrial processes	Lime production	CO ₂	40.17	0.0008	0.9925
Agriculture	Agriculture soils, direct emissions , Sludge	N ₂ O	38.76	0.0007	0.9933
Waste	N2O indirect from human sewage	N ₂ O	35.17	0.0007	0.9939
Industrial processes	Other, lubricants	CO ₂	31.70	0.0006	0.9946
Energy	Combustion excluding transport, Solid Fuels	N ₂ O	27.45	0.0005	0.9951
Industrial processes	Limestone and dolomite use	CO ₂	25.64	0.0005	0.9956
Energy	Combustion excluding transport, Gaseous Fuels	N ₂ O	24.45	0.0005	0.9960
Energy	Fugitive emissions , 1B2c2i, Flaring oil	CO ₂	22.33	0.0004	0.9965
Industrial processes	Yellow bricks	CO ₂	17.48	0.0003	0.9968
Waste	Waste, other	CO ₂	16.36	0.0003	0.9971
Industrial processes	Aerosols	HFC	15.83	0.0003	0.9974
Industrial processes	Electrical equipment	SF6	13.27	0.0003	0.9977
Solvents and other product use	Chemical Products, Manufacture and Processing	CO ₂	12.17	0.0002	0.9979
Solvents and other product use	Other solvent	N ₂ O	11.93	0.0002	0.9981
Energy	Road transportation	CH ₄	11.40	0.0002	0.9983
Industrial processes	Glass production	CO ₂	9.69	0.0002	0.9985
Energy	Combustion excluding transport, Liquid Fuels	CH ₄	9.59	0.0002	0.9987
Energy	Domestic navigation	N ₂ O	8.94	0.0002	0.9989
Energy	Combustion excluding transport, Other Fuels	N ₂ O	7.23	0.0001	0.9990
Solvents and other product use	Paint application	CO ₂	7.23	0.0001	0.9992
Industrial processes	Expanded clay	CO ₂	5.89	0.0001	0.9993

IPCC Source Categories (LULUCF excluded)		GHG	Reporting Year Estimate Ex,t Gg CO ₂ eqv	Reporting Year Level Assessment Lx,t	Reporting Year Cumulative total
<i>Continued</i>					
Industrial processes	Other i.e. Fibre Optics	HFC and PFC	4.31	0.0001	0.9994
Energy	Fugitive emissions , 1B2aii Oil Production	CO ₂	3.71	0.0001	0.9994
Waste	Waste incineration	CO ₂	3.13	0.0001	0.9995
Energy	Combustion excluding transport, Solid Fuels	CH ₄	3.12	0.0001	0.9996
Energy	Fugitive emissions , 1B2biv, Gas distribution	CH ₄	3.08	0.0001	0.9996
Energy	Civil aviation	N ₂ O	2.48	0.0000	0.9997
Agriculture	Field Burning of Agricultural Residues	CH ₄	2.43	0.0000	0.9997
Industrial processes	Food and drink	CO ₂	2.24	0.0000	0.9997
Energy	Railways	N ₂ O	2.13	0.0000	0.9998
Industrial processes	Road Paving with asphalt	CO ₂	1.77	0.0000	0.9998
Waste	Waste incineration	CH ₄	1.61	0.0000	0.9999
Industrial processes	Catalysts/Fertilizers, Pesticides and Sulphuric acid	CO ₂	1.41	0.0000	0.9999
Energy	Fugitive emissions , 1B2c Venting	CH ₄	1.21	0.0000	0.9999
Waste	Waste incineration	N ₂ O	1.08	0.0000	0.9999
Agriculture	Field Burning of Agricultural Residues	N ₂ O	0.93	0.0000	0.9999
Energy	Combustion excluding transport, Other Fuels	CH ₄	0.77	0.0000	1.0000
Energy	Domestic navigation	CH ₄	0.75	0.0000	1.0000
Energy	Fugitive emissions , 1B2c2ii, Flaring gas	N ₂ O	0.47	0.0000	1.0000
Energy	Fugitive emissions , 1B2c2ii, Flaring gas	CH ₄	0.32	0.0000	1.0000
Energy	Fugitive emissions , 1B2biii, Gas transmission	CH ₄	0.29	0.0000	1.0000
Energy	Fugitive emissions , 1B2c2i, Flaring oil	CH ₄	0.16	0.0000	1.0000
Energy	Railways	CH ₄	0.15	0.0000	1.0000
Energy	Fugitive emissions , 1B2c2i, Flaring oil	N ₂ O	0.06	0.0000	1.0000
Energy	Civil aviation	CH ₄	0.06	0.0000	1.0000
Industrial processes	Asphalt roofing	CO ₂	0.02	0.0000	1.0000
Energy	Fugitive emissions , 1B2b Natural gas transmission and distribution	CO ₂	0.01	0.0000	1.0000
Energy	Fugitive emissions , 1B2c Venting	CO ₂	0.00	0.0000	1.0000
Solvents and other product use	Degreasing and Dry Cleaning	CO ₂	0.00	0.0000	1.0000
Industrial processes	Magnesium Production	SF ₆	0.00	0.0000	1.0000
Energy	Fugitive emissions , 1B2ai Oil Exploration	CH ₄	0.00	0.0000	1.0000
Energy	Fugitive emissions , 1B2ai Oil Exploration	CO ₂	0.00	0.0000	1.0000
Energy	Fugitive emissions , 1B2ai Oil Exploration	N ₂ O	0.00	0.0000	1.0000
Energy	Fugitive emissions , 1B2aii Oil Production	N ₂ O	0.00	0.0000	1.0000
Industrial processes	Nitric acid production	N ₂ O	0.00	0.0000	1.0000
Industrial processes	Iron and steel production	CO ₂	0.00	0.0000	1.0000
Total			52247.69	1.000	

Table 17.7 Key Category Analysis for Denmark and Greenland, level assessment for the reporting year, incl. LULUCF.

IPCC Source Categories (LULUCF included)		GHG	Reporting Year Estimate	Reporting Year Level	Reporting Year Cumulative total
			Ex,t Gg CO ₂ eqv	Assessment Lx,t	
Energy	Road transportation	CO ₂	11257.01	0.1864	0.1864
Energy	Combustion excluding transport, Solid Fuels	CO ₂	10080.46	0.1669	0.3533
Energy	Combustion excluding transport, Gaseous Fuels	CO ₂	8293.11	0.1373	0.4906
Energy	Combustion excluding transport, Liquid Fuels	CO ₂	6602.22	0.1093	0.5999
LULUCF	Forest Land remaining Forest L., (blank)	CO ₂	4491.41	0.0744	0.6743
Agriculture	Enteric Fermentation	CH ₄	2909.18	0.0482	0.7225
LULUCF	Cropland, 5.B Cropland, Organic soils	CO ₂	1980.76	0.0328	0.7553
Agriculture	Agriculture soils, indirect, Nitrogen Leaching and Run-off	N ₂ O	1430.55	0.0237	0.7790
Energy	Combustion excluding transport, Other Fuels	CO ₂	1403.99	0.0232	0.8022
Agriculture	Manure Management	CH ₄	1297.30	0.0215	0.8237
Agriculture	Agriculture soils, direct emissions , Animal Manure Appl. to Soils	N ₂ O	1161.67	0.0192	0.8429
Agriculture	Agriculture soils, direct emissions , Synthetic Fertilizers	N ₂ O	1104.26	0.0183	0.8612
Industrial processes	Cement production	CO ₂	871.08	0.0144	0.8756
Waste	Solid Waste Disposal Sites	CH ₄	701.94	0.0116	0.8873
Industrial processes	Refrigeration and AC Equipment	HFC and PFC	586.60	0.0097	0.8970
LULUCF	Cropland, 5.B Cropland, Mineral soils	CO ₂	577.08	0.0096	0.9065
Energy	Domestic navigation	CO ₂	527.38	0.0087	0.9153
LULUCF	Grassland, 5.C Grassland, Living biomass	CO ₂	463.33	0.0077	0.9229
Agriculture	Manure Management	N ₂ O	391.42	0.0065	0.9294
Agriculture	Agriculture soils, direct emissions , Crop Residue	N ₂ O	311.30	0.0052	0.9346
Agriculture	Agriculture soils, indirect, Atmospheric Deposition	N ₂ O	294.70	0.0049	0.9394
Agriculture	Agriculture soils, direct emissions , N-fixing Crops	N ₂ O	256.21	0.0042	0.9437
Energy	Railways	CO ₂	249.33	0.0041	0.9478
Agriculture	Agriculture soils, pasture, range and paddock	N ₂ O	211.53	0.0035	0.9513
Agriculture	Agriculture soils, direct emissions , Cultivation of Histosols	N ₂ O	198.45	0.0033	0.9546
Energy	Fugitive emissions , 1B2c2ii, Flaring gas	CO ₂	194.57	0.0032	0.9578
LULUCF	Cropland, 5.B Cropland, Living biomass	CO ₂	192.99	0.0032	0.9610
LULUCF	Agricultural lime application, 5IV Cropland Limestone	CO ₂	192.30	0.0032	0.9642
Energy	Civil aviation	CO ₂	179.28	0.0030	0.9672
Energy	Combustion excluding transport, Biomass	CH ₄	167.06	0.0028	0.9699
Waste	Waste, other	N ₂ O	126.83	0.0021	0.9720
Energy	Combustion excluding transport, Gaseous Fuels	CH ₄	126.02	0.0021	0.9741
Solvents and other product use	Other solvent	CO ₂	125.22	0.0021	0.9762
Energy	Road transportation	N ₂ O	116.53	0.0019	0.9781
Industrial processes	Other emissions of SF6 i.e. from double glaze windows and laboratories	SF6	104.59	0.0017	0.9799
Energy	Combustion excluding transport, Biomass	N ₂ O	99.62	0.0016	0.9815
Waste	Waste, other	CH ₄	91.97	0.0015	0.9830
LULUCF	Settlements, 5E Total settlements	CO ₂	90.81	0.0015	0.9845
LULUCF	Grassland, 5.C Grassland, Organic soils	CO ₂	79.91	0.0013	0.9859
Waste	Waste Water Handling	CH ₄	74.00	0.0012	0.9871
Industrial processes	Foam Blowing	HFC	66.18	0.0011	0.9882
Energy	Combustion excluding transport, Liquid Fuels	N ₂ O	55.36	0.0009	0.9891
Energy	Fugitive emissions , 1B2aiv Oil Production	CH ₄	54.20	0.0009	0.9900
Waste	N ₂ O direct, Domestic and Commercial Wastewater	N ₂ O	49.46	0.0008	0.9908
Energy	Fugitive emissions , 1B2aiv, Oil refining and storage	CH ₄	46.76	0.0008	0.9916
Industrial processes	Lime production	CO ₂	40.17	0.0007	0.9923
Agriculture	Agriculture soils, direct emissions , Sludge	N ₂ O	38.76	0.0006	0.9929
LULUCF	Land converted to Forest L., (blank)	CO ₂	37.83	0.0006	0.9935
Waste	N ₂ O indirect from human sewage	N ₂ O	35.17	0.0006	0.9941
Industrial processes	Other, lubricants	CO ₂	31.70	0.0005	0.9946
Energy	Combustion excluding transport, Solid Fuels	N ₂ O	27.45	0.0005	0.9951
Industrial processes	Limestone and dolomite use	CO ₂	25.64	0.0004	0.9955
Energy	Combustion excluding transport, Gaseous Fuels	N ₂ O	24.45	0.0004	0.9959
Energy	Fugitive emissions , 1B2c2i, Flaring oil	CO ₂	22.33	0.0004	0.9963
Industrial processes	Yellow bricks	CO ₂	17.48	0.0003	0.9966
Waste	Waste, other	CO ₂	16.36	0.0003	0.9968
Industrial processes	Aerosols	HFC	15.83	0.0003	0.9971
Industrial processes	Electrical equipment	SF6	13.27	0.0002	0.9973
Solvents and other product use	Chemical Products, Manufacture and Processing	CO2	12.17	0.0002	0.9975
LULUCF	Cropland, 5.B Cropland, Dead organic matter	CO ₂	12.16	0.0002	0.9977
LULUCF	Non-CO ₂ drainage of soils and wetlands, 5IID Forest Land.	N ₂ O	12.07	0.0002	0.9979

IPCC Source Categories (LULUCF included)		GHG	Reporting Year Estimate	Reporting Year Level	Reporting Year Cumulative total
			Ex,t Gg CO ₂ eqv	Assessment Lx,t	
<i>Continued</i>					
Solvents and other product use	Other solvent	N ₂ O	11.93	0.0002	0.9981
Energy	Road transportation	CH ₄	11.40	0.0002	0.9983
Industrial processes	Glass production	CO ₂	9.69	0.0002	0.9985
Energy	Combustion excluding transport, Liquid Fuels	CH ₄	9.59	0.0002	0.9986
Energy	Domestic navigation	N ₂ O	8.94	0.0001	0.9988
Energy	Combustion excluding transport, Other Fuels	N ₂ O	7.23	0.0001	0.9989
Solvents and other product use	Paint application	CO ₂	7.23	0.0001	0.9990
LULUCF	Grassland, 5.C Grassland, Dead organic matter	CO ₂	7.04	0.0001	0.9991
Industrial processes	Expanded clay	CO ₂	5.89	0.0001	0.9992
LULUCF	Grassland, 5.C Grassland, Mineral soils	CO ₂	4.54	0.0001	0.9993
Industrial processes	Other i.e. Fibre Optics	HFC and PFC	4.31	0.0001	0.9994
Energy	Fugitive emissions , 1B2a ⁱⁱ Oil Production	CO ₂	3.71	0.0001	0.9994
Waste	Waste incineration	CO ₂	3.13	0.0001	0.9995
Energy	Combustion excluding transport, Solid Fuels	CH ₄	3.12	0.0001	0.9995
Energy	Fugitive emissions , 1B2b ^{iv} , Gas distribution	CH ₄	3.08	0.0001	0.9996
LULUCF	Wetlands, 5D Wetlands, soils	CO ₂	2.50	0.0000	0.9996
Energy	Civil aviation	N ₂ O	2.48	0.0000	0.9997
Agriculture	Field Burning of Agricultural Residues	CH ₄	2.43	0.0000	0.9997
Industrial processes	Food and drink	CO ₂	2.24	0.0000	0.9998
Energy	Railways	N ₂ O	2.13	0.0000	0.9998
Industrial processes	Road Paving with asphalt	CO ₂	1.77	0.0000	0.9998
Waste	Waste incineration	CH ₄	1.61	0.0000	0.9998
Industrial processes	Catalysts/Fertilizers, Pesticides and Sulphuric acid	CO ₂	1.41	0.0000	0.9999
Energy	Fugitive emissions , 1B2c Venting	CH ₄	1.21	0.0000	0.9999
Waste	Waste incineration	N ₂ O	1.08	0.0000	0.9999
LULUCF	N ₂ O Disturbance, Land converted to cropland, 5I ^{ll} Cropland	N ₂ O	1.01	0.0000	0.9999
Agriculture	Field Burning of Agricultural Residues	N ₂ O	0.93	0.0000	0.9999
Energy	Combustion excluding transport, Other Fuels	CH ₄	0.77	0.0000	1.0000
Energy	Domestic navigation	CH ₄	0.75	0.0000	1.0000
Energy	Fugitive emissions , 1B2c ²ⁱⁱ , Flaring gas	N ₂ O	0.47	0.0000	1.0000
Energy	Fugitive emissions , 1B2c ²ⁱⁱ , Flaring gas	CH ₄	0.32	0.0000	1.0000
Energy	Fugitive emissions , 1B2b ⁱⁱⁱ , Gas transmission	CH ₄	0.29	0.0000	1.0000
LULUCF	Wetlands, 5D Wetlands Living biomass	CO ₂	0.22	0.0000	1.0000
Energy	Fugitive emissions , 1B2c ²ⁱ , Flaring oil	CH ₄	0.16	0.0000	1.0000
Energy	Railways	CH ₄	0.15	0.0000	1.0000
LULUCF	Non-CO ₂ drainage of soils and wetlands, 5I ^{lD} Wetlands. Peatland	N ₂ O	0.13	0.0000	1.0000
Energy	Fugitive emissions , 1B2c ²ⁱ , Flaring oil	N ₂ O	0.06	0.0000	1.0000
Energy	Civil aviation	CH ₄	0.06	0.0000	1.0000
LULUCF	Biomass burning, 5V Biomass burning	N ₂ O	0.03	0.0000	1.0000
LULUCF	Biomass burning, 5V Biomass burning	CH ₄	0.02	0.0000	1.0000
Industrial processes	Asphalt roofing	CO ₂	0.02	0.0000	1.0000
Energy	Fugitive emissions , 1B2b Natural gas transmission and distribution	CO ₂	0.01	0.0000	1.0000
Energy	Fugitive emissions , 1B2c Venting	CO ₂	0.00	0.0000	1.0000
Solvents and other product use	Degreasing and Dry Cleaning	CO ₂	0.00	0.0000	1.0000
Industrial processes	Magnesium Production	SF ₆	0.00	0.0000	1.0000
Energy	Fugitive emissions , 1B2a ⁱ Oil Exploration	CH ₄	0.00	0.0000	1.0000
Energy	Fugitive emissions , 1B2a ⁱ Oil Exploration	CO ₂	0.00	0.0000	1.0000
Energy	Fugitive emissions , 1B2a ⁱ Oil Exploration	N ₂ O	0.00	0.0000	1.0000
Energy	Fugitive emissions , 1B2a ⁱⁱ Oil Production	N ₂ O	0.00	0.0000	1.0000
LULUCF	Wetlands, 5D Wetlands, dead org. matter	CO ₂	0.00	0.0000	1.0000
Industrial processes	Nitric acid production	N ₂ O	0.00	0.0000	1.0000
Industrial processes	Iron and steel production	CO ₂	0.00	0.0000	1.0000
Total			60393.85	1.000	

Table 17.8 Key Category Analysis for Denmark and Greenland, trend assessment, excl. LULUCF.

IPCC Source Categories (LULUCF excluded)		GHG	Base Year Estimate	Year 2012 Estimate	Trend Assessment	Contribution to Trend	Cumulative total
			Ex,o Gg CO ₂ eq	Ex,t Gg CO ₂ eq	Tx,t		
Energy	Combustion excluding transport, Solid Fuels	CO ₂	23982.67	10080.46	0.2661	0.4093	0.4093
Energy	Combustion excluding transport, Liquid Fuels	CO ₂	12310.84	6602.22	0.1093	0.1681	0.5774
Energy	Combustion excluding transport, Gaseous Fuels	CO ₂	4335.40	8293.11	0.0757	0.1165	0.6939
Energy	Road transportation	CO ₂	9319.58	11257.01	0.0371	0.0570	0.7509
Agriculture	Agriculture soils, direct emissions , Synthetic Fertilizers	N ₂ O	2353.58	1104.26	0.0239	0.0368	0.7877
Industrial processes	Nitric acid production	N ₂ O	1042.90	0.00	0.0200	0.0307	0.8184
Agriculture	Agriculture soils, indirect, Nitrogen Leaching and Run-off	N ₂ O	2447.73	1430.55	0.0195	0.0299	0.8484
Energy	Combustion excluding transport, Other Fuels	CO ₂	575.14	1403.99	0.0159	0.0244	0.8728
Waste	Solid Waste Disposal Sites	CH ₄	1369.41	701.94	0.0128	0.0197	0.8924
Industrial processes	Refrigeration and AC Equipment	HFC and PFC	35.68	586.60	0.0105	0.0162	0.9087
Agriculture	Enteric Fermentation	CH ₄	3252.95	2909.18	0.0066	0.0101	0.9188
Agriculture	Manure Management	CH ₄	985.30	1297.30	0.0060	0.0092	0.9280
Energy	Domestic navigation	CO ₂	816.48	527.38	0.0055	0.0085	0.9365
Agriculture	Manure Management	N ₂ O	600.42	391.42	0.0040	0.0062	0.9426
Agriculture	Agriculture soils, indirect, Atmospheric Deposition	N ₂ O	496.50	294.70	0.0039	0.0059	0.9486
Agriculture	Agriculture soils, pasture, range and paddock	N ₂ O	335.05	211.53	0.0024	0.0036	0.9522
Energy	Combustion excluding transport, Gaseous Fuels	CH ₄	7.77	126.02	0.0023	0.0035	0.9557
Industrial processes	Foam Blowing	HFC	182.58	66.18	0.0022	0.0034	0.9591
Waste	Waste, other	N ₂ O	12.85	126.83	0.0022	0.0034	0.9625
Energy	Fugitive emissions, 1B2c2ii, Flaring gas	CO ₂	301.85	194.57	0.0021	0.0032	0.9656
Energy	Civil aviation	CO ₂	281.01	179.28	0.0019	0.0030	0.9686
Agriculture	Agriculture soils, direct emissions , Cultivation of Histosols	N ₂ O	290.29	198.45	0.0018	0.0027	0.9713
Industrial processes	Lime production	CO ₂	115.53	40.17	0.0014	0.0022	0.9736
Energy	Combustion excluding transport, Biomass	CH ₄	103.75	167.06	0.0012	0.0019	0.9754
Waste	Waste, other	CH ₄	30.72	91.97	0.0012	0.0018	0.9772
Energy	Combustion excluding transport, Biomass	N ₂ O	41.58	99.62	0.0011	0.0017	0.9789
Waste	N ₂ O indirect from human sewage	N ₂ O	85.23	35.17	0.0010	0.0015	0.9804
Agriculture	Agriculture soils, direct emissions , Crop Residue	N ₂ O	361.27	311.30	0.0010	0.0015	0.9819
Agriculture	Agriculture soils, direct emissions , Animal Manure Appl. to Soils	N ₂ O	1112.06	1161.67	0.0009	0.0015	0.9833
Energy	Railways	CO ₂	296.75	249.33	0.0009	0.0014	0.9847
Energy	Fugitive emissions , 1B2aiv, Oil refining and storage	CH ₄	0.78	46.76	0.0009	0.0014	0.9861
Solvents and other product use	Other solvent	CO ₂	82.61	125.22	0.0008	0.0013	0.9873
Energy	Combustion excluding transport, Solid Fuels	N ₂ O	68.11	27.45	0.0008	0.0012	0.9885
Industrial processes	Other emissions of SF ₆ i.e. from double glaze windows and laboratories	SF ₆	67.62	104.59	0.0007	0.0011	0.9896
Energy	Road transportation	CH ₄	46.97	11.40	0.0007	0.0010	0.9907
Energy	Combustion excluding transport, Liquid Fuels	N ₂ O	84.59	55.36	0.0006	0.0009	0.9915
Industrial processes	Iron and steel production	CO ₂	28.45	0.00	0.0005	0.0008	0.9924
Energy	Road transportation	N ₂ O	90.74	116.53	0.0005	0.0008	0.9931
Energy	Fugitive emissions , 1B2aii Oil Production	CH ₄	32.03	54.20	0.0004	0.0007	0.9938
Industrial processes	Other, lubricants	CO ₂	49.71	31.70	0.0003	0.0005	0.9943
Industrial processes	Aerosols	HFC	0.00	15.83	0.0003	0.0005	0.9948
Waste	N ₂ O direct, Domestic and Commercial Wastewater	N ₂ O	34.60	49.46	0.0003	0.0004	0.9952
Agriculture	Agriculture soils, direct emissions , N-fixing Crops	N ₂ O	269.47	256.21	0.0003	0.0004	0.9956
Industrial processes	Limestone and dolomite use	CO ₂	13.69	25.64	0.0002	0.0004	0.9960

IPCC Source Categories (LULUCF excluded)		GHG	Base Year Estimate	Year 2012 Estimate	Trend Assessment	Contribution to Trend	Cumulative total
			Ex,o Gg CO ₂ eq	Ex,t Gg CO ₂ eq	Tx,t		
<i>Continued</i>							
Industrial processes	Cement production	CO ₂	882.40	871.08	0.0002	0.0003	0.9963
Agriculture	Agriculture soils, direct emissions , Sludge	N ₂ O	27.92	38.76	0.0002	0.0003	0.9966
Solvents and other product use	Other solvent	N ₂ O	1.14	11.93	0.0002	0.0003	0.9969
Energy	Combustion excluding transport, Solid Fuels	CH ₄	12.88	3.12	0.0002	0.0003	0.9972
Industrial processes	Electrical equipment	SF ₆	3.91	13.27	0.0002	0.0003	0.9975
Industrial processes	Expanded clay	CO ₂	14.93	5.89	0.0002	0.0003	0.9978
Waste	Waste Water Handling	CH ₄	65.49	74.00	0.0002	0.0003	0.9980
Energy	Combustion excluding transport, Gaseous Fuels	N ₂ O	16.38	24.45	0.0002	0.0002	0.9982
Industrial processes	Glass production	CO ₂	17.41	9.69	0.0001	0.0002	0.9985
Solvents and other product use	Chemical Products, Manufacture and Processing	CO ₂	19.42	12.17	0.0001	0.0002	0.9987
Energy	Domestic navigation	N ₂ O	15.05	8.94	0.0001	0.0002	0.9989
Solvents and other product use	Paint application	CO ₂	13.23	7.23	0.0001	0.0002	0.9990
Industrial processes	Yellow bricks	CO ₂	23.02	17.48	0.0001	0.0002	0.9992
Industrial processes	Other i.e. Fibre Optics	HFC and PFC	0.00	4.31	0.0001	0.0001	0.9993
Energy	Combustion excluding transport, Other Fuels	N ₂ O	2.99	7.23	0.0001	0.0001	0.9995
Energy	Fugitive emissions , 1B2biii, Gas transmission	CH ₄	3.99	0.29	0.0001	0.0001	0.9996
Energy	Fugitive emissions , 1B2biv, Gas distribution	CH ₄	5.42	3.08	0.0000	0.0001	0.9996
Industrial processes	Food and drink	CO ₂	4.45	2.24	0.0000	0.0001	0.9997
Energy	Combustion excluding transport, Liquid Fuels	CH ₄	11.10	9.59	0.0000	0.0000	0.9997
Energy	Fugitive emissions , 1B2aii Oil Production	CO ₂	2.38	3.71	0.0000	0.0000	0.9998
Waste	Waste, other	CO ₂	17.54	16.36	0.0000	0.0000	0.9998
Energy	Civil aviation	N ₂ O	3.53	2.48	0.0000	0.0000	0.9998
Waste	Waste incineration	CH ₄	2.27	1.61	0.0000	0.0000	0.9999
Energy	Fugitive emissions , 1B2c2i, Flaring oil	CO ₂	22.95	22.33	0.0000	0.0000	0.9999
Industrial processes	Catalysts/Fertilizers, Pesticides and Sulphuric acid	CO ₂	0.80	1.41	0.0000	0.0000	0.9999
Agriculture	Field Burning of Agricultural Residues	CH ₄	1.82	2.43	0.0000	0.0000	0.9999
Waste	Waste incineration	CO ₂	2.55	3.13	0.0000	0.0000	0.9999
Energy	Combustion excluding transport, Other Fuels	CH ₄	0.36	0.77	0.0000	0.0000	0.9999
Energy	Fugitive emissions , 1B2c2ii, Flaring gas	CH ₄	0.73	0.32	0.0000	0.0000	1.0000
Energy	Railways	N ₂ O	2.54	2.13	0.0000	0.0000	1.0000
Energy	Fugitive emissions , 1B2c2ii, Flaring gas	N ₂ O	0.71	0.47	0.0000	0.0000	1.0000
Agriculture	Field Burning of Agricultural Residues	N ₂ O	0.70	0.93	0.0000	0.0000	1.0000
Waste	Waste incineration	N ₂ O	1.30	1.08	0.0000	0.0000	1.0000
Energy	Railways	CH ₄	0.26	0.15	0.0000	0.0000	1.0000
Energy	Domestic navigation	CH ₄	0.71	0.75	0.0000	0.0000	1.0000
Energy	Civil aviation	CH ₄	0.10	0.06	0.0000	0.0000	1.0000
Energy	Fugitive emissions , 1B2c Venting	CH ₄	1.23	1.21	0.0000	0.0000	1.0000
Industrial processes	Road Paving with asphalt	CO ₂	1.76	1.77	0.0000	0.0000	1.0000
Energy	Fugitive emissions , 1B2c2i, Flaring oil	CH ₄	0.16	0.16	0.0000	0.0000	1.0000
Energy	Fugitive emissions , 1B2b Natural gas transmission and distribution	CO ₂	0.01	0.01	0.0000	0.0000	1.0000
Energy	Fugitive emissions , 1B2c2i, Flaring oil	N ₂ O	0.06	0.06	0.0000	0.0000	1.0000
Solvents and other product use	Degreasing and Dry Cleaning	CO ₂	0.00	0.00	0.0000	0.0000	1.0000
Industrial processes	Asphalt roofing	CO ₂	0.02	0.02	0.0000	0.0000	1.0000
Energy	Fugitive emissions , 1B2c Venting	CO ₂	0.00	0.00	0.0000	0.0000	1.0000
Industrial processes	Magnesium Production	SF ₆	0.00	0.00	0.0000	0.0000	1.0000

IPCC Source Categories (LULUCF excluded)		GHG	Base Year Estimate	Year 2012 Estimate	Trend Asses- sment	Contribution to Trend	Cumulative total
			Ex,o Gg CO ₂ eq	Ex,t Gg CO ₂ eq	Tx,t		
<i>Continued</i>							
Energy	Fugitive emissions , 1B2ai Oil Exploration	CH ₄	0.00	0.00	0.0000	0.0000	1.0000
Energy	Fugitive emissions , 1B2ai Oil Exploration	CO ₂	0.00	0.00	0.0000	0.0000	1.0000
Energy	Fugitive emissions , 1B2ai Oil Exploration	N ₂ O	0.00	0.00	0.0000	0.0000	1.0000
Energy	Fugitive emissions , 1B2aii Oil Production	N ₂ O	0.00	0.00	0.0000	0.0000	1.0000
Total			69557.81	52247.69		1.0000	

Table 17.9 Key Category Analysis for Denmark and Greenland, trend assessment, incl. LULUCF.

IPCC Source Categories (LULUCF included)		GHG	Base Year Estimate Ex,o Gg CO ₂ eq	Year 2012 Estimate Ex,t Gg CO ₂ eq	Trend Assessment Tx,t	Contribution to Trend	Cumulative total
Energy	Combustion excluding transport, Solid Fuels	CO ₂	23982.671	10080.462	0.2310	0.3348	0.3348
Energy	Combustion excluding transport, Liquid Fuels	CO ₂	12310.843	6602.223	0.0949	0.1376	0.4723
LULUCF	Forest Land remaining Forest L., (blank)	CO ₂	49.604	-4491.413	0.0752	0.1090	0.5813
Energy	Combustion excluding transport, Gaseous Fuels	CO ₂	4335.403	8293.108	0.0654	0.0948	0.6761
Energy	Road transportation	CO ₂	9319.578	11257.010	0.0318	0.0461	0.7221
Agriculture	Agriculture soils, direct emissions , Synthetic Fertilizers	N ₂ O	2353.581	1104.262	0.0208	0.0301	0.7522
Industrial processes	Nitric acid production	N ₂ O	1042.902	0.000	0.0173	0.0251	0.7773
Agriculture	Agriculture soils, indirect, Nitrogen Leaching and Run-off	N ₂ O	2447.727	1430.546	0.0169	0.0245	0.8018
LULUCF	Cropland, 5.B Cropland, Organic soils	CO ₂	2887.334	1980.757	0.0151	0.0219	0.8237
LULUCF	Cropland, 5.B Cropland, Mineral soils	CO ₂	1415.313	577.084	0.0139	0.0202	0.8439
Energy	Combustion excluding transport, Other Fuels	CO ₂	575.139	1403.989	0.0137	0.0199	0.8638
Waste	Solid Waste Disposal Sites	CH ₄	1369.408	701.938	0.0111	0.0161	0.8799
Industrial processes	Refrigeration and AC Equipment	HFC and PFC	35.677	586.597	0.0091	0.0132	0.8931
LULUCF	Agricultural lime application, 5IV Cropland Limestone	CO ₂	622.931	192.305	0.0072	0.0104	0.9034
LULUCF	Grassland, 5.C Grassland, Living biomass	CO ₂	74.899	463.334	0.0064	0.0093	0.9128
Agriculture	Enteric Fermentation	CH ₄	3252.947	2909.179	0.0058	0.0084	0.9212
Agriculture	Manure Management	CH ₄	985.298	1297.300	0.0051	0.0074	0.9286
Energy	Domestic navigation	CO ₂	816.479	527.378	0.0048	0.0070	0.9356
LULUCF	Cropland, 5.B Cropland, Living biomass	CO ₂	-73.524	192.986	0.0044	0.0064	0.9420
Agriculture	Manure Management	N ₂ O	600.423	391.416	0.0035	0.0050	0.9470
Agriculture	Agriculture soils, indirect, Atmospheric Deposition	N ₂ O	496.501	294.695	0.0034	0.0049	0.9519
Agriculture	Agriculture soils, pasture, range and paddock	N ₂ O	335.049	211.525	0.0021	0.0030	0.9549
Energy	Combustion excluding transport, Gaseous Fuels	CH ₄	7.769	126.015	0.0020	0.0028	0.9577
Industrial processes	Foam Blowing	HFC	182.578	66.176	0.0019	0.0028	0.9605
Waste	Waste, other	N ₂ O	12.853	126.828	0.0019	0.0027	0.9632
Energy	Fugitive emissions , 1B2c2ii, Flaring gas	CO ₂	301.847	194.572	0.0018	0.0026	0.9658
Energy	Civil aviation	CO ₂	281.013	179.283	0.0017	0.0025	0.9683
Agriculture	Agriculture soils, direct emissions , Cultivation of Histosols	N ₂ O	290.294	198.450	0.0015	0.0022	0.9705
LULUCF	Wetlands, 5D Wetlands, soils	CO ₂	84.667	2.497	0.0014	0.0020	0.9725
LULUCF	Settlements, 5E Total settlements	CO ₂	12.921	90.806	0.0013	0.0019	0.9743
Industrial processes	Lime production	CO ₂	115.532	40.166	0.0013	0.0018	0.9762
Energy	Combustion excluding transport, Biomass	CH ₄	103.754	167.060	0.0010	0.0015	0.9777
Waste	Waste, other	CH ₄	30.723	91.971	0.0010	0.0015	0.9791
Energy	Combustion excluding transport, Biomass	N ₂ O	41.576	99.615	0.0010	0.0014	0.9805
Agriculture	Agriculture soils, direct emissions , Crop Residue	N ₂ O	361.266	311.296	0.0008	0.0012	0.9817
Waste	N2O indirect from human sewage	N ₂ O	85.233	35.172	0.0008	0.0012	0.9830
Energy	Railways	CO ₂	296.745	249.325	0.0008	0.0012	0.9841
Agriculture	Agriculture soils, direct emissions , Animal Manure Appl. to Soils	N ₂ O	1112.058	1161.673	0.0008	0.0011	0.9852
Energy	Fugitive emissions , 1B2aiv, Oil refining and storage	CH ₄	0.778	46.761	0.0008	0.0011	0.9863
Solvents and other product use	Other solvent	CO ₂	82.610	125.216	0.0007	0.0010	0.9874
Energy	Combustion excluding transport, Solid Fuels	N ₂ O	68.113	27.452	0.0007	0.0010	0.9883
LULUCF	Land converted to Forest L., (blank)	CO ₂	76.621	37.833	0.0006	0.0009	0.9893
Industrial processes	Other emissions of SF6 i.e. from double glaze windows and laboratories	SF ₆	67.616	104.589	0.0006	0.0009	0.9902
Energy	Road transportation	CH ₄	46.967	11.398	0.0006	0.0009	0.9910
Energy	Combustion excluding transport, Liquid Fuels	N ₂ O	84.591	55.364	0.0005	0.0007	0.9917
Industrial processes	Iron and steel production	CO ₂	28.447	0.000	0.0005	0.0007	0.9924

IPCC Source Categories (LULUCF included)		GHG	Base Year Estimate Ex,o Gg CO ₂ eq	Year 2012 Estimate Ex,t Gg CO ₂ eq	Trend Assessment Tx,t	Contribution to Trend	Cumulative total
<i>Continued</i>							
LULUCF	Grassland, 5.C Grassland, Organic soils	CO ₂	107.143	79.915	0.0005	0.0007	0.9931
Energy	Road transportation	N ₂ O	90.736	116.531	0.0004	0.0006	0.9937
Energy	Fugitive emissions , 1B2aii Oil Production	CH ₄	32.030	54.200	0.0004	0.0005	0.9942
Industrial processes	Other, lubricants	CO ₂	49.706	31.696	0.0003	0.0004	0.9946
Industrial processes	Aerosols	HFC	0.000	15.832	0.0003	0.0004	0.9950
Waste	N2O direct, Domestic and Commercial Wastewater	N ₂ O	34.604	49.460	0.0002	0.0004	0.9954
Agriculture	Agriculture soils, direct emissions , N-fixing Crops	N ₂ O	269.467	256.206	0.0002	0.0003	0.9957
Industrial processes	Cement production	CO ₂	882.402	871.083	0.0002	0.0003	0.9960
Industrial processes	Limestone and dolomite use	CO ₂	13.692	25.645	0.0002	0.0003	0.9963
Agriculture	Agriculture soils, direct emissions , Sludge	N ₂ O	27.923	38.762	0.0002	0.0003	0.9966
Solvents and other product use	Other solvent	N ₂ O	1.142	11.928	0.0002	0.0003	0.9968
Energy	Combustion excluding transport, Solid Fuels	CH ₄	12.876	3.121	0.0002	0.0002	0.9971
Industrial processes	Electrical equipment	SF ₆	3.908	13.266	0.0002	0.0002	0.9973
LULUCF	Cropland, 5.B Cropland, Dead organic matter	CO ₂	2.930	12.159	0.0002	0.0002	0.9975
Industrial processes	Expanded clay	CO ₂	14.929	5.887	0.0002	0.0002	0.9977
Waste	Waste Water Handling	CH ₄	65.487	74.001	0.0001	0.0002	0.9979
Energy	Combustion excluding transport, Gaseous Fuels	N ₂ O	16.380	24.446	0.0001	0.0002	0.9981
Industrial processes	Glass production	CO ₂	17.407	9.691	0.0001	0.0002	0.9983
Solvents and other product use	Chemical Products, Manufacture and Processing	CO ₂	19.424	12.175	0.0001	0.0002	0.9985
Energy	Domestic navigation	N ₂ O	15.047	8.943	0.0001	0.0001	0.9986
Solvents and other product use	Paint application	CO ₂	13.229	7.230	0.0001	0.0001	0.9988
Industrial processes	Yellow bricks	CO ₂	23.016	17.483	0.0001	0.0001	0.9989
LULUCF	Grassland, 5.C Grassland, Dead organic matter	CO ₂	1.633	7.042	0.0001	0.0001	0.9990
LULUCF	Grassland, 5.C Grassland, Mineral soils	CO ₂	0.156	4.537	0.0001	0.0001	0.9991
Industrial processes	Other i.e. Fibre Optics	HFC and PFC	0.000	4.314	0.0001	0.0001	0.9992
Energy	Combustion excluding transport, Other Fuels	N ₂ O	2.988	7.234	0.0001	0.0001	0.9993
Energy	Fugitive emissions , 1B2biii, Gas transmission	CH ₄	3.995	0.292	0.0001	0.0001	0.9994
LULUCF	Non-CO2 drainage of soils and wetlands, 5IID Forest Land.	N ₂ O	15.643	12.066	0.0001	0.0001	0.9995
LULUCF	Wetlands, 5D Wetlands Living biomass	CO ₂	2.567	-0.222	0.0000	0.0001	0.9996
Energy	Fugitive emissions , 1B2biv, Gas distribution	CH ₄	5.415	3.077	0.0000	0.0001	0.9996
Industrial processes	Food and drink	CO ₂	4.450	2.241	0.0000	0.0001	0.9997
Energy	Combustion excluding transport, Liquid Fuels	CH ₄	11.100	9.591	0.0000	0.0000	0.9997
Energy	Fugitive emissions , 1B2aii Oil Production	CO ₂	2.381	3.707	0.0000	0.0000	0.9998
Waste	Waste, other	CO ₂	17.538	16.356	0.0000	0.0000	0.9998
Energy	Civil aviation	N ₂ O	3.525	2.478	0.0000	0.0000	0.9998
LULUCF	N2O Disturbance, Land converted to cropland, 5III Cropland	N ₂ O	0.253	1.006	0.0000	0.0000	0.9998
Waste	Waste incineration	CH ₄	2.272	1.611	0.0000	0.0000	0.9999
Energy	Fugitive emissions , 1B2c2i, Flaring oil	CO ₂	22.950	22.333	0.0000	0.0000	0.9999
Industrial processes	Catalysts/Fertilizers, Pesticides and Sulphuric acid	CO ₂	0.800	1.407	0.0000	0.0000	0.9999
Agriculture	Field Burning of Agricultural Residues	CH ₄	1.824	2.429	0.0000	0.0000	0.9999
Waste	Waste incineration	CO ₂	2.550	3.135	0.0000	0.0000	0.9999
LULUCF	Wetlands, 5D Wetlands, dead org. matter	CO ₂	0.541	0.000	0.0000	0.0000	0.9999
LULUCF	Biomass burning, 5V Biomass burning	CH ₄	0.549	0.025	0.0000	0.0000	0.9999
LULUCF	Biomass burning, 5V Biomass burning	N ₂ O	0.449	0.034	0.0000	0.0000	0.9999
Energy	Combustion excluding transport, Other Fuels	CH ₄	0.361	0.774	0.0000	0.0000	1.0000
Energy	Railways	N ₂ O	2.535	2.131	0.0000	0.0000	1.0000
Energy	Fugitive emissions , 1B2c2ii, Flaring gas	CH ₄	0.730	0.325	0.0000	0.0000	1.0000

IPCC Source Categories (LULUCF included)		GHG	Base Year Estimate Ex,o Gg CO ₂ eq	Year 2012 Estimate Ex,t Gg CO ₂ eq	Trend Assessment Tx,t	Contribution to Trend	Cumulative total
<i>Continued</i>							
Energy	Fugitive emissions , 1B2c2ii, Flaring gas	N ₂ O	0.710	0.468	0.0000	0.0000	1.0000
Agriculture	Field Burning of Agricultural Residues	N ₂ O	0.698	0.930	0.0000	0.0000	1.0000
Waste	Waste incineration	N ₂ O	1.296	1.083	0.0000	0.0000	1.0000
Energy	Railways	CH ₄	0.259	0.153	0.0000	0.0000	1.0000
Energy	Civil aviation	CH ₄	0.097	0.056	0.0000	0.0000	1.0000
Energy	Domestic navigation	CH ₄	0.710	0.752	0.0000	0.0000	1.0000
Energy	Fugitive emissions , 1B2c Venting	CH ₄	1.226	1.206	0.0000	0.0000	1.0000
Energy	Fugitive emissions , 1B2b Natural gas transmission and distribution	CO ₂	0.008	0.005	0.0000	0.0000	1.0000
Energy	Fugitive emissions , 1B2c2i, Flaring oil	CH ₄	0.157	0.160	0.0000	0.0000	1.0000
Energy	Fugitive emissions , 1B2c2i, Flaring oil	N ₂ O	0.060	0.061	0.0000	0.0000	1.0000
Industrial processes	Road Paving with asphalt	CO ₂	1.762	1.766	0.0000	0.0000	1.0000
LULUCF	Non-CO2 drainage of soils and wetlands, 5IID Wetlands. Peatland	N ₂ O	0.135	0.135	0.0000	0.0000	1.0000
Solvents and other product use	Degreasing and Dry Cleaning	CO ₂	0.000	0.000	0.0000	0.0000	1.0000
Industrial processes	Asphalt roofing	CO ₂	0.019	0.019	0.0000	0.0000	1.0000
Energy	Fugitive emissions , 1B2c Venting	CO ₂	0.001	0.001	0.0000	0.0000	1.0000
Industrial processes	Magnesium Production	SF ₆	0.000	0.000	0.0000	0.0000	1.0000
Energy	Fugitive emissions , 1B2ai Oil Exploration	CH ₄	0.000	0.000	0.0000	0.0000	1.0000
Energy	Fugitive emissions , 1B2ai Oil Exploration	CO ₂	0.000	0.000	0.0000	0.0000	1.0000
Energy	Fugitive emissions , 1B2ai Oil Exploration	N ₂ O	0.000	0.000	0.0000	0.0000	1.0000
Energy	Fugitive emissions , 1B2aii Oil Production	N ₂ O	0.000	0.000	0.0000	0.0000	1.0000
Total			74840.574	51410.578		1.000	

17.4.1 Key category analysis for KP-LULUCF

The contribution from Greenland to the KP-LULUCF inventory is miniscule and the same categories are therefore identified as key as for the submission from Denmark, see Chapter 11.9 for more information.

17.5 Recalculations

17.5.1 Implications for emission levels

For the national total CO₂ equivalent emissions without Land-Use, Land-Use Change and Forestry, the general impact of the improvements and recalculations performed is small and the changes for the whole time series are between -0.10 % (1990) and 0.46 % (2011). Therefore, the implications of the recalculations on the level and on the trend, 1990-2011, of the recalculations are small, refer Table 17.10.

For the national total CO₂ equivalent emissions with Land-Use, Land-Use Change and Forestry, the general impact of the recalculations is larger due to recalculations in the LULUCF sector. The changes vary between -0.34 % (1990) and 0.54 % (2010), refer Table 17.10.

The impact of recalculation in the Greenlandic inventory is insignificant compared to the recalculations in the Danish inventory. Therefore the explanations and justifications are not repeated in this Chapter. Detailed information on the recalculations in the Danish inventory is provided in Chapter 10 and in the sectoral Chapters 3-8. The recalculations carried out for the Greenlandic inventory are described in Chapter 16.

Table 17.10 Recalculation performed on national total for 1990-2011. Differences in pct of CO₂ eqv between this submission and the May 2013 submission for Denmark under the Kyoto Protocol, i.e. Denmark & Greenland.

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
Total CO ₂ eqv. Emissions with Land-Use Change and Forestry	-0.34	-0.04	-0.03	-0.02	0.00	0.02	0.03	0.05	0.09	0.16	0.45
Total CO ₂ eqv. Emissions without Land-Use Change and Forestry	-0.10	-0.08	-0.07	-0.05	-0.03	-0.01	0.00	0.03	0.07	0.15	0.44
<i>Continued</i>	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011
Total CO ₂ eqv. Emissions with Land-Use Change and Forestry	0.17	0.28	0.34	0.16	-0.11	-0.06	0.26	-0.26	0.52	0.54	0.34
Total CO ₂ eqv. Emissions without Land-Use Change and Forestry	0.22	0.34	0.38	0.20	0.22	0.23	0.28	0.21	0.36	0.29	0.46

Table 17.11 Recalculation for CO₂ performed in the 2014 submission for 1990-2011. Differences in Gg CO₂ eqv. between this and the May 2013 submission for Denmark under the Kyoto Protocol, i.e. Denmark & Greenland.

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
Total National Emissions and Removals	-136	81	80	81	80	77	75	78	76	77	223
1. Energy	55	55	55	58	58	55	55	56	56	57	202
1.A. Fuel Combustion Activities	53	53	53	55	58	55	55	56	56	57	202
1.A.1. Energy Industries	-3	-3	-3	-1	66	4	10	9	10	10	10
1.A.2. Manufacturing Industries and Construction	59	59	59	59	47	55	49	51	51	51	51
1.A.3. Transport	-1	-1	-1	-1	-1	-1	-1	-1	-1	-1	-1
1.A.4. Other Sectors	-2	-3	-3	-3	-55	-3	-4	-4	-4	-4	142
1.A.5. Other	0	0	0	0	0	0	0	0	0	0	0
1.B. Fugitive Emissions from Fuels	2	3	2	3	0	0	0	0	0	0	0
2. Industrial Processes	-	-	-	-	-	1	0	3	2	3	5
2.A. Mineral Products	-	-	-	-	-	1	1	2	2	3	5
2.B. Chemical Industry	-	-	-	-	-	-	-1	1	-	-	0
2.C. Metal Production	-	-	-	-	-	-	-	-	-	-	-
2.D. Other Production	-	-	-	-	-	-	-	-	-	-	-
2.G. Other	-	-	-	-	-	-	-	-	-	-	-
3. Solvent and Other Product Use	-	-	-	-	-	-	-	-	-	-	-
4. Agriculture	-	-	-	-	-	-	-	-	-	-	-
5. Land Use, Land-Use Change and Forestry (net)	-190	26	25	24	23	22	21	20	19	18	17
5.A. Forest Land	7	7	7	7	7	7	7	7	7	7	7
5.B. Cropland	-192	26	26	27	27	27	27	27	27	27	28
5.C. Grassland	0	0	0	0	0	0	0	0	0	0	0
5.D. Wetlands	-3	-5	-6	-7	-8	-9	-10	-11	-12	-13	-14
5.E. Settlements	-3	-3	-3	-3	-4	-4	-4	-4	-4	-4	-4
5.F. Other Land	-	-	-	-	-	-	-	-	-	-	-
5.G. Other	-	-	-	-	-	-	-	-	-	-	-
6. Waste	-1	-1	-1	-1	-1	-1	-1	0	0	0	0
6.C. Waste Incineration	-	-	-	-	-	-	-	-	-	-	-
6.D. Other	-1	-1	-1	-1	-1	-1	-1	0	0	0	0
<i>Continued</i>	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011
Total National Emissions and Removals	33	35	74	75	-126	-112	94	-253	234	242	60
1. Energy	57	57	94	94	94	102	102	45	131	94	139
1.A. Fuel Combustion Activities	57	57	94	94	94	102	102	45	131	94	139
1.A.1. Energy Industries	9	9	32	11	10	6	5	12	32	27	5
1.A.2. Manufacturing Industries and Construction	53	52	84	87	89	85	124	110	107	152	132
1.A.3. Transport	-1	-1	-1	-1	-1	-1	0	-70	0	-4	-13
1.A.4. Other Sectors	-4	-4	-21	-3	-5	11	-26	-7	-8	-81	15
1.A.5. Other	0	0	0	0	0	0	0	0	0	0	0
1.B. Fugitive Emissions from Fuels	0	0	0	0	0	0	0	0	0	0	0
2. Industrial Processes	2	1	1	-1	-1	3	1	1	1	4	-2
2.A. Mineral Products	2	1	1	1	1	3	2	3	2	4	-2
2.B. Chemical Industry	0	0	-1	-2	-2	-1	-1	-1	-1	-	-1
2.C. Metal Production	-	-	-	-	-	-	-	-	-	-	-
2.D. Other Production	-	-	-	-	-	-	-	-	-	-	-
2.G. Other	-	-	-	-	-	-	-	-	-	-	-
3. Solvent and Other Product Use	-	-	-	-	-	-	-	-	-12	-7	1
4. Agriculture	-	-	-	-	-	-	-	-	-	-	-
5. Land Use, Land-Use Change and Forestry (net)	-25	-23	-21	-19	-219	-216	-10	-299	114	150	-78
5.A. Forest Land	-34	-30	-27	-23	-51	-49	157	-132	280	316	88
5.B. Cropland	28	27	27	27	-181	-181	-181	-181	-181	-181	-181
5.C. Grassland	0	0	0	0	32	31	31	31	30	30	29
5.D. Wetlands	-15	-16	-17	-18	-14	-13	-13	-12	-11	-11	-10
5.E. Settlements	-4	-4	-4	-4	-5	-5	-5	-4	-4	-4	-4

<i>Continued</i>												
5.F. Other Land	-	-	-	-	-	-	-	-	-	-	-	-
5.G. Other	-	-	-	-	-	-	-	-	-	-	-	-
6. Waste	0	0	0	0	0	0	0	0	0	0	0	0
6.C. Waste Incineration	-	-	-	-	-	-	-	-	-	-	-	-
6.D. Other	0	0	0	0	0	0	0	0	0	0	0	0

Table 17.12 Recalculation for CH₄ performed in the 2014 submission for 1990-2011. Differences in Gg CO₂ eqv. between this and the May 2013 submission for Denmark under the Kyoto Protocol, i.e. Denmark & Greenland.

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
Total National Emissions and Removals	-113.9	-108.2	-101.0	-89.3	-74.7	-60.7	-50.6	-37.5	-28.3	-20.5	-16.8
1. Energy	5.6	6.8	6.4	7.3	7.3	8.6	7.7	7.0	6.1	4.7	6.0
1.A. Fuel Combustion Activities	4.8	5.6	5.8	6.4	6.8	7.1	7.5	6.6	5.7	6.2	7.8
1.A.1. Energy Industries	0.0	0.0	0.0	0.0	0.6	1.9	3.4	2.2	2.0	2.2	2.3
1.A.2. Manufacturing Industries and Construction	0.0	0.0	0.0	0.0	0.0	-1.2	-2.8	-2.5	-2.3	-2.4	-2.5
1.A.3. Transport	-0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
1.A.4. Other Sectors	4.9	5.7	5.9	6.5	6.2	6.4	6.9	6.9	6.1	6.3	8.1
1.A.5. Other	-	-	-	-	-	-	-	-	-	0.0	0.0
1.B. Fugitive Emissions from Fuels	0.8	1.2	0.6	0.9	0.5	1.5	0.2	0.4	0.4	-1.5	-1.8
2. Industrial Processes	-	-	-	-	-	-	-	-	-	-	-
3. Solvent and Other Product Use	-	-	-	-	-	-	-	-	-	-	-
4. Agriculture	-8.0	-8.9	-9.6	-10.4	-10.8	-11.6	-12.7	-12.7	-12.6	-12.3	-12.4
4.A. Enteric Fermentation	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
4.B. Manure Management	-8.0	-8.9	-9.6	-10.4	-10.8	-11.6	-12.7	-12.7	-12.6	-12.3	-12.4
4.F. Field Burning of Agricultural Residues	-	-	-	-	-	-	-	-	-	-	-
5. Land Use, Land-Use Change and Forestry (net)	-	-	-	-	-	-	-	-	-	-	-
6. Waste	-111.5	-106.2	-97.8	-86.2	-71.3	-57.7	-45.6	-31.8	-21.8	-12.9	-10.5
6.A. Solid Waste Disposal on Land	-112.0	-106.9	-98.9	-87.5	-72.7	-60.1	-47.6	-34.3	-24.5	-16.4	-14.9
6.B. Waste-water Handling	-0.7	-0.7	-0.7	-0.7	-0.8	-0.8	-1.0	-1.2	-1.1	-1.3	-0.9
6.C. Waste Incineration	-	-	-	-	-	-	-	-	-	-	-
6.D. Other	1.2	1.5	1.8	2.0	2.2	3.2	3.0	3.7	3.8	4.9	5.4
<i>Continued</i>	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011
Total National Emissions and Removals	-14.3	-5.7	14.7	16.8	22.6	28.5	37.4	36.7	42.3	22.2	53.4
1. Energy	8.7	9.1	10.8	12.3	14.7	16.2	20.6	18.3	16.7	18.7	17.0
1.A. Fuel Combustion Activities	8.8	9.1	10.6	12.2	14.6	16.1	20.3	18.5	16.9	19.1	16.9
1.A.1. Energy Industries	1.9	2.1	1.9	2.2	2.0	2.0	1.8	1.7	2.0	2.0	2.1
1.A.2. Manufacturing Industries and Construction	-2.1	-2.3	-2.1	-2.3	-2.0	-2.3	-1.8	-2.0	-2.2	-0.9	-0.8
1.A.3. Transport	0.0	0.0	0.0	0.0	0.1	0.0	0.1	0.0	0.3	0.3	0.3
1.A.4. Other Sectors	9.0	9.3	10.8	12.3	14.6	16.4	20.2	18.7	16.9	17.8	15.3
1.A.5. Other	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
1.B. Fugitive Emissions from Fuels	0.0	0.0	0.1	0.1	0.1	0.1	0.3	-0.2	-0.2	-0.4	0.0
2. Industrial Processes	-	-	-	-	-	-	-	-	-	-	-
3. Solvent and Other Product Use	-	-	-	-	-	-	-	-	-	-	-
4. Agriculture	-13.0	-12.4	0.0	0.0	0.0	0.0	0.0	0.0	0.0	-1.0	-0.2
4.A. Enteric Fermentation	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	-0.1
4.B. Manure Management	-13.0	-12.4	0.0	0.0	0.0	0.0	0.0	0.0	0.0	-1.0	-0.2
4.F. Field Burning of Agricultural Residues	-	-	-	-	-	-	-	-	-	-	-
5. Land Use, Land-Use Change and Forestry (net)	-	-	-	-	-	-	-	-	-	-	-
6. Waste	-10.0	-2.4	4.0	4.5	7.9	12.3	16.8	18.5	25.6	4.4	36.7
6.A. Solid Waste Disposal on Land	-14.4	-8.2	-1.9	2.0	5.9	9.6	13.9	17.1	21.2	0.2	32.3
6.B. Waste-water Handling	-1.4	-1.8	-1.9	-2.1	-1.9	-1.6	-1.3	-3.0	-1.9	-1.2	-1.3
6.C. Waste Incineration	-	-	-	-	-	-	-	-	-	-	-
6.D. Other	5.7	7.7	7.8	4.6	3.9	4.3	4.2	4.3	6.3	5.5	5.7

Table 17.13 Recalculation for N₂O performed in the 2014 submission for 1990-2011. Differences in Gg CO₂ eqv. between this and the May 2013 submission for Denmark under the Kyoto Protocol, i.e. Denmark & Greenland.

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
Total National Emissions and Removals	-7.6	-6.9	-6.1	-7.4	-6.8	-2.3	-1.2	0.7	26.1	69.6	115.9
1. Energy	1.7	2.3	1.9	1.4	1.4	0.9	1.6	1.8	0.6	2.2	3.6
1.A. Fuel Combustion Activities	1.8	2.4	1.9	1.5	1.4	0.9	1.6	1.8	0.6	2.2	3.6
1.A.1. Energy Industries	0.7	1.2	0.8	0.4	0.4	0.3	0.9	1.2	0.0	1.5	0.0
1.A.2. Manufacturing Industries and Construction	1.1	1.1	1.1	1.1	1.1	0.6	0.7	0.6	0.6	0.6	0.7
1.A.3. Transport	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
1.A.4. Other Sectors	-	0.0	0.0	0.0	-0.1	0.0	0.0	0.1	0.0	0.0	2.9
1.A.5. Other	-	-	-	-	-	-	-	-	-	0.0	0.0
1.B. Fugitive Emissions from Fuels	-0.1	-0.1	-0.1	-0.1	-	-	-	-	-	-	-
2. Industrial Processes	-	-	-	-	-	-	-	-	-	-	-
3. Solvent and Other Product Use	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	-
4. Agriculture	-10.5	-10.5	-9.7	-10.7	-10.3	-9.8	-8.7	-8.2	-7.6	-6.8	-6.1
4.A. Enteric Fermentation	-	-	-	-	-	-	-	-	-	-	-
4.B. Manure Management	-	-	-	-	-	-	-	-	-	-	-
4.D. Agricultural Soils	-10.5	-10.5	-9.7	-10.7	-10.3	-9.8	-8.7	-8.2	-7.6	-6.8	-6.1
4.F. Field Burning of Agricultural Residues	-	-	-	-	-	-	-	-	-	-	-
5. Land Use, Land-Use Change and Forestry (net)	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.1	0.1
5.A. Forest Land	-0.2	-0.2	-0.2	-0.2	-0.2	-0.2	-0.2	-0.2	-0.2	-0.2	-0.2
5.B. Cropland	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
5.C. Grassland	-	-	-	-	-	-	-	-	-	-	-
5.D. Wetlands	-	-	-	-	-	-	-	-	-	-	-
5.E. Settlements	-	-	-	-	-	-	-	-	-	-	-
5.F. Other Land	-	-	-	-	-	-	-	-	-	-	-
5.G. Other	-	-	-	-	-	-	-	-	-	-	-
6. Waste	1.1	1.3	1.6	1.8	2.1	6.5	5.9	7.0	33.1	74.2	118.4
6.B. Waste-water Handling	-	-	-	-	-	-	-	-	-	-	-
6.C. Waste Incineration	-	-	-	-	-	-	-	-	-	-	-
6.D. Other	1.1	1.3	1.6	1.8	2.1	6.5	5.9	7.0	33.1	74.2	118.4
<i>Continued</i>	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011
Total National Emissions and Removals	110.3	184.0	178.3	28.8	27.3	36.5	48.7	49.7	59.7	65.2	73.4
1. Energy	0.8	0.8	1.2	2.3	2.7	2.0	1.9	1.5	1.9	2.1	2.3
1.A. Fuel Combustion Activities	0.8	0.8	1.2	2.3	2.7	2.0	1.9	1.5	1.9	2.1	2.3
1.A.1. Energy Industries	0.1	0.0	0.4	0.7	0.8	0.6	0.0	0.1	0.3	0.0	0.0
1.A.2. Manufacturing Industries and Construction	0.7	0.7	1.3	1.4	1.5	1.4	2.2	1.7	1.9	3.8	3.3
1.A.3. Transport	0.0	0.1	0.1	0.2	0.4	-0.2	-0.1	-0.6	0.0	0.0	0.0
1.A.4. Other Sectors	0.0	0.0	-0.6	0.0	0.0	0.2	-0.4	0.3	-0.3	-1.8	-1.0
1.A.5. Other	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
1.B. Fugitive Emissions from Fuels	-	-	-	-	-	-	-	-	-	-	-
2. Industrial Processes	-	-	-	-	-	-	-	-	-	-	-
3. Solvent and Other Product Use	-	-	-	-	-	-	-	-	0.0	0.0	0.0
4. Agriculture	-5.6	-5.0	-5.0	-5.2	-4.8	-4.2	-4.2	-4.6	-3.5	-4.3	-2.8
4.A. Enteric Fermentation	-	-	-	-	-	-	-	-	-	-	-
4.B. Manure Management	-	-	-	-	-	-	-	-	-	-0.6	0.0
4.D. Agricultural Soils	-5.6	-5.0	-5.0	-5.2	-4.8	-4.2	-4.2	-4.6	-3.5	-3.7	-2.8
4.F. Field Burning of Agricultural Residues	-	-	-	-	-	-	-	-	-	-	-
5. Land Use, Land-Use Change and Forestry (net)	0.1	0.1	0.1	0.1	-0.1	-0.1	-0.1	0.0	0.0	0.0	0.0
5.A. Forest Land	-0.2	-0.2	-0.2	-0.2	-0.2	-0.2	-0.2	-0.2	-0.2	-0.2	-0.2
5.B. Cropland	0.2	0.2	0.3	0.3	0.1	0.1	0.1	0.1	0.1	0.2	0.2
5.C. Grassland	-	-	-	-	-	-	-	-	-	-	-

<i>Continued</i>											
5.D. Wetlands	-	-	-	-	-	-	-	-	-	-	-
5.E. Settlements	-	-	-	-	-	-	-	-	-	-	-
5.F. Other Land	-	-	-	-	-	-	-	-	-	-	-
5.G. Other	-	-	-	-	-	-	-	-	-	-	-
6. Waste	115.0	188.1	182.0	31.7	29.6	38.8	51.1	52.9	61.4	67.4	73.9
6.B. Waste-water Handling	-	-	-	-	-	-	-	-	-	-	-
6.C. Waste Incineration	-	-	-	-	-	-	-	-	-	-	-
6.D. Other	115.0	188.1	182.0	31.7	29.6	38.8	51.1	52.9	61.4	67.4	73.9

Table 17.14 Recalculation for HFCs, PFCs and SF₆ performed in the 2014 submission for 1990-2011. Differences in Gg CO₂ eqv. between this and the May 2013 submission for Denmark under the Kyoto Protocol, i.e. Denmark & Greenland.

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
HFC	-	-	-	-	-	-	-	-	-	-	-
PFC	-	-	-	-	-	-	-	-	-	-	-
SF ₆	-	-	-	-	-	-	-	-	-	-	-
<i>Continued</i>	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011
HFC	-	-	-	-	-	-	-	-	0.0	0.0	-
PFC	-	-	-	-	-	-	-	-	-	-	-
SF ₆	-	-	-	-	-	-	-	-	-	-0.4	-

17.6 Technical description of the aggregation of the emission inventories of Denmark and Greenland

In order to accommodate the request of the ERT of full inclusion of the Greenlandic emission data in the full CRF format, Denmark operates separate installations for Denmark and Greenland (and the Faroe Islands). The country identification codes provided by the UNFCCC secretariat are DNM for Denmark and GRL for Greenland (FRO for the Faroe Islands). Two additional installations are necessary to enable the submission of aggregated submissions under the Kyoto Protocol (Denmark and Greenland) and under UNFCCC (Denmark, Greenland and the Faroe Islands). The country identification codes provided by the UNFCCC secretariat are DKE for the submission under the Kyoto Protocol (Denmark and Greenland) and DNK for the UNFCCC submission (Denmark, Greenland and the Faroe Islands).

These five versions of CRF Reporter are installed on separate virtual MS Windows XP machines. The installations are at the AU VMWare environment, which is operated and maintained by the IT department at AU. As such backups of these systems are performed routinely on a daily basis.

For the aggregation of the submissions three IT tools are used.

- EU CRF Aggregator developed by the European Environment Agency – Aggregation of global CRF variables
- NERI CRF Aggregator developed by NERI (Now DCE) – Aggregation of local CRF variables
- MS Excel

The three main work processes in connection with the aggregation of the submissions are:

- In the EU CRF Aggregator the following work processes take place:
 - Aggregation of global variables; sum of emissions and activity data, notation keys and comments.
 - As input data the xml submission files from the CRF Reporter installations for DNM (Denmark), GRL (Greenland) and FRO (Faroe Islands) are used.
 - As output file a CRF Reporter xml import file is generated. This file is then imported in the installation for the aggregated submission, DKE (KP) or DNK (UNFCCC).
- In NERI CRF Aggregator the following work processes take place:
 - Aggregation of local variables; sum of emissions and activity data, notation keys and comments. Aggregation of additional information variables either as sums or uniform values.
 - As input data the simple CRF Reporter xml files from the CRF Reporter installations for DNM (Denmark), GRL (Greenland) and FRO (Faroe Islands) are used.
 - As output file a CRF Reporter simple xml import file is generated. This file is then imported in the installation for the aggregated submission, DKE (KP) or DNK (UNFCCC).
- In MS Excel the following work processes take place:
 - Aggregation of additional information variables where average values or weighted average values are used.
 - Aggregation of KP-LULUCF/NIR-1 and KP-LULUCF/NIR-2.
 - The aggregated data is at the moment copy/pasted from the CRF Reporter installations of Denmark and Greenland to Excel aggregated and copy/pasted back to the CRF Reporter installations of the KP submission (DKE).

Efforts are ongoing to ensure the highest possible degree of automation to avoid the risk of errors during the manual work processes.

17.7 QA/QC of the aggregated submission for Denmark and Greenland

The QA/QC procedures for the Danish inventory are described in Chapter 1.6 and the sectoral chapters. Please refer to Chapter 1.6 for a general description of the QA/QC system, and the structural setup of the Danish QA/QC system for the greenhouse gas inventory. The QA/QC procedures carried out by Greenlandic authorities for the Greenlandic inventory are described in Chapter 16. The following focuses on the specific QA/QC measures carried out at DCE both on the data (CRF tables and documentation) received from Greenland and the QC checks carried out for the aggregated versions of the inventory for reporting to the Kyoto Protocol and the UNFCCC. The PM's relevant for this are listed in Table 17.15.

Table 17.15 PM's specific to the handling of Greenlandic emission data and the aggregated submissions.

Data Storage level 4	3.Completeness	DS.4.3.3	Check that no sources where methodology exists in the IPCC guidelines are reported as NE by Greenland.
	4.Consistency	DS.4.4.2	Check time series consistency of the reporting by Greenland prior to aggregating the final submissions.
	5.Correctness	DS.4.5.1	Check that the aggregated submissions for Denmark under the Kyoto Protocol and the UNFCCC match the sum of the individual submissions.
		DS.4.5.2	Check that additional information and information related to land-use changes has been correctly aggregated compared to the individual submissions of Denmark and Greenland.
	7.Transparency	DS.4.7.2	Perform QA on the documentation report provided by the Government of Greenland.

Data Storage level 4	3.Completeness	DS.4.3.3	Check that no sources where a methodology exists in the IPCC guidelines or good practice guidance are reported as NE by Greenland
----------------------	----------------	----------	---

A check is made to filter any NE's from the CRF tables. If any greenhouse gas emissions are reported as NE, it is checked whether methodologies exist in the IPCC guidelines or the IPCC good practice guidance. If methodologies do exist efforts are made to quickly estimate and report emissions. No categories where methodology exists were identified for the submission of Denmark and Greenland.

Data Storage level 4	4.Consistency	DS.4.4.2	Check time series consistency of the reporting of Greenland and the Faroe Islands prior to aggregating the final submissions
----------------------	---------------	----------	--

The time series for all pollutants in the submissions from Greenland and the Faroe Islands are checked at the CRF 3 level for large variations in the time series. Any large variations are explained or corrected in cooperation with the authorities in Greenland and the Faroe Islands.

Data Storage level 4	5.Correctness	DS.4.5.1	Check that the aggregated submissions for Denmark under the Kyoto Protocol and the UNFCCC matches the sum of the individual submissions
----------------------	---------------	----------	---

To ensure that the submission for Denmark under the Kyoto Protocol matches the sum of the submissions of Denmark and Greenland a spreadsheet check has been implemented to ensure complete correctness of the submitted inventory. The same procedure is followed for the submission under the UNFCCC, where it is ensured that the submitted emissions equate to the sum of Denmark, Greenland and the Faroe Islands. Special attention is paid to the additional information provided in the CRF, e.g. for the agricultural sector. Certain parameters cannot simply be added, e.g. animal weights. In these cases a weighted average is reported in the CRF tables.

Since the 2012 submission the checks have been extended to also cover area information reported in the KP-LULUCF tables (NIR-2).

Data Storage level 4	5. Correctness	DS.4.5.2	Check that additional information and information related to land-use changes has been correctly aggregated compared to the individual submissions of Denmark and Greenland.
----------------------	----------------	----------	--

The CRF submission for Denmark and Greenland is checked to see if the additional information has been aggregated correctly. The additional information is mainly related to the agricultural and waste sectors.

Data Storage level 4	7. Transparency	DS.4.7.2	Perform QA on the documentation report provided by the Government of Greenland
----------------------	-----------------	----------	--

The documentation report is received by DCE from the Government of Greenland in the early spring every year. The documentation report is included in the NIR as Chapter 16. DCE experts read and provide comments on the report to the Government of Greenland, so that any questions are resolved prior to the UNFCCC reporting deadline of April 15.

Annex 1 - Key category analysis

Description of the methodology used for identifying key categories

Key Category Analysis (KCA) tier 1 and 2 for year 1990 and 2012 for Denmark (excluding Greenland and Faroe Islands) has been carried out in accordance with the IPCC Good Practice Guidance / IPCC Guidelines (2006). The KCA has been carried out excluding and including the LULUCF sector. A tier 1 KCA has also been worked out for Greenland and for Denmark and Greenland; refer to Chapter 16 and Chapter 17, respectively.

The base year in the analysis is the year 1990 for the greenhouse gases CO₂, CH₄, N₂O and 1995 for the F-gases HFC, PFC and SF₆. The KCA approaches are

- a tier 1 quantitative analysis and
- a tier 2 approach using tier 1 uncertainties.

The level assessment of the tier 1 KCA is a ranking of the source categories in accordance to their relative contribution to the national total of greenhouse gases calculated in CO₂ equivalent units. The level key categories are found from the list of source categories ranked according to their contribution in descending order. Level key categories are those from the top of the list and of which the sum constitutes 95 % of the national total.

The trend assessment of the tier 1 KCA is a ranking of the source categories according to their contribution to the trend of the national total of greenhouse gases, calculated in CO₂ equivalents, from the base year to the year under consideration. The trend of the source category is calculated relative to that of the national totals and the trend is then weighted with the contribution, according to the level assessment. The ranking is in descending order. As for the level assessment, the cut-off point for the sum of contribution to the trend is 95 % and the source categories from the top of the list to the cut-off line are trend key categories.

In addition, a tier 2 KCA has been carried out to provide additional insight into categories being key sources. The categorisation used is as for the tier 1 analysis and the uncertainties used are tier 1 uncertainties as listed in Annex 7.

The level tier 2 KCA is a ranking of the categories according to their relative contribution to the national total multiplied by the uncertainty of the emission of the category as the combined uncertainty on activity data and on emission factor. Chosen for cut of for key categories in the analysis is 90 %.

The trend tier 2 KCA is a ranking of the categories according to their relative contribution to the trend 1990-2012 of the national total multiplied by the uncertainty of the emission of the category. Chosen for cut of for key categories in the analysis is 90 %.

Since the level KCA is carried out for 1990 and 2012 (exclusive and inclusive LULUCF) and for tier 1 and 2 a total 12 KCA tables for Denmark (excluding Greenland and Faroe Islands) has been worked out following the sugges-

tions in the GPG Tables 7.A1-2. Further, two overview tables based on the GPG Table 7.A3 (exclusive and inclusive LULUCF) are shown. The overview table shows summary results of the KCA for 1990, for 2012, and for the trend 1990-2012.

The inclusion of the LULUCF sector in the level analysis implies that the emissions in this sector are all calculated positive, i.e. the absolute value of removals are included. Note also that according to the GPG, the analysis implies that contributions to the trend are all calculated as mathematically positive to be able to perform the ranking. The LULUCF activities are in the table included with their sign, i.e. emissions: +, removals: -.

The level of disaggregation

The starting-point for source categories is GPG Table 7.1. This table constitutes a suggested list of source categories for the KCA. It is mentioned in the GPG that categories for the KCA should be chosen in a way so that emissions from a single category are estimated with the same method and the same emission factor. Therefore, for categories in Table 7.1, which in our Corinair database are composed of activities with different emission factors or estimated with different methods, splits were made accordingly.

The categorisation has been somewhat revised compared to the 2011 submission. The categories follow the categorisation used for the uncertainty analyses, cf. Annex 7.

The source categories in the KCA for stationary combustion are defined according to the greenhouse gas and fuel. For CH₄ and N₂O fuels are aggregated to the fuel categories solid, liquid, gas, waste and biomass.

Table A1.1 KCA source categories for stationary combustion.

CRF, part of category	KCA category	GHG
1A1, 1A2 and 1A4	Stationary Combustion, Coal	CO ₂
1A1, 1A2 and 1A4	Stationary Combustion, BKB	CO ₂
1A1, 1A2 and 1A4	Stationary Combustion, Coke	CO ₂
1A1, 1A2 and 1A4	Stationary Combustion, Fossil waste	CO ₂
1A1, 1A2 and 1A4	Stationary Combustion, Petroleum coke	CO ₂
1A1, 1A2 and 1A4	Stationary Combustion, Residual oil	CO ₂
1A1, 1A2 and 1A4	Stationary Combustion, Gas oil	CO ₂
1A1, 1A2 and 1A4	Stationary Combustion, Kerosene	CO ₂
1A1, 1A2 and 1A4	Stationary Combustion, LPG	CO ₂
1A1, 1A2 and 1A4	Stationary Combustion, Refinery gas	CO ₂
1A1, 1A2 and 1A4	Stationary Combustion, Natural gas	CO ₂
1A1, 1A2 and 1A4	Stationary Combustion, SOLID	CH ₄
1A1, 1A2 and 1A4	Stationary Combustion, LIQUID	CH ₄
1A1, 1A2 and 1A4	Stationary Combustion, GAS	CH ₄
1A1, 1A2 and 1A4	Natural gas fuelled engines, GAS	CH ₄
1A1, 1A2 and 1A4	Stationary Combustion, WASTE	CH ₄
1A1, 1A2 and 1A4	Stationary Combustion, BIOMASS	CH ₄
1A1, 1A2 and 1A4	Biogas fuelled engines, BIOMASS	CH ₄
1A1, 1A2 and 1A4	Stationary Combustion, SOLID	N ₂ O
1A1, 1A2 and 1A4	Stationary Combustion, LIQUID	N ₂ O
1A1, 1A2 and 1A4	Stationary Combustion, GAS	N ₂ O
1A1, 1A2 and 1A4	Stationary Combustion, WASTE	N ₂ O
1A1, 1A2 and 1A4	Stationary Combustion, BIOMASS	N ₂ O

KCA source categories for mobile combustion are shown in Table A1-2. The categorisation is used for CO₂, CH₄ and N₂O.

Table A1.2 KCA source categories for mobile combustion.

CRF, part of category	KCA category	GHG
1.A.3.b	Transport, Road transport	CO ₂
1.A.5.b	Transport, Military	CO ₂
1.A.3.c	Transport, Railways	CO ₂
1.A.3.d (part)	Transport, Navigation (small boats)	CO ₂
1.A.3.d (part)	Transport, Navigation (large vessels)	CO ₂
1.A.4.c (part)	Transport, Fisheries	CO ₂
1.A.4.c (part)	Transport, Agriculture	CO ₂
1.A.4.c (part)	Transport, Forestry	CO ₂
1.A.2.f (part)	Transport, Industry (mobile)	CO ₂
1.A.4.b (part)	Transport, Residential	CO ₂
1.A.4.a (part)	Transport, Commercial/institutional	CO ₂
1.A.3.a	Transport, Civil aviation	CO ₂
1.A.3.b	Transport, Road transport	CH ₄
1.A.5.b	Transport, Military	CH ₄
1.A.3.c	Transport, Railways	CH ₄
1.A.3.d (part)	Transport, Navigation (small boats)	CH ₄
1.A.3.d (part)	Transport, Navigation (large vessels)	CH ₄
1.A.4.c (part)	Transport, Fisheries	CH ₄
1.A.4.c (part)	Transport, Agriculture	CH ₄
1.A.4.c (part)	Transport, Forestry	CH ₄
1.A.2.f (part)	Transport, Industry (mobile)	CH ₄
1.A.4.b (part)	Transport, Residential	CH ₄
1.A.4.a (part)	Transport, Commercial/institutional	CH ₄
1.A.3.a	Transport, Civil aviation	CH ₄
1.A.3.b	Transport, Road transport	N ₂ O
1.A.5.b	Transport, Military	N ₂ O
1.A.3.c	Transport, Railways	N ₂ O
1.A.3.d (part)	Transport, Navigation (small boats)	N ₂ O
1.A.3.d (part)	Transport, Navigation (large vessels)	N ₂ O
1.A.4.c (part)	Transport, Fisheries	N ₂ O
1.A.4.c (part)	Transport, Agriculture	N ₂ O
1.A.4.c (part)	Transport, Forestry	N ₂ O
1.A.2.f (part)	Transport, Industry (mobile)	N ₂ O
1.A.4.b (part)	Transport, Residential	N ₂ O
1.A.4.a (part)	Transport, Commercial/institutional	N ₂ O
1.A.3.a	Transport, Civil aviation	N ₂ O

For fugitive emissions, the categorisation used in the KCA is shown in Table A1-3.

Table A1-3 KCA source categories for fugitive emissions.

CRF and KCA category	GHG
1.B.2 Flaring in refinery	CO ₂
1.B.2 Flaring off-shore	CO ₂
1.B.2 Land based activities	CO ₂
1.B.2 Off-shore activities	CO ₂
1.B.2 Transmission of natural gas	CO ₂
1.B.2 Distribution of natural gas	CO ₂
1.B.2 Venting in gas storage	CO ₂
1.B.2. Flaring in refinery	CH ₄
1.B.2. Flaring off-shore	CH ₄
1.B.2 Refinery processes	CH ₄
1.B.2 Land based activities	CH ₄
1.B.2 Off-shore activities	CH ₄
1.B.2 Transmission of natural gas	CH ₄
1.B.2 Distribution of natural gas	CH ₄
1.B.2 Venting in gas storage	CH ₄
1.B.2 Flaring in refinery	N ₂ O
1.B.2 Flaring off-shore	N ₂ O

KCA categories for industry are shown in Table A1-4. All data can be found in CRF. Base year for the consumption of HFC, PFC and SF₆ is 1995.

Table A1-4 KCA source categories for industry.

CRF and KCA category	GHG
2A1 Cement production	CO ₂
2A2 Lime production	CO ₂
2A3 Limestone and dolomite use	CO ₂
2A5 Asphalt roofing	CO ₂
2A6 Road paving with asphalt	CO ₂
2A7a Glass and Glass wool	CO ₂
2A7b Yellow bricks	CO ₂
2A7c Expanded clay	CO ₂
2B5 Catalysts/Fertilizers, Pesticides and Sulphuric acid	CO ₂
2C1 Iron and steel production	CO ₂
2D2 Food and Drink	CO ₂
2G Lubricants	CO ₂
2B2 Nitric acid production	N ₂ O
2F Consumption of HFC	HFC
2F Consumption of PFC	PFC
2F Consumption of SF ₆	SF ₆

KCA categories for solvents are shown in Table A1-5. All data can be found in CRF.

Table A1-5 KCA source categories for solvents.

CRF and KCA category	GHG
3A Paint application	CO ₂
3B Degreasing and dry cleaning	CO ₂
3C Chemical products, manufacturing and processing	CO ₂
3D5 Other	CO ₂
3D5 Consumption of fireworks	CO ₂
3D5 Use of candles	CO ₂
3D1 Other - Use of N ₂ O for Anaesthesia	N ₂ O
3D5 Use of tobacco	N ₂ O
3D5 Use of charcoal for BBQ	N ₂ O
3D5 Consumption of fireworks	N ₂ O
3D5 Use of candles	N ₂ O

KCA categories for agriculture are shown in Table A1-6. All data can be found in CRF.

Table A1-6 KCA source categories for agriculture.

CRF and KCA category	GHG
4A Enteric Fermentation	CH ₄
4B Manure Management	CH ₄
4F Field burning of agricultural residues	CH ₄
4.B Manure Management	N ₂ O
4.D1.1 Synthetic Fertilizer	N ₂ O
4.D1.2 Animal waste applied to soils	N ₂ O
4.D1.3 N-fixing crops	N ₂ O
4.D1.4 Crop Residue	N ₂ O
4.D1.5 Cultivation of histosols	N ₂ O
4.D.2 Grassing animals	N ₂ O
4.D3 Atmospheric deposition	N ₂ O
4.D3 Leaching	N ₂ O
4.D1.6 Sewage sludge and Industrial waste used as fertiliser	N ₂ O
4.F Field Burning of Agricultural Residues	N ₂ O

For LULUCF the categorisation used for this KCA is according to Table A1-7. KCA have been estimated both including and excluding LULUCF.

Table A1-7 KCA source categories for LULUCF.

CRF and KCA category	GHG
5.A.1 Forest remaining forest	CO ₂
5.A.2 Land converted to forest	CO ₂
5(II) Forest Land.	N ₂ O
5.B Cropland, Living biomass	CO ₂
5.B Cropland, Dead organic matter	CO ₂
5.B Cropland, Mineral soils	CO ₂
5.B Cropland, Organic soils	CO ₂
5(III) Disturbance, Land converted to cropland	N ₂ O
5.C Grassland, Living biomass	CO ₂
5.C Grassland, Dead organic matter	CO ₂
5.C Grassland, Mineral soils	CO ₂
5.C Grassland, Organic soils	CO ₂
5.D Wetlands, Living biomass	CO ₂
5.D Wetlands, Dead organic matter	CO ₂
5.D Wetlands, Soils	CO ₂
5(II) Wetlands	N ₂ O
5.E Settlements, Living biomass	CO ₂
5.E Settlements, Dead organic matter	CO ₂
5.E Settlements, Soils	CO ₂
5(IV) Cropland Limestone	CO ₂
5(V) Biomass Burning	CH ₄
5(V) Biomass Burning	N ₂ O

KCA categories for the waste sector are shown in Table A1-8.

Table A1-8 KCA source categories for the waste sector.

CRF and KCA category	GHG
6 A. Solid Waste Disposal on Land	CH ₄
6 B. Wastewater Handling	CH ₄
6 B. Wastewater Handling - Direct	N ₂ O
6 B. Wastewater Handling - Indirect	N ₂ O
6.D Accidental fires, buildings	CO ₂
6.D Accidental fires, vehicles	CO ₂
6.C Incineration of corpses	CH ₄
6.C Incineration of carcasses	CH ₄
6.D Compost production	CH ₄
6.D Accidental fires, buildings	CH ₄
6.D Accidental fires, vehicles	CH ₄
6.C Incineration of corpses	N ₂ O
6.C Incineration of carcasses	N ₂ O
6.D Compost production	N ₂ O

The choice of categories identifies 131 categories for the analysis excluding LULUCF and 153 categories for the analysis including LULUCF.

The result of the Key Category Analysis for Denmark for the year 1990 and 2012

The entries for the KCA are composed from the databases producing the CRF inventory and from CRFs.

An overview of results of the KCA excluding LULUCF are shown in Table A1-9 and results of the KCA including LULUCF in Table A1-10. The number of key source categories for each of the KCA are shown in Table A1-11. The 12 different KCA for Denmark point out 25-35 key source categories. Ten source categories are key in all 12 KCA:

- Stationary combustion, Coal, CO₂
- Transport, Road transport, CO₂
- Transport, Agriculture, CO₂
- Transport, Transport industry (mobile), CO₂
- 4A Enteric Fermentation, CH₄
- 4B Manure Management, CH₄
- 4D1.1 Synthetic Fertilizer, N₂O
- 4D1.2 Animal waste applied to soils, N₂O
- 4D3 Leaching, N₂O
- 6A Solid waste disposal on land, CH₄

The 12 different KCA point out a total of 52 different key source categories (out of the total 153 source categories), see Table A1-9 and Table A1-10.

The tier 1 approach point out mainly the large emission sources as key categories and thus CO₂ emission from stationary and mobile combustion are important key categories. The tier 2 approach point out some of the sources with larger uncertainty rates.

The list below gives an overview of the different KCA for Denmark (not including Greenland) that are presented in Table A1-12 – Table A1-23.

Table A1-12 KCA for Denmark, level assessment base year excl. LULUCF, tier 1.

Table A1-13 KCA for Denmark, level assessment base year incl. LULUCF, tier 1.

Table A1-14 KCA for Denmark, level assessment 2012 excl. LULUCF, tier 1.

Table A1-15 KCA for Denmark, level assessment 2012 incl. LULUCF, tier 1.

Table A1-16 KCA for Denmark, trend assessment 1990-2012 excl. LULUCF, tier 1.

Table A1-17 KCA for Denmark, trend assessment 1990-2012 incl. LULUCF, tier 1.

Table A1-18 KCA for Denmark, level assessment base year excl. LULUCF, tier 2.

Table A1-19 KCA for Denmark, level assessment base year incl. LULUCF, tier 2.

Table A1-20 KCA for Denmark, level assessment 2012 excl. LULUCF, tier 2.

Table A1-21 KCA for Denmark, level assessment 2012 incl. LULUCF, tier 2.

Table A1-22 KCA for Denmark, trend assessment 1990-2012 excl. LULUCF, tier 2.

Table A1-23 KCA for Denmark, trend assessment 1990-2012 incl. LULUCF, tier 2.

Table A1-9 Summary of KCA for Denmark, level and trend for 1990-2012, excl. LULUCF, tier 1 and tier 2.

IPCC Source Categories (LULUCF excluded)			Key categories with number according to ranking in analysis					
			Identification criteria					
GHG			Level Tier 1	Level Tier 1	Trend Tier 1	Level Tier 2	Level Tier 2	Trend Tier 2
			1990	2012	1990-2012	1990	2012	1990-2012
Energy	Stationary Combustion, Coal	CO ₂	1	2	1	19	26	19
Energy	Stationary Combustion, BKB	CO ₂						
Energy	Stationary Combustion, Coke	CO ₂						
Energy	Stationary Combustion, Fossil waste	CO ₂	20	6	6		20	14
Energy	Stationary Combustion, Petroleum coke	CO ₂	22	17	17			
Energy	Stationary Combustion, Residual oil	CO ₂	6	18	5			
Energy	Stationary Combustion, Gas oil	CO ₂	3	14	4	20		12
Energy	Stationary Combustion, Kerosene	CO ₂	23		19			
Energy	Stationary Combustion, LPG	CO ₂						
Energy	Stationary Combustion, Refinery gas	CO ₂	16	12	18			
Energy	Stationary Combustion, Natural gas	CO ₂	4	3	2			26
Energy	Stationary Combustion, SOLID	CH ₄						
Energy	Stationary Combustion, LIQUID	CH ₄						
Energy	Stationary Combustion, GAS	CH ₄						
Energy	Natural gas fuelled engines, GAS	CH ₄			23			
Energy	Stationary Combustion, WASTE	CH ₄						
Energy	Stationary Combustion, BIOMASS	CH ₄					22	24
Energy	Biogas fuelled engines, BIOMASS	CH ₄						
Energy	Stationary Combustion, SOLID	N ₂ O				17	25	17
Energy	Stationary Combustion, LIQUID	N ₂ O				9	24	8
Energy	Stationary Combustion, GAS	N ₂ O				24	18	18
Energy	Stationary Combustion, WASTE	N ₂ O						29
Energy	Stationary Combustion, BIOMASS	N ₂ O				10	4	2
Energy	Transport, Road transport	CO ₂	2	1	3	8	6	7
Energy	Transport, Military	CO ₂						
Energy	Transport, Railways	CO ₂		25				
Energy	Transport, Navigation (small boats)	CO ₂						
Energy	Transport, Navigation (large vessels)	CO ₂	17	20	21			
Energy	Transport, Fisheries	CO ₂	19	19				
Energy	Transport, Agriculture	CO ₂	10	7	14	14	11	15
Energy	Transport, Forestry	CO ₂						
Energy	Transport, Industry (mobile)	CO ₂	15	11	13	12	8	10
Energy	Transport, Residential	CO ₂						
Energy	Transport, Commercial/institutional	CO ₂			24			
Energy	Transport, Civil aviation	CO ₂						
Energy	Transport, Road transport	CH ₄						
Energy	Transport, Military	CH ₄						

IPCC Source Categories (LULUCF excluded)			Key categories with number according to ranking in analysis					
		GHG	Identification criteria					
			Level Tier 1 1990	Level Tier 1 2012	Trend Tier 1 1990-2012	Level Tier 2 1990	Level Tier 2 2012	Trend Tier 2 1990-2012
Energy	Transport, Railways	CH ₄						
Energy	Transport, Navigation (small boats)	CH ₄						
Energy	Transport, Navigation (large vessels)	CH ₄						
Energy	Transport, Fisheries	CH ₄						
Energy	Transport, Agriculture	CH ₄						
Energy	Transport, Forestry	CH ₄						
Energy	Transport, Industry (mobile)	CH ₄						
Energy	Transport, Residential	CH ₄						
Energy	Transport, Commercial/institutional	CH ₄						
Energy	Transport, Civil aviation	CH ₄						
Energy	Transport, Road transport	N ₂ O						
Energy	Transport, Military	N ₂ O						
Energy	Transport, Railways	N ₂ O						
Energy	Transport, Navigation (small boats)	N ₂ O						
Energy	Transport, Navigation (large vessels)	N ₂ O				23		
Energy	Transport, Fisheries	N ₂ O					28	
Energy	Transport, Agriculture	N ₂ O				22	19	22
Energy	Transport, Forestry	N ₂ O						
Energy	Transport, Industry (mobile)	N ₂ O					23	28
Energy	Transport, Residential	N ₂ O						
Energy	Transport, Commercial/institutional	N ₂ O						
Energy	Transport, Civil aviation	N ₂ O						
Energy	1.B.2 Flaring in refinery	CO ₂						
Energy	1.B.2 Flaring off-shore	CO ₂	26	28				
Energy	1.B.2 Land based activities	CO ₂						
Energy	1.B.2 Off-shore activities	CO ₂						
Energy	1.B.2 Transmission of natural gas	CO ₂						
Energy	1.B.2 Distribution of natural gas	CO ₂						
Energy	1.B.2 Venting in gas storage	CO ₂						
Energy	1.B.2. Flaring in refinery	CH ₄						
Energy	1.B.2. Flaring off-shore	CH ₄						
Energy	1.B.2 Refinery processes	CH ₄						25
Energy	1.B.2 Land based activities	CH ₄						
Energy	1.B.2 Off-shore activities	CH ₄						
Energy	1.B.2 Transmission of natural gas	CH ₄						
Energy	1.B.2 Distribution of natural gas	CH ₄						
Energy	1.B.2 Venting in gas storage	CH ₄						
Energy	1.B.2 Flaring in refinery	N ₂ O						

IPCC Source Categories (LULUCF excluded)			Key categories with number according to ranking in analysis					
			Identification criteria					
			Level Tier 1	Level Tier 1	Trend Tier 1	Level Tier 2	Level Tier 2	Trend Tier 2
			1990	2012	1990-2012	1990	2012	1990-2012
Energy	1.B.2 Flaring off-shore	N ₂ O						
Industrial Proc.	2A1 Cement production	CO ₂	14	13	20			
Industrial Proc.	2A2 Lime production	CO ₂						
Industrial Proc.	2A3 Limestone and dolomite use	CO ₂						
Industrial Proc.	2A5 Asphalt roofing	CO ₂						
Industrial Proc.	2A6 Road paving with asphalt	CO ₂						
Industrial Proc.	2A7a Glass and Glass wool	CO ₂						
Industrial Proc.	2A7b Yellow bricks	CO ₂						
Industrial Proc.	2A7c Expanded clay	CO ₂						
Industrial Proc.	2B5 Catalysts/Fertilizers, Pesticides and Sulphuric acid	CO ₂						
Industrial Proc.	2C1 Iron and steel production	CO ₂						
Industrial Proc.	2D2 Food and Drink	CO ₂						
Industrial Proc.	2G Lubricants	CO ₂						
Industrial Proc.	2B2 Nitric acid production	N ₂ O	12		7	18		9
Industrial Proc.	2F Consumption of HFC	HFC		16	10		10	6
Industrial Proc.	2F Consumption of PFC	PFC						
Industrial Proc.	2F Consumption of SF ₆	SF ₆						
Solvent and Other Prod. Use	3A Paint application	CO ₂						
Solvent and Other Prod. Use	3B Degreasing and dry cleaning	CO ₂						
Solvent and Other Prod. Use	3C Chemical products, manufacturing and processing	CO ₂						
Solvent and Other Prod. Use	3D5 Other	CO ₂						
Solvent and Other Prod. Use	3D5 Consumption of fireworks	CO ₂						
Solvent and Other Prod. Use	3D5 Use of candles	CO ₂						
Solvent and Other Prod. Use	3D1 Other - Use of N ₂ O for Anaesthesia	N ₂ O						
Solvent and Other Prod. Use	3D5 Use of tobacco	N ₂ O						
Solvent and Other Prod. Use	3D5 Use of charcoal for BBQ	N ₂ O						
Solvent and Other Prod. Use	3D5 Consumption of fireworks	N ₂ O						
Solvent and Other Prod. Use	3D5 Use of candles	N ₂ O						
Agriculture	4A Enteric Fermentation	CH ₄	5	4	11	5	7	16
Agriculture	4B Manure Management	CH ₄	13	8	9	21	14	13
Agriculture	4F Field burning of agricultural residues	CH ₄						
Agriculture	4.B Manure Management	N ₂ O	18	21		6	9	23
Agriculture	4.D1.1 Synthetic Fertilizer	N ₂ O	8	10	8	2	3	1
Agriculture	4.D1.2 Animal waste applied to soils	N ₂ O	11	9	15	4	2	5
Agriculture	4.D1.3 N-fixing crops	N ₂ O		24		16	15	27
Agriculture	4.D1.4 Crop Residue	N ₂ O	24	22		11	12	
Agriculture	4.D1.5 Cultivation of histosols	N ₂ O		27		15	17	
Agriculture	4.D.2 Grassing animals	N ₂ O	25	26		13	16	

IPCC Source Categories (LULUCF excluded)			Key categories with number according to ranking in analysis					
			Identification criteria					
			Level Tier 1	Level Tier 1	Trend Tier 1	Level Tier 2	Level Tier 2	Trend Tier 2
			1990	2012	1990-2012	1990	2012	1990-2012
Agriculture	4.D3 Atmospheric deposition	N ₂ O	21	23		7	13	20
Agriculture	4.D3 Leaching	N ₂ O	7	5	12	1	1	3
Agriculture	4.D1.6 Sewage sludge and Industrial waste used as fertiliser	N ₂ O						
Agriculture	4.F Field Burning of Agricultural Residues	N ₂ O						
Waste	6 A. Solid Waste Disposal on Land	CH ₄	9	15	16	3	5	4
Waste	6 B. Wastewater Handling	CH ₄						
Waste	6 B. Wastewater Handling - Direct	N ₂ O						
Waste	6 B. Wastewater Handling - Indirect	N ₂ O						
Waste	6.D Accidental fires, buildings	CO ₂						
Waste	6.D Accidental fires, vehicles	CO ₂						
Waste	6.C Incineration of corpses	CH ₄						
Waste	6.C Incineration of carcasses	CH ₄						
Waste	6.D Compost production	CH ₄					27	21
Waste	6.D Accidental fires, buildings	CH ₄						
Waste	6.D Accidental fires, vehicles	CH ₄						
Waste	6.C Incineration of corpses	N ₂ O						
Waste	6.C Incineration of carcasses	N ₂ O						
Waste	6.D Compost production	N ₂ O			22		21	11

Table A1-10 Summary of KCA for Denmark, level and trend for 1990-2012, incl. LULUCF, tier 1 and tier 2.

IPCC Source Categories (LULUCF included)			Key categories with number according to ranking in analysis					
			Identification criteria					
GHG			Level Tier 1	Level Tier 1	Trend Tier 1	Level Tier 2	Level Tier 2	Trend Tier 2
			1990	2012	1990-2012	1990	2012	1990-2012
Energy	Stationary Combustion, Coal	CO ₂	1	2	1	22	31	27
Energy	Stationary Combustion, BKB	CO ₂						
Energy	Stationary Combustion, Coke	CO ₂						
Energy	Stationary Combustion, Fossil waste	CO ₂	23	8	7		25	20
Energy	Stationary Combustion, Petroleum coke	CO ₂	25	19	19			
Energy	Stationary Combustion, Residual oil	CO ₂	7	21	6			
Energy	Stationary Combustion, Gas oil	CO ₂	3	16	5	23		21
Energy	Stationary Combustion, Kerosene	CO ₂	26		22			
Energy	Stationary Combustion, LPG	CO ₂						
Energy	Stationary Combustion, Refinery gas	CO ₂	18	14	18			
Energy	Stationary Combustion, Natural gas	CO ₂	4	3	2			31
Energy	Stationary Combustion, SOLID	CH ₄						
Energy	Stationary Combustion, LIQUID	CH ₄						
Energy	Stationary Combustion, GAS	CH ₄						
Energy	Natural gas fuelled engines, GAS	CH ₄			28			
Energy	Stationary Combustion, WASTE	CH ₄						
Energy	Stationary Combustion, BIOMASS	CH ₄					27	28
Energy	Biogas fuelled engines, BIOMASS	CH ₄						
Energy	Stationary Combustion, SOLID	N ₂ O				20	30	23
Energy	Stationary Combustion, LIQUID	N ₂ O				11	29	12
Energy	Stationary Combustion, GAS	N ₂ O					23	22
Energy	Stationary Combustion, WASTE	N ₂ O						
Energy	Stationary Combustion, BIOMASS	N ₂ O				12	5	2
Energy	Transport, Road transport	CO ₂	2	1	3	10	8	9
Energy	Transport, Military	CO ₂						
Energy	Transport, Railways	CO ₂	30	29				
Energy	Transport, Navigation (small boats)	CO ₂						
Energy	Transport, Navigation (large vessels)	CO ₂	19	24				
Energy	Transport, Fisheries	CO ₂	22	22				
Energy	Transport, Agriculture	CO ₂	12	9	13	17	15	19
Energy	Transport, Forestry	CO ₂						

IPCC Source Categories (LULUCF included)			Key categories with number according to ranking in analysis					
			Identification criteria					
			Level Tier 1	Level Tier 1	Trend Tier 1	Level Tier 2	Level Tier 2	Trend Tier 2
			1990	2012	1990-2012	1990	2012	1990-2012
Energy	Transport, Industry (mobile)	CO ₂	17	13	14	14	11	13
Energy	Transport, Residential	CO ₂						
Energy	Transport, Commercial/institutional	CO ₂			26			
Energy	Transport, Civil aviation	CO ₂						
Energy	Transport, Road transport	CH ₄						
Energy	Transport, Military	CH ₄						
Energy	Transport, Railways	CH ₄						
Energy	Transport, Navigation (small boats)	CH ₄						
Energy	Transport, Navigation (large vessels)	CH ₄						
Energy	Transport, Fisheries	CH ₄						
Energy	Transport, Agriculture	CH ₄						
Energy	Transport, Forestry	CH ₄						
Energy	Transport, Industry (mobile)	CH ₄						
Energy	Transport, Residential	CH ₄						
Energy	Transport, Commercial/institutional	CH ₄						
Energy	Transport, Civil aviation	CH ₄						
Energy	Transport, Road transport	N ₂ O						
Energy	Transport, Military	N ₂ O						
Energy	Transport, Railways	N ₂ O						
Energy	Transport, Navigation (small boats)	N ₂ O						
Energy	Transport, Navigation (large vessels)	N ₂ O				26		
Energy	Transport, Fisheries	N ₂ O						
Energy	Transport, Agriculture	N ₂ O				25	24	25
Energy	Transport, Forestry	N ₂ O						
Energy	Transport, Industry (mobile)	N ₂ O					28	30
Energy	Transport, Residential	N ₂ O						
Energy	Transport, Commercial/institutional	N ₂ O						
Energy	Transport, Civil aviation	N ₂ O						
Energy	1.B.2 Flaring in refinery	CO ₂						
Energy	1.B.2 Flaring off-shore	CO ₂	29	32				
Energy	1.B.2 Land based activities	CO ₂						
Energy	1.B.2 Off-shore activities	CO ₂						

IPCC Source Categories (LULUCF included)			Key categories with number according to ranking in analysis					
		GHG	Identification criteria					
			Level Tier 1 1990	Level Tier 1 2012	Trend Tier 1 1990-2012	Level Tier 2 1990	Level Tier 2 2012	Trend Tier 2 1990-2012
Energy	1.B.2 Transmission of natural gas	CO ₂						
Energy	1.B.2 Distribution of natural gas	CO ₂						
Energy	1.B.2 Venting in gas storage	CO ₂						
Energy	1.B.2. Flaring in refinery	CH ₄						
Energy	1.B.2. Flaring off-shore	CH ₄						
Energy	1.B.2 Refinery processes	CH ₄						32
Energy	1.B.2 Land based activities	CH ₄						
Energy	1.B.2 Off-shore activities	CH ₄						
Energy	1.B.2 Transmission of natural gas	CH ₄						
Energy	1.B.2 Distribution of natural gas	CH ₄						
Energy	1.B.2 Venting in gas storage	CH ₄						
Energy	1.B.2 Flaring in refinery	N ₂ O						
Energy	1.B.2 Flaring off-shore	N ₂ O						
Industrial Proc.	2A1 Cement production	CO ₂	16	15	21			
Industrial Proc.	2A2 Lime production	CO ₂						
Industrial Proc.	2A3 Limestone and dolomite use	CO ₂						
Industrial Proc.	2A5 Asphalt roofing	CO ₂						
Industrial Proc.	2A6 Road paving with asphalt	CO ₂						
Industrial Proc.	2A7a Glass and Glass wool	CO ₂						
Industrial Proc.	2A7b Yellow bricks	CO ₂						
Industrial Proc.	2A7c Expanded clay	CO ₂						
Industrial Proc.	2B5 Catalysts/Fertilizers, Pesticides and Sulphuric acid	CO ₂						
Industrial Proc.	2C1 Iron and steel production	CO ₂						
Industrial Proc.	2D2 Food and Drink	CO ₂						
Industrial Proc.	2G Lubricants	CO ₂						
Industrial Proc.	2B2 Nitric acid production	N ₂ O	14		8	21		14
Industrial Proc.	2F Consumption of HFC	HFC		18	11		14	10
Industrial Proc.	2F Consumption of PFC	PFC						
Industrial Proc.	2F Consumption of SF ₆	SF ₆						
Solvent and Other Prod. Use	3A Paint application	CO ₂						
Solvent and Other Prod. Use	3B Degreasing and dry cleaning	CO ₂						
Solvent and Other Prod. Use	3C Chemical products, manufacturing and processing	CO ₂						

IPCC Source Categories (LULUCF included)			Key categories with number according to ranking in analysis						
			GHG	Identification criteria					
				Level Tier 1 1990	Level Tier 1 2012	Trend Tier 1 1990-2012	Level Tier 2 1990	Level Tier 2 2012	Trend Tier 2 1990-2012
Solvent and Other Prod. Use	3D5 Other	CO ₂							
Solvent and Other Prod. Use	3D5 Consumption of fireworks	CO ₂							
Solvent and Other Prod. Use	3D5 Use of candles	CO ₂							
Solvent and Other Prod. Use	3D1 Other - Use of N2O for Anaesthesia	N ₂ O							
Solvent and Other Prod. Use	3D5 Use of tobacco	N ₂ O							
Solvent and Other Prod. Use	3D5 Use of charcoal for BBQ	N ₂ O							
Solvent and Other Prod. Use	3D5 Consumption of fireworks	N ₂ O							
Solvent and Other Prod. Use	3D5 Use of candles	N ₂ O							
Agriculture	4A Enteric Fermentation	CH ₄	5	5	9	7	9	15	
Agriculture	4B Manure Management	CH ₄	15	10	10	24	18	17	
Agriculture	4F Field burning of agricultural residues	CH ₄							
Agriculture	4.B Manure Management	N ₂ O	21	25		8	12		
Agriculture	4.D1.1 Synthetic Fertilizer	N ₂ O	9	12	12	3	4	3	
Agriculture	4.D1.2 Animal waste applied to soils	N ₂ O	13	11	16	5	3	4	
Agriculture	4.D1.3 N-fixing crops	N ₂ O		28		19	19	26	
Agriculture	4.D1.4 Crop Residue	N ₂ O	27	26		13	16	29	
Agriculture	4.D1.5 Cultivation of histosols	N ₂ O		31		18	22		
Agriculture	4.D.2 Grassing animals	N ₂ O	28	30		15	21		
Agriculture	4.D3 Atmospheric deposition	N ₂ O	24	27		9	17	33	
Agriculture	4.D3 Leaching	N ₂ O	8	7	23	2	2	8	
Agriculture	4.D1.6 Sewage sludge and Industrial waste used as fertiliser	N ₂ O							
Agriculture	4.F Field Burning of Agricultural Residues	N ₂ O							
LULUCF	5.A.1 Forest remaining forest	CO ₂		4	4		7	1	
LULUCF	5.A.2 Land converted to forest	CO ₂							
LULUCF	5(II) Forest Land.	N ₂ O							
LULUCF	5.B Cropland, Living biomass	CO ₂		33	20		13	5	
LULUCF	5.B Cropland, Dead organic matter	CO ₂							
LULUCF	5.B Cropland, Mineral soils	CO ₂	10	20	17	6	10	6	
LULUCF	5.B Cropland, Organic soils	CO ₂	6	6		1	1		
LULUCF	5(III) Disturbance, Land converted to cropland	N ₂ O							
LULUCF	5.C Grassland, Living biomass	CO ₂		23	15		20	11	

IPCC Source Categories (LULUCF included)		GHG	Key categories with number according to ranking in analysis					
			Identification criteria					
			Level Tier 1 1990	Level Tier 1 2012	Trend Tier 1 1990-2012	Level Tier 2 1990	Level Tier 2 2012	Trend Tier 2 1990-2012
LULUCF	5.C Grassland, Dead organic matter	CO ₂						
LULUCF	5.C Grassland, Mineral soils	CO ₂						
LULUCF	5.C Grassland, Organic soils	CO ₂						
LULUCF	5.D Wetlands, Living biomass	CO ₂						
LULUCF	5.D Wetlands, Dead organic matter	CO ₂						
LULUCF	5.D Wetlands, Soils	CO ₂						34
LULUCF	5(II) Wetlands	N ₂ O						
LULUCF	5.E Settlements, Living biomass	CO ₂						
LULUCF	5.E Settlements, Dead organic matter	CO ₂						
LULUCF	5.E Settlements, Soils	CO ₂						
LULUCF	5(IV) Cropland Limestone	CO ₂	20		25	16		18
LULUCF	5(V) Biomass Burning	CH ₄						
LULUCF	5(V) Biomass Burning	N ₂ O						
Waste	6 A. Solid Waste Disposal on Land	CH ₄	11	17	24	4	6	7
Waste	6 B. Wastewater Handling	CH ₄						
Waste	6 B. Wastewater Handling - Direct	N ₂ O						
Waste	6 B. Wastewater Handling - Indirect	N ₂ O						
Waste	6.D Accidental fires, buildings	CO ₂						
Waste	6.D Accidental fires, vehicles	CO ₂						
Waste	6.C Incineration of corpses	CH ₄						
Waste	6.C Incineration of carcasses	CH ₄						
Waste	6.D Compost production	CH ₄					32	24
Waste	6.D Accidental fires, buildings	CH ₄						
Waste	6.D Accidental fires, vehicles	CH ₄						
Waste	6.C Incineration of corpses	N ₂ O						
Waste	6.C Incineration of carcasses	N ₂ O						
Waste	6.D Compost production	N ₂ O			27		26	16

Table A1-11 Summary of KCA for Denmark, number of key source categories in each of the KCA.

	Level Tier 1 1990	Level Tier 1 2012	Trend Tier 1 1990-2012	Level Tier 2 1990	Level Tier 2 2012	Trend Tier 2 1990-2012
Excluding LULUCF	28	30	25	26	30	30
Including LULUCF	32	35	29	28	34	35

Table A1-12 KCA for Denmark, level assessment base year excl. LULUCF, tier 1.

Tier 1 Analysis IPCC Source Categories (LULUCF excluded)		DK GHG	Base Year Estimate Ex,o Mt CO ₂ eq	Base Year Level Assessment Lx,o	Base Year Cumulative Total of Lx,o
Energy	Stationary Combustion, Coal	CO ₂	23.834	0.346	0.346
Energy	Transport, Road transport	CO ₂	9.284	0.135	0.480
Energy	Stationary Combustion, Gas oil	CO ₂	4.547	0.066	0.546
Energy	Stationary Combustion, Natural gas	CO ₂	4.335	0.063	0.609
Agriculture	4A Enteric Fermentation	CH ₄	3.247	0.047	0.656
Energy	Stationary Combustion, Residual oil	CO ₂	2.496	0.036	0.692
Agriculture	4.D3 Leaching	N ₂ O	2.447	0.035	0.728
Agriculture	4.D1.1 Synthetic Fertilizer	N ₂ O	2.354	0.034	0.762
Waste	6 A. Solid Waste Disposal on Land	CH ₄	1.366	0.020	0.782
Energy	Transport, Agriculture	CO ₂	1.272	0.018	0.800
Agriculture	4.D1.2 Animal waste applied to soils	N ₂ O	1.112	0.016	0.817
Industrial Proc.	2B2 Nitric acid production	N ₂ O	1.043	0.015	0.832
Agriculture	4B Manure Management	CH ₄	0.985	0.014	0.846
Industrial Proc.	2A1 Cement production	CO ₂	0.882	0.013	0.859
Energy	Transport, Industry (mobile)	CO ₂	0.839	0.012	0.871
Energy	Stationary Combustion, Refinery gas	CO ₂	0.816	0.012	0.883
Energy	Transport, Navigation (large vessels)	CO ₂	0.748	0.011	0.894
Agriculture	4.B Manure Management	N ₂ O	0.600	0.009	0.902
Energy	Transport, Fisheries	CO ₂	0.591	0.009	0.911
Energy	Stationary Combustion, Fossil waste	CO ₂	0.573	0.008	0.919
Agriculture	4.D3 Atmospheric deposition	N ₂ O	0.496	0.007	0.926
Energy	Stationary Combustion, Petroleum coke	CO ₂	0.415	0.006	0.932
Energy	Stationary Combustion, Kerosene	CO ₂	0.366	0.005	0.938
Agriculture	4.D1.4 Crop Residue	N ₂ O	0.361	0.005	0.943
Agriculture	4.D.2 Grassing animals	N ₂ O	0.334	0.005	0.948
Energy	1.B.2 Flaring off-shore	CO ₂	0.302	0.004	0.952
Energy	Transport, Railways	CO ₂	0.297	0.004	0.956
Agriculture	4.D1.5 Cultivation of histosols	N ₂ O	0.290	0.004	0.961
Agriculture	4.D1.3 N-fixing crops	N ₂ O	0.269	0.004	0.965
Energy	Transport, Civil aviation	CO ₂	0.243	0.004	0.968
Industrial Proc.	2F Consumption of HFC	HFC	0.218	0.003	0.971
Energy	Stationary Combustion, LPG	CO ₂	0.184	0.003	0.974
Energy	Stationary Combustion, Coke	CO ₂	0.138	0.002	0.976
Energy	Transport, Military	CO ₂	0.119	0.002	0.978
Industrial Proc.	2A2 Lime production	CO ₂	0.116	0.002	0.979
Industrial Proc.	2F Consumption of SF ₆	SF ₆	0.107	0.002	0.981
Energy	Stationary Combustion, BIOMASS	CH ₄	0.102	0.001	0.982
Energy	Transport, Road transport	N ₂ O	0.091	0.001	0.984
Waste	6 B. Wastewater Handling - Indirect	N ₂ O	0.082	0.001	0.985
Energy	Transport, Commercial/institutional	CO ₂	0.074	0.001	0.986
Energy	Stationary Combustion, SOLID	N ₂ O	0.068	0.001	0.987
Waste	6 B. Wastewater Handling	CH ₄	0.065	0.001	0.988
Solvent and Other Prod. Use	3D5 Other	CO ₂	0.061	0.001	0.989
Industrial Proc.	2G Lubricants	CO ₂	0.050	0.001	0.989
Energy	Transport, Navigation (small boats)	CO ₂	0.048	0.001	0.990
Energy	Transport, Road transport	CH ₄	0.047	0.001	0.991
Energy	Stationary Combustion, LIQUID	N ₂ O	0.044	0.001	0.991
Energy	Transport, Residential	CO ₂	0.039	0.001	0.992
Energy	Stationary Combustion, BIOMASS	N ₂ O	0.038	0.001	0.993
Energy	Transport, Forestry	CO ₂	0.036	0.001	0.993
Waste	6.D Compost production	CH ₄	0.029	0.000	0.994
Industrial Proc.	2C1 Iron and steel production	CO ₂	0.028	0.000	0.994
Agriculture	4.D1.6 Sewage sludge and Industrial waste used as fertiliser	N ₂ O	0.028	0.000	0.994

Tier 1 Analysis IPCC Source Categories (LULUCF excluded)		DK GHG	Base Year Estimate	Base Year Level Assessment Lx,o	Base Year Cumulative Total of Lx,o
			Ex,o Mt CO ₂ eq		
Industrial Proc.	2A7b Yellow bricks	CO ₂	0.023	0.000	0.995
Energy	1.B.2 Flaring in refinery	CO ₂	0.023	0.000	0.995
Waste	6 B. Wastewater Handling - Direct	N ₂ O	0.023	0.000	0.995
Solvent and Other Prod. Use	3D5 Use of candles	CO ₂	0.022	0.000	0.996
Solvent and Other Prod. Use	3C Chemical products, manufacturing and processing	CO ₂	0.019	0.000	0.996
Industrial Proc.	2A7a Glass and Glass wool	CO ₂	0.017	0.000	0.996
Energy	1.B.2 Land based activities	CH ₄	0.017	0.000	0.996
Energy	Stationary Combustion, GAS	N ₂ O	0.016	0.000	0.997
Energy	Transport, Agriculture	N ₂ O	0.015	0.000	0.997
Industrial Proc.	2A7c Expanded clay	CO ₂	0.015	0.000	0.997
Energy	1.B.2 Off-shore activities	CH ₄	0.015	0.000	0.997
Energy	Transport, Navigation (large vessels)	N ₂ O	0.015	0.000	0.998
Industrial Proc.	2A3 Limestone and dolomite use	CO ₂	0.014	0.000	0.998
Solvent and Other Prod. Use	3A Paint application	CO ₂	0.013	0.000	0.998
Energy	Stationary Combustion, SOLID	CH ₄	0.013	0.000	0.998
Waste	6.D Compost production	N ₂ O	0.013	0.000	0.998
Energy	Transport, Fisheries	N ₂ O	0.011	0.000	0.998
Waste	6.D Accidental fires, buildings	CO ₂	0.011	0.000	0.999
Energy	Stationary Combustion, BKB	CO ₂	0.011	0.000	0.999
Energy	Transport, Industry (mobile)	N ₂ O	0.011	0.000	0.999
Energy	Stationary Combustion, WASTE	N ₂ O	0.007	0.000	0.999
Waste	6.D Accidental fires, vehicles	CO ₂	0.006	0.000	0.999
Energy	1.B.2 Distribution of natural gas	CH ₄	0.005	0.000	0.999
Energy	Natural gas fuelled engines, GAS	CH ₄	0.005	0.000	0.999
Industrial Proc.	2D2 Food and Drink	CO ₂	0.004	0.000	0.999
Energy	1.B.2 Transmission of natural gas	CH ₄	0.004	0.000	0.999
Energy	Transport, Civil aviation	N ₂ O	0.003	0.000	0.999
Energy	Stationary Combustion, GAS	CH ₄	0.003	0.000	1.000
Energy	Stationary Combustion, LIQUID	CH ₄	0.003	0.000	1.000
Energy	Transport, Railways	N ₂ O	0.003	0.000	1.000
Energy	1.B.2 Off-shore activities	CO ₂	0.002	0.000	1.000
Energy	Transport, Agriculture	CH ₄	0.002	0.000	1.000
Energy	Transport, Commercial/institutional	CH ₄	0.002	0.000	1.000
Agriculture	4F Field burning of agricultural residues	CH ₄	0.002	0.000	1.000
Industrial Proc.	2A6 Road paving with asphalt	CO ₂	0.002	0.000	1.000
Energy	Biogas fuelled engines, BIOMASS	CH ₄	0.001	0.000	1.000
Waste	6.D Accidental fires, buildings	CH ₄	0.001	0.000	1.000
Energy	Transport, Industry (mobile)	CH ₄	0.001	0.000	1.000
Energy	1.B.2 Venting in gas storage	CH ₄	0.001	0.000	1.000
Energy	Transport, Military	N ₂ O	0.001	0.000	1.000
Energy	Transport, Residential	CH ₄	0.001	0.000	1.000
Industrial Proc.	2B5 Catalysts/Fertilizers, Pesticides and Sulphuric acid	CO ₂	0.001	0.000	1.000
Energy	1.B.2 Refinery processes	CH ₄	0.001	0.000	1.000
Energy	Stationary Combustion, WASTE	CH ₄	0.001	0.000	1.000
Solvent and Other Prod. Use	3D5 Consumption of fireworks	N ₂ O	0.001	0.000	1.000
Energy	1.B.2. Flaring off-shore	CH ₄	0.001	0.000	1.000
Energy	1.B.2 Flaring off-shore	N ₂ O	0.001	0.000	1.000
Agriculture	4.F Field Burning of Agricultural Residues	N ₂ O	0.001	0.000	1.000
Industrial Proc.	2F Consumption of PFC	PFC	0.001	0.000	1.000
Energy	Transport, Forestry	CH ₄	0.000	0.000	1.000
Energy	Transport, Navigation (small boats)	N ₂ O	0.000	0.000	1.000
Energy	Transport, Navigation (small boats)	CH ₄	0.000	0.000	1.000
Energy	Transport, Commercial/institutional	N ₂ O	0.000	0.000	1.000
Energy	Transport, Navigation (large vessels)	CH ₄	0.000	0.000	1.000
Energy	Transport, Fisheries	CH ₄	0.000	0.000	1.000
Waste	6.D Accidental fires, vehicles	CH ₄	0.000	0.000	1.000
Energy	Transport, Railways	CH ₄	0.000	0.000	1.000
Solvent and Other Prod. Use	3D5 Use of tobacco	N ₂ O	0.000	0.000	1.000
Energy	Transport, Residential	N ₂ O	0.000	0.000	1.000
Waste	6.C Incineration of corpses	N ₂ O	0.000	0.000	1.000
Energy	Transport, Forestry	N ₂ O	0.000	0.000	1.000
Energy	1.B.2. Flaring in refinery	CH ₄	0.000	0.000	1.000

Tier 1 Analysis IPCC Source Categories (LULUCF excluded)		DK GHG	Base Year Estimate	Base Year Level Assessment	Base Year Cumulative Total of Lx,o
			Ex,o Mt CO ₂ eq	Lx,o	
Energy	Transport, Military	CH ₄	0.000	0.000	1.000
Energy	Transport, Civil aviation	CH ₄	0.000	0.000	1.000
Solvent and Other Prod. Use	3D5 Use of charcoal for BBQ	N ₂ O	0.000	0.000	1.000
Energy	1.B.2 Flaring in refinery	N ₂ O	0.000	0.000	1.000
Solvent and Other Prod. Use	3D5 Use of candles	N ₂ O	0.000	0.000	1.000
Solvent and Other Prod. Use	3D5 Consumption of fireworks	CO ₂	0.000	0.000	1.000
Industrial Proc.	2A5 Asphalt roofing	CO ₂	0.000	0.000	1.000
Waste	6.C Incineration of carcasses	N ₂ O	0.000	0.000	1.000
Waste	6.C Incineration of corpses	CH ₄	0.000	0.000	1.000
Energy	1.B.2 Distribution of natural gas	CO ₂	0.000	0.000	1.000
Energy	1.B.2 Transmission of natural gas	CO ₂	0.000	0.000	1.000
Energy	1.B.2 Land based activities	CO ₂	0.000	0.000	1.000
Energy	1.B.2 Venting in gas storage	CO ₂	0.000	0.000	1.000
Waste	6.C Incineration of carcasses	CH ₄	0.000	0.000	1.000
Solvent and Other Prod. Use	3B Degreasing and dry cleaning	CO ₂	0.000	0.000	1.000
Solvent and Other Prod. Use	3D1 Other - Use of N2O for Anaesthesia	N ₂ O	0.000	0.000	1.000
Total			68.943	1.000	

Table A1-13 KCA for Denmark, level assessment base year incl. LULUCF, tier 1.

Tier 1 Analysis		DK			
IPCC Source Categories (LULUCF included)		GHG	Base Year Estimate	Base Year Level Assessment	Base Year Cumulative Total of Lx,o
			Ex,o	Lx,o	
			Mt CO ₂ eq		
Energy	Stationary Combustion, Coal	CO ₂	23.834	0.320	0.320
Energy	Transport, Road transport	CO ₂	9.284	0.125	0.445
Energy	Stationary Combustion, Gas oil	CO ₂	4.547	0.061	0.506
Energy	Stationary Combustion, Natural gas	CO ₂	4.335	0.058	0.565
Agriculture	4A Enteric Fermentation	CH ₄	3.247	0.044	0.608
LULUCF	5.B Cropland, Organic soils	CO ₂	2.887	0.039	0.647
Energy	Stationary Combustion, Residual oil	CO ₂	2.496	0.034	0.681
Agriculture	4.D3 Leaching	N ₂ O	2.447	0.033	0.714
Agriculture	4.D1.1 Synthetic Fertilizer	N ₂ O	2.354	0.032	0.745
LULUCF	5.B Cropland, Mineral soils	CO ₂	1.415	0.019	0.764
Waste	6 A. Solid Waste Disposal on Land	CH ₄	1.366	0.018	0.783
Energy	Transport, Agriculture	CO ₂	1.272	0.017	0.800
Agriculture	4.D1.2 Animal waste applied to soils	N ₂ O	1.112	0.015	0.815
Industrial Proc.	2B2 Nitric acid production	N ₂ O	1.043	0.014	0.829
Agriculture	4B Manure Management	CH ₄	0.985	0.013	0.842
Industrial Proc.	2A1 Cement production	CO ₂	0.882	0.012	0.854
Energy	Transport, Industry (mobile)	CO ₂	0.839	0.011	0.865
Energy	Stationary Combustion, Refinery gas	CO ₂	0.816	0.011	0.876
Energy	Transport, Navigation (large vessels)	CO ₂	0.748	0.010	0.886
LULUCF	5(IV) Cropland Limestone	CO ₂	0.623	0.008	0.895
Agriculture	4.B Manure Management	N ₂ O	0.600	0.008	0.903
Energy	Transport, Fisheries	CO ₂	0.591	0.008	0.911
Energy	Stationary Combustion, Fossil waste	CO ₂	0.573	0.008	0.918
Agriculture	4.D3 Atmospheric deposition	N ₂ O	0.496	0.007	0.925
Energy	Stationary Combustion, Petroleum coke	CO ₂	0.415	0.006	0.931
Energy	Stationary Combustion, Kerosene	CO ₂	0.366	0.005	0.935
Agriculture	4.D1.4 Crop Residue	N ₂ O	0.361	0.005	0.940
Agriculture	4.D.2 Grassing animals	N ₂ O	0.334	0.004	0.945
Energy	1.B.2 Flaring off-shore	CO ₂	0.302	0.004	0.949
Energy	Transport, Railways	CO ₂	0.297	0.004	0.953
Agriculture	4.D1.5 Cultivation of histosols	N ₂ O	0.290	0.004	0.957
Agriculture	4.D1.3 N-fixing crops	N ₂ O	0.269	0.004	0.960
Energy	Transport, Civil aviation	CO ₂	0.243	0.003	0.964
Industrial Proc.	2F Consumption of HFC	HFC	0.218	0.003	0.967
Energy	Stationary Combustion, LPG	CO ₂	0.184	0.002	0.969
Energy	Stationary Combustion, Coke	CO ₂	0.138	0.002	0.971
Energy	Transport, Military	CO ₂	0.119	0.002	0.973
Industrial Proc.	2A2 Lime production	CO ₂	0.116	0.002	0.974
Industrial Proc.	2F Consumption of SF ₆	SF ₆	0.107	0.001	0.976
LULUCF	5.C Grassland, Organic soils	CO ₂	0.107	0.001	0.977
Energy	Stationary Combustion, BIOMASS	CH ₄	0.102	0.001	0.978
Energy	Transport, Road transport	N ₂ O	0.091	0.001	0.980
LULUCF	5.D Wetlands, Soils	CO ₂	0.085	0.001	0.981
Waste	6 B. Wastewater Handling - Indirect	N ₂ O	0.082	0.001	0.982
LULUCF	5.A.2 Land converted to forest	CO ₂	0.077	0.001	0.983
LULUCF	5.C Grassland, Living biomass	CO ₂	0.075	0.001	0.984
Energy	Transport, Commercial/institutional	CO ₂	0.074	0.001	0.985
LULUCF	5.B Cropland, Living biomass	CO ₂	0.074	0.001	0.986
Energy	Stationary Combustion, SOLID	N ₂ O	0.068	0.001	0.987
Waste	6 B. Wastewater Handling	CH ₄	0.065	0.001	0.988
Solvent and Other Prod. Use	3D5 Other	CO ₂	0.061	0.001	0.988
Industrial Proc.	2G Lubricants	CO ₂	0.050	0.001	0.989
LULUCF	5.A.1 Forest remaining forest	CO ₂	0.050	0.001	0.990
Energy	Transport, Navigation (small boats)	CO ₂	0.048	0.001	0.990
Energy	Transport, Road transport	CH ₄	0.047	0.001	0.991
Energy	Stationary Combustion, LIQUID	N ₂ O	0.044	0.001	0.992
Energy	Transport, Residential	CO ₂	0.039	0.001	0.992
Energy	Stationary Combustion, BIOMASS	N ₂ O	0.038	0.001	0.993
Energy	Transport, Forestry	CO ₂	0.036	0.000	0.993
Waste	6.D Compost production	CH ₄	0.029	0.000	0.994
Industrial Proc.	2C1 Iron and steel production	CO ₂	0.028	0.000	0.994

Tier 1 Analysis		DK			
IPCC Source Categories (LULUCF included)		GHG	Base Year Estimate	Base Year Level Assessment	Base Year Cumulative Total of Lx,o
			Ex,o Mt CO ₂ eq	Lx,o	
Agriculture	4.D1.6 Sewage sludge and Industrial waste used as fertiliser	N ₂ O	0.028	0.000	0.994
Industrial Proc.	2A7b Yellow bricks	CO ₂	0.023	0.000	0.995
Energy	1.B.2 Flaring in refinery	CO ₂	0.023	0.000	0.995
Waste	6 B. Wastewater Handling - Direct	N ₂ O	0.023	0.000	0.995
Solvent and Other Prod. Use	3D5 Use of candles	CO ₂	0.022	0.000	0.995
Solvent and Other Prod. Use	3C Chemical products, manufacturing and processing	CO ₂	0.019	0.000	0.996
Industrial Proc.	2A7a Glass and Glass wool	CO ₂	0.017	0.000	0.996
Energy	1.B.2 Land based activities	CH ₄	0.017	0.000	0.996
Energy	Stationary Combustion, GAS	N ₂ O	0.016	0.000	0.996
LULUCF	5(II) Forest Land.	N ₂ O	0.016	0.000	0.997
Energy	Transport, Agriculture	N ₂ O	0.015	0.000	0.997
Industrial Proc.	2A7c Expanded clay	CO ₂	0.015	0.000	0.997
Energy	1.B.2 Off-shore activities	CH ₄	0.015	0.000	0.997
Energy	Transport, Navigation (large vessels)	N ₂ O	0.015	0.000	0.997
Industrial Proc.	2A3 Limestone and dolomite use	CO ₂	0.014	0.000	0.998
Solvent and Other Prod. Use	3A Paint application	CO ₂	0.013	0.000	0.998
Energy	Stationary Combustion, SOLID	CH ₄	0.013	0.000	0.998
Waste	6.D Compost production	N ₂ O	0.013	0.000	0.998
Energy	Transport, Fisheries	N ₂ O	0.011	0.000	0.998
LULUCF	5.E Settlements, Living biomass	CO ₂	0.011	0.000	0.998
Waste	6.D Accidental fires, buildings	CO ₂	0.011	0.000	0.999
Energy	Stationary Combustion, BKB	CO ₂	0.011	0.000	0.999
Energy	Transport, Industry (mobile)	N ₂ O	0.011	0.000	0.999
Energy	Stationary Combustion, WASTE	N ₂ O	0.007	0.000	0.999
Waste	6.D Accidental fires, vehicles	CO ₂	0.006	0.000	0.999
Energy	1.B.2 Distribution of natural gas	CH ₄	0.005	0.000	0.999
Energy	Natural gas fuelled engines, GAS	CH ₄	0.005	0.000	0.999
Industrial Proc.	2D2 Food and Drink	CO ₂	0.004	0.000	0.999
Energy	1.B.2 Transmission of natural gas	CH ₄	0.004	0.000	0.999
Energy	Transport, Civil aviation	N ₂ O	0.003	0.000	0.999
Energy	Stationary Combustion, GAS	CH ₄	0.003	0.000	0.999
LULUCF	5.B Cropland, Dead organic matter	CO ₂	0.003	0.000	0.999
Energy	Stationary Combustion, LIQUID	CH ₄	0.003	0.000	0.999
LULUCF	5.D Wetlands, Living biomass	CO ₂	0.003	0.000	1.000
Energy	Transport, Railways	N ₂ O	0.003	0.000	1.000
Energy	1.B.2 Off-shore activities	CO ₂	0.002	0.000	1.000
Energy	Transport, Agriculture	CH ₄	0.002	0.000	1.000
Energy	Transport, Commercial/institutional	CH ₄	0.002	0.000	1.000
Agriculture	4F Field burning of agricultural residues	CH ₄	0.002	0.000	1.000
Industrial Proc.	2A6 Road paving with asphalt	CO ₂	0.002	0.000	1.000
LULUCF	5.C Grassland, Dead organic matter	CO ₂	0.002	0.000	1.000
Energy	Biogas fuelled engines, BIOMASS	CH ₄	0.001	0.000	1.000
Waste	6.D Accidental fires, buildings	CH ₄	0.001	0.000	1.000
Energy	Transport, Industry (mobile)	CH ₄	0.001	0.000	1.000
Energy	1.B.2 Venting in gas storage	CH ₄	0.001	0.000	1.000
Energy	Transport, Military	N ₂ O	0.001	0.000	1.000
Energy	Transport, Residential	CH ₄	0.001	0.000	1.000
LULUCF	5.E Settlements, Soils	CO ₂	0.001	0.000	1.000
Industrial Proc.	2B5 Catalysts/Fertilizers, Pesticides and Sulphuric acid	CO ₂	0.001	0.000	1.000
Energy	1.B.2 Refinery processes	CH ₄	0.001	0.000	1.000
Energy	Stationary Combustion, WASTE	CH ₄	0.001	0.000	1.000
Solvent and Other Prod. Use	3D5 Consumption of fireworks	N ₂ O	0.001	0.000	1.000
Energy	1.B.2. Flaring off-shore	CH ₄	0.001	0.000	1.000
Energy	1.B.2 Flaring off-shore	N ₂ O	0.001	0.000	1.000
Agriculture	4.F Field Burning of Agricultural Residues	N ₂ O	0.001	0.000	1.000
LULUCF	5.E Settlements, Dead organic matter	CO ₂	0.001	0.000	1.000
LULUCF	5(V) Biomass Burning	CH ₄	0.001	0.000	1.000
LULUCF	5.D Wetlands, Dead organic matter	CO ₂	0.001	0.000	1.000

Tier 1 Analysis		DK			
IPCC Source Categories (LULUCF included)		GHG	Base Year Estimate	Base Year Level Assessment	Base Year Cumulative Total of Lx,o
			Ex,o	Lx,o	
			Mt CO ₂ eq		
Industrial Proc.	2F Consumption of PFC	PFC	0.001	0.000	1.000
LULUCF	5(V) Biomass Burning	N ₂ O	0.000	0.000	1.000
Energy	Transport, Forestry	CH ₄	0.000	0.000	1.000
Energy	Transport, Navigation (small boats)	N ₂ O	0.000	0.000	1.000
Energy	Transport, Navigation (small boats)	CH ₄	0.000	0.000	1.000
Energy	Transport, Commercial/institutional	N ₂ O	0.000	0.000	1.000
Energy	Transport, Navigation (large vessels)	CH ₄	0.000	0.000	1.000
Energy	Transport, Fisheries	CH ₄	0.000	0.000	1.000
Waste	6.D Accidental fires, vehicles	CH ₄	0.000	0.000	1.000
Energy	Transport, Railways	CH ₄	0.000	0.000	1.000
Solvent and Other Prod. Use	3D5 Use of tobacco	N ₂ O	0.000	0.000	1.000
LULUCF	5(III) Disturbance, Land converted to cropland	N ₂ O	0.000	0.000	1.000
Energy	Transport, Residential	N ₂ O	0.000	0.000	1.000
Waste	6.C Incineration of corpses	N ₂ O	0.000	0.000	1.000
Energy	Transport, Forestry	N ₂ O	0.000	0.000	1.000
Energy	1.B.2. Flaring in refinery	CH ₄	0.000	0.000	1.000
LULUCF	5.C Grassland, Mineral soils	CO ₂	0.000	0.000	1.000
LULUCF	5(II) Wetlands	N ₂ O	0.000	0.000	1.000
Energy	Transport, Military	CH ₄	0.000	0.000	1.000
Energy	Transport, Civil aviation	CH ₄	0.000	0.000	1.000
Solvent and Other Prod. Use	3D5 Use of charcoal for BBQ	N ₂ O	0.000	0.000	1.000
Energy	1.B.2 Flaring in refinery	N ₂ O	0.000	0.000	1.000
Solvent and Other Prod. Use	3D5 Use of candles	N ₂ O	0.000	0.000	1.000
Solvent and Other Prod. Use	3D5 Consumption of fireworks	CO ₂	0.000	0.000	1.000
Industrial Proc.	2A5 Asphalt roofing	CO ₂	0.000	0.000	1.000
Waste	6.C Incineration of carcasses	N ₂ O	0.000	0.000	1.000
Waste	6.C Incineration of corpses	CH ₄	0.000	0.000	1.000
Energy	1.B.2 Distribution of natural gas	CO ₂	0.000	0.000	1.000
Energy	1.B.2 Transmission of natural gas	CO ₂	0.000	0.000	1.000
Energy	1.B.2 Land based activities	CO ₂	0.000	0.000	1.000
Energy	1.B.2 Venting in gas storage	CO ₂	0.000	0.000	1.000
Waste	6.C Incineration of carcasses	CH ₄	0.000	0.000	1.000
Solvent and Other Prod. Use	3B Degreasing and dry cleaning	CO ₂	0.000	0.000	1.000
Solvent and Other Prod. Use	3D1 Other - Use of N ₂ O for Anaesthesia	N ₂ O	0.000	0.000	1.000
Total			74.372	1.000	

¹⁾ The Estimates include signs, where + : emission - : removal, although in the level analyses only the absolute values are used.

Table A1-14 KCA for Denmark, level assessment 2012 excl. LULUCF, tier 1.

Tier 1 Analysis IPCC Source Categories (LULUCF excluded)		DK GHG	Latest Year Estimate Ex,t Mt CO ₂ eq	Latest Year Level Assessment Lx,t	Latest Year Cumulative Total of Lx,t
Energy	Transport, Road transport	CO ₂	11.224	0.217	0.217
Energy	Stationary Combustion, Coal	CO ₂	10.005	0.194	0.411
Energy	Stationary Combustion, Natural gas	CO ₂	8.293	0.161	0.572
Agriculture	4A Enteric Fermentation	CH ₄	2.904	0.056	0.628
Agriculture	4.D3 Leaching	N ₂ O	1.430	0.028	0.656
Energy	Stationary Combustion, Fossil waste	CO ₂	1.397	0.027	0.683
Energy	Transport, Agriculture	CO ₂	1.343	0.026	0.709
Agriculture	4B Manure Management	CH ₄	1.297	0.025	0.734
Agriculture	4.D1.2 Animal waste applied to soils	N ₂ O	1.161	0.022	0.756
Agriculture	4.D1.1 Synthetic Fertilizer	N ₂ O	1.103	0.021	0.778
Energy	Transport, Industry (mobile)	CO ₂	1.021	0.020	0.797
Energy	Stationary Combustion, Refinery gas	CO ₂	0.906	0.018	0.815
Industrial Proc.	2A1 Cement production	CO ₂	0.871	0.017	0.832
Energy	Stationary Combustion, Gas oil	CO ₂	0.742	0.014	0.846
Waste	6 A. Solid Waste Disposal on Land	CH ₄	0.698	0.014	0.860
Industrial Proc.	2F Consumption of HFC	HFC	0.657	0.013	0.872
Energy	Stationary Combustion, Petroleum coke	CO ₂	0.628	0.012	0.885
Energy	Stationary Combustion, Residual oil	CO ₂	0.571	0.011	0.896
Energy	Transport, Fisheries	CO ₂	0.479	0.009	0.905
Energy	Transport, Navigation (large vessels)	CO ₂	0.399	0.008	0.913
Agriculture	4.B Manure Management	N ₂ O	0.391	0.008	0.920
Agriculture	4.D1.4 Crop Residue	N ₂ O	0.311	0.006	0.926
Agriculture	4.D3 Atmospheric deposition	N ₂ O	0.295	0.006	0.932
Agriculture	4.D1.3 N-fixing crops	N ₂ O	0.256	0.005	0.937
Energy	Transport, Railways	CO ₂	0.249	0.005	0.942
Agriculture	4.D.2 Grassing animals	N ₂ O	0.211	0.004	0.946
Agriculture	4.D1.5 Cultivation of histosols	N ₂ O	0.198	0.004	0.950
Energy	1.B.2 Flaring off-shore	CO ₂	0.195	0.004	0.953
Energy	Transport, Commercial/institutional	CO ₂	0.171	0.003	0.957
Energy	Stationary Combustion, BIOMASS	CH ₄	0.135	0.003	0.959
Energy	Transport, Civil aviation	CO ₂	0.133	0.003	0.962
Waste	6.D Compost production	N ₂ O	0.127	0.002	0.964
Energy	Natural gas fuelled engines, GAS	CH ₄	0.120	0.002	0.967
Industrial Proc.	2F Consumption of SF ₆	SF ₆	0.118	0.002	0.969
Energy	Transport, Road transport	N ₂ O	0.116	0.002	0.971
Energy	Transport, Military	CO ₂	0.116	0.002	0.974
Energy	Transport, Navigation (small boats)	CO ₂	0.099	0.002	0.975
Energy	Stationary Combustion, LPG	CO ₂	0.091	0.002	0.977
Energy	Stationary Combustion, BIOMASS	N ₂ O	0.091	0.002	0.979
Waste	6.D Compost production	CH ₄	0.090	0.002	0.981
Solvent and Other Prod. Use	3D5 Use of candles	CO ₂	0.081	0.002	0.982
Waste	6 B. Wastewater Handling	CH ₄	0.074	0.001	0.984
Energy	Stationary Combustion, Coke	CO ₂	0.074	0.001	0.985
Energy	Transport, Residential	CO ₂	0.062	0.001	0.986
Energy	1.B.2 Refinery processes	CH ₄	0.047	0.001	0.987
Solvent and Other Prod. Use	3D5 Other	CO ₂	0.044	0.001	0.988
Waste	6 B. Wastewater Handling - Direct	N ₂ O	0.041	0.001	0.989
Industrial Proc.	2A2 Lime production	CO ₂	0.040	0.001	0.990
Agriculture	4.D1.6 Sewage sludge and Industrial waste used as fertiliser	N ₂ O	0.039	0.001	0.990
Energy	1.B.2 Off-shore activities	CH ₄	0.037	0.001	0.991
Waste	6 B. Wastewater Handling - Indirect	N ₂ O	0.032	0.001	0.992
Industrial Proc.	2G Lubricants	CO ₂	0.032	0.001	0.992
Energy	Biogas fuelled engines, BIOMASS	CH ₄	0.031	0.001	0.993
Energy	Stationary Combustion, SOLID	N ₂ O	0.027	0.001	0.994
Industrial Proc.	2A3 Limestone and dolomite use	CO ₂	0.026	0.000	0.994
Energy	Stationary Combustion, GAS	N ₂ O	0.024	0.000	0.995
Energy	1.B.2 Flaring in refinery	CO ₂	0.022	0.000	0.995
Energy	Transport, Agriculture	N ₂ O	0.018	0.000	0.995
Energy	1.B.2 Land based activities	CH ₄	0.018	0.000	0.996
Industrial Proc.	2A7b Yellow bricks	CO ₂	0.017	0.000	0.996
Energy	Transport, Forestry	CO ₂	0.017	0.000	0.996
Energy	Stationary Combustion, WASTE	N ₂ O	0.016	0.000	0.997

Tier 1 Analysis IPCC Source Categories (LULUCF excluded)		DK GHG	Latest Year Estimate	Latest Year Level Assessment	Latest Year Cumulative Total of Lx,t
			Ex,t Mt CO ₂ eq	Lx,t	
Energy	Transport, Industry (mobile)	N ₂ O	0.013	0.000	0.997
Solvent and Other Prod. Use	3C Chemical products, manufacturing and processing	CO ₂	0.012	0.000	0.997
Energy	Transport, Road transport	CH ₄	0.011	0.000	0.997
Energy	Stationary Combustion, LIQUID	N ₂ O	0.011	0.000	0.998
Waste	6.D Accidental fires, buildings	CO ₂	0.011	0.000	0.998
Industrial Proc.	2A7a Glass and Glass wool	CO ₂	0.010	0.000	0.998
Energy	Transport, Fisheries	N ₂ O	0.009	0.000	0.998
Solvent and Other Prod. Use	3D1 Other - Use of N ₂ O for Anaesthesia	N ₂ O	0.009	0.000	0.998
Industrial Proc.	2F Consumption of PFC	PFC	0.009	0.000	0.998
Energy	Transport, Navigation (large vessels)	N ₂ O	0.008	0.000	0.999
Solvent and Other Prod. Use	3A Paint application	CO ₂	0.007	0.000	0.999
Industrial Proc.	2A7c Expanded clay	CO ₂	0.006	0.000	0.999
Energy	Stationary Combustion, GAS	CH ₄	0.006	0.000	0.999
Waste	6.D Accidental fires, vehicles	CO ₂	0.006	0.000	0.999
Energy	1.B.2 Off-shore activities	CO ₂	0.004	0.000	0.999
Energy	Transport, Commercial/institutional	CH ₄	0.003	0.000	0.999
Energy	Stationary Combustion, SOLID	CH ₄	0.003	0.000	0.999
Energy	1.B.2 Distribution of natural gas	CH ₄	0.003	0.000	0.999
Agriculture	4F Field burning of agricultural residues	CH ₄	0.002	0.000	0.999
Industrial Proc.	2D2 Food and Drink	CO ₂	0.002	0.000	0.999
Energy	Transport, Railways	N ₂ O	0.002	0.000	0.999
Solvent and Other Prod. Use	3D5 Consumption of fireworks	N ₂ O	0.002	0.000	0.999
Energy	Transport, Civil aviation	N ₂ O	0.002	0.000	1.000
Energy	Transport, Agriculture	CH ₄	0.002	0.000	1.000
Industrial Proc.	2A6 Road paving with asphalt	CO ₂	0.002	0.000	1.000
Energy	Stationary Combustion, Kerosene	CO ₂	0.002	0.000	1.000
Energy	Stationary Combustion, WASTE	CH ₄	0.002	0.000	1.000
Energy	Stationary Combustion, BKB	CO ₂	0.001	0.000	1.000
Industrial Proc.	2B5 Catalysts/Fertilizers, Pesticides and Sulphuric acid	CO ₂	0.001	0.000	1.000
Energy	Transport, Residential	CH ₄	0.001	0.000	1.000
Waste	6.D Accidental fires, buildings	CH ₄	0.001	0.000	1.000
Energy	Transport, Military	N ₂ O	0.001	0.000	1.000
Energy	1.B.2 Venting in gas storage	CH ₄	0.001	0.000	1.000
Energy	Stationary Combustion, LIQUID	CH ₄	0.001	0.000	1.000
Energy	Transport, Navigation (small boats)	N ₂ O	0.001	0.000	1.000
Agriculture	4.F Field Burning of Agricultural Residues	N ₂ O	0.001	0.000	1.000
Energy	Transport, Commercial/institutional	N ₂ O	0.001	0.000	1.000
Energy	Transport, Industry (mobile)	CH ₄	0.001	0.000	1.000
Energy	Transport, Navigation (small boats)	CH ₄	0.001	0.000	1.000
Energy	1.B.2 Flaring off-shore	N ₂ O	0.000	0.000	1.000
Energy	Transport, Residential	N ₂ O	0.000	0.000	1.000
Energy	1.B.2. Flaring off-shore	CH ₄	0.000	0.000	1.000
Energy	1.B.2 Transmission of natural gas	CH ₄	0.000	0.000	1.000
Energy	Transport, Fisheries	CH ₄	0.000	0.000	1.000
Waste	6.D Accidental fires, vehicles	CH ₄	0.000	0.000	1.000
Solvent and Other Prod. Use	3D5 Use of candles	N ₂ O	0.000	0.000	1.000
Energy	Transport, Navigation (large vessels)	CH ₄	0.000	0.000	1.000
Waste	6.C Incineration of corpses	N ₂ O	0.000	0.000	1.000
Energy	Transport, Forestry	N ₂ O	0.000	0.000	1.000
Solvent and Other Prod. Use	3D5 Use of tobacco	N ₂ O	0.000	0.000	1.000
Energy	1.B.2. Flaring in refinery	CH ₄	0.000	0.000	1.000
Energy	Transport, Railways	CH ₄	0.000	0.000	1.000
Solvent and Other Prod. Use	3D5 Consumption of fireworks	CO ₂	0.000	0.000	1.000
Solvent and Other Prod. Use	3D5 Use of charcoal for BBQ	N ₂ O	0.000	0.000	1.000
Waste	6.C Incineration of carcasses	N ₂ O	0.000	0.000	1.000
Energy	Transport, Military	CH ₄	0.000	0.000	1.000
Energy	1.B.2 Flaring in refinery	N ₂ O	0.000	0.000	1.000
Energy	Transport, Civil aviation	CH ₄	0.000	0.000	1.000
Energy	Transport, Forestry	CH ₄	0.000	0.000	1.000
Industrial Proc.	2A5 Asphalt roofing	CO ₂	0.000	0.000	1.000
Waste	6.C Incineration of corpses	CH ₄	0.000	0.000	1.000
Energy	1.B.2 Land based activities	CO ₂	0.000	0.000	1.000

Tier 1 Analysis IPCC Source Categories (LULUCF excluded)		DK GHG	Latest Year Estimate	Latest Year Level Assessment	Latest Year Cumulative Total of Lx,t
			Ex,t Mt CO ₂ eq	Lx,t	
Energy	1.B.2 Distribution of natural gas	CO ₂	0.000	0.000	1.000
Waste	6.C Incineration of carcasses	CH ₄	0.000	0.000	1.000
Energy	1.B.2 Venting in gas storage	CO ₂	0.000	0.000	1.000
Energy	1.B.2 Transmission of natural gas	CO ₂	0.000	0.000	1.000
Solvent and Other Prod. Use	3B Degreasing and dry cleaning	CO ₂	0.000	0.000	1.000
Industrial Proc.	2C1 Iron and steel production	CO ₂	0.000	0.000	1.000
Industrial Proc.	2B2 Nitric acid production	N ₂ O	0.000	0.000	1.000
Total			51.637	1.000	

Table A1-15 KCA for Denmark, level assessment 2012 incl. LULUCF, tier 1.

Tier 1 Analysis IPCC Source Categories (LU- LUCF included)		DK GHG	Latest Year Estimate Ex,t Mt CO ₂ eq	Latest Year Level Assessment Lx,t	Latest Year Cumulative Total of Lx,t
Energy	Transport, Road transport	CO ₂	11.224	0.188	0.188
Energy	Stationary Combustion, Coal	CO ₂	10.005	0.167	0.355
Energy	Stationary Combustion, Natural gas	CO ₂	8.293	0.139	0.494
LULUCF	5.A.1 Forest remaining forest	CO ₂	4.491	0.075	0.569
Agriculture	4A Enteric Fermentation	CH ₄	2.904	0.049	0.618
LULUCF	5.B Cropland, Organic soils	CO ₂	1.981	0.033	0.651
Agriculture	4.D3 Leaching	N ₂ O	1.430	0.024	0.675
Energy	Stationary Combustion, Fossil waste	CO ₂	1.397	0.023	0.698
Energy	Transport, Agriculture	CO ₂	1.343	0.022	0.720
Agriculture	4B Manure Management	CH ₄	1.297	0.022	0.742
Agriculture	4.D1.2 Animal waste applied to soils	N ₂ O	1.161	0.019	0.762
Agriculture	4.D1.1 Synthetic Fertilizer	N ₂ O	1.103	0.018	0.780
Energy	Transport, Industry (mobile)	CO ₂	1.021	0.017	0.797
Energy	Stationary Combustion, Refinery gas	CO ₂	0.906	0.015	0.812
Industrial Proc.	2A1 Cement production	CO ₂	0.871	0.015	0.827
Energy	Stationary Combustion, Gas oil	CO ₂	0.742	0.012	0.839
Waste	6 A. Solid Waste Disposal on Land	CH ₄	0.698	0.012	0.851
Industrial Proc.	2F Consumption of HFC	HFC	0.657	0.011	0.862
Energy	Stationary Combustion, Petroleum coke	CO ₂	0.628	0.010	0.872
LULUCF	5.B Cropland, Mineral soils	CO ₂	0.577	0.010	0.882
Energy	Stationary Combustion, Residual oil	CO ₂	0.571	0.010	0.892
Energy	Transport, Fisheries	CO ₂	0.479	0.008	0.900
LULUCF	5.C Grassland, Living biomass	CO ₂	0.463	0.008	0.907
Energy	Transport, Navigation (large vessels)	CO ₂	0.399	0.007	0.914
Agriculture	4.B Manure Management	N ₂ O	0.391	0.007	0.921
Agriculture	4.D1.4 Crop Residue	N ₂ O	0.311	0.005	0.926
Agriculture	4.D3 Atmospheric deposition	N ₂ O	0.295	0.005	0.931
Agriculture	4.D1.3 N-fixing crops	N ₂ O	0.256	0.004	0.935
Energy	Transport, Railways	CO ₂	0.249	0.004	0.939
Agriculture	4.D.2 Grassing animals	N ₂ O	0.211	0.004	0.943
Agriculture	4.D1.5 Cultivation of histosols	N ₂ O	0.198	0.003	0.946
Energy	1.B.2 Flaring off-shore	CO ₂	0.195	0.003	0.949
LULUCF	5.B Cropland, Living biomass	CO ₂	0.193	0.003	0.952
LULUCF	5(IV) Cropland Limestone	CO ₂	0.192	0.003	0.956
Energy	Transport, Commercial/institutional	CO ₂	0.171	0.003	0.959
Energy	Stationary Combustion, BIOMASS	CH ₄	0.135	0.002	0.961
Energy	Transport, Civil aviation	CO ₂	0.133	0.002	0.963
Waste	6.D Compost production	N ₂ O	0.127	0.002	0.965
Energy	Natural gas fuelled engines, GAS	CH ₄	0.120	0.002	0.967
Industrial Proc.	2F Consumption of SF ₆	SF ₆	0.118	0.002	0.969
Energy	Transport, Road transport	N ₂ O	0.116	0.002	0.971
Energy	Transport, Military	CO ₂	0.116	0.002	0.973
Energy	Transport, Navigation (small boats)	CO ₂	0.099	0.002	0.975
Energy	Stationary Combustion, LPG	CO ₂	0.091	0.002	0.976
Energy	Stationary Combustion, BIOMASS	N ₂ O	0.091	0.002	0.978
Waste	6.D Compost production	CH ₄	0.090	0.002	0.979
Solvent and Other Prod. Use	3D5 Use of candles	CO ₂	0.081	0.001	0.981
LULUCF	5.C Grassland, Organic soils	CO ₂	0.079	0.001	0.982
Waste	6 B. Wastewater Handling	CH ₄	0.074	0.001	0.983
Energy	Stationary Combustion, Coke	CO ₂	0.074	0.001	0.984
Energy	Transport, Residential	CO ₂	0.062	0.001	0.985
LULUCF	5.E Settlements, Living biomass	CO ₂	0.056	0.001	0.986
Energy	1.B.2 Refinery processes	CH ₄	0.047	0.001	0.987
Solvent and Other Prod. Use	3D5 Other	CO ₂	0.044	0.001	0.988
Waste	6 B. Wastewater Handling - Direct	N ₂ O	0.041	0.001	0.989
Industrial Proc.	2A2 Lime production	CO ₂	0.040	0.001	0.989
Agriculture	4.D1.6 Sewage sludge and Industrial waste used as fertiliser	N ₂ O	0.039	0.001	0.990
LULUCF	5.A.2 Land converted to forest	CO ₂	0.038	0.001	0.991
Energy	1.B.2 Off-shore activities	CH ₄	0.037	0.001	0.991
LULUCF	5.E Settlements, Soils	CO ₂	0.035	0.001	0.992

Tier 1 Analysis IPCC Source Categories (LU- LUCF included)		DK GHG	Latest Year Estimate	Latest Year Level Assessment Lx,t	Latest Year Cumulative Total of Lx,t
			Ex,t Mt CO ₂ eq		
Waste	6 B. Wastewater Handling - Indirect	N ₂ O	0.032	0.001	0.992
Industrial Proc.	2G Lubricants	CO ₂	0.032	0.001	0.993
Energy	Biogas fuelled engines, BIOMASS	CH ₄	0.031	0.001	0.993
Energy	Stationary Combustion, SOLID	N ₂ O	0.027	0.000	0.994
Industrial Proc.	2A3 Limestone and dolomite use	CO ₂	0.026	0.000	0.994
Energy	Stationary Combustion, GAS	N ₂ O	0.024	0.000	0.995
Energy	1.B.2 Flaring in refinery	CO ₂	0.022	0.000	0.995
Energy	Transport, Agriculture	N ₂ O	0.018	0.000	0.995
Energy	1.B.2 Land based activities	CH ₄	0.018	0.000	0.996
Industrial Proc.	2A7b Yellow bricks	CO ₂	0.017	0.000	0.996
Energy	Transport, Forestry	CO ₂	0.017	0.000	0.996
Energy	Stationary Combustion, WASTE	N ₂ O	0.016	0.000	0.996
Energy	Transport, Industry (mobile)	N ₂ O	0.013	0.000	0.997
Solvent and Other Prod. Use	3C Chemical products, manufacturing and processing	CO ₂	0.012	0.000	0.997
LULUCF	5.B Cropland, Dead organic matter	CO ₂	0.012	0.000	0.997
LULUCF	5(II) Forest Land.	N ₂ O	0.012	0.000	0.997
Energy	Transport, Road transport	CH ₄	0.011	0.000	0.997
Energy	Stationary Combustion, LIQUID	N ₂ O	0.011	0.000	0.998
Waste	6.D Accidental fires, buildings	CO ₂	0.011	0.000	0.998
Industrial Proc.	2A7a Glass and Glass wool	CO ₂	0.010	0.000	0.998
Energy	Transport, Fisheries	N ₂ O	0.009	0.000	0.998
Solvent and Other Prod. Use	3D1 Other - Use of N ₂ O for Anaesthesia	N ₂ O	0.009	0.000	0.998
Industrial Proc.	2F Consumption of PFC	PFC	0.009	0.000	0.998
Energy	Transport, Navigation (large vessels)	N ₂ O	0.008	0.000	0.999
Solvent and Other Prod. Use	3A Paint application	CO ₂	0.007	0.000	0.999
LULUCF	5.C Grassland, Dead organic matter	CO ₂	0.007	0.000	0.999
Industrial Proc.	2A7c Expanded clay	CO ₂	0.006	0.000	0.999
Energy	Stationary Combustion, GAS	CH ₄	0.006	0.000	0.999
Waste	6.D Accidental fires, vehicles	CO ₂	0.006	0.000	0.999
LULUCF	5.C Grassland, Mineral soils	CO ₂	0.005	0.000	0.999
Energy	1.B.2 Off-shore activities	CO ₂	0.004	0.000	0.999
Energy	Transport, Commercial/institutional	CH ₄	0.003	0.000	0.999
Energy	Stationary Combustion, SOLID	CH ₄	0.003	0.000	0.999
Energy	1.B.2 Distribution of natural gas	CH ₄	0.003	0.000	0.999
LULUCF	5.D Wetlands, Soils	CO ₂	0.002	0.000	0.999
Agriculture	4F Field burning of agricultural residues	CH ₄	0.002	0.000	0.999
Industrial Proc.	2D2 Food and Drink	CO ₂	0.002	0.000	0.999
Energy	Transport, Railways	N ₂ O	0.002	0.000	1.000
Solvent and Other Prod. Use	3D5 Consumption of fireworks	N ₂ O	0.002	0.000	1.000
Energy	Transport, Civil aviation	N ₂ O	0.002	0.000	1.000
Energy	Transport, Agriculture	CH ₄	0.002	0.000	1.000
Industrial Proc.	2A6 Road paving with asphalt	CO ₂	0.002	0.000	1.000
Energy	Stationary Combustion, Kerosene	CO ₂	0.002	0.000	1.000
Energy	Stationary Combustion, WASTE	CH ₄	0.002	0.000	1.000
Energy	Stationary Combustion, BKB	CO ₂	0.001	0.000	1.000
Industrial Proc.	2B5 Catalysts/Fertilizers, Pesticides and Sulphuric acid	CO ₂	0.001	0.000	1.000
Energy	Transport, Residential	CH ₄	0.001	0.000	1.000
Waste	6.D Accidental fires, buildings	CH ₄	0.001	0.000	1.000
Energy	Transport, Military	N ₂ O	0.001	0.000	1.000
Energy	1.B.2 Venting in gas storage	CH ₄	0.001	0.000	1.000
Energy	Stationary Combustion, LIQUID	CH ₄	0.001	0.000	1.000
Energy	Transport, Navigation (small boats)	N ₂ O	0.001	0.000	1.000
LULUCF	5(III) Disturbance, Land converted to cropland	N ₂ O	0.001	0.000	1.000
Agriculture	4.F Field Burning of Agricultural Residues	N ₂ O	0.001	0.000	1.000
Energy	Transport, Commercial/institutional	N ₂ O	0.001	0.000	1.000
Energy	Transport, Industry (mobile)	CH ₄	0.001	0.000	1.000
Energy	Transport, Navigation (small boats)	CH ₄	0.001	0.000	1.000
Energy	1.B.2 Flaring off-shore	N ₂ O	0.000	0.000	1.000
Energy	Transport, Residential	N ₂ O	0.000	0.000	1.000

Tier 1 Analysis IPCC Source Categories (LU- LUCF included)		DK GHG	Latest Year Estimate Ex,t Mt CO ₂ eq	Latest Year Level Asses- ment Lx,t	Latest Year Cumulative Total of Lx,t
Energy	1.B.2. Flaring off-shore	CH ₄	0.000	0.000	1.000
Energy	1.B.2 Transmission of natural gas	CH ₄	0.000	0.000	1.000
Energy	Transport, Fisheries	CH ₄	0.000	0.000	1.000
Waste	6.D Accidental fires, vehicles	CH ₄	0.000	0.000	1.000
LULUCF	5.D Wetlands, Living biomass	CO ₂	0.000	0.000	1.000
Solvent and Other Prod. Use	3D5 Use of candles	N ₂ O	0.000	0.000	1.000
Energy	Transport, Navigation (large vessels)	CH ₄	0.000	0.000	1.000
Waste	6.C Incineration of corpses	N ₂ O	0.000	0.000	1.000
Energy	Transport, Forestry	N ₂ O	0.000	0.000	1.000
Solvent and Other Prod. Use	3D5 Use of tobacco	N ₂ O	0.000	0.000	1.000
Energy	1.B.2. Flaring in refinery	CH ₄	0.000	0.000	1.000
Energy	Transport, Railways	CH ₄	0.000	0.000	1.000
Solvent and Other Prod. Use	3D5 Consumption of fireworks	CO ₂	0.000	0.000	1.000
LULUCF	5(II) Wetlands	N ₂ O	0.000	0.000	1.000
Solvent and Other Prod. Use	3D5 Use of charcoal for BBQ	N ₂ O	0.000	0.000	1.000
Waste	6.C Incineration of carcasses	N ₂ O	0.000	0.000	1.000
Energy	Transport, Military	CH ₄	0.000	0.000	1.000
Energy	1.B.2 Flaring in refinery	N ₂ O	0.000	0.000	1.000
Energy	Transport, Civil aviation	CH ₄	0.000	0.000	1.000
Energy	Transport, Forestry	CH ₄	0.000	0.000	1.000
LULUCF	5(V) Biomass Burning	N ₂ O	0.000	0.000	1.000
LULUCF	5.E Settlements, Dead organic matter	CO ₂	0.000	0.000	1.000
LULUCF	5(V) Biomass Burning	CH ₄	0.000	0.000	1.000
Industrial Proc.	2A5 Asphalt roofing	CO ₂	0.000	0.000	1.000
Waste	6.C Incineration of corpses	CH ₄	0.000	0.000	1.000
Energy	1.B.2 Land based activities	CO ₂	0.000	0.000	1.000
Energy	1.B.2 Distribution of natural gas	CO ₂	0.000	0.000	1.000
Waste	6.C Incineration of carcasses	CH ₄	0.000	0.000	1.000
Energy	1.B.2 Venting in gas storage	CO ₂	0.000	0.000	1.000
Energy	1.B.2 Transmission of natural gas	CO ₂	0.000	0.000	1.000
Solvent and Other Prod. Use	3B Degreasing and dry cleaning	CO ₂	0.000	0.000	1.000
Industrial Proc.	2C1 Iron and steel production	CO ₂	0.000	0.000	1.000
Industrial Proc.	2B2 Nitric acid production	N ₂ O	0.000	0.000	1.000
LULUCF	5.D Wetlands, Dead organic matter	CO ₂	0.000	0.000	1.000
Total			59.782	1.000	

¹⁾ The Estimates include signs, where + : emission - : removal, although in the level analyses only the absolute values are used.

Table A1-16 KCA for Denmark, trend assessment 1990-2012 excl. LULUCF, tier 1.

Tier 1 Analysis IPCC Source Categories (LULUCF excluded)		DK GHG	Base Year Estimate	Latest Year Estimate	Trend Assessment	Contribution to Trend	Cumulative
			Ex,o Mt CO ₂ eq	Ex,t Mt CO ₂ eq	Tx,t		
Energy	Stationary Combustion, Coal	CO ₂	23.834	10.005	0.1138	0.262	0.262
Energy	Stationary Combustion, Natural gas	CO ₂	4.335	8.293	0.0732	0.168	0.430
Energy	Transport, Road transport	CO ₂	9.284	11.224	0.0619	0.143	0.573
Energy	Stationary Combustion, Gas oil	CO ₂	4.547	0.742	0.0386	0.089	0.662
Energy	Stationary Combustion, Residual oil	CO ₂	2.496	0.571	0.0188	0.043	0.705
Energy	Stationary Combustion, Fossil waste	CO ₂	0.573	1.397	0.0140	0.032	0.737
Industrial Proc.	2B2 Nitric acid production	N ₂ O	1.043	0.000	0.0113	0.026	0.763
Agriculture	4.D1.1 Synthetic Fertilizer	N ₂ O	2.354	1.103	0.0096	0.022	0.785
Agriculture	4B Manure Management	CH ₄	0.985	1.297	0.0081	0.019	0.804
Industrial Proc.	2F Consumption of HFC	HFC	0.218	0.657	0.0072	0.016	0.821
Agriculture	4A Enteric Fermentation	CH ₄	3.247	2.904	0.0068	0.016	0.836
Agriculture	4.D3 Leaching	N ₂ O	2.447	1.430	0.0059	0.013	0.850
Energy	Transport, Industry (mobile)	CO ₂	0.839	1.021	0.0057	0.013	0.863
Energy	Transport, Agriculture	CO ₂	1.272	1.343	0.0057	0.013	0.876
Agriculture	4.D1.2 Animal waste applied to soils	N ₂ O	1.112	1.161	0.0048	0.011	0.887
Waste	6 A. Solid Waste Disposal on Land	CH ₄	1.366	0.698	0.0047	0.011	0.898
Energy	Stationary Combustion, Petroleum coke	CO ₂	0.415	0.628	0.0046	0.011	0.908
Energy	Stationary Combustion, Refinery gas	CO ₂	0.816	0.906	0.0043	0.010	0.918
Energy	Stationary Combustion, Kerosene	CO ₂	0.366	0.002	0.0039	0.009	0.927
Industrial Proc.	2A1 Cement production	CO ₂	0.882	0.871	0.0030	0.007	0.934
Energy	Transport, Navigation (large vessels)	CO ₂	0.748	0.399	0.0023	0.005	0.940
Waste	6.D Compost production	N ₂ O	0.013	0.127	0.0017	0.004	0.943
Energy	Natural gas fuelled engines, GAS	CH ₄	0.005	0.120	0.0017	0.004	0.947
Energy	Transport, Commercial/institutional	CO ₂	0.074	0.171	0.0017	0.004	0.951
Agriculture	4.D3 Atmospheric deposition	N ₂ O	0.496	0.295	0.0011	0.003	0.954
Waste	6.D Compost production	CH ₄	0.029	0.090	0.0010	0.002	0.956
Solvent and Other Prod. Use	3D5 Use of candles	CO ₂	0.022	0.081	0.0009	0.002	0.958
Energy	Transport, Navigation (small boats)	CO ₂	0.048	0.099	0.0009	0.002	0.960
Energy	Stationary Combustion, BIOMASS	N ₂ O	0.038	0.091	0.0009	0.002	0.962
Energy	Stationary Combustion, BIOMASS	CH ₄	0.102	0.135	0.0008	0.002	0.964
Agriculture	4.B Manure Management	N ₂ O	0.600	0.391	0.0008	0.002	0.966
Agriculture	4.D1.3 N-fixing crops	N ₂ O	0.269	0.256	0.0008	0.002	0.968
Energy	Transport, Civil aviation	CO ₂	0.243	0.133	0.0007	0.002	0.970
Energy	Transport, Road transport	N ₂ O	0.091	0.116	0.0007	0.002	0.971
Energy	Stationary Combustion, LPG	CO ₂	0.184	0.091	0.0007	0.002	0.973
Industrial Proc.	2A2 Lime production	CO ₂	0.116	0.040	0.0007	0.002	0.975
Energy	1.B.2 Refinery processes	CH ₄	0.001	0.047	0.0007	0.002	0.976
Agriculture	4.D1.4 Crop Residue	N ₂ O	0.361	0.311	0.0006	0.001	0.977
Agriculture	4.D.2 Grassing animals	N ₂ O	0.334	0.211	0.0006	0.001	0.979
Industrial Proc.	2F Consumption of SF ₆	SF ₆	0.107	0.118	0.0005	0.001	0.980
Energy	Transport, Fisheries	CO ₂	0.591	0.479	0.0005	0.001	0.981
Energy	Transport, Residential	CO ₂	0.039	0.062	0.0005	0.001	0.982
Energy	1.B.2 Flaring off-shore	CO ₂	0.302	0.195	0.0005	0.001	0.983
Energy	Biogas fuelled engines, BIOMASS	CH ₄	0.001	0.031	0.0004	0.001	0.984
Waste	6 B. Wastewater Handling - Indirect	N ₂ O	0.082	0.032	0.0004	0.001	0.985
Energy	Stationary Combustion, Coke	CO ₂	0.138	0.074	0.0004	0.001	0.986
Energy	Transport, Railways	CO ₂	0.297	0.249	0.0004	0.001	0.987
Energy	Transport, Military	CO ₂	0.119	0.116	0.0004	0.001	0.988
Energy	1.B.2 Off-shore activities	CH ₄	0.015	0.037	0.0004	0.001	0.989
Waste	6 B. Wastewater Handling	CH ₄	0.065	0.074	0.0004	0.001	0.990
Energy	Transport, Road transport	CH ₄	0.047	0.011	0.0003	0.001	0.991
Waste	6 B. Wastewater Handling - Direct	N ₂ O	0.023	0.041	0.0003	0.001	0.991
Energy	Stationary Combustion, SOLID	N ₂ O	0.068	0.027	0.0003	0.001	0.992
Energy	Stationary Combustion, LIQUID	N ₂ O	0.044	0.011	0.0003	0.001	0.993
Industrial Proc.	2C1 Iron and steel production	CO ₂	0.028	0.000	0.0003	0.001	0.994
Agriculture	4.D1.5 Cultivation of histosols	N ₂ O	0.290	0.198	0.0003	0.001	0.994
Agriculture	4.D1.6 Sewage sludge and Industrial waste used as fertiliser	N ₂ O	0.028	0.039	0.0003	0.001	0.995
Industrial Proc.	2A3 Limestone and dolomite use	CO ₂	0.014	0.026	0.0002	0.001	0.995
Energy	Stationary Combustion, GAS	N ₂ O	0.016	0.024	0.0002	0.000	0.996
Energy	Stationary Combustion, WASTE	N ₂ O	0.007	0.016	0.0002	0.000	0.996

Tier 1 Analysis IPCC Source Categories (LULUCF excluded)		DK GHG	Base Year Estimate	Latest Year Estimate	Trend Assessment	Contribution on to Trend	Cumulative
			Ex,o Mt CO ₂ eq	Ex,t Mt CO ₂ eq	Tx,t		
Energy	Transport, Forestry	CO ₂	0.036	0.017	0.0001	0.000	0.997
Solvent and Other Prod. Use	3D1 Other - Use of N ₂ O for Anaesthesia	N ₂ O	0.000	0.009	0.0001	0.000	0.997
Industrial Proc.	2F Consumption of PFC	PFC	0.001	0.009	0.0001	0.000	0.997
Energy	Stationary Combustion, BKB	CO ₂	0.011	0.001	0.0001	0.000	0.997
Energy	Stationary Combustion, SOLID	CH ₄	0.013	0.003	0.0001	0.000	0.998
Energy	Transport, Agriculture	N ₂ O	0.015	0.018	0.0001	0.000	0.998
Industrial Proc.	2G Lubricants	CO ₂	0.050	0.032	0.0001	0.000	0.998
Energy	Transport, Industry (mobile)	N ₂ O	0.011	0.013	0.0001	0.000	0.998
Industrial Proc.	2A7c Expanded clay	CO ₂	0.015	0.006	0.0001	0.000	0.998
Energy	1.B.2 Flaring in refinery	CO ₂	0.023	0.022	0.0001	0.000	0.998
Energy	1.B.2 Land based activities	CH ₄	0.017	0.018	0.0001	0.000	0.999
Industrial Proc.	2A7a Glass and Glass wool	CO ₂	0.017	0.010	0.0000	0.000	0.999
Energy	Stationary Combustion, GAS	CH ₄	0.003	0.006	0.0000	0.000	0.999
Energy	Transport, Navigation (large vessels)	N ₂ O	0.015	0.008	0.0000	0.000	0.999
Energy	1.B.2 Transmission of natural gas	CH ₄	0.004	0.000	0.0000	0.000	0.999
Solvent and Other Prod. Use	3A Paint application	CO ₂	0.013	0.007	0.0000	0.000	0.999
Solvent and Other Prod. Use	3C Chemical products, manufacturing and processing	CO ₂	0.019	0.012	0.0000	0.000	0.999
Waste	6.D Accidental fires, buildings	CO ₂	0.011	0.011	0.0000	0.000	0.999
Energy	1.B.2 Off-shore activities	CO ₂	0.002	0.004	0.0000	0.000	0.999
Solvent and Other Prod. Use	3D5 Other	CO ₂	0.061	0.044	0.0000	0.000	0.999
Energy	Transport, Commercial/institutional	CH ₄	0.002	0.003	0.0000	0.000	0.999
Solvent and Other Prod. Use	3D5 Consumption of fireworks	N ₂ O	0.001	0.002	0.0000	0.000	1.000
Industrial Proc.	2D2 Food and Drink	CO ₂	0.004	0.002	0.0000	0.000	1.000
Agriculture	4F Field burning of agricultural residues	CH ₄	0.002	0.002	0.0000	0.000	1.000
Energy	1.B.2 Distribution of natural gas	CH ₄	0.005	0.003	0.0000	0.000	1.000
Waste	6.D Accidental fires, vehicles	CO ₂	0.006	0.006	0.0000	0.000	1.000
Energy	Stationary Combustion, WASTE	CH ₄	0.001	0.002	0.0000	0.000	1.000
Energy	Stationary Combustion, LIQUID	CH ₄	0.003	0.001	0.0000	0.000	1.000
Industrial Proc.	2B5 Catalysts/Fertilizers, Pesticides and Sulphuric acid	CO ₂	0.001	0.001	0.0000	0.000	1.000
Energy	Transport, Navigation (small boats)	N ₂ O	0.000	0.001	0.0000	0.000	1.000
Energy	Transport, Fisheries	N ₂ O	0.011	0.009	0.0000	0.000	1.000
Energy	Transport, Commercial/institutional	N ₂ O	0.000	0.001	0.0000	0.000	1.000
Energy	Transport, Residential	CH ₄	0.001	0.001	0.0000	0.000	1.000
Industrial Proc.	2A6 Road paving with asphalt	CO ₂	0.002	0.002	0.0000	0.000	1.000
Energy	Transport, Military	N ₂ O	0.001	0.001	0.0000	0.000	1.000
Agriculture	4.F Field Burning of Agricultural Residues	N ₂ O	0.001	0.001	0.0000	0.000	1.000
Energy	Transport, Agriculture	CH ₄	0.002	0.002	0.0000	0.000	1.000
Energy	Transport, Civil aviation	N ₂ O	0.003	0.002	0.0000	0.000	1.000
Energy	1.B.2 Venting in gas storage	CH ₄	0.001	0.001	0.0000	0.000	1.000
Waste	6.D Accidental fires, buildings	CH ₄	0.001	0.001	0.0000	0.000	1.000
Energy	Transport, Forestry	CH ₄	0.000	0.000	0.0000	0.000	1.000
Energy	Transport, Navigation (small boats)	CH ₄	0.000	0.001	0.0000	0.000	1.000
Industrial Proc.	2A7b Yellow bricks	CO ₂	0.023	0.017	0.0000	0.000	1.000
Energy	Transport, Railways	N ₂ O	0.003	0.002	0.0000	0.000	1.000
Energy	1.B.2. Flaring off-shore	CH ₄	0.001	0.000	0.0000	0.000	1.000
Energy	Transport, Residential	N ₂ O	0.000	0.000	0.0000	0.000	1.000
Energy	Transport, Industry (mobile)	CH ₄	0.001	0.001	0.0000	0.000	1.000
Solvent and Other Prod. Use	3D5 Use of candles	N ₂ O	0.000	0.000	0.0000	0.000	1.000
Solvent and Other Prod. Use	3D5 Consumption of fireworks	CO ₂	0.000	0.000	0.0000	0.000	1.000
Solvent and Other Prod. Use	3D5 Use of charcoal for BBQ	N ₂ O	0.000	0.000	0.0000	0.000	1.000
Waste	6.C Incineration of carcasses	N ₂ O	0.000	0.000	0.0000	0.000	1.000
Energy	1.B.2 Flaring off-shore	N ₂ O	0.001	0.000	0.0000	0.000	1.000
Energy	Transport, Navigation (large vessels)	CH ₄	0.000	0.000	0.0000	0.000	1.000
Waste	6.C Incineration of corpses	N ₂ O	0.000	0.000	0.0000	0.000	1.000

Tier 1 Analysis IPCC Source Categories (LULUCF excluded)		DK GHG	Base Year Estimate	Latest Year Estimate	Trend Assessment	Contribution to Trend	Cumulative
			Ex,o Mt CO ₂ eq	Ex,t Mt CO ₂ eq	Tx,t		
Waste	6.D Accidental fires, vehicles	CH ₄	0.000	0.000	0.0000	0.000	1.000
Energy	1.B.2. Flaring in refinery	CH ₄	0.000	0.000	0.0000	0.000	1.000
Energy	Transport, Fisheries	CH ₄	0.000	0.000	0.0000	0.000	1.000
Energy	Transport, Railways	CH ₄	0.000	0.000	0.0000	0.000	1.000
Energy	Transport, Forestry	N ₂ O	0.000	0.000	0.0000	0.000	1.000
Solvent and Other Prod. Use	3D5 Use of tobacco	N ₂ O	0.000	0.000	0.0000	0.000	1.000
Energy	Transport, Civil aviation	CH ₄	0.000	0.000	0.0000	0.000	1.000
Energy	Transport, Military	CH ₄	0.000	0.000	0.0000	0.000	1.000
Energy	1.B.2 Flaring in refinery	N ₂ O	0.000	0.000	0.0000	0.000	1.000
Industrial Proc.	2A5 Asphalt roofing	CO ₂	0.000	0.000	0.0000	0.000	1.000
Waste	6.C Incineration of carcasses	CH ₄	0.000	0.000	0.0000	0.000	1.000
Energy	1.B.2 Land based activities	CO ₂	0.000	0.000	0.0000	0.000	1.000
Waste	6.C Incineration of corpses	CH ₄	0.000	0.000	0.0000	0.000	1.000
Energy	1.B.2 Transmission of natural gas	CO ₂	0.000	0.000	0.0000	0.000	1.000
Energy	1.B.2 Distribution of natural gas	CO ₂	0.000	0.000	0.0000	0.000	1.000
Energy	1.B.2 Venting in gas storage	CO ₂	0.000	0.000	0.0000	0.000	1.000
Solvent and Other Prod. Use	3B Degreasing and dry cleaning	CO ₂	0.000	0.000	0.0000	0.000	1.000
Total			68.943	51.637			

Table A1-17 KCA for Denmark, trend assessment 1990-2012 incl. LULUCF, tier 1.

Tier 1 Analysis IPCC Source Categories (LULUCF included)		DK GHG	Base Year Estimate Ex,o Mt CO ₂ eq	Latest Year Estimate Ex,t Mt CO ₂ eq	Trend Asses- ment Tx,t	Contri- bution to Trend	Cumula- tive
Energy	Stationary Combustion, Coal	CO ₂	23.834	10.005	0.0848	0.180	0.180
Energy	Stationary Combustion, Natural gas	CO ₂	4.335	8.293	0.0716	0.152	0.333
Energy	Transport, Road transport	CO ₂	9.284	11.224	0.0655	0.139	0.472
LULUCF	5.A.1 Forest remaining forest	CO ₂	0.050	-4.491	0.0608	0.129	0.601
Energy	Stationary Combustion, Gas oil	CO ₂	4.547	0.742	0.0319	0.068	0.669
Energy	Stationary Combustion, Residual oil	CO ₂	2.496	0.571	0.0153	0.033	0.701
Energy	Stationary Combustion, Fossil waste	CO ₂	0.573	1.397	0.0135	0.029	0.730
Industrial Proc.	2B2 Nitric acid production	N ₂ O	1.043	0.000	0.0096	0.020	0.751
Agriculture	4A Enteric Fermentation	CH ₄	3.247	2.904	0.0092	0.019	0.770
Agriculture	4B Manure Management	CH ₄	0.985	1.297	0.0084	0.018	0.788
Industrial Proc.	2F Consumption of HFC	HFC	0.218	0.657	0.0068	0.015	0.802
Agriculture	4.D1.1 Synthetic Fertilizer	N ₂ O	2.354	1.103	0.0068	0.015	0.817
Energy	Transport, Agriculture	CO ₂	1.272	1.343	0.0064	0.014	0.830
Energy	Transport, Industry (mobile)	CO ₂	0.839	1.021	0.0060	0.013	0.843
LULUCF	5.C Grassland, Living biomass	CO ₂	0.075	0.463	0.0055	0.012	0.855
Agriculture	4.D1.2 Animal waste applied to soils	N ₂ O	1.112	1.161	0.0054	0.011	0.866
LULUCF	5.B Cropland, Mineral soils	CO ₂	1.415	0.577	0.0053	0.011	0.878
Energy	Stationary Combustion, Refinery gas	CO ₂	0.816	0.906	0.0047	0.010	0.888
Energy	Stationary Combustion, Petroleum coke	CO ₂	0.415	0.628	0.0046	0.010	0.897
LULUCF	5.B Cropland, Living biomass	CO ₂	-0.074	0.193	0.0039	0.008	0.906
Industrial Proc.	2A1 Cement production	CO ₂	0.882	0.871	0.0036	0.008	0.913
Energy	Stationary Combustion, Kerosene	CO ₂	0.366	0.002	0.0033	0.007	0.920
Agriculture	4.D3 Leaching	N ₂ O	2.447	1.430	0.0033	0.007	0.927
Waste	6 A. Solid Waste Disposal on Land	CH ₄	1.366	0.698	0.0032	0.007	0.934
LULUCF	5(IV) Cropland Limestone	CO ₂	0.623	0.192	0.0031	0.007	0.941
Energy	Transport, Commercial/institutional	CO ₂	0.074	0.171	0.0016	0.003	0.944
Waste	6.D Compost production	N ₂ O	0.013	0.127	0.0016	0.003	0.948
Energy	Natural gas fuelled engines, GAS	CH ₄	0.005	0.120	0.0016	0.003	0.951
Energy	Transport, Navigation (large vessels)	CO ₂	0.748	0.399	0.0015	0.003	0.954
Energy	Transport, Fisheries	CO ₂	0.591	0.479	0.0010	0.002	0.956
Agriculture	4.D1.3 N-fixing crops	N ₂ O	0.269	0.256	0.0010	0.002	0.958
Waste	6.D Compost production	CH ₄	0.029	0.090	0.0009	0.002	0.960
Solvent and Other Prod. Use	3D5 Use of candles	CO ₂	0.022	0.081	0.0009	0.002	0.962
Energy	Transport, Navigation (small boats)	CO ₂	0.048	0.099	0.0009	0.002	0.964
Energy	Stationary Combustion, BIOMASS	CH ₄	0.102	0.135	0.0009	0.002	0.966
Energy	Stationary Combustion, BIOMASS	N ₂ O	0.038	0.091	0.0009	0.002	0.968
Agriculture	4.D1.4 Crop Residue	N ₂ O	0.361	0.311	0.0009	0.002	0.970
LULUCF	5.D Wetlands, Soils	CO ₂	0.085	0.002	0.0007	0.002	0.971
Energy	Transport, Road transport	N ₂ O	0.091	0.116	0.0007	0.002	0.973
LULUCF	5.E Settlements, Living biomass	CO ₂	0.011	0.056	0.0006	0.001	0.974
Energy	Transport, Railways	CO ₂	0.297	0.249	0.0006	0.001	0.976
Energy	1.B.2 Refinery processes	CH ₄	0.001	0.047	0.0006	0.001	0.977
Agriculture	4.D3 Atmospheric deposition	N ₂ O	0.496	0.295	0.0006	0.001	0.978
Industrial Proc.	2F Consumption of SF ₆	SF ₆	0.107	0.118	0.0006	0.001	0.979
Industrial Proc.	2A2 Lime production	CO ₂	0.116	0.040	0.0005	0.001	0.981
Energy	Transport, Residential	CO ₂	0.039	0.062	0.0005	0.001	0.982
Energy	Stationary Combustion, LPG	CO ₂	0.184	0.091	0.0005	0.001	0.983
Energy	Transport, Military	CO ₂	0.119	0.116	0.0005	0.001	0.984
LULUCF	5.E Settlements, Soils	CO ₂	0.001	0.035	0.0005	0.001	0.985
Energy	Transport, Civil aviation	CO ₂	0.243	0.133	0.0005	0.001	0.986
Energy	Biogas fuelled engines, BIOMASS	CH ₄	0.001	0.031	0.0004	0.001	0.986
Waste	6 B. Wastewater Handling	CH ₄	0.065	0.074	0.0004	0.001	0.987
Energy	1.B.2 Off-shore activities	CH ₄	0.015	0.037	0.0004	0.001	0.988
Waste	6 B. Wastewater Handling - Direct	N ₂ O	0.023	0.041	0.0003	0.001	0.989
Waste	6 B. Wastewater Handling - Indirect	N ₂ O	0.082	0.032	0.0003	0.001	0.989
Energy	Transport, Road transport	CH ₄	0.047	0.011	0.0003	0.001	0.990
Energy	Stationary Combustion, Coke	CO ₂	0.138	0.074	0.0003	0.001	0.991
Agriculture	4.B Manure Management	N ₂ O	0.600	0.391	0.0003	0.001	0.991
Agriculture	4.D1.6 Sewage sludge and Industrial waste used as fertiliser	N ₂ O	0.028	0.039	0.0003	0.001	0.992
Industrial Proc.	2C1 Iron and steel production	CO ₂	0.028	0.000	0.0003	0.001	0.992
Energy	Stationary Combustion, SOLID	N ₂ O	0.068	0.027	0.0003	0.001	0.993

Tier 1 Analysis IPCC Source Categories (LULUCF included)		DK GHG	Base Year Estimate Ex,o Mt CO ₂ eq	Latest Year Estimate Ex,t Mt CO ₂ eq	Trend Asses- ment Tx,t	Contri- bution to Trend	Cumula- tive
Energy	Stationary Combustion, LIQUID	N ₂ O	0.044	0.011	0.0003	0.001	0.993
Agriculture	4.D.2 Grassing animals	N ₂ O	0.334	0.211	0.0002	0.001	0.994
Industrial Proc.	2A3 Limestone and dolomite use	CO ₂	0.014	0.026	0.0002	0.000	0.994
LULUCF	5.A.2 Land converted to forest	CO ₂	0.077	0.038	0.0002	0.000	0.995
Energy	Stationary Combustion, GAS	N ₂ O	0.016	0.024	0.0002	0.000	0.995
Energy	1.B.2 Flaring off-shore	CO ₂	0.302	0.195	0.0002	0.000	0.995
Energy	Stationary Combustion, WASTE	N ₂ O	0.007	0.016	0.0002	0.000	0.996
LULUCF	5.B Cropland, Dead organic matter	CO ₂	0.003	0.012	0.0001	0.000	0.996
Solvent and Other Prod. Use	3D1 Other - Use of N2O for Anaesthesia	N ₂ O	0.000	0.009	0.0001	0.000	0.996
Industrial Proc.	2F Consumption of PFC	PFC	0.001	0.009	0.0001	0.000	0.997
Energy	Transport, Forestry	CO ₂	0.036	0.017	0.0001	0.000	0.997
Energy	Transport, Agriculture	N ₂ O	0.015	0.018	0.0001	0.000	0.997
Energy	1.B.2 Flaring in refinery	CO ₂	0.023	0.022	0.0001	0.000	0.997
Energy	Transport, Industry (mobile)	N ₂ O	0.011	0.013	0.0001	0.000	0.997
Energy	Stationary Combustion, BKB	CO ₂	0.011	0.001	0.0001	0.000	0.998
LULUCF	5.C Grassland, Dead organic matter	CO ₂	0.002	0.007	0.0001	0.000	0.998
Energy	1.B.2 Land based activities	CH ₄	0.017	0.018	0.0001	0.000	0.998
LULUCF	5.C Grassland, Organic soils	CO ₂	0.107	0.079	0.0001	0.000	0.998
Energy	Stationary Combustion, SOLID	CH ₄	0.013	0.003	0.0001	0.000	0.998
LULUCF	5.B Cropland, Organic soils	CO ₂	2.887	1.981	0.0001	0.000	0.998
LULUCF	5.C Grassland, Mineral soils	CO ₂	0.000	0.005	0.0001	0.000	0.998
Industrial Proc.	2A7c Expanded clay	CO ₂	0.015	0.006	0.0001	0.000	0.999
Energy	Stationary Combustion, GAS	CH ₄	0.003	0.006	0.0000	0.000	0.999
Waste	6.D Accidental fires, buildings	CO ₂	0.011	0.011	0.0000	0.000	0.999
Energy	1.B.2 Transmission of natural gas	CH ₄	0.004	0.000	0.0000	0.000	0.999
Industrial Proc.	2G Lubricants	CO ₂	0.050	0.032	0.0000	0.000	0.999
Solvent and Other Prod. Use	3D5 Other	CO ₂	0.061	0.044	0.0000	0.000	0.999
Industrial Proc.	2A7a Glass and Glass wool	CO ₂	0.017	0.010	0.0000	0.000	0.999
Energy	Transport, Navigation (large vessels)	N ₂ O	0.015	0.008	0.0000	0.000	0.999
Energy	1.B.2 Off-shore activities	CO ₂	0.002	0.004	0.0000	0.000	0.999
LULUCF	5.D Wetlands, Living biomass	CO ₂	0.003	0.000	0.0000	0.000	0.999
Solvent and Other Prod. Use	3A Paint application	CO ₂	0.013	0.007	0.0000	0.000	0.999
Energy	Transport, Commercial/institutional	CH ₄	0.002	0.003	0.0000	0.000	0.999
Industrial Proc.	2A7b Yellow bricks	CO ₂	0.023	0.017	0.0000	0.000	0.999
Solvent and Other Prod. Use	3D5 Consumption of fireworks	N ₂ O	0.001	0.002	0.0000	0.000	0.999
Energy	Transport, Fisheries	N ₂ O	0.011	0.009	0.0000	0.000	0.999
Waste	6.D Accidental fires, vehicles	CO ₂	0.006	0.006	0.0000	0.000	1.000
LULUCF	5(II) Forest Land.	N ₂ O	0.016	0.012	0.0000	0.000	1.000
Agriculture	4F Field burning of agricultural residues	CH ₄	0.002	0.002	0.0000	0.000	1.000
Solvent and Other Prod. Use	3C Chemical products, manufacturing and processing	CO ₂	0.019	0.012	0.0000	0.000	1.000
Energy	Stationary Combustion, WASTE	CH ₄	0.001	0.002	0.0000	0.000	1.000
Industrial Proc.	2B5 Catalysts/Fertilizers, Pesticides and Sulphuric acid	CO ₂	0.001	0.001	0.0000	0.000	1.000
LULUCF	5(III) Disturbance, Land converted to cropland	N ₂ O	0.000	0.001	0.0000	0.000	1.000
Energy	Transport, Navigation (small boats)	N ₂ O	0.000	0.001	0.0000	0.000	1.000
Industrial Proc.	2D2 Food and Drink	CO ₂	0.004	0.002	0.0000	0.000	1.000
Energy	Stationary Combustion, LIQUID	CH ₄	0.003	0.001	0.0000	0.000	1.000
Energy	1.B.2 Distribution of natural gas	CH ₄	0.005	0.003	0.0000	0.000	1.000
Energy	Transport, Residential	CH ₄	0.001	0.001	0.0000	0.000	1.000
Energy	Transport, Commercial/institutional	N ₂ O	0.000	0.001	0.0000	0.000	1.000
Industrial Proc.	2A6 Road paving with asphalt	CO ₂	0.002	0.002	0.0000	0.000	1.000
Energy	Transport, Agriculture	CH ₄	0.002	0.002	0.0000	0.000	1.000
Energy	Transport, Military	N ₂ O	0.001	0.001	0.0000	0.000	1.000
Agriculture	4.F Field Burning of Agricultural Residues	N ₂ O	0.001	0.001	0.0000	0.000	1.000
Energy	Transport, Railways	N ₂ O	0.003	0.002	0.0000	0.000	1.000
Waste	6.D Accidental fires, buildings	CH ₄	0.001	0.001	0.0000	0.000	1.000
LULUCF	5.D Wetlands, Dead organic matter	CO ₂	0.001	0.000	0.0000	0.000	1.000
Energy	1.B.2 Venting in gas storage	CH ₄	0.001	0.001	0.0000	0.000	1.000

Tier 1 Analysis IPCC Source Categories (LULUCF included)		DK GHG	Base Year Estimate Ex,o Mt CO ₂ eq	Latest Year Estimate Ex,t Mt CO ₂ eq	Trend Asses- sment Tx,t	Contri- bution to Trend	Cumula- tive
LULUCF	5(V) Biomass Burning	CH ₄	0.001	0.000	0.0000	0.000	1.000
LULUCF	5.E Settlements, Dead organic matter	CO ₂	0.001	0.000	0.0000	0.000	1.000
Agriculture	4.D1.5 Cultivation of histosols	N ₂ O	0.290	0.198	0.0000	0.000	1.000
LULUCF	5(V) Biomass Burning	N ₂ O	0.000	0.000	0.0000	0.000	1.000
Energy	Transport, Navigation (small boats)	CH ₄	0.000	0.001	0.0000	0.000	1.000
Energy	Transport, Forestry	CH ₄	0.000	0.000	0.0000	0.000	1.000
Energy	Transport, Residential	N ₂ O	0.000	0.000	0.0000	0.000	1.000
Energy	1.B.2. Flaring off-shore	CH ₄	0.001	0.000	0.0000	0.000	1.000
Solvent and Other Prod. Use	3D5 Use of candles	N ₂ O	0.000	0.000	0.0000	0.000	1.000
Energy	Transport, Civil aviation	N ₂ O	0.003	0.002	0.0000	0.000	1.000
Solvent and Other Prod. Use	3D5 Consumption of fireworks	CO ₂	0.000	0.000	0.0000	0.000	1.000
Energy	Transport, Industry (mobile)	CH ₄	0.001	0.001	0.0000	0.000	1.000
Solvent and Other Prod. Use	3D5 Use of charcoal for BBQ	N ₂ O	0.000	0.000	0.0000	0.000	1.000
Waste	6.C Incineration of carcasses	N ₂ O	0.000	0.000	0.0000	0.000	1.000
Waste	6.D Accidental fires, vehicles	CH ₄	0.000	0.000	0.0000	0.000	1.000
Energy	Transport, Fisheries	CH ₄	0.000	0.000	0.0000	0.000	1.000
Waste	6.C Incineration of corpses	N ₂ O	0.000	0.000	0.0000	0.000	1.000
Energy	1.B.2. Flaring in refinery	CH ₄	0.000	0.000	0.0000	0.000	1.000
Energy	Transport, Forestry	N ₂ O	0.000	0.000	0.0000	0.000	1.000
LULUCF	5(II) Wetlands	N ₂ O	0.000	0.000	0.0000	0.000	1.000
Energy	Transport, Navigation (large vessels)	CH ₄	0.000	0.000	0.0000	0.000	1.000
Energy	Transport, Railways	CH ₄	0.000	0.000	0.0000	0.000	1.000
Energy	1.B.2 Flaring in refinery	N ₂ O	0.000	0.000	0.0000	0.000	1.000
Energy	1.B.2 Flaring off-shore	N ₂ O	0.001	0.000	0.0000	0.000	1.000
Energy	Transport, Civil aviation	CH ₄	0.000	0.000	0.0000	0.000	1.000
Energy	Transport, Military	CH ₄	0.000	0.000	0.0000	0.000	1.000
Solvent and Other Prod. Use	3D5 Use of tobacco	N ₂ O	0.000	0.000	0.0000	0.000	1.000
Industrial Proc.	2A5 Asphalt roofing	CO ₂	0.000	0.000	0.0000	0.000	1.000
Waste	6.C Incineration of carcasses	CH ₄	0.000	0.000	0.0000	0.000	1.000
Waste	6.C Incineration of corpses	CH ₄	0.000	0.000	0.0000	0.000	1.000
Energy	1.B.2 Land based activities	CO ₂	0.000	0.000	0.0000	0.000	1.000
Energy	1.B.2 Transmission of natural gas	CO ₂	0.000	0.000	0.0000	0.000	1.000
Energy	1.B.2 Distribution of natural gas	CO ₂	0.000	0.000	0.0000	0.000	1.000
Energy	1.B.2 Venting in gas storage	CO ₂	0.000	0.000	0.0000	0.000	1.000
Solvent and Other Prod. Use	3B Degreasing and dry cleaning	CO ₂	0.000	0.000	0.0000	0.000	1.000
Total			74.225	50.799			

Table A1-18 KCA for Denmark, level assessment base year excl. LULUCF, tier 2.

Tier 2 Analysis IPCC Source Categories (LULUCF excluded)		DK GHG	Base Year Estimate	Base Year Level Assessment	Base Year Cumulative Total of Lx,o
			Ex,o Mt CO ₂ eq	Lx,o	
Agriculture	4.D3 Leaching	N ₂ O	2.496	0.157	0.157
Agriculture	4.D1.1 Synthetic Fertilizer	N ₂ O	2.427	0.153	0.310
Waste	6 A. Solid Waste Disposal on Land	CH ₄	1.616	0.102	0.412
Agriculture	4.D1.2 Animal waste applied to soils	N ₂ O	1.161	0.073	0.485
Agriculture	4A Enteric Fermentation	CH ₄	0.653	0.041	0.526
Agriculture	4.B Manure Management	N ₂ O	0.615	0.039	0.564
Agriculture	4.D3 Atmospheric deposition	N ₂ O	0.505	0.032	0.596
Energy	Transport, Road transport	CO ₂	0.500	0.031	0.628
Energy	Stationary Combustion, LIQUID	N ₂ O	0.440	0.028	0.655
Energy	Stationary Combustion, BIOMASS	N ₂ O	0.379	0.024	0.679
Agriculture	4.D1.4 Crop Residue	N ₂ O	0.368	0.023	0.702
Energy	Transport, Industry (mobile)	CO ₂	0.347	0.022	0.724
Agriculture	4.D.2 Grassing animals	N ₂ O	0.345	0.022	0.746
Energy	Transport, Agriculture	CO ₂	0.312	0.020	0.766
Agriculture	4.D1.5 Cultivation of histosols	N ₂ O	0.296	0.019	0.784
Agriculture	4.D1.3 N-fixing crops	N ₂ O	0.275	0.017	0.802
Energy	Stationary Combustion, SOLID	N ₂ O	0.272	0.017	0.819
Industrial Proc.	2B2 Nitric acid production	N ₂ O	0.262	0.016	0.835
Energy	Stationary Combustion, Coal	CO ₂	0.245	0.015	0.851
Energy	Stationary Combustion, Gas oil	CO ₂	0.209	0.013	0.864
Agriculture	4B Manure Management	CH ₄	0.203	0.013	0.877
Energy	Transport, Agriculture	N ₂ O	0.153	0.010	0.886
Energy	Transport, Navigation (large vessels)	N ₂ O	0.146	0.009	0.895
Energy	Stationary Combustion, GAS	N ₂ O	0.123	0.008	0.903
Energy	Transport, Fisheries	N ₂ O	0.115	0.007	0.910
Industrial Proc.	2F Consumption of HFC	HFC	0.111	0.007	0.917
Energy	Transport, Industry (mobile)	N ₂ O	0.106	0.007	0.924
Energy	Stationary Combustion, BIOMASS	CH ₄	0.103	0.006	0.930
Energy	Transport, Navigation (large vessels)	CO ₂	0.090	0.006	0.936
Energy	Stationary Combustion, Fossil waste	CO ₂	0.064	0.004	0.940
Waste	6 B. Wastewater Handling - Indirect	N ₂ O	0.060	0.004	0.944
Energy	Stationary Combustion, Residual oil	CO ₂	0.058	0.004	0.948
Industrial Proc.	2F Consumption of SF ₆	SF ₆	0.055	0.003	0.951
Energy	Stationary Combustion, Natural gas	CO ₂	0.048	0.003	0.954
Energy	Transport, Road transport	N ₂ O	0.045	0.003	0.957
Waste	6.D Accidental fires, buildings	CO ₂	0.034	0.002	0.959
Energy	Transport, Civil aviation	N ₂ O	0.032	0.002	0.961
Energy	Transport, Fisheries	CO ₂	0.032	0.002	0.963
Waste	6.D Compost production	CH ₄	0.031	0.002	0.965
Waste	6.D Accidental fires, vehicles	CO ₂	0.031	0.002	0.967
Agriculture	4.D1.6 Sewage sludge and Industrial waste used as fertiliser	N ₂ O	0.028	0.002	0.969
Energy	Transport, Civil aviation	CO ₂	0.027	0.002	0.971
Energy	Stationary Combustion, WASTE	N ₂ O	0.026	0.002	0.972
Energy	Transport, Commercial/institutional	CO ₂	0.026	0.002	0.974
Waste	6 B. Wastewater Handling	CH ₄	0.026	0.002	0.975
Energy	Transport, Railways	N ₂ O	0.025	0.002	0.977
Energy	1.B.2 Flaring off-shore	CO ₂	0.023	0.001	0.979
Energy	Stationary Combustion, Petroleum coke	CO ₂	0.022	0.001	0.980
Energy	Transport, Navigation (small boats)	CO ₂	0.020	0.001	0.981
Industrial Proc.	2A1 Cement production	CO ₂	0.020	0.001	0.982
Energy	Stationary Combustion, Kerosene	CO ₂	0.020	0.001	0.984
Energy	Transport, Road transport	CH ₄	0.019	0.001	0.985
Energy	Stationary Combustion, Refinery gas	CO ₂	0.018	0.001	0.986
Energy	Transport, Railways	CO ₂	0.016	0.001	0.987
Waste	6.D Compost production	N ₂ O	0.014	0.001	0.988
Energy	Transport, Residential	CO ₂	0.014	0.001	0.989
Solvent and Other Prod. Use	3D5 Other	CO ₂	0.014	0.001	0.990
Energy	Stationary Combustion, SOLID	CH ₄	0.013	0.001	0.990
Waste	6 B. Wastewater Handling - Direct	N ₂ O	0.012	0.001	0.991
Energy	Transport, Military	N ₂ O	0.011	0.001	0.992
Energy	Transport, Forestry	CO ₂	0.011	0.001	0.993

Tier 2 Analysis IPCC Source Categories (LULUCF excluded)		DK GHG	Base Year Estimate	Base Year Level Assessment	Base Year Cumulative Total of Lx,o
			Ex,o Mt CO ₂ eq	Lx,o	
Energy	Stationary Combustion, LPG	CO ₂	0.010	0.001	0.993
Industrial Proc.	2A2 Lime production	CO ₂	0.008	0.001	0.994
Energy	Stationary Combustion, Coke	CO ₂	0.007	0.000	0.994
Energy	1.B.2 Flaring off-shore	N ₂ O	0.007	0.000	0.995
Energy	1.B.2 Land based activities	CH ₄	0.007	0.000	0.995
Waste	6.D Accidental fires, buildings	CH ₄	0.007	0.000	0.996
Energy	Transport, Military	CO ₂	0.006	0.000	0.996
Solvent and Other Prod. Use	3D5 Use of candles	CO ₂	0.005	0.000	0.996
Energy	1.B.2 Off-shore activities	CH ₄	0.004	0.000	0.996
Energy	Transport, Navigation (small boats)	N ₂ O	0.004	0.000	0.997
Solvent and Other Prod. Use	3C Chemical products, manufacturing and processing	CO ₂	0.004	0.000	0.997
Energy	Transport, Commercial/institutional	N ₂ O	0.003	0.000	0.997
Energy	Stationary Combustion, GAS	CH ₄	0.003	0.000	0.997
Energy	Stationary Combustion, LIQUID	CH ₄	0.003	0.000	0.998
Industrial Proc.	2G Lubricants	CO ₂	0.003	0.000	0.998
Energy	1.B.2 Flaring in refinery	CO ₂	0.003	0.000	0.998
Solvent and Other Prod. Use	3A Paint application	CO ₂	0.002	0.000	0.998
Energy	Transport, Agriculture	CH ₄	0.002	0.000	0.998
Energy	Transport, Commercial/institutional	CH ₄	0.002	0.000	0.998
Industrial Proc.	2C1 Iron and steel production	CO ₂	0.002	0.000	0.998
Energy	Transport, Residential	N ₂ O	0.002	0.000	0.999
Waste	6.D Accidental fires, vehicles	CH ₄	0.002	0.000	0.999
Energy	Transport, Forestry	N ₂ O	0.002	0.000	0.999
Energy	1.B.2 Distribution of natural gas	CH ₄	0.001	0.000	0.999
Energy	Transport, Industry (mobile)	CH ₄	0.001	0.000	0.999
Industrial Proc.	2A7b Yellow bricks	CO ₂	0.001	0.000	0.999
Energy	Transport, Residential	CH ₄	0.001	0.000	0.999
Agriculture	4F Field burning of agricultural residues	CH ₄	0.001	0.000	0.999
Energy	1.B.2 Refinery processes	CH ₄	0.001	0.000	0.999
Industrial Proc.	2A3 Limestone and dolomite use	CO ₂	0.001	0.000	0.999
Industrial Proc.	2A7a Glass and Glass wool	CO ₂	0.001	0.000	0.999
Energy	1.B.2. Flaring off-shore	CH ₄	0.001	0.000	0.999
Industrial Proc.	2A7c Expanded clay	CO ₂	0.001	0.000	0.999
Energy	Stationary Combustion, WASTE	CH ₄	0.001	0.000	1.000
Solvent and Other Prod. Use	3D5 Consumption of fireworks	N ₂ O	0.001	0.000	1.000
Energy	1.B.2 Off-shore activities	CO ₂	0.001	0.000	1.000
Energy	Stationary Combustion, BKB	CO ₂	0.001	0.000	1.000
Energy	1.B.2 Transmission of natural gas	CH ₄	0.001	0.000	1.000
Energy	1.B.2 Flaring in refinery	N ₂ O	0.001	0.000	1.000
Energy	Transport, Forestry	CH ₄	0.000	0.000	1.000
Industrial Proc.	2A6 Road paving with asphalt	CO ₂	0.000	0.000	1.000
Agriculture	4.F Field Burning of Agricultural Residues	N ₂ O	0.000	0.000	1.000
Energy	Transport, Navigation (small boats)	CH ₄	0.000	0.000	1.000
Energy	Transport, Navigation (large vessels)	CH ₄	0.000	0.000	1.000
Industrial Proc.	2D2 Food and Drink	CO ₂	0.000	0.000	1.000
Waste	6.C Incineration of corpses	N ₂ O	0.000	0.000	1.000
Energy	Transport, Fisheries	CH ₄	0.000	0.000	1.000
Energy	Transport, Railways	CH ₄	0.000	0.000	1.000
Industrial Proc.	2F Consumption of PFC	PFC	0.000	0.000	1.000
Energy	1.B.2 Venting in gas storage	CH ₄	0.000	0.000	1.000
Energy	Biogas fuelled engines, BIOMASS	CH ₄	0.000	0.000	1.000
Energy	Transport, Military	CH ₄	0.000	0.000	1.000
Energy	Natural gas fuelled engines, GAS	CH ₄	0.000	0.000	1.000
Energy	Transport, Civil aviation	CH ₄	0.000	0.000	1.000
Solvent and Other Prod. Use	3D5 Use of tobacco	N ₂ O	0.000	0.000	1.000
Solvent and Other Prod. Use	3D5 Use of charcoal for BBQ	N ₂ O	0.000	0.000	1.000
Industrial Proc.	2B5 Catalysts/Fertilizers, Pesticides and Sulphuric acid	CO ₂	0.000	0.000	1.000
Solvent and Other Prod. Use	3D5 Consumption of fireworks	CO ₂	0.000	0.000	1.000
Energy	1.B.2. Flaring in refinery	CH ₄	0.000	0.000	1.000
Waste	6.C Incineration of carcasses	N ₂ O	0.000	0.000	1.000
Waste	6.C Incineration of corpses	CH ₄	0.000	0.000	1.000
Solvent and Other Prod. Use	3D5 Use of candles	N ₂ O	0.000	0.000	1.000

Tier 2 Analysis IPCC Source Categories (LULUCF excluded)		DK GHG	Base Year Estimate	Base Year Level Assessment	Base Year Cumulative Total of Lx,o
			Ex,o Mt CO ₂ eq	Lx,o	
Industrial Proc.	2A5 Asphalt roofing	CO ₂	0.000	0.000	1.000
Energy	1.B.2 Land based activities	CO ₂	0.000	0.000	1.000
Energy	1.B.2 Distribution of natural gas	CO ₂	0.000	0.000	1.000
Waste	6.C Incineration of carcasses	CH ₄	0.000	0.000	1.000
Energy	1.B.2 Transmission of natural gas	CO ₂	0.000	0.000	1.000
Energy	1.B.2 Venting in gas storage	CO ₂	0.000	0.000	1.000
Solvent and Other Prod. Use	3B Degreasing and dry cleaning	CO ₂	0.000	0.000	1.000
Solvent and Other Prod. Use	3D1 Other - Use of N ₂ O for Anaesthesia	N ₂ O	0.000	0.000	1.000
Total			15.886	1.000	

Table A1-19 KCA for Denmark, level assessment base year incl. LULUCF, tier 2.

Tier 2 Analysis IPCC Source Categories (LU- LUCF included)		DK GHG	Base Year Estimate Ex,o Mt CO ₂ eq	Base Year Level Assessment Lx,o	Base Year Cumulative Total of Lx,o
LULUCF	5.B Cropland, Organic soils	CO ₂	2.615	0.130	0.130
Agriculture	4.D3 Leaching	N ₂ O	2.496	0.124	0.253
Agriculture	4.D1.1 Synthetic Fertilizer	N ₂ O	2.427	0.120	0.374
Waste	6 A. Solid Waste Disposal on Land	CH ₄	1.616	0.080	0.454
Agriculture	4.D1.2 Animal waste applied to soils	N ₂ O	1.161	0.058	0.511
LULUCF	5.B Cropland, Mineral soils	CO ₂	1.071	0.053	0.564
Agriculture	4A Enteric Fermentation	CH ₄	0.653	0.032	0.597
Agriculture	4.B Manure Management	N ₂ O	0.615	0.030	0.627
Agriculture	4.D3 Atmospheric deposition	N ₂ O	0.505	0.025	0.652
Energy	Transport, Road transport	CO ₂	0.500	0.025	0.677
Energy	Stationary Combustion, LIQUID	N ₂ O	0.440	0.022	0.699
Energy	Stationary Combustion, BIOMASS	N ₂ O	0.379	0.019	0.717
Agriculture	4.D1.4 Crop Residue	N ₂ O	0.368	0.018	0.736
Energy	Transport, Industry (mobile)	CO ₂	0.347	0.017	0.753
Agriculture	4.D.2 Grassing animals	N ₂ O	0.345	0.017	0.770
LULUCF	5(IV) Cropland Limestone	CO ₂	0.313	0.016	0.785
Energy	Transport, Agriculture	CO ₂	0.312	0.015	0.801
Agriculture	4.D1.5 Cultivation of histosols	N ₂ O	0.296	0.015	0.816
Agriculture	4.D1.3 N-fixing crops	N ₂ O	0.275	0.014	0.829
Energy	Stationary Combustion, SOLID	N ₂ O	0.272	0.014	0.843
Industrial Proc.	2B2 Nitric acid production	N ₂ O	0.262	0.013	0.856
Energy	Stationary Combustion, Coal	CO ₂	0.245	0.012	0.868
Energy	Stationary Combustion, Gas oil	CO ₂	0.209	0.010	0.878
Agriculture	4B Manure Management	CH ₄	0.203	0.010	0.888
Energy	Transport, Agriculture	N ₂ O	0.153	0.008	0.896
Energy	Transport, Navigation (large vessels)	N ₂ O	0.146	0.007	0.903
Energy	Stationary Combustion, GAS	N ₂ O	0.123	0.006	0.909
Energy	Transport, Fisheries	N ₂ O	0.115	0.006	0.915
Industrial Proc.	2F Consumption of HFC	HFC	0.111	0.006	0.920
Energy	Transport, Industry (mobile)	N ₂ O	0.106	0.005	0.926
Energy	Stationary Combustion, BIOMASS	CH ₄	0.103	0.005	0.931
LULUCF	5.C Grassland, Organic soils	CO ₂	0.097	0.005	0.935
Energy	Transport, Navigation (large vessels)	CO ₂	0.090	0.004	0.940
LULUCF	5.D Wetlands, Soils	CO ₂	0.085	0.004	0.944
Energy	Stationary Combustion, Fossil waste	CO ₂	0.064	0.003	0.947
Waste	6 B. Wastewater Handling - Indirect	N ₂ O	0.060	0.003	0.950
Energy	Stationary Combustion, Residual oil	CO ₂	0.058	0.003	0.953
Industrial Proc.	2F Consumption of SF ₆	SF ₆	0.055	0.003	0.956
Energy	Stationary Combustion, Natural gas	CO ₂	0.048	0.002	0.958
Energy	Transport, Road transport	N ₂ O	0.045	0.002	0.961
LULUCF	5.C Grassland, Living biomass	CO ₂	0.038	0.002	0.962
LULUCF	5.B Cropland, Living biomass	CO ₂	0.037	0.002	0.964
Waste	6.D Accidental fires, buildings	CO ₂	0.034	0.002	0.966
Energy	Transport, Civil aviation	N ₂ O	0.032	0.002	0.968
Energy	Transport, Fisheries	CO ₂	0.032	0.002	0.969
Waste	6.D Compost production	CH ₄	0.031	0.002	0.971
Waste	6.D Accidental fires, vehicles	CO ₂	0.031	0.002	0.972
Agriculture	4.D1.6 Sewage sludge and Industrial waste used as fertiliser	N ₂ O	0.028	0.001	0.974
Energy	Transport, Civil aviation	CO ₂	0.027	0.001	0.975
Energy	Stationary Combustion, WASTE	N ₂ O	0.026	0.001	0.976
Energy	Transport, Commercial/institutional	CO ₂	0.026	0.001	0.978
Waste	6 B. Wastewater Handling	CH ₄	0.026	0.001	0.979
Energy	Transport, Railways	N ₂ O	0.025	0.001	0.980
Energy	1.B.2 Flaring off-shore	CO ₂	0.023	0.001	0.981
Energy	Stationary Combustion, Petroleum coke	CO ₂	0.022	0.001	0.982
Energy	Transport, Navigation (small boats)	CO ₂	0.020	0.001	0.983
Industrial Proc.	2A1 Cement production	CO ₂	0.020	0.001	0.984
Energy	Stationary Combustion, Kerosene	CO ₂	0.020	0.001	0.985
Energy	Transport, Road transport	CH ₄	0.019	0.001	0.986
Energy	Stationary Combustion, Refinery gas	CO ₂	0.018	0.001	0.987
Energy	Transport, Railways	CO ₂	0.016	0.001	0.988
Waste	6.D Compost production	N ₂ O	0.014	0.001	0.989

Tier 2 Analysis IPCC Source Categories (LU- LUCF included)		DK GHG	Base Year Estimate	Base Year Level Assessment Lx,o	Base Year Cumulative Total of Lx,o
			Ex,o Mt CO ₂ eq		
Energy	Transport, Residential	CO ₂	0.014	0.001	0.989
Solvent and Other Prod. Use	3D5 Other	CO ₂	0.014	0.001	0.990
LULUCF	5.A.2 Land converted to forest	CO ₂	0.013	0.001	0.991
Energy	Stationary Combustion, SOLID	CH ₄	0.013	0.001	0.991
Waste	6 B. Wastewater Handling - Direct	N ₂ O	0.012	0.001	0.992
Energy	Transport, Military	N ₂ O	0.011	0.001	0.992
Energy	Transport, Forestry	CO ₂	0.011	0.001	0.993
Energy	Stationary Combustion, LPG	CO ₂	0.010	0.000	0.993
Industrial Proc.	2A2 Lime production	CO ₂	0.008	0.000	0.994
LULUCF	5.A.1 Forest remaining forest	CO ₂	0.008	0.000	0.994
Energy	Stationary Combustion, Coke	CO ₂	0.007	0.000	0.995
Energy	1.B.2 Flaring off-shore	N ₂ O	0.007	0.000	0.995
Energy	1.B.2 Land based activities	CH ₄	0.007	0.000	0.995
Waste	6.D Accidental fires, buildings	CH ₄	0.007	0.000	0.996
Energy	Transport, Military	CO ₂	0.006	0.000	0.996
LULUCF	5.E Settlements, Living biomass	CO ₂	0.006	0.000	0.996
LULUCF	5(II) Forest Land.	N ₂ O	0.005	0.000	0.996
Solvent and Other Prod. Use	3D5 Use of candles	CO ₂	0.005	0.000	0.997
Energy	1.B.2 Off-shore activities	CH ₄	0.004	0.000	0.997
Energy	Transport, Navigation (small boats)	N ₂ O	0.004	0.000	0.997
Solvent and Other Prod. Use	3C Chemical products, manufacturing and processing	CO ₂	0.004	0.000	0.997
Energy	Transport, Commercial/institutional	N ₂ O	0.003	0.000	0.997
Energy	Stationary Combustion, GAS	CH ₄	0.003	0.000	0.998
Energy	Stationary Combustion, LIQUID	CH ₄	0.003	0.000	0.998
Industrial Proc.	2G Lubricants	CO ₂	0.003	0.000	0.998
Energy	1.B.2 Flaring in refinery	CO ₂	0.003	0.000	0.998
Solvent and Other Prod. Use	3A Paint application	CO ₂	0.002	0.000	0.998
Energy	Transport, Agriculture	CH ₄	0.002	0.000	0.998
Energy	Transport, Commercial/institutional	CH ₄	0.002	0.000	0.998
Industrial Proc.	2C1 Iron and steel production	CO ₂	0.002	0.000	0.998
Energy	Transport, Residential	N ₂ O	0.002	0.000	0.999
Waste	6.D Accidental fires, vehicles	CH ₄	0.002	0.000	0.999
Energy	Transport, Forestry	N ₂ O	0.002	0.000	0.999
LULUCF	5.B Cropland, Dead organic matter	CO ₂	0.001	0.000	0.999
Energy	1.B.2 Distribution of natural gas	CH ₄	0.001	0.000	0.999
Energy	Transport, Industry (mobile)	CH ₄	0.001	0.000	0.999
LULUCF	5.D Wetlands, Living biomass	CO ₂	0.001	0.000	0.999
Industrial Proc.	2A7b Yellow bricks	CO ₂	0.001	0.000	0.999
Energy	Transport, Residential	CH ₄	0.001	0.000	0.999
Agriculture	4F Field burning of agricultural residues	CH ₄	0.001	0.000	0.999
Energy	1.B.2 Refinery processes	CH ₄	0.001	0.000	0.999
Industrial Proc.	2A3 Limestone and dolomite use	CO ₂	0.001	0.000	0.999
Industrial Proc.	2A7a Glass and Glass wool	CO ₂	0.001	0.000	0.999
Energy	1.B.2. Flaring off-shore	CH ₄	0.001	0.000	0.999
LULUCF	5.C Grassland, Dead organic matter	CO ₂	0.001	0.000	0.999
Industrial Proc.	2A7c Expanded clay	CO ₂	0.001	0.000	0.999
Energy	Stationary Combustion, WASTE	CH ₄	0.001	0.000	0.999
Solvent and Other Prod. Use	3D5 Consumption of fireworks	N ₂ O	0.001	0.000	1.000
Energy	1.B.2 Off-shore activities	CO ₂	0.001	0.000	1.000
Energy	Stationary Combustion, BKB	CO ₂	0.001	0.000	1.000
Energy	1.B.2 Transmission of natural gas	CH ₄	0.001	0.000	1.000
Energy	1.B.2 Flaring in refinery	N ₂ O	0.001	0.000	1.000
LULUCF	5.D Wetlands, Dead organic matter	CO ₂	0.001	0.000	1.000
LULUCF	5.E Settlements, Soils	CO ₂	0.000	0.000	1.000
Energy	Transport, Forestry	CH ₄	0.000	0.000	1.000
Industrial Proc.	2A6 Road paving with asphalt	CO ₂	0.000	0.000	1.000
Agriculture	4.F Field Burning of Agricultural Residues	N ₂ O	0.000	0.000	1.000
Energy	Transport, Navigation (small boats)	CH ₄	0.000	0.000	1.000
Energy	Transport, Navigation (large vessels)	CH ₄	0.000	0.000	1.000

Tier 2 Analysis IPCC Source Categories (LU- LUCF included)		DK GHG	Base Year Estimate Ex,o Mt CO ₂ eq	Base Year Level Asses- sment Lx,o	Base Year Cumulative Total of Lx,o
LULUCF	5(V) Biomass Burning	CH ₄	0.000	0.000	1.000
Industrial Proc.	2D2 Food and Drink	CO ₂	0.000	0.000	1.000
LULUCF	5.E Settlements, Dead organic matter	CO ₂	0.000	0.000	1.000
Waste	6.C Incineration of corpses	N ₂ O	0.000	0.000	1.000
Energy	Transport, Fisheries	CH ₄	0.000	0.000	1.000
LULUCF	5(V) Biomass Burning	N ₂ O	0.000	0.000	1.000
Energy	Transport, Railways	CH ₄	0.000	0.000	1.000
Industrial Proc.	2F Consumption of PFC	PFC	0.000	0.000	1.000
LULUCF	5(III) Disturbance, Land converted to cropland	N ₂ O	0.000	0.000	1.000
Energy	1.B.2 Venting in gas storage	CH ₄	0.000	0.000	1.000
Energy	Biogas fuelled engines, BIOMASS	CH ₄	0.000	0.000	1.000
LULUCF	5(II) Wetlands	N ₂ O	0.000	0.000	1.000
LULUCF	5.C Grassland, Mineral soils	CO ₂	0.000	0.000	1.000
Energy	Transport, Military	CH ₄	0.000	0.000	1.000
Energy	Natural gas fuelled engines, GAS	CH ₄	0.000	0.000	1.000
Energy	Transport, Civil aviation	CH ₄	0.000	0.000	1.000
Solvent and Other Prod. Use	3D5 Use of tobacco	N ₂ O	0.000	0.000	1.000
Solvent and Other Prod. Use	3D5 Use of charcoal for BBQ	N ₂ O	0.000	0.000	1.000
Industrial Proc.	2B5 Catalysts/Fertilizers, Pesticides and Sulphuric acid	CO ₂	0.000	0.000	1.000
Solvent and Other Prod. Use	3D5 Consumption of fireworks	CO ₂	0.000	0.000	1.000
Energy	1.B.2. Flaring in refinery	CH ₄	0.000	0.000	1.000
Waste	6.C Incineration of carcasses	N ₂ O	0.000	0.000	1.000
Waste	6.C Incineration of corpses	CH ₄	0.000	0.000	1.000
Solvent and Other Prod. Use	3D5 Use of candles	N ₂ O	0.000	0.000	1.000
Industrial Proc.	2A5 Asphalt roofing	CO ₂	0.000	0.000	1.000
Energy	1.B.2 Land based activities	CO ₂	0.000	0.000	1.000
Energy	1.B.2 Distribution of natural gas	CO ₂	0.000	0.000	1.000
Waste	6.C Incineration of carcasses	CH ₄	0.000	0.000	1.000
Energy	1.B.2 Transmission of natural gas	CO ₂	0.000	0.000	1.000
Energy	1.B.2 Venting in gas storage	CO ₂	0.000	0.000	1.000
Solvent and Other Prod. Use	3B Degreasing and dry cleaning	CO ₂	0.000	0.000	1.000
Solvent and Other Prod. Use	3D1 Other - Use of N ₂ O for Anaesthesia	N ₂ O	0.000	0.000	1.000
Total			20.179	1.000	

¹⁾ The Estimates include signs, where + : emission - : removal, although in the level analyses only the absolute values are used.

Table A1-20 KCA for Denmark, level assessment 2012 excl. LULUCF, tier 2.

Tier 2 Analysis IPCC Source Categories (LULUCF excluded)		DK GHG	Latest Year Estimate	Latest Year Level Assessment	Latest Year Cumulative Total of Lx,t
			Ex,t Mt CO ₂ eq	Lx,t	
Agriculture	4.D3 Leaching	N ₂ O	1.458	0.118	0.118
Agriculture	4.D1.2 Animal waste applied to soils	N ₂ O	1.212	0.098	0.216
Agriculture	4.D1.1 Synthetic Fertilizer	N ₂ O	1.138	0.092	0.307
Energy	Stationary Combustion, BIOMASS	N ₂ O	0.909	0.073	0.381
Waste	6 A. Solid Waste Disposal on Land	CH ₄	0.826	0.067	0.448
Energy	Transport, Road transport	CO ₂	0.604	0.049	0.496
Agriculture	4A Enteric Fermentation	CH ₄	0.584	0.047	0.543
Energy	Transport, Industry (mobile)	CO ₂	0.422	0.034	0.578
Agriculture	4.B Manure Management	N ₂ O	0.400	0.032	0.610
Industrial Proc.	2F Consumption of HFC	HFC	0.335	0.027	0.637
Energy	Transport, Agriculture	CO ₂	0.329	0.027	0.663
Agriculture	4.D1.4 Crop Residue	N ₂ O	0.317	0.026	0.689
Agriculture	4.D3 Atmospheric deposition	N ₂ O	0.300	0.024	0.713
Agriculture	4B Manure Management	CH ₄	0.267	0.022	0.735
Agriculture	4.D1.3 N-fixing crops	N ₂ O	0.261	0.021	0.756
Agriculture	4.D.2 Grassing animals	N ₂ O	0.218	0.018	0.774
Agriculture	4.D1.5 Cultivation of histosols	N ₂ O	0.202	0.016	0.790
Energy	Stationary Combustion, GAS	N ₂ O	0.183	0.015	0.805
Energy	Transport, Agriculture	N ₂ O	0.177	0.014	0.819
Energy	Stationary Combustion, Fossil waste	CO ₂	0.156	0.013	0.832
Waste	6.D Compost production	N ₂ O	0.137	0.011	0.843
Energy	Stationary Combustion, BIOMASS	CH ₄	0.136	0.011	0.854
Energy	Transport, Industry (mobile)	N ₂ O	0.134	0.011	0.865
Energy	Stationary Combustion, LIQUID	N ₂ O	0.111	0.009	0.873
Energy	Stationary Combustion, SOLID	N ₂ O	0.110	0.009	0.882
Energy	Stationary Combustion, Coal	CO ₂	0.103	0.008	0.891
Waste	6.D Compost production	CH ₄	0.097	0.008	0.899
Energy	Transport, Fisheries	N ₂ O	0.094	0.008	0.906
Energy	Stationary Combustion, Natural gas	CO ₂	0.093	0.007	0.914
Energy	Transport, Navigation (large vessels)	N ₂ O	0.078	0.006	0.920
Energy	Stationary Combustion, WASTE	N ₂ O	0.063	0.005	0.925
Energy	Transport, Commercial/institutional	CO ₂	0.061	0.005	0.930
Industrial Proc.	2F Consumption of SF ₆	SF ₆	0.060	0.005	0.935
Energy	1.B.2 Refinery processes	CH ₄	0.058	0.005	0.939
Energy	Transport, Road transport	N ₂ O	0.058	0.005	0.944
Energy	Transport, Navigation (large vessels)	CO ₂	0.048	0.004	0.948
Energy	Transport, Navigation (small boats)	CO ₂	0.041	0.003	0.951
Agriculture	4.D1.6 Sewage sludge and Industrial waste used as fertiliser	N ₂ O	0.040	0.003	0.954
Energy	Stationary Combustion, Gas oil	CO ₂	0.034	0.003	0.957
Energy	Stationary Combustion, Petroleum coke	CO ₂	0.034	0.003	0.960
Waste	6.D Accidental fires, buildings	CO ₂	0.032	0.003	0.963
Waste	6 B. Wastewater Handling	CH ₄	0.029	0.002	0.965
Waste	6.D Accidental fires, vehicles	CO ₂	0.028	0.002	0.967
Energy	Transport, Fisheries	CO ₂	0.026	0.002	0.969
Waste	6 B. Wastewater Handling - Indirect	N ₂ O	0.023	0.002	0.971
Energy	Transport, Residential	CO ₂	0.022	0.002	0.973
Waste	6 B. Wastewater Handling - Direct	N ₂ O	0.022	0.002	0.975
Energy	Transport, Railways	N ₂ O	0.021	0.002	0.976
Energy	Transport, Civil aviation	N ₂ O	0.021	0.002	0.978
Energy	Stationary Combustion, Refinery gas	CO ₂	0.020	0.002	0.980
Industrial Proc.	2A1 Cement production	CO ₂	0.019	0.002	0.981
Solvent and Other Prod. Use	3D5 Use of candles	CO ₂	0.018	0.001	0.983
Energy	1.B.2 Flaring off-shore	CO ₂	0.015	0.001	0.984
Energy	Transport, Civil aviation	CO ₂	0.015	0.001	0.985
Energy	Transport, Railways	CO ₂	0.013	0.001	0.986
Energy	Stationary Combustion, Residual oil	CO ₂	0.013	0.001	0.987
Energy	Transport, Military	N ₂ O	0.013	0.001	0.988
Energy	1.B.2 Off-shore activities	CH ₄	0.011	0.001	0.989
Energy	Transport, Navigation (small boats)	N ₂ O	0.011	0.001	0.990
Solvent and Other Prod. Use	3D5 Other	CO ₂	0.010	0.001	0.991
Energy	Transport, Commercial/institutional	N ₂ O	0.008	0.001	0.992
Energy	1.B.2 Land based activities	CH ₄	0.007	0.001	0.992

Tier 2 Analysis IPCC Source Categories (LULUCF excluded)		DK GHG	Latest Year Estimate	Latest Year Level Assessment	Latest Year Cumulative Total of Lx,t
			Ex,t Mt CO ₂ eq	Lx,t	
Waste	6.D Accidental fires, buildings	CH ₄	0.006	0.001	0.993
Energy	Transport, Military	CO ₂	0.006	0.001	0.993
Energy	Stationary Combustion, GAS	CH ₄	0.006	0.000	0.994
Energy	Transport, Forestry	CO ₂	0.005	0.000	0.994
Energy	Stationary Combustion, LPG	CO ₂	0.005	0.000	0.994
Energy	1.B.2 Flaring off-shore	N ₂ O	0.005	0.000	0.995
Energy	Transport, Road transport	CH ₄	0.005	0.000	0.995
Industrial Proc.	2F Consumption of PFC	PFC	0.004	0.000	0.996
Energy	Stationary Combustion, Coke	CO ₂	0.004	0.000	0.996
Energy	Transport, Commercial/institutional	CH ₄	0.003	0.000	0.996
Energy	Biogas fuelled engines, BIOMASS	CH ₄	0.003	0.000	0.996
Energy	Transport, Residential	N ₂ O	0.003	0.000	0.997
Energy	Stationary Combustion, SOLID	CH ₄	0.003	0.000	0.997
Industrial Proc.	2A2 Lime production	CO ₂	0.003	0.000	0.997
Energy	Natural gas fuelled engines, GAS	CH ₄	0.003	0.000	0.997
Energy	1.B.2 Flaring in refinery	CO ₂	0.002	0.000	0.998
Solvent and Other Prod. Use	3C Chemical products, manufacturing and processing	CO ₂	0.002	0.000	0.998
Solvent and Other Prod. Use	3D5 Consumption of fireworks	N ₂ O	0.002	0.000	0.998
Energy	Transport, Agriculture	CH ₄	0.002	0.000	0.998
Industrial Proc.	2A3 Limestone and dolomite use	CO ₂	0.002	0.000	0.998
Industrial Proc.	2G Lubricants	CO ₂	0.002	0.000	0.998
Waste	6.D Accidental fires, vehicles	CH ₄	0.002	0.000	0.999
Energy	Transport, Forestry	N ₂ O	0.002	0.000	0.999
Energy	Stationary Combustion, WASTE	CH ₄	0.002	0.000	0.999
Energy	Transport, Residential	CH ₄	0.001	0.000	0.999
Agriculture	4F Field burning of agricultural residues	CH ₄	0.001	0.000	0.999
Solvent and Other Prod. Use	3A Paint application	CO ₂	0.001	0.000	0.999
Energy	1.B.2 Off-shore activities	CO ₂	0.001	0.000	0.999
Energy	Stationary Combustion, LIQUID	CH ₄	0.001	0.000	0.999
Industrial Proc.	2A7b Yellow bricks	CO ₂	0.001	0.000	0.999
Energy	1.B.2 Distribution of natural gas	CH ₄	0.001	0.000	0.999
Energy	Transport, Industry (mobile)	CH ₄	0.001	0.000	0.999
Solvent and Other Prod. Use	3D1 Other - Use of N2O for Anaesthesia	N ₂ O	0.001	0.000	1.000
Energy	1.B.2 Flaring in refinery	N ₂ O	0.001	0.000	1.000
Energy	Transport, Navigation (small boats)	CH ₄	0.001	0.000	1.000
Industrial Proc.	2A7a Glass and Glass wool	CO ₂	0.001	0.000	1.000
Agriculture	4.F Field Burning of Agricultural Residues	N ₂ O	0.001	0.000	1.000
Industrial Proc.	2A6 Road paving with asphalt	CO ₂	0.000	0.000	1.000
Energy	1.B.2. Flaring off-shore	CH ₄	0.000	0.000	1.000
Industrial Proc.	2A7c Expanded clay	CO ₂	0.000	0.000	1.000
Waste	6.C Incineration of corpses	N ₂ O	0.000	0.000	1.000
Energy	Transport, Fisheries	CH ₄	0.000	0.000	1.000
Energy	Transport, Navigation (large vessels)	CH ₄	0.000	0.000	1.000
Energy	1.B.2 Venting in gas storage	CH ₄	0.000	0.000	1.000
Industrial Proc.	2D2 Food and Drink	CO ₂	0.000	0.000	1.000
Energy	Transport, Railways	CH ₄	0.000	0.000	1.000
Solvent and Other Prod. Use	3D5 Consumption of fireworks	CO ₂	0.000	0.000	1.000
Waste	6.C Incineration of carcasses	N ₂ O	0.000	0.000	1.000
Solvent and Other Prod. Use	3D5 Use of charcoal for BBQ	N ₂ O	0.000	0.000	1.000
Industrial Proc.	2B5 Catalysts/Fertilizers, Pesticides and Sulphuric acid	CO ₂	0.000	0.000	1.000
Energy	Stationary Combustion, Kerosene	CO ₂	0.000	0.000	1.000
Energy	Stationary Combustion, BKB	CO ₂	0.000	0.000	1.000
Energy	Transport, Military	CH ₄	0.000	0.000	1.000
Solvent and Other Prod. Use	3D5 Use of tobacco	N ₂ O	0.000	0.000	1.000
Energy	Transport, Forestry	CH ₄	0.000	0.000	1.000
Energy	Transport, Civil aviation	CH ₄	0.000	0.000	1.000
Solvent and Other Prod. Use	3D5 Use of candles	N ₂ O	0.000	0.000	1.000
Energy	1.B.2 Transmission of natural gas	CH ₄	0.000	0.000	1.000
Energy	1.B.2. Flaring in refinery	CH ₄	0.000	0.000	1.000
Waste	6.C Incineration of corpses	CH ₄	0.000	0.000	1.000
Waste	6.C Incineration of carcasses	CH ₄	0.000	0.000	1.000
Industrial Proc.	2A5 Asphalt roofing	CO ₂	0.000	0.000	1.000

Tier 2 Analysis IPCC Source Categories (LULUCF excluded)		DK GHG	Latest Year Estimate	Latest Year Level Assessment	Latest Year Cumulative Total of Lx,t
			Ex,t Mt CO ₂ eq	Lx,t	
Energy	1.B.2 Land based activities	CO ₂	0.000	0.000	1.000
Energy	1.B.2 Distribution of natural gas	CO ₂	0.000	0.000	1.000
Energy	1.B.2 Venting in gas storage	CO ₂	0.000	0.000	1.000
Energy	1.B.2 Transmission of natural gas	CO ₂	0.000	0.000	1.000
Solvent and Other Prod. Use	3B Degreasing and dry cleaning	CO ₂	0.000	0.000	1.000
Industrial Proc.	2C1 Iron and steel production	CO ₂	0.000	0.000	1.000
Industrial Proc.	2B2 Nitric acid production	N ₂ O	0.000	0.000	1.000
Total			12.386	1.000	

Table A1-21 KCA for Denmark, level assessment 2012 incl. LULUCF, tier 2.

Tier 2 Analysis IPCC Source Categories (LU- LUCF included)		DK GHG	Latest Year Estimate Ex,t Mt CO ₂ eq	Latest Year Level Assessment Lx,t	Latest Year Cumulative Total of Lx,t
LULUCF	5.B Cropland, Organic soils	CO ₂	1.794	0.113	0.113
Agriculture	4.D3 Leaching	N ₂ O	1.458	0.092	0.205
Agriculture	4.D1.2 Animal waste applied to soils	N ₂ O	1.212	0.076	0.281
Agriculture	4.D1.1 Synthetic Fertilizer	N ₂ O	1.138	0.072	0.353
Energy	Stationary Combustion, BIOMASS	N ₂ O	0.909	0.057	0.410
Waste	6 A. Solid Waste Disposal on Land	CH ₄	0.826	0.052	0.462
LULUCF	5.A.1 Forest remaining forest	CO ₂	0.680	0.043	0.505
Energy	Transport, Road transport	CO ₂	0.604	0.038	0.543
Agriculture	4A Enteric Fermentation	CH ₄	0.584	0.037	0.580
LULUCF	5.B Cropland, Mineral soils	CO ₂	0.437	0.028	0.607
Energy	Transport, Industry (mobile)	CO ₂	0.422	0.027	0.634
Agriculture	4.B Manure Management	N ₂ O	0.400	0.025	0.659
Industrial Proc.	2F Consumption of HFC	HFC	0.335	0.021	0.680
Energy	Transport, Agriculture	CO ₂	0.329	0.021	0.701
Agriculture	4.D1.4 Crop Residue	N ₂ O	0.317	0.020	0.721
Agriculture	4.D3 Atmospheric deposition	N ₂ O	0.300	0.019	0.740
Agriculture	4B Manure Management	CH ₄	0.267	0.017	0.757
Agriculture	4.D1.3 N-fixing crops	N ₂ O	0.261	0.016	0.773
LULUCF	5.C Grassland, Living biomass	CO ₂	0.236	0.015	0.788
Agriculture	4.D.2 Grassing animals	N ₂ O	0.218	0.014	0.802
Agriculture	4.D1.5 Cultivation of histosols	N ₂ O	0.202	0.013	0.815
Energy	Stationary Combustion, GAS	N ₂ O	0.183	0.012	0.826
Energy	Transport, Agriculture	N ₂ O	0.177	0.011	0.837
Energy	Stationary Combustion, Fossil waste	CO ₂	0.156	0.010	0.847
Waste	6.D Compost production	N ₂ O	0.137	0.009	0.856
Energy	Stationary Combustion, BIOMASS	CH ₄	0.136	0.009	0.864
Energy	Transport, Industry (mobile)	N ₂ O	0.134	0.008	0.873
Energy	Stationary Combustion, LIQUID	N ₂ O	0.111	0.007	0.880
Energy	Stationary Combustion, SOLID	N ₂ O	0.110	0.007	0.887
Energy	Stationary Combustion, Coal	CO ₂	0.103	0.006	0.893
LULUCF	5.B Cropland, Living biomass	CO ₂	0.098	0.006	0.899
Waste	6.D Compost production	CH ₄	0.097	0.006	0.906
LULUCF	5(IV) Cropland Limestone	CO ₂	0.097	0.006	0.912
Energy	Transport, Fisheries	N ₂ O	0.094	0.006	0.918
Energy	Stationary Combustion, Natural gas	CO ₂	0.093	0.006	0.923
Energy	Transport, Navigation (large vessels)	N ₂ O	0.078	0.005	0.928
LULUCF	5.C Grassland, Organic soils	CO ₂	0.071	0.004	0.933
Energy	Stationary Combustion, WASTE	N ₂ O	0.063	0.004	0.937
Energy	Transport, Commercial/institutional	CO ₂	0.061	0.004	0.941
Industrial Proc.	2F Consumption of SF ₆	SF ₆	0.060	0.004	0.944
Energy	1.B.2 Refinery processes	CH ₄	0.058	0.004	0.948
Energy	Transport, Road transport	N ₂ O	0.058	0.004	0.952
Energy	Transport, Navigation (large vessels)	CO ₂	0.048	0.003	0.955
Energy	Transport, Navigation (small boats)	CO ₂	0.041	0.003	0.957
Agriculture	4.D1.6 Sewage sludge and Industrial waste used as fertiliser	N ₂ O	0.040	0.002	0.960
Energy	Stationary Combustion, Gas oil	CO ₂	0.034	0.002	0.962
Energy	Stationary Combustion, Petroleum coke	CO ₂	0.034	0.002	0.964
Waste	6.D Accidental fires, buildings	CO ₂	0.032	0.002	0.966
Waste	6 B. Wastewater Handling	CH ₄	0.029	0.002	0.968
LULUCF	5.E Settlements, Living biomass	CO ₂	0.029	0.002	0.970
Waste	6.D Accidental fires, vehicles	CO ₂	0.028	0.002	0.972
Energy	Transport, Fisheries	CO ₂	0.026	0.002	0.973
Waste	6 B. Wastewater Handling - Indirect	N ₂ O	0.023	0.001	0.975
Energy	Transport, Residential	CO ₂	0.022	0.001	0.976
Waste	6 B. Wastewater Handling - Direct	N ₂ O	0.022	0.001	0.977
Energy	Transport, Railways	N ₂ O	0.021	0.001	0.979
Energy	Transport, Civil aviation	N ₂ O	0.021	0.001	0.980
Energy	Stationary Combustion, Refinery gas	CO ₂	0.020	0.001	0.981
Industrial Proc.	2A1 Cement production	CO ₂	0.019	0.001	0.983
Solvent and Other Prod. Use	3D5 Use of candles	CO ₂	0.018	0.001	0.984
LULUCF	5.E Settlements, Soils	CO ₂	0.018	0.001	0.985

Tier 2 Analysis IPCC Source Categories (LU- LUCF included)		DK GHG	Latest Year Estimate	Latest Year Level Asses- ment Lx,t	Latest Year Cumulative Total of Lx,t
			Ex,t Mt CO ₂ eq		
Energy	1.B.2 Flaring off-shore	CO ₂	0.015	0.001	0.986
Energy	Transport, Civil aviation	CO ₂	0.015	0.001	0.987
Energy	Transport, Railways	CO ₂	0.013	0.001	0.988
Energy	Stationary Combustion, Residual oil	CO ₂	0.013	0.001	0.988
Energy	Transport, Military	N ₂ O	0.013	0.001	0.989
Energy	1.B.2 Off-shore activities	CH ₄	0.011	0.001	0.990
Energy	Transport, Navigation (small boats)	N ₂ O	0.011	0.001	0.991
Solvent and Other Prod. Use	3D5 Other	CO ₂	0.010	0.001	0.991
Energy	Transport, Commercial/institutional	N ₂ O	0.008	0.001	0.992
Energy	1.B.2 Land based activities	CH ₄	0.007	0.000	0.992
LULUCF	5.A.2 Land converted to forest	CO ₂	0.007	0.000	0.993
Waste	6.D Accidental fires, buildings	CH ₄	0.006	0.000	0.993
Energy	Transport, Military	CO ₂	0.006	0.000	0.993
LULUCF	5.B Cropland, Dead organic matter	CO ₂	0.006	0.000	0.994
Energy	Stationary Combustion, GAS	CH ₄	0.006	0.000	0.994
Energy	Transport, Forestry	CO ₂	0.005	0.000	0.994
Energy	Stationary Combustion, LPG	CO ₂	0.005	0.000	0.995
Energy	1.B.2 Flaring off-shore	N ₂ O	0.005	0.000	0.995
Energy	Transport, Road transport	CH ₄	0.005	0.000	0.995
Industrial Proc.	2F Consumption of PFC	PFC	0.004	0.000	0.996
Energy	Stationary Combustion, Coke	CO ₂	0.004	0.000	0.996
LULUCF	5(II) Forest Land.	N ₂ O	0.004	0.000	0.996
LULUCF	5.C Grassland, Dead organic matter	CO ₂	0.004	0.000	0.996
LULUCF	5.C Grassland, Mineral soils	CO ₂	0.003	0.000	0.997
Energy	Transport, Commercial/institutional	CH ₄	0.003	0.000	0.997
Energy	Biogas fuelled engines, BIOMASS	CH ₄	0.003	0.000	0.997
Energy	Transport, Residential	N ₂ O	0.003	0.000	0.997
Energy	Stationary Combustion, SOLID	CH ₄	0.003	0.000	0.997
Industrial Proc.	2A2 Lime production	CO ₂	0.003	0.000	0.998
Energy	Natural gas fuelled engines, GAS	CH ₄	0.003	0.000	0.998
LULUCF	5.D Wetlands, Soils	CO ₂	0.003	0.000	0.998
Energy	1.B.2 Flaring in refinery	CO ₂	0.002	0.000	0.998
Solvent and Other Prod. Use	3C Chemical products, manufacturing and pro- cessing	CO ₂	0.002	0.000	0.998
Solvent and Other Prod. Use	3D5 Consumption of fireworks	N ₂ O	0.002	0.000	0.998
Energy	Transport, Agriculture	CH ₄	0.002	0.000	0.998
Industrial Proc.	2A3 Limestone and dolomite use	CO ₂	0.002	0.000	0.999
Industrial Proc.	2G Lubricants	CO ₂	0.002	0.000	0.999
Waste	6.D Accidental fires, vehicles	CH ₄	0.002	0.000	0.999
Energy	Transport, Forestry	N ₂ O	0.002	0.000	0.999
Energy	Stationary Combustion, WASTE	CH ₄	0.002	0.000	0.999
Energy	Transport, Residential	CH ₄	0.001	0.000	0.999
Agriculture	4F Field burning of agricultural residues	CH ₄	0.001	0.000	0.999
Solvent and Other Prod. Use	3A Paint application	CO ₂	0.001	0.000	0.999
Energy	1.B.2 Off-shore activities	CO ₂	0.001	0.000	0.999
Energy	Stationary Combustion, LIQUID	CH ₄	0.001	0.000	0.999
Industrial Proc.	2A7b Yellow bricks	CO ₂	0.001	0.000	0.999
LULUCF	5(III) Disturbance, Land converted to cropland	N ₂ O	0.001	0.000	0.999
Energy	1.B.2 Distribution of natural gas	CH ₄	0.001	0.000	1.000
Energy	Transport, Industry (mobile)	CH ₄	0.001	0.000	1.000
Solvent and Other Prod. Use	3D1 Other - Use of N ₂ O for Anaesthesia	N ₂ O	0.001	0.000	1.000
Energy	1.B.2 Flaring in refinery	N ₂ O	0.001	0.000	1.000
Energy	Transport, Navigation (small boats)	CH ₄	0.001	0.000	1.000
Industrial Proc.	2A7a Glass and Glass wool	CO ₂	0.001	0.000	1.000
Agriculture	4.F Field Burning of Agricultural Residues	N ₂ O	0.001	0.000	1.000
Industrial Proc.	2A6 Road paving with asphalt	CO ₂	0.000	0.000	1.000
Energy	1.B.2. Flaring off-shore	CH ₄	0.000	0.000	1.000
Industrial Proc.	2A7c Expanded clay	CO ₂	0.000	0.000	1.000
Waste	6.C Incineration of corpses	N ₂ O	0.000	0.000	1.000
Energy	Transport, Fisheries	CH ₄	0.000	0.000	1.000

Tier 2 Analysis IPCC Source Categories (LU- LUCF included)		DK GHG	Latest Year Estimate	Latest Year Level Assessment Lx,t	Latest Year Cumulative Total of Lx,t
			Ex,t Mt CO ₂ eq		
Energy	Transport, Navigation (large vessels)	CH ₄	0.000	0.000	1.000
Energy	1.B.2 Venting in gas storage	CH ₄	0.000	0.000	1.000
Industrial Proc.	2D2 Food and Drink	CO ₂	0.000	0.000	1.000
Energy	Transport, Railways	CH ₄	0.000	0.000	1.000
Solvent and Other Prod. Use	3D5 Consumption of fireworks	CO ₂	0.000	0.000	1.000
LULUCF	5(II) Wetlands	N ₂ O	0.000	0.000	1.000
Waste	6.C Incineration of carcasses	N ₂ O	0.000	0.000	1.000
Solvent and Other Prod. Use	3D5 Use of charcoal for BBQ	N ₂ O	0.000	0.000	1.000
LULUCF	5.D Wetlands, Living biomass	CO ₂	0.000	0.000	1.000
Industrial Proc.	2B5 Catalysts/Fertilizers, Pesticides and Sulphuric acid	CO ₂	0.000	0.000	1.000
Energy	Stationary Combustion, Kerosene	CO ₂	0.000	0.000	1.000
Energy	Stationary Combustion, BKB	CO ₂	0.000	0.000	1.000
Energy	Transport, Military	CH ₄	0.000	0.000	1.000
Solvent and Other Prod. Use	3D5 Use of tobacco	N ₂ O	0.000	0.000	1.000
Energy	Transport, Forestry	CH ₄	0.000	0.000	1.000
Energy	Transport, Civil aviation	CH ₄	0.000	0.000	1.000
Solvent and Other Prod. Use	3D5 Use of candles	N ₂ O	0.000	0.000	1.000
Energy	1.B.2 Transmission of natural gas	CH ₄	0.000	0.000	1.000
Energy	1.B.2. Flaring in refinery	CH ₄	0.000	0.000	1.000
LULUCF	5(V) Biomass Burning	N ₂ O	0.000	0.000	1.000
LULUCF	5.E Settlements, Dead organic matter	CO ₂	0.000	0.000	1.000
Waste	6.C Incineration of corpses	CH ₄	0.000	0.000	1.000
LULUCF	5(V) Biomass Burning	CH ₄	0.000	0.000	1.000
Waste	6.C Incineration of carcasses	CH ₄	0.000	0.000	1.000
Industrial Proc.	2A5 Asphalt roofing	CO ₂	0.000	0.000	1.000
Energy	1.B.2 Land based activities	CO ₂	0.000	0.000	1.000
Energy	1.B.2 Distribution of natural gas	CO ₂	0.000	0.000	1.000
Energy	1.B.2 Venting in gas storage	CO ₂	0.000	0.000	1.000
Energy	1.B.2 Transmission of natural gas	CO ₂	0.000	0.000	1.000
Solvent and Other Prod. Use	3B Degreasing and dry cleaning	CO ₂	0.000	0.000	1.000
Industrial Proc.	2C1 Iron and steel production	CO ₂	0.000	0.000	1.000
Industrial Proc.	2B2 Nitric acid production	N ₂ O	0.000	0.000	1.000
LULUCF	5.D Wetlands, Dead organic matter	CO ₂	0.000	0.000	1.000
Total			15.872	1.000	

¹⁾ The Estimates include signs, where + : emission - : removal, although in the level analyses only the absolute values are used.

Table A1-22 KCA for Denmark, trend assessment 1990-2012 excl. LULUCF, tier 2.

Tier 2 Analysis IPCC Source Categories (LULUCF excluded)		DK GHG	Base Year Estimate	Latest Year Estimate	Trend Assessment	Contribution to Trend	Cumulative
			Ex,o Mt CO ₂ eq	Ex,t Mt CO ₂ eq	Tx,t		
Agriculture	4.D1.1 Synthetic Fertilizer	N ₂ O	2.354	1.103	0.9862	0.122	0.122
Energy	Stationary Combustion, BIOMASS	N ₂ O	0.038	0.091	0.9064	0.112	0.234
Agriculture	4.D3 Leaching	N ₂ O	2.447	1.430	0.5966	0.074	0.307
Waste	6 A. Solid Waste Disposal on Land	CH ₄	1.366	0.698	0.5576	0.069	0.376
Agriculture	4.D1.2 Animal waste applied to soils	N ₂ O	1.112	1.161	0.4977	0.061	0.438
Industrial Proc.	2F Consumption of HFC	HFC	0.218	0.657	0.3655	0.045	0.483
Energy	Transport, Road transport	CO ₂	9.284	11.224	0.3336	0.041	0.524
Energy	Stationary Combustion, LIQUID	N ₂ O	0.044	0.011	0.3175	0.039	0.563
Industrial Proc.	2B2 Nitric acid production	N ₂ O	1.043	0.000	0.2842	0.035	0.598
Energy	Transport, Industry (mobile)	CO ₂	0.839	1.021	0.2353	0.029	0.627
Waste	6.D Compost production	N ₂ O	0.013	0.127	0.1831	0.023	0.650
Energy	Stationary Combustion, Gas oil	CO ₂	4.547	0.742	0.1772	0.022	0.672
Agriculture	4B Manure Management	CH ₄	0.985	1.297	0.1672	0.021	0.692
Energy	Stationary Combustion, Fossil waste	CO ₂	0.573	1.397	0.1569	0.019	0.712
Energy	Transport, Agriculture	CO ₂	1.272	1.343	0.1388	0.017	0.729
Agriculture	4A Enteric Fermentation	CH ₄	3.247	2.904	0.1376	0.017	0.746
Energy	Stationary Combustion, SOLID	N ₂ O	0.068	0.027	0.1367	0.017	0.763
Energy	Stationary Combustion, GAS	N ₂ O	0.016	0.024	0.1325	0.016	0.779
Energy	Stationary Combustion, Coal	CO ₂	23.834	10.005	0.1172	0.014	0.794
Agriculture	4.D3 Atmospheric deposition	N ₂ O	0.496	0.295	0.1139	0.014	0.808
Waste	6.D Compost production	CH ₄	0.029	0.090	0.1072	0.013	0.821
Energy	Transport, Agriculture	N ₂ O	0.015	0.018	0.0909	0.011	0.832
Agriculture	4.B Manure Management	N ₂ O	0.600	0.391	0.0869	0.011	0.843
Energy	Stationary Combustion, BIOMASS	CH ₄	0.102	0.135	0.0860	0.011	0.853
Energy	1.B.2 Refinery processes	CH ₄	0.001	0.047	0.0837	0.010	0.864
Energy	Stationary Combustion, Natural gas	CO ₂	4.335	8.293	0.0818	0.010	0.874
Agriculture	4.D1.3 N-fixing crops	N ₂ O	0.269	0.256	0.0804	0.010	0.884
Energy	Transport, Industry (mobile)	N ₂ O	0.011	0.013	0.0796	0.010	0.894
Energy	Stationary Combustion, WASTE	N ₂ O	0.007	0.016	0.0626	0.008	0.901
Agriculture	4.D1.4 Crop Residue	N ₂ O	0.361	0.311	0.0601	0.007	0.909
Energy	Transport, Commercial/institutional	CO ₂	0.074	0.171	0.0596	0.007	0.916
Agriculture	4.D.2 Grassing animals	N ₂ O	0.334	0.211	0.0591	0.007	0.923
Energy	Transport, Navigation (large vessels)	N ₂ O	0.015	0.008	0.0458	0.006	0.929
Energy	Stationary Combustion, Residual oil	CO ₂	2.496	0.571	0.0440	0.005	0.935
Energy	Transport, Navigation (small boats)	CO ₂	0.048	0.099	0.0376	0.005	0.939
Energy	Transport, Road transport	N ₂ O	0.091	0.116	0.0352	0.004	0.944
Waste	6 B. Wastewater Handling - Indirect	N ₂ O	0.082	0.032	0.0310	0.004	0.947
Energy	Transport, Navigation (large vessels)	CO ₂	0.748	0.399	0.0282	0.003	0.951
Agriculture	4.D1.5 Cultivation of histosols	N ₂ O	0.290	0.198	0.0282	0.003	0.954
Industrial Proc.	2F Consumption of SF ₆	SF ₆	0.107	0.118	0.0277	0.003	0.958
Agriculture	4.D1.6 Sewage sludge and Industrial waste used as fertiliser	N ₂ O	0.028	0.039	0.0264	0.003	0.961
Energy	Stationary Combustion, Petroleum coke	CO ₂	0.415	0.628	0.0248	0.003	0.964
Energy	Stationary Combustion, Kerosene	CO ₂	0.366	0.002	0.0213	0.003	0.967
Solvent and Other Prod. Use	3D5 Use of candles	CO ₂	0.022	0.081	0.0210	0.003	0.969
Waste	6 B. Wastewater Handling - Direct	N ₂ O	0.023	0.041	0.0186	0.002	0.972
Energy	Transport, Residential	CO ₂	0.039	0.062	0.0170	0.002	0.974
Waste	6 B. Wastewater Handling	CH ₄	0.065	0.074	0.0144	0.002	0.975
Energy	Transport, Road transport	CH ₄	0.047	0.011	0.0138	0.002	0.977
Energy	Transport, Navigation (small boats)	N ₂ O	0.000	0.001	0.0115	0.001	0.979
Energy	Transport, Fisheries	N ₂ O	0.011	0.009	0.0112	0.001	0.980
Energy	1.B.2 Off-shore activities	CH ₄	0.015	0.037	0.0111	0.001	0.981
Waste	6.D Accidental fires, buildings	CO ₂	0.011	0.011	0.0098	0.001	0.983
Energy	Stationary Combustion, Refinery gas	CO ₂	0.816	0.906	0.0096	0.001	0.984
Energy	Stationary Combustion, SOLID	CH ₄	0.013	0.003	0.0095	0.001	0.985
Energy	Transport, Commercial/institutional	N ₂ O	0.000	0.001	0.0082	0.001	0.986
Energy	Transport, Civil aviation	CO ₂	0.243	0.133	0.0080	0.001	0.987
Waste	6.D Accidental fires, vehicles	CO ₂	0.006	0.006	0.0071	0.001	0.988
Industrial Proc.	2A1 Cement production	CO ₂	0.882	0.871	0.0068	0.001	0.989
Industrial Proc.	2F Consumption of PFC	PFC	0.001	0.009	0.0060	0.001	0.989
Energy	Transport, Military	N ₂ O	0.001	0.001	0.0059	0.001	0.990

Tier 2 Analysis IPCC Source Categories (LULUCF excluded)		DK GHG	Base Year Estimate	Latest Year Estimate	Trend Assessment	Contribution to Trend	Cumulative
			Ex,o Mt CO ₂ eq	Ex,t Mt CO ₂ eq	Tx,t		
Industrial Proc.	2A2 Lime production	CO ₂	0.116	0.040	0.0048	0.001	0.991
Energy	Biogas fuelled engines, BIOMASS	CH ₄	0.001	0.031	0.0047	0.001	0.991
Energy	Stationary Combustion, GAS	CH ₄	0.003	0.006	0.0047	0.001	0.992
Energy	Transport, Civil aviation	N ₂ O	0.003	0.002	0.0046	0.001	0.992
Energy	Transport, Forestry	CO ₂	0.036	0.017	0.0043	0.001	0.993
Energy	Natural gas fuelled engines, GAS	CH ₄	0.005	0.120	0.0038	0.000	0.993
Energy	Stationary Combustion, LPG	CO ₂	0.184	0.091	0.0036	0.000	0.994
Energy	1.B.2 Flaring off-shore	CO ₂	0.302	0.195	0.0035	0.000	0.994
Energy	Transport, Railways	N ₂ O	0.003	0.002	0.0034	0.000	0.995
Energy	Transport, Fisheries	CO ₂	0.591	0.479	0.0029	0.000	0.995
Energy	Transport, Residential	N ₂ O	0.000	0.000	0.0028	0.000	0.995
Energy	1.B.2 Land based activities	CH ₄	0.017	0.018	0.0027	0.000	0.996
Energy	Transport, Commercial/institutional	CH ₄	0.002	0.003	0.0025	0.000	0.996
Energy	Stationary Combustion, Coke	CO ₂	0.138	0.074	0.0023	0.000	0.996
Solvent and Other Prod. Use	3D5 Consumption of fireworks	N ₂ O	0.001	0.002	0.0022	0.000	0.997
Industrial Proc.	2C1 Iron and steel production	CO ₂	0.028	0.000	0.0022	0.000	0.997
Energy	Transport, Railways	CO ₂	0.297	0.249	0.0021	0.000	0.997
Energy	Transport, Military	CO ₂	0.119	0.116	0.0021	0.000	0.997
Waste	6.D Accidental fires, buildings	CH ₄	0.001	0.001	0.0021	0.000	0.998
Industrial Proc.	2A3 Limestone and dolomite use	CO ₂	0.014	0.026	0.0016	0.000	0.998
Energy	Stationary Combustion, WASTE	CH ₄	0.001	0.002	0.0014	0.000	0.998
Energy	Stationary Combustion, LIQUID	CH ₄	0.003	0.001	0.0014	0.000	0.998
Solvent and Other Prod. Use	3D1 Other - Use of N2O for Anaesthesia	N ₂ O	0.000	0.009	0.0010	0.000	0.998
Energy	1.B.2 Flaring off-shore	N ₂ O	0.001	0.000	0.0009	0.000	0.998
Agriculture	4F Field burning of agricultural residues	CH ₄	0.002	0.002	0.0009	0.000	0.999
Energy	Transport, Residential	CH ₄	0.001	0.001	0.0009	0.000	0.999
Energy	1.B.2 Off-shore activities	CO ₂	0.002	0.004	0.0008	0.000	0.999
Energy	1.B.2 Flaring in refinery	CO ₂	0.023	0.022	0.0008	0.000	0.999
Solvent and Other Prod. Use	3A Paint application	CO ₂	0.013	0.007	0.0007	0.000	0.999
Solvent and Other Prod. Use	3C Chemical products, manufacturing and processing	CO ₂	0.019	0.012	0.0006	0.000	0.999
Energy	1.B.2 Transmission of natural gas	CH ₄	0.004	0.000	0.0006	0.000	0.999
Energy	Stationary Combustion, BKB	CO ₂	0.011	0.001	0.0006	0.000	0.999
Energy	Transport, Agriculture	CH ₄	0.002	0.002	0.0006	0.000	0.999
Energy	Transport, Forestry	N ₂ O	0.000	0.000	0.0006	0.000	0.999
Solvent and Other Prod. Use	3D5 Other	CO ₂	0.061	0.044	0.0005	0.000	0.999
Industrial Proc.	2G Lubricants	CO ₂	0.050	0.032	0.0004	0.000	0.999
Waste	6.D Accidental fires, vehicles	CH ₄	0.000	0.000	0.0004	0.000	0.999
Energy	Transport, Forestry	CH ₄	0.000	0.000	0.0004	0.000	1.000
Industrial Proc.	2A7c Expanded clay	CO ₂	0.015	0.006	0.0004	0.000	1.000
Energy	1.B.2. Flaring off-shore	CH ₄	0.001	0.000	0.0004	0.000	1.000
Energy	Transport, Navigation (small boats)	CH ₄	0.000	0.001	0.0004	0.000	1.000
Energy	1.B.2 Distribution of natural gas	CH ₄	0.005	0.003	0.0004	0.000	1.000
Agriculture	4.F Field Burning of Agricultural Residues	N ₂ O	0.001	0.001	0.0003	0.000	1.000
Energy	Transport, Industry (mobile)	CH ₄	0.001	0.001	0.0003	0.000	1.000
Industrial Proc.	2A7a Glass and Glass wool	CO ₂	0.017	0.010	0.0003	0.000	1.000
Energy	1.B.2 Flaring in refinery	N ₂ O	0.000	0.000	0.0002	0.000	1.000
Waste	6.C Incineration of carcasses	N ₂ O	0.000	0.000	0.0002	0.000	1.000
Industrial Proc.	2A6 Road paving with asphalt	CO ₂	0.002	0.002	0.0002	0.000	1.000
Solvent and Other Prod. Use	3D5 Consumption of fireworks	CO ₂	0.000	0.000	0.0002	0.000	1.000
Solvent and Other Prod. Use	3D5 Use of charcoal for BBQ	N ₂ O	0.000	0.000	0.0001	0.000	1.000
Industrial Proc.	2D2 Food and Drink	CO ₂	0.004	0.002	0.0001	0.000	1.000
Waste	6.C Incineration of corpses	N ₂ O	0.000	0.000	0.0001	0.000	1.000
Industrial Proc.	2B5 Catalysts/Fertilizers, Pesticides and Sulphuric acid	CO ₂	0.001	0.001	0.0001	0.000	1.000
Energy	Transport, Navigation (large vessels)	CH ₄	0.000	0.000	0.0001	0.000	1.000
Energy	1.B.2 Venting in gas storage	CH ₄	0.001	0.001	0.0001	0.000	1.000

Tier 2 Analysis IPCC Source Categories (LULUCF excluded)		DK GHG	Base Year Estimate	Latest Year Estimate	Trend Assessment	Contribution to Trend	Cumulative
			Ex,o Mt CO ₂ eq	Ex,t Mt CO ₂ eq	Tx,t		
Energy	Transport, Fisheries	CH ₄	0.000	0.000	0.0001	0.000	1.000
Energy	Transport, Railways	CH ₄	0.000	0.000	0.0001	0.000	1.000
Solvent and Other Prod. Use	3D5 Use of candles	N ₂ O	0.000	0.000	0.0001	0.000	1.000
Energy	Transport, Civil aviation	CH ₄	0.000	0.000	0.0000	0.000	1.000
Energy	Transport, Military	CH ₄	0.000	0.000	0.0000	0.000	1.000
Industrial Proc.	2A7b Yellow bricks	CO ₂	0.023	0.017	0.0000	0.000	1.000
Solvent and Other Prod. Use	3D5 Use of tobacco	N ₂ O	0.000	0.000	0.0000	0.000	1.000
Energy	1.B.2. Flaring in refinery	CH ₄	0.000	0.000	0.0000	0.000	1.000
Waste	6.C Incineration of carcasses	CH ₄	0.000	0.000	0.0000	0.000	1.000
Waste	6.C Incineration of corpses	CH ₄	0.000	0.000	0.0000	0.000	1.000
Industrial Proc.	2A5 Asphalt roofing	CO ₂	0.000	0.000	0.0000	0.000	1.000
Energy	1.B.2 Land based activities	CO ₂	0.000	0.000	0.0000	0.000	1.000
Energy	1.B.2 Distribution of natural gas	CO ₂	0.000	0.000	0.0000	0.000	1.000
Energy	1.B.2 Transmission of natural gas	CO ₂	0.000	0.000	0.0000	0.000	1.000
Energy	1.B.2 Venting in gas storage	CO ₂	0.000	0.000	0.0000	0.000	1.000
Solvent and Other Prod. Use	3B Degreasing and dry cleaning	CO ₂	0.000	0.000	0.0000	0.000	1.000
Total			68.943	51.637			

Table A1-23 KCA for Denmark, trend assessment 1990-2012 incl. LULUCF, tier 2.

Tier 2 Analysis IPCC Source Categories (LU- LUCF included)		DK GHG	Base Year Estimate Ex,o Mt CO ₂ eq	Latest Year Estimate Ex,t Mt CO ₂ eq	Trend Asses- ment Tx,t	Contri- bution to Trend	Cumula- tive
LULUCF	5.A.1 Forest remaining forest	CO ₂	0.050	-4.491	0.9208	0.100	0.100
Energy	Stationary Combustion, BIOMASS	N ₂ O	0.038	0.091	0.8731	0.095	0.194
Agriculture	4.D1.1 Synthetic Fertilizer	N ₂ O	2.354	1.103	0.7034	0.076	0.271
Agriculture	4.D1.2 Animal waste applied to soils	N ₂ O	1.112	1.161	0.5622	0.061	0.332
LULUCF	5.B Cropland, Mineral soils	CO ₂	1.415	0.577	0.3983	0.043	0.375
Waste	6 A. Solid Waste Disposal on Land	CH ₄	1.366	0.698	0.3765	0.041	0.416
Energy	Transport, Road transport	CO ₂	9.284	11.224	0.3526	0.038	0.454
Industrial Proc.	2F Consumption of HFC	HFC	0.218	0.657	0.3484	0.038	0.492
Agriculture	4.D3 Leaching	N ₂ O	2.447	1.430	0.3363	0.036	0.528
LULUCF	5.C Grassland, Living biomass	CO ₂	0.075	0.463	0.2823	0.031	0.559
Energy	Stationary Combustion, LIQUID	N ₂ O	0.044	0.011	0.2561	0.028	0.586
Energy	Transport, Industry (mobile)	CO ₂	0.839	1.021	0.2482	0.027	0.613
Industrial Proc.	2B2 Nitric acid production	N ₂ O	1.043	0.000	0.2407	0.026	0.639
LULUCF	5.B Cropland, Living biomass	CO ₂	-0.074	0.193	0.1986	0.022	0.661
Agriculture	4A Enteric Fermentation	CH ₄	3.247	2.904	0.1842	0.020	0.681
Agriculture	4B Manure Management	CH ₄	0.985	1.297	0.1727	0.019	0.700
Waste	6.D Compost production	N ₂ O	0.013	0.127	0.1709	0.019	0.718
LULUCF	5(IV) Cropland Limestone	CO ₂	0.623	0.192	0.1581	0.017	0.735
Energy	Transport, Agriculture	CO ₂	1.272	1.343	0.1558	0.017	0.752
Energy	Stationary Combustion, Fossil waste	CO ₂	0.573	1.397	0.1510	0.016	0.768
Energy	Stationary Combustion, Gas oil	CO ₂	4.547	0.742	0.1461	0.016	0.784
Energy	Stationary Combustion, GAS	N ₂ O	0.016	0.024	0.1335	0.014	0.799
Energy	Stationary Combustion, SOLID	N ₂ O	0.068	0.027	0.1031	0.011	0.810
Waste	6.D Compost production	CH ₄	0.029	0.090	0.1021	0.011	0.821
Agriculture	4.D1.3 N-fixing crops	N ₂ O	0.269	0.256	0.0984	0.011	0.832
Energy	Transport, Agriculture	N ₂ O	0.015	0.018	0.0975	0.011	0.842
Energy	Stationary Combustion, BIOMASS	CH ₄	0.102	0.135	0.0887	0.010	0.852
Agriculture	4.D1.4 Crop Residue	N ₂ O	0.361	0.311	0.0877	0.010	0.861
Energy	Stationary Combustion, Coal	CO ₂	23.834	10.005	0.0873	0.009	0.871
Energy	Transport, Industry (mobile)	N ₂ O	0.011	0.013	0.0830	0.009	0.880
Energy	Stationary Combustion, Natural gas	CO ₂	4.335	8.293	0.0801	0.009	0.889
Energy	1.B.2 Refinery processes	CH ₄	0.001	0.047	0.0777	0.008	0.897
LULUCF	5.D Wetlands, Soils	CO ₂	0.085	0.002	0.0749	0.008	0.905
Agriculture	4.D3 Atmospheric deposition	N ₂ O	0.496	0.295	0.0617	0.007	0.912
Energy	Stationary Combustion, WASTE	N ₂ O	0.007	0.016	0.0603	0.007	0.918
Energy	Transport, Commercial/institutional	CO ₂	0.074	0.171	0.0575	0.006	0.925
Energy	Transport, Road transport	N ₂ O	0.091	0.116	0.0366	0.004	0.928
Energy	Transport, Navigation (small boats)	CO ₂	0.048	0.099	0.0366	0.004	0.932
Energy	Stationary Combustion, Residual oil	CO ₂	2.496	0.571	0.0357	0.004	0.936
LULUCF	5.E Settlements, Living biomass	CO ₂	0.011	0.056	0.0330	0.004	0.940
Industrial Proc.	2F Consumption of SF ₆	SF ₆	0.107	0.118	0.0304	0.003	0.943
Energy	Transport, Navigation (large vessels)	N ₂ O	0.015	0.008	0.0297	0.003	0.946
Agriculture	4.B Manure Management	N ₂ O	0.600	0.391	0.0272	0.003	0.949
Agriculture	4.D1.6 Sewage sludge and Industrial waste used as fertiliser	N ₂ O	0.028	0.039	0.0269	0.003	0.952
Energy	Stationary Combustion, Petroleum coke	CO ₂	0.415	0.628	0.0249	0.003	0.955
Agriculture	4.D.2 Grassing animals	N ₂ O	0.334	0.211	0.0248	0.003	0.958
Waste	6 B. Wastewater Handling - Indirect	N ₂ O	0.082	0.032	0.0236	0.003	0.960
LULUCF	5.E Settlements, Soils	CO ₂	0.001	0.035	0.0234	0.003	0.963
Energy	Transport, Fisheries	N ₂ O	0.011	0.009	0.0204	0.002	0.965
Solvent and Other Prod. Use	3D5 Use of candles	CO ₂	0.022	0.081	0.0199	0.002	0.967
Energy	Transport, Navigation (large vessels)	CO ₂	0.748	0.399	0.0183	0.002	0.969
Waste	6 B. Wastewater Handling - Direct	N ₂ O	0.023	0.041	0.0183	0.002	0.971
Energy	Stationary Combustion, Kerosene	CO ₂	0.366	0.002	0.0180	0.002	0.973
Energy	Transport, Residential	CO ₂	0.039	0.062	0.0170	0.002	0.975
Waste	6 B. Wastewater Handling	CH ₄	0.065	0.074	0.0156	0.002	0.977
Waste	6.D Accidental fires, buildings	CO ₂	0.011	0.011	0.0120	0.001	0.978
Energy	Transport, Road transport	CH ₄	0.047	0.011	0.0112	0.001	0.979
Energy	Transport, Navigation (small boats)	N ₂ O	0.000	0.001	0.0110	0.001	0.980
Energy	1.B.2 Off-shore activities	CH ₄	0.015	0.037	0.0107	0.001	0.981
Energy	Stationary Combustion, Refinery gas	CO ₂	0.816	0.906	0.0104	0.001	0.983
Waste	6.D Accidental fires, vehicles	CO ₂	0.006	0.006	0.0092	0.001	0.984

Tier 2 Analysis IPCC Source Categories (LU- LUCF included)		DK GHG	Base Year Estimate Ex,o Mt CO ₂ eq	Latest Year Estimate Ex,t Mt CO ₂ eq	Trend Asses- ment Tx,t	Contribu- tion to Trend	Cumula- tive
Industrial Proc.	2A1 Cement production	CO ₂	0.882	0.871	0.0080	0.001	0.984
Energy	Transport, Commercial/institutional	N ₂ O	0.000	0.001	0.0079	0.001	0.985
Energy	Stationary Combustion, SOLID	CH ₄	0.013	0.003	0.0077	0.001	0.986
LULUCF	5.C Grassland, Organic soils	CO ₂	0.107	0.079	0.0070	0.001	0.987
LULUCF	5.B Cropland, Dead organic matter	CO ₂	0.003	0.012	0.0070	0.001	0.988
Energy	Transport, Military	N ₂ O	0.001	0.001	0.0065	0.001	0.988
LULUCF	5.B Cropland, Organic soils	CO ₂	2.887	1.981	0.0057	0.001	0.989
Industrial Proc.	2F Consumption of PFC	PFC	0.001	0.009	0.0056	0.001	0.990
Energy	Transport, Fisheries	CO ₂	0.591	0.479	0.0054	0.001	0.990
Energy	Transport, Railways	N ₂ O	0.003	0.002	0.0053	0.001	0.991
Energy	Transport, Civil aviation	CO ₂	0.243	0.133	0.0050	0.001	0.991
Energy	Stationary Combustion, GAS	CH ₄	0.003	0.006	0.0046	0.001	0.992
LULUCF	5.C Grassland, Mineral soils	CO ₂	0.000	0.005	0.0045	0.000	0.992
Energy	Biogas fuelled engines, BIOMASS	CH ₄	0.001	0.031	0.0044	0.000	0.993
LULUCF	5.C Grassland, Dead organic matter	CO ₂	0.002	0.007	0.0041	0.000	0.993
Industrial Proc.	2A2 Lime production	CO ₂	0.116	0.040	0.0037	0.000	0.994
Energy	Natural gas fuelled engines, GAS	CH ₄	0.005	0.120	0.0035	0.000	0.994
LULUCF	5.A.2 Land converted to forest	CO ₂	0.077	0.038	0.0034	0.000	0.994
Energy	Transport, Railways	CO ₂	0.297	0.249	0.0033	0.000	0.995
Energy	1.B.2 Land based activities	CH ₄	0.017	0.018	0.0031	0.000	0.995
Energy	Transport, Forestry	CO ₂	0.036	0.017	0.0031	0.000	0.995
Energy	Transport, Residential	N ₂ O	0.000	0.000	0.0027	0.000	0.996
Waste	6.D Accidental fires, buildings	CH ₄	0.001	0.001	0.0025	0.000	0.996
Energy	Transport, Commercial/institutional	CH ₄	0.002	0.003	0.0025	0.000	0.996
Energy	Stationary Combustion, LPG	CO ₂	0.184	0.091	0.0025	0.000	0.996
Energy	Transport, Military	CO ₂	0.119	0.116	0.0025	0.000	0.997
Solvent and Other Prod. Use	3D5 Consumption of fireworks	N ₂ O	0.001	0.002	0.0021	0.000	0.997
Industrial Proc.	2C1 Iron and steel production	CO ₂	0.028	0.000	0.0019	0.000	0.997
Industrial Proc.	2A3 Limestone and dolomite use	CO ₂	0.014	0.026	0.0015	0.000	0.997
Energy	Transport, Civil aviation	N ₂ O	0.003	0.002	0.0015	0.000	0.998
Energy	Stationary Combustion, Coke	CO ₂	0.138	0.074	0.0015	0.000	0.998
Energy	Stationary Combustion, WASTE	CH ₄	0.001	0.002	0.0014	0.000	0.998
LULUCF	5.D Wetlands, Living biomass	CO ₂	0.003	0.000	0.0014	0.000	0.998
Energy	1.B.2 Flaring off-shore	CO ₂	0.302	0.195	0.0013	0.000	0.998
Energy	Stationary Combustion, LIQUID	CH ₄	0.003	0.001	0.0010	0.000	0.998
LULUCF	5(III) Disturbance, Land converted to cropland	N ₂ O	0.000	0.001	0.0010	0.000	0.998
Energy	1.B.2 Flaring in refinery	CO ₂	0.023	0.022	0.0010	0.000	0.998
Solvent and Other Prod. Use	3D1 Other - Use of N ₂ O for Anaesthesia	N ₂ O	0.000	0.009	0.0009	0.000	0.999
Agriculture	4F Field burning of agricultural residues	CH ₄	0.002	0.002	0.0009	0.000	0.999
Energy	Transport, Residential	CH ₄	0.001	0.001	0.0009	0.000	0.999
Energy	1.B.2 Off-shore activities	CO ₂	0.002	0.004	0.0008	0.000	0.999
Energy	Transport, Agriculture	CH ₄	0.002	0.002	0.0007	0.000	0.999
Solvent and Other Prod. Use	3D5 Other	CO ₂	0.061	0.044	0.0007	0.000	0.999
Energy	Transport, Forestry	N ₂ O	0.000	0.000	0.0007	0.000	0.999
LULUCF	5(II) Forest Land.	N ₂ O	0.016	0.012	0.0006	0.000	0.999
Waste	6.D Accidental fires, vehicles	CH ₄	0.000	0.000	0.0006	0.000	0.999
LULUCF	5.D Wetlands, Dead organic matter	CO ₂	0.001	0.000	0.0005	0.000	0.999
Energy	1.B.2 Transmission of natural gas	CH ₄	0.004	0.000	0.0005	0.000	0.999
Energy	Stationary Combustion, BKB	CO ₂	0.011	0.001	0.0005	0.000	0.999
Agriculture	4.D1.5 Cultivation of histosols	N ₂ O	0.290	0.198	0.0004	0.000	0.999
Solvent and Other Prod. Use	3A Paint application	CO ₂	0.013	0.007	0.0004	0.000	0.999
Energy	Transport, Navigation (small boats)	CH ₄	0.000	0.001	0.0004	0.000	0.999
Energy	Transport, Forestry	CH ₄	0.000	0.000	0.0004	0.000	1.000
Agriculture	4.F Field Burning of Agricultural Residues	N ₂ O	0.001	0.001	0.0003	0.000	1.000
Industrial Proc.	2A7c Expanded clay	CO ₂	0.015	0.006	0.0003	0.000	1.000
Energy	1.B.2. Flaring off-shore	CH ₄	0.001	0.000	0.0003	0.000	1.000
LULUCF	5(V) Biomass Burning	CH ₄	0.001	0.000	0.0003	0.000	1.000
Solvent and Other Prod. Use	3C Chemical products, manufacturing and processing	CO ₂	0.019	0.012	0.0003	0.000	1.000

Tier 2 Analysis IPCC Source Categories (LU- LUCF included)		DK GHG	Base Year Estimate Ex,o Mt CO ₂ eq	Latest Year Estimate Ex,t Mt CO ₂ eq	Trend Asses- sment Tx,t	Contribu- tion to Trend	Cumula- tive
Energy	1.B.2 Flaring in refinery	N ₂ O	0.000	0.000	0.0003	0.000	1.000
LULUCF	5.E Settlements, Dead organic matter	CO ₂	0.001	0.000	0.0002	0.000	1.000
Energy	1.B.2 Flaring off-shore	N ₂ O	0.001	0.000	0.0002	0.000	1.000
Energy	1.B.2 Distribution of natural gas	CH ₄	0.005	0.003	0.0002	0.000	1.000
LULUCF	5(V) Biomass Burning	N ₂ O	0.000	0.000	0.0002	0.000	1.000
Industrial Proc.	2A6 Road paving with asphalt	CO ₂	0.002	0.002	0.0002	0.000	1.000
Industrial Proc.	2G Lubricants	CO ₂	0.050	0.032	0.0002	0.000	1.000
Waste	6.C Incineration of carcasses	N ₂ O	0.000	0.000	0.0002	0.000	1.000
Industrial Proc.	2A7a Glass and Glass wool	CO ₂	0.017	0.010	0.0002	0.000	1.000
Energy	Transport, Industry (mobile)	CH ₄	0.001	0.001	0.0002	0.000	1.000
Solvent and Other Prod. Use	3D5 Consumption of fireworks	CO ₂	0.000	0.000	0.0002	0.000	1.000
Industrial Proc.	2A7b Yellow bricks	CO ₂	0.023	0.017	0.0001	0.000	1.000
Waste	6.C Incineration of corpses	N ₂ O	0.000	0.000	0.0001	0.000	1.000
Solvent and Other Prod. Use	3D5 Use of charcoal for BBQ	N ₂ O	0.000	0.000	0.0001	0.000	1.000
Industrial Proc.	2B5 Catalysts/Fertilizers, Pesticides and Sulphuric acid	CO ₂	0.001	0.001	0.0001	0.000	1.000
Energy	Transport, Fisheries	CH ₄	0.000	0.000	0.0001	0.000	1.000
Industrial Proc.	2D2 Food and Drink	CO ₂	0.004	0.002	0.0001	0.000	1.000
Energy	1.B.2 Venting in gas storage	CH ₄	0.001	0.001	0.0001	0.000	1.000
LULUCF	5(II) Wetlands	N ₂ O	0.000	0.000	0.0001	0.000	1.000
Solvent and Other Prod. Use	3D5 Use of candles	N ₂ O	0.000	0.000	0.0001	0.000	1.000
Energy	Transport, Navigation (large vessels)	CH ₄	0.000	0.000	0.0000	0.000	1.000
Energy	Transport, Railways	CH ₄	0.000	0.000	0.0000	0.000	1.000
Energy	Transport, Civil aviation	CH ₄	0.000	0.000	0.0000	0.000	1.000
Energy	Transport, Military	CH ₄	0.000	0.000	0.0000	0.000	1.000
Energy	1.B.2. Flaring in refinery	CH ₄	0.000	0.000	0.0000	0.000	1.000
Waste	6.C Incineration of carcasses	CH ₄	0.000	0.000	0.0000	0.000	1.000
Waste	6.C Incineration of corpses	CH ₄	0.000	0.000	0.0000	0.000	1.000
Solvent and Other Prod. Use	3D5 Use of tobacco	N ₂ O	0.000	0.000	0.0000	0.000	1.000
Industrial Proc.	2A5 Asphalt roofing	CO ₂	0.000	0.000	0.0000	0.000	1.000
Energy	1.B.2 Land based activities	CO ₂	0.000	0.000	0.0000	0.000	1.000
Energy	1.B.2 Distribution of natural gas	CO ₂	0.000	0.000	0.0000	0.000	1.000
Energy	1.B.2 Transmission of natural gas	CO ₂	0.000	0.000	0.0000	0.000	1.000
Energy	1.B.2 Venting in gas storage	CO ₂	0.000	0.000	0.0000	0.000	1.000
Solvent and Other Prod. Use	3B Degreasing and dry cleaning	CO ₂	0.000	0.000	0.0000	0.000	1.000
Total			74.225	50.799			

¹⁾ The Estimates include signs, where + : emission - : removal, although in the level analyses only the absolute values are used.

Annex 2 - Detailed discussion of methodology and data for estimation of CO₂ emission from fossil fuel combustion

Please refer to Annex 3A and 3B.

Annex 3 - Other detailed methodological descriptions for individual source or sink categories (where relevant)

Annex 3A – Stationary Combustion

Annex 3B – Transport

Annex 3C – Industrial Processes

Annex 3D – Solvents and Other Product Use

Annex 3E – Agriculture

Annex 3F – LULUCF

Annex 3G – Waste

Annex 3A - Stationary combustion

Annex 3A-1:	Correspondence list between SNAP and CRF source categories
Annex 3A-2:	Fuel rate
Annex 3A-3:	Default Lower Calorific Value (LCV) of fuels and fuel correspondence list
Annex 3A-4:	Emission factors
Annex 3A-5:	Large point sources
Annex 3A-6:	Adjustment of CO ₂ emission
Annex 3A-7:	Uncertainty estimates
Annex 3A-8:	Emission inventory 2012 based on SNAP sectors
Annex 3A-9:	Description of the Danish energy statistics
Annex 3A-10	EU ETS data

Annex 3A-1 Correspondence list between SNAP and CRF source categories

Table 3A-1.1 Correspondence list between SNAP and CRF source categories for stationary combustion.

SNAP_id	SNAP	CRF_id	CRF_name
010100	Public power	1A1a	Electricity and heat production
010101	Combustion plants >= 300 MW (boilers)	1A1a	Electricity and heat production
010102	Combustion plants >= 50 and < 300 MW (boilers)	1A1a	Electricity and heat production
010103	Combustion plants < 50 MW (boilers)	1A1a	Electricity and heat production
010104	Gas turbines	1A1a	Electricity and heat production
010105	Stationary engines	1A1a	Electricity and heat production
010200	District heating plants	1A1a	Electricity and heat production
010201	Combustion plants >= 300 MW (boilers)	1A1a	Electricity and heat production
010202	Combustion plants >= 50 and < 300 MW (boilers)	1A1a	Electricity and heat production
010203	Combustion plants < 50 MW (boilers)	1A1a	Electricity and heat production
010204	Gas turbines	1A1a	Electricity and heat production
010205	Stationary engines	1A1a	Electricity and heat production
010300	Petroleum refining plants	1A1b	Petroleum refining
010301	Combustion plants >= 300 MW (boilers)	1A1b	Petroleum refining
010302	Combustion plants >= 50 and < 300 MW (boilers)	1A1b	Petroleum refining
010303	Combustion plants < 50 MW (boilers)	1A1b	Petroleum refining
010304	Gas turbines	1A1b	Petroleum refining
010305	Stationary engines	1A1b	Petroleum refining
010306	Process furnaces	1A1b	Petroleum refining
010400	Solid fuel transformation plants	1A1c	Manufacture of solid fuels and other energy industries
010401	Combustion plants >= 300 MW (boilers)	1A1c	Manufacture of solid fuels and other energy industries
010402	Combustion plants >= 50 and < 300 MW (boilers)	1A1c	Manufacture of solid fuels and other energy industries
010403	Combustion plants < 50 MW (boilers)	1A1c	Manufacture of solid fuels and other energy industries
010404	Gas turbines	1A1c	Manufacture of solid fuels and other energy industries
010405	Stationary engines	1A1c	Manufacture of solid fuels and other energy industries
010406	Coke oven furnaces	1A1c	Manufacture of solid fuels and other energy industries
010407	Other (coal gasification, liquefaction, ...)	1A1c	Manufacture of solid fuels and other energy industries
010500	Coal mining, oil/gas extraction, pipeline compressors	1A1c	Manufacture of solid fuels and other energy industries
010501	Combustion plants >= 300 MW (boilers)	1A1c	Manufacture of solid fuels and other energy industries
010502	Combustion plants >= 50 and < 300 MW (boilers)	1A1c	Manufacture of solid fuels and other energy industries
010503	Combustion plants < 50 MW (boilers)	1A1c	Manufacture of solid fuels and other energy industries
010504	Gas turbines	1A1c	Manufacture of solid fuels and other energy industries
010505	Stationary engines	1A1c	Manufacture of solid fuels and other energy industries
020100	Commercial and institutional plants (t)	1A4a i	Commercial/Institutional plants
020101	Combustion plants >= 300 MW (boilers)	1A4a i	Commercial/Institutional plants
020102	Combustion plants >= 50 and < 300 MW (boilers)	1A4a i	Commercial/Institutional plants
020103	Combustion plants < 50 MW (boilers)	1A4a i	Commercial/Institutional plants
020104	Stationary gas turbines	1A4a i	Commercial/Institutional plants
020105	Stationary engines	1A4a i	Commercial/Institutional plants
020106	Other stationary equipments (n)	1A4a i	Commercial/Institutional plants
020200	Residential plants	1A4b i	Residential plants
020201	Combustion plants >= 50 MW (boilers)	1A4b i	Residential plants
020202	Combustion plants < 50 MW (boilers)	1A4b i	Residential plants
020203	Gas turbines	1A4b i	Residential plants
020204	Stationary engines	1A4b i	Residential plants
020205	Other equipments (stoves, fireplaces, cooking,...) ¹⁾	1A4b i	Residential plants
020300	Plants in agriculture, forestry and aquaculture	1A4c i	Agriculture/Forestry/Fishing, Stationary
020301	Combustion plants >= 50 MW (boilers)	1A4c i	Agriculture/Forestry/Fishing, Stationary
020302	Combustion plants < 50 MW (boilers)	1A4c i	Agriculture/Forestry/Fishing, Stationary
020303	Stationary gas turbines	1A4c i	Agriculture/Forestry/Fishing, Stationary
020304	Stationary engines	1A4c i	Agriculture/Forestry/Fishing, Stationary
020305	Other stationary equipments (n)	1A4c i	Agriculture/Forestry/Fishing, Stationary
030100	Comb. in boilers, gas turbines and stationary	1A2f i	Industry-Other
030101	Combustion plants >= 300 MW (boilers)	1A2f i	Industry-Other
030102	Combustion plants >= 50 and < 300 MW (boilers)	1A2f i	Industry-Other
030103	Combustion plants < 50 MW (boilers)	1A2f i	Industry-Other
030104	Gas turbines	1A2f i	Industry-Other
030105	Stationary engines	1A2f i	Industry-Other
030106	Other stationary equipments (n)	1A2f i	Industry-Other
030200	Process furnaces without contact	1A2f i	Industry-Other
030203	Blast furnace cowpers	1A2a	Industry-Iron and steel
030204	Plaster furnaces	1A2f i	Industry-Other
030205	Other furnaces	1A2f i	Industry-Other
030400	Iron and steel	1A2a	Iron and steel
030401	Combustion plants >= 300 MW (boilers)	1A2a	Iron and steel
030402	Combustion plants >= 50 and < 300 MW (boilers)	1A2a	Iron and steel
030403	Combustion plants < 50 MW (boilers)	1A2a	Iron and steel
030404	Gas turbines	1A2a	Iron and steel
030405	Stationary engines	1A2a	Iron and steel

SNAP_id	SNAP	CRF_id	CRF_name
030406	Other stationary equipments (n)	1A2a	Iron and steel
030500	Non-ferrous metals	1A2b	Non-ferrous metals
030501	Combustion plants >= 300 MW (boilers)	1A2b	Non-ferrous metals
030502	Combustion plants >= 50 and < 300 MW (boilers)	1A2b	Non-ferrous metals
030503	Combustion plants < 50 MW (boilers)	1A2b	Non-ferrous metals
030504	Gas turbines	1A2b	Non-ferrous metals
030505	Stationary engines	1A2b	Non-ferrous metals
030506	Other stationary equipments (n)	1A2b	Non-ferrous metals
030600	Chemical and petrochemical	1A2c	Chemicals
030601	Combustion plants >= 300 MW (boilers)	1A2c	Chemicals
030602	Combustion plants >= 50 and < 300 MW (boilers)	1A2c	Chemicals
030603	Combustion plants < 50 MW (boilers)	1A2c	Chemicals
030604	Gas turbines	1A2c	Chemicals
030605	Stationary engines	1A2c	Chemicals
030606	Other stationary equipments (n)	1A2c	Chemicals
030700	Non-metallic minerals	1A2f i	Industry-Other
030701	Combustion plants >= 300 MW (boilers)	1A2f i	Industry-Other
030702	Combustion plants >= 50 and < 300 MW (boilers)	1A2f i	Industry-Other
030703	Combustion plants < 50 MW (boilers)	1A2f i	Industry-Other
030704	Gas turbines	1A2f i	Industry-Other
030705	Stationary engines	1A2f i	Industry-Other
030706	Other stationary equipments (n)	1A2f i	Industry-Other
030800	Mining and Quarrying	1A2f i	Industry-Other
030801	Combustion plants >= 300 MW (boilers)	1A2f i	Industry-Other
030802	Combustion plants >= 50 and < 300 MW (boilers)	1A2f i	Industry-Other
030803	Combustion plants < 50 MW (boilers)	1A2f i	Industry-Other
030804	Gas turbines	1A2f i	Industry-Other
030805	Stationary engines	1A2f i	Industry-Other
030806	Other stationary equipments (n)	1A2f i	Industry-Other
030900	Food and tobacco	1A2e	Food processing, beverages and tobacco
030901	Combustion plants >= 300 MW (boilers)	1A2e	Food processing, beverages and tobacco
030902	Combustion plants >= 50 and < 300 MW (boilers)	1A2e	Food processing, beverages and tobacco
030903	Combustion plants < 50 MW (boilers)	1A2e	Food processing, beverages and tobacco
030904	Gas turbines	1A2e	Food processing, beverages and tobacco
030905	Stationary engines	1A2e	Food processing, beverages and tobacco
030906	Other stationary equipments (n)	1A2e	Food processing, beverages and tobacco
031000	Textile and leather	1A2f i	Industry-Other
031001	Combustion plants >= 300 MW (boilers)	1A2f i	Industry-Other
031002	Combustion plants >= 50 and < 300 MW (boilers)	1A2f i	Industry-Other
031003	Combustion plants < 50 MW (boilers)	1A2f i	Industry-Other
031004	Gas turbines	1A2f i	Industry-Other
031005	Stationary engines	1A2f i	Industry-Other
031006	Other stationary equipments (n)	1A2f i	Industry-Other
031100	Paper, pulp and print	1A2d	Pulp, Paper and Print
031101	Combustion plants >= 300 MW (boilers)	1A2d	Pulp, Paper and Print
031102	Combustion plants >= 50 and < 300 MW (boilers)	1A2d	Pulp, Paper and Print
031103	Combustion plants < 50 MW (boilers)	1A2d	Pulp, Paper and Print
031104	Gas turbines	1A2d	Pulp, Paper and Print
031105	Stationary engines	1A2d	Pulp, Paper and Print
031106	Other stationary equipments (n)	1A2d	Pulp, Paper and Print
031200	Transport equipment	1A2f i	Industry-Other
031201	Combustion plants >= 300 MW (boilers)	1A2f i	Industry-Other
031202	Combustion plants >= 50 and < 300 MW (boilers)	1A2f i	Industry-Other
031203	Combustion plants < 50 MW (boilers)	1A2f i	Industry-Other
031204	Gas turbines	1A2f i	Industry-Other
031205	Stationary engines	1A2f i	Industry-Other
031206	Other stationary equipments (n)	1A2f i	Industry-Other
031300	Machinery	1A2f i	Industry-Other
031301	Combustion plants >= 300 MW (boilers)	1A2f i	Industry-Other
031302	Combustion plants >= 50 and < 300 MW (boilers)	1A2f i	Industry-Other
031303	Combustion plants < 50 MW (boilers)	1A2f i	Industry-Other
031304	Gas turbines	1A2f i	Industry-Other
031305	Stationary engines	1A2f i	Industry-Other
031306	Other stationary equipments (n)	1A2f i	Industry-Other
031400	Wood and wood products	1A2f i	Industry-Other
031401	Combustion plants >= 300 MW (boilers)	1A2f i	Industry-Other
031402	Combustion plants >= 50 and < 300 MW (boilers)	1A2f i	Industry-Other
031403	Combustion plants < 50 MW (boilers)	1A2f i	Industry-Other
031404	Gas turbines	1A2f i	Industry-Other
031405	Stationary engines	1A2f i	Industry-Other
031406	Other stationary equipments (n)	1A2f i	Industry-Other
031500	Construction	1A2f i	Industry-Other
031501	Combustion plants >= 300 MW (boilers)	1A2f i	Industry-Other

SNAP_id	SNAP	CRF_id	CRF_name
031502	Combustion plants >= 50 and < 300 MW (boilers)	1A2f i	Industry-Other
031503	Combustion plants < 50 MW (boilers)	1A2f i	Industry-Other
031504	Gas turbines	1A2f i	Industry-Other
031505	Stationary engines	1A2f i	Industry-Other
031506	Other stationary equipments (n)	1A2f i	Industry-Other
031600	Cement production	1A2f i	Industry-Other
031601	Combustion plants >= 300 MW (boilers)	1A2f i	Industry-Other
031602	Combustion plants >= 50 and < 300 MW (boilers)	1A2f i	Industry-Other
031603	Combustion plants < 50 MW (boilers)	1A2f i	Industry-Other
031604	Gas turbines	1A2f i	Industry-Other
031605	Stationary engines	1A2f i	Industry-Other
031606	Other stationary equipments (n)	1A2f i	Industry-Other
032000	Non-specified (industry)	1A2f i	Industry-Other
032001	Combustion plants >= 300 MW (boilers)	1A2f i	Industry-Other
032002	Combustion plants >= 50 and < 300 MW (boilers)	1A2f i	Industry-Other
032003	Combustion plants < 50 MW (boilers)	1A2f i	Industry-Other
032004	Gas turbines	1A2f i	Industry-Other
032005	Stationary engines	1A2f i	Industry-Other
032006	Other stationary equipments (n)	1A2f i	Industry-Other

¹⁾ Stoves, fireplaces and cooking is included in the sector 020200 or 020202 in the Danish inventory.

Annex 3A-2 Fuel rate

Table 3A-2.1 Fuel consumption rate for stationary combustion plants 1990-2012, PJ.

Sum of Fuel_rate_PJ			Year									
fuel_type	fuel_id	fuel_gr_abbr	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
SOLID	102A	COAL	253.4	344.3	286.8	300.8	323.4	270.3	371.9	276.3	234.3	196.5
	106A	BROWN COAL BRI.	0.1	0.2	0.1	0.1	0.1	0.1	0.1	0.1	0.0	0.0
	107A	COKE OVEN COKE	1.3	1.4	1.2	1.2	1.2	1.3	1.2	1.3	1.3	1.4
LIQUID	110A	PETROLEUM COKE	4.5	4.4	4.3	5.7	7.5	5.3	5.9	6.0	5.3	6.8
	203A	RESIDUAL OIL	32.1	38.3	38.5	32.8	46.2	33.0	37.8	26.6	30.0	23.7
	204A	GAS OIL	61.4	65.0	56.1	62.0	53.9	53.7	58.0	51.1	48.4	47.5
	206A	KEROSENE	5.1	0.9	0.8	0.8	0.6	0.6	0.5	0.4	0.4	0.3
	225A	ORIMULSION						19.9	36.8	40.5	32.6	34.2
	303A	LPG	2.9	2.7	2.4	2.5	2.5	2.7	3.0	2.6	2.8	2.5
	308A	REFINERY GAS	14.2	14.5	14.9	15.4	16.4	20.8	21.4	16.9	15.2	15.7
GAS	301A	NATURAL GAS	76.1	86.1	90.5	102.5	114.6	132.7	156.3	164.5	178.7	187.9
WASTE	114A	WASTE	15.5	16.7	17.8	19.4	20.3	22.9	25.0	26.8	26.6	29.1
BIOMASS	111A	WOOD	18.2	20.0	21.0	22.2	21.9	21.8	23.4	23.4	22.9	24.4
	117A	STRAW	12.5	13.3	13.9	13.4	12.7	13.1	13.5	13.9	13.9	13.7
	215A	BIO OIL	0.7	0.7	0.7	0.8	0.2	0.3	0.1	0.0	0.0	0.0
	309A	BIOGAS	0.8	0.9	0.9	1.1	1.3	1.8	2.0	2.4	2.6	2.6
	310A	BIO PROD GAS					0.1	0.0	0.0	0.0	0.0	0.0
Grand Total			498.8	609.6	549.9	580.6	623.0	600.2	756.8	652.7	615.2	586.4
Sum of Fuel_rate_PJ			Year									
fuel_type	fuel_id	fuel_gr_abbr	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
SOLID	101A	ANODE CARBON										0.0
	102A	COAL	164.7	174.3	174.7	239.0	182.5	154.0	232.0	194.1	170.5	167.7
	106A	BROWN COAL BRI.	0.0	0.0	0.0	0.0					0.0	0.0
	107A	COKE OVEN COKE	1.2	1.1	1.1	1.0	1.1	1.0	1.0	1.1	1.0	0.8
LIQUID	110A	PETROLEUM COKE	6.8	7.8	7.8	8.0	8.4	8.1	8.5	9.2	6.9	5.9
	203A	RESIDUAL OIL	18.8	21.1	26.2	28.6	24.5	21.9	26.1	19.8	15.8	14.7
	204A	GAS OIL	41.3	43.6	38.6	38.9	35.8	31.6	26.5	21.3	20.4	23.3
	206A	KEROSENE	0.2	0.3	0.3	0.3	0.2	0.3	0.2	0.1	0.1	0.1
	225A	ORIMULSION	34.1	30.2	23.8	1.9	0.0					
	303A	LPG	2.4	2.1	2.0	2.1	2.1	2.2	2.2	1.9	1.7	1.5
	308A	REFINERY GAS	15.6	15.8	15.2	16.6	15.9	15.3	16.1	15.9	14.8	15.4
GAS	301A	NATURAL GAS	186.1	193.8	193.6	195.9	195.1	187.4	191.1	171.0	171.9	164.9
WASTE	114A	WASTE	29.8	31.3	33.3	35.1	35.3	35.8	36.9	38.1	39.6	37.6
	115A	INDUSTR. WASTES	0.5	1.4	1.9	1.5	2.0	2.0	1.5	1.6	2.0	1.7
BIOMASS	111A	WOOD	27.5	30.8	31.6	38.9	43.9	49.7	52.1	60.3	63.6	66.0
	117A	STRAW	12.2	13.7	15.7	16.9	17.9	18.5	18.5	18.8	15.9	17.4
	215A	BIO OIL	0.0	0.2	0.1	0.4	0.6	0.8	1.1	1.2	1.8	1.7
	309A	BIOGAS	2.9	3.0	3.3	3.5	3.7	3.8	3.9	3.9	3.9	4.2
	310A	BIO PROD GAS	0.0	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.3
Grand Total			544.3	570.7	569.1	628.6	569.2	532.5	618.0	558.4	530.1	523.2
Sum of Fuel_rate_PJ			Year									
fuel_type	fuel_id	fuel_gr_abbr	2010	2011	2012							
SOLID	101A	ANODE CARBON	0.0	0.0	0.0							
	102A	COAL	163.0	135.5	106.2							
	103A	SUB-BITUMINOUS		0.0	0.1							
	106A	BROWN COAL BRI.	0.0	0.0	0.0							
	107A	COKE OVEN COKE	0.7	0.7	0.6							
LIQUID	110A	PETROLEUM COKE	5.1	6.5	6.7							
	203A	RESIDUAL OIL	13.0	8.0	7.3							
	204A	GAS OIL	21.5	14.6	10.0							
	206A	KEROSENE	0.1	0.0	0.0							
	303A	LPG	1.4	1.3	1.4							
	308A	REFINERY GAS	14.3	15.0	15.6							
GAS	301A	NATURAL GAS	184.9	157.7	145.2							
WASTE	114A	WASTE	36.4	36.9	36.0							
	115A	INDUSTR. WASTES	1.4	1.7	1.5							
BIOMASS	111A	WOOD	81.3	79.3	84.7							
	117A	STRAW	23.3	19.4	17.5							
	215A	BIO OIL	2.0	0.8	1.1							
	309A	BIOGAS	4.4	4.2	4.4							
	310A	BIO PROD GAS	0.2	0.3	0.4							
Grand Total			553.0	481.9	438.8							

Table 3A-2.2 Detailed fuel consumption data for stationary combustion plants, 1990-2012, PJ.

Sum of Fuel_rate_PJ				Year												
fuel_type	fuel_gr_abbr	nfr_id	nfr_name	snap_id	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999		
SOLID	COAL	1A1a	Electricity and heat production	010100	8.5	12.9	10.2	8.2								
				010101	207.9	294.7	241.8	256.3	284.7	233.2	333.6	244.3	206.2	172.0		
				010102	14.0	11.0	13.2	15.4	18.9	19.4	22.6	17.1	14.2	12.8		
				010103					0.5	0.4	0.1					
				010104					0.3	0.3	0.3	0.1				
				010105					0.0							
				010200	6.0	6.6	5.2	3.6								
				010202					1.1	0.7						
				010203					1.4	1.0	0.7	0.2	0.1	0.0		
		1A2c	Chemicals	030600	0.1	0.1	0.1	0.7	0.7	0.6	0.6	0.5	0.5	0.5		
		1A2d	Pulp, Paper and Print	031100	1.3	1.7	1.1	0.7	0.7	0.0						
				031102						0.1	0.0					
		1A2e	Food processing, beverages and tobacco	030900	4.0	4.0	3.1	3.4	2.5	2.3	1.6	1.4	1.8	1.0		
				030902					0.5	1.0	1.4	1.5	1.4	1.4		
				030903					0.3	0.4	0.4	0.5	0.3	0.2		
		1A2f i	Industry-Other	030700	0.2			0.2	0.3	0.3	0.8	0.6	0.7	1.1		
				030800	1.6	1.9	1.7	1.9	1.6	1.3	1.3	1.5	1.4	0.9		
				031200	0.0	0.0	0.0	0.0								
				031300	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0			
				031600	5.0	6.0	6.6	6.6	6.9	7.2	7.1	7.2	6.6	5.6		
				032000	1.6	1.2	0.7	0.8								
				1A4a i	Commercial/Institutional plants	020100	0.1	0.0	0.1	0.1	0.1	0.1	0.0	0.0	0.0	
		1A4b i	Residential plants	020200	0.6	1.1	0.9	0.8	0.6	0.4	0.1	0.1	0.1	0.1		
		1A4c i	Agriculture/Forestry/Fishing, Stationary	020300	2.5	2.9	2.2	2.1	2.3	1.8	1.4	1.2	0.9	0.7		
		BROWN COAL BRI.	1A2f i	Industry-Other	030800	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0			
			1A4a i	Commercial/Institutional plants	020100	0.0	0.0		0.0	0.0	0.0	0.0	0.0			
			1A4b i	Residential plants	020200	0.1	0.1	0.0	0.1	0.1	0.1	0.0	0.0	0.0	0.0	
			1A4c i	Agriculture/Forestry/Fishing, Stationary	020300	0.1	0.1	0.1	0.0	0.0	0.0	0.0	0.0	0.0		
		COKE OVEN COKE	1A2a	Iron and steel	030400	0.0	0.0								0.0	
			1A2e	Food processing, beverages and tobacco	030900	0.2	0.2	0.2	0.2	0.2	0.1	0.2	0.2	0.2	0.2	
		LIQUID	PETROLEUM COKE	1A2f i	Industry-Other	030700	0.8	1.0	0.9	0.8	0.9	0.1	0.1	0.1	0.1	0.1
						030800	0.0								0.0	0.0
						031200	0.0	0.0				0.0	0.0	0.0	0.0	
				031300	0.1	0.1	0.1	0.0	0.0	0.0	0.0	0.0	0.0			
				031400	0.0	0.0	0.0	0.0	0.0	0.0						
				032000	0.0	0.0	0.0	0.0		0.9	0.9	0.9	1.0	1.0		
				1A4b i	Residential plants	020200	0.1	0.1	0.1	0.1	0.1	0.0	0.0	0.0	0.0	0.0
				1A1a	Electricity and heat production	010100				1.2						
						010102					3.7	0.9				
				1A2a	Iron and steel	030400										0.0
				1A2c	Chemicals	030600				0.0						
				1A2d	Pulp, Paper and Print	031100	0.0	0.0	0.0							

Sum of					Year										
Fuel_rate_PJ															
fuel_type	fuel_gr_abbr	nfr_id	nfr_name	snap_id	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	
RESIDUAL OIL		1A2e	Food processing, beverages and tobacco	030900				0.1							
		1A2f i	Industry-Other	030700	0.2						0.1	0.0	0.0	0.0	
				030800	0.1	0.1	0.1	0.0		0.2				0.0	
				031000	0.0	0.0	0.0			0.0					
				031300	0.0	0.0	0.0			0.0	0.0		0.0	0.0	
				031400	0.0	0.0	0.0	0.0		0.0					
				031600	2.5	3.0	3.2	3.2	3.5	3.7	5.0	5.2	4.8	6.4	
				032000						0.0					
		1A4a i	Commercial/Institutional plants	020100	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
		1A4b i	Residential plants	020200	0.8	0.7	0.5	0.5	0.2	0.2	0.4	0.3	0.2	0.2	0.2
		1A4c i	Agriculture/Forestry/Fishing, Stationary	020300	0.8	0.5	0.4	0.4		0.1	0.3	0.3	0.2	0.1	
		1A1a	Electricity and heat production	010100	0.8	0.4	1.8	0.8							
				010101	6.5	9.6	8.3	7.8	21.5	8.5	11.6	5.2	8.9	6.0	
				010102	0.7	0.4	0.5	0.7	0.7	2.5	4.5	2.7	2.8	1.6	
				010103					0.3	0.3	0.0	0.2	0.2	0.0	
				010104					0.0	0.0	0.0	0.0	0.0	0.0	
				010105					0.0	0.0	0.0	0.0	0.0	0.0	
				010200	2.0	2.2	1.1	0.9							
				010202					0.1	0.5	0.5	0.4	0.2	0.1	
				010203					1.1	1.1	1.6	1.3	1.5	1.6	
		1A1b	Petroleum refining	010306	1.3	2.0	3.6	3.5	3.3	2.3	2.2	1.6	1.1	1.1	
		1A2a	Iron and steel	030400	0.0	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
		1A2b	Non-ferrous metals	030500	0.0	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	
		1A2c	Chemicals	030600	2.4	2.6	2.7	2.0	1.9	1.7	1.7	1.6	1.5	0.8	
		1A2d	Pulp, Paper and Print	031100	1.0	1.0	0.7	0.8	0.7	0.6	0.2	0.2	0.2	0.1	
				031102						0.0	0.1				
		1A2e	Food processing, beverages and tobacco	030900	7.4	7.7	7.6	7.0	7.1	5.7	5.8	5.0	5.5	5.5	
				030902					0.8	0.8	0.7	0.7	0.6	0.6	
				030903					0.1	0.2	0.1	0.1	0.1	0.2	
				030904								0.1			
		1A2f i	Industry-Other	030700	1.0	2.8	2.4	0.8	0.9	0.4	0.8	0.5			0.9
				030800	0.4	0.4	0.4	0.4	0.5	0.7	0.6	0.7	0.7	0.6	
				031000	0.5	0.3	0.3	0.2	0.2	0.2	0.2	0.1	0.1	0.0	
				031200	0.2	0.2	0.2	0.2	0.1	0.1	0.0			0.0	
				031300	0.8	0.6	0.7	0.6	0.4	0.2	0.2	0.1	0.2	0.1	
				031400	0.4	0.4	0.3	0.4	0.4	0.4	0.5	0.6	0.5	0.5	
				031403					0.0	0.0					
				031500	1.0	1.5	1.6	0.5	0.2	0.2	0.4	0.2	0.2	0.1	
				031503						0.0					
				031600	1.8	2.2	2.4	2.4	2.6	2.8	1.8	1.9	2.5	0.9	
				032000	1.5	1.5	1.5	1.5	0.2	0.2	0.1	0.1	0.1	0.1	
		1A4a i	Commercial/Institutional plants	020100	1.1	0.9	0.6	0.5	0.7	0.7	0.7	0.7	0.4	0.5	
				020103					0.1	0.1					

Sum of Fuel_rate_PJ		Year											
fuel_type	fuel_gr_abbr	nfr_id nfr_name	snap_id	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
GAS OIL	1A4b i Residential plants	1A4c i Agriculture/Forestry/Fishing, Stationary	020200	0.2	0.2	0.2	0.1	0.1	0.1	0.1	0.0	0.0	0.1
			020300	1.2	1.3	1.6	1.7	1.9	2.6	3.1	2.5	2.6	2.4
			020304									0.0	0.0
	1A1a Electricity and heat production		010100	0.3	0.5	0.7	0.3						
			010101					0.0	0.1	0.0	0.1	0.1	0.3
			010102					0.0	0.0	0.0	0.0	0.0	0.1
			010103						0.0	0.0	0.0	0.0	0.0
			010104		0.0	0.0	0.0	0.0	0.1	0.1	0.0	0.1	0.0
			010105					0.1	0.1	0.1	0.1	0.1	0.1
			010200	1.9	0.8	0.7	0.9						
			010202					0.1	0.2	0.8	0.5	0.4	0.3
			010203					1.0	0.7	0.8	0.7	0.8	0.4
			1A1b Petroleum refining		0.0	0.0	0.0	0.0	0.0	0.0	0.1		
			1A2a Iron and steel	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
			1A2b Non-ferrous metals	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
			1A2c Chemicals	0.0	0.1	0.1	0.0	0.0	0.1	0.1	0.1	0.1	0.1
			1A2d Pulp, Paper and Print	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.0	0.1
			031103							0.0	0.0	0.0	
			1A2e Food processing, beverages and tobacco	0.1	0.4	0.4	0.3	0.2	0.4	0.5	0.4	0.3	0.6
			030902					0.0	0.0	0.0	0.0	0.0	0.0
			030903					0.0	0.0				0.0
			030904								0.0	0.0	0.0
	1A2f i Industry-Other		030700	0.1	0.2	0.2	0.1	0.1	0.2	0.4	0.5	0.5	0.2
			030800	0.0	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.2
			031000	0.0	0.1	0.1	0.0	0.0	0.0	0.1	0.0	0.0	0.0
			031200	0.0	0.0	0.1	0.0	0.0	0.1	0.1	0.1	0.1	0.1
			031300	0.1	0.3	0.3	0.2	0.2	0.4	0.5	0.4	0.4	0.6
			031305									0.0	0.0
			031400	0.0	0.0	0.0	0.0	0.0	0.1	0.1	0.1	0.0	0.1
			031403					0.0	0.0		0.0	0.0	0.0
			031505					0.0					
			032000	0.1	0.1	0.1	0.1	0.1	0.1	0.2	0.2	0.2	0.3
			032005					0.0					
			1A4a i Commercial/Institutional plants	11.8	10.6	9.1	9.0	7.2	6.6	6.6	6.1	5.4	5.8
			020103					0.2		0.1	0.1	0.1	0.0
			020105					0.0	0.0	0.0	0.0	0.0	0.0
			1A4b i Residential plants	46.5	50.6	42.9	50.0	43.7	43.3	45.3	39.6	37.8	35.7
			1A4c i Agriculture/Forestry/Fishing, Stationary	0.4	1.0	1.2	0.8	0.7	1.2	1.9	1.8	1.7	2.3
			020302								0.0		
			020304							0.0	0.0		
KEROSENE	1A2f i Industry-Other		031500	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
			032000	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	1A4a i Commercial/Institutional plants		020100	0.6	0.2	0.2	0.2	0.2	0.1	0.1	0.1	0.1	0.1

Sum of Fuel_rate_PJ					Year											
fuel_type	fuel_gr_abbr	nfr_id	nfr_name	snap_id	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999		
GAS	NATURAL GAS	1A4b i	Residential plants	020200	4.4	0.7	0.5	0.5	0.4	0.4	0.4	0.3	0.3	0.1		
		1A4c i	Agriculture/Forestry/Fishing, Stationary	020300	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0		
		ORIMULSION	1A1a	Electricity and heat production	010101						19.9	36.8	40.5	32.6	34.2	
		LPG	1A1a	Electricity and heat production	010100		0.0	0.0	0.0							
					010200	0.0	0.0	0.0								
					010203					0.0	0.0				0.0	
		1A1b	Petroleum refining	010306				0.0		0.0	0.0	0.0	0.0			
		1A2a	Iron and steel	030400	0.1	0.1	0.0	0.0	0.0	0.1	0.1	0.0	0.0	0.0	0.0	
		1A2b	Non-ferrous metals	030500	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
		1A2c	Chemicals	030600	0.0	0.1	0.0	0.1	0.1	0.1	0.1	0.1	0.1	0.0	0.0	
					030602							0.0	0.0	0.0	0.0	
		1A2d	Pulp, Paper and Print	031100	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
		1A2e	Food processing, beverages and tobacco	030900	0.3	0.2	0.2	0.2	0.1	0.1	0.1	0.1	0.1	0.1	0.1	
		1A2f i	Industry-Other	030700	0.2	0.2	0.2	0.2	0.3	0.4	0.6	0.4	0.4	0.4	0.4	
				030800	0.1	0.1	0.1	0.1	0.1	0.1	0.0	0.0	0.0	0.0	0.0	
				031000	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
				031200	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
				031300	0.2	0.3	0.2	0.2	0.3	0.3	0.3	0.3	0.3	0.3	0.2	
				031400	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
				031500	0.5	0.6	0.7	0.6	0.5	0.5	0.5	0.5	0.5	0.6	0.4	
				032000	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.1	0.1	0.1	0.1	
			1A4a i	Commercial/Institutional plants	020100	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
					020105											0.0
			1A4b i	Residential plants	020200	1.0	0.7	0.5	0.8	0.7	0.7	0.8	0.7	0.9	0.9	1.0
			1A4c i	Agriculture/Forestry/Fishing, Stationary	020300	0.3	0.2	0.2	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
			REFINERY GAS	1A1a	Electricity and heat production	010203						0.0	0.0			
				1A1b	Petroleum refining	010300	0.5	0.9	1.5	0.0						
					010304				2.1	2.4	2.3	2.7	2.3	2.5	2.5	2.7
					010306	13.5	13.5	13.2	13.2	14.0	18.5	18.7	14.5	12.7	12.7	13.1
				1A2f i	Industry-Other	032000	0.2	0.1	0.1	0.1		0.0	0.1	0.0		
				1A1a	Electricity and heat production	010101	4.0	4.4	3.3	4.4	6.4	7.8	9.5	8.4	17.5	17.3
					010102						2.0	2.8	4.1	8.1	9.3	6.5
					010103						0.0	0.1	0.1	0.1	0.1	0.1
					010104	2.5	3.9	5.7	7.5	7.6	8.2	15.1	17.7	12.6	12.6	23.6
					010105	0.7	1.3	2.2	4.2	8.6	16.9	22.2	23.6	26.5	26.5	26.8
					010200	10.0	11.8	11.1	10.9							
					010202						0.3	0.4	0.4	0.5	0.5	0.2
					010203						9.3	7.9	5.0	2.0	3.1	0.4
				1A1c	Manufacture of solid fuels and other energy industries	010504	9.5	9.7	11.1	11.2	12.3	13.0	15.3	20.0	22.1	24.1
				1A2a	Iron and steel	030400	1.7	1.5	1.5	1.5	1.6	0.1	0.1			
						030402						1.6	1.6	1.9	1.9	2.1
				1A2b	Non-ferrous metals	030500	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.2	0.1

Sum of		Year												
Fuel_rate_PJ														
fuel_type	fuel_gr_abbr	nfr_id	nfr_name	snap_id	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
	1A2c	Chemicals		030600	1.0	1.3	1.5	1.2	1.4	1.2	1.1	1.4	2.8	3.3
				030602	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.3	0.3	0.3
				030603					0.2	0.3	0.5	0.6	0.5	0.3
				030604	0.5	0.6	0.7	0.7	0.8	0.9	1.3	1.3	1.2	1.3
				030605								0.0	0.1	0.1
	1A2d	Pulp, Paper and Print		031100	2.3	1.8	1.6	1.2	1.3	1.3	1.5	1.4	1.6	1.8
				031102					0.7	1.1	0.9	1.1	1.1	1.0
				031103					0.0	0.1	0.1	0.1	0.1	0.1
				031104						0.1	0.9	1.0	1.0	1.0
	1A2e	Food processing, beverages and tobacco		030900	8.1	9.2	9.5	11.2	12.7	14.0	12.2	13.4	12.2	11.8
				030902						0.0	0.0	0.0	0.0	0.0
				030903					0.4	0.5	0.4	0.3	0.5	0.4
				030904					0.1	0.3	0.9	1.5	2.3	3.5
				030905					0.0	0.1	0.6	0.6	0.6	0.5
	1A2f i	Industry-Other		030106	0.1	0.0	0.0	0.1	0.1	0.0	0.0	0.0	0.0	0.1
				030700	4.2	4.2	4.1	4.5	5.0	5.9	5.2	5.6	5.7	6.4
				030705						0.0	0.0	0.0	0.0	0.0
				030800	0.2	0.3	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.7
				031000	1.2	1.4	1.4	1.3	1.3	1.2	1.2	1.2	1.0	1.3
				031005							0.0	0.0	0.0	0.0
				031200	0.2	0.2	0.3	0.4	0.5	0.7	0.7	0.7	0.7	0.6
				031300	1.4	2.0	2.2	2.5	2.8	2.9	3.5	3.2	3.2	3.8
				031305							0.0	0.0	0.1	0.1
				031400	0.0	0.0	0.0	0.1	0.1	0.1	0.0	0.0	0.0	0.2
				031405						0.0	0.1	0.1	0.1	0.1
				031500	0.1	0.1	0.1	0.2	0.2	0.2	0.2	0.2	0.2	0.2
				031503						0.0				
				032000	1.9	1.9	2.3	2.4	1.6	1.7	1.9	1.8	2.0	2.1
				032003					0.1		0.0	0.0	0.0	0.0
				032004							0.1	0.2	0.2	0.2
				032005	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.1	0.1	0.1
	1A4a i	Commercial/Institutional plants		020100	6.3	6.8	7.1	8.6	7.3	8.4	11.2	9.1	8.7	7.5
				020103					0.0	0.0	0.0	0.0	0.2	0.0
				020104						0.0		0.0		
				020105	0.0	0.1	0.3	0.4	0.6	0.7	0.8	1.0	1.0	1.1
	1A4b i	Residential plants		020200	17.4	20.4	20.9	24.1	24.7	26.9	30.4	28.4	29.1	29.0
				020202					0.0	0.0	0.1	0.0	0.0	0.0
				020204		0.0	0.5	0.8	1.0	1.0	1.4	1.5	1.5	1.5
	1A4c i	Agriculture/Forestry/Fishing, Stationary		020300	2.1	2.6	2.2	2.3	2.5	2.6	2.7	2.6	2.5	2.2
				020304	0.1	0.1	0.1	0.2	0.3	1.2	2.2	3.0	3.7	3.7
WASTE	WASTE	1A1a	Electricity and heat production	010100	1.0	3.6	5.6	8.4						
				010101									1.3	1.3
				010102					5.1	4.4	6.3	7.7	8.1	14.5

Sum of					Year											
Fuel_rate_PJ																
fuel_type	fuel_gr_abbr	nfr_id	nfr_name	snap_id	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999		
BIOMASS	WOOD			010103					4.1	5.3	6.0	5.6	4.7	1.1		
				010104					0.6	0.9	1.9	1.9	1.6	1.5		
				010200	13.6	12.1	11.1	9.8								
				010202						3.3	4.6	4.6	4.6			
				010203					9.3	7.8	4.8	5.7	5.6	9.2		
				1A2a	Iron and steel	030400					0.0	0.0				
				1A2c	Chemicals	030600	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
				1A2d	Pulp, Paper and Print	031100	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
				1A2e	Food processing, beverages and tobacco	030900	0.0	0.0	0.0	0.0	0.0	0.0	0.0		0.0	0.0
				1A2f i	Industry-Other	030700	0.0	0.0	0.0	0.0	0.0	0.0			0.0	
						031000	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
						031200	0.0	0.0			0.0	0.0				
						031300	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0		
						031400			0.0	0.0	0.0	0.0	0.0	0.0	0.0	
						032000	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
		1A4a i	Commercial/Institutional plants	020100	0.9	1.0	1.1	1.1	1.2	1.3	1.2	1.2	0.7	1.5		
				020103						0.0	0.0	0.0	0.0	0.0		
			1A1a	Electricity and heat production	010100			0.2	0.5							
					010101						0.0			0.3		
					010102						1.7	1.6	1.6	1.7	2.0	2.9
					010103						0.0	0.0	0.0	0.1	0.1	0.3
					010104							0.0				
					010200	3.2	3.6	4.1	3.8							
					010203						3.3	3.5	3.9	3.9	4.1	4.0
			1A2a	Iron and steel	030400	0.0	0.0	0.0	0.0	0.0						
			1A2b	Non-ferrous metals	030500	0.0										
			1A2c	Chemicals	030600	0.0										
			1A2d	Pulp, Paper and Print	031100	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.0	0.0	0.0
			1A2e	Food processing, beverages and tobacco	030900	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.2
					030902										0.0	0.0
			1A2f i	Industry-Other	030700	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
					031000						0.0	0.0				0.0
					031200	0.0	0.0	0.0			0.0	0.0	0.0	0.0	0.0	0.0
					031300	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.2
					031400	3.2	3.0	3.0	3.0	3.0	3.0	3.0	2.7	2.8	2.9	2.9
					031403						0.4	0.3	0.5	0.4	0.3	0.4
					032000	2.4	2.5	2.5	2.6	1.3	1.1	1.2	1.2	1.2	1.2	0.9
					032003						0.0	0.0	0.0	0.0	0.0	0.0
			1A4a i	Commercial/Institutional plants	020100	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.4	0.5	0.5	0.6
					020103						0.0	0.0	0.0	0.0		
			1A4b i	Residential plants	020200	9.0	10.4	10.7	11.9	11.6	11.8	12.7	12.6	11.1	11.6	
			1A4c i	Agriculture/Forestry/Fishing, Stationary	020300	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.2	0.2
					020304									0.0	0.0	0.0

Sum of Fuel_rate_PJ					Year									
fuel_type	fuel_gr_abbr	nfr_id	nfr_name	snap_id	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
STRAW	1A1a Electricity and heat production	010100		0.5	1.0	1.5	1.6							
			010101				0.1	0.1	0.4	0.7	1.0	1.3		
			010102				0.6	1.1	1.5	1.3	1.3	1.3		
			010103				0.7	1.0	1.4	1.5	1.5	1.3		
			010200	3.5	3.8	3.9	3.8							
		010203				3.9	4.0	4.2	3.9	3.9	3.9			
		1A2f i Industry-Other	031305									0.0	0.0	
			032003					0.0						
		1A4b i Residential plants	020200	5.1	5.1	5.1	4.8	4.4	4.1	3.6	3.9	3.8	3.4	
		1A4c i Agriculture/Forestry/Fishing, Stationary	020300	3.4	3.4	3.4	3.2	2.9	2.7	2.4	2.6	2.5	2.3	
020302						0.0	0.0	0.0	0.0	0.0	0.0			
BIO OIL	1A1a Electricity and heat production	010200	0.7	0.7	0.7	0.8								
		010203					0.2	0.3	0.1	0.0	0.0	0.0		
BIOGAS	1A1a Electricity and heat production	010100		0.1	0.2	0.0	0.0							
			010101				0.0	0.0	0.0	0.0				
		010102				0.0		0.1	0.0	0.1	0.0			
		010104				0.0	0.0	0.0						
		010105	0.1	0.2	0.3	0.5	0.5	0.6	0.6	0.8	1.0	1.0		
		010200	0.0	0.0	0.1	0.1								
		010203				0.2	0.2	0.2	0.2	0.2	0.2			
		1A2a Iron and steel	030400	0.0										
		1A2e Food processing, beverages and tobacco	030900	0.0			0.0	0.1	0.1	0.1	0.1	0.0		
			030903				0.0	0.0	0.0	0.0	0.0	0.0		
		1A2f i Industry-Other	032000	0.3	0.3	0.4	0.4							
		1A4a i Commercial/Institutional plants	020100				0.1	0.2	0.2	0.3	0.2	0.3		
			020103					0.0	0.0	0.0	0.1	0.1		
			020105	0.2	0.2	0.1	0.1	0.4	0.6	0.5	0.8	0.9	0.8	
		1A4c i Agriculture/Forestry/Fishing, Stationary	020300				0.0	0.0	0.1	0.0	0.0	0.0		
			020304				0.0	0.0	0.0	0.0	0.0	0.0		
BIO PROD GAS	1A1a Electricity and heat production	010105				0.1	0.0	0.0	0.0	0.0	0.0			
	1A4a i Commercial/Institutional plants	020105									0.0	0.0		
Grand Total					498.8	609.6	549.9	580.6	623.0	600.2	756.8	652.7	615.2	586.4

Sum of Fuel_rate_PJ		Year												
fuel_type	fuel_gr_abbr	nfr_id	nfr_name	snap_id	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
SOLID	ANODE CARBON	1A2f i	Industry-Other	032000										0.0
		1A1a	Electricity and heat production	010101	143.8	156.2	158.3	223.5	167.9	140.0	218.4	180.9	159.4	161.9
	COAL			010102	9.3	7.7	8.0	6.4	4.5	4.0	3.3	3.1	2.8	2.0
				010104										0.0
				010203	0.0	0.0	0.0	0.0	0.0	0.0		0.1	0.1	
	1A2a	Iron and steel		030400									0.0	0.0
	1A2b	Non-ferrous metals		030500									0.0	0.0
	1A2c	Chemicals		030600	0.5	0.5	0.4	0.6	0.6	0.5	0.2			
	1A2e	Food processing, beverages and tobacco		030900	1.5	1.8	1.4	0.4	0.7	0.4	0.6	0.6	1.1	0.1
				030902	1.1	1.0	1.0	1.6	1.5	1.5	1.2	1.2	1.2	1.2
				030903	0.4	0.4	0.1	0.1	0.1	0.1	0.2	0.2	0.3	0.3
	1A2f i	Industry-Other		030700	0.3	0.3		1.6	1.8	1.6	1.8	1.9	0.0	0.4
				030800	0.8	0.6	0.3	0.1	0.1	0.1	0.1	0.1	0.2	0.1
				031300	0.0	0.0	0.0	0.0	0.0				0.0	0.0
				031400	0.1			0.0	0.0					
				031600	5.7	4.5	4.3	3.4	3.8	3.9	4.3	4.0	3.5	1.1
				032000										0.1
	1A4a i	Commercial/Institutional plants		020100					0.0					
	1A4b i	Residential plants		020200	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	1A4c i	Agriculture/Forestry/Fishing, Stationary		020300	1.1	1.2	0.9	1.2	1.4	1.8	2.0	2.1	1.8	0.5
				020304					0.0	0.0				0.0
	BROWN COAL BRI.	1A4b i	Residential plants	020200	0.0	0.0	0.0	0.0					0.0	0.0
	COKE OVEN COKE	1A2a	Iron and steel	030400	0.0	0.0	0.0							
				030900	0.2	0.2	0.1	0.2	0.2	0.1	0.1	0.1	0.1	
				030902									0.1	0.1
				030903								0.0	0.0	0.1
		1A2f i	Industry-Other	030700	0.0	0.0	0.1	0.1	0.1	0.1	0.1	0.1	0.0	0.0
				030800	0.0									
				031200									0.0	0.0
				031300				0.0	0.0			0.0	0.0	0.0
				032000	0.9	0.9	0.8	0.7	0.8	0.7	0.8	1.0	0.9	0.6
		1A4b i	Residential plants	020200	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	LIQUID	PETROLEUM COKE	1A1a	Electricity and heat production	010102				0.0	0.0				0.0
			1A2e	Food processing, beverages and tobacco	030900								0.0	
			1A2f i	Industry-Other	030700	0.2	0.1	0.1	0.1	0.1				
					030800	0.0	0.1	0.1	0.1	0.1				
					031300								0.0	
					031600	6.5	7.7	7.5	7.7	8.2	7.8	8.5	9.1	6.8
			1A4a i	Commercial/Institutional plants	020100	0.0	0.0	0.0	0.0		0.1	0.0	0.0	
			1A4b i	Residential plants	020200	0.0	0.0	0.0	0.0		0.1		0.0	0.1
			1A4c i	Agriculture/Forestry/Fishing, Stationary	020300	0.0	0.0	0.0	0.0					
	RESIDUAL OIL	1A1a	Electricity and heat production	010101	3.4	3.5	3.7	5.8	4.6	4.3	3.3	5.4	2.8	3.6
				010102	0.7	2.3	1.2	1.7	1.3	1.5	1.8	0.3	0.9	1.9

Sum of Fuel_rate_PJ				Year														
fuel_type	fuel_gr_abbr	nfr_id	nfr_name	snap_id	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009				
				010103	0.3	0.1	0.1	0.1	0.2	0.2	0.1	0.6	0.2	0.1				
				010104		1.7	6.6	9.3	7.4	6.3	8.4	4.5	4.5	2.9				
				010105	0.0	0.0	0.0	0.0	0.0	0.0	0.0							
				010202							0.1							
				010203	1.1	1.0	1.0	0.6	0.3	0.5	0.3			0.1	0.1			
				1A1b Petroleum refining	010306	1.3	1.4	1.4	0.9	1.1	0.7	0.6	0.8	0.9	0.7			
				1A2a Iron and steel	030400	0.0	0.0	0.0					0.0	0.0	0.0			
					030403						0.0	0.0						
				1A2b Non-ferrous metals	030500	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0			
				1A2c Chemicals	030600	0.9	1.0	1.1	0.8	0.9	0.4	0.6	0.5	0.3	0.2			
				1A2d Pulp, Paper and Print	031100	0.1	0.2	0.2	0.1	0.1	0.1	0.1	0.1	0.1	0.0			
				1A2e Food processing, beverages and tobacco	030900	5.4	5.1	5.6	4.9	4.5	3.4	5.2	3.3	1.9	1.2			
					030902	0.6	0.6	0.5	0.9	0.9	1.1	0.8	0.6	1.9	2.0			
					030903	0.2	0.3	0.3	0.7	0.8	0.8	0.8	0.8	1.0	1.1			
				1A2f i Industry-Other	030700	0.6	0.6	0.7	0.3	0.1		0.2	0.0					
					030800	0.5	0.4	0.4	0.3	0.3	0.2	0.3	0.2	0.1	0.1			
					031000	0.0	0.0	0.0					0.0	0.0	0.0			
					031200	0.0	0.0	0.0			0.0	0.0	0.0	0.0	0.0			
					031300	0.1	0.1	0.2	0.1	0.1	0.4	0.6	0.2	0.2	0.1			
					031305		0.0	0.0										
					031400	0.4	0.3	0.3	0.2	0.2	0.4	0.5	0.2	0.2	0.1			
					031500	0.0	0.0	0.0										
					031600	0.9	0.5	0.6	0.6	0.8	0.7	1.0	1.1	0.5	0.2			
					032000	0.1	0.1	0.1	0.0	0.0	0.0	0.0	0.2	0.0	0.0			
					032003		0.0											
						032005				0.0	0.0	0.0	0.0		0.0			
					1A4a i Commercial/Institutional plants	020100	0.3	0.2	0.5	0.2	0.1	0.1	0.3	0.2	0.1	0.0		
					1A4b i Residential plants	020200	0.0	0.0	0.1	0.0	0.0	0.0	0.2	0.0	0.0	0.0		
					1A4c i Agriculture/Forestry/Fishing, Stationary	020300	1.8	1.6	1.4	0.9	0.7	0.8	0.9	0.6		0.1		
						020302					0.0	0.0	0.0	0.0	0.0	0.0		
						020304	0.0	0.0	0.0	0.0								
				GAS OIL	1A1a	Electricity and heat production	010101	0.1	0.1	0.1	1.0	0.2	0.2	0.5	0.5	0.9	2.3	
							010102	0.1	0.1	0.1	0.0	0.1	0.0	0.0	0.1	0.0	0.1	
							010103	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	
							010104	0.1	0.0	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	
							010105	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	
							010202	0.5	0.9	0.2	0.4	0.5	0.2	0.2	0.2	0.3	0.4	
							010203	0.6	0.5	0.4	1.0	0.6	0.5	0.4	0.4	0.8	1.0	
							1A1b Petroleum refining	010306				0.0	0.0	0.0	0.0	0.0	0.0	0.0
							1A2a Iron and steel	030400	0.0	0.0	0.0	0.0	0.0	0.0	0.0			
	030403									0.0								

Sum of Fuel_rate_PJ			Year										
fuel_type	fuel_gr_abbr	nfr_id nfr_name	snap_id	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
KEROSENE	1A2b Non-ferrous metals	1A2c Chemicals	030500	0.0	0.0	0.0	0.0	0.0	0.0	0.0			
			030600	0.1	0.2	0.1	0.1	0.1	0.1	0.0			
			030602							0.0	0.0	0.0	0.0
	1A2d Pulp, Paper and Print		030604	0.0					0.0	0.0	0.0	0.0	0.0
			031100	0.1	0.1	0.1	0.1	0.1	0.0	0.0			
			031102							0.0	0.0		
	1A2e Food processing, beverages and tobacco		031103			0.0	0.0	0.0	0.0	0.0	0.0		
			030900	0.5	0.7	0.6	0.5	0.5	0.4	0.1			
			030902				0.0	0.0	0.0	0.0	0.0	0.0	0.0
			030903	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	1A2f i Industry-Other		030904			0.0							
			030700	0.2	0.3	0.2	0.3	0.2	0.2	0.1			
			030800	0.3	0.5	0.4	0.5	0.4	0.3	0.1			
			031000	0.0	0.1	0.0	0.1	0.1	0.0	0.0			
			031200	0.1	0.1	0.1	0.1	0.1	0.1	0.0			
			031205									0.0	0.0
			031300	0.6	0.7	0.5	0.6	0.6	0.3	0.1			
			031305	0.0	0.0								
			031400	0.1	0.1	0.1	0.1	0.1	0.0	0.0			
			031403	0.1	0.0	0.0							
			031600							0.0	0.0	0.0	0.0
			032000	0.2	0.3	0.2	0.3	0.3	0.1	0.0	0.0	0.0	0.0
			032003			0.0							
	1A4a i Commercial/Institutional plants		020100	5.0	4.7	4.0	4.3	4.4	3.8	3.0	2.6	2.8	2.8
			020103	0.1	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1
			020105	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	1A4b i Residential plants		020200	30.3	31.5	29.0	27.0	25.3	23.9	21.2	17.2	15.2	16.4
			020204									0.0	0.0
	1A4c i Agriculture/Forestry/Fishing, Stationary		020300	2.2	2.6	2.2	2.3	2.0	1.2	0.5			
			020302								0.0	0.0	0.0
			020304	0.0	0.0	0.0	0.0		0.0	0.0			0.0
	1A2f i Industry-Other		031500	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
			032000	0.0	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0
			1A4a i Commercial/Institutional plants	020100	0.1	0.1	0.1	0.1	0.1	0.1	0.0	0.0	0.0
			1A4b i Residential plants	020200	0.1	0.2	0.1	0.2	0.1	0.2	0.1	0.1	0.1
	1A4c i Agriculture/Forestry/Fishing, Stationary		020300	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
ORIMULSION	1A1a Electricity and heat production		010101	34.1	30.2	23.8	1.9	0.0					
LPG	1A1a Electricity and heat production		010101							0.0		0.0	0.0
			010102									0.0	0.0
			010202						0.0	0.0	0.0		
			010203	0.0					0.0		0.0	0.0	0.0
			1A2a Iron and steel	030400	0.1	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0

Sum of Fuel_rate_PJ		Year											
fuel_type	fuel_gr_abbr	nfr_id nfr_name	snap_id	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
GAS	NATURAL GAS	1A2b Non-ferrous metals	030500	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
		1A2c Chemicals	030600	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
			030602	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
		1A2d Pulp, Paper and Print	031100	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
		1A2e Food processing, beverages and tobacco	030900	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.0	0.1
		1A2f i Industry-Other	030700	0.3	0.1	0.1	0.1	0.1	0.2	0.2	0.1	0.0	0.0
			030800	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
			031000	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
			031200	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
			031300	0.2	0.1	0.1	0.1	0.1	0.1	0.2	0.1	0.1	0.1
			031400	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
			031500	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.1
			032000	0.0	0.0	0.0	0.0	0.0	0.1	0.1	0.0	0.0	0.0
		1A4a i Commercial/Institutional plants	020100	0.1	0.1	0.1	0.2	0.2	0.3	0.3	0.3	0.3	0.3
			020105						0.0	0.0	0.0	0.0	0.0
		1A4b i Residential plants	020200	1.2	1.2	1.1	1.1	1.1	1.1	1.1	1.1	1.0	0.8
		1A4c i Agriculture/Forestry/Fishing, Stationary	020300	0.1	0.1	0.1	0.1	0.1	0.0	0.0	0.0	0.0	0.0
	REFINERY GAS	1A1b Petroleum refining	010304	2.4	2.4	2.5	2.7	2.4	2.0	2.2	2.3	1.8	1.9
			010306	13.2	13.3	12.7	13.9	13.4	13.4	13.9	13.6	12.9	13.5
		1A1a Electricity and heat production	010101	18.4	18.2	16.5	17.9	17.3	17.2	19.0	13.9	10.9	13.4
			010102	6.5	6.4	5.5	3.9	3.3	3.0	2.6	0.9	3.8	2.7
			010103	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.1	0.0
			010104	22.8	24.9	30.0	29.7	32.0	27.3	33.5	26.2	27.8	24.6
			010105	25.7	28.0	27.8	26.9	27.1	24.2	21.6	17.3	18.4	15.4
			010202	0.1	0.1	0.2	0.2	0.3			0.2	0.3	0.5
			010203	2.3	2.9	2.3	3.1	1.2	2.8	3.1	6.1	6.0	8.1
		1A1c Manufacture of solid fuels and other energy industries	010504	25.4	24.9	26.6	26.6	27.5	28.2	28.8	28.6	28.2	26.7
		1A2a Iron and steel	030400	0.1	0.0	1.7	0.3	0.1	0.0			0.1	0.0
			030402	1.6	1.8		1.2	1.2	1.2	1.3	1.4	1.4	0.7
		1A2b Non-ferrous metals	030500	0.2	0.2	0.2	0.1	0.1	0.1	0.1	0.1	0.1	0.1
		1A2c Chemicals	030600	3.0	3.2	2.7	3.1	2.3	2.7	2.5	1.9	2.3	2.7
			030602	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.6	0.5
			030603	0.5									
			030604	1.4	1.7	1.8	1.7	1.6	1.3	1.2	1.1	1.1	1.1
			030605	0.1	0.1	0.1	0.1	0.1	0.0				
		1A2d Pulp, Paper and Print	031100	1.3	1.8	1.2	1.9	1.5	2.0	2.5	1.9	1.5	1.4
			031102	1.1	1.1	1.2	1.0	1.0	1.0	1.0	0.2	0.1	0.1
			031103	0.1	0.1	0.0	0.0	0.0	0.0	0.0	0.0		
			031104	1.0	0.7	1.0	0.9	0.9	1.0	0.9	0.9	0.9	0.7
		1A2e Food processing, beverages and tobacco	030900	11.1	11.7	10.2	9.0	9.9	11.6	11.3	12.3	13.0	12.1
			030902	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
			030903	0.1	0.1	0.4	0.6	0.6	0.6	0.6	0.6	0.6	0.6
			030904	3.8	3.5	3.8	3.8	4.2	3.6	2.6	2.4	1.4	1.2

Sum of Fuel_rate_PJ				Year												
fuel_type	fuel_gr_abbr	nfr_id	nfr_name	snap_id	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009		
		1A2f i	Industry-Other	030905	0.9	1.1	1.0	1.0	1.1	1.0	0.6	0.1	0.3	0.3		
				030106	0.1	0.1	0.0	0.0	0.0	0.0	0.0	0.1	0.1			
				030700	5.8	5.8	5.4	5.5	5.0	3.9	4.1	4.8	4.0	3.7		
				030705	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0		
				030800	0.6	0.9	0.6	0.7	0.8	1.4	1.3	1.1	0.6	0.5		
				031000	1.2	1.2	1.2	1.0	0.9	0.5	0.5	0.3	0.4	0.4		
				031005	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0		
				031200	0.5	0.5	0.4	0.5	0.5	0.4	0.5	0.6	0.6	0.6		
				031300	3.6	4.1	3.7	4.1	3.8	2.8	2.9	3.0	3.9	3.4		
				031305	0.2	0.1	0.1	0.2	0.1	0.1	0.1	0.1	0.1	0.1		
				031400	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.2	0.2		
				031405	0.1	0.0	0.1	0.1	0.0	0.0	0.0	0.0	0.0	0.0		
				031500	0.2	0.2	0.2	0.2	0.2	0.3	0.4	0.4	0.3	0.4		
				031503			0.0									
				032000	2.1	2.3	2.2	2.3	2.1	1.5	1.4	1.5	1.2	1.0		
				032003	0.0	0.0	0.0	0.0								
				032004	0.2	0.2	0.2	0.1	0.1	0.1		0.0	0.0	0.0		
				032005	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.0	0.0		
		1A4a i	Commercial/Institutional plants	020100	7.2	7.3	7.6	9.2	9.2	9.7	10.8	10.1	10.0	10.1		
				020103	0.2	0.2	0.2	0.0	0.1	0.0	0.1	0.0	0.1	0.0		
				020104	0.0	0.0	0.0	0.0	0.0	0.0	0.0					
				020105	1.1	1.1	1.2	1.1	1.1	1.0	0.9	0.8	0.8			
		1A4b i	Residential plants	020200	27.6	29.3	28.1	30.0	29.9	29.5	28.6	26.6	26.5	26.7		
				020202	0.1	0.1	0.0	0.1	0.1	0.0	0.1	0.1	0.1	0.1		
				020204	1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.3	1.2	1.1		
		1A4c i	Agriculture/Forestry/Fishing, Stationary	020300	2.4	2.7	2.5	2.3	2.3	2.2	2.2	1.9	1.7	1.7		
				020304	3.3	3.1	3.4	3.2	3.3	2.9	2.0	1.4	1.1	0.9		
				010101		0.2	0.9	0.1					0.0	0.0		
		WASTE	WASTE	1A1a	Electricity and heat production	010102	12.1	13.0	13.9	14.1	16.6	19.3	20.1	20.3	21.1	21.1
						010103	9.0	9.0	9.2	9.1	9.1	8.8	9.6	9.4	9.6	8.8
						010104	2.2	2.5	2.6	3.0	2.9	2.6	3.1	3.3	3.3	3.2
						010203	6.4	6.6	6.6	7.4	6.5	4.3	3.8	4.7	5.0	4.0
						030600	0.0					0.0		0.0		
				1A2d	Pulp, Paper and Print	031100	0.0					0.0		0.0	0.0	0.0
				1A2e	Food processing, beverages and tobacco	030900	0.0					0.0		0.1	0.1	0.1
						030902					0.0		0.0	0.0		
1A2f i	Industry-Other			030800						0.1		0.0	0.1	0.1		
				031000	0.0							0.0				
				031300	0.0					0.0		0.0	0.0	0.0		
				031400	0.0					0.0		0.0				
				032000	0.0					0.1		0.1				
1A4a i	Commercial/Institutional plants			020100				1.3	0.1	0.4	0.2		0.1			
				020103	0.0	0.0	0.0	0.1	0.1	0.2	0.2	0.0	0.1	0.1		

Sum of		Year												
Fuel_rate_PJ														
fuel_type	fuel_gr_abbr	nfr_id	nfr_name	snap_id	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
	INDUSTR. WASTES	1A2f i	Industry-Other	031600	0.5	1.4	1.9	1.5	2.0	2.0	1.5	1.6	2.0	1.7
BIOMASS	WOOD	1A1a	Electricity and heat production	010101		0.0	0.1	0.3	0.2	0.2	0.3	0.2	0.3	0.5
				010102	2.7	2.5	3.2	5.3	5.4	6.5	6.5	6.3	5.8	7.1
				010103	0.4	0.5	0.6	0.7	0.7	0.6	0.5	0.6	0.6	0.7
				010104			0.1	1.6	4.5	4.5	2.6	3.8	6.0	6.3
				010203	3.9	4.5	5.0	5.6	6.2	6.6	7.0	7.1	7.9	8.6
		1A2a	Iron and steel	030400				0.0	0.0					
		1A2d	Pulp, Paper and Print	031100	0.0	0.0	0.0	0.0	0.0	0.0			0.7	1.4
				031102							0.0	1.1	1.2	1.2
		1A2e	Food processing, beverages and tobacco	030900	0.1	0.1	0.1	0.0	0.1	0.3	0.3	0.3	0.2	0.0
				030902	0.0	0.0								0.0
				030903					0.0	0.1	0.1	0.1	0.1	0.0
		1A2f i	Industry-Other	030700	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
				030800								0.1	0.9	0.8
				031000	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
				031200	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
				031300	0.1	0.1	0.0	0.1	0.1	0.1	0.2	0.3	0.2	0.2
				031305								0.0	0.0	0.0
				031400	3.0	3.1	2.5	1.8	1.7	2.3	2.6	2.7	2.9	2.4
				031403	0.4	0.4	0.4	0.3	0.4	0.3	0.4	0.4	0.4	0.5
				032000	1.2	1.3	0.7	1.6	1.6	0.9	1.1	0.8	0.8	0.8
				032003	0.0	0.0	0.0							
		1A4a i	Commercial/Institutional plants	020100	0.8	0.7	0.7	0.7	0.7	0.8	1.0	1.0	1.1	1.0
		1A4b i	Residential plants	020200	14.6	17.5	18.1	20.9	22.3	26.4	29.4	35.5	34.5	34.3
				020202						0.0	0.0	0.0	0.0	0.0
				020204									0.0	0.0
		1A4c i	Agriculture/Forestry/Fishing, Stationary	020300	0.2	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.2
				020304	0.0	0.0								
STRAW		1A1a	Electricity and heat production	010101	1.1	1.6	2.6	3.2	3.7	3.3	3.7	3.6	2.4	2.8
				010102	1.3	1.3	1.2	1.3	2.1	2.0	1.7	1.9	1.7	1.9
				010103	0.7	2.1	1.9	2.1	2.1	2.1	2.1	2.1	2.1	2.2
				010104		0.1	1.2	1.7	1.9	2.4	2.5	2.5	0.8	1.5
				010203	3.8	3.8	3.8	3.8	3.4	3.7	3.7	3.8	3.9	4.1
		1A2f i	Industry-Other	031305	0.0	0.0								
		1A4b i	Residential plants	020200	3.1	2.9	2.9	2.9	2.9	2.9	2.9	2.9	2.9	2.9
		1A4c i	Agriculture/Forestry/Fishing, Stationary	020300	2.1	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9
				020302	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
BIO OIL		1A1a	Electricity and heat production	010101				0.1				0.0	0.0	
				010105					0.0			0.0	0.0	0.0
				010202				0.0	0.0	0.0	0.0	0.0	0.4	0.2
				010203	0.0	0.2	0.1	0.3	0.6	0.7	1.1	1.1	1.4	1.4
		1A2e	Food processing, beverages and tobacco	030903			0.1							
		1A2f i	Industry-Other	031305			0.0	0.0	0.0	0.0	0.0			

Sum of Fuel_rate_PJ				Year										
fuel_type	fuel_gr_abbr	nfr_id	nfr_name	snap_id	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
				032005					0.0	0.0	0.0	0.0	0.0	0.0
		1A4a i	Commercial/Institutional plants	020105								0.0		
		1A4b i	Residential plants	020200								0.0	0.0	0.0
		1A4c i	Agriculture/Forestry/Fishing, Stationary	020304	0.0	0.0	0.0							
	BIOGAS	1A1a	Electricity and heat production	010102	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
				010105	1.1	1.1	1.3	1.3	1.4	1.6	1.6	1.7	1.6	1.9
				010203	0.3	0.3	0.2	0.3	0.1	0.1	0.1	0.1	0.2	0.1
		1A2e	Food processing, beverages and tobacco	030900	0.0	0.0	0.0	0.0	0.0	0.1	0.1	0.1	0.0	0.1
				030902	0.0	0.0	0.0	0.0	0.1	0.0	0.0	0.0	0.1	0.1
				030903	0.0	0.1	0.1	0.1	0.1	0.0	0.0	0.0	0.0	0.1
		1A2f i	Industry-Other	031300									0.0	0.0
				032005		0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
		1A4a i	Commercial/Institutional plants	020100	0.3	0.4	0.4	0.3	0.4	0.4	0.5	0.4	0.4	0.3
				020103	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
				020105	0.9	0.8	0.8	0.8	0.8	0.8	0.6	0.6	0.5	0.6
		1A4c i	Agriculture/Forestry/Fishing, Stationary	020300	0.1	0.1	0.1	0.1	0.2	0.1	0.3	0.3	0.4	0.2
				020304	0.1	0.1	0.2	0.4	0.5	0.6	0.5	0.6	0.5	0.6
	BIO PROD GAS	1A1a	Electricity and heat production	010105	0.0	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.3
		1A4a i	Commercial/Institutional plants	020105		0.0	0.0		0.0	0.0	0.0	0.0	0.0	0.0
Grand Total					544.3	570.7	569.1	628.6	569.2	532.5	618.0	558.4	530.1	523.2

Sum of Fuel_rate_PJ					Year		
fuel_type	fuel_gr_abbr	nfr_id	nfr_name	snap_id	2010	2011	2012
SOLID	ANODE CARBON	1A2f i	Industry-Other	032000	0.0	0.0	0.0
		1A1a	Electricity and heat production	010101	155.9	128.1	100.7
	COAL		Electricity and heat production	010102	1.7	1.1	1.0
				010104	0.0	0.7	
				010203	0.1	0.1	0.1
		1A2a	Iron and steel	030400	0.0	0.0	0.0
		1A2b	Non-ferrous metals	030500	0.0	0.0	0.0
		1A2e	Food processing, beverages and tobacco	030900	0.5	0.5	0.0
				030902	1.0	1.2	1.2
				030903	0.2	0.2	0.3
		1A2f i	Industry-Other	030700		0.6	0.3
				030703		0.2	0.3
				030800	0.0	0.1	0.1
				031300	0.0	0.0	0.0
				031600	2.0	1.4	0.9
				032000	0.1	0.1	0.1
		1A4b i	Residential plants	020200	0.0	0.0	0.0
		1A4c i	Agriculture/Forestry/Fishing, Stationary	020300	1.3	1.2	1.2
	SUB-BITUMINOUS	1A1a	Electricity and heat production	010104		0.0	0.1
	BROWN COAL BRI.	1A4b i	Residential plants	020200	0.0	0.0	0.0
	COKE OVEN COKE	1A2e	Food processing, beverages and tobacco	030900	0.0	0.0	0.0
				030902	0.0	0.0	0.0
				030903	0.1	0.1	0.1
		1A2f i	Industry-Other	030700	0.0	0.0	0.1
				031200	0.0	0.0	0.0
				031300	0.0	0.0	0.0
				032000	0.6	0.5	0.4
		1A4b i	Residential plants	020200	0.0	0.0	0.0
LIQUID	PETROLEUM COKE	1A1a	Electricity and heat production	010102	0.0	0.0	
		1A2e	Food processing, beverages and tobacco	030900	0.0	0.0	0.0
		1A2f i	Industry-Other	030700	0.0	0.1	0.1
				031300	0.0	0.0	0.0
				031600	5.1	6.4	6.6
		1A4a i	Commercial/Institutional plants	020100	0.0	0.0	
		1A4b i	Residential plants	020200	0.0	0.0	
	RESIDUAL OIL	1A1a	Electricity and heat production	010101	4.9	1.8	1.5
				010102	0.2	0.2	0.1
				010103	0.1	0.1	0.1
				010104	0.2	0.1	0.0
				010105	0.0	0.0	
				010202	0.0	0.0	
				010203	0.4	0.1	0.1
				010306	0.5	0.5	0.8

Sum of Fuel_rate_PJ				Year						
fuel_type	fuel_gr_abbr	nfr_id	nfr_name	snap_id	2010	2011	2012			
		1A2a	Iron and steel	030400	0.0	0.0	0.0			
		1A2b	Non-ferrous metals	030500	0.0	0.0	0.0			
		1A2c	Chemicals	030600	0.3	0.2				
				030603			0.2			
		1A2d	Pulp, Paper and Print	031100	0.0	0.0	0.0			
		1A2e	Food processing, beverages and tobacco	030900	2.0	0.9	0.2			
				030902	1.9	1.9	1.9			
				030903	1.0	1.0	1.2			
		1A2f i	Industry-Other	030800	0.1	0.1	0.1			
				031000	0.0	0.0	0.0			
				031200	0.0	0.0	0.0			
				031300	0.1	0.1	0.1			
				031400	0.2	0.1	0.1			
				031600	0.3	0.3	0.2			
				032000	0.0	0.0	0.0			
				032005		0.0	0.0			
		1A4a i	Commercial/Institutional plants	020100	0.0	0.0	0.2			
		1A4b i	Residential plants	020200	0.0	0.0	0.1			
		1A4c i	Agriculture/Forestry/Fishing, Stationary	020300	0.6	0.4	0.2			
				020302	0.0	0.0	0.0			
				020304	0.0	0.0				
		GAS OIL		1A1a	Electricity and heat production	010101	1.3	1.3	1.2	
						010102	0.2	0.1	0.1	
						010103	0.0	0.1	0.0	
						010104	0.1	0.0	0.1	
						010105	0.1	0.1	0.1	
						010202	1.2	0.3	0.2	
						010203	1.4	0.9	0.9	
				1A1b	Petroleum refining	010306	0.0	0.0	0.0	
				1A1c	Manufacture of solid fuels and other energy industries	010504		0.0	0.0	
				1A2a	Iron and steel	030402		0.0	0.0	
				1A2c	Chemicals	030602	0.0	0.0	0.0	
				030604	0.0	0.0	0.0			
1A2d	Pulp, Paper and Print			031102		0.0	0.0			
1A2e	Food processing, beverages and tobacco			030902	0.0	0.0	0.0			
				030903	0.0					
1A2f i	Industry-Other			030703		0.0	0.0			
				031205	0.0	0.0	0.0			
				031600	0.0	0.0	0.0			
				032000	0.0	0.0	0.0			
1A4a i	Commercial/Institutional plants			020100	2.7	2.1	2.4			
				020103	0.1	0.1	0.0			
				020105	0.0	0.0	0.0			

Sum of Fuel_rate_PJ				Year		
fuel_type	fuel_gr_abbr	nfr_id nfr_name	snap_id	2010	2011	2012
KEROSENE		1A4b i Residential plants	020200	14.5	9.6	5.0
			020204	0.0		
		1A4c i Agriculture/Forestry/Fishing, Stationary	020302	0.0	0.0	0.0
		1A2f i Industry-Other	031500	0.0	0.0	0.0
			032000	0.0	0.0	0.0
		1A4a i Commercial/Institutional plants	020100	0.0	0.0	0.0
		1A4b i Residential plants	020200	0.0	0.0	0.0
		1A4c i Agriculture/Forestry/Fishing, Stationary	020300	0.0	0.0	0.0
		1A1a Electricity and heat production	010101	0.0	0.0	0.0
			010102	0.1	0.0	0.0
LPG			010103	0.0	0.0	0.0
			010104			0.0
			010203	0.0	0.0	0.0
		1A1b Petroleum refining	010306		0.4	0.2
		1A2a Iron and steel	030400	0.0		0.0
			030402		0.0	0.0
		1A2b Non-ferrous metals	030500	0.0	0.0	0.0
		1A2c Chemicals	030600	0.0	0.0	0.0
			030602	0.0	0.0	0.0
		1A2d Pulp, Paper and Print	031100	0.0	0.0	0.0
		1A2e Food processing, beverages and tobacco	030900	0.0	0.0	0.0
		1A2f i Industry-Other	030700	0.0	0.0	0.0
			030800	0.0	0.0	0.0
			031000	0.0	0.0	0.0
			031200	0.0	0.0	0.0
			031300	0.1	0.1	0.1
			031400	0.0	0.0	0.0
			031500	0.1	0.1	0.1
			032000	0.0	0.0	0.0
		1A4a i Commercial/Institutional plants	020100	0.3		0.1
			020105	0.0	0.0	0.0
		1A4b i Residential plants	020200	0.8	0.7	0.7
		1A4c i Agriculture/Forestry/Fishing, Stationary	020300	0.0	0.0	0.0
REFINERY GAS		1A1b Petroleum refining	010304	1.5	1.5	1.7
			010306	12.7	13.4	13.9
GAS	NATURAL GAS	1A1a Electricity and heat production	010101	14.4	9.1	8.8
			010102	4.6	3.1	3.4
			010103	0.0	0.0	0.0
			010104	28.6	21.8	15.5
			010105	19.7	16.2	10.1
			010202	1.2	0.9	1.0
			010203	8.9	7.7	11.4
		1A1c Manufacture of solid fuels and other energy industries	010503		0.2	0.2

Sum of Fuel_rate_PJ				Year			
fuel_type	fuel_gr_abbrev	nfr_id	nfr_name	snap_id	2010	2011	2012
				010504	26.0	24.5	24.8
		1A2a	Iron and steel	030402	1.2	1.3	1.0
		1A2b	Non-ferrous metals	030500	0.1	0.1	0.1
		1A2c	Chemicals	030600	2.9	3.0	2.7
				030602	0.6	0.5	0.6
				030603			0.0
				030604	1.0	0.8	0.8
		1A2d	Pulp, Paper and Print	031100	1.5	1.5	1.4
				031102	0.1	0.1	0.0
				031104	0.7	0.6	0.5
		1A2e	Food processing, beverages and tobacco	030900	12.5	12.3	11.8
				030902	0.1	0.0	0.0
				030903	0.6	0.6	0.4
				030904	2.4	1.2	1.3
				030905	0.4	0.3	0.2
		1A2f i	Industry-Other	030106	0.1	0.0	0.0
				030700	4.0	4.0	3.7
				030703		0.0	0.0
				030705	0.0	0.0	0.0
				030800	0.6	0.5	0.5
				031000	0.4	0.4	0.4
				031005	0.0	0.0	0.0
				031200	0.6	0.6	0.6
				031300	3.7	3.6	3.4
				031305	0.1	0.1	0.0
				031400	0.2	0.2	0.2
				031405	0.1	0.0	0.0
				031500	0.4	0.6	0.6
				032000	1.2	1.2	1.2
				032004	0.0	0.0	0.0
				032005	0.0	0.0	0.0
		1A4a i	Commercial/Institutional plants	020100	10.0	9.4	7.4
				020103	0.1	0.1	0.0
				020105	0.8	0.6	0.4
		1A4b i	Residential plants	020200	31.5	27.1	27.7
				020202	0.1	0.1	0.2
				020204	1.1	0.9	0.7
		1A4c i	Agriculture/Forestry/Fishing, Stationary	020300	1.8	1.7	1.5
				020304	1.0	0.8	0.5
WASTE	WASTE	1A1a	Electricity and heat production	010102	20.0	20.2	19.6
				010103	8.5	8.6	8.3
				010104	3.3	3.2	3.2
				010203	4.2	4.3	4.4

Sum of Fuel_rate_PJ				Year				
fuel_type	fuel_gr_abbr	nfr_id	nfr_name	snap_id	2010	2011	2012	
BIOMASS	WOOD	1A2d	Pulp, Paper and Print	031100	0.0	0.0	0.0	
		1A2e	Food processing, beverages and tobacco	030900	0.1	0.1	0.1	
		1A2f i	Industry-Other	030800	0.1	0.1	0.1	
				031300	0.0	0.0	0.0	
		1A4a i	Commercial/Institutional plants	020100	0.1	0.1	0.2	
				020103	0.1	0.1	0.0	
		INDUSTR. WASTES	1A2f i	Industry-Other	031600	1.4	1.7	1.5
			1A1a	Electricity and heat production	010101	3.3	4.7	5.6
				010102	9.3	8.8	10.3	
				010103	1.3	1.3	1.3	
				010104	11.3	11.8	13.9	
				010203	10.3	10.4	11.8	
			1A2d	Pulp, Paper and Print	031100	1.5	1.9	2.0
				031102	1.2	1.3	1.3	
			1A2e	Food processing, beverages and tobacco	030900	0.0		
				030902	0.0	0.1	0.1	
				030903	0.0	0.0	0.0	
			1A2f i	Industry-Other	030700	0.0	0.0	0.0
				030800	0.8	0.9	0.9	
				031000	0.0	0.0	0.0	
			031200	0.0	0.0	0.0		
			031300	0.2	0.2	0.2		
			031305	0.0				
			031400	2.5	2.6	2.6		
			031403	0.4	0.4	0.3		
			032000	0.8	0.8	0.8		
		1A4a i	Commercial/Institutional plants	020100	1.0	1.0	0.9	
		1A4b i	Residential plants	020200	36.9	33.0	32.6	
				020202	0.0	0.0	0.0	
				020204	0.0			
		1A4c i	Agriculture/Forestry/Fishing, Stationary	020300	0.2	0.2	0.2	
				020302	0.0	0.0	0.0	
		STRAW	1A1a	Electricity and heat production	010101	5.5	4.0	2.5
				010102	3.9	3.2	2.9	
				010103	2.2	1.6	1.5	
				010104	2.0	1.3	1.2	
				010203	4.9	4.6	4.5	
			1A4b i	Residential plants	020200	2.9	2.9	2.9
			1A4c i	Agriculture/Forestry/Fishing, Stationary	020300	1.9	1.9	1.9
				020302		0.0	0.0	
		BIO OIL	1A1a	Electricity and heat production	010102	0.1	0.0	0.1
			010105	0.0	0.0	0.0		

Sum of Fuel_rate_PJ				Year			
fuel_type	fuel_gr_abbr	nfr_id	nfr_name	snap_id	2010	2011	2012
BIOGAS				010202	0.0	0.0	0.1
				010203	1.9	0.7	0.8
		1A2f i	Industry-Other	031600		0.0	0.0
				032005	0.0		
		1A4b i	Residential plants	020200	0.0	0.0	0.1
		1A1a	Electricity and heat production	010101	0.0	0.0	0.0
				010102	0.0	0.0	0.0
				010105	2.0	2.0	2.3
				010203	0.1	0.1	0.1
		1A2e	Food processing, beverages and tobacco	030900	0.2	0.1	0.1
				030902	0.0	0.0	0.0
				030903	0.1	0.1	0.1
		1A2f i	Industry-Other	031300	0.0	0.0	0.0
				032005	0.0	0.0	0.0
		1A4a i	Commercial/Institutional plants	020100	0.3	0.4	0.3
				020103	0.1	0.1	0.1
				020105	0.6	0.6	0.6
		1A4c i	Agriculture/Forestry/Fishing, Stationary	020300	0.2	0.2	0.1
				020304	0.7	0.7	0.6
BIO PROD GAS		1A1a	Electricity and heat production	010105	0.2	0.3	0.4
		1A4a i	Commercial/Institutional plants	020105	0.0	0.0	0.0
Grand Total					553.0	481.9	438.8

Annex 3A-3 Default Lower Calorific Value (LCV) of fuels and fuel correspondence list

Table 3A-3.1a Time-series for calorific values of fuels (DEA 2013a).

		1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
Crude Oil, Average	GJ pr tonne	42.40	42.40	42.40	42.70	42.70	42.70	42.70	43.00	43.00	43.00
Crude Oil, Golf	GJ pr tonne	41.80	41.80	41.80	41.80	41.80	41.80	41.80	41.80	41.80	41.80
Crude Oil, North Sea	GJ pr tonne	42.70	42.70	42.70	42.70	42.70	42.70	42.70	43.00	43.00	43.00
Refinery Feedstocks	GJ pr tonne	41.60	41.60	41.60	41.60	41.60	41.60	41.60	42.70	42.70	42.70
Refinery Gas	GJ pr tonne	52.00	52.00	52.00	52.00	52.00	52.00	52.00	52.00	52.00	52.00
LPG	GJ pr tonne	46.00	46.00	46.00	46.00	46.00	46.00	46.00	46.00	46.00	46.00
Naphtha (LVN)	GJ pr tonne	44.50	44.50	44.50	44.50	44.50	44.50	44.50	44.50	44.50	44.50
Motor Gasoline	GJ pr tonne	43.80	43.80	43.80	43.80	43.80	43.80	43.80	43.80	43.80	43.80
Aviation Gasoline	GJ pr tonne	43.80	43.80	43.80	43.80	43.80	43.80	43.80	43.80	43.80	43.80
JP4	GJ pr tonne	43.80	43.80	43.80	43.80	43.80	43.80	43.80	43.80	43.80	43.80
Other Kerosene	GJ pr tonne	43.50	43.50	43.50	43.50	43.50	43.50	43.50	43.50	43.50	43.50
JP1	GJ pr tonne	43.50	43.50	43.50	43.50	43.50	43.50	43.50	43.50	43.50	43.50
Gas/Diesel Oil	GJ pr tonne	42.70	42.70	42.70	42.70	42.70	42.70	42.70	42.70	42.70	42.70
Fuel Oil	GJ pr tonne	40.40	40.40	40.40	40.40	40.40	40.40	40.70	40.65	40.65	40.65
Orimulsion	GJ pr tonne	27.60	27.60	27.60	27.60	27.60	28.13	28.02	27.72	27.84	27.58
Petroleum Coke	GJ pr tonne	31.40	31.40	31.40	31.40	31.40	31.40	31.40	31.40	31.40	31.40
Waste Oil	GJ pr tonne	41.90	41.90	41.90	41.90	41.90	41.90	41.90	41.90	41.90	41.90
White Spirit	GJ pr tonne	43.50	43.50	43.50	43.50	43.50	43.50	43.50	43.50	43.50	43.50
Bitumen	GJ pr tonne	39.80	39.80	39.80	39.80	39.80	39.80	39.80	39.80	39.80	39.80
Lubricants	GJ pr tonne	41.90	41.90	41.90	41.90	41.90	41.90	41.90	41.90	41.90	41.90
Natural Gas	GJ pr 1000 Nm ³	39.00	39.00	39.00	39.30	39.30	39.30	39.30	39.60	39.90	40.00
Town Gas	GJ pr 1000 m ³							17.00	17.00	17.00	17.00
Electricity Plant Coal	GJ pr tonne	25.30	25.40	25.80	25.20	24.50	24.50	24.70	24.96	25.00	25.00
Other Hard Coal	GJ pr tonne	26.10	26.50	26.50	26.50	26.50	26.50	26.50	26.50	26.50	26.50
Coke	GJ pr tonne	31.80	29.30	29.30	29.30	29.30	29.30	29.30	29.30	29.30	29.30
Brown Coal Briquettes	GJ pr tonne	18.30	18.30	18.30	18.30	18.30	18.30	18.30	18.30	18.30	18.30
Straw	GJ pr tonne	14.50	14.50	14.50	14.50	14.50	14.50	14.50	14.50	14.50	14.50
Wood Chips	GJ pr Cubic metre	2.80	2.80	2.80	2.80	2.80	2.80	2.80	2.80	2.80	2.80
Wood Chips	GJ pr m ³	9.30	9.30	9.30	9.30	9.30	9.30	9.30	9.30	9.30	9.30
Firewood, Hardwood	GJ pr m ³	10.40	10.40	10.40	10.40	10.40	10.40	10.40	10.40	10.40	10.40
Firewood, Conifer	GJ pr tonne	7.60	7.60	7.60	7.60	7.60	7.60	7.60	7.60	7.60	7.60
Wood Pellets	GJ pr tonne	17.50	17.50	17.50	17.50	17.50	17.50	17.50	17.50	17.50	17.50
Wood Waste	GJ pr Cubic metre	14.70	14.70	14.70	14.70	14.70	14.70	14.70	14.70	14.70	14.70
Wood Waste	GJ pr 1000 m ³	3.20	3.20	3.20	3.20	3.20	3.20	3.20	3.20	3.20	3.20
Biogas	GJ pr tonne								23.00	23.00	23.00
Wastes	GJ pr tonne	8.20	8.20	9.00	9.40	9.40	10.00	10.50	10.50	10.50	10.50
Bioethanol	GJ pr tonne	26.70	26.70	26.70	26.70	26.70	26.70	26.70	26.70	26.70	26.70
Liquid Biofuels	GJ pr tonne	37.60	37.60	37.60	37.60	37.60	37.60	37.60	37.60	37.60	37.60
Bio Oil	GJ pr tonne	37.20	37.20	37.20	37.20	37.20	37.20	37.20	37.20	37.20	37.20

Table 3A-3.1b Time-series for calorific values of fuels (DEA 2013a).

<i>Continued</i>		2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Crude Oil, Average	GJ pr tonne	43.00	43.00	43.00	43.00	43.00	43.00	43.00	43.00	43.00	43.00
Crude Oil, Golf	GJ pr tonne	41.80	41.80	41.80	41.80	41.80	41.80	41.80	41.80	41.80	41.80
Crude Oil, North Sea	GJ pr tonne	43.00	43.00	43.00	43.00	43.00	43.00	43.00	43.00	43.00	43.00
Refinery Feedstocks	GJ pr tonne	42.70	42.70	42.70	42.70	42.70	42.70	42.70	42.70	42.70	42.70
Refinery Gas	GJ pr tonne	52.00	52.00	52.00	52.00	52.00	52.00	52.00	52.00	52.00	52.00
LPG	GJ pr tonne	46.00	46.00	46.00	46.00	46.00	46.00	46.00	46.00	46.00	46.00
Naphtha (LVN)	GJ pr tonne	44.50	44.50	44.50	44.50	44.50	44.50	44.50	44.50	44.50	44.50
Motor Gasoline	GJ pr tonne	43.80	43.80	43.80	43.80	43.80	43.80	43.80	43.80	43.80	43.80
Aviation Gasoline	GJ pr tonne	43.80	43.80	43.80	43.80	43.80	43.80	43.80	43.80	43.80	43.80
JP4	GJ pr tonne	43.80	43.80	43.80	43.80	43.80	43.80	43.80	43.80	43.80	43.80
Other Kerosene	GJ pr tonne	43.50	43.50	43.50	43.50	43.50	43.50	43.50	43.50	43.50	43.50
JP1	GJ pr tonne	43.50	43.50	43.50	43.50	43.50	43.50	43.50	43.50	43.50	43.50
Gas/Diesel Oil	GJ pr tonne	42.70	42.70	42.70	42.70	42.70	42.70	42.70	42.70	42.70	42.70
Fuel Oil	GJ pr tonne	40.65	40.65	40.65	40.65	40.65	40.65	40.65	40.65	40.65	40.65
Orimulsion	GJ pr tonne	27.62	27.64	27.71	27.65	27.65	27.65	27.65	27.65	27.65	27.65
Petroleum Coke	GJ pr tonne	31.40	31.40	31.40	31.40	31.40	31.40	31.40	31.40	31.40	31.40
Waste Oil	GJ pr tonne	41.90	41.90	41.90	41.90	41.90	41.90	41.90	41.90	41.90	41.90
White Spirit	GJ pr tonne	43.50	43.50	43.50	43.50	43.50	43.50	43.50	43.50	43.50	43.50
Bitumen	GJ pr tonne	39.80	39.80	39.80	39.80	39.80	39.80	39.80	39.80	39.80	39.80
Lubricants	GJ pr tonne	41.90	41.90	41.90	41.90	41.90	41.90	41.90	41.90	41.90	41.90
Natural Gas	GJ pr 1000 Nm ³	40.15	39.99	40.06	39.94	39.77	39.67	39.54	39.59	39.48	39.46
Town Gas	GJ pr 1000 m ³	17.01	16.88	17.39	16.88	17.58	17.51	17.20	17.14	15.50	21.29
Electricity Plant Coal	GJ pr tonne	24.80	24.90	25.15	24.73	24.60	24.40	24.80	24.40	24.30	24.60
Other Hard Coal	GJ pr tonne	26.50	26.50	26.50	26.50	26.50	26.50	26.50	26.50	25.81	25.13
Coke	GJ pr tonne	29.30	29.30	29.30	29.30	29.30	29.30	29.30	29.30	29.30	29.30
Brown Coal Briquettes	GJ pr tonne	18.30	18.30	18.30	18.30	18.30	18.30	18.30	18.30	18.30	18.30
Straw	GJ pr tonne	14.50	14.50	14.50	14.50	14.50	14.50	14.50	14.50	14.50	14.50
Wood Chips	GJ pr Cubic metre	2.80	2.80	2.80	2.80	2.80	2.80	2.80	2.80	2.80	2.80
Wood Chips	GJ pr m ³	9.30	9.30	9.30	9.30	9.30	9.30	9.30	9.30	9.30	9.30
Firewood, Hardwood	GJ pr m ³	10.40	10.40	10.40	10.40	10.40	10.40	10.40	10.40	10.40	10.40
Firewood, Conifer	GJ pr tonne	7.60	7.60	7.60	7.60	7.60	7.60	7.60	7.60	7.60	7.60
Wood Pellets	GJ pr tonne	17.50	17.50	17.50	17.50	17.50	17.50	17.50	17.50	17.50	17.50
Wood Waste	GJ pr Cubic metre	14.70	14.70	14.70	14.70	14.70	14.70	14.70	14.70	14.70	14.70
Wood Waste	GJ pr 1000 m ³	3.20	3.20	3.20	3.20	3.20	3.20	3.20	3.20	3.20	3.20
Biogas	GJ pr tonne	23.00	23.00	23.00	23.00	23.00	23.00	23.00	23.00	23.00	23.00
Wastes	GJ pr tonne	10.50	10.50	10.50	10.50	10.50	10.50	10.50	10.50	10.50	10.50
Bioethanol	GJ pr tonne	26.70	26.70	26.70	26.70	26.70	26.70	26.70	26.70	26.70	26.70
Liquid Biofuels	GJ pr tonne	37.60	37.60	37.60	37.60	37.60	37.60	37.60	37.60	37.50	37.50
Bio Oil	GJ pr tonne	37.20	37.20	37.20	37.20	37.20	37.20	37.20	37.20	37.20	37.20

Table 3A-3.1c Time-series for calorific values of fuels (DEA 2013a).

<i>Continued</i>		2010	2011	2012
Crude Oil, Average	GJ pr tonne	43.00	43.00	43.00
Crude Oil, Golf	GJ pr tonne	41.80	41.80	41.80
Crude Oil, North Sea	GJ pr tonne	43.00	43.00	43.00
Refinery Feedstocks	GJ pr tonne	42.70	42.70	42.70
Refinery Gas	GJ pr tonne	52.00	52.00	52.00
LPG	GJ pr tonne	46.00	46.00	46.00
Naphtha (LVN)	GJ pr tonne	44.50	44.50	44.50
Motor Gasoline	GJ pr tonne	43.80	43.80	43.80
Aviation Gasoline	GJ pr tonne	43.80	43.80	43.80
JP4	GJ pr tonne	43.80	43.80	43.80
Other Kerosene	GJ pr tonne	43.50	43.50	43.50
JP1	GJ pr tonne	43.50	43.50	43.50
Gas/Diesel Oil	GJ pr tonne	42.70	42.70	42.70
Fuel Oil	GJ pr tonne	40.65	40.65	40.65
Orimulsion	GJ pr tonne	27.65	27.65	27.65
Petroleum Coke	GJ pr tonne	31.40	31.40	31.40
Waste Oil	GJ pr tonne	41.90	41.90	41.90
White Spirit	GJ pr tonne	43.50	43.50	43.50
Bitumen	GJ pr tonne	39.80	39.80	39.80
Lubricants	GJ pr tonne	41.90	41.90	41.90
Natural Gas	GJ pr 1000 Nm3	39.46	39.51	39.55
Town Gas	GJ pr 1000 m3	21.35	21.37	21.37
Electricity Plant Coal	GJ pr tonne	24.44	24.38	24.23
Other Hard Coal	GJ pr tonne	24.44	24.38	24.23
Coke	GJ pr tonne	29.30	29.30	29.30
Brown Coal Briquettes	GJ pr tonne	18.30	18.30	18.30
Straw	GJ pr tonne	14.50	14.50	14.50
Wood Chips	GJ pr Cubic metre	2.80	2.80	2.80
Wood Chips	GJ pr m3	9.30	9.30	9.30
Firewood, Hardwood	GJ pr m3	10.40	10.40	10.40
Firewood, Conifer	GJ pr tonne	7.60	7.60	7.60
Wood Pellets	GJ pr tonne	17.50	17.50	17.50
Wood Waste	GJ pr Cubic metre	14.70	14.70	14.70
Wood Waste	GJ pr 1000 m3	3.20	3.20	3.20
Biogas	GJ pr tonne	23.00	23.00	23.00
Wastes	GJ pr tonne	10.50	10.50	10.50
Bioethanol	GJ pr tonne	26.70	26.70	26.70
Liquid Biofuels	GJ pr tonne	37.50	37.50	37.50
Bio Oil	GJ pr tonne	37.20	37.20	37.20

Table 3A-3.2 category correspondence list, DEA, DCE and Climate Convention reportings (CRF).

Danish Energy Agency	DCE Emission database	IPCC fuel category
Other Hard Coal	Coal	Solid
Coke	Coke oven coke	Solid
Electricity Plant Coal	Coal	Solid
Brown Coal Briquettes	Brown coal briq.	Solid
-	Anode carbon	Solid
-	Fly ash	Solid
Orimulsion	Orimulsion	Liquid
Petroleum Coke	Petroleum coke	Liquid
Fuel Oil	Residual oil	Liquid
Waste Oil	Residual oil	Liquid
Gas/Diesel Oil	Gas oil	Liquid
Other Kerosene	Kerosene	Liquid
LPG	LPG	Liquid
Refinery Gas	Refinery gas	Liquid
Town Gas	Natural gas	Gas
Natural Gas	Natural gas	Gas
Straw	Straw	Biomass
Wood Waste	Wood and simil.	Biomass
Wood Pellets	Wood and simil.	Biomass
Wood Chips	Wood and simil.	Biomass
Firewood, Hardwood & Conifer	Wood and simil.	Biomass
Waste Combustion (biomass)	Municip. wastes	Biomass
Bio Oil	Bio oil	Biomass
Biogas	Biogas	Biomass
Biogas, other	Biogas	Biomass
Biogas, landfill	Biogas	Biomass
Biogas, sewage sludge	Biogas	Biomass
(Wood applied in gas engines)	Biomass producer gas	Biomass
Waste Combustion (fossil)	Fossil waste	Other fuel

Annex 3A-4 Emission factors

Table 3A-4.1 CO₂ emission factors, 2012.

Fuel	Emission factor kg per GJ		Reference type	IPCC fuel category
	Biomass	Fossil fuel		
Coal, source category 1A1a Public electricity and heat production		94.25 ¹⁾	Country specific	Solid
Coal, Other source categories		94.6 ³⁾	IPCC (1997)	Solid
Brown coal briquettes		94.6	IPCC (1997)	Solid
Coke oven coke		108	IPCC (1997)	Solid
Other solid fossil fuels ⁶⁾		108 ¹⁾	IPCC (1997)	Solid
Fly ash fossil (from coal)		93.6	Country specific	Solid
Petroleum coke		93 ³⁾	Country specific	Liquid
Residual oil, source category 1A1a Public electricity and heat production		79.21 ¹⁾	Country specific	Liquid
Residual oil, other source categories		77.4 ³⁾	IPCC (1997)	Liquid
Gas oil		74 ¹⁾	EEA (2007)	Liquid
Kerosene		71.9	IPCC (1997)	Liquid
Orimulsion		80 ²⁾	Country specific	Liquid
LPG		63.1	IPCC (1997)	Liquid
Refinery gas		58.099	Country specific	Liquid
Natural gas, off shore gas turbines		57.423	Country specific	Gas
Natural gas, other		57.03	Country specific	Gas
Waste	75.1 ³⁾⁴⁾	+ 37 ³⁾⁴⁾	Country specific	Biomass and Other fuels
Straw	110		IPCC (1997)	Biomass
Wood	110		IPCC (1997)	Biomass
Bio oil	74		Country specific	Biomass
Biogas	83.6		Country specific	Biomass
Biomass producer gas	142.9 ⁵⁾		Country specific	Biomass

1) Plant specific data from EU ETS incorporated for individual plants.

2) Not applied in 2012. Orimulsion was applied in Denmark in 1995 – 2004.

3) Plant specific data from EU ETS incorporated for cement industry and sugar, lime and glass wool production.

4) The emission factor for waste is (37+75.1) kg CO₂ per GJ waste. The fuel consumption and the CO₂ emission have been disaggregated to the two IPCC fuel categories *Biomass* and *Other fuels* in CRF. The IEF¹ for CO₂, Other fuels is 82.22 kg CO₂ per GJ fossil waste.

5) Includes a high content of CO₂ in the gas.

6) Anodic carbon

Time series have been estimated for coal (1A1a), residual oil (1A1a), refinery gas, natural gas fuelled off shore gas turbines, other natural gas consumption and the biomass part of industrial waste. For all other fuels the same emission factor has been applied for 1990-2012.

¹ Not including cement production.

Table 3A-4.2 CO₂ emission factors, time-series.

Year	Coal, sector 1A1a, kg per GJ	Residual oil, sector 1A1a, kg per GJ	Refinery gas, kg per GJ	Natural gas, off shore gas turbines, kg per GJ	Natural gas, other, kg per GJ	Industrial waste, biomass part
1990	94	78.4	57.6	57.469	56.9	86.7
1991	94	78.4	57.6	57.469	56.9	86.7
1992	94	78.4	57.6	57.469	56.9	84.2
1993	94	78.4	57.6	57.469	56.9	83
1994	94	78.4	57.6	57.469	56.9	83
1995	94	78.4	57.6	57.469	56.9	81.1
1996	94	78.4	57.6	57.469	56.9	79.6
1997	94	78.4	57.6	57.469	56.9	79.6
1998	94	78.4	57.6	57.469	56.9	79.6
1999	94	78.4	57.6	57.469	56.9	79.6
2000	94	78.4	57.6	57.469	57.1	79.6
2001	94	78.4	57.6	57.469	57.25	79.6
2002	94	78.4	57.6	57.469	57.28	79.6
2003	94	78.4	57.6	57.469	57.19	79.6
2004	94	78.4	57.6	57.469	57.12	79.6
2005	94	78.4	57.6	57.469	56.96	79.6
2006	94.4	78.2	57.812	57.879	56.78	79.6
2007	94.3	78.1	57.848	57.784	56.78	79.6
2008	94	78.5	57.948	56.959	56.77	79.6
2009	93.6	78.9	56.814	57.254	56.69	79.6
2010	93.6	79.2	57.134	57.314	56.74	79.6
2011	93.73	79.25	57.881	57.379	56.97	79.6
2012	93.25	79.21	58.099	57.423	57.03	79.6

Table 3A-4.3 CH₄ emission factors and references, 2012.

Fuel group	Fuel	CRF source category	CRF source category	SNAP	Emission factor, g pr GJ	Reference
SOLID	COAL	1A1a	Public electricity and heat production	0101 0102	0.9	IPCC (1997), Tier 2, Table 1-15, Utility Boiler, Pulverised Bituminous Combustion, Wet bottom.
		1A2 a-f	Industry	03	10	IPCC (1997), Tier 2, Table 1-19, Commercial coal boilers.
		1A4b i	Residential	0202	300	IPCC (1997), Tier 1, Table 1-7, Residential, coal.
		1A4c i	Agriculture/ Forestry	0203	10	IPCC (1997), Tier 2, Table 1-19, Commercial coal boilers. ¹⁾
	BROWN COAL BRI.	1A4b i	Residential	0202	300	IPCC (1997), Tier 1, Table 1-7, Residential, coal.
	COKE OVEN COKE	1A2 A-f	Industry	03	10	IPCC (1997), Tier 2, Table 1-19, Commercial coal boilers.
		1A4b i	Residential	0202	300	IPCC (1997), Tier 1, Table 1-7, Residential, coal.
	ANODIC CARBON	1A2a-f	Industry	03	10	IPCC (1997), Tier 2, Table 1-19, Commercial coal boilers.
	FOSSIL FLY ASH	1A1a	Public electricity and heat production	0101	0.9	IPCC (1997), Tier 2, Table 1-15, Utility Boiler, Pulverised Bituminous Combustion, Wet bottom.
LIQUID	PETROLEUM COKE	1A2a-f	Industry	03	2	IPCC (1997), Tier 1, Table 1-7, Industry, oil.
	RESIDUAL OIL	1A1a	Public electricity and heat production	010101	0.9	IPCC (1997), Tier 2, Table 1-15, Utility Boiler, Residual fuel oil.
				010102	1.3	Nielsen et al. (2010)
				010103		
				010104	3	IPCC (1997), Tier 1, Table 1-7, Energy industries, oil.
				010203	0.9	IPCC (1997), Tier 2, Table 1-15, Utility Boiler, Residual fuel oil.
		1A1b	Petroleum refining	010306	3	IPCC (1997), Tier 1, Table 1-7, Energy industries, oil.
		1A2 a-f	Industry	03	1.3	Nielsen et al. (2010)
				Engines	4	IPCC (1997), Tier 2, Table 1-15, Utility, Large diesel engines
		1A4a	Commercial/ Institutional	0201	1.4	IPCC (1997), Tier 2, Table 1-19, Commercial, residual fuel oil ¹⁾ .
		1A4b	Residential	0202	1.4	IPCC (1997), Tier 2, Table 1-19, Commercial, residual fuel oil ¹⁾ .
		1A4c	Agriculture/ Forestry	0203	1.4	IPCC (1997), Tier 2, Table 1-19, Commercial, residual fuel oil ¹⁾ .
	GAS OIL	1A1a	Public electricity and heat production	010101	0.9	IPCC (1997), Tier 2, Table 1-15, Utility Boiler, distillate fuel oil.
				010102		
				010103		
				010104	3	IPCC (1997), Tier 1, Table 1-7, Energy industries, oil.
				010105	24	Nielsen et al. (2010)
				010202	0.9	IPCC (1997), Tier 2, Table 1-15, Utility Boiler, distillate fuel oil.
				010203		
		1A1b	Petroleum refining	010306	3	IPCC (1997), Tier 1, Table 1-7, Energy industries, oil.
		1A1c	Other energy industries	010504	3	IPCC (1997), Tier 1, Table 1-7, Energy industries, oil.
		1A2 a-f	Industry	03	0.2	IPCC (1997), Tier 2, Table 1-16, Industry, distillate fuel oil.
				Turbines	2	IPCC (1997), Tier 1, Table 1-7, Industry, oil.
				Engines	24	Nielsen et al. (2010)

Fuel group	Fuel	CRF source category	CRF source category	SNAP	Emission factor, g pr GJ	Reference
	KEROSENE	1A4a	Commercial/ Institutional	0201	0.7	IPCC (1997), Tier 2, Table 1-19, Commercial, distillate fuel oil.
				020105	24	Nielsen et al. (2010)
		1A4b i	Residential	0202	0.7	IPCC (1997), Tier 2, Table 1-18, Residential, distillate fuel oil.
		1A4c	Agriculture/ Forestry	0203	0.7	IPCC (1997), Tier 2, Table 1-19, Commercial, distillate fuel oil.
		1A2 a-f	Industry	all	0.2	IPCC (1997), Tier 2, Table 1-16, Industry, distillate fuel oil.
		1A4a	Commercial/ Institutional	0201	0.7	IPCC (1997), Tier 2, Table 1-19, Commercial, distillate fuel oil.
		1A4b i	Residential	0202	0.7	IPCC (1997), Tier 2, Table 1-18, Residential, distillate fuel oil.
		1A4c i	Agriculture/ Forestry	0203	0.7	IPCC (1997), Tier 2, Table 1-19, Commercial, distillate fuel oil ¹⁾ .
		1A1a	Public electricity and heat production	0101 0102	3	IPCC (1997), Tier 1, Table 1-7, Energy Industries, oil.
		1A1b	Petroleum refining	0103	3	IPCC (1997), Tier 1, Table 1-7, Energy Industries, oil.
	LPG	1A2 a-f	Industry	03	2	IPCC (1997), Tier 1, Table 1-7, Industry, oil
		1A4a	Commercial/ Institutional	0201	10	IPCC (1997), Tier 1, Table 1-7, Commercial, oil.
		1A4b i	Residential	0202	1.1	IPCC (1997), Tier 2, Table 1-18, Residential propane/butane furnaces.
		1A4c i	Agriculture/ Forestry	0203	10	IPCC (1997), Tier 1, Table 1-7, Agriculture, oil.
	REFINERY GAS	1A1b	Petroleum refining	010304	1.7	Assumed equal to natural gas fuelled gas turbines. Nielsen et al. (2010)
				010306	1	Assumed equal to natural gas fuelled plants. IPCC (1997), Tier 1, Table 1-7, Natural gas
GAS	NATURAL GAS	1A1a	Public electricity and heat production	010101	0.1	IPCC (1997), Tier 2, Table 1-15, Utility Boiler, natural gas.
				010102		
				010103		
				010104	1.7	Nielsen et al. (2010)
				010105	481	Nielsen et al. (2010)
				010202	0.1	IPCC (1997), Tier 2, Table 1-15, Utility Boiler, natural gas.
				010203		
		1A1c	Other energy industries	010504	1.7	Nielsen et al. (2010)
		1A2 a-f	Industry	Other	1.4	IPCC (1997), Tier 2, Table 1-16, Industry, natural gas boilers.
				Gas turbines	1.7	Nielsen et al. (2010)
				Engines	481	Nielsen et al. (2010)
		1A4a	Commercial/ Institutional	0201	1.2	IPCC (1997), Tier 2, Table 1-19, Commercial, natural gas boilers.
				020105	481	Nielsen et al. (2010)
		1A4b i	Residential	0202	5	IPCC (1997), Tier 1, Table 1-7, Residential, natural gas.
				020204	481	Nielsen et al. (2010)
		1A4c i	Agriculture/ Forestry	0203	1.2	IPCC (1997), Tier 2, Table 1-19, Commercial, natural gas boilers ¹⁾ .
				020304	481	Nielsen et al. (2010)
WASTE	WASTE	1A1a	Public electricity and heat production	0101 0102	0.34	Nielsen et al. (2010)
		1A2a-f	Industry	03	30	IPCC (1997), Tier 1, Table 1-7, Industry, wastes.

Fuel group	Fuel	CRF source category	CRF source category	SNAP	Emission factor, g pr GJ	Reference	
BIO-MASS	INDUSTRIAL WASTE	1A4a	Commercial/ Institutional	0201	30	IPCC (1997), Tier 1, Table 1-7, Industry, wastes.	
		1A2f	Industry	0316	30	IPCC (1997), Tier 1, Table 1-7, Industry, wastes.	
	WOOD	1A1a	Public electricity and heat production	0101	3.1	Nielsen et al. (2010)	
				0102	30	IPCC (1997), Tier 1, Table 1-7, Energy industries, wood	
		1A2 a-f	Industry	03	15	IPCC (1997), Tier 2, Table 1-16, Industry, wood stoker boilers.	
		1A4a	Commercial/ Institutional	0201	30	IPCC (1997), Tier 1, Table 1-7, Industry, wood ²⁾ .	
		1A4b i	Residential	0202	129.3	DCE estimate based on technology distribution ³⁾	
		1A4c i	Agriculture/ Forestry	0203	30	IPCC (1997), Tier 1, Table 1-7, Industry, wood ²⁾ .	
	STRAW	1A1a	Public electricity and heat production	0101	0.47	Nielsen et al. (2010)	
				0102	30	IPCC (1997), Tier 1, Table 1-7, Energy industries, other biomass	
		1A4b i	Residential	0202	300	IPCC (1997), Tier 1, Table 1-7, Residential, other biomass.	
		1A4c i	Agriculture/ Forestry	020300	300	IPCC (1997), Tier 1, Table 1-7, Agriculture, other biomass.	
				020302	30	IPCC (1997), Tier 1, Table 1-7, Energy industries, other biomass	
BIO OIL		1A1a	Public electricity and heat production	010102	0.9	IPCC (1997), Tier 2, Table 1-15, Utility Boiler, distillate fuel oil.	
	010105			24	Nielsen et al. (2010) assumed same emission factor as for gas oil fuelled engines.		
	0102			0.7	IPCC (1997), Tier 2, Table 1-19, Commercial, distillate fuel oil.		
	1A2 a-f			Industry	03	0.7	Assumed equal to district heating plants.
	1A4b i			Residential	0202	0.7	IPCC (1997), Tier 2, Table 1-18, Residential, distillate fuel oil.
	BIOGAS			1A1a	Public electricity and heat production	0101	1
010105		434	Nielsen et al. (2010)				
0102		1	IPCC (1997), Tier 1, Table 1-7, Energy industries, natural gas. Assumed similar to natural gas (DCE assumption).				
1A2 a-f		Industry	03	5	IPCC (1997), Tier 1, Table 1-7, Industry, natural gas. Assumed similar to natural gas (DCE assumption).		
			Engines	434	Nielsen et al. (2010)		
1A4a		Commercial/ Institutional	0201	5	IPCC (1997), Tier 1, Table 1-7, Commercial, natural gas. Assumed similar to natural gas (DCE assumption).		
			020105	434	Nielsen et al. (2010)		
1A4c i		Agriculture/ Forestry	0203	5	IPCC (1997), Tier 1, Table 1-7, Agriculture, natural gas. Assumed similar to natural gas (DCE assumption).		
			020304	434	Nielsen et al. (2010)		
BIO PROD GAS		1A1a	Public electricity and heat production	010105	13	Nielsen et al. (2010)	
	1A4a	Commercial/Institutional	020105	13	Nielsen et al. (2010)		

- 1) Assumed same emission factors as for commercial plants. Plant capacity and technology are similar for Danish plants.
- 2) Assumed same emission factor as for industrial plants. Plant capacity and technology is similar to industrial plants rather than to residential plants.
- 3) Aggregated emission factor based on the technology distribution in the sector (DEPA, 2013) and technology specific emission factors that refer to: Paulrud et al. (2005), Johansson et al. (2004), Olsson & Kjällstrand (2005) and Johansson et al. (2005).

In general, the same emission factors have been applied for 1990-2012. However, time series have been estimated for both natural gas fuelled engines and biogas fuelled engines, residential wood combustion, natural gas fuelled gas turbines² and waste incineration plants².

Table 3A-4.4 CH₄ emission factors, time-series.

Year	Natural gas fuelled engines Emission factor, g per GJ	Biogas fuelled engines Emission factor, g per GJ	Residential wood combustion, g per GJ	Waste incineration g per GJ	Natural gas fuelled gas turbines, g per GJ
1990	266	239	224.0	0.59	1.5
1991	309	251	224.0	0.59	1.5
1992	359	264	224.0	0.59	1.5
1993	562	276	224.0	0.59	1.5
1994	623	289	224.0	0.59	1.5
1995	632	301	224.0	0.59	1.5
1996	616	305	224.0	0.59	1.5
1997	551	310	224.0	0.59	1.5
1998	542	314	224.0	0.59	1.5
1999	541	318	224.0	0.59	1.5
2000	537	323	224.0	0.59	1.5
2001	522	342	199.3	0.59	1.5
2002	508	360	189.6	0.59	1.6
2003	494	379	187.2	0.59	1.6
2004	479	397	184.5	0.51	1.7
2005	465	416	175.5	0.42	1.7
2006	473	434	165.2	0.34	1.7
2007	481	434	166.5	0.34	1.7
2008	481	434	156.5	0.34	1.7
2009	481	434	143.8	0.34	1.7
2010	481	434	136.6	0.34	1.7
2011	481	434	129.3	0.34	1.7
2012	481	434	123.7	0.34	1.7

² A minor emission source.

Table 3A-4.5 N₂O emission factors and references 2012.

Fuel group	Fuel	CRF source category	CRF source category	SNAP	Emission factor, g per GJ	Reference
SOLID	COAL	1A1a	Public electricity and heat production	0101	0.8	Elsam (2005)
				0102	1.6	IPCC (1997), Tier 2, Table 1-15, Utility boiler, pulverised bituminous coal.
		1A2 a-f	Industry	03	1.4	IPCC (1997), Tier 1, Table 1-8, Industry, coal
		1A4b i	Residential	0202	1.4	IPCC (1997), Tier 1, Table 1-8, Residential, coal
		1A4c i	Agriculture/ Forestry	0203	1.4	IPCC (1997), Tier 1, Table 1-8, Commercial, coal
	BROWN COAL BRI.	1A4b i	Residential	0202	1.4	IPCC (1997), Tier 1, Table 1-8, Residential, coal
	COKE OVEN COKE	1A2 a-f	Industry	03	1.4	IPCC (1997), Tier 1, Table 1-8, Industry, coal
		1A4b i	Residential	020200	1.4	IPCC (1997), Tier 1, Table 1-8, Residential, coal
	ANODIC CARBON	1A2 a-f	Industry	03	1.4	IPCC (1997), Tier 1, Table 1-8, Industry, coal
LI-QUID	PETROLEUM COKE	1A2a-f	Industry – other	03	0.6	IPCC (1997), Tier 1, Table 1-8, Industry, oil
	RESIDUAL OIL	1A1a	Public electricity and heat production	010101	0.3	IPCC (1997), Tier 2, Table 1-15, Utility, residual fuel oil
				010102	5	Nielsen et al. (2010)
				010103		
				010104	0.6	IPCC (1997), Tier 1, Table 1-8, Energy industries, oil
		010203			0.3	IPCC (1997), Tier 2, Table 1-15, Utility, residual fuel oil
		1A1b	Petroleum refining	010306	0.6	IPCC (1997), Tier 1, Table 1-8, Energy industries, oil
		1A2 a-f	Industry	03	5	Nielsen et al. (2010)
				Engines	0.6	IPCC (1997), Tier 2, Table 1-15, Utility, residual fuel oil
		1A4a	Commercial/ Institutional	0201	0.3	IPCC (1997), Tier 2, Table 1-19, Commercial, fuel oil
		1A4b i	Residential	0202	0.6	IPCC (1997), Tier 1, Table 1-8, Residential, oil
		1A4c i	Agriculture/ Forestry	0203	0.3	IPCC (1997), Tier 2, Table 1-19, Commercial, fuel oil
	GAS OIL	1A1a	Public electricity and heat production	010101	0.4	IPCC (1997), Tier 2, Table 1-15, Utility, distillate fuel oil
				010102		
				010103		
				010104	0.6	IPCC (1997), Tier 1, Table 1-8, Energy industries, oil
				010105	2.1	Nielsen et al. (2010)
				0102	0.4	IPCC (1997), Tier 2, Table 1-15, Utility, distillate fuel oil
		1A1b	Petroleum refining	010306	0.6	IPCC (1997), Tier 1, Table 1-8, Energy industries, oil
		1A2 a-f	Industry	03	0.4	IPCC (1997), Tier 2, Table 1-16, Industry, distillate fuel oil boilers
		Turbin- Engines			0.6	IPCC (1997), Tier 1, Table 1-8, Industry, oil
		1A4a	Commercial/ Institutional	0201	0.4	IPCC (1997), Tier 2, Table 1-19, Commercial, distillate fuel oil
		Engines			2.1	Nielsen et al. (2010)
		1A4b i	Residential	0202	0.6	IPCC (1997), Tier 1, Table 1-8, Residential, oil

Fuel group	Fuel	CRF source category	CRF source category	SNAP	Emission factor, g per GJ	Reference
GAS	KEROSENE	1A4c	Agriculture/ Forestry	0203	0.4	IPCC (1997), Tier 2, Table 1-19, Commercial, distillate fuel oil
		1A2 a-f	Industry	03	0.4	IPCC (1997), Tier 2, Table 1-16, Industry, distillate fuel oil boilers
		1A4a	Commercial/ Institutional	0201	0.4	IPCC (1997), Tier 2, Table 1-19, Commercial, distillate fuel oil
		1A4b i	Residential	0202	0.6	IPCC (1997), Tier 1, Table 1-8, Residential, oil
		1A4c i	Agriculture/ Forestry	0203	0.4	IPCC (1997), Tier 2, Table 1-19, Commercial, distillate fuel oil ¹⁾
		1A1a	Public electricity and heat production	0101 0102	0.6	IPCC (1997), Tier 1, Table 1-8, Energy industries, oil
		1A1b	Petroleum refining	010306	0.6	IPCC (1997), Tier 1, Table 1-8, Energy industries, oil
		1A2 a-f	Industry	03	0.6	IPCC (1997), Tier 1, Table 1-8, Industry, oil
		1A4a	Commercial/ Institutional	0201	0.6	IPCC (1997), Tier 1, Table 1-8, Commercial, oil
		1A4b i	Residential	0202	0.6	IPCC (1997), Tier 1, Table 1-8, Residential, oil
		1A4c i	Agriculture/ Forestry	0203	0.6	IPCC (1997), Tier 1, Table 1-8, Agriculture, oil
	REFINERY GAS	1A1b	Petroleum refining	010304	1	Assumed equal to natural gas fuelled turbines. Based on Nielsen et al. (2010).
				010306	0.1	IPCC (1997), Tier 1, Table 1-8, Energy industries, natural gas
	NATURAL GAS	1A1a	Public electricity and heat production	010101	0.1	IPCC (1997), Tier 1, Table 1-8, Energy industries, natural gas
				010102		
				010103		
				010104	1	Nielsen et al. (2010)
				010105	0.58	Nielsen et al. (2010)
				0102	0.1	IPCC (1997), Tier 1, Table 1-8, Energy industries, natural gas
		1A1c	Other energy industries	010504	1	Nielsen et al. (2010)
		1A2 a-f	Industry	03	0.1	IPCC (1997), Tier 1, Table 1-8, Industry, natural gas
				Gas turbines	1	Nielsen et al. (2010)
				Engines	0.58	Nielsen et al. (2010)
		1A4a	Commercial/ Institutional	020100	2.3	IPCC (1997), Tier 2, Table 1-19, Commercial, natural gas boilers
				020103		
				Engines	0.58	Nielsen et al. (2010)
		1A4b i	Residential	0202	0.1	IPCC (1997), Tier 1, Table 1-8, Residential, natural gas
				Engines	0.58	Nielsen et al. (2010)
		1A4c i	Agriculture/ Forestry	0203	2.3	IPCC (1997), Tier 2, Table 1-19, Commercial, natural gas boilers ¹⁾
				Engines	0.58	Nielsen et al. (2010)
WA-STE	WASTE	1A1a	Public electricity and heat production	0101 0102	1.2	Nielsen et al. (2010)
		1A2 a-f	Industry	03	4	IPCC (1997), Tier 1, Table 1-8, Industry, wastes
		1A4a	Commercial/ Institutional	0201	4	IPCC (1997), Tier 1, Table 1-8, Commercial, wastes
	INDUSTR. WA-STE	1A2a-f	Industry	03	4	IPCC (1997), Tier 1, Table 1-8, Industry, waste
BIO-MASS	WOOD	1A1a	Public electricity and heat production	0101	0.8	Nielsen et al. (2010)
				0102	4	IPCC (1997), Tier 1, Table 1-8, Energy industries, wood

Fuel group	Fuel	CRF source category	CRF source category	SNAP	Emission factor, g per GJ	Reference
		1A2 a-f	Industry	03	4	IPCC (1997), Tier 1, Table 1-8, Industry, wood
		1A4a	Commercial/ Institutional	0201	4	IPCC (1997), Tier 1, Table 1-8, Commercial, wood
		1A4b i	Residential	0202	4	IPCC (1997), Tier 1, Table 1-8, Residential, wood
		1A4c i	Agriculture/ Forestry	0203	4	IPCC (1997), Tier 1, Table 1-8, Agriculture, wood
	STRAW	1A1a	Public electricity and heat production	0101	1.1	Nielsen et al. (2010)
				0102	4	IPCC (1997), Tier 1, Table 1-8, Energy industries, other biomass
		1A4b i	Residential	0202	4	IPCC (1997), Tier 1, Table 1-8, Residential, other biomass
		1A4c i	Agriculture/ Forestry	0203	4	IPCC (1997), Tier 1, Table 1-8, Agriculture, other biomass
	BIO OIL	1A1a	Public electricity and heat production	0101 0102 Engines	0.4 2.1	IPCC (1997), Tier 2, Table 1-15, Utility, distillate fuel oil Assumed equal to gas oil. Based on Nielsen et al. (2010)
		1A2 a-f	Industry	03	0.4	IPCC (1997), Tier 1, Table 1-8, Industry, oil
		1A4b i	Residential	0202	0.6	IPCC (1997), Tier 1, Table 1-8, Residential, oil
	BIOGAS	1A1a	Public electricity and heat production	0101 0102 Engines	0.1 1.6	IPCC (1997), Tier 1, Table 1-8, Energy industries, natural gas Nielsen et al. (2010)
		1A2 a-f	Industry	03 Engines	0.1 1.6	IPCC (1997), Tier 1, Table 1-8, Industry, natural gas Nielsen et al. (2010)
		1A4a	Commercial/ Institutional	0201 Engines	0.1 1.6	IPCC (1997), Tier 1, Table 1-8, Commercial, natural gas Nielsen et al. (2010)
		1A4c i	Agriculture/ Forestry	0203 Engines	0.1 1.6	IPCC (1997), Tier 1, Table 1-8, Agriculture, natural gas Nielsen et al. (2010)
	BIO PROD GAS	1A1a	Public electricity and heat production	010105	2.7	Nielsen et al. (2010)
		1A4a	Commercial/ Institutional	020105	2.7	Nielsen et al. (2010)

1) In Denmark, plants in Agriculture/Forestry are similar to Commercial plants.

Time-series have been estimated for natural gas fuelled gas turbines and refinery gas fuelled gas turbines. All other emission factors have been applied unchanged for 1990-2012.

Table 3A-4.6 N₂O emission factors, time-series.

Year	Natural gas fuelled gas turbines. Emission factor, g per GJ	Refinery gas fuelled gas turbines. Emission factor, g per GJ
1990	2.2	2.2
1991	2.2	2.2
1992	2.2	2.2
1993	2.2	2.2
1994	2.2	2.2
1995	2.2	2.2
1996	2.2	2.2
1997	2.2	2.2
1998	2.2	2.2
1999	2.2	2.2
2000	2.2	2.2
2001	2.0	2.0
2002	1.9	1.9
2003	1.7	1.7
2004	1.5	1.5
2005	1.4	1.4
2006	1.2	1.2
2007	1.0	1.0
2008	1.0	1.0
2009	1.0	1.0
2010	1.0	1.0
2011	1.0	1.0
2012	1.0	1.0

Table 3A-4.7 SO₂ emission factors and references 2012.

Fuel type	Fuel	NFR	NFR_name	SNAP	SO ₂ emission factor, g/GJ	Reference
SOLID	ANODE CARBON	1A2f	Industry - other	032000	574	Assumed equal to coal. DCE assumption.
	COAL	1A1a	Public electricity and heat production	0101	11	DCE estimate based on data reported by plant owners to the electricity transmission company, Energinet.dk (Energinet.dk, 2013)
				0102	574	DCE calculation based on DEPA (2010c), DEA (2012a) and EMEP (2006)
		1A2a-f	Industry	03	574	DCE calculation based on DEPA (2010c), DEA (2012a) and EMEP (2006)
		1A4b i	Residential	020200	574	DCE calculation based on DEPA (2010c), DEA (2012a) and EMEP (2006)
		1A4c i	Agriculture/ Forestry	0203	574	DCE calculation based on DEPA (2010c), DEA (2012a) and EMEP (2006)
	FLY ASH FOSSIL	1A1a	Public electricity and heat production	010104	10	Assumed equal to the emission factor for coal in 2010. DCE assumption.
	BROWN COAL BRI.	1A4b	Residential	0202	574	Assumed equal to coal. DCE assumption.
	COKE OVEN COKE	1A2a-f	Industry	03	574	Assumed equal to coal. DCE assumption.
		1A4b i	Residential	020200	574	Assumed equal to coal. DCE assumption.
LIQUID	PETROLEUM COKE	1A2a-f	Industry	03	605	DCE calculation based on DEPA (2001b), DEA (2012a) and EMEP (2006).
	RESIDUAL OIL	1A1a	Electricity and heat production	0101	100	DCE estimate based on plant specific data for 2008 and 2009.
				0102	344	DCE estimate based on EOF (2013) and DEA (2012a)
		1A1b	Petroleum refining	010306	537	DCE calculation based on plant specific data for year 2003.
		1A2a-f	Industry	03	344	DCE estimate based on EOF (2013) and DEA (2012a)
		1A4a	Commercial/ Institutional	0201	344	DCE estimate based on EOF (2013) and DEA (2012a)
		1A4b	Residential	0202	344	DCE estimate based on EOF (2013) and DEA (2012a)
		1A4c i	Agriculture/ Forestry	0203	344	DCE estimate based on EOF (2013) and DEA (2012a)
	GAS OIL	1A1a	Public electricity and heat production	0101	23	DCE estimate based on DEPA (1998), Miljø- og planlægningsudvalget (1998) and DEA (2012a).
				0102		
		1A1b	Petroleum refining	010306	23	DCE estimate based on DEPA (1998), Miljø- og planlægningsudvalget (1998) and DEA (2012a).
		1A1c	Other energy industries	0105	23	DCE estimate based on DEPA (1998), Miljø- og planlægningsudvalget (1998) and DEA (2012a).
		1A2a-f	Industry	03	23	DCE estimate based on DEPA (1998), Miljø- og planlægningsudvalget (1998) and DEA (2012a).
		1A4a	Commercial/ Institutional	0201	23	DCE estimate based on DEPA (1998), Miljø- og planlægningsudvalget (1998) and DEA (2012a).
		1A4b i	Residential	0202	23	DCE estimate based on DEPA (1998), Miljø- og planlægningsudvalget (1998) and DEA (2012a).
		1A4c	Agriculture/Forestry	0203	23	DCE estimate based on DEPA (1998), Miljø- og planlægningsudvalget (1998) and DEA (2012a).
	KEROSENE	1A2f	Industry - other	03	5	DCE estimate based on Tønder (2004) and Shell (2013).
		1A4a	Commercial/ Institutional	0201	5	DCE estimate based on Tønder (2004) and Shell (2013).
		1A4b i	Residential	0202	5	DCE estimate based on Tønder (2004) and Shell (2013).
		1A4c i	Agriculture/ Forestry	0203	5	DCE estimate based on Tønder (2004) and Shell (2013).
	LPG	1A1a	Public electricity and heat production	All	0.13	DCE estimate based on Augustesen (2003) and DEA (2012a).

Fuel type	Fuel	NFR	NFR_name	SNAP	SO ₂ emission factor, g/GJ	Reference
		1A2a-f	Industry	03	0.13	DCE estimate based on Augustesen (2003) and DEA (2012a).
		1A4a	Commercial/ Institutional	0201	0.13	DCE estimate based on Augustesen (2003) and DEA (2012a).
		1A4b i	Residential	0202	0.13	DCE estimate based on Augustesen (2003) and DEA (2012a).
		1A4c i	Agriculture/ Forestry	0203	0.13	DCE estimate based on Augustesen (2003) and DEA (2012a).
	REFINERY GAS	1A1b	Petroleum refining	0103	1	DCE estimate based on plant specific data for one plant, average value for 1995-2002.
GAS	NATURAL GAS	1A1a	Public electricity and heat production	0101, 0102, except engines	0.43	DCE estimate based on data from Energinet.dk (2013b)
				010105, engines	0.5	Kristensen (2003)
		1A1c	Other energy industries	0105	0.43	DCE estimate based on data from Energinet.dk (2013b)
		1A2a-f	Industry	03 except engines	0.43	DCE estimate based on data from Energinet.dk (2013b)
				Engines	0.5	Kristensen (2003)
		1A4a	Commercial/ Institutional	0201 except engines	0.43	DCE estimate based on data from Energinet.dk (2013b)
				Engines	0.5	Kristensen (2003)
		1A4b i	Residential	0202 except engines	0.43	DCE estimate based on data from Energinet.dk (2013b)
				Engines	0.5	Kristensen (2003)
		1A4c i	Agriculture/ Forestry	0203 except engines	0.43	DCE estimate based on data from Energinet.dk (2013b)
				Engines	0.5	Kristensen (2003)
WASTE	WASTE	1A1a	Public electricity and heat production	0101	8.3	Nielsen et al. (2010a)
				0102	14	DCE estimate based on plant specific data for four plants, 2009 data.
		1A2a-f	Industry	03	14	Assumed equal to district heating plants (DCE assumption).
		1A4a	Commercial/ Institutional	0201	14	Assumed equal to district heating plants (DCE assumption).
	INDU-STRIAL WASTE	1A2f	Industry - Other	031600	14	Assumed equal to waste. DCE assumption.
BIOMASS	WOOD	1A1a	Public electricity and heat production	0101	1.9	Nielsen et al. (2010a)
				0102	11	EEA (2013)
		1A2a-f	Industry	03	11	EEA (2013)
		1A4a	Commercial/ Institutional	0201	11	EEA (2013)
		1A4b i	Residential	0202	11	EEA (2013)
		1A4c i	Agriculture/ Forestry	0203	11	EEA (2013)
	STRAW	1A1a	Public electricity and heat production	0101	49	Nielsen et al. (2010a)
				0102	130	Nikolaisen et al. (1998)
		1A4b i	Residential	0202	130	Assumed equal to district heating plants. DCE assumption.
		1A4c i	Agriculture/ Forestry	0203	130	Assumed equal to district heating plants. DCE assumption.
	BIO OIL	1A1a	Public electricity and heat production	0101	0.1	DCE estimate based on Folkecenter for Vedvarende Energi (2000) and DEA (2012a).
		1A2a-f	Industry	03	0.1	DCE estimate based on Folkecenter for Vedvarende Energi (2000) and DEA (2012a).
		1A4b i	Residential	0202	0.1	DCE estimate based on Folkecenter for Vedvarende Energi (2000) and DEA (2012a).

Fuel type	Fuel	NFR	NFR_name	SNAP	SO ₂ emission factor, g/GJ	Reference
	BIOGAS	1A1a	Public electricity and heat production	0101, except engines	25	DCE estimate based on Christiansen (2003), Hjort-Gregersen (1999) and DEA (2012a).
				Engines	19.2	Nielsen & Illerup (2003)
				0102	25	DCE estimate based on Christiansen (2003), Hjort-Gregersen (1999) and DEA (2012a).
		1A2a-f	Industry	03, except engines	25	DCE estimate based on Christiansen (2003), Hjort-Gregersen (1999) and DEA (2012a).
				03, engines	19.2	Nielsen & Illerup (2003)
		1A4a	Commercial/ Institutional	0201, except engines	25	DCE estimate based on Christiansen (2003), Hjort-Gregersen (1999) and DEA (2012a).
				020105	19.2	Nielsen & Illerup (2003)
		1A4c i	Agriculture/ Forestry	0203, except engines	25	DCE estimate based on Christiansen (2003), Hjort-Gregersen (1999) and DEA (2012a).
				020304	19.2	Nielsen & Illerup (2003)
	BIO PROD GAS	1A1a	Public electricity and heat production	010105	1.9	Assumed equal to wood. DCE assumption.

Table 3A-4.8 NOx emission factors and references 2012.

Fuel type	Fuel	NFR	NFR_name	SNAP	NO _x emission factor, g/GJ	Reference	
SOLID	ANODE CARBON	1A2f	Industry - other	032000	132	Assumed equal to coal. DCE assumption.	
	COAL	1A1a	Public electricity and heat production	0101	32	DCE estimate based on Energinet.dk (2013) and EU ETS (2013)	
				0102	95	DEPA (2001a)	
		1A2a-f	Industry	03	132	DCE estimate based on plant specific data for 2011.	
				except cement production			
		1A2f	Industry, cement production	0316	176	DCE estimate based on plant specific data for 2012.	
		1A4b i	Residential	020200	95	DEPA (2001a)	
		1A4c i	Agriculture/ Forestry	0203	95	DEPA (2001a)	
	FLY ASH FOSSIL	1A1a	Public electricity and heat production	0101	30	Assumed equal to the emission factor for coal in 2010.	
	BROWN COAL BRI.	1A4b	Residential	0202	95	Assumed equal to coal. DCE assumption.	
	COKE OVEN COKE	1A2a-f	Industry	03	132	Assumed equal to coal. DCE assumption.	
1A4b i		Residential	020200	95	Assumed equal to coal. DCE assumption.		
LIQUID	PETROLEUM COKE	1A2a-f	Industry	03	132	Assumed equal to coal. DCE assumption.	
	RESIDUAL OIL	1A1a	Public electricity and heat production	0101	138	DCE estimate based on Energinet.dk (2009); Energinet.dk (2010); Energinet.dk (2011); EU ETS (2009-2011)	
				0102	142	DEPA (2001a)	
		1A1b	Petroleum refining	010306	200	IPCC (1997)	
		1A2a-f	Industry	03	129	DCE estimate based on plant specific data from two plants, 2011.	
		1A4a	Commercial/ Institutional	0201	142	DEPA (2001a)	
		1A4b	Residential	0202	142	DEPA (2001a)	
		1A4c i	Agriculture/ Forestry	0203	142	DEPA (2001a)	
		GAS OIL	1A1a	Public electricity and heat production	010101, 010102	114	DCE estimate based on plant specific data for 2011. Data from Energinet.dk (2011) and EU ETS (2011).
					010103	130	DEPA (2012b), DEPA (2003b) and DEPA (1990)
					0102		
					010104	350	DCE estimate based on Eltra & Elkraft System, (2001) and DEA (2012b)
					010105	942	Nielsen et al. (2010a)
			1A1b	Petroleum refining	010306	65	DEPA (1990)
			1A1c	Other energy industries	010504	350	(?)
			1A2a-f	Industry	03	130	DEPA (2012b), DEPA (2003b) and DEPA (1990)
				except engines			
	1A2a-f		Industry	Engines	942	Nielsen et al. (2010a)	
		1A4a	Commercial/ Institutional	0201	52	DEPA (2001a)	
				020105	942	Nielsen et al. (2010a)	
		1A4b i	Residential	0202	52	DEPA (2001a)	
				Engines	942	Nielsen et al. (2010a)	
		1A4c	Agriculture/Forestry	0203	52	DEPA (2001a)	
				Engines	942	Nielsen et al. (2010a)	
		KEROSENE	1A2f	Industry - other	03	50	EEA (2009)

Fuel type	Fuel	NFR	NFR_name	SNAP	NO _x emission factor, g/GJ	Reference
		1A4a	Commercial/ Institutional	0201	50	EEA (2009)
		1A4b i	Residential	0202	50	EEA (2009)
		1A4c i	Agriculture/ Forestry	0203	50	EEA (2009)
	LPG	1A1a	Public electricity and heat production	All	96	IPCC (1997)
		1A1b	Petroleum refining	0103	96	IPCC (1997)
		1A2a-f	Industry	03	96	IPCC (1997)
		1A4a	Commercial/ Institutional	0201	71	IPCC (1997)
		1A4b i	Residential	0202	47	IPCC (1997)
		1A4c i	Agriculture/ Forestry	0203	71	IPCC (1997)
	REFINERY GAS	1A1b	Petroleum refining	010304	170	DCE estimate based on plant specific data for a gas turbine in year 2000.
				010306	84	DCE estimate based on plant specific data for years 2011.
GAS	NATURAL GAS	1A1a	Public electricity and heat production	010101, 010102	55	DEPA (2003b)
				010103	42	Larsen (2009)
				010104	48	Nielsen et al. (2010a)
				010105	135	Nielsen et al. (2010a)
				0102	42	Larsen (2009)
		1A1c	Other energy industries	010504	250	Kristensen (2004)
		1A2a-f	Industry	03	42	Larsen (2009)
			Engines		135	Nielsen et al. (2010a)
			Turbines		48	Nielsen et al. (2010a)
				030700	87	DCE estimate based on plant specific data for 11 clay production plants, EU ETS (2011-2012); DEPA (2012)
		1A4a	Commercial/ Institutional	0201	30	Larsen (2009); DEPA (2001a)
			Engines		135	Nielsen et al. (2010a)
		1A4b i	Residential	0202	30	Larsen (2009); DEPA (2001a)
			Engines		135	Nielsen et al. (2010a)
		1A4c i	Agriculture/ Forestry	0203	30	Larsen (2009); DEPA (2001a)
			Engines		135	Nielsen et al. (2010a)
WASTE	WASTE	1A1a	Public electricity and heat production	0101	102	Nielsen et al. (2010a)
				0102	164	DCE estimate based on plant specific data for year 2000.
		1A2a-f	Industry	03	164	DCE estimate based on plant specific data for district heating plants in year 2000.
		1A4a	Commercial/ Institutional	0201	164	DCE estimate based on plant specific data for district heating plants in year 2000.
	INDUSTRIAL WASTE	1A2f	Industry - Other	031600	164	Assumed equal to waste. DCE assumption.
BIOMASS	WOOD	1A1a	Public electricity and heat production	0101	81	Nielsen et al. (2010a)
				0102	90	Serup et al. (1999)
		1A2a-f	Industry	All	90	Serup et al. (1999)
		1A4a	Commercial/ Institutional	0201	90	Serup et al. (1999)
		1A4b i	Residential	0202	73.9	DCE estimate based on DEA (2013a), DEPA (2013) and EEA (2013),
		1A4c i	Agriculture/ Forestry	0203	90	Serup et al. (1999)

Fuel type	Fuel	NFR	NFR_name	SNAP	NO _x emission factor, g/GJ	Reference
STRAW		1A1a	Public electricity and heat production	0101	125	Nielsen et al. (2010a)
				0102	90	Nikolaisen et al. (1998)
		1A4b i	Residential	0202	90	Assumed equal to district heating plants. DCE assumption.
		1A4c i	Agriculture/ Forestry	0203	90	Assumed equal to district heating plants. DCE assumption.
BIO OIL		1A1a	Public electricity and heat production	0101	249	Assumed equal to gas oil. DCE assumption. The emission factor for gas oil have been changed and the emission factor for biooil will also be changed in future inventories.
				010105	700	Assumed equal to gas oil. DCE assumption.
				0102	65	Assumed equal to gas oil. DCE assumption.
		1A2a-f	Industry	03	65	Assumed equal to gas oil. DCE assumption.
				Engines	700	Assumed equal to gas oil. DCE assumption.
		1A4b i	Residential	0202	65	Assumed equal to gas oil. DCE assumption.
BIOGAS		1A1a	Public electricity and heat production	0101, not engines	28	DEPA (2001a)
				Engines	202	Nielsen et al. (2010a)
				0102	28	DEPA (2001a)
		1A2a-f	Industry	03, not engines	28	DEPA (2001a)
				03, engines	202	Nielsen et al. (2010a)
				030902	59	DEPA (1990); DEPA (1995)
		1A4a	Commercial/ Institutional	0201, not engines	28	DEPA (2001a)
				020105	202	Nielsen et al. (2010a)
		1A4c i	Agriculture/ Forestry	0203, not engines	28	DEPA (2001a)
				020304	202	Nielsen et al. (2010a)
BIO PROD GAS		1A1a	Public electricity and heat production	010105	173	Nielsen et al. (2010a)
		1A4a	Commercial/ Institutional	020105	173	Nielsen et al. (2010a)

Table 3A-4.9 NMVOC emission factors and references 2012.

Fuel type	Fuel	NFR	NFR_name	SNAP	NMVOC, g/GJ	Reference
SOLID	ANODE CARBON COAL	1A2f	Industry - other	032000	10	Assumed equal to coal. DCE assumption.
		1A1a	Electricity and heat production	0101 0102	1.2	EEA (2009)
		1A2a-f	Industry	(all)	10	EEA (2009)
		1A4b i	Residential	020200	484	EEA (2009)
		1A4c i	Agriculture/ Forestry	020300	88.8	EEA (2009)
	BROWN COAL BRI.	1A4b i	Residential	020200	484	EEA (2009)
	COKE OVEN COKE	1A2a-f	Industry	03	10	EEA (2009)
		1A4b i	Residential	020200	484	EEA (2009)
	PETROLEUM COKE	1A2a-f	Industry	03	10	Assumed equal to coal. DCE assumption.
		1A1a	Public electricity and heat production	010101 010102 010103 010104	0.8	Nielsen et al. (2010)
LIQUID	RESIDUAL OIL	1A1b	Petroleum refining	010203	2.3	EEA (2009)
				010306	2.3	EEA (2009)
				03 except engines	0.8	Nielsen et al. (2010)
				Engines	10	EEA (2009)
				1A4a	5	EEA (2009)
		1A4b	Residential	0202	15	EEA (2009)
		1A4c i	Agriculture/ Forestry	0203	5	EEA (2009)
		1A1a	Public electricity and heat production	010101 010102 010103 010104 010105 0102	0.8	EEA (2009)
				010306	0.8	EEA (2009)
				010504	0.2	EEA (2009)
				03 boilers > 50 MW	5	EEA (2009)
				03 boilers < 50 MW	10	EEA (2009)
				Gas turbines	0.2	EEA (2009)
				Engines	37	EEA (2009)
				0201 except engines	5	EEA (2009)
				Engines	37	EEA (2009)
		1A4b i	Residential	0202	15	EEA (2009)
	GAS OIL	1A4c	Agriculture/ Forestry	0203	5	EEA (2009)
		1A2a-f	Industry	03	10	EEA (2009)
		1A4a	Commercial/ Institutional	0201	5	EEA (2009)
		1A4b i	Residential	0202	15	EEA (2009)
		1A4c i	Agriculture/ Forestry	0203	5	EEA (2009)
		KEROSENE		0201	5	EEA (2009)
				0202	15	EEA (2009)
				0203	5	EEA (2009)

Fuel type	Fuel	NFR	NFR_name	SNAP	NMVOC, g/GJ	Reference
GAS	LPG	1A1a	Public electricity and heat production	0101	0.8	EEA (2009)
				0102		
		1A1b	Petroleum refining	0103	0.8	EEA (2009)
		1A2a-f	Iron and steel	03	5	EEA (2009)
		1A4a	Commercial/ Institutional	0201	5	EEA (2009)
		1A4b i	Residential	0202	10	EEA (2009)
		1A4c i	Agriculture/ Forestry	0203	5	EEA (2009)
	REFINERY GAS	1A1b	Petroleum refining	0103	1.4	Assumed equal to natural gas fuelled gas turbines. DCE assumption.
	NATURAL GAS	1A1a	Public electricity and heat production	010101	2	Danish Gas Technology Centre (2001).
				010102		
				010103		
				010104	1.6	Nielsen et al. (2010)
				010105	92	Nielsen et al. (2010)
				0102	2	Danish Gas Technology Centre (2001).
		1A1c	Other energy industries	010504	1.6	Nielsen et al. (2010)
		1A2a-f	Industry	03 except engines and turbines	2	Danish Gas Technology Centre (2001).
				Turbines	1.6	Nielsen et al. (2010)
				Engines	92	Nielsen et al. (2010)
		1A4a	Commercial/ Institutional	0201 except engines	2	Danish Gas Technology Centre (2001).
				Engines	92	Nielsen et al. (2010)
		1A4b i	Residential	0202 except engines	4	Gruijthuijsen & Jensen (2000)
				Engines	92	Nielsen et al. (2010)
		1A4c i	Agriculture/ Forestry	0203 except engines	2	Danish Gas Technology Centre (2001).
				Engines	92	Nielsen et al. (2010)
WASTE	WASTE	1A1a	Public electricity and heat production	0101	0.56	Nielsen et al. (2010)
				0102	2	EEA (2009)
		1A2a-f	Industry	03	2	EEA (2009)
		1A4a	Commercial/ Institutional	0201	2	EEA (2009)
	INDISTRIAL WASTE	1A2f	Industry - Other	0316	2	EEA (2009)
BIOMASS	WOOD	1A1a	Public electricity and heat production	0101	5.1	Nielsen et al. (2010)
				0102	7.3	EEA (2009)
		1A2a-f	Industry	03	10	EEA (2009)
		1A4a	Commercial/ Institutional	0201	146	EEA (2009)
		1A4b i	Residential	0202	363.4	DCE estimate based on DEA (2013a), DEPA (2013) and EEA (2013),
		1A4c i	Agriculture/ Forestry	0203	146	EEA (2009)
	STRAW	1A1a	Public electricity and heat production	0101	0.78	Nielsen et al. (2010)
				0102	7.3	EEA (2009)
		1A4b i	Residential	0202	400	EEA (2009)
		1A4c i	Agriculture/ Forestry	0203	146	EEA (2009)
				020302	10	EEA (2009)

Fuel type	Fuel	NFR	NFR_name	SNAP	NMVOC, g/GJ	Reference
	BIO OIL	1A1a	Public electricity and heat production	010102	0.8	EEA (2009)
				010105	37	EEA (2009)
				0102	0.8	EEA (2009)
		1A2a-f	Industry	03	0.8	EEA (2009)
		1A4b i	Residential	0202	15	EEA (2009)
	BIOGAS	1A1a	Public electricity and heat production	0101	2	Assumed equal to natural gas. DCE assumption.
				010105	10	Nielsen et al. (2010)
				0102	2	Assumed equal to natural gas. DCE assumption.
		1A2a-f	Industry	03 except engines	2	Assumed equal to natural gas. DCE assumption.
				Engines	10	Nielsen et al. (2010)
				0201 except engines	2	Assumed equal to natural gas. DCE assumption.
		1A4a	Commercial/ Institutional	Engines	10	Nielsen et al. (2010)
				0203 except engines	2	Assumed equal to natural gas. DCE assumption.
				Engines	10	Nielsen et al. (2010)
	BIO PROD GAS	1A1a	Public electricity and heat production	010105	2	Nielsen et al. (2010)
		1A4a	Commercial/ Institutional	020105	2	Nielsen et al. (2010)

Table 3A-4.10 CO emission factors and references 2012.

Fuel type	Fuel	NFR	NFR_name	SNAP	CO emission factor g/GJ	Reference
SOLID	ANODE CARBON	1A2a-f	Industry	03	10	Assumed the same emission factor as for coal. DCE assumption.
	COAL	1A1a	Public electricity and heat production	0101 and 0102	10	Sander (2002)
		1A2a-f	Industry	03	10	Assumed equal to boilers in public electricity and heat production. DCE assumption.
		1A4b i	Residential	0202	2000	IPCC (1997)
		1A4c i	Agriculture/ Forestry	020300	931	EEA (2009)
	FLY ASH FOSSIL	1A1a	Public electricity and heat production	0101	10	Assumed equal to coal. DCE assumption.
	BROWN COAL BRI.	1A4b i	Residential	020200	2000	Assumed the same emission factor as for coal. DCE assumption.
	COKE OVEN COKE	1A2a-f	Industry	03	10	Assumed the same emission factor as for coal. DCE assumption.
		1A4b i	Residential	0202	2000	Assumed the same emission factor as for coal. DCE assumption.
LIQUID	PETROLEUM COKE	1A2a-f	Industry	03	61	Unknown – the emission factor will be updated according to EEA (2013)
	RESIDUAL OIL	1A1a	Electricity and heat production	0101	15	Sander (2002)
				010203	30	EEA (2007)
		1A1b	Petroleum refining	010306	30	EEA (2007)
		1A2a-f	Industry	03 except engines	2.8	Nielsen et al. (2010)
				Engines	100	EEA (2009)
		1A4a	Commercial/Institutional	0201	30	EEA (2007)
		1A4b	Residential	0202	30	EEA (2007)
		1A4c i	Agriculture/ Forestry	0203	30	EEA (2007)
		1A1a	Public electricity and heat production	0101 except engines	15	Sander (2002)
				Engines	130	Nielsen et al. (2010)
	GAS OIL			0102	30	EEA (2007)
		1A1b	Petroleum refining	010306	30	EEA (2007)
		1A2a-f	Industry	03 except gas turbines and engines	30	EEA (2007)
				Gas turbines	15	Sander (2002)
				Engines	130	Nielsen et al. (2010)
		1A4a	Commercial/ Institutional	0201 except engines	30	EEA (2007)
				Engines	130	Nielsen et al. (2010)
		1A4b i	Residential	0202	43	EEA (2007)
		1A4c	Agriculture/Forestry	0203	30	EEA (2007)
	KEROSENE	1A2a-f	Industry	03	20	EEA (2007)
		1A4a	Commercial/ Institutional	0201	20	EEA (2007)
		1A4b i	Residential	0202	20	EEA (2007)
		1A4c i	Agriculture/ Forestry	0203	20	EEA (2007)
	LPG	1A1a	Public electricity and heat production	0101 and 0102	25	EEA (2007)

Fuel type	Fuel	NFR	NFR_name	SNAP	CO emission factor g/GJ	Reference
GAS	REFINERY GAS	1A2a-f	Industry	03	25	EEA (2007)
		1A4a	Commercial/ Institutional	0201	25	EEA (2007)
		1A4b i	Residential	0202	25	EEA (2007)
		1A4c i	Agriculture/ Forestry	0203	25	EEA (2007)
		1A1b	Petroleum refining	0103	6.2	Assumed same emission factor as for natural gas fuelled gas turbines. DCE assumption.
		1A1a	Public electricity and heat production	010101 and 010102	15	Sander (2002)
				010103	28	DEPA (2001)
				010104	4.8	Nielsen et al. (2010)
				010105	58	Nielsen et al. (2010)
				0102	28	DEPA (2001)
		1A1c	Other energy industries	010504	4.8	Nielsen et al. (2010)
		1A2a-f	Industry	03 except gas turbines and engines	28	DEPA (2001)
				Gas turbines	4.8	Nielsen et al. (2010)
				Engines	58	Nielsen et al. (2010)
		1A4a	Commercial/ Institutional	0201 except engines	28	DEPA (2001)
				Engines	58	Nielsen et al. (2010)
		1A4b i	Residential	0202 except engines	20	Gruijthuijsen & Jensen (2000)
				Engines	58	Nielsen et al. (2010)
		1A4c i	Agriculture/ Forestry	0203 except engines	28	DEPA (2001)
				Engines	58	Nielsen et al. (2010)
WASTE	WASTE	1A1a	Public electricity and heat production	0101	3.9	Nielsen et al. (2010)
				0102	10	DCE calculation based on annual environmental reports for Danish plants year 2000.
		1A2a-f	Industry	03	10	Assumed equal to district heating plants. DCE assumption.
		1A4a	Commercial/ Institutional	0201	10	Assumed equal to district heating plants. DCE assumption.
	INDISTRIAL WASTE	1A2f	Industry - Other	0316	10	Assumed equal to waste, district heating plants. DCE assumption.
BIOMASS	WOOD	1A1a	Public electricity and heat production	0101	90	Nielsen et al. (2010)
				010203	240	DEPA (2001)
		1A2a-f	Industry	03	240	DEPA (2001)
		1A4a	Commercial/ Institutional	020100	240	DEPA (2001)
		1A4b i	Residential	0202	2676	DCE estimate based on DEA (2013a), DEPA (2013) and EEA (2013),
		1A4c i	Agriculture/ Forestry	020300	240	DEPA (2001)
	STRAW	1A1a	Public electricity and heat production	0101	67	Nielsen et al. (2010)
				0102	325	DEPA (2001); Nikolaisen et al (1998)
		1A4b i	Residential	0202	4000	EEA (2007); Jensen & Nielsen (1990) and Bjerrum (2002)
		1A4c i	Agriculture/ Forestry	0203	4000	EEA (2007); Jensen & Nielsen (1990) and Bjerrum (2002)
				020302	325	DEPA (2001); Nikolaisen et al (1998)

Fuel type	Fuel	NFR	NFR_name	SNAP	CO emission factor g/GJ	Reference
	BIO OIL	1A1a	Public electricity and heat production	0101 and 0102	15	Assumed same emission factor as for gas oil. DCE assumption.
		1A2a-f			15	Assumed same emission factor as for gas oil. DCE assumption.
		1A4b i	Residential	0202	100	Assumed same emission factor as for gas oil. DCE assumption.
BIOGAS		1A1a	Public electricity and heat production	0101 except engines	36	DEPA (2001)
				Engines	310	Nielsen et al. (2010)
				0102	36	DEPA (2001)
		1A2a-f	Industry	03 except engines	36	DEPA (2001)
				Engines	310	Nielsen et al. (2010)
		1A4a	Commercial/ Institutional	0201 except engines	36	DEPA (2001)
				Engines	310	Nielsen et al. (2010)
		1A4c i	Agriculture/ Forestry	0203 except engines	36	DEPA (2001)
				Engines	310	Nielsen et al. (2010)
BIO PROD GAS		1A1a	Public electricity and heat production	010105	586	Nielsen et al. (2010)
		1A4a	Commercial/ Institutional	020105	586	Nielsen et al. (2010)

Table 3A-4.11 SO₂ emission factors time-series, g per GJ for the years 1990 to 2012.

fuel_type	fuel_id	fuel_gr_abbr	nfr_id_EA	snap_id	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
SOLID	102A	COAL	1A1a	010100	506	571	454	386																			
SOLID	102A	COAL	1A1a	010101	506	571	454	386	343	312	420	215	263	193	64	47	45	61	42	41	37	40	26	14	10	9	11
SOLID	102A	COAL	1A1a	010102	506	571	454	386	343	312	420	215	263	193	64	47	45	61	42	41	37	40	26	14	10	9	11
SOLID	102A	COAL	1A1a	010103					343	312	420																
SOLID	102A	COAL	1A1a	010104					343	312	420	215													10	9	
LIQUID	110A	PETROLEUM COKE	1A1a	010102					787	787															605	605	
LIQUID	110A	PETROLEUM COKE	1A2e	030900					787														605		605	605	605
LIQUID	110A	PETROLEUM COKE	1A2f i	030700	787						787	787	787	787	787	605	605	605	605	605					605	605	605
LIQUID	110A	PETROLEUM COKE	1A2f i	030800	787	787	787	787		787					787	787	605	605	605	605	605						
LIQUID	110A	PETROLEUM COKE	1A2f i	031300	787	787	787			787	787		787	787									605		605	605	605
LIQUID	110A	PETROLEUM COKE	1A2f i	031600	787	787	787	787	787	787	787	787	787	787	787	605	605	605	605	605	605	605	605	605	605	605	605
LIQUID	110A	PETROLEUM COKE	1A4a i	020100	787	787	787	787	787	787	787	787	787	787	787	605	605	605		605	605	605	605		605	605	
LIQUID	110A	PETROLEUM COKE	1A4b i	020200	787	787	787	787	787	787	787	787	787	787	787	605	605	605		605		605	605		605	605	
LIQUID	110A	PETROLEUM COKE	1A4c i	020300	787	787	787	787		787	787	787	787	787	787	605	605	605									
LIQUID	203A	RESIDUAL OIL	1A1a	010100	446	470	490	475																			
LIQUID	203A	RESIDUAL OIL	1A1a	010101	446	470	490	475	543	351	408	344	369	369	403	315	290	334	349	283	308	206	100	100	100	100	100
LIQUID	203A	RESIDUAL OIL	1A1a	010102	446	470	490	475	543	351	408	344	369	369	403	315	290	334	349	283	308	206	100	100	100	100	100
LIQUID	203A	RESIDUAL OIL	1A1a	010103					543	351	408	344	369	369	403	315	290	334	349				100	100	100	100	100
LIQUID	203A	RESIDUAL OIL	1A1a	010104					543	351	408	344	369	369		315	290	334	349	283	308	206	100	100	100	100	100
LIQUID	203A	RESIDUAL OIL	1A1a	010105					543	351	408	344	369	369	403	315	290	334	349	283	308				100	100	
LIQUID	203A	RESIDUAL OIL	1A1a	010202					495	495	495	344	344	344							344				344	344	
LIQUID	203A	RESIDUAL OIL	1A1a	010203					495	495	495	344	344	344	344	344	344	344	344	344	344		344	344	344	344	344
LIQUID	203A	RESIDUAL OIL	1A1b	010306	798	798	798	798	537	537	537	537	537	537	537	537	537	537	537	537	537	537	537	537	537	537	537
LIQUID	203A	RESIDUAL OIL	1A2a	030400	495	495	495	495	495	495	495	344	344	344	344	344	344					344	344	344	344	344	344
LIQUID	203A	RESIDUAL OIL	1A2b	030500	495	495	495	495	495	495	495	344	344	344	344	344	344	344	344	344	344	344	344	344	344	344	344
LIQUID	203A	RESIDUAL OIL	1A2c	030600	495	495	495	495	495	495	495	344	344	344	344	344	344	344	344	344	344	344	344	344	344	344	344
LIQUID	203A	RESIDUAL OIL	1A2d	031100	495	495	495	495	495	495	495	344	344	344	344	344	344	344	344	344	344	344	344	344	344	344	344
LIQUID	203A	RESIDUAL OIL	1A2e	030900	495	495	495	495	495	495	495	344	344	344	344	344	344	344	344	344	344	344	344	344	344	344	344
LIQUID	203A	RESIDUAL OIL	1A2e	030902					495	495	495	344	344	344	344	344	344	344	344	344	344	344	344	344	344	344	344
LIQUID	203A	RESIDUAL OIL	1A2e	030903					495	495	495	344	344	344	344	344	344	344	344	344	344	344	344	344	344	344	344
LIQUID	203A	RESIDUAL OIL	1A2f i	030700	495	495	495	495	495	495	495	344		344	344	344	344	344	344		344	344					
LIQUID	203A	RESIDUAL OIL	1A2f i	030800	495	495	495	495	495	495	495	344	344	344	344	344	344	344	344	344	344	344	344	344	344	344	344
LIQUID	203A	RESIDUAL OIL	1A2f i	031000	495	495	495	495	495	495	495	344	344	344	344	344	344					344	344	344	344	344	344
LIQUID	203A	RESIDUAL OIL	1A2f i	031200	495	495	495	495	495	495	495		344	344	344	344	344				344	344	344	344	344	344	344
LIQUID	203A	RESIDUAL OIL	1A2f i	031300	495	495	495	495	495	495	495	344	344	344	344	344	344	344	344	344	344	344	344	344	344	344	344
LIQUID	203A	RESIDUAL OIL	1A2f i	031400	495	495	495	495	495	495	495	344	344	344	344	344	344	344	344	344	344	344	344	344	344	344	344
LIQUID	203A	RESIDUAL OIL	1A2f i	031500	495	495	495	495	495	495	495	344	344	344	344	344	344										
LIQUID	203A	RESIDUAL OIL	1A2f i	031600	495	495	495	495	495	495	495	344	344	344	344	344	344	344	344	344	344	344	344	344	344	344	344
LIQUID	203A	RESIDUAL OIL	1A2f i	032000	495	495	495	495	495	495	495	344	344	344	344	344	344	344	344	344	344	344	344	344	344	344	344

fuel_type	fuel_id	fuel_gr_abbr	nfr_id_EA	snap_id	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
LIQUID	203A	RESIDUAL OIL	1A4a i	020100	495	495	495	495	495	495	495	344	344	344	344	344	344	344	344	344	344	344	344	344	344	344	
LIQUID	203A	RESIDUAL OIL	1A4b i	020200	495	495	495	495	495	495	495	344	344	344	344	344	344	344	344	344	344	344	344	344	344	344	
LIQUID	203A	RESIDUAL OIL	1A4c i	020300	495	495	495	495	495	495	495	344	344	344	344	344	344	344	344	344	344		344	344	344	344	
LIQUID	204A	GAS OIL	1A1a	010101					94	23	23	23	23	23	23	23	23	23	23	23	23	23	23	23	23	23	
LIQUID	204A	GAS OIL	1A1a	010102					94	23	23	23	23	23	23	23	23	23	23	23	23	23	23	23	23	23	
LIQUID	204A	GAS OIL	1A1a	010104		94	94	94	94	23	23	23	23	23	23	23	23	23	23	23	23	23	23	23	23	23	
LIQUID	204A	GAS OIL	1A1a	010105					94	23	23	23	23	23	23	23	23	23	23	23	23	23	23	23	23	23	
LIQUID	204A	GAS OIL	1A1a	010202					94	23	23	23	23	23	23	23	23	23	23	23	23	23	23	23	23	23	
LIQUID	204A	GAS OIL	1A1a	010203					94	23	23	23	23	23	23	23	23	23	23	23	23	23	23	23	23	23	
LIQUID	204A	GAS OIL	1A1b	010306		94	94	94	94	23	23	23						23	23	23	23	23	23	23	23	23	
LIQUID	204A	GAS OIL	1A2a	030400	94	94	94	94	94	23	23	23	23	23	23	23	23	23	23	23	23						
LIQUID	204A	GAS OIL	1A2b	030500	94	94	94	94	94	23	23	23	23	23	23	23	23	23	23	23	23						
LIQUID	204A	GAS OIL	1A2c	030600	94	94	94	94	94	23	23	23	23	23	23	23	23	23	23	23	23						
LIQUID	204A	GAS OIL	1A2d	031100	94	94	94	94	94	23	23	23	23	23	23	23	23	23	23	23	23						
LIQUID	204A	GAS OIL	1A2e	030900	94	94	94	94	94	23	23	23	23	23	23	23	23	23	23	23	23						
LIQUID	204A	GAS OIL	1A2e	030902					94	23	23	23	23	23				23	23	23	23	23	23	23	23	23	
LIQUID	204A	GAS OIL	1A2e	030903					94	23					23	23	23	23	23	23	23	23	23	23	23	23	
LIQUID	204A	GAS OIL	1A2f i	030700	94	94	94	94	94	23	23	23	23	23	23	23	23	23	23	23	23						
LIQUID	204A	GAS OIL	1A2f i	030800	94	94	94	94	94	23	23	23	23	23	23	23	23	23	23	23	23						
LIQUID	204A	GAS OIL	1A2f i	031000	94	94	94	94	94	23	23	23	23	23	23	23	23	23	23	23	23						
LIQUID	204A	GAS OIL	1A2f i	031200	94	94	94	94	94	23	23	23	23	23	23	23	23	23	23	23	23						
LIQUID	204A	GAS OIL	1A2f i	031300	94	94	94	94	94	23	23	23	23	23	23	23	23	23	23	23	23						
LIQUID	204A	GAS OIL	1A2f i	031400	94	94	94	94	94	23	23	23	23	23	23	23	23	23	23	23	23						
LIQUID	204A	GAS OIL	1A2f i	031403					94	23		23	23	23	23	23	23										
LIQUID	204A	GAS OIL	1A2f i	032000	94	94	94	94	94	23	23	23	23	23	23	23	23	23	23	23	23	23	23	23	23	23	
LIQUID	204A	GAS OIL	1A4a i	020100	94	94	94	94	94	23	23	23	23	23	23	23	23	23	23	23	23	23	23	23	23	23	
LIQUID	204A	GAS OIL	1A4a i	020103					94		23	23	23	23	23	23	23	23	23	23	23	23	23	23	23	23	
LIQUID	204A	GAS OIL	1A4a i	020105					94	23	23	23	23	23	23	23	23	23	23	23	23	23	23	23	23	23	
LIQUID	204A	GAS OIL	1A4b i	020200	94	94	94	94	94	23	23	23	23	23	23	23	23	23	23	23	23	23	23	23	23	23	
LIQUID	204A	GAS OIL	1A4c i	020300	94	94	94	94	94	23	23	23	23	23	23	23	23	23	23	23	23						
LIQUID	225A	ORIMULSION	1A1a	010101						149	149	149	149	149		12	12	12	12								
WASTE	114A	WASTE	1A1a	010100	138	116	95	73																			
WASTE	114A	WASTE	1A1a	010102					52	30	29	28	26	25	24	24	24	24	19	14	8.3	8.3	8.3	8.3	8.3	8.3	
WASTE	114A	WASTE	1A1a	010103					52	30	29	28	26	25	24	24	24	24	19	14	8.3	8.3	8.3	8.3	8.3	8.3	
WASTE	114A	WASTE	1A1a	010104					52	30	29	28	26	25	24			24			8.3				8.3	8.3	
WASTE	114A	WASTE	1A1a	010200	138	131	124	117																	8.3	8.3	
WASTE	114A	WASTE	1A1a	010202						103	95																
WASTE	114A	WASTE	1A1a	010203					110	103	95	88	81	74	67	60	52	45	37	30	22	14	14	14	14	14	
WASTE	114A	WASTE	1A2a	030400					110	103																	
WASTE	114A	WASTE	1A2c	030600	138	131	124	117	110	103	95	88	81		67					30		14					
WASTE	114A	WASTE	1A2d	031100	138	131	124	117	110	103	95	88	81	74	67					30		14	14	14	14	14	
WASTE	114A	WASTE	1A2e	030900	138	131	124	117	110	103	95		81	74	67					30		14	14	14	14	14	
WASTE	114A	WASTE	1A2e	030902																30		14	14				
WASTE	114A	WASTE	1A2f i	030700	138	131	124	117	110	103			81														
WASTE	114A	WASTE	1A2f i	030800																30		14	14	14	14	14	
WASTE	114A	WASTE	1A2f i	031000	138	131	124	117	110	103	95	88	81	74	67							14					
WASTE	114A	WASTE	1A2f i	031200	138	131			110	103																	
WASTE	114A	WASTE	1A2f i	031300	138	131	124	117	110	103	95	88	81		67					30		14	14	14	14	14	

fuel_type	fuel_id	fuel_gr_abbr	nfr_id	EA	snap_id	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
WASTE	114A	WASTE	1A2f i		031400			124	117	110	103	95	88	81	74	67					30		14					
WASTE	114A	WASTE	1A2f i		032000	138	131	124	117	110	103	95	88	81	74	67					30		14					
WASTE	114A	WASTE	1A4a i		020100	138	131	124	117	110	103	95	88	81	74				45	37	30	22		14		14	14	14
WASTE	114A	WASTE	1A4a i		020103					110	103	95	88	81	74	67	60	52	45	37	30	22	14	14	14	14	14	14
WASTE	115A	INDUSTR. WASTE	1A2f i		031600											67	60	52	45	37	30	22	14	14	14	14	14	14

Table 3A-4.12 NO_x emission factors time-series, g per GJ for the years 1990 to 2012.

fuel_type	fuel_id	fuel_gr_abbr	nfr_id	EA	snap_id	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
SOLID	102A	COAL	1A1a		010100	342	384	294	289																			
SOLID	102A	COAL	1A1a		010101	342	384	294	289	267	239	250	200	177	152	129	122	130	144	131	127	109	98	59	39	30	30	32
SOLID	102A	COAL	1A1a		010102	342	384	294	289	267	239	250	200	177	152	129	122	130	144	131	127	109	98	59	39	30	30	32
SOLID	102A	COAL	1A1a		010103					267	239	250																
SOLID	102A	COAL	1A1a		010104					267	239	250	200													30	30	
SOLID	102A	COAL	1A1a		010203					200	200	200	200	200	200	95	95	95	95	95	95		95	95		95	95	95
																					516.		389.		262.			
SOLID	102A	COAL	1A2f i		031600	715	715	715	715	715	715	715	715	715	715	715	688	661	634	607	580	5	453	5	326	5	199	176
SOLID	102A	COAL	1A4a i		020100	200	200	200	200	200	200	200	200	200	200				95									
SOLID	102A	COAL	1A4b i		020200	200	200	200	200	200	200	200	200	200	200	95	95	95	95	95	95	95	95	95	95	95	95	95
SOLID	102A	COAL	1A4c i		020300	200	200	200	200	200	200	200	200	200	200	95	95	95	95	95	95	95	95	95	95	95	95	95
		BROWN COAL																										
SOLID	106A	BRI.	1A4b i		020200	200	200	200	200	200	200	200	200	200	200	95	95	95	95					95	95	95	95	95
		COKE OVEN																										
SOLID	107A	COKE	1A4b i		020200	200	200	200	200	200	200	200	200	200	200	95	95	95	95	95	95	95	95	95	95	95	95	95
		PETROLEUM																										
LIQUID	110A	COKE	1A1a		010102					267	239															138	138	
LIQUID	203A	RESIDUAL OIL	1A1a		010100	342	384	294	289																			
LIQUID	203A	RESIDUAL OIL	1A1a		010101	342	384	294	289	267	239	250	200	177	152	129	122	130	144	131	127	109	98	138	138	138	138	138
LIQUID	203A	RESIDUAL OIL	1A1a		010102	342	384	294	289	267	239	250	200	177	152	129	122	130	144	131	127	109	98	138	138	138	138	138
LIQUID	203A	RESIDUAL OIL	1A1a		010103					267	239	250	200	177	152	129	122	130	144	131	127	109	98	138	138	138	138	138
LIQUID	203A	RESIDUAL OIL	1A1a		010104					267	239	250	200	177	152		122	130	144	131	127	109	98	138	138	138	138	138
LIQUID	203A	RESIDUAL OIL	1A1a		010105					267	239	250	200	177	152	129	122	130	144	131	127	109				138	138	
LIQUID	203A	RESIDUAL OIL	1A4c i		020304									142	142	142	142	142	142							130	130	
LIQUID	204A	GAS OIL	1A1a		010101					249	249	249	249	249	249	249	249	249	249	232	215	198	182	165	148	131	114	114
LIQUID	204A	GAS OIL	1A1a		010102					249	249	249	249	249	249	249	249	249	249	232	215	198	182	165	148	131	114	114
LIQUID	204A	GAS OIL	1A1a		010105					1247	1196	1145	1094	1044	993	942	942	942	942	942	942	942	942	942	942	942	942	942
LIQUID	204A	GAS OIL	1A1b		010306		95	90	85	80	75	70	65						65	65	65	65	65	65	65	65	65	65
LIQUID	204A	GAS OIL	1A4a i		020105					1247	1196	1145	1094	1044	993	942	942	942	942	942	942	942	942	942	942	942	942	942
LIQUID	204A	GAS OIL	1A4c i		020304									1145	1094													
LIQUID	225A	ORIMULSION	1A1a		010101						138	139	138	138	138		88	86	86	86								
LIQUID	308A	REFINERY GAS	1A1b		010306	83	83	83	83	83	83	83	83	83	83	83	83	83	83	83	83	83	83	83	94	94	94	94
GAS	301A	NATURAL GAS	1A1a		010101					115	115		115				115	115	115	115	97	97	97	97	55	55	55	55
GAS	301A	NATURAL GAS	1A1a		010102					115	115	115		115	115	115	115	115	115	115	97	97	97	97	55	55	55	55
GAS	301A	NATURAL GAS	1A1a		010104	161	157	153	149	145	141	138	134	131	127	124	119	113	108	103	98	73	48	48	48	48	48	48
GAS	301A	NATURAL GAS	1A1a		010105	276	241	235	214	199	194	193	170	167	167	168	163	158	153	148	143	139	135	135	135	135	135	135
GAS	301A	NATURAL GAS	1A2c		030604	161	157	153	149	145	141	138	134	131	127	124	119	113	108	103	98	73	48	48	48	48	48	48
GAS	301A	NATURAL GAS	1A2c		030605								170	167	167	168	163	158	153	148	143							

fuel_type	fuel_id	fuel_gr_abbr	nfr_id	EA	snap_id	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
GAS	301A	NATURAL GAS	1A2d		031104						141	138	134	131	127	124	119	113	108	103	98	73	48	48	48	48	48	48
GAS	301A	NATURAL GAS	1A2e		030904					145	141	138	134	131	127	124	119	113	108	103	98	73	48	48	48	48	48	48
GAS	301A	NATURAL GAS	1A2e		030905					199	194	193	170	167	167	168	163	158	153	148	143	139	135	135	135	135	135	135
GAS	301A	NATURAL GAS	1A2f i		030705						194	193	170	167	167	168	163	158	153	148	143	139	135	135	135	135	135	135
GAS	301A	NATURAL GAS	1A2f i		031005							193	170	167	167	168	163	158	153	148	143	139	135	135	135	135	135	135
GAS	301A	NATURAL GAS	1A2f i		031305							193	170	167	167	168	163	158	153	148	143	139	135	135	135	135	135	135
GAS	301A	NATURAL GAS	1A2f i		031405						194	193	170	167	167	168	163	158	153	148	143	139	135	135	135	135	135	135
GAS	301A	NATURAL GAS	1A2f i		032004							138	134	131	127	124	119	113	108	103	98		48	48	48	48	48	48
GAS	301A	NATURAL GAS	1A2f i		032005	276	241	235	214	199	194	193	170	167	167	168	163	158	153	148	143	139	135	135	135	135	135	135
GAS	301A	NATURAL GAS	1A4a i		020104						141		134			124	119	113	108	103	98	73	48					
GAS	301A	NATURAL GAS	1A4a i		020105	276	241	235	214	199	194	193	170	167	167	168	163	158	153	148	143	139	135	135	135	135	135	135
GAS	301A	NATURAL GAS	1A4b i		020204		241	235	214	199	194	193	170	167	167	168	163	158	153	148	143	139	135	135	135	135	135	135
GAS	301A	NATURAL GAS	1A4c i		020304	276	241	235	214	199	194	193	170	167	167	168	163	158	153	148	143	139	135	135	135	135	135	135
WASTE	114A	WASTE	1A1a		010102					134	134	134	134	134	129	124	124	124	124	117	110	102	102	102	102	102	102	102
WASTE	114A	WASTE	1A1a		010103					134	134	134	134	134	129	124	124	124	124	117	110	102	102	102	102	102	102	102
WASTE	114A	WASTE	1A1a		010104					134	134	134	134	134	129	124			124			102				102	102	102
BIO- MASS	111A	WOOD	1A4b i		020200	63.4	63.4	63.4	63.4	63.4	63.4	63.4	63.4	63.4	63.4	63.4	65.4	66.2	66.5	66.8	67.6	68.5	68.8	70.1	71.5	72.3	73.2	73.9
BIO- MASS	111A	WOOD	1A4b i		020202																67.6	68.5	68.8	70.1	71.5	72.3	73.2	73.9
BIO- MASS	111A	WOOD	1A4b i		020204																			70.1	71.5	72.3		
BIO- MASS	215A	BIO OIL	1A1a		010200	100	95	90	85																			
BIO- MASS	215A	BIO OIL	1A1a		010203					80	75	70	65	65	65	65	65	65	65	65	65	65	65	65	65	65	65	65
BIO- MASS	309A	BIOGAS	1A1a		010105	711	696	681	665	650	635	616	597	578	559	540	484	427	371	315	259	202	202	202	202	202	202	202
BIO- MASS	309A	BIOGAS	1A2f i		032005												484	427	371	315	259	202	202	202	202	202	202	202
BIO- MASS	309A	BIOGAS	1A4a i		020105	711	696	681	665	650	635	616	597	578	559	540	484	427	371	315	259	202	202	202	202	202	202	202
BIO- MASS	309A	BIOGAS	1A4c i		020304					650	635	616	597	578	559	540	484	427	371	315	259	202	202	202	202	202	202	202

Table 3A-4.13 NMVOC emission factors time-series, g per GJ for the years 1990 to 2012.

fuel_type	fuel_id	fuel_gr_abbr	nfr_id_EA	snap_id	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
GAS	301A	NATURAL GAS	1A1a	010104	1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.5	1.5	1.6	1.6	1.6	1.6	1.6	1.6	1.6	1.6	1.6
GAS	301A	NATURAL GAS	1A1a	010105	60	69	81	127	140	142	138	124	122	122	121	114	108	101	95	88	90	92	92	92	92	92	92
GAS	301A	NATURAL GAS	1A1c	010504	1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.5	1.5	1.6	1.6	1.6	1.6	1.6	1.6	1.6	1.6	1.6
GAS	301A	NATURAL GAS	1A2c	030604	1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.5	1.5	1.6	1.6	1.6	1.6	1.6	1.6	1.6	1.6	1.6
GAS	301A	NATURAL GAS	1A2c	030605								124	122	122	121	114	108	101	95	88							
GAS	301A	NATURAL GAS	1A2d	031104						1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.5	1.5	1.6	1.6	1.6	1.6	1.6	1.6	1.6	1.6	1.6
GAS	301A	NATURAL GAS	1A2e	030904					1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.5	1.5	1.6	1.6	1.6	1.6	1.6	1.6	1.6	1.6	1.6
GAS	301A	NATURAL GAS	1A2e	030905					140	142	138	124	122	122	121	114	108	101	95	88	90	92	92	92	92	92	92
GAS	301A	NATURAL GAS	1A2f i	030705						142	138	124	122	122	121	114	108	101	95	88	90	92	92	92	92	92	92
GAS	301A	NATURAL GAS	1A2f i	031005							138	124	122	122	121	114	108	101	95	88	90	92	92	92	92	92	92
GAS	301A	NATURAL GAS	1A2f i	031305							138	124	122	122	121	114	108	101	95	88	90	92	92	92	92	92	92
GAS	301A	NATURAL GAS	1A2f i	031405						142	138	124	122	122	121	114	108	101	95	88	90	92	92	92	92	92	92
GAS	301A	NATURAL GAS	1A2f i	032004							1.4	1.4	1.4	1.4	1.4	1.4	1.5	1.5	1.6	1.6		1.6	1.6	1.6	1.6	1.6	1.6
GAS	301A	NATURAL GAS	1A2f i	032005	60	69	81	127	140	142	138	124	122	122	121	114	108	101	95	88	90	92	92	92	92	92	92
GAS	301A	NATURAL GAS	1A4a i	020104						1.4		1.4			1.4	1.4	1.5	1.5	1.6	1.6	1.6	1.6					
GAS	301A	NATURAL GAS	1A4a i	020105	60	69	81	127	140	142	138	124	122	122	121	114	108	101	95	88	90	92	92	92	92	92	92
GAS	301A	NATURAL GAS	1A4b i	020204		69	81	127	140	142	138	124	122	122	121	114	108	101	95	88	90	92	92	92	92	92	92
GAS	301A	NATURAL GAS	1A4c i	020304	60	69	81	127	140	142	138	124	122	122	121	114	108	101	95	88	90	92	92	92	92	92	92
WASTE	114A	WASTE	1A1a	010101									0.98	0.98		0.98	0.98	0.98					0.56	0.56			
WASTE	114A	WASTE	1A1a	010102					0.98	0.98	0.98	0.98	0.98	0.98	0.98	0.98	0.98	0.98	0.84	0.7	0.56	0.56	0.56	0.56	0.56	0.56	0.56
WASTE	114A	WASTE	1A1a	010103					0.98	0.98	0.98	0.98	0.98	0.98	0.98	0.98	0.98	0.98	0.84	0.7	0.56	0.56	0.56	0.56	0.56	0.56	0.56
WASTE	114A	WASTE	1A1a	010104					0.98	0.98	0.98	0.98	0.98	0.98	0.98	0.98	0.98	0.98	0.84	0.7	0.56	0.56	0.56	0.56	0.56	0.56	0.56
BIO-MASS	111A	WOOD	1A2a	030400	146	132	119	105	92									10	10								
BIO-MASS	111A	WOOD	1A2d	031100	146	132	119	105	92	78	64	51	37	24	10	10	10	10	10	10			10	10	10	10	10
BIO-MASS	111A	WOOD	1A2e	030900	146	132	119	105	92	78	64	51	37	24	10	10	10	10	10	10	10	10	10	10	10		
BIO-MASS	111A	WOOD	1A2e	030902									37	24	10	10								10	10	10	10

fuel_type	fuel_id	fuel_gr_abbr	nfr_id	EA	snap_id	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	
BIO-MASS	111A	WOOD	1A2f i	030700	146	132	119	105	92	78	64	51	37	24	10	10	10	10	10	10	10	10	10	10	10	10	10	10	
BIO-MASS	111A	WOOD	1A2f i	031000					92	78					24	10	10	10	10	10	10	10	10	10	10	10	10	10	
BIO-MASS	111A	WOOD	1A2f i	031200	146	132	119		92	78	64	51	37	24	10	10	10	10	10	10	10	10	10	10	10	10	10	10	
BIO-MASS	111A	WOOD	1A2f i	031300	146	132	119	105	92	78	64	51	37	24	10	10	10	10	10	10	10	10	10	10	10	10	10	10	
BIO-MASS	111A	WOOD	1A2f i	031400	146	132	119	105	92	78	64	51	37	24	10	10	10	10	10	10	10	10	10	10	10	10	10	10	
BIO-MASS	111A	WOOD	1A2f i	031403					92	78	64	51	37	24	10	10	10	10	10	10	10	10	10	10	10	10	10	10	
BIO-MASS	111A	WOOD	1A2f i	032000	146	132	119	105	92	78	64	51	37	24	10	10	10	10	10	10	10	10	10	10	10	10	10	10	
BIO-MASS	111A	WOOD	1A2f i	032003					92	78	64	51	37	24	10	10	10												
BIO-MASS	111A	WOOD	1A4b i	020200	584.9	584.9	584.9	584.9	584.9	584.9	584.9	584.9	584.9	584.9	584.9	584.9	522.2	498.0	493.2	487.3	464.8	438.7	448.3	428.7	400.6	386.7	372.7	363.4	
BIO-MASS	111A	WOOD	1A4b i	020202	464.8 438.7 448.3 428.7 400.6 386.7 372.7 363.4																								
BIO-MASS	111A	WOOD	1A4b i	020204	428.7 400.6 386.7																								
BIO-MASS	117A	STRAW	1A4b i	020200	925	872.5	820	767	715	663	610	558	505	453	400	400	400	400	400	400	400	400	400	400	400	400	400		
BIO-MASS	309A	BIOGAS	1A1a	010105	14	14	14	14	14	14	14	14	14	14	14	14	13	13	12	11	10	10	10	10	10	10	10		
BIO-MASS	309A	BIOGAS	1A2f i	032005													13	13	12	11	10	10	10	10	10	10	10	10	10
BIO-MASS	309A	BIOGAS	1A4a i	020105	14	14	14	14	14	14	14	14	14	14	14	14	13	13	12	11	10	10	10	10	10	10	10		
BIO-MASS	309A	BIOGAS	1A4c i	020304	14					14	14	14	14	14	14	14	13	13	12	11	10	10	10	10	10	10	10	10	

Table 3A-4.14 CO emission factors time-series, g per GJ for the years 1990 to 2012.

fuel_type	fuel	fuel_gr_abbr	nfr_id	snap	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
GAS	301A	NATURAL GAS	1A1a	010104	6.2	6.2	6.2	6.2	6.2	6.2	6.2	6.2	6.2	6.2	6.2	6.2	6.2	6.2	6.2	6.2	5.5	4.8	4.8	4.8	4.8	4.8	4.8
GAS	301A	NATURAL GAS	1A1a	010105	189	211	212	227	226	222	221	182	182	182	183	163	142	122	101	81	70	58	58	58	58	58	58
GAS	301A	NATURAL GAS	1A1c	010504	6.2	6.2	6.2	6.2	6.2	6.2	6.2	6.2	6.2	6.2	6.2	6.2	6.2	6.2	6.2	6.2	5.5	4.8	4.8	4.8	4.8	4.8	4.8
GAS	301A	NATURAL GAS	1A2c	030604	6.2	6.2	6.2	6.2	6.2	6.2	6.2	6.2	6.2	6.2	6.2	6.2	6.2	6.2	6.2	6.2	5.5	4.8	4.8	4.8	4.8	4.8	4.8
GAS	301A	NATURAL GAS	1A2c	030605								182	182	182	183	163	142	122	101	81							
GAS	301A	NATURAL GAS	1A2d	031104						6.2	6.2	6.2	6.2	6.2	6.2	6.2	6.2	6.2	6.2	6.2	5.5	4.8	4.8	4.8	4.8	4.8	4.8
GAS	301A	NATURAL GAS	1A2e	030904					6.2	6.2	6.2	6.2	6.2	6.2	6.2	6.2	6.2	6.2	6.2	6.2	5.5	4.8	4.8	4.8	4.8	4.8	4.8
GAS	301A	NATURAL GAS	1A2e	030905					226	222	221	182	182	182	183	163	142	122	101	81	70	58	58	58	58	58	58
GAS	301A	NATURAL GAS	1A2f i	030705						222	221	182	182	182	183	163	142	122	101	81	70	58	58	58	58	58	58
GAS	301A	NATURAL GAS	1A2f i	031005							221	182	182	182	183	163	142	122	101	81	70	58	58	58	58	58	58
GAS	301A	NATURAL GAS	1A2f i	031305							221	182	182	182	183	163	142	122	101	81	70	58	58	58	58	58	58
GAS	301A	NATURAL GAS	1A2f i	031405						222	221	182	182	182	183	163	142	122	101	81	70	58	58	58	58	58	58
GAS	301A	NATURAL GAS	1A2f i	032004							6.2	6.2	6.2	6.2	6.2	6.2	6.2	6.2	6.2	6.2		4.8	4.8	4.8	4.8	4.8	4.8
GAS	301A	NATURAL GAS	1A2f i	032005	189	211	212	227	226	222	221	182	182	182	183	163	142	122	101	81	70	58	58	58	58	58	58
GAS	301A	NATURAL GAS	1A4a i	020104						6.2		6.2			6.2	6.2	6.2	6.2	6.2	6.2	5.5	4.8					
GAS	301A	NATURAL GAS	1A4a i	020105	189	211	212	227	226	222	221	182	182	182	183	163	142	122	101	81	70	58	58	58	58	58	58
GAS	301A	NATURAL GAS	1A4b i	020204		211	212	227	226	222	221	182	182	182	183	163	142	122	101	81	70	58	58	58	58	58	58
GAS	301A	NATURAL GAS	1A4c i	020304	189	211	212	227	226	222	221	182	182	182	183	163	142	122	101	81	70	58	58	58	58	58	58
WASTE	114A	WASTE	1A1a	010101									7.4	7.4		8	8	8						3.9			
WASTE	114A	WASTE	1A1a	010102					7.4	7.4	7.4	7.4	7.4	7.4	8	8	8	8	6.6	5.3	3.9	3.9	3.9	3.9	3.9	3.9	3.9
WASTE	114A	WASTE	1A1a	010103					7.4	7.4	7.4	7.4	7.4	7.4	8	8	8	8	6.6	5.3	3.9	3.9	3.9	3.9	3.9	3.9	3.9
WASTE	114A	WASTE	1A1a	010104					7.4	7.4	7.4	7.4	7.4	7.4	8		8	8			3.9			3.9	3.9	3.9	3.9
WASTE	114A	WASTE	1A1a	010200	100	85	70	55																			
WASTE	114A	WASTE	1A1a	010202						25	10	10	10														
WASTE	114A	WASTE	1A1a	010203					40	25	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10
WASTE	114A	WASTE	1A2a	030400					40	25																	
WASTE	114A	WASTE	1A2c	030600	100	85	70	55	40	25	10	10	10		10					10		10					
WASTE	114A	WASTE	1A2d	031100	100	85	70	55	40	25	10	10	10	10	10					10		10	10	10	10	10	10
WASTE	114A	WASTE	1A2e	030900	100	85	70	55	40	25	10		10	10	10					10		10	10	10	10	10	10
WASTE	114A	WASTE	1A2f i	030700	100	85	70	55	40	25			10									10	10	10	10	10	10
WASTE	114A	WASTE	1A2f i	031000	100	85	70	55	40	25	10	10	10	10	10							10					
WASTE	114A	WASTE	1A2f i	031200	100	85			40	25																	
WASTE	114A	WASTE	1A2f i	031300	100	85	70	55	40	25	10	10	10		10					10		10	10	10	10	10	10
WASTE	114A	WASTE	1A2f i	031400			70	55	40	25	10	10	10	10	10					10		10					
WASTE	114A	WASTE	1A2f i	032000	100	85	70	55	40	25	10	10	10	10	10					10		10					
WASTE	114A	WASTE	1A4a i	020100	100	85	70	55	40	25	10	10	10	10				10	10	10	10		10		10	10	10
WASTE	114A	WASTE	1A4a i	020103					40	25	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10
BIOMASS	111A	WOOD	1A1a	010200	400	373	347	320																			
BIOMASS	111A	WOOD	1A1a	010203					293	267	240	240	240	240	240	240	240	240	240	240	240	240	240	240	240	240	240
BIOMASS	111A	WOOD	1A2a	030400	400	373	347	320	293									240	240	240							
BIOMASS	111A	WOOD	1A2d	031100	400	373	347	320	293	267	240	240	240	240	240	240	240	240	240	240			240	240	240	240	240
BIOMASS	111A	WOOD	1A2e	030900	400	373	347	320	293	267	240	240	240	240	240	240	240	240	240	240	240	240	240	240	240	240	240
BIOMASS	111A	WOOD	1A2f i	030700	400	373	347	320	293	267	240	240	240	240	240	240	240	240	240	240	240	240	240	240	240	240	240
BIOMASS	111A	WOOD	1A2f i	031000					293	267					240	240	240	240	240	240	240	240	240	240	240	240	240
BIOMASS	111A	WOOD	1A2f i	031200	400	373	347		293	267	240	240	240	240	240	240	240	240	240	240	240	240	240	240	240	240	240
BIOMASS	111A	WOOD	1A2f i	031300	400	373	347	320	293	267	240	240	240	240	240	240	240	240	240	240	240	240	240	240	240	240	240
BIOMASS	111A	WOOD	1A2f i	031400	400	373	347	320	293	267	240	240	240	240	240	240	240	240	240	240	240	240	240	240	240	240	240
BIOMASS	111A	WOOD	1A2f i	031403					293	267	240	240	240	240	240	240	240	240	240	240	240	240	240	240	240	240	240
BIOMASS	111A	WOOD	1A2f i	032000	400	373	347	320	293	267	240	240	240	240	240	240	240	240	240	240	240	240	240	240	240	240	240
BIOMASS	111A	WOOD	1A2f i	032003					293	267	240	240	240	240	240	240	240										
BIOMASS	111A	WOOD	1A4a i	020100	400	373	347	320	293	267	240	240	240	240	240	240	240	240	240	240	240	240	240	240	240	240	240
BIOMASS	111A	WOOD	1A4a i	020103					293	267	240	240															
BIOMASS	111A	WOOD	1A4b i	020200	4227.8	4227.8	4227.8	4227.8	4227.8	4227.8	4227.8	4227.8	4227.8	4227.8	4227.8	3810.0	3654.9	3633.0	3604.7	3461.1	3296.7	3349.5	3194.1	2982.1	2868.0	2755.7	2675.8

fuel_type	fuel	fuel_gr_abbr	nfr_id	snap	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
BIOMASS	111A	WOOD	1A4b i	020202																3461.1	3296.7	3349.5	3194.1	2982.1	2868.0	2755.7	2675.8
BIOMASS	111A	WOOD	1A4b i	020204																			3194.1	2982.1	2868.0		
BIOMASS	111A	WOOD	1A4c i	020300	400	373	347	320	293	267	240	240	240	240	240	240	240	240	240	240	240	240	240	240	240	240	240
BIOMASS	117A	STRAW	1A1a	010200	600	554	508	463																			
BIOMASS	117A	STRAW	1A1a	010203					417	371	325	325	325	325	325	325	325	325	325	325	325	325	325	325	325	325	325
BIOMASS	117A	STRAW	1A4b i	020200	8500	8500	8500	8500	8500	7500	6500	5500	4500	4000	4000	4000	4000	4000	4000	4000	4000	4000	4000	4000	4000	4000	4000
BIOMASS	117A	STRAW	1A4c i	020300	8500	8500	8500	8500	8500	7500	6500	5500	4500	4000	4000	4000	4000	4000	4000	4000	4000	4000	4000	4000	4000	4000	4000
BIOMASS	309A	BIOGAS	1A1a	010105	230	234	239	243	248	252	256	260	265	269	273	279	285	292	298	304	310	310	310	310	310	310	310
BIOMASS	309A	BIOGAS	1A2f i	032005												279	285	292	298	304	310	310	310	310	310	310	310
BIOMASS	309A	BIOGAS	1A4a i	020105	230	234	239	243	248	252	256	260	265	269	273	279	285	292	298	304	310	310	310	310	310	310	310
BIOMASS	309A	BIOGAS	1A4c i	020304					248	252	256	260	265	269	273	279	285	292	298	304	310	310	310	310	310	310	310

Table 3A-4.15 Technology specific emission factors for residential wood combustion.

Emission factor (g/GJ)	Pollutant	Emission factor, Reference	
		g/GJ	
Old stove	NOx	50	EEA (2013), conventional stoves
New stove	NOx	50	EEA (2013), conventional stoves
Stove according to resent Danish legislation (2008)	NOx	80	EEA (2013), energy efficient stoves
Eco labelled stove	NOx	95	EEA (2013), advanced / ecolabelled stoves
Other stove	NOx	50	EEA (2013), conventional stoves
Old boilers with hot water storage	NOx	80	EEA (2013), conventional boilers
Old boilers without hot water storage	NOx	80	EEA (2013), conventional boilers
New boilers with hot water storage	NOx	95	EEA (2013), advanced / ecolabelled stoves
New boilers without hot water storage	NOx	95	EEA (2013), advanced / ecolabelled stoves
Pellet boilers/stoves	NOx	80	EEA (2013), pellet stoves / boilers
Old stove	NM VOC	1200	Assumed two times conventional stoves. EEA (2013) for conventional stoves 20-3000 g/GJ.
New stove	NM VOC	600	EEA (2013), conventional stoves
Stove according to resent Danish legislation (2008)	NM VOC	350	EEA (2013), energy efficient stoves
Eco labelled stove	NM VOC	175	Assumed ½ modern stove. The EEA (2013) emission factor for ecolabelled stoves is 250 g/GJ, but this emission factor have not been revised since the 2009 version of the Guide-book.
Other stove	NM VOC	600	EEA (2013), conventional stoves
Old boilers with hot water storage	NM VOC	350	EEA (2013), conventional boilers
Old boilers without hot water storage	NM VOC	350	EEA (2013), conventional boilers
New boilers with hot water storage	NM VOC	175	Assumed equal to ecolabelled stoves.
New boilers without hot water storage	NM VOC	350	Assumed 2 times the emission from new boilers with heat accumulation tank
Pellet boilers/stoves	NM VOC	10	EEA (2013), pellet stoves / boilers
Old stove	CO	8000	Assumed two times conventional stoves. EEA (2013) for conventional stoves 1,000-10,000 g/GJ.
New stove	CO	4000	EEA (2013), conventional stoves
Stove according to resent Danish legislation (2008)	CO	4000	EEA (2013), energy efficient stoves
Eco labelled stove	CO	1117	Nordic Swan label limit. The EEA (2013) emission factor for advanced / ecolabelled stoves is 2000 g/GJ.
Other stove	CO	4000	EEA (2013), conventional stoves
Old boilers with hot water storage	CO	4000	EEA (2013), conventional boilers
Old boilers without hot water storage	CO	4000	EEA (2013), conventional boilers
New boilers with hot water storage	CO	1117	Assumed equal to ecolabelled stoves.
New boilers without hot water storage	CO	2234	Assumed 2 times the emission from new boilers with heat accumulation tank
Pellet boilers/stoves	CO	300	EEA (2013), pellet stoves and boilers

Annex 3A-5 Large point sources

Table 3A-5.1 Large point sources, fuel consumption in 2012 (1A1, 1A2 and 1A4).

nfr_id_EA	lps_id	lps_name	fuel_id	fuel_gr_abbr	Fuel_rate_TJ
1A1a	001	Amagervaerket	111A	WOOD AND SIMIL.	5332
1A1a	001	Amagervaerket	117A	STRAW	622
1A1a	001	Amagervaerket	203A	RESIDUAL OIL	85
1A1a	001	Amagervaerket	102A	COAL	9951
1A1a	001	Amagervaerket	203A	RESIDUAL OIL	335
1A1a	002	Svanemoellevaerket	301A	NATURAL GAS	2573
1A1a	002	Svanemoellevaerket	204A	GAS OIL	6
1A1a	003	H.C.Oerstedsvaerket	203A	RESIDUAL OIL	1
1A1a	003	H.C.Oerstedsvaerket	301A	NATURAL GAS	2898
1A1a	003	H.C.Oerstedsvaerket	301A	NATURAL GAS	1082
1A1a	003	H.C.Oerstedsvaerket	204A	GAS OIL	232
1A1a	004	Kyndbyvaerket	204A	GAS OIL	937
1A1a	005	Masnadoevaerket	204A	GAS OIL	40
1A1a	007	Stigsnaesvaerket	203A	RESIDUAL OIL	99
1A1a	007	Stigsnaesvaerket	203A	RESIDUAL OIL	61
1A1a	007	Stigsnaesvaerket	204A	GAS OIL	5
1A1a	008	Asnaesvaerket	102A	COAL	7300
1A1a	008	Asnaesvaerket	117A	STRAW	7
1A1a	008	Asnaesvaerket	111A	WOOD AND SIMIL.	5
1A1a	008	Asnaesvaerket	204A	GAS OIL	0
1A1a	008	Asnaesvaerket	303A	LPG	1
1A1a	008	Asnaesvaerket	203A	RESIDUAL OIL	28
1A1a	008	Asnaesvaerket	102A	COAL	257
1A1a	008	Asnaesvaerket	203A	RESIDUAL OIL	24
1A1a	008	Asnaesvaerket	204A	GAS OIL	31
1A1a	010	Avedoerevaerket	203A	RESIDUAL OIL	46
1A1a	010	Avedoerevaerket	117A	STRAW	1230
1A1a	010	Avedoerevaerket	111A	WOOD AND SIMIL.	13735
1A1a	010	Avedoerevaerket	103A	FLY ASH	61
1A1a	010	Avedoerevaerket	204A	GAS OIL	14
1A1a	010	Avedoerevaerket	203A	RESIDUAL OIL	286
1A1a	010	Avedoerevaerket	102A	COAL	8364
1A1a	010	Avedoerevaerket	301A	NATURAL GAS	3162
1A1a	011	Fynsvaerket	111A	WOOD AND SIMIL.	308
1A1a	011	Fynsvaerket	309A	BIOGAS	46
1A1a	011	Fynsvaerket	117A	STRAW	1959
1A1a	011	Fynsvaerket	102A	COAL	17425
1A1a	011	Fynsvaerket	203A	RESIDUAL OIL	147
1A1a	011	Fynsvaerket	204A	GAS OIL	0
1A1a	012	Studstrupvaerket	204A	GAS OIL	6
1A1a	012	Studstrupvaerket	111A	WOOD AND SIMIL.	27
1A1a	012	Studstrupvaerket	117A	STRAW	62
1A1a	012	Studstrupvaerket	203A	RESIDUAL OIL	150
1A1a	012	Studstrupvaerket	102A	COAL	16363
1A1a	012	Studstrupvaerket	117A	STRAW	777
1A1a	012	Studstrupvaerket	203A	RESIDUAL OIL	141
1A1a	012	Studstrupvaerket	102A	COAL	6675
1A1a	014	Nordjyllandsvaerket	303A	LPG	0
1A1a	014	Nordjyllandsvaerket	204A	GAS OIL	11
1A1a	014	Nordjyllandsvaerket	204A	GAS OIL	8
1A1a	014	Nordjyllandsvaerket	203A	RESIDUAL OIL	127
1A1a	014	Nordjyllandsvaerket	204A	GAS OIL	4
1A1a	014	Nordjyllandsvaerket	102A	COAL	14945
1A1a	018	Skaerbaekvaerket	204A	GAS OIL	12
1A1a	018	Skaerbaekvaerket	301A	NATURAL GAS	8783
1A1a	019	Enstedvaerket	102A	COAL	2358
1A1a	019	Enstedvaerket	203A	RESIDUAL OIL	24
1A1a	019	Enstedvaerket	111A	WOOD AND SIMIL.	169
1A1a	019	Enstedvaerket	117A	STRAW	1016
1A1a	019	Enstedvaerket	204A	GAS OIL	7
1A1a	019	Enstedvaerket	303A	LPG	0
1A1a	020	Esbjergvaerket	102A	COAL	17081
1A1a	020	Esbjergvaerket	203A	RESIDUAL OIL	122
1A1a	020	Esbjergvaerket	303A	LPG	0
1A1a	022	Oestkraft	102A	COAL	457
1A1a	022	Oestkraft	111A	WOOD AND SIMIL.	213
1A1a	022	Oestkraft	203A	RESIDUAL OIL	81
1A1a	022	Oestkraft	204A	GAS OIL	14
1A1a	022	Oestkraft	303A	LPG	0

nfr_id_EA	lps_id	lps_name	fuel_id	fuel_gr_abbr	Fuel_rate_TJ
1A1a	025	Horsens Kraftvarmevaerk	114A	MUNICIP. WASTES	1013
1A1a	025	Horsens Kraftvarmevaerk	111A	WOOD AND SIMIL.	13
1A1a	025	Horsens Kraftvarmevaerk	301A	NATURAL GAS	567
1A1a	025	Horsens Kraftvarmevaerk	303A	LPG	1
1A1a	026	Herningvaerket	111A	WOOD AND SIMIL.	4168
1A1a	026	Herningvaerket	203A	RESIDUAL OIL	0
1A1a	026	Herningvaerket	204A	GAS OIL	0
1A1a	026	Herningvaerket	215A	BIO OIL	5
1A1a	026	Herningvaerket	301A	NATURAL GAS	454
1A1a	026	Herningvaerket	303A	LPG	0
1A1a	027	I/S Vestforbraending	114A	MUNICIP. WASTES	5349
1A1a	027	I/S Vestforbraending	303A	LPG	0
1A1a	027	I/S Vestforbraending	204A	GAS OIL	1
1A1a	027	I/S Vestforbraending	301A	NATURAL GAS	17
1A1a	028	Amagerforbraending	111A	WOOD AND SIMIL.	541
1A1a	028	Amagerforbraending	114A	MUNICIP. WASTES	3901
1A1a	028	Amagerforbraending	303A	LPG	31
1A1a	029	Energi Randers Produktion	309A	BIOGAS	10
1A1a	029	Energi Randers Produktion	215A	BIO OIL	49
1A1a	029	Energi Randers Produktion	204A	GAS OIL	4
1A1a	029	Energi Randers Produktion	102A	COAL	2
1A1a	029	Energi Randers Produktion	111A	WOOD AND SIMIL.	2775
1A1a	030	Grenaa Kraftvarmevaerk	117A	STRAW	497
1A1a	030	Grenaa Kraftvarmevaerk	102A	COAL	528
1A1a	030	Grenaa Kraftvarmevaerk	111A	WOOD AND SIMIL.	261
1A1a	030	Grenaa Kraftvarmevaerk	203A	RESIDUAL OIL	30
1A1a	030	Grenaa Kraftvarmevaerk	204A	GAS OIL	3
1A1a	030	Grenaa Kraftvarmevaerk	303A	LPG	0
1A1a	031	Hilleroed Kraftvarmevaerk	301A	NATURAL GAS	1502
1A1a	032	Helsingoer Kraftvarmevaerk	301A	NATURAL GAS	1533
1A1a	032	Helsingoer Kraftvarmevaerk	301A	NATURAL GAS	1533
1A1a	036	Kolding Forbraendingsanlaeg TAS	203A	RESIDUAL OIL	0
1A1a	036	Kolding Forbraendingsanlaeg TAS	114A	MUNICIP. WASTES	530
1A1a	036	Kolding Forbraendingsanlaeg TAS	111A	WOOD AND SIMIL.	331
1A1a	036	Kolding Forbraendingsanlaeg TAS	111A	WOOD AND SIMIL.	348
1A1a	036	Kolding Forbraendingsanlaeg TAS	117A	STRAW	2
1A1a	036	Kolding Forbraendingsanlaeg TAS	117A	STRAW	1
1A1a	036	Kolding Forbraendingsanlaeg TAS	114A	MUNICIP. WASTES	504
1A1a	037	Maabjergvaerket	114A	MUNICIP. WASTES	1757
1A1a	037	Maabjergvaerket	301A	NATURAL GAS	55
1A1a	037	Maabjergvaerket	117A	STRAW	458
1A1a	037	Maabjergvaerket	111A	WOOD AND SIMIL.	419
1A1a	037	Maabjergvaerket	309A	BIOGAS	15
1A1a	038	Soenderborg Kraftvarmevaerk	301A	NATURAL GAS	86
1A1a	038	Soenderborg Kraftvarmevaerk	111A	WOOD AND SIMIL.	77
1A1a	038	Soenderborg Kraftvarmevaerk	114A	MUNICIP. WASTES	676
1A1a	038	Soenderborg Kraftvarmevaerk	117A	STRAW	0
1A1a	038	Soenderborg Kraftvarmevaerk	204A	GAS OIL	0
1A1a	039	I/S Kara Affaldsforbraendingsanlaeg	301A	NATURAL GAS	10
1A1a	039	I/S Kara Affaldsforbraendingsanlaeg	111A	WOOD AND SIMIL.	48
1A1a	039	I/S Kara Affaldsforbraendingsanlaeg	114A	MUNICIP. WASTES	2156
1A1a	040	Viborg Kraftvarme	204A	GAS OIL	1
1A1a	040	Viborg Kraftvarme	301A	NATURAL GAS	1921
1A1a	042	I/S Nordforbraending	111A	WOOD AND SIMIL.	446
1A1a	042	I/S Nordforbraending	301A	NATURAL GAS	9
1A1a	042	I/S Nordforbraending	114A	MUNICIP. WASTES	1068
1A1a	046	Affaldscenter aarhus - Forbraendsanlaegget	114A	MUNICIP. WASTES	2705
1A1a	047	I/S Reno Nord	114A	MUNICIP. WASTES	1860
1A1a	047	I/S Reno Nord	204A	GAS OIL	2
1A1a	047	I/S Reno Nord	111A	WOOD AND SIMIL.	30
1A1a	048	Silkeborg Kraftvarmevaerk	204A	GAS OIL	0
1A1a	048	Silkeborg Kraftvarmevaerk	301A	NATURAL GAS	1818
1A1a	050	AffaldPlus+, Naestved Forbraendingsanlaeg	114A	MUNICIP. WASTES	1169
1A1a	051	AVV Forbraendingsanlaeg	204A	GAS OIL	2
1A1a	051	AVV Forbraendingsanlaeg	111A	WOOD AND SIMIL.	23
1A1a	051	AVV Forbraendingsanlaeg	114A	MUNICIP. WASTES	737
1A1a	051	AVV Forbraendingsanlaeg	117A	STRAW	0
1A1a	051	AVV Forbraendingsanlaeg	203A	RESIDUAL OIL	7
1A1a	052	Affaldsforbraendingsanlaeg I/S REFA	114A	MUNICIP. WASTES	1233
1A1a	053	Svendborg Kraftvarmevaerk	114A	MUNICIP. WASTES	516
1A1a	053	Svendborg Kraftvarmevaerk	111A	WOOD AND SIMIL.	10
1A1a	053	Svendborg Kraftvarmevaerk	301A	NATURAL GAS	6
1A1a	053	Svendborg Kraftvarmevaerk	204A	GAS OIL	0

nfr_id_EA	lps_id	lps_name	fuel_id	fuel_gr_abbr	Fuel_rate_TJ
1A1a	053	Svendborg Kraftvarmevaerk	117A	STRAW	0
1A1a	054	Kommunekemi	204A	GAS OIL	22
1A1a	054	Kommunekemi	203A	RESIDUAL OIL	57
1A1a	054	Kommunekemi	114A	MUNICIP. WASTES	1906
1A1a	055	I/S Faelles Forbraending	114A	MUNICIP. WASTES	257
1A1a	058	I/S Reno Syd	204A	GAS OIL	2
1A1a	058	I/S Reno Syd	117A	STRAW	0
1A1a	058	I/S Reno Syd	114A	MUNICIP. WASTES	620
1A1a	058	I/S Reno Syd	111A	WOOD AND SIMIL.	91
1A1a	059	I/S Kraftvarmevaerk Thisted	117A	STRAW	1
1A1a	059	I/S Kraftvarmevaerk Thisted	114A	MUNICIP. WASTES	516
1A1a	059	I/S Kraftvarmevaerk Thisted	111A	WOOD AND SIMIL.	49
1A1a	061	Affaldplus+, Slagelse Forbr. and DONG Slagelse KVV	114A	MUNICIP. WASTES	566
1A1a	061	Affaldplus+, Slagelse Forbr. and DONG Slagelse KVV	117A	STRAW	474
1A1a	065	Haderslev Kraftvarmevaerk	301A	NATURAL GAS	11
1A1a	065	Haderslev Kraftvarmevaerk	111A	WOOD AND SIMIL.	27
1A1a	065	Haderslev Kraftvarmevaerk	114A	MUNICIP. WASTES	595
1A1a	065	Haderslev Kraftvarmevaerk	117A	STRAW	1
1A1a	066	Frederikshavn Affaldskraftvarmevaerk	204A	GAS OIL	1
1A1a	066	Frederikshavn Affaldskraftvarmevaerk	111A	WOOD AND SIMIL.	41
1A1a	066	Frederikshavn Affaldskraftvarmevaerk	114A	MUNICIP. WASTES	359
1A1a	068	Bofa I/S	114A	MUNICIP. WASTES	193
1A1a	069	DTU	301A	NATURAL GAS	1035
1A1a	070	AffaldPlus+, Naestved Kraftvarmevaerk	301A	NATURAL GAS	143
1A1a	072	Hjoerring Varmeforsyning	301A	NATURAL GAS	24
1A1a	072	Hjoerring Varmeforsyning	111A	WOOD AND SIMIL.	536
1A1a	085	L90 Affaldsforbraending	114A	MUNICIP. WASTES	2333
1A1a	085	L90 Affaldsforbraending	204A	GAS OIL	7
1A1a	086	Hammel Fjernvarmeselskab	114A	MUNICIP. WASTES	293
1A1a	086	Hammel Fjernvarmeselskab	215A	BIO OIL	14
1A1a	086	Hammel Fjernvarmeselskab	111A	WOOD AND SIMIL.	35
1A1a	087	Koege Kraftvarmevaerk	111A	WOOD AND SIMIL.	1098
1A1a	087	Koege Kraftvarmevaerk	203A	RESIDUAL OIL	1
1A1a	087	Koege Kraftvarmevaerk	204A	GAS OIL	0
1A1a	088	Skagen Forbraending	114A	MUNICIP. WASTES	114
1A1a	088	Skagen Forbraending	111A	WOOD AND SIMIL.	25
1A1a	090	Odense Kraftvarmevaerk	204A	GAS OIL	22
1A1a	090	Odense Kraftvarmevaerk	117A	STRAW	5
1A1a	090	Odense Kraftvarmevaerk	111A	WOOD AND SIMIL.	12
1A1a	090	Odense Kraftvarmevaerk	114A	MUNICIP. WASTES	2239
1A1a	091	Centralkommunernes Transmissionsselskab F.berg	204A	GAS OIL	170
1A1a	092	Frederikshavn Kraftvarmevaerk	301A	NATURAL GAS	61
1A1a	093	Fjernvarme Fyn, Centrum Varmecentral	301A	NATURAL GAS	28
1A1a	094	Special Waste System	114A	MUNICIP. WASTES	37
1A1a	095	Grenaa Forbraending	111A	WOOD AND SIMIL.	8
1A1a	095	Grenaa Forbraending	114A	MUNICIP. WASTES	247
1A1a	098	Vordingborg Kraftvarme	111A	WOOD AND SIMIL.	162
1A1a	098	Vordingborg Kraftvarme	117A	STRAW	472
1A1a	098	Vordingborg Kraftvarme	204A	GAS OIL	1
1A1b	009	Statoil Raffinaderi	303A	LPG	193
1A1b	009	Statoil Raffinaderi	308A	REFINERY GAS	8549
1A1b	009	Statoil Raffinaderi	204A	GAS OIL	1
1A1b	017	Shell Raffinaderi	308A	REFINERY GAS	4365
1A1b	017	Shell Raffinaderi	308A	REFINERY GAS	1734
1A1b	017	Shell Raffinaderi	203A	RESIDUAL OIL	808
1A1c	024	Nybro Gasbehandlingsanlaeg	204A	GAS OIL	0
1A1c	024	Nybro Gasbehandlingsanlaeg	301A	NATURAL GAS	0
1A1c	024	Nybro Gasbehandlingsanlaeg	301A	NATURAL GAS	175
1A4a i	049	Rensningsanlaegget Lynetten	114A	MUNICIP. WASTES	47
1A4a i	049	Rensningsanlaegget Lynetten	309A	BIOGAS	144
1A4a i	049	Rensningsanlaegget Lynetten	204A	GAS OIL	4
Total					225087

Table 3A-5.2 Large point sources, plant specific emissions (IPCC 1A1, 1A2 and 1A4)¹⁾.

nfr_id	lps_id	lps_name	SO ₂	NO _x	NMVOC	CO
1A1a	001	Amagervaerket	x	x		
1A1a	002	Svanemoellevaerket	x	x		x
1A1a	003	H.C.Oerstedsvaerket	x	x		
1A1a	004	Kyndbyvaerket	x	x		
1A1a	005	Masnedoevaerket	x	x		
1A1a	007	Stigsnaesvaerket	x	x		
1A1a	008	Asnaesvaerket	x	x		
1A1a	010	Avedoerevaerket	x	x		x
1A1a	011	Fynsvaerket	x	x		
1A1a	012	Studstrupvaerket	x	x		
1A1a	014	Nordjyllandsvaerket	x	x		
1A1a	018	Skaerbaekvaerket	x	x		
1A1a	019	Enstedvaerket	x	x		
1A1a	020	Esbjergvaerket	x	x		
1A1a	022	Oestkraft	x	x		
1A1a	025	Horsens Kraftvarmevaerk	x	x		
1A1a	026	Herningvaerket	x	x		x
1A1a	027	I/S Vestforbraending	x	x		
1A1a	028	Amagerforbraending	x	x		x
1A1a	029	Energi Randers Produktion	x	x		
1A1a	030	Grenaa Kraftvarmevaerk	x	x		x
1A1a	031	Hilleroed Kraftvarmevaerk		x		
1A1a	032	Helsingoer Kraftvarmevaerk		x		
1A1a	036	Kolding Forbraendingsanlaeg TAS	x	x	x	x
1A1a	037	Maabjergvaerket	x	x		x
1A1a	038	Soenderborg Kraftvarmevaerk	x	x	x	x
1A1a	039	I/S Kara Affaldsforbraendingsanlaeg	x	x		x
1A1a	040	Viborg Kraftvarme		x		
1A1a	042	I/S Nordforbraending	x	x		x
1A1a	046	Affaldscenter aarhus - Forbraendsanlaegget	x	x	x	
1A1a	047	I/S Reno Nord	x	x	x	x
1A1a	048	Silkeborg Kraftvarmevaerk		x		
1A1a	050	AffaldPlus+, Naestved Forbraendingsanlaeg	x	x	x	x
1A1a	051	AVV Forbraendingsanlaeg	x	x	x	x
1A1a	052	Affaldsforbraendingsanlaeg I/S REFA	x	x		x
1A1a	053	Svendborg Kraftvarmevaerk	x	x	x	x
1A1a	054	Kommunekemi	x	x	x	x
1A1a	055	I/S Faelles Forbraending	x	x		x
1A1a	058	I/S Reno Syd	x	x	x	x
1A1a	059	I/S Kraftvarmevaerk Thisted	x	x	x	x
1A1a	061	Affaldplus+, Slagelse Forbr. and DONG Slagelse KVV	x	x		x
1A1a	065	Haderslev Kraftvarmevaerk	x	x		x
1A1a	066	Frederikshavn Affaldskraftvarmevaerk	x	x	x	x
1A1a	068	Bofa I/S	x	x		x
1A1a	069	DTU		x		x
1A1a	070	AffaldPlus+, Naestved Kraftvarmevaerk		x		x
1A1a	072	Hjoerring Varmeforsyning		x		x
1A1a	085	L90 Affaldsforbraending	x	x		x
1A1a	086	Hammel Fjernvarmeselskab	x	x	x	x
1A1a	087	Koege Kraftvarmevaerk	x	x		
1A1a	088	Skagen Forbraending	x	x	x	
1A1a	090	Odense Kraftvarmevaerk	x	x		x
1A1a	091	Centralkommunernes Transmissionsselskab F_berg	x	x		
1A1a	092	Frederikshavn Kraftvarmevaerk	x	x		
1A1a	093	Fjernvarme Fyn, Centrum Varmecentral		x		
1A1a	094	Special Waste System	x	x	x	x
1A1b	009	Statoil Raffinaderi	x	x		
1A1b	017	Shell Raffinaderi	x	x		
1A1c	024	Nybro Gasbehandlingsanlaeg		x		
1A2a	033	DanSteel		x		
1A2c	081	Haldor Topsoee		x		
1A2c	084	Cheminova		x		
1A2c	097	Koppers	x	x	x	

nfr_id	lps_id	lps_name	SO ₂	NO _x	NMVOC	CO
1A2d	034	Dalum Papir		x		
1A2e	023	Danisco Grindsted		x		
1A2e	071	Maricogen		x		
1A2e	082	Nordic Sugar Nakskov	x	x		
1A2e	083	Nordic Sugar Nykoebing	x			
1A2e	089	AarhusKarlshamn Denmark A/S	x	x		
1A2f i	045	Aalborg Portland	x	x		x
1A2f i	076	Rockwool A/S Vamdrup	x	x		
1A2f i	077	Rockwool A/S Doense	x	x		
1A2f i	078	Ardagh Glass Holmegaard A/S		x		x
1A2f i	080	Saint-Gobain Isover A/S	x	x		
1A2f i	096	Faxe Kalk	x	x		
1A4a i	049	Rensningsanlaegget Lynetten	x	x		x
TOTAL			4157	14036	4	3412
Share of total emission from stationary combustion, %			52%	43%	0.02%	3%

1) Emissions of the pollutants marked with "x" are plant specific. Emission of other pollutants is estimated based on emission factors.
The total shown *in this table* only includes plant specific data.

Annex 3A-6 Adjustment of CO₂ emission

Table 3A-6.1 Adjustment of CO₂ emission (ref. DEA, 2013a).

		1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
Actual Degree Days	Degree days	2857	3284	3022	3434	3148	3297	3837	3236	3217	3056
Normal Degree Days	Degree days	3379	3380	3 359	3 365	3 366	3 378	3 395	3 389	3 375	3 339
Net electricity import	PJ	25.4	-7.1	13.5	4.3	-17.4	-2.9	-55.4	-26.1	-15.6	-8.3
Actual CO ₂ emission	1 000 000 tonnes	37.7	47.3	41.5	43.7	47.3	44.0	57.1	47.3	43.4	40.2
Adjusted CO ₂ emission	1 000 000 tonnes	43.9	45.8	44.4	44.8	43.5	43.3	44.1	41.3	39.8	38.3
Continued		2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Actual Degree Days	Degree days	2902	3279	3011	3150	3113	3068	2908	2807	2853	3061
Normal Degree Days	Degree days	3304	3289	3273	3271	3261	3224	3188	3136	3120	3127
Net electricity import	PJ	2.4	-2.1	-7.5	-30.8	-10.3	4.9	-25.0	-3.4	5.2	1.2
Actual CO ₂ emission	1 000 000 tonnes	36.3	37.9	37.5	42.3	36.2	32.5	40.2	34.6	31.8	31.0
Adjusted CO ₂ emission	1 000 000 tonnes	37.0	37.5	35.9	35.4	34.0	33.7	34.6	33.9	33.0	31.3
Continued		2010	2011	2012							
Actual Degree Days	Degree days	3742	2970	3234							
Normal Degree Days	Degree days	3171	3156	3166							
Net electricity import	PJ	-4.1	4.7	18.8							
Actual CO ₂ emission	1 000 000 tonnes	31.3	26.6	22.7							
Adjusted CO ₂ emission	1 000 000 tonnes	30.4	27.7	26.9							

Annex 3A-7 Uncertainty estimates

Table 3A-7.1 Uncertainty estimation, tier 1.

IPCC Source category	Gas	Base year emission	Year t emission	Activity data uncertainty	Emission factor uncertainty	Combined uncertainty	Combined uncertainty as % of total na- tional emissions in year t	Type A sensitivity	Type B sensitivity	Uncertainty i trend in national emissions introduced by emis- sion factor uncertainty	Uncertainty in trend in national emissions introduced by activity data uncer- tainty	Uncertainty introduced into the trend in total na- tional emissions	
		Input data Gg CO ₂ eq	Input data Gg CO ₂ eq	Input data %	Input data %	%	%	%	%	%	%	%	
Stationary Combustion, Coal	CO ₂	23834	10005	0.9	0.5	1.030	0.444	-0.118	0.263	-0.059	0.335	0.340	
Stationary Combustion, BKB	CO ₂	11	1	3.0	5	5.831	0.000	-0.000	0.000	-0.001	0.000	0.001	
Stationary Combustion, Coke	CO ₂	138	74	1.8	5	5.301	0.017	-0.000	0.002	-0.001	0.005	0.005	
Stationary Combustion, Fossil waste	CO ₂	573	1397	5.0	10	11.180	0.674	0.028	0.037	0.275	0.260	0.379	
Stationary Combustion, Petroleum coke	CO ₂	415	628	2.0	5	5.385	0.146	0.010	0.017	0.049	0.047	0.068	
Stationary Combustion, Residual oil	CO ₂	2496	571	1.2	2	2.334	0.057	-0.025	0.015	-0.050	0.026	0.056	
Stationary Combustion, Gas oil	CO ₂	4547	742	2.2	4	4.587	0.147	-0.053	0.020	-0.213	0.062	0.222	
Stationary Combustion, Kerosene	CO ₂	366	2	2.0	5	5.393	0.000	-0.006	0.000	-0.029	0.000	0.029	
Stationary Combustion, LPG	CO ₂	184	91	1.8	5	5.317	0.021	-0.001	0.002	-0.003	0.006	0.007	
Stationary Combustion, Refinery gas	CO ₂	816	906	1.0	2	2.236	0.087	0.011	0.024	0.021	0.034	0.040	
Stationary Combustion, Natural gas	CO ₂	4335	8293	1.0	0	1.118	0.400	0.148	0.218	0.059	0.322	0.327	
Stationary Combustion, SOLID, CH ₄	CH ₄	13	3	1.0	100	100.005	0.013	-0.000	0.000	-0.012	0.000	0.012	
Stationary Combustion, LIQUID, CH ₄	CH ₄	3	1	0.8	100	100.003	0.005	-0.000	0.000	-0.002	0.000	0.002	
Stationary Combustion, GAS, CH ₄	CH ₄	3	6	1.0	100	100.005	0.024	0.000	0.000	0.010	0.000	0.010	
Natural gas fuelled engines, GAS, CH ₄	CH ₄	5	120	1.0	2	2.236	0.012	0.003	0.003	0.006	0.004	0.008	
Stationary Combustion, WASTE, CH ₄	CH ₄	1	2	5.0	100	100.125	0.007	0.000	0.000	0.003	0.000	0.003	
Stationary Combustion, BIOMASS, CH ₄	CH ₄	102	135	15.7	100	101.233	0.589	0.002	0.004	0.191	0.079	0.207	
Biogas fuelled engines, BIOMASS, CH ₄	CH ₄	1	31	3.9	10	10.728	0.015	0.001	0.001	0.008	0.005	0.009	
Stationary Combustion, SOLID, N ₂ O	N ₂ O	68	27	1.0	400	400.001	0.474	-0.000	0.001	-0.148	0.001	0.148	
Stationary Combustion, LIQUID, N ₂ O	N ₂ O	44	11	0.8	1000	1000.000	0.478	-0.000	0.000	-0.415	0.000	0.415	
Stationary Combustion, GAS, N ₂ O	N ₂ O	16	24	1.0	750	750.001	0.791	0.000	0.001	0.285	0.001	0.285	
Stationary Combustion, WASTE, N ₂ O	N ₂ O	7	16	5.0	400	400.031	0.271	0.000	0.000	0.123	0.003	0.123	
Stationary Combustion, BIOMASS, N ₂ O	N ₂ O	38	91	15.3	400	400.292	1.570	0.002	0.002	0.713	0.052	0.715	
Total		38015.248	23177.199				4.831					1.271	
Total uncertainties				Overall uncertainty i the year (%):			2.198			Trend uncertainty (%):			1.127

IPCC Source category	Gas	Base year emission	Year t emission	Activity data uncertainty	Emission factor uncertainty	Combined uncertainty	Combined uncertainty as % of total na- tional emissions in year t	Type A sensitivity	Type B sensitivity	Uncertainty i trend in national emissions introduced by emis- sion factor uncertainty	Uncertainty in trend in national emissions introduced by activity data uncer- tainty	Uncertainty introduced into the trend in total na- tional emissions
		Input data Gg CO ₂	Input data Gg CO ₂	Input data %	Input data %	%	%	%	%	%	%	%
Stationary Combustion, Coal	CO ₂	23834	10005	0.9	0.5	1.030	0.454	-0.115	0.265	-0.057	0.338	0.342
Stationary Combustion, BKB	CO ₂	11	1	3.0	5	5.831	0.000	-0.000	0.000	-0.001	0.000	0.001
Stationary Combustion, Coke	CO ₂	138	74	1.8	5	5.301	0.017	-0.000	0.002	-0.001	0.005	0.005
Stationary Combustion, Fosssil waste	CO ₂	573	1397	5.0	10	11.180	0.688	0.028	0.037	0.279	0.262	0.383
Stationary Combustion, Petroleum coke	CO ₂	415	628	2.0	5	5.385	0.149	0.010	0.017	0.050	0.047	0.069
Stationary Combustion, Residual oil	CO ₂	2496	571	1.2	2	2.334	0.059	-0.025	0.015	-0.049	0.026	0.056
Stationary Combustion, Gas oil	CO ₂	4547	742	2.2	4	4.587	0.150	-0.053	0.020	-0.211	0.062	0.220
Stationary Combustion, Kerosene	CO ₂	366	2	2.0	5	5.393	0.000	-0.006	0.000	-0.029	0.000	0.029
Stationary Combustion, LPG	CO ₂	184	91	1.8	5	5.317	0.021	-0.001	0.002	-0.003	0.006	0.007
Stationary Combustion, Refinery gas	CO ₂	816	906	1.0	2	2.236	0.089	0.011	0.024	0.022	0.034	0.040
Stationary Combustion, Natural gas	CO ₂	4335	8293	1.0	0	1.118	0.408	0.150	0.220	0.060	0.325	0.330
Total	CO ₂	37715	22710				0.902					0.432
Total uncertainties				Overall uncertainty i the year (%):			0.950	Trend uncertainty (%):				0.657

IPCC Source category	Gas	Base year emission	Year t emission	Activity data uncertainty	Emission factor uncertainty	Combined uncertainty	Combined uncertainty as % of total na- tional emissions in year t	Type A sensitivity	Type B sensitivity	Uncertainty i trend in national emissions introduced by emis- sion factor uncertainty	Uncertainty in trend in national emissions introduced by activity data uncer- tainty	Uncertainty introduced into the trend in total na- tional emissions
		Input data Mg CH ₄	Input data Mg CH ₄	Input data %	Input data %	%	%	%	%	%	%	%
Stationary Combustion, SOLID, CH ₄	CH ₄	613	149	1.0	100	100.005	1.048	-0.211	0.024	-21.136	0.033	21.136
Stationary Combustion, LIQUID, CH ₄	CH ₄	131	53	0.8	100	100.003	0.372	-0.042	0.009	-4.183	0.010	4.183
Stationary Combustion, GAS, CH ₄	CH ₄	149	266	1.0	100	100.005	1.876	-0.014	0.044	-1.359	0.062	1.360
Natural gas fuelled engines, GAS, CH ₄	CH ₄	221	5735	1.0	2	2.236	0.904	0.859	0.945	1.719	1.336	2.177
Stationary Combustion, WASTE, CH ₄	CH ₄	37	73	5.0	100	100.125	0.518	-0.002	0.012	-0.209	0.086	0.226
Stationary Combustion, BIOMASS, CH ₄	CH ₄	4849	6421	15.7	100	101.233	45.809	-0.803	1.058	-80.286	23.555	83.670
Biogas fuelled engines, BIOMASS, CH ₄	CH ₄	70	1492	3.9	10	10.728	1.128	0.219	0.246	2.187	1.351	2.571
Total	CH₄	6071	14189				2105.583					7478.193
Total uncertainties				Overall uncertainty i the year (%):			45.887			Trend uncertainty (%):		86.477

IPCC Source category	Gas	Base year emission	Year t emission	Activity data uncertainty	Emission factor uncertainty	Combined uncertainty	Combined uncertainty as % of total na- tional emissions in year t	Type A sensitivity	Type B sensitivity	Uncertainty i trend in national emissions introduced by emis- sion factor uncertainty	Uncertainty in trend in national emissions introduced by activity data uncer- tainty	Uncertainty introduced into the trend in total na- tional emissions
		Input data Mg N ₂ O	Input data Mg N ₂ O	Input data %	Input data %	%	%	%	%	%	%	%
Stationary Combustion, SOLID, N ₂ O	N ₂ O	220	89	1.0	400	400.001	64.748	-0.226	0.159	-90.530	0.214	90.530
Stationary Combustion, LIQUID, N ₂ O	N ₂ O	142	36	0.8	1000	1000.000	65.321	-0.185	0.064	-184.872	0.070	184.872
Stationary Combustion, GAS, N ₂ O	N ₂ O	53	79	1.0	750	750.001	108.108	0.048	0.141	36.334	0.200	36.334
Stationary Combustion, WASTE, N ₂ O	N ₂ O	21	51	5.0	400	400.031	37.091	0.054	0.091	21.427	0.643	21.437
Stationary Combustion, BIOMASS, N ₂ O	N ₂ O	122	293	15.3	400	400.292	214.535	0.310	0.525	123.920	11.359	124.440
Total	N₂O	558	547				67547.291					59638.222
Total uncertainties				Overall uncertainty i the year (%):			259.899			Trend uncertainty (%):		244.209

SNAP	Gas	Base year emission	Year t emission	Activity data uncertainty	Emission factor uncertainty	Combined uncertainty	Combined uncertainty as % of total na- tional emissions in year t	Type A sensitivity	Type B sensitivity	Uncertainty i trend in national emissions introduced by emis- sion factor uncertainty	Uncertainty in trend in national emissions introduced by activity data uncer- tainty	Uncertainty introduced into the trend in total na- tional emissions
		Input data Mg SO ₂	Input data Mg SO ₂	Input data %	Input data %	%	%	%	%	%	%	%
01	SO ₂	127207	3066	2	10	10.198	3.878	-0.023	0.020	-0.227	0.056	0.234
02	SO ₂	11359	2092	2	20	20.100	5.217	0.010	0.013	0.194	0.038	0.197
03	SO ₂	16437	2903	2	10	10.198	3.673	0.013	0.019	0.132	0.053	0.142
Total	SO₂	155003	8061				55.745					0.114
Total uncertainties						Overall uncertainty i the year (%):		7.466		Trend uncertainty (%):		0.337

SNAP	Gas	Base year emission	Year t emission	Activity data uncertainty	Emission factor uncertainty	Combined uncertainty	Combined uncertainty as % of total na- tional emissions in year t	Type A sensitivity	Type B sensitivity	Uncertainty i trend in national emissions introduced by emis- sion factor uncertainty	Uncertainty in trend in national emissions introduced by activity data uncer- tainty	Uncertainty introduced into the trend in total na- tional emissions
		Input data Mg NO _x	Input data Mg NO _x	Input data %	Input data %	%	%	%	%	%	%	%
01	NO _x	94566	21846	2	20	20.100	13.574	-0.043	0.192	-0.869	0.542	1.024
02	NO _x	6949	5186	2	50	50.040	8.023	0.028	0.046	1.409	0.129	1.415
03	NO _x	12457	5316	2	20	20.100	3.303	0.016	0.047	0.312	0.132	0.339
Total	NO_x	113972	32349				259.527					3.167
Total uncertainties						Overall uncertainty i the year (%):		16.110		Trend uncertainty (%):		1.779

SNAP	Gas	Base year emission	Year t emission	Activity data uncertainty	Emission factor uncertainty	Combined uncertainty	Combined uncertainty as % of total na- tional emissions in year t	Type A sensitivity	Type B sensitivity	Uncertainty i trend in national emissions introduced by emis- sion factor uncertainty	Uncertainty in trend in national emissions introduced by activity data uncer- tainty	Uncertainty introduced into the trend in total na- tional emissions
		Input data Mg NMVOC	Input data Mg NMVOC	Input data %	Input data %	%	%	%	%	%	%	%
01	NMVOC	486	1509	2	50	50.040	4.796	0.069	0.107	3.430	0.304	3.443
02	NMVOC	12475	13966	2	50	50.040	44.383	-0.000	0.993	-0.012	2.809	2.809
03	NMVOC	1100	271	2	50	50.040	0.861	-0.068	0.019	-3.416	0.055	3.417
Total	NMVOC	14061	15746				1993.576					31.422
Total uncertainties						Overall uncertainty i the year (%):		44.649		Trend uncertainty (%):		5.606

SNAP	Gas	Base year emission	Year t emission	Activity data uncertainty	Emission factor uncertainty	Combined uncertainty	Combined uncertainty as % of total na- tional emissions in year t	Type A sensitivity	Type B sensitivity	Uncertainty i trend in national emissions introduced by emis- sion factor uncertainty	Uncertainty in trend in national emissions introduced by activity data uncer- tainty	Uncertainty introduced into the trend in total na- tional emissions
		Input data Mg CO	Input data Mg CO	Input data %	Input data %	%	%	%	%	%	%	%
01	CO	8105	10974	2	20	20.100	1.767	0.025	0.083	0.500	0.236	0.553
02	CO	118884	109626	2	50	50.040	43.955	-0.023	0.832	-1.142	2.354	2.617
03	CO	4706	4201	2	20	20.100	0.677	-0.002	0.032	-0.039	0.090	0.098
Total	CO	131695	124802				1935.658					7.162
Total uncertainties						Overall uncertainty i the year (%):		43.996		Trend uncertainty (%):		2.676

Table 3A-7.2 Uncertainty estimation for GHG 2012, tier 2.

	Activity			Emission Factor			Emissions			
Parameter	Below 2.5%	Above 97.5%	Difference	Below 2.5%	Above 97.5%	Difference	Median	Below 2.5%	Above 97.5%	Difference
all							23285.44	22957.32	23854.53	897.21
all							244.86	63.07	792.41	729.34
all							22711.13	22502.56	22937.38	434.81
all							300.23	217.24	506.12	288.88
Stationary Combustion, BIOMASS, N ₂ O	93.05	125.47	32.42	0.06	5.77	5.70	84.51	6.89	624.10	617.21
Stationary Combustion, Fossil waste, CO ₂	16.12	17.75	1.64	74.82	90.99	16.17	1396.99	1251.03	1559.84	308.81
Stationary Combustion, BIOMASS, CH ₄	89.70	122.34	32.64	0.52	3.23	2.71	135.78	52.79	342.10	289.31
Stationary Combustion, GAS, N ₂ O	143.85	146.66	2.81	0.01	1.89	1.88	21.91	0.95	272.80	271.86
Stationary Combustion, Coal, CO ₂	105.23	107.08	1.86	93.81	94.71	0.90	10004.67	9904.99	10104.74	199.75
Stationary Combustion, SOLID, N ₂ O	105.88	107.87	1.99	0.02	1.76	1.74	26.24	2.37	188.02	185.65
Stationary Combustion, Natural gas, CO ₂	143.75	146.77	3.02	56.88	57.32	0.45	8293.45	8202.23	8386.32	184.10
Stationary Combustion, LIQUID, N ₂ O	40.80	41.43	0.63	0.01	3.97	3.96	9.90	0.32	163.14	162.82
Stationary Combustion, WASTE, N ₂ O	35.69	39.37	3.68	0.03	2.79	2.75	14.99	1.29	104.24	102.95
Stationary Combustion, Refinery gas, CO ₂	15.48	15.79	0.31	55.19	60.81	5.62	905.69	862.24	951.47	89.23
Stationary Combustion, Gas oil, CO ₂	9.81	10.24	0.44	71.18	76.98	5.79	741.68	709.68	775.56	65.88
Stationary Combustion, Petroleum coke, CO ₂	6.58	6.85	0.26	89.07	98.13	9.06	627.64	595.97	661.37	65.40
Stationary Combustion, Residual oil, CO ₂	7.20	7.37	0.17	76.88	79.94	3.06	570.90	557.69	584.13	26.44
Stationary Combustion, GAS, CH ₄	143.81	146.70	2.89	0.02	0.10	0.08	5.67	2.25	14.52	12.27
Stationary Combustion, LPG, CO ₂	1.40	1.46	0.05	60.64	66.84	6.21	91.08	86.45	95.94	9.50
Stationary Combustion, Coke, CO ₂	0.10	0.10	0.00	713.63	786.54	72.91	73.80	70.10	77.73	7.63
Stationary Combustion, SOLID, CH ₄	105.88	107.87	1.99	0.01	0.07	0.06	3.11	1.23	7.83	6.60
Biogas fuelled engines, BIOMASS, CH ₄	3.31	3.57	0.26	8.25	10.03	1.78	31.36	28.24	34.82	6.57
Natural gas fuelled engines, GAS, CH ₄	11.80	12.04	0.24	9.90	10.30	0.40	120.43	117.78	123.12	5.33
Stationary Combustion, WASTE, CH ₄	35.70	39.37	3.67	0.02	0.10	0.09	1.54	0.60	3.87	3.28
Stationary Combustion, LIQUID, CH ₄	40.79	41.41	0.61	0.01	0.07	0.06	1.11	0.44	2.78	2.35
Stationary Combustion, Kerosene, CO ₂	0.02	0.02	0.00	68.43	75.55	7.12	1.69	1.60	1.78	0.18
Stationary Combustion, BKB, CO ₂	0.01	0.02	0.00	90.10	99.42	9.32	1.41	1.33	1.49	0.16

Annex 3A-8 Emission inventory 2012 based on SNAP sectors

Table 3A-8.1 Emission inventory 2012 based on SNAP sectors.

SNAP	SO ₂ [Mg]	NO _x [Mg]	NMVOC [Mg]	CH ₄ [Mg]	CO [Mg]	CO ₂ [Gg] ¹⁾	N ₂ O [Mg]
Total:	10964.09	37665.07	16017.48	14597.32	129003	41472.48	622.75
1	3065.62	21846.17	1509.22	6609.77	10973.86	25623.37	283.32
101	1908.04	11350.17	1300.29	6052.78	6077.9	20038.57	181.79
10100	0	0	0	0	0	0	0
10101	1131.63	3897.19	171.02	112.45	1848.68	11093.11	89.59
10102	530.78	2984.69	74.07	41.41	1095.7	3955.57	36.69
10103	117.78	1060.02	7.82	7.68	159.14	1244.5	12.95
10104	77.61	1476.07	96.99	71.54	1455.03	2923.21	31.96
10105	50.25	1932.21	950.39	5819.7	1519.35	822.17	10.61
102	873.68	2644.26	144.99	494.59	4649.73	3164.48	72.78
10200	0	0	0	0	0	0	0
10201	0	0	0	0	0	0	0
10202	4.41	56.27	2.21	0.29	34.37	74.52	0.19
10203	869.27	2587.98	142.78	494.3	4615.36	3089.97	72.59
10204	0	0	0	0	0	0	0
10205	0	0	0	0	0	0	0
103	273.13	1604.15	23.91	19.86	126.13	983.65	3.73
10300	0	0	0	0	0	0	0
10301	0	0	0	0	0	0	0
10302	0	0	0	0	0	0	0
10303	0	0	0	0	0	0	0
10304	5.24	270	2.43	2.95	10.75	100.47	1.73
10305	0	0	0	0	0	0	0
10306	267.89	1334.15	21.48	16.92	115.38	883.18	1.99
104	0	0	0	0	0	0	0
10400	0	0	0	0	0	0	0
10401	0	0	0	0	0	0	0
10402	0	0	0	0	0	0	0
10403	0	0	0	0	0	0	0
10404	0	0	0	0	0	0	0
10405	0	0	0	0	0	0	0
10406	0	0	0	0	0	0	0
10407	0	0	0	0	0	0	0
105	10.76	6247.59	40.03	42.53	120.1	1436.67	25.02
10500	0	0	0	0	0	0	0
10501	0	0	0	0	0	0	0
10502	0	0	0	0	0	0	0
10503	0.08	36.26	0.28	0.3	0.84	9.97	0.17
10504	10.68	6211.34	39.75	42.24	119.26	1426.7	24.85
10505	0	0	0	0	0	0	0
10506	0	0	0	0	0	0	0
2	2092.06	5186.26	13966.27	7170.22	109626.4	7382.39	188.07
201	146.75	677.6	201.33	498.12	714.86	856.8	23.94
20100	133.98	495.36	156.65	45.6	509.15	762.16	22.45
20101	0	0	0	0	0	0	0
20102	0	0	0	0	0	0	0
20103	1.47	8.87	0.66	2.21	2.57	22.63	0.32
20104	0	0	0	0	0	0	0
20105	11.3	173.36	44.03	450.31	203.13	72.01	1.17
20106	0	0	0	0	0	0	0
202	922.22	3927.72	13288.78	5560.88	99781.02	5979.45	148.94
20200	921.76	3831.03	13224.92	5236.25	99728.8	5929.87	148.51
20201	0	0	0	0	0	0	0
20202	0.12	5.98	2.04	1.41	13.25	11.26	0.03
20203	0	0	0	0	0	0	0
20204	0.34	90.72	61.82	323.22	38.97	38.32	0.39
20205	0	0	0	0	0	0	0

SNAP	SO ₂ [Mg]	NO _x [Mg]	NMVOC [Mg]	CH ₄ [Mg]	CO [Mg]	CO ₂ [Gg] ¹⁾	N ₂ O [Mg]
203	1023.1	580.94	476.16	1111.22	9130.54	546.15	15.19
20300	1005.38	386.41	420.19	599.26	8908.43	462.76	13.83
20301	0	0	0	0	0	0	0
20302	6.16	3.9	1.16	0.85	8.89	3.88	0.12
20303	0	0	0	0	0	0	0
20304	11.56	190.63	54.8	511.1	213.22	79.51	1.25
20305	0	0	0	0	0	0	0
3	5806.41	10632.64	541.99	817.33	8402.69	8466.72	151.36
301	0	0.16	0.01	0.01	0.11	0.22	0
30100	0	0	0	0	0	0	0
30101	0	0	0	0	0	0	0
30102	0	0	0	0	0	0	0
30103	0	0	0	0	0	0	0
30104	0	0	0	0	0	0	0
30105	0	0	0	0	0	0	0
30106	0	0.16	0.01	0.01	0.11	0.22	0
302	0	0	0	0	0	0	0
30200	0	0	0	0	0	0	0
30203	0	0	0	0	0	0	0
30204	0	0	0	0	0	0	0
30205	0	0	0	0	0	0	0
304	1.39	53.54	1.94	1.35	26.78	54.83	0.11
30400	0.98	0.55	0.01	0.01	0.06	0.35	0.01
30401	0	0	0	0	0	0	0
30402	0.41	52.99	1.92	1.34	26.72	54.48	0.1
30403	0	0	0	0	0	0	0
30404	0	0	0	0	0	0	0
30405	0	0	0	0	0	0	0
30406	0	0	0	0	0	0	0
305	2.84	3.37	0.12	0.09	1.56	3.77	0.05
30500	2.84	3.37	0.12	0.09	1.56	3.77	0.05
30501	0	0	0	0	0	0	0
30502	0	0	0	0	0	0	0
30503	0	0	0	0	0	0	0
30504	0	0	0	0	0	0	0
30505	0	0	0	0	0	0	0
30506	0	0	0	0	0	0	0
306	12.63	270.22	8.78	6.25	97.34	249.83	2.13
30600	1.15	112.67	5.37	3.75	75.02	152.83	0.27
30601	0	0	0	0	0	0	0
30602	0.28	4.75	1.26	0.88	17.66	35.99	0.06
30603	11.2	36.6	0.89	0.28	0.89	16.28	1.01
30604	0	116.2	1.25	1.33	3.77	44.73	0.78
30605	0	0	0	0	0	0	0
30606	0	0	0	0	0	0	0
307	280.57	453.62	16.1	17.25	118.66	286.84	1.37
30700	280.49	392.55	12.52	9.43	115.28	262.45	1.01
30701	0	0	0	0	0	0	0
30702	0	0	0	0	0	0	0
30703	0.07	59.6	2.58	2.57	2.75	23.77	0.36
30704	0	0	0	0	0	0	0
30705	0.01	1.47	1	5.25	0.63	0.62	0.01
30706	0	0	0	0	0	0	0
308	92.47	144.15	10.97	18.76	223.59	154.52	4.54
30800	92.47	144.15	10.97	18.76	223.59	154.52	4.54
30801	0	0	0	0	0	0	0
30802	0	0	0	0	0	0	0
30803	0	0	0	0	0	0	0
30804	0	0	0	0	0	0	0
30805	0	0	0	0	0	0	0
30806	0	0	0	0	0	0	0
309	1365.35	1348.75	59.88	136.62	415.65	1240.61	22.49
30900	72.08	547.45	24.5	21.22	336.42	711.37	2.6

SNAP	SO ₂ [Mg]	NO _x [Mg]	NM VOC [Mg]	CH ₄ [Mg]	CO [Mg]	CO ₂ [Gg] ¹⁾	N ₂ O [Mg]
30901	0	0	0	0	0	0	0
30902	960.1	485.09	10.09	15.95	33.18	282.7	11.69
30903	332.53	228.52	5.91	6.6	29	163.17	6.81
30904	0.55	62.24	2.04	2.16	6.11	72.62	1.27
30905	0.09	25.45	17.35	90.69	10.94	10.75	0.11
30906	0	0	0	0	0	0	0
310	1.44	19.13	1.62	4.57	13.09	24.36	0.09
31000	1.43	18.03	0.87	0.65	12.62	23.9	0.08
31001	0	0	0	0	0	0	0
31002	0	0	0	0	0	0	0
31003	0	0	0	0	0	0	0
31004	0	0	0	0	0	0	0
31005	0	1.1	0.75	3.92	0.47	0.46	0
31006	0	0	0	0	0	0	0
311	47.36	373.69	36.23	52	820.64	473.52	13.85
31100	33.03	242.51	22.51	31.88	510.25	300.04	8.2
31101	0	0	0	0	0	0	0
31102	14.1	105.7	12.87	19.22	307.84	143.2	5.11
31103	0	0	0	0	0	0	0
31104	0.23	25.48	0.85	0.9	2.55	30.28	0.53
31105	0	0	0	0	0	0	0
31106	0	0	0	0	0	0	0
312	3.83	26.69	1.3	1.01	19.02	34.86	0.16
31200	3.82	26.64	1.29	1	18.97	34.83	0.15
31201	0	0	0	0	0	0	0
31202	0	0	0	0	0	0	0
31203	0	0	0	0	0	0	0
31204	0	0	0	0	0	0	0
31205	0.01	0.05	0.01	0.01	0.05	0.03	0
31206	0	0	0	0	0	0	0
313	39.7	195.2	10.31	11.75	156.54	240.63	1.99
31300	39.7	194.66	9.94	9.83	156.31	240.4	1.99
31301	0	0	0	0	0	0	0
31302	0	0	0	0	0	0	0
31303	0	0	0	0	0	0	0
31304	0	0	0	0	0	0	0
31305	0	0.54	0.37	1.91	0.23	0.23	0
31306	0	0	0	0	0	0	0
314	66.39	288.46	31.08	50.59	710.51	342.59	12.27
31400	62.86	257.91	26.7	39.62	632.91	306.63	10.98
31401	0	0	0	0	0	0	0
31402	0	0	0	0	0	0	0
31403	3.52	28.82	3.2	4.8	76.85	35.23	1.28
31404	0	0	0	0	0	0	0
31405	0.01	1.73	1.18	6.16	0.74	0.73	0.01
31406	0	0	0	0	0	0	0
315	0.29	34.08	1.72	1	18.59	39.33	0.12
31500	0.29	34.08	1.72	1	18.59	39.33	0.12
31501	0	0	0	0	0	0	0
31502	0	0	0	0	0	0	0
31503	0	0	0	0	0	0	0
31504	0	0	0	0	0	0	0
31505	0	0	0	0	0	0	0
31506	0	0	0	0	0	0	0
316	504	1621	78.43	66.72	1372	865.82	12.25
31600	504	1621	78.43	66.72	1372	865.82	12.25
31601	0	0	0	0	0	0	0
31602	0	0	0	0	0	0	0
31603	0	0	0	0	0	0	0
31604	0	0	0	0	0	0	0
31605	0	0	0	0	0	0	0
31606	0	0	0	0	0	0	0
320	484.95	484.26	12.51	40.71	207.27	221.63	4.28

SNAP	SO ₂ [Mg]	NO _x [Mg]	NMVOG [Mg]	CH ₄ [Mg]	CO [Mg]	CO ₂ [Gg] ¹⁾	N ₂ O [Mg]
32000	484.5	476.44	10.11	19.49	198.84	218.33	4.23
32001	0	0	0	0	0	0	0
32002	0	0	0	0	0	0	0
32003	0	0	0	0	0	0	0
32004	0	0.04	0	0	0	0.05	0
32005	0.45	7.78	2.4	21.22	8.43	3.25	0.05
32006	0	0	0	0	0	0	0

¹⁾ Including CO₂ emission from biomass.

Annex 3A-9 Description of the Danish energy statistics

This description of the Danish energy statistics has been prepared by Denmark's National Environmental Research Institute, NERI (now DCE) in cooperation with the Danish Energy Agency (DEA) as background information to the Danish National Inventory Report (NIR).

The Danish energy statistics system

DEA is responsible for the Danish energy balance. Main contributors to the energy statistics outside DEA are Statistics Denmark and Danish Energy Association (before Association of Danish Energy Companies). The statistics is performed using an integrated statistical system building on an Access database and Excel spreadsheets.

The DEA follows the recommendations of the International Energy Agency as well as Eurostat.

The national energy statistics is updated annually and all revisions are immediately included in the published statistics, which can be found on the DEA homepage³. It is an easy task to check for breaks in a series because the statistics is 100% time-series oriented.

The national energy statistics does not include Greenland and Faroe Islands.

For historical reasons, DEA receive monthly information from the Danish oil companies regarding Danish deliveries of oil products to Greenland and Faroe Islands. However, the monthly (MOS) and annual (AOS) reporting of oil statistics to Eurostat and IEA exclude Greenland and Faroe Islands. For all other energy products, the Danish figures are also excluding Greenland and Faroe Islands.

Reporting to the Danish Energy Agency

The Danish Energy Agency receives monthly statistics for the following fuel groups:

- Crude oil and oil products.
- Monthly data from 46 oil companies, the main purpose is monitoring oil stocks according to the oil preparedness system.
- Natural gas.
- Fuel/flare from platforms in the North Sea.
- Natural gas balance from the regulator Energinet.dk (National monopoly).
- Coal and coke.
- Power plants (94 %).
- Industry companies (4 %).
- Coal and coke traders (2 %).
- Electricity.
- Monthly reporting by e-mail from the regulator Energinet.dk (National monopoly).
- The statistics covers:
 - Production by type of producer.
 - Own use of electricity.
 - Import and export by country.

³ http://www.ens.dk/EN-US/INFO/FACTSANDFIGURES/ENERGY_STATISTICS_AND_INDICATORS/ANNUAL%20STATISTICS/Sider/Forside.aspx

- Domestic supply (consumption + distribution loss).
- Town gas (quarterly) from two town gas producers.
- The large central power plants also report monthly consumption of biomass.

Annual data includes renewable energy including waste. The DEA conducts a biannual survey on wood pellets and wood fuel. Statistics Denmark conducts biannual surveys on the energy consumption in the service and industrial sectors. Statistics Denmark prepares annual surveys on forest (wood fuel) & straw.

Other annual data sources include:

- DEA.
- Survey on production of electricity and heat and fuels used.
- Survey on end use of oil.
- Survey on end use of natural gas.
- Survey on end use of coal and coke.
- DCE, Aarhus University.
- Energy consumption for domestic air transport.
- Danish Energy Association (Association of Danish Energy companies).
- Survey on electricity consumption.
- Ministry of Taxation.
- Border trade.
- Centre for Biomass Technology.
- Annual estimates of final consumption of straw and wood chips.

Annual revisions

In general, DEA follows the same procedures as in the Danish national account. This means that normally only figures for the last two years are revised.

Aggregating the energy statistics on SNAP level

The sectors used in the official energy statistics have been mapped to SNAP categories, used in the Danish emission database. DCE aggregates the official energy statistics to SNAP level based on a source correspondence table.

In cooperation between DEA and DCE, a fuel correspondence table has been developed mapping the fuels used by the DEA in the official energy statistics with the fuel codes used in the Danish national emission database. The fuel correspondence table between fuel categories used by the DEA, DCE and IPCC is presented in Annex 3A-3.

The mapping between the energy statistics and the SNAP and fuel codes used by DCE can be seen in the table below.

Table 3A-9.1 Correspondence between the Danish national energy statistics and the SNAP nomenclature (only stationary combustion part shown).

Unit: TJ	Enduse		Transformation	
	SNAP	Fuel	SNAP	Fuel
Energy Sector				
Extraction and Gasification				
- Extraction				
- - Natural Gas	010504	301A		
- Gasification				
- - Biogas, Landfill				
- - Biogas, Other				
- - Electricity				
Refineries				
- Used for Refining				
- - Crude Oil				
- - Refinery Feedstocks				
- - Electricity				
- - District Heating				
- Own Use				
- - Refinery Gas	010306	308A		
- - LPG	010306	303A		
- - Gas-/Diesel Oil	010306	204A		
- - Fuel Oil	010306	203A		
- Net Production				
- - Refinery Gas				
- - LPG				
- - Naphtha (LVN)				
- - Aviation Gasoline				
- - Motor Gasoline				
- - JP4				
- - Other Kerosene				
- - JP1				
- - Gas-/Diesel Oil				
- - Fuel Oil				
- - Petroleum Coke				
- - White Spirit				
- - Lubricants				
- - Bitumen				
Distribution				
- Electricity Used in Distribution				
- - Electricity Distribution				
- - District Heating Distribution				
- - Gas Distribution				
Transformation Sector				
Large-scale Power Units				
- Fuels Used for Power Production				
- - Gas-/Diesel Oil			010100	204A
- - Fuel Oil			010100	203A
- - Electricity Plant Coal			010100	102A
- - Straw			010100	117A
- Own Use				
- - Electricity				
- Gross Production				
- - Electricity				
Large-Scale CHP Units				
- Fuels Used for Power Production				
- - Refinery Gas			010300	308A
- - LPG			010100	303A
- - Naphtha (LVN)			010100	210A
- - Gas-/Diesel Oil			010100	204A
- - Fuel Oil			010100	203A
- - Petroleum Coke			010100	110A
- - Orimulsion			010100	225A
- - Natural Gas			010100	301A
- - Electricity Plant Coal			010100	102A
- - Straw			010100	117A
- - Wood Chips			010100	111A
- - Wood Pellets			010100	111A
- - Wood Waste			010100	111A
- - Biogas, Landfill			010100	309A
- - Biogas, Sludge			010100	309A
- - Biogas, Others			010100	309A

- - Waste, Non-renewable		010100	114A
- - Wastes, Renewable		010100	114A
<i>Continued</i>			
- Fuels Used for Heat Production			
- - Refinery Gas		010300	308A
- - LPG		010100	303A
- - Naphtha (LVN)		010100	210A
- - Gas-/Diesel Oil		010100	204A
- - Fuel Oil		010100	203A
- - Petroleum Coke		010100	110A
- - Orimulsion		010100	225A
- - Natural Gas		010100	301A
- - Electricity Plant Coal		010100	102A
- - Straw		010100	117A
- - Wood Chips		010100	111A
- - Wood Pellets		010100	111A
- - Wood Waste		010100	111A
- - Biogas, Landfill		010100	309A
- - Biogas, Sludge		010100	309A
- - Biogas, Other		010100	309A
- - Wastes, Non-renewable		010100	114A
- - Wastes, Renewable		010100	114A
- Own Use			
- - Electricity			
- - District Heating			
- Production			
- - Electricity, Gross			
- - District Heating, Net			
Small-Scale CHP Units			
- Fuels Used for Power Production			
- - Gas-/Diesel Oil		010100	204A
- - Fuel Oil		010100	203A
- - Natural Gas		010100	301A
- - Hard Coal		010100	102A
- - Straw		010100	117A
- - Wood Chips		010100	111A
- - Wood Pellets		010100	111A
- - Wood Waste		010100	111A
- - Biogas, Landfill		010100	309A
- - Biogas, Sludge		010100	309A
- - Biogas, Other		010100	309A
- - Waste, Non-renewable		010100	114A
- - Wastes, Renewable		010100	114A
- Fuels Used for Heat Production			
- - Gas-/Diesel Oil		010100	204A
- - Fuel Oil		010100	203A
- - Natural Gas		010100	301A
- - Coal		010100	102A
- - Straw		010100	117A
- - Wood Chips		010100	111A
- - Wood Pellets		010100	111A
- - Wood Waste		010100	111A
- - Biogas, Landfill		010100	309A
- - Biogas, Sludge		010100	309A
- - Biogas, Other		010100	309A
- - Wastes, Non-renewable		010100	114A
- - Wastes, Renewable		010100	114A
- Own Use			
- - Electricity			
- - District Heating			
- Production			
- - Electricity, Gross			
- - District Heating, Net			
Wind Turbines			
- Used for Power Production			
- - Wind Power			
- Gross Production			
- - Electricity			
Hydro Power Units			
- Used for Power Production			
- - Hydro Power			
- Gross Production			
- - Electricity			

District Heating Units			
- Fuels Used for Heat Production			
- - Refinery Gas		010300	308A
- - LPG		010200	303A
<i>Continued</i>			
- - Gas-/Diesel Oil		010200	204A
- - Fuel Oil		010200	203A
- - Waste Oil		010200	203A
- - Petroleum Coke		010200	110A
- - Natural Gas		010200	301A
- - Electricity Plant Coal		010200	102A
- - Coal		010200	102A
- - Solar Energy			
- - Geothermal Energy			
- - Straw		010200	117A
- - Wood Chips		010200	111A
- - Wood Pellets		010200	111A
- - Wood Waste		010200	111A
- - Biogas, Landfill		010200	309A
- - Biogas, Sludge		010200	309A
- - Biogas, Other		010200	309A
- - Wastes, Non-renewable		010200	114A
- - Wastes, Renewable		010200	114A
- - Fish Oil		010200	215A
- - Electricity for Heat Pumps			
- Own Use			
- - District Heating			
- Net Production			
- - District Heating			
Autoproducers, Electricity Only			
- Fuels Used for Power Production			
- - Natural Gas		030100	301A
- - Solar Energy			
- - Biogas, Landfill		030100	309A
- - Biogas, Sewage Sludge		030100	309A
- - Biogas, Other		030100	309A
- Gross Production			
- - Electricity			
Autoproducers, CHP Units			
- Fuels Used for Power Production			
- - Refinery Gas		010300	308A
- - Gas-/Diesel Oil		030100	204A
- - Fuel Oil		030100	203A
- - Waste Oil		030100	203A
- - Natural Gas		030100	301A
- - Coal		030100	102A
- - Straw		030100	117A
- - Wood Chips		030100	111A
- - Wood Pellets		030100	111A
- - Wood Waste		030100	111A
- - Biogas, Landfill		030100	309A
- - Biogas, Sludge		030100	309A
- - Biogas, Other		030100	309A
- - Fish Oil		030100	215A
- - Wastes, Non-renewable		010100	114A
- - Wastes, Renewable		010100	114A
- Fuels Used for Heat Production		030100	114A
- - Refinery Gas		010300	308A
- - Gas-/Diesel Oil		030100	204A
- - Fuel Oil		030100	203A
- - Waste Oil		030100	203A
- - Natural Gas		030100	301A
- - Coal		030100	102A
- - Wood Chips		030100	111A
- - Wood Waste		030100	111A
- - Biogas, Landfill		030100	309A
- - Biogas, Sludge		030100	309A
- - Biogas, Other		030100	309A
- - Wastes, Non-renewable		010100	114A
- - Wastes, Renewable		010100	114A
- Production			
- - Electricity, Gross			
- - District Heating, Net			

Autoproducers, Heat Only		
- Fuels Used for Heat Production		
- - Gas-/Diesel Oil		030100 204A
- - Fuel Oil		030100 203A
- - Waste Oil		030100 203A
- - Natural Gas		030100 301A
- - Straw		030100 117A
<i>Continued</i>		
- - Wood Chips		030100 111A
- - Wood Chips		030100 111A
- - Wood Waste		030100 111A
- - Biogas, Landfill		030100 309A
- - Biogas, Sludge		030100 309A
- - Biogas, Other		030100 309A
- - Wastes, Non-renewable		010200 114A
- - Wastes, Renewable		010200 114A
- - Heat Pumps		
- Net Production		
- - District Heating		
Gas Works Gas Units	030106	301A
- Fuels Used for Production of District Heating		
- - Refinery Gas		
- - LPG		
- - Naphtha (LVN)		
- - Gas-/Diesel Oil		
- - Natural Gas		
- - Hard Coal		
- Production		
- - Gas Works Gas		
- - Coke		
Distribution Losses		
- Distribution Losses etc.		
- - Natural Gas		
- - Electricity		
- - District Heating		
- - Gas Works Gas		
Consumption Sector		
- Non-energy Use		
- - White Spirit		
- - Lubricants		
- - Bitumen		
Transport		
Military Transport		
- Aviation Gasoline		
- Motor Gasoline		
- JP4		
- JP1		
- Gas-/Diesel Oil		
Road		
- LPG		
- Motor Gasoline		
- Other Kerosene	020200	206A
- Gas-/Diesel Oil		
- Fuel Oil		
- Bioethanol		
- Biodiesel		
Rail		
- Motor Gasoline		
- Other Kerosene		
- Gas-/Diesel Oil		
- Electricity		
Domestic Sea Transport		
- LPG	Transport	
- Other Kerosene	Transport	
- Gas-/Diesel Oil	Transport	
- Fuel Oil	Transport	
Domestic Aviation		
- LPG	Transport	
- Aviation Gasoline	Transport	
- Motor Gasoline	Transport	
- Other Kerosene	020100	206A

- JP1	Transport	
International Aviation		
- Aviation Gasoline	Transport	
- JP1	Transport	
Agriculture and Forestry		
- LPG	Transport	
- Motor Gasoline	Transport	
- Other Kerosene	020300	206A
- Gas-/Diesel Oil	Transport	
- Fuel Oil	020300	203A
- Petroleum Coke	020300	110A
<i>Continued</i>		
- Natural Gas	020300	301A
- Coal	020300	102A
- Brown Coal Briquettes	020300	106A
- Straw	020300	117A
- Wood Chips	020300	111A
- Wood Waste	020300	111A
- Biogas, Other	020300	309A
- Heat Pumps		
- Electricity		
Horticulture		
- LPG	Transport	
- Motor Gasoline	Transport	
- Gas-/Diesel Oil	Transport	
- Fuel Oil	020300	203A
- Petroleum Coke	020300	110A
- Natural Gas	020300	301A
- Coal	020300	102A
- Wood Waste	020300	111A
- Electricity		
- District Heating		
Fishing		
- LPG	Transport	
- Motor Gasoline	Transport	
- Other Kerosene	Transport	
- Gas-/Diesel Oil	Transport	
- Fuel Oil	Transport	
Manufacturing Industry		
- Refinery Gas	030100	308A
- LPG	Transport	
- Naphtha (LVN)	Transport	
- Motor Gasoline	Transport	
- Other Kerosene	030100	206A
- Gas-/Diesel Oil	Transport	
- Fuel Oil	030100	203A
- Waste Oil	030100	203A
- Petroleum Coke	030100	110A
- Natural Gas	030100	301A
- Coal	030100	102A
- Coke	030100	107A
- Brown Coal Briquettes	030100	106A
- Wood Pellets	030100	111A
- Wood Waste	030100	111A
- Biogas, Landfill	030100	111A
- Biogas, Sludge	030100	309A
- Biogas, Other	030100	309A
- Wastes, Non-renewable	030100	114A
- Wastes, Renewable	030100	114A
- Heat Pumps		
- Electricity		
- District Heating		
- Gas Works Gas	030100	301A
Construction		
- LPG	031500	303A
- Motor Gasoline	Transport	
- Other Kerosene	031500	206A
- Gas-/Diesel Oil	Transport	
- Fuel Oil	031500	203A
- Natural Gas	031500	301A
- Electricity		
Wholesale		
- LPG	020100	303A

- Motor Gasoline	020100	206A	
- Other Kerosene	020100	204A	
- Gas-/Diesel Oil	020100	203A	
- Petroleum Coke	020100	110A	
- Natural Gas	020100	301A	
- Wood Waste	020100	111A	
- Electricity			
- District Heating			
Retail Trade			
- LPG	020100	303A	
- Other Kerosene	020100	206A	
- Gas-/Diesel Oil	020100	204A	
- Fuel Oil	020100	203A	
- Petroleum Coke	020100	110A	
<i>Continued</i>			
- Natural Gas	020100	301A	
- Electricity			
- District Heating			
Private Service			
- LPG	020100	303A	
- Other Kerosene	020100	206A	
- Gas-/Diesel Oil	020100	204A	
- Fuel Oil	020100	203A	
- Waste Oil	020100	203A	
- Petroleum Coke	020100	110A	
- Natural Gas	020100	301A	
- Wood Chips	020100	111A	
- Wood Waste	020100	111A	
- Biogas, Landfill	020100	309A	
- Biogas, Sludge	020100	309A	
- Biogas, Other	020100	309A	
- Wastes, Non-renewable	020100	114A	
- Wastes, Renewable	020100	114A	
- Electricity			
- District Heating			
- Gas Works Gas	020100	301A	
Public Service			
- LPG	020100	303A	
- Other Kerosene	020100	206A	
- Gas-/Diesel Oil	020100	204A	
- Fuel Oil	020100	203A	
- Petroleum Coke	020100	110A	
- Natural Gas	020100	301A	
- Coal	020100	102A	
- Brown Coal Briquettes	020100	106A	
- Solar Energy			
- Wood Chips	020100	111A	
- Wood Pellets	020100	111A	
- Electricity			
- District Heating			
- Gas Works Gas	020100	301A	
Single Family Houses			
- LPG	020200	303A	
- Motor Gasoline	Transport		
- Other Kerosene	020200	206A	
- Gas-/Diesel Oil	020200	204A	
- Fuel Oil	020200	203A	
- Petroleum Coke	020200	110A	
- Natural Gas	020200	301A	
- Coal	020200	102A	
- Coke	020200	107A	
- Brown Coal Briquettes	020200	106A	
- Solar Energy			
- Straw	020200	117A	
- Firewood	020200	111A	
- Wood Chips	020200	111A	
- Wood Pellets	020200	111A	
- Biodiesel	020200	215A	
- Heat Pumps			
- Electricity			
- District Heating			
- Gas Works Gas	020200	301A	

Multi-family Houses			
- LPG	020200	303A	
- Other Kerosene	020200	206A	
- Gas-/Diesel Oil	020200	204A	
- Fuel Oil	020200	203A	
- Petroleum Coke	020200	110A	
- Natural Gas	020200	301A	
- Coal	020200	102A	
- Coke	020200	107A	
- Brown Coal Briquettes	020200	106A	
- Solar Energy			
- Electricity			
- District Heating			
- Gas Works Gas	020200	301A	

Annex 3A-10 EU ETS data for coal

EU ETS data are available for the years 2006-2012. Corresponding values for lower calorific value (LCV) and implied emission factor (IEF) for CO₂ for 2006-2009 are shown in Figure 3A-10.1. The IEF factors include the oxidation factors.




Figure 3A-10.1 EU ETS data for LCV and CO₂ IEF (including oxidation factor) for coal. Data for the years 2006-2009.

Annex 3B - Transport

List of content

Annex 3B-1: Fleet data 1985-2012 for road transport (No. vehicles)

Annex 3B-2: Mileage data 1985-2012 for road transport (km)

Annex 3B-3: EU directive emission limits for road transportation vehicles

Annex 3B-4: Basis emission factors (g/km)

Annex 3B-5: Reduction factors for road transport emission factors

Annex 3B-6: Deterioration factors for road transport emission factors

Annex 3B-7: Final fuel consumption factors (MJ/km) and emission factors (g/km) in 2012

Annex 3B-8: Fuel consumption (GJ) and emissions (tons) per vehicle category and as totals

Annex 3B-9: COPERT IV:DEA statistics fuel use ratios and mileage adjustment factors

Annex 3B-10: Actual vs. representative aircraft types, no. of LTO's from Danish airports, no. of flights between Denmark and Greenland/Faroe Islands, LTO and average cruise fuel consumption and emission factors

Annex 3B-11: Basis fuel consumption and emission factors, deterioration factors, transient factors, stock and activity data for non road working machinery and equipment, and recreational craft

Annex 3B-12: Traffic data and different technical and operational data for Danish domestic ferries

Annex 3B-13: Fuel consumption and emission factors, engine specific (NO_x , CO, VOC (NMVOC and CH_4)), and fuel type specific (S-%, SO_2 , PM) for ship engines

Annex 3B-14: Fuel sales figures from DEA, and further processed fuel consumption data suited for the Danish inventory

Annex 3B-15: Emission factors and total emissions in CollectER format

Annex 3B-16: Fuel consumption and emissions in CRF format

Annex 3B-17: Uncertainty estimates

Annex 3B-1 Fleet data 1985-2012 for road transport (No. vehicles)

Sector	Subsector	Tech 2	FYear	LYear	1985	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
Passenger Cars	Gasoline 0,8 - 1,4 l	PRE ECE	0	1969	75564	16627	13368	10706	8571	7246	6992	6618	6159	5646	5194
Passenger Cars	Gasoline 0,8 - 1,4 l	ECE 15/00-01	1970	1978	404440	179963	156167	134583	102209	66638	55669	43359	30440	19722	12950
Passenger Cars	Gasoline 0,8 - 1,4 l	ECE 15/02	1979	1980	97500	87416	63723	53008	61799	45282	38690	30726	21910	14275	8539
Passenger Cars	Gasoline 0,8 - 1,4 l	ECE 15/03	1981	1985	152241	318620	330061	307288	254029	235151	221927	204913	179982	150784	119474
Passenger Cars	Gasoline 0,8 - 1,4 l	ECE 15/04	1986	1990		165103	178393	209260	261579	258381	253651	249450	243072	232062	220895
Passenger Cars	Gasoline 0,8 - 1,4 l	Euro I	1991	1996			28375	60724	96922	141546	180780	219477	218990	216002	214711
Passenger Cars	Gasoline 0,8 - 1,4 l	Euro II	1997	2000									39547	74071	106936
Passenger Cars	Gasoline 0,8 - 1,4 l	Euro III	2001	2005											
Passenger Cars	Gasoline 0,8 - 1,4 l	Euro IV	2006	2010											
Passenger Cars	Gasoline 0,8 - 1,4 l	Euro V	2011	2014											
Passenger Cars	Gasoline 1,4 - 2,0 l	PRE ECE	0	1969	90872	28856	23474	19524	15744	13167	12527	11642	10624	9570	8659
Passenger Cars	Gasoline 1,4 - 2,0 l	ECE 15/00-01	1970	1978	344507	171158	152919	137410	110813	76213	63961	50125	35583	23605	15800
Passenger Cars	Gasoline 1,4 - 2,0 l	ECE 15/02	1979	1980	87587	74393	54644	44813	52998	40866	35395	28785	21181	14516	9144
Passenger Cars	Gasoline 1,4 - 2,0 l	ECE 15/03	1981	1985	210664	276844	281145	261224	218177	205239	196226	184150	165329	142253	115689
Passenger Cars	Gasoline 1,4 - 2,0 l	ECE 15/04	1986	1990		221807	211098	215194	242500	240697	238039	236139	232642	225250	217019
Passenger Cars	Gasoline 1,4 - 2,0 l	Euro I	1991	1996			51521	101611	148509	235536	319571	414973	413070	407030	404816
Passenger Cars	Gasoline 1,4 - 2,0 l	Euro II	1997	2000									105322	217501	303755
Passenger Cars	Gasoline 1,4 - 2,0 l	Euro III	2001	2005											
Passenger Cars	Gasoline 1,4 - 2,0 l	Euro IV	2006	2010											
Passenger Cars	Gasoline 1,4 - 2,0 l	Euro V	2011	2014											
Passenger Cars	Gasoline >2,0 l	PRE ECE	0	1969	3246	1388	1186	1033	897	911	945	971	986	987	989
Passenger Cars	Gasoline >2,0 l	ECE 15/00-01	1970	1978	3113	3661	3581	3373	3096	2800	2589	2352	2039	1657	1381
Passenger Cars	Gasoline >2,0 l	ECE 15/02	1979	1980	1078	564	531	687	859	865	865	846	773	702	599
Passenger Cars	Gasoline >2,0 l	ECE 15/03	1981	1985	4087	2263	2037	1700	1575	1659	1801	1950	2055	2081	2018
Passenger Cars	Gasoline >2,0 l	ECE 15/04	1986	1990		4323	3630	3161	2668	2810	3052	3331	3638	3874	4089
Passenger Cars	Gasoline >2,0 l	Euro I	1991	1996			1263	2350	3350	5384	7888	10682	11000	11250	11334
Passenger Cars	Gasoline >2,0 l	Euro II	1997	2000									3980	8667	14014
Passenger Cars	Gasoline >2,0 l	Euro III	2001	2005											
Passenger Cars	Gasoline >2,0 l	Euro IV	2006	2010											
Passenger Cars	Gasoline >2,0 l	Euro V	2011	2014											
Passenger Cars	Diesel 1,4 - 2,0 l	Conventional	0	1990	69406	71018	70198	69500	68720	65169	62762	59117	54631	50590	48238
Passenger Cars	Diesel 1,4 - 2,0 l	Euro I	1991	1996			979	2163	3799	6613	9919	13122	13689	14318	15305
Passenger Cars	Diesel 1,4 - 2,0 l	Euro II	1997	2000									3064	8535	18211
Passenger Cars	Diesel 1,4 - 2,0 l	Euro III	2001	2005											

Passenger Cars	Diesel 1,4 - 2,0 l	Euro IV	2006	2010											
Passenger Cars	Diesel 1,4 - 2,0 l	Euro V	2011	2014											
Passenger Cars	Diesel >2,0 l	Conventional	0	1990	14055	14871	13888	13012	12136	11757	11413	10708	10043	9269	8435
Passenger Cars	Diesel >2,0 l	Euro I	1991	1996			1017	1988	3035	4323	5638	7401	7600	7595	7716
Passenger Cars	Diesel >2,0 l	Euro II	1997	2000									2079	5072	9083
Passenger Cars	Diesel >2,0 l	Euro III	2001	2005											
Passenger Cars	Diesel >2,0 l	Euro IV	2006	2010											
Passenger Cars	Diesel >2,0 l	Euro V	2011	2014											
Passenger Cars	LPG cars	Conventional	0	1990	1136	1163	1166	1173	1184	734	495	310	171	96	56
Passenger Cars	LPG cars	Euro I	1991	1996				1	4	4	3	1	1	1	3
Passenger Cars	LPG cars	Euro II	1997	2000											
Passenger Cars	LPG cars	Euro III	2001	2005											
Passenger Cars	LPG cars	Euro IV	2006	2010											
Passenger Cars	2-Stroke	Conventional	0	9999	4823	5417	4804	4308	3747	3029	2443	1824	1248	761	400
Passenger Cars	Electric cars	Conventional	0	9999	130	133	133	134	136	155	163	187	230	292	298
Passenger Cars	Gasoline <0,8 l	PRE ECE	0	1969											
Passenger Cars	Gasoline <0,8 l	ECE 15/00-01	1970	1978											
Passenger Cars	Gasoline <0,8 l	ECE 15/02	1979	1980											
Passenger Cars	Gasoline <0,8 l	ECE 15/03	1981	1985											
Passenger Cars	Gasoline <0,8 l	ECE 15/04	1986	1990											
Passenger Cars	Gasoline <0,8 l	Euro I	1991	1996											
Passenger Cars	Gasoline <0,8 l	Euro II	1997	2000											40
Passenger Cars	Gasoline <0,8 l	Euro III	2001	2005											
Passenger Cars	Gasoline <0,8 l	Euro IV	2006	2010											
Passenger Cars	Gasoline <0,8 l	Euro V	2011	2014											
Passenger Cars	Diesel <1,4 l	Conventional	0	1990											
Passenger Cars	Diesel <1,4 l	Euro I	1991	1996											
Passenger Cars	Diesel <1,4 l	Euro II	1997	2000											361
Passenger Cars	Diesel <1,4 l	Euro III	2001	2005											
Passenger Cars	Diesel <1,4 l	Euro IV	2006	2010											
Passenger Cars	Diesel <1,4 l	Euro V	2011	2014											
Light Duty Vehicles	Gasoline <3,5t	Conventional	0	1994	34172	44442	45625	46865	48934	49865	46712	42710	37987	34274	30224
Light Duty Vehicles	Gasoline <3,5t	Euro I	1995	1998							3773	7509	12025	17550	17352
Light Duty Vehicles	Gasoline <3,5t	Euro II	1999	2001											5272
Light Duty Vehicles	Gasoline <3,5t	Euro III	2002	2006											
Light Duty Vehicles	Gasoline <3,5t	Euro IV	2007	2011											
Light Duty Vehicles	Gasoline <3,5t	Euro V	2012	2015											
Light Duty Vehicles	Diesel <3,5t	Conventional	0	1994	113019	146986	150898	154999	161842	169142	160228	148520	133718	120795	105967

Light Duty Vehicles	Diesel <3,5t	Euro I	1995	1998							16899	35370	56836	76717	75753
Light Duty Vehicles	Diesel <3,5t	Euro II	1999	2001											24555
Light Duty Vehicles	Diesel <3,5t	Euro III	2002	2006											
Light Duty Vehicles	Diesel <3,5t	Euro IV	2007	2011											
Light Duty Vehicles	Diesel <3,5t	Euro V	2012	2015											
Light Duty Vehicles	LPG <3,5t	Conventional	0	1994	684	889	913	938	979	632	462	295	196	125	90
Light Duty Vehicles	LPG <3,5t	Euro I	1995	1998										1	1
Light Duty Vehicles	LPG <3,5t	Euro II	1999	2001											
Light Duty Vehicles	LPG <3,5t	Euro III	2002	2006											
Light Duty Vehicles	LPG <3,5t	Euro IV	2007	2011											
Light Duty Vehicles	Electric <3,5t	Conventional	0	9999	3	4	4	4	4	3	2	2	1	1	1
Heavy Duty Vehicles	Gasoline >3,5t	Conventional	0	9999	621	530	510	497	503	455	412	365	326	336	318
Heavy Duty Vehicles	Diesel RT 3,5 - 7,5t	Conventional	0	1993	8686	7049	6675	6430	6419	6194	5738	5137	4646	4156	3518
Heavy Duty Vehicles	Diesel RT 3,5 - 7,5t	Euro I	1994	1996					66	376	711	976	973	967	906
Heavy Duty Vehicles	Diesel RT 3,5 - 7,5t	Euro II	1997	2001								89	521	1236	1782
Heavy Duty Vehicles	Diesel RT 3,5 - 7,5t	Euro III	2002	2006											
Heavy Duty Vehicles	Diesel RT 3,5 - 7,5t	Euro IV	2007	2009											
Heavy Duty Vehicles	Diesel RT 3,5 - 7,5t	Euro V	2010	2013											
Heavy Duty Vehicles	Diesel RT 3,5 - 7,5t	Euro VI	2014	9999											
Heavy Duty Vehicles	Diesel RT 7,5 - 12t	Conventional	0	1993	7266	5897	5584	5379	5375	5316	5373	5207	4854	4491	4116
Heavy Duty Vehicles	Diesel RT 7,5 - 12t	Euro I	1994	1996					51	298	671	968	1002	1081	1102
Heavy Duty Vehicles	Diesel RT 7,5 - 12t	Euro II	1997	2001								94	429	798	1200
Heavy Duty Vehicles	Diesel RT 7,5 - 12t	Euro III	2002	2006											
Heavy Duty Vehicles	Diesel RT 7,5 - 12t	Euro IV	2007	2009											
Heavy Duty Vehicles	Diesel RT 7,5 - 12t	Euro V	2010	2013											
Heavy Duty Vehicles	Diesel RT 7,5 - 12t	Euro VI	2014	9999											
Heavy Duty Vehicles	Diesel RT 12 - 14 t	Conventional	0	1993	4984	4519	4461	4388	4454	3991	3248	2731	2360	1984	1623
Heavy Duty Vehicles	Diesel RT 12 - 14 t	Euro I	1994	1996					37	156	234	285	283	286	289
Heavy Duty Vehicles	Diesel RT 12 - 14 t	Euro II	1997	2001								21	126	216	262
Heavy Duty Vehicles	Diesel RT 12 - 14 t	Euro III	2002	2006											
Heavy Duty Vehicles	Diesel RT 12 - 14 t	Euro IV	2007	2009											
Heavy Duty Vehicles	Diesel RT 12 - 14 t	Euro V	2010	2013											
Heavy Duty Vehicles	Diesel RT 14 - 20t	Conventional	0	1993	5171	4689	4628	4552	4601	4348	4047	3669	3316	2924	2537
Heavy Duty Vehicles	Diesel RT 14 - 20t	Euro I	1994	1996					58	334	708	1001	1007	985	963
Heavy Duty Vehicles	Diesel RT 14 - 20t	Euro II	1997	2001								98	535	937	1371
Heavy Duty Vehicles	Diesel RT 14 - 20t	Euro III	2002	2006											
Heavy Duty Vehicles	Diesel RT 14 - 20t	Euro IV	2007	2009											
Heavy Duty Vehicles	Diesel RT 14 - 20t	Euro V	2010	2013											

Heavy Duty Vehicles	Diesel RT 14 - 20t	Euro VI	2014	9999											
Heavy Duty Vehicles	Diesel RT 20 - 26t	Conventional	0	1993	4307	5179	5237	5326	5315	5031	4565	4059	3536	3067	2596
Heavy Duty Vehicles	Diesel RT 20 - 26t	Euro I	1994	1996					67	469	1003	1452	1442	1400	1322
Heavy Duty Vehicles	Diesel RT 20 - 26t	Euro II	1997	2001								152	748	1330	1898
Heavy Duty Vehicles	Diesel RT 20 - 26t	Euro III	2002	2006											
Heavy Duty Vehicles	Diesel RT 20 - 26t	Euro IV	2007	2009											
Heavy Duty Vehicles	Diesel RT 20 - 26t	Euro V	2010	2013											
Heavy Duty Vehicles	Diesel RT 20 - 26t	Euro VI	2014	9999											
Heavy Duty Vehicles	Diesel RT 26 - 28t	Conventional	0	1993	7	8	8	9	9	7	6	6	6	6	6
Heavy Duty Vehicles	Diesel RT 26 - 28t	Euro I	1994	1996							0	1	1	1	1
Heavy Duty Vehicles	Diesel RT 26 - 28t	Euro II	1997	2001								0	1	2	3
Heavy Duty Vehicles	Diesel RT 26 - 28t	Euro III	2002	2006											
Heavy Duty Vehicles	Diesel RT 26 - 28t	Euro IV	2007	2009											
Heavy Duty Vehicles	Diesel RT 26 - 28t	Euro V	2010	2013											
Heavy Duty Vehicles	Diesel RT 28 - 32t	Conventional	0	1993	271	326	329	335	327	326	329	321	300	262	231
Heavy Duty Vehicles	Diesel RT 28 - 32t	Euro I	1994	1996					11	62	152	239	246	252	253
Heavy Duty Vehicles	Diesel RT 28 - 32t	Euro II	1997	2001								28	147	289	455
Heavy Duty Vehicles	Diesel RT 28 - 32t	Euro III	2002	2006											
Heavy Duty Vehicles	Diesel RT 28 - 32t	Euro IV	2007	2009											
Heavy Duty Vehicles	Diesel RT 28 - 32t	Euro V	2010	2013											
Heavy Duty Vehicles	Diesel RT 28 - 32t	Euro VI	2014	9999											
Heavy Duty Vehicles	Diesel RT >32t	Conventional	0	1993	0	0	0	0	0	0	1	0	0	1	1
Heavy Duty Vehicles	Diesel RT >32t	Euro I	1994	1996							0	1	1	1	1
Heavy Duty Vehicles	Diesel RT >32t	Euro II	1997	2001								0	1	0	0
Heavy Duty Vehicles	Diesel RT >32t	Euro III	2002	2006											
Heavy Duty Vehicles	Diesel RT >32t	Euro IV	2007	2009											
Heavy Duty Vehicles	Diesel RT >32t	Euro V	2010	2013											
Heavy Duty Vehicles	Diesel TT/AT 28 - 34t	Conventional	0	1993	5617	5132	5080	5011	5065	4783	4448	4025	3645	3208	2772
Heavy Duty Vehicles	Diesel TT/AT 28 - 34t	Euro I	1994	1996					63	356	759	1069	1076	1051	1028
Heavy Duty Vehicles	Diesel TT/AT 28 - 34t	Euro II	1997	2001								104	570	1000	1467
Heavy Duty Vehicles	Diesel TT/AT 28 - 34t	Euro III	2002	2006											
Heavy Duty Vehicles	Diesel TT/AT 28 - 34t	Euro IV	2007	2009											
Heavy Duty Vehicles	Diesel TT/AT 28 - 34t	Euro V	2010	2013											
Heavy Duty Vehicles	Diesel TT/AT 28 - 34t	Euro VI	2014	9999											
Heavy Duty Vehicles	Diesel TT/AT 34 - 40t	Conventional	0	1993	8359	10252	10740	11202	11174	10480	8917	7262	5877	4730	3842
Heavy Duty Vehicles	Diesel TT/AT 34 - 40t	Euro I	1994	1996					204	1616	3609	4958	4683	4110	3555
Heavy Duty Vehicles	Diesel TT/AT 34 - 40t	Euro II	1997	2001								495	2223	4240	5939
Heavy Duty Vehicles	Diesel TT/AT 34 - 40t	Euro III	2002	2006											

Heavy Duty Vehicles	Diesel TT/AT 34 - 40t	Euro IV	2007	2009											
Heavy Duty Vehicles	Diesel TT/AT 34 - 40t	Euro V	2010	2013											
Heavy Duty Vehicles	Diesel TT/AT 34 - 40t	Euro VI	2014	9999											
Heavy Duty Vehicles	Diesel TT/AT 40 - 50t	Conventional	0	1993	1672	2083	2242	2382	2379	2398	2257	2045	1799	1469	1240
Heavy Duty Vehicles	Diesel TT/AT 40 - 50t	Euro I	1994	1996					49	333	888	1316	1327	1314	1305
Heavy Duty Vehicles	Diesel TT/AT 40 - 50t	Euro II	1997	2001								143	778	1564	2540
Heavy Duty Vehicles	Diesel TT/AT 40 - 50t	Euro III	2002	2006											
Heavy Duty Vehicles	Diesel TT/AT 40 - 50t	Euro IV	2007	2009											
Heavy Duty Vehicles	Diesel TT/AT 40 - 50t	Euro V	2010	2013											
Heavy Duty Vehicles	Diesel TT/AT 40 - 50t	Euro VI	2014	9999											
Heavy Duty Vehicles	Diesel TT/AT 50 - 60t	Euro I	1994	1996								1	1	1	1
Heavy Duty Vehicles	Diesel TT/AT 50 - 60t	Euro II	1997	2001										1	1
Heavy Duty Vehicles	Diesel TT/AT 50 - 60t	Euro IV	2007	2009											
Heavy Duty Vehicles	Diesel TT/AT >60t	Euro III	2002	2006											
Heavy Duty Vehicles	Diesel TT/AT >60t	Euro V	2010	2013											
Buses	Gasoline Urban Buses	Conventional	0	9999	8	8	9	11	14	11	11	16	17	17	15
Buses	Diesel Urban Buses <15t	Conventional	0	1993	347	352	433	488	639	558	494	411	335	281	250
Buses	Diesel Urban Buses <15t	Euro I	1994	1996						49	81	122	130	132	124
Buses	Diesel Urban Buses <15t	Euro II	1997	2001									103	295	438
Buses	Diesel Urban Buses <15t	Euro III	2002	2006											
Buses	Diesel Urban Buses <15t	Euro IV	2007	2009											
Buses	Diesel Urban Buses <15t	Euro V	2010	2013											
Buses	Diesel Urban Buses 15 - 18t	Conventional	0	1993	2083	2109	2597	2928	3833	3475	3205	2861	2691	2353	2012
Buses	Diesel Urban Buses 15 - 18t	Euro I	1994	1996						397	632	985	989	891	891
Buses	Diesel Urban Buses 15 - 18t	Euro II	1997	2001									183	568	817
Buses	Diesel Urban Buses 15 - 18t	Euro III	2002	2006											
Buses	Diesel Urban Buses 15 - 18t	Euro IV	2007	2009											
Buses	Diesel Urban Buses 15 - 18t	Euro V	2010	2013											
Buses	Diesel Urban Buses >18t	Conventional	0	1993	5	5	6	7	9	8	6	7	6	3	2
Buses	Diesel Urban Buses >18t	Euro I	1994	1996						1	1	3	3	3	2
Buses	Diesel Urban Buses >18t	Euro II	1997	2001										6	20
Buses	Diesel Urban Buses >18t	Euro III	2002	2006											
Buses	Diesel Urban Buses >18t	Euro IV	2007	2009											
Buses	Diesel Urban Buses >18t	Euro V	2010	2013											
Buses	Gasoline Coaches	Conventional	0	9999	931	942	1161	1309	1508	1762	1775	1786	1791	1808	1810
Buses	Diesel Coaches <15t	Conventional	0	1993	3710	3756	4627	5215	6010	5926	5739	5506	5208	4941	4629
Buses	Diesel Coaches <15t	Euro I	1994	1996						420	682	1113	1103	1091	1056
Buses	Diesel Coaches <15t	Euro II	1997	2001									370	695	1039

Buses	Diesel Coaches <15t	Euro III	2002	2006											
Buses	Diesel Coaches <15t	Euro IV	2007	2009											
Buses	Diesel Coaches <15t	Euro V	2010	2013											
Buses	Diesel Coaches 15 - 18t	Conventional	0	1993	804	814	1003	1131	1303	1389	1393	1342	1253	1241	1184
Buses	Diesel Coaches 15 - 18t	Euro I	1994	1996						35	89	153	162	163	159
Buses	Diesel Coaches 15 - 18t	Euro II	1997	2001									44	77	119
Buses	Diesel Coaches 15 - 18t	Euro III	2002	2006											
Buses	Diesel Coaches 15 - 18t	Euro IV	2007	2009											
Buses	Diesel Coaches 15 - 18t	Euro V	2010	2013											
Buses	Diesel Coaches >18t	Conventional	0	1993	122	123	152	171	197	210	221	211	193	193	206
Buses	Diesel Coaches >18t	Euro I	1994	1996						20	42	78	84	82	81
Buses	Diesel Coaches >18t	Euro II	1997	2001									25	54	99
Buses	Diesel Coaches >18t	Euro III	2002	2006											
Buses	Diesel Coaches >18t	Euro IV	2007	2009											
Buses	Diesel Coaches >18t	Euro V	2010	2013											
Mopeds	2-stroke <50 cm³	Conventional	0	1999	151000	120000	118000	113000	109000	105000	114167	123333	132496	141636	150802
Mopeds	2-stroke <50 cm³	Euro I	2000	2003											
Mopeds	2-stroke <50 cm³	Euro II	2004	2014											
Mopeds	4-stroke <50 cm³	Euro II	2004	2014											
Motorcycles	2-stroke >50 cm³	Conventional	0	1999	6072	6470	6653	6737	6949	7255	7666	8228	8891	9524	10316
Motorcycles	2-stroke >50 cm³	Euro I	2000	2003											
Motorcycles	2-stroke >50 cm³	Euro II	2004	2006											
Motorcycles	2-stroke >50 cm³	Euro III	2007	9999											
Motorcycles	4-stroke <250 cm³	Conventional	0	1999	6881	7333	7541	7635	7875	8222	8688	9325	10077	10794	11692
Motorcycles	4-stroke <250 cm³	Euro I	2000	2003											
Motorcycles	4-stroke <250 cm³	Euro II	2004	2006											
Motorcycles	4-stroke <250 cm³	Euro III	2007	9999											
Motorcycles	4-stroke 250 - 750 cm³	Conventional	0	1999	18923	20165	20737	20996	21657	22611	23892	25645	27712	29683	32152
Motorcycles	4-stroke 250 - 750 cm³	Euro I	2000	2003											
Motorcycles	4-stroke 250 - 750 cm³	Euro II	2004	2006											
Motorcycles	4-stroke 250 - 750 cm³	Euro III	2007	9999											
Motorcycles	4-stroke >750 cm³	Conventional	0	1999	8601	9166	9426	9544	9844	10278	10860	11657	12596	13492	14615
Motorcycles	4-stroke >750 cm³	Euro I	2000	2003											
Motorcycles	4-stroke >750 cm³	Euro II	2004	2006											
Motorcycles	4-stroke >750 cm³	Euro III	2007	9999											

Sector	Subsector	Tech 2	FYear	LYear	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
Passenger Cars	Gasoline 0,8 - 1,4 l	PRE ECE ECE 15/00-01	0	1969	4994	4949	4963	5045	5223	5417	5688	6017	6367	6599	6784	6867	6833
Passenger Cars	Gasoline 0,8 - 1,4 l	ECE 15/02	1970	1978	9402	7791	6441	5527	4770	4352	4057	4054	4021	4051	3997	4014	4024
Passenger Cars	Gasoline 0,8 - 1,4 l	ECE 15/03	1979	1980	5582	4146	3061	2228	1672	1270	1027	853	726	631	566	520	467
Passenger Cars	Gasoline 0,8 - 1,4 l	ECE 15/04	1981	1985	95486	78149	62695	47507	35638	25238	18614	13038	9396	6522	4731	3282	2418
Passenger Cars	Gasoline 0,8 - 1,4 l	Euro I	1986	1990	203911	188827	166452	145685	119763	96437	73954	56825	40791	29907	20894	14633	9806
Passenger Cars	Gasoline 0,8 - 1,4 l	Euro II	1991	1996	212883	211037	207661	203273	197813	189157	177729	161931	144861	127430	107614	87661	67243
Passenger Cars	Gasoline 0,8 - 1,4 l	Euro III	1997	2000	132750	131463	130039	129601	128725	127432	126887	122693	118998	115799	111468	104577	96701
Passenger Cars	Gasoline 0,8 - 1,4 l	Euro IV	2001	2005		20346	43295	64167	93617	135259	136188	133545	132050	131454	129736	127743	125511
Passenger Cars	Gasoline 0,8 - 1,4 l	Euro V	2006	2010							46062	87481	131346	170512	229373	227730	225805
Passenger Cars	Gasoline 0,8 - 1,4 l		2011	2014												67894	155289
Passenger Cars	Gasoline 1,4 - 2,0 l	PRE ECE ECE 15/00-01	0	1969	8291	8215	8200	8321	8638	9068	9590	10256	10933	11397	11659	11779	11667
Passenger Cars	Gasoline 1,4 - 2,0 l	ECE 15/02	1970	1978	11566	9555	7938	6866	5944	5373	5152	5259	5418	5580	5670	5749	5743
Passenger Cars	Gasoline 1,4 - 2,0 l	ECE 15/03	1979	1980	6258	4775	3690	2780	2170	1670	1386	1183	1020	895	801	724	636
Passenger Cars	Gasoline 1,4 - 2,0 l	ECE 15/04	1981	1985	94495	78552	64108	49671	37838	27501	20745	15213	11502	8470	6409	4690	3594
Passenger Cars	Gasoline 1,4 - 2,0 l	Euro I	1986	1990	203365	190772	171667	153308	129614	107639	85477	67960	51214	39588	29267	21797	15650
Passenger Cars	Gasoline 1,4 - 2,0 l	Euro II	1991	1996	402938	402008	397847	391776	383213	370015	348989	317437	286224	256621	220110	181444	140026
Passenger Cars	Gasoline 1,4 - 2,0 l	Euro III	1997	2000	363391	359754	355763	355858	352963	349516	350200	334149	320149	310654	298510	280024	258140
Passenger Cars	Gasoline 1,4 - 2,0 l	Euro IV	2001	2005		51742	107686	149320	197612	252007	258765	251961	248489	247402	243897	238197	233396
Passenger Cars	Gasoline 1,4 - 2,0 l	Euro V	2006	2010							55865	102438	130442	145805	160847	156500	155094
Passenger Cars	Gasoline 1,4 - 2,0 l		2011	2014												15850	25993
Passenger Cars	Gasoline >2,0 l	PRE ECE ECE 15/00-01	0	1969	1024	1079	1128	1237	1391	1600	2061	2629	3223	3590	3776	3921	4420
Passenger Cars	Gasoline >2,0 l	ECE 15/02	1970	1978	1181	1034	936	859	830	841	1031	1313	1734	2009	2238	2457	2858
Passenger Cars	Gasoline >2,0 l	ECE 15/03	1979	1980	520	479	444	399	369	318	311	330	319	297	271	243	268
Passenger Cars	Gasoline >2,0 l	ECE 15/04	1981	1985	1904	1798	1696	1572	1431	1299	1182	1129	1031	935	835	734	743
Passenger Cars	Gasoline >2,0 l	Euro I	1986	1990	4161	4188	4196	4099	3992	3847	3772	3641	3404	3151	2818	2454	2316
Passenger Cars	Gasoline >2,0 l	Euro II	1991	1996	11470	11572	11776	11983	12425	12702	13046	13207	12846	12337	11594	10655	9901
Passenger Cars	Gasoline >2,0 l	Euro III	1997	2000	18870	18780	18760	18987	19329	19850	21109	21230	20991	20706	20140	19257	18272
Passenger Cars	Gasoline >2,0 l	Euro IV	2001	2005		4624	9882	14671	21339	29801	32250	33685	34091	34561	34549	33571	32596
Passenger Cars	Gasoline >2,0 l	Euro V	2006	2010							7894	14607	18442	20013	21352	20826	20873
Passenger Cars	Gasoline >2,0 l		2011	2014												846	1125
Passenger Cars	Diesel 1,4 - 2,0 l	Conventional	0	1990	46384	44480	41523	38006	34340	30088	26004	22021	17989	14348	10977	8074	5859
Passenger Cars	Diesel 1,4 - 2,0 l	Euro I	1991	1996	16471	17245	18106	19220	20895	21613	21522	20504	19072	17688	15816	13503	10845
Passenger Cars	Diesel 1,4 - 2,0 l	Euro II	1997	2000	28716	28688	28633	28948	29194	29373	31180	32603	31362	30703	30387	29751	29009
Passenger Cars	Diesel 1,4 - 2,0 l	Euro III	2001	2005		10980	25186	38421	57411	82415	88649	104131	104503	105907	107353	107313	110697
Passenger Cars	Diesel 1,4 - 2,0 l	Euro IV	2006	2010							27710	68948	93912	103857	111254	110710	112787

Passenger Cars	Diesel 1,4 - 2,0 l	Euro V	2011	2014							3079	12844	38640	66553	119156	179520	226628
Passenger Cars	Diesel >2,0 l	Conventional	0	1990	7728	7120	6345	5723	5039	4460	3895	3402	2908	2516	2095	1656	1376
Passenger Cars	Diesel >2,0 l	Euro I	1991	1996	7698	7640	7463	7353	7287	7147	6944	6585	6018	5575	5022	4380	3823
Passenger Cars	Diesel >2,0 l	Euro II	1997	2000	13125	13236	13141	13296	13564	13885	14771	14907	14386	13960	13559	12861	12557
Passenger Cars	Diesel >2,0 l	Euro III	2001	2005		3884	8635	12968	18885	25776	27913	29844	30102	30677	30829	30074	28854
Passenger Cars	Diesel >2,0 l	Euro IV	2006	2010							6563	15768	18658	20592	22307	22792	23793
Passenger Cars	Diesel >2,0 l	Euro V	2011	2014							729	2860	5908	9868	14513	20115	26047
Passenger Cars	LPG cars	Conventional	0	1990	30	24	17	11	10	10	10	7	8	7	6	6	5
Passenger Cars	LPG cars	Euro I	1991	1996	2	2	3	2	4	4	3	2	2	2	3	2	2
Passenger Cars	LPG cars	Euro II	1997	2000			1	2	1	1	1			1	1	4	4
Passenger Cars	LPG cars	Euro III	2001	2005								1	2	4	3	3	3
Passenger Cars	LPG cars	Euro IV	2006	2010										1	1	4	4
Passenger Cars	2-Stroke	Conventional	0	1999	300	200	150	100	50								
Passenger Cars	Electric cars	Conventional	0	1999	322	301	280	250	211	183	185	188	191	276	350	800	1307
Passenger Cars	Gasoline <0,8 l	PRE ECE ECE 15/00- 01	0	1969							33	65	98	124	142	157	214
Passenger Cars	Gasoline <0,8 l		1970	1978							17	49	71	95	117	127	164
Passenger Cars	Gasoline <0,8 l	ECE 15/02	1979	1980								3	2	3	4	3	4
Passenger Cars	Gasoline <0,8 l	ECE 15/03	1981	1985						1	4	10	13	13	13	15	32
Passenger Cars	Gasoline <0,8 l	ECE 15/04	1986	1990							13	20	27	30	31	33	35
Passenger Cars	Gasoline <0,8 l	Euro I	1991	1996						4	21	38	47	59	60	56	65
Passenger Cars	Gasoline <0,8 l	Euro II	1997	2000	97	96	95	94	94	95	117	141	159	171	185	198	239
Passenger Cars	Gasoline <0,8 l	Euro III	2001	2005		52	118	193	324	553	573	607	626	650	657	685	743
Passenger Cars	Gasoline <0,8 l	Euro IV	2006	2010							257	645	1596	2031	2091	2067	2012
Passenger Cars	Gasoline <0,8 l	Euro V	2011	2014												113	207
Passenger Cars	Diesel <1,4 l	Conventional	0	1990						1	5	8	11	13	11	9	11
Passenger Cars	Diesel <1,4 l	Euro I	1991	1996						3	45	85	96	98	93	92	86
Passenger Cars	Diesel <1,4 l	Euro II	1997	2000	1372	1407	1402	1401	1403	1406	1462	1395	1388	1404	1378	1347	1363
Passenger Cars	Diesel <1,4 l	Euro III	2001	2005		1751	4929	8243	12614	17773	17986	17827	17580	17685	17471	17086	16976
Passenger Cars	Diesel <1,4 l	Euro IV	2006	2010							4746	13769	21404	23476	24508	24289	24022
Passenger Cars	Diesel <1,4 l	Euro V	2011	2014							527	2784	10485	16524	26729	37853	50905
Light Duty Vehicles	Gasoline <3,5t	Conventional	0	1994	27140	23832	21083	18787	16405	14063	11895	9933	7994	6336	4955	3852	3038
Light Duty Vehicles	Gasoline <3,5t	Euro I	1995	1998	17103	16862	16703	16454	16011	15464	14730	13333	12215	11198	10027	8622	7236
Light Duty Vehicles	Gasoline <3,5t	Euro II	1999	2001	9655	14319	14153	14012	13791	13616	13436	10305	9611	8985	8074	6752	5253
Light Duty Vehicles	Gasoline <3,5t	Euro III	2002	2006			3784	8014	13934	20623	26285	19003	18316	17583	15860	13792	11006
Light Duty Vehicles	Gasoline <3,5t	Euro IV	2007	2011								3187	3814	3801	4055	4105	3680
Light Duty Vehicles	Gasoline <3,5t	Euro V	2012	2015													233
Light Duty Vehicles	Diesel <3,5t	Conventional	0	1994	94102	80466	67925	56940	46624	37412	29739	24091	18856	14741	11426	9018	7286

Light Duty Vehicles	Diesel <3,5t	Euro I	1995	1998	74373	72684	71182	69081	66775	63284	58503	52349	46834	41796	36667	31364	26257
Light Duty Vehicles	Diesel <3,5t	Euro II	1999	2001	49951	74831	73532	72069	70326	68384	65694	55259	49908	45261	40307	34072	27892
Light Duty Vehicles	Diesel <3,5t	Euro III	2002	2006			27192	54236	92157	139815	191494	165460	156173	147683	134874	120633	103085
Light Duty Vehicles	Diesel <3,5t	Euro IV	2007	2011								37697	54077	54534	62080	78410	73628
Light Duty Vehicles	Diesel <3,5t	Euro V	2012	2015							2832	11924	20902	21750	34043	32718	50059
Light Duty Vehicles	LPG <3,5t	Conventional	0	1994	60	36	27	21	14	10	9	7	5	4	4	4	4
Light Duty Vehicles	LPG <3,5t	Euro I	1995	1998	1	1											
Light Duty Vehicles	LPG <3,5t	Euro II	1999	2001	1				1	3	3	2	2	3	3	2	2
Light Duty Vehicles	LPG <3,5t	Euro III	2002	2006							5	7	7	8	8	7	7
Light Duty Vehicles	LPG <3,5t	Euro IV	2007	2011								1	3	4	3	4	3
Light Duty Vehicles	Electric <3,5t	Conventional	0	9999	1								1	7	4	17	128
Heavy Duty Vehicles	Gasoline >3,5t	Conventional	0	9999	307	295	291	283	268	287	297	328	325	340	344	347	335
Heavy Duty Vehicles	Diesel RT 3,5 - 7,5t	Conventional	0	1993	3011	2552	2088	1709	1430	1244	1075	937	793	653	540	481	440
Heavy Duty Vehicles	Diesel RT 3,5 - 7,5t	Euro I	1994	1996	834	769	715	656	594	492	434	359	290	234	191	157	141
Heavy Duty Vehicles	Diesel RT 3,5 - 7,5t	Euro II	1997	2001	2136	2254	2161	2078	2003	1901	1728	1509	1250	1060	893	750	642
Heavy Duty Vehicles	Diesel RT 3,5 - 7,5t	Euro III	2002	2006		166	460	755	1049	1437	1674	1661	1576	1450	1315	1209	1100
Heavy Duty Vehicles	Diesel RT 3,5 - 7,5t	Euro IV	2007	2009							52	363	758	911	968	972	941
Heavy Duty Vehicles	Diesel RT 3,5 - 7,5t	Euro V	2010	2013								2	5	27	155	322	496
Heavy Duty Vehicles	Diesel RT 3,5 - 7,5t	Euro VI	2014	9999													1
Heavy Duty Vehicles	Diesel RT 7,5 - 12t	Conventional	0	1993	3782	3406	3069	2766	2503	2241	2084	1901	1683	1419	1250	1128	1030
Heavy Duty Vehicles	Diesel RT 7,5 - 12t	Euro I	1994	1996	1099	1070	1040	985	948	885	828	748	667	545	481	418	359
Heavy Duty Vehicles	Diesel RT 7,5 - 12t	Euro II	1997	2001	1575	1783	1840	1884	1858	1838	1706	1584	1352	1201	1079	951	856
Heavy Duty Vehicles	Diesel RT 7,5 - 12t	Euro III	2002	2006		155	443	713	1061	1501	1949	2011	1924	1798	1631	1529	1390
Heavy Duty Vehicles	Diesel RT 7,5 - 12t	Euro IV	2007	2009		2	2	2	2	3	91	424	824	889	935	937	941
Heavy Duty Vehicles	Diesel RT 7,5 - 12t	Euro V	2010	2013				1	1	1	1	41	181	352	551	724	889
Heavy Duty Vehicles	Diesel RT 7,5 - 12t	Euro VI	2014	9999													1
Heavy Duty Vehicles	Diesel RT 12 - 14 t	Conventional	0	1993	1368	1094	896	734	612	500	441	372	299	228	187	139	112
Heavy Duty Vehicles	Diesel RT 12 - 14 t	Euro I	1994	1996	278	274	248	203	174	152	138	115	100	86	67	57	48
Heavy Duty Vehicles	Diesel RT 12 - 14 t	Euro II	1997	2001	298	312	291	285	278	273	267	243	205	162	142	122	109
Heavy Duty Vehicles	Diesel RT 12 - 14 t	Euro III	2002	2006		10	32	46	58	82	100	109	108	104	95	77	71
Heavy Duty Vehicles	Diesel RT 12 - 14 t	Euro IV	2007	2009					1	1	3	28	52	65	63	58	57
Heavy Duty Vehicles	Diesel RT 12 - 14 t	Euro V	2010	2013									8	11	35	52	57
Heavy Duty Vehicles	Diesel RT 14 - 20t	Conventional	0	1993	2143	1897	1382	1158	1003	884	902	730	536	430	351	289	259
Heavy Duty Vehicles	Diesel RT 14 - 20t	Euro I	1994	1996	905	983	787	701	638	562	571	457	330	242	205	169	150
Heavy Duty Vehicles	Diesel RT 14 - 20t	Euro II	1997	2001	1642	1926	1653	1586	1587	1564	1718	1459	1083	862	733	629	539
Heavy Duty Vehicles	Diesel RT 14 - 20t	Euro III	2002	2006		194	389	665	919	1245	1745	1658	1469	1330	1207	1098	940
Heavy Duty Vehicles	Diesel RT 14 - 20t	Euro IV	2007	2009		4	4	6	7	14	101	457	699	747	757	748	707
Heavy Duty Vehicles	Diesel RT 14 - 20t	Euro V	2010	2013						3	21	106	254	415	572	717	867

Heavy Duty Vehicles	Diesel RT 14 - 20t	Euro VI	2014	9999													2
Heavy Duty Vehicles	Diesel RT 20 - 26t	Conventional	0	1993	2097	1769	1231	984	797	655	626	464	307	217	163	140	114
Heavy Duty Vehicles	Diesel RT 20 - 26t	Euro I	1994	1996	1204	1206	935	815	728	643	649	511	356	267	204	164	129
Heavy Duty Vehicles	Diesel RT 20 - 26t	Euro II	1997	2001	2179	2589	2176	2053	1970	1846	1974	1670	1245	986	838	697	561
Heavy Duty Vehicles	Diesel RT 20 - 26t	Euro III	2002	2006		197	487	803	1143	1583	2274	2161	1907	1748	1590	1407	1188
Heavy Duty Vehicles	Diesel RT 20 - 26t	Euro IV	2007	2009		3	3	3	3	26	128	595	910	988	990	964	942
Heavy Duty Vehicles	Diesel RT 20 - 26t	Euro V	2010	2013						7	24	124	293	493	696	920	1160
Heavy Duty Vehicles	Diesel RT 20 - 26t	Euro VI	2014	9999													1
Heavy Duty Vehicles	Diesel RT 26 - 28t	Conventional	0	1993	4	4	4	4	4	4	4	4	3	2	2	2	1
Heavy Duty Vehicles	Diesel RT 26 - 28t	Euro I	1994	1996	1	2	1	1	1	0	1	1	1	0	1		1
Heavy Duty Vehicles	Diesel RT 26 - 28t	Euro II	1997	2001	3	3	2	2	2	2	2	2	1	1	1	1	1
Heavy Duty Vehicles	Diesel RT 26 - 28t	Euro III	2002	2006				0	2	2	3	3	3	3	3	3	2
Heavy Duty Vehicles	Diesel RT 26 - 28t	Euro IV	2007	2009								3	3	1	2	1	1
Heavy Duty Vehicles	Diesel RT 26 - 28t	Euro V	2010	2013								1	1	1	1	2	3
Heavy Duty Vehicles	Diesel RT 28 - 32t	Conventional	0	1993	185	139	93	70	50	42	36	22	13	9	6	6	5
Heavy Duty Vehicles	Diesel RT 28 - 32t	Euro I	1994	1996	239	241	190	157	134	114	95	67	40	26	20	15	13
Heavy Duty Vehicles	Diesel RT 28 - 32t	Euro II	1997	2001	618	792	670	641	637	639	702	589	439	327	279	231	181
Heavy Duty Vehicles	Diesel RT 28 - 32t	Euro III	2002	2006		82	193	341	509	747	1178	1146	1016	924	873	815	734
Heavy Duty Vehicles	Diesel RT 28 - 32t	Euro IV	2007	2009			0	1	1	21	97	412	619	674	686	677	668
Heavy Duty Vehicles	Diesel RT 28 - 32t	Euro V	2010	2013							10	69	157	255	341	504	723
Heavy Duty Vehicles	Diesel RT 28 - 32t	Euro VI	2014	9999													1
Heavy Duty Vehicles	Diesel RT >32t	Conventional	0	1993	2	2	1	2	2	2	1	1	1				
Heavy Duty Vehicles	Diesel RT >32t	Euro I	1994	1996	0	1	1	1	1	1	1	1	1	1	1	1	1
Heavy Duty Vehicles	Diesel RT >32t	Euro II	1997	2001	1	1	0										
Heavy Duty Vehicles	Diesel RT >32t	Euro III	2002	2006		1	1	2	1	2	3	3	3	3	3	2	3
Heavy Duty Vehicles	Diesel RT >32t	Euro IV	2007	2009								1	1	1			1
Heavy Duty Vehicles	Diesel RT >32t	Euro V	2010	2013										1	2	4	4
Heavy Duty Vehicles	Diesel TT/AT 28 - 34t	Conventional	0	1993	2481	1887	1804	1515	1250	1033	763	659	552	445	365	304	279
Heavy Duty Vehicles	Diesel TT/AT 28 - 34t	Euro I	1994	1996	1025	954	1006	898	781	648	473	404	333	244	207	170	151
Heavy Duty Vehicles	Diesel TT/AT 28 - 34t	Euro II	1997	2001	1862	1872	2119	2035	1942	1802	1414	1281	1087	865	736	633	541
Heavy Duty Vehicles	Diesel TT/AT 28 - 34t	Euro III	2002	2006		188	497	852	1123	1432	1438	1457	1473	1333	1211	1103	946
Heavy Duty Vehicles	Diesel TT/AT 28 - 34t	Euro IV	2007	2009		3	6	8	8	15	83	405	707	754	764	754	709
Heavy Duty Vehicles	Diesel TT/AT 28 - 34t	Euro V	2010	2013						3	17	93	255	416	573	718	870
Heavy Duty Vehicles	Diesel TT/AT 28 - 34t	Euro VI	2014	9999													2
Heavy Duty Vehicles	Diesel TT/AT 34 - 40t	Conventional	0	1993	3173	2250	1980	1585	1255	973	712	583	456	328	253	223	188
Heavy Duty Vehicles	Diesel TT/AT 34 - 40t	Euro I	1994	1996	2884	2100	1834	1472	1214	979	714	597	465	345	271	224	178
Heavy Duty Vehicles	Diesel TT/AT 34 - 40t	Euro II	1997	2001	7098	7055	6586	5636	4638	3653	2749	2277	1781	1351	1128	937	752
Heavy Duty Vehicles	Diesel TT/AT 34 - 40t	Euro III	2002	2006		1009	2342	3625	4439	5378	5569	4881	4150	3380	2811	2234	1802

Heavy Duty Vehicles	Diesel TT/AT 34 - 40t	Euro IV	2007	2009		4	7	6	10	76	214	994	1635	1720	1738	1605	1442
Heavy Duty Vehicles	Diesel TT/AT 34 - 40t	Euro V	2010	2013		1	1	1		27	151	675	1162	1550	2018	2802	3305
Heavy Duty Vehicles	Diesel TT/AT 34 - 40t	Euro VI	2014	9999													7
Heavy Duty Vehicles	Diesel TT/AT 40 - 50t	Conventional	0	1993	1029	708	549	388	287	219	170	123	95	67	61	58	50
Heavy Duty Vehicles	Diesel TT/AT 40 - 50t	Euro I	1994	1996	1215	1060	967	781	616	482	351	285	176	114	101	84	69
Heavy Duty Vehicles	Diesel TT/AT 40 - 50t	Euro II	1997	2001	3548	4062	4016	3731	3293	2841	2242	1791	1225	823	654	538	426
Heavy Duty Vehicles	Diesel TT/AT 40 - 50t	Euro III	2002	2006		552	1706	3011	4472	6217	7578	7024	5985	4772	3954	3229	2562
Heavy Duty Vehicles	Diesel TT/AT 40 - 50t	Euro IV	2007	2009		1	5	6	6	82	340	2129	3557	3680	3845	3624	3280
Heavy Duty Vehicles	Diesel TT/AT 40 - 50t	Euro V	2010	2013		1	2	2	2	1	68	724	1428	1909	2680	3768	4838
Heavy Duty Vehicles	Diesel TT/AT 40 - 50t	Euro VI	2014	9999													16
Heavy Duty Vehicles	Diesel TT/AT 50 - 60t	Euro I	1994	1996	1	1	1	1	1	1							
Heavy Duty Vehicles	Diesel TT/AT 50 - 60t	Euro II	1997	2001	1	1	1	1	1	1	1	1	1	1	1	1	1
Heavy Duty Vehicles	Diesel TT/AT 50 - 60t	Euro IV	2007	2009								1	1	1	1		1
Heavy Duty Vehicles	Diesel TT/AT >60t	Euro III	2002	2006													1
Heavy Duty Vehicles	Diesel TT/AT >60t	Euro V	2010	2013									1	3		1	
Buses	Gasoline Urban Buses	Conventional	0	9999	11	9	7	1	2	2	2	4	7	9	9	10	11
Buses	Diesel Urban Buses <15t	Conventional	0	1993	200	183	154	123	101	80	68	56	49	33	25	16	7
Buses	Diesel Urban Buses <15t	Euro I	1994	1996	118	118	96	106	88	84	75	57	53	28	16	15	10
Buses	Diesel Urban Buses <15t	Euro II	1997	2001	525	542	553	569	535	545	494	427	367	221	117	90	53
Buses	Diesel Urban Buses <15t	Euro III	2002	2006			56	155	248	378	461	439	433	416	363	332	255
Buses	Diesel Urban Buses <15t	Euro IV	2007	2009								119	262	434	425	461	491
Buses	Diesel Urban Buses <15t	Euro V	2010	2013											165	266	397
Buses	Diesel Urban Buses 15 - 18t	Conventional	0	1993	1701	1506	1175	1030	880	758	629	544	460	336	276	217	151
Buses	Diesel Urban Buses 15 - 18t	Euro I	1994	1996	845	810	749	691	620	561	477	399	338	296	253	180	158
Buses	Diesel Urban Buses 15 - 18t	Euro II	1997	2001	1049	1165	1156	1136	1066	1061	1033	1002	919	851	744	636	519
Buses	Diesel Urban Buses 15 - 18t	Euro III	2002	2006			288	456	596	733	1007	1008	989	962	969	951	870
Buses	Diesel Urban Buses 15 - 18t	Euro IV	2007	2009								110	327	624	628	629	628
Buses	Diesel Urban Buses 15 - 18t	Euro V	2010	2013											217	404	562
Buses	Diesel Urban Buses >18t	Conventional	0	1993	37	47	45	25	24	23	16	7	6	6	2	2	1
Buses	Diesel Urban Buses >18t	Euro I	1994	1996	28	44	52	51	42	44	44	23	6	4	2	1	1
Buses	Diesel Urban Buses >18t	Euro II	1997	2001	106	220	225	224	218	217	215	213	161	148	142	105	48
Buses	Diesel Urban Buses >18t	Euro III	2002	2006			135	228	337	388	448	439	414	398	389	377	334
Buses	Diesel Urban Buses >18t	Euro IV	2007	2009								125	247	338	340	333	333
Buses	Diesel Urban Buses >18t	Euro V	2010	2013											97	162	280
Buses	Gasoline Coaches	Conventional	0	9999	1796	1788	1763	1722	1663	1586	1522	1421	1306	1186	1052	913	769
Buses	Diesel Coaches <15t	Conventional	0	1993	4340	3989	3649	3360	3029	2726	2438	2163	1927	1662	1439	1184	996
Buses	Diesel Coaches <15t	Euro I	1994	1996	1079	1053	1031	982	956	920	873	814	733	664	614	545	488
Buses	Diesel Coaches <15t	Euro II	1997	2001	1347	1658	1694	1740	1908	2023	2162	2144	2077	2011	1914	1801	1689

Buses	Diesel Coaches <15t	Euro III	2002	2006			253	482	751	1052	1361	1423	1439	1461	1454	1365	1215
Buses	Diesel Coaches <15t	Euro IV	2007	2009								228	480	793	822	798	742
Buses	Diesel Coaches <15t	Euro V	2010	2013											204	328	382
Buses	Diesel Coaches 15 - 18t	Conventional	0	1993	1133	1061	1013	957	914	847	768	693	609	540	463	377	319
Buses	Diesel Coaches 15 - 18t	Euro I	1994	1996	148	161	173	176	176	184	176	177	178	193	179	154	135
Buses	Diesel Coaches 15 - 18t	Euro II	1997	2001	173	208	221	220	230	240	237	236	226	245	258	267	284
Buses	Diesel Coaches 15 - 18t	Euro III	2002	2006			19	46	61	71	90	81	99	106	107	109	119
Buses	Diesel Coaches 15 - 18t	Euro IV	2007	2009								11	38	69	66	65	64
Buses	Diesel Coaches 15 - 18t	Euro V	2010	2013											41	48	59
Buses	Diesel Coaches >18t	Conventional	0	1993	192	177	157	142	138	121	92	77	56	48	38	31	26
Buses	Diesel Coaches >18t	Euro I	1994	1996	78	76	79	74	70	65	60	56	49	46	36	26	20
Buses	Diesel Coaches >18t	Euro II	1997	2001	145	190	196	201	192	192	202	199	173	165	156	141	134
Buses	Diesel Coaches >18t	Euro III	2002	2006			32	92	152	230	293	302	312	321	322	309	322
Buses	Diesel Coaches >18t	Euro IV	2007	2009								55	114	180	194	197	193
Buses	Diesel Coaches >18t	Euro V	2010	2013											39	70	112
Mopeds	2-stroke <50 cm ³	Conventional	0	1999	143569	136233	128203	120288	112245	103814	95123	86611	78807	71061	63625	56546	50013
Mopeds	2-stroke <50 cm ³	Euro I	2000	2003	16403	28734	42762	48678	46056	43440	40735	37815	35222	32562	29999	27566	25345
Mopeds	2-stroke <50 cm ³	Euro II	2004	2014					7996	16278	24801	33341	38136	42179	44312	46905	53316
Mopeds	4-stroke <50 cm ³	Euro II	2004	2014					2665	5426	8267	11114	12712	14060	14771	15635	17772
Motorcycles	2-stroke >50 cm ³	Conventional	0	1999	10536	10539	10440	10400	10480	10665	10952	11137	11004	10594	10060	9465	8930
Motorcycles	2-stroke >50 cm ³	Euro I	2000	2003	464	779	1097	1409	1356	1339	1361	1368	1357	1308	1250	1190	1136
Motorcycles	2-stroke >50 cm ³	Euro II	2004	2006					450	1115	1957	1864	1769	1655	1543	1425	1326
Motorcycles	2-stroke >50 cm ³	Euro III	2007	9999								1042	1673	1927	2033	2060	2049
Motorcycles	4-stroke <250 cm ³	Conventional	0	1999	11941	12429	12827	13327	14028	14931	16063	17142	17806	18061	18108	18033	18059
Motorcycles	4-stroke <250 cm ³	Euro I	2000	2003	526	918	1348	1805	1814	1874	1996	2106	2196	2230	2250	2267	2297
Motorcycles	4-stroke <250 cm ³	Euro II	2004	2006					603	1560	2870	2870	2863	2822	2777	2715	2682
Motorcycles	4-stroke <250 cm ³	Euro III	2007	9999								1604	2707	3285	3659	3924	4143
Motorcycles	4-stroke 250 - 750 cm ³	Conventional	0	1999	32838	34180	35273	36650	38576	41061	44174	47140	48967	49667	49797	49591	49662
Motorcycles	4-stroke 250 - 750 cm ³	Euro I	2000	2003	1447	2525	3707	4964	4990	5154	5489	5792	6038	6133	6188	6234	6318
Motorcycles	4-stroke 250 - 750 cm ³	Euro II	2004	2006					1657	4291	7892	7891	7873	7761	7638	7467	7376
Motorcycles	4-stroke 250 - 750 cm ³	Euro III	2007	9999								4410	7445	9033	10061	10791	11394
Motorcycles	4-stroke >750 cm ³	Conventional	0	1999	14926	15536	16033	16659	17535	18664	20079	21427	22258	22576	22635	22541	22574
Motorcycles	4-stroke >750 cm ³	Euro I	2000	2003	658	1148	1685	2257	2268	2343	2495	2633	2745	2788	2813	2834	2872
Motorcycles	4-stroke >750 cm ³	Euro II	2004	2006					753	1950	3587	3587	3578	3528	3472	3394	3353
Motorcycles	4-stroke >750 cm ³	Euro III	2007	9999								2005	3384	4106	4573	4905	5179

Annex 3B-2 Mileage data 1985-2012 for road transport (km)

Sector	Subsector	Tech 2	FYear	LYear	1985	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
Passenger Cars	Gasoline 0,8 - 1,4 l	PRE ECE	0	1969	10987	10126	10512	10813	10717	10998	10369	9770	9516	9206	8738
Passenger Cars	Gasoline 0,8 - 1,4 l	ECE 15/00-01	1970	1978	13512	12672	13276	13818	13785	14238	13441	12686	12361	11935	11315
Passenger Cars	Gasoline 0,8 - 1,4 l	ECE 15/02	1979	1980	15074	14174	14709	15129	15002	15478	14621	13813	13487	13071	12472
Passenger Cars	Gasoline 0,8 - 1,4 l	ECE 15/03	1981	1985	15016	15385	16091	16689	16643	17192	16255	15377	15047	14633	13973
Passenger Cars	Gasoline 0,8 - 1,4 l	ECE 15/04	1986	1990		15922	18158	18617	18423	18995	17947	16963	16578	16099	15347
Passenger Cars	Gasoline 0,8 - 1,4 l	Euro I	1991	1996			9822	15599	17451	17935	18227	17893	19347	18764	17873
Passenger Cars	Gasoline 0,8 - 1,4 l	Euro II	1997	2000									10518	15722	16679
Passenger Cars	Gasoline 0,8 - 1,4 l	Euro III	2001	2005											
Passenger Cars	Gasoline 0,8 - 1,4 l	Euro IV	2006	2010											
Passenger Cars	Gasoline 0,8 - 1,4 l	Euro V	2011	2014											
Passenger Cars	Gasoline 1,4 - 2,0 l	PRE ECE	0	1969	13245	12360	12855	13288	13221	13566	12782	12036	11712	11322	10737
Passenger Cars	Gasoline 1,4 - 2,0 l	ECE 15/00-01	1970	1978	16412	15390	16109	16770	16715	17277	16317	15407	15023	14509	13761
Passenger Cars	Gasoline 1,4 - 2,0 l	ECE 15/02	1979	1980	18351	17283	17934	18482	18333	18919	17876	16896	16503	15993	15257
Passenger Cars	Gasoline 1,4 - 2,0 l	ECE 15/03	1981	1985	17890	18791	19630	20355	20297	20965	19820	18752	18345	17823	16991
Passenger Cars	Gasoline 1,4 - 2,0 l	ECE 15/04	1986	1990		19005	22385	22891	22603	23307	22019	20810	20338	19743	18813
Passenger Cars	Gasoline 1,4 - 2,0 l	Euro I	1991	1996			12083	18712	20806	21397	21850	21413	23821	23096	21991
Passenger Cars	Gasoline 1,4 - 2,0 l	Euro II	1997	2000									12863	18602	20691
Passenger Cars	Gasoline 1,4 - 2,0 l	Euro III	2001	2005											
Passenger Cars	Gasoline 1,4 - 2,0 l	Euro IV	2006	2010											
Passenger Cars	Gasoline 1,4 - 2,0 l	Euro V	2011	2014											
Passenger Cars	Gasoline >2,0 l	PRE ECE	0	1969	14692	13694	14191	14612	14494	14941	14084	13263	12911	12478	11825
Passenger Cars	Gasoline >2,0 l	ECE 15/00-01	1970	1978	17959	16981	17820	18516	18543	19119	18039	17008	16558	16029	15211
Passenger Cars	Gasoline >2,0 l	ECE 15/02	1979	1980	20799	19449	20135	20768	20669	21331	20133	19014	18551	17943	17090
Passenger Cars	Gasoline >2,0 l	ECE 15/03	1981	1985	20329	21465	22268	22895	22702	23415	22104	20877	20384	19776	18841
Passenger Cars	Gasoline >2,0 l	ECE 15/04	1986	1990		21408	25931	26252	25851	26645	25127	23708	23150	22469	21401
Passenger Cars	Gasoline >2,0 l	Euro I	1991	1996			14128	21423	23687	24914	24459	24235	26905	26023	24747
Passenger Cars	Gasoline >2,0 l	Euro II	1997	2000									14567	20725	22239
Passenger Cars	Gasoline >2,0 l	Euro III	2001	2005											
Passenger Cars	Gasoline >2,0 l	Euro IV	2006	2010											
Passenger Cars	Gasoline >2,0 l	Euro V	2011	2014											
Passenger Cars	Diesel 1,4 - 2,0 l	Conventional	0	1990	35362	42261	43428	41168	40097	42647	40813	39720	39124	37694	36387
Passenger Cars	Diesel 1,4 - 2,0 l	Euro I	1991	1996			67316	85071	76115	63947	57531	55204	56392	48899	44835
Passenger Cars	Diesel 1,4 - 2,0 l	Euro II	1997	2000									35632	44754	42660
Passenger Cars	Diesel 1,4 - 2,0 l	Euro III	2001	2005											

Passenger Cars	Diesel 1,4 - 2,0 l	Euro IV	2006	2010											
Passenger Cars	Diesel 1,4 - 2,0 l	Euro V	2011	2014											
Passenger Cars	Diesel >2,0 l	Conventional	0	1990	46795	53584	53600	45969	41568	42757	40159	38685	37779	36369	35252
Passenger Cars	Diesel >2,0 l	Euro I	1991	1996			75908	100414	95448	94680	87668	79953	77912	58365	47751
Passenger Cars	Diesel >2,0 l	Euro II	1997	2000									62706	82066	72553
Passenger Cars	Diesel >2,0 l	Euro III	2001	2005											
Passenger Cars	Diesel >2,0 l	Euro IV	2006	2010											
Passenger Cars	Diesel >2,0 l	Euro V	2011	2014											
Passenger Cars	LPG cars	Conventional	0	1990	25575	26636	28202	29712	30233	31206	29655	28181	27555	26612	25314
Passenger Cars	LPG cars	Euro I	1991	1996				38174	37417	38309	37546	35227	34476	33513	33997
Passenger Cars	LPG cars	Euro II	1997	2000											
Passenger Cars	LPG cars	Euro III	2001	2005											
Passenger Cars	LPG cars	Euro IV	2006	2010											
Passenger Cars	2-Stroke	Conventional	0	9999	14642	15402	16008	16500	16419	16953	16011	15123	14770	14328	13647
Passenger Cars	Electric cars	Conventional	0	9999	10260	10737	11377	11996	12218	15192	14786	13930	13775	13126	13594
Passenger Cars	Gasoline <0,8 l	PRE ECE	0	1969											
Passenger Cars	Gasoline <0,8 l	ECE 15/00-01	1970	1978											
Passenger Cars	Gasoline <0,8 l	ECE 15/02	1979	1980											
Passenger Cars	Gasoline <0,8 l	ECE 15/03	1981	1985											
Passenger Cars	Gasoline <0,8 l	ECE 15/04	1986	1990											
Passenger Cars	Gasoline <0,8 l	Euro I	1991	1996											
Passenger Cars	Gasoline <0,8 l	Euro II	1997	2000											10225
Passenger Cars	Gasoline <0,8 l	Euro III	2001	2005											
Passenger Cars	Gasoline <0,8 l	Euro IV	2006	2010											
Passenger Cars	Gasoline <0,8 l	Euro V	2011	2014											
Passenger Cars	Diesel <1,4 l	Conventional	0	1990											
Passenger Cars	Diesel <1,4 l	Euro I	1991	1996											
Passenger Cars	Diesel <1,4 l	Euro II	1997	2000											23322
Passenger Cars	Diesel <1,4 l	Euro III	2001	2005											
Passenger Cars	Diesel <1,4 l	Euro IV	2006	2010											
Passenger Cars	Diesel <1,4 l	Euro V	2011	2014											
Light Duty Vehicles	Gasoline <3,5t	Conventional	0	1994	16590	15777	16152	16764	16864	17510	17508	16793	16443	16079	15437
Light Duty Vehicles	Gasoline <3,5t	Euro I	1995	1998							11236	16067	17035	17363	19669
Light Duty Vehicles	Gasoline <3,5t	Euro II	1999	2001											10620
Light Duty Vehicles	Gasoline <3,5t	Euro III	2002	2006											
Light Duty Vehicles	Gasoline <3,5t	Euro IV	2007	2011											
Light Duty Vehicles	Gasoline <3,5t	Euro V	2012	2015											
Light Duty Vehicles	Diesel <3,5t	Conventional	0	1994	29423	32398	32514	31159	30456	31409	31302	30275	29362	28039	26954

Light Duty Vehicles	Diesel <3,5t	Euro I	1995	1998							20709	29650	31926	32950	36390
Light Duty Vehicles	Diesel <3,5t	Euro II	1999	2001											20043
Light Duty Vehicles	Diesel <3,5t	Euro III	2002	2006											
Light Duty Vehicles	Diesel <3,5t	Euro IV	2007	2011											
Light Duty Vehicles	Diesel <3,5t	Euro V	2012	2015											
Light Duty Vehicles	LPG <3,5t	Conventional	0	1994	20338	19332	19790	20538	20660	20626	19555	18924	18421	17853	16944
Light Duty Vehicles	LPG <3,5t	Euro I	1995	1998										16428	30783
Light Duty Vehicles	LPG <3,5t	Euro II	1999	2001											
Light Duty Vehicles	LPG <3,5t	Euro III	2002	2006											
Light Duty Vehicles	LPG <3,5t	Euro IV	2007	2011											
Light Duty Vehicles	Electric <3,5t	Conventional	0	9999	12109	11522	11796	12244	12317	12509	11218	10676	12192	11836	11262
Heavy Duty Vehicles	Gasoline >3,5t	Conventional	0	9999	20176	18562	19950	20541	19045	18224	18229	17308	16292	15404	15444
Heavy Duty Vehicles	Diesel RT 3,5 - 7,5t	Conventional	0	1993	31593	30572	33046	29469	24828	23910	25188	22923	20954	19357	17443
Heavy Duty Vehicles	Diesel RT 3,5 - 7,5t	Euro I	1994	1996					23054	26299	36682	37885	39907	36293	32356
Heavy Duty Vehicles	Diesel RT 3,5 - 7,5t	Euro II	1997	2001								24587	27344	31676	34883
Heavy Duty Vehicles	Diesel RT 3,5 - 7,5t	Euro III	2002	2006											
Heavy Duty Vehicles	Diesel RT 3,5 - 7,5t	Euro IV	2007	2009											
Heavy Duty Vehicles	Diesel RT 3,5 - 7,5t	Euro V	2010	2013											
Heavy Duty Vehicles	Diesel RT 3,5 - 7,5t	Euro VI	2014	9999											
Heavy Duty Vehicles	Diesel RT 7,5 - 12t	Conventional	0	1993	37315	36012	38915	34698	29224	28272	29993	27218	24915	23136	20843
Heavy Duty Vehicles	Diesel RT 7,5 - 12t	Euro I	1994	1996					27938	31807	42654	45806	48666	44243	39322
Heavy Duty Vehicles	Diesel RT 7,5 - 12t	Euro II	1997	2001								29796	34383	41420	41778
Heavy Duty Vehicles	Diesel RT 7,5 - 12t	Euro III	2002	2006											
Heavy Duty Vehicles	Diesel RT 7,5 - 12t	Euro IV	2007	2009											
Heavy Duty Vehicles	Diesel RT 7,5 - 12t	Euro V	2010	2013											
Heavy Duty Vehicles	Diesel RT 7,5 - 12t	Euro VI	2014	9999											
Heavy Duty Vehicles	Diesel RT 12 - 14 t	Conventional	0	1993	27513	30299	32856	31321	26652	28985	28652	23167	21675	21127	22644
Heavy Duty Vehicles	Diesel RT 12 - 14 t	Euro I	1994	1996					23844	32017	43687	39124	40920	39134	41056
Heavy Duty Vehicles	Diesel RT 12 - 14 t	Euro II	1997	2001								24863	28125	38236	47872
Heavy Duty Vehicles	Diesel RT 12 - 14 t	Euro III	2002	2006											
Heavy Duty Vehicles	Diesel RT 12 - 14 t	Euro IV	2007	2009											
Heavy Duty Vehicles	Diesel RT 12 - 14 t	Euro V	2010	2013											
Heavy Duty Vehicles	Diesel RT 14 - 20t	Conventional	0	1993	49555	54586	59192	56427	48056	53001	52789	42730	40125	39183	41634
Heavy Duty Vehicles	Diesel RT 14 - 20t	Euro I	1994	1996					41509	54202	69364	66546	72839	69704	72988
Heavy Duty Vehicles	Diesel RT 14 - 20t	Euro II	1997	2001								43283	49885	65693	77120
Heavy Duty Vehicles	Diesel RT 14 - 20t	Euro III	2002	2006											
Heavy Duty Vehicles	Diesel RT 14 - 20t	Euro IV	2007	2009											
Heavy Duty Vehicles	Diesel RT 14 - 20t	Euro V	2010	2013											

Heavy Duty Vehicles	Diesel RT 14 - 20t	Euro VI	2014	9999											
Heavy Duty Vehicles	Diesel RT 20 - 26t	Conventional	0	1993	77104	85021	80265	84890	81445	80397	77249	73888	64558	62670	54436
Heavy Duty Vehicles	Diesel RT 20 - 26t	Euro I	1994	1996					67784	77139	97621	109399	112920	107957	92590
Heavy Duty Vehicles	Diesel RT 20 - 26t	Euro II	1997	2001								71827	78066	100546	97933
Heavy Duty Vehicles	Diesel RT 20 - 26t	Euro III	2002	2006											
Heavy Duty Vehicles	Diesel RT 20 - 26t	Euro IV	2007	2009											
Heavy Duty Vehicles	Diesel RT 20 - 26t	Euro V	2010	2013											
Heavy Duty Vehicles	Diesel RT 20 - 26t	Euro VI	2014	9999											
Heavy Duty Vehicles	Diesel RT 26 - 28t	Conventional	0	1993	74213	81865	77431	81805	78420	81812	79350	79054	67611	64481	54746
Heavy Duty Vehicles	Diesel RT 26 - 28t	Euro I	1994	1996							121173	94707	117422	111012	95023
Heavy Duty Vehicles	Diesel RT 26 - 28t	Euro II	1997	2001								72706	86469	88492	100154
Heavy Duty Vehicles	Diesel RT 26 - 28t	Euro III	2002	2006											
Heavy Duty Vehicles	Diesel RT 26 - 28t	Euro IV	2007	2009											
Heavy Duty Vehicles	Diesel RT 26 - 28t	Euro V	2010	2013											
Heavy Duty Vehicles	Diesel RT 28 - 32t	Conventional	0	1993	85446	94256	89150	94186	91028	93121	88668	84213	73749	72366	63114
Heavy Duty Vehicles	Diesel RT 28 - 32t	Euro I	1994	1996					69026	80737	95841	108625	114746	109433	93722
Heavy Duty Vehicles	Diesel RT 28 - 32t	Euro II	1997	2001								72706	78416	98957	96107
Heavy Duty Vehicles	Diesel RT 28 - 32t	Euro III	2002	2006											
Heavy Duty Vehicles	Diesel RT 28 - 32t	Euro IV	2007	2009											
Heavy Duty Vehicles	Diesel RT 28 - 32t	Euro V	2010	2013											
Heavy Duty Vehicles	Diesel RT 28 - 32t	Euro VI	2014	9999											
Heavy Duty Vehicles	Diesel RT >32t	Conventional	0	1993	56752	62603	59212	62557	59968	56125	80742	49186	42044	75528	64649
Heavy Duty Vehicles	Diesel RT >32t	Euro I	1994	1996							71581	107606	120397	114825	89619
Heavy Duty Vehicles	Diesel RT >32t	Euro II	1997	2001								72706	86871	118546	65155
Heavy Duty Vehicles	Diesel RT >32t	Euro III	2002	2006											
Heavy Duty Vehicles	Diesel RT >32t	Euro IV	2007	2009											
Heavy Duty Vehicles	Diesel RT >32t	Euro V	2010	2013											
Heavy Duty Vehicles	Diesel TT/AT 28 - 34t	Conventional	0	1993	53602	59453	64508	61415	52944	58371	58054	47394	44784	43638	46146
Heavy Duty Vehicles	Diesel TT/AT 28 - 34t	Euro I	1994	1996					45559	59585	76332	73342	80388	77115	80613
Heavy Duty Vehicles	Diesel TT/AT 28 - 34t	Euro II	1997	2001								47629	54989	72613	85120
Heavy Duty Vehicles	Diesel TT/AT 28 - 34t	Euro III	2002	2006											
Heavy Duty Vehicles	Diesel TT/AT 28 - 34t	Euro IV	2007	2009											
Heavy Duty Vehicles	Diesel TT/AT 28 - 34t	Euro V	2010	2013											
Heavy Duty Vehicles	Diesel TT/AT 28 - 34t	Euro VI	2014	9999											
Heavy Duty Vehicles	Diesel TT/AT 34 - 40t	Conventional	0	1993	84648	90436	86534	82989	83916	86825	79943	79896	73919	69127	62201
Heavy Duty Vehicles	Diesel TT/AT 34 - 40t	Euro I	1994	1996					61500	71410	83419	102475	117340	106705	99631
Heavy Duty Vehicles	Diesel TT/AT 34 - 40t	Euro II	1997	2001								66415	81057	94399	102498
Heavy Duty Vehicles	Diesel TT/AT 34 - 40t	Euro III	2002	2006											

Heavy Duty Vehicles	Diesel TT/AT 34 - 40t	Euro IV	2007	2009											
Heavy Duty Vehicles	Diesel TT/AT 34 - 40t	Euro V	2010	2013											
Heavy Duty Vehicles	Diesel TT/AT 34 - 40t	Euro VI	2014	9999											
Heavy Duty Vehicles	Diesel TT/AT 40 - 50t	Conventional	0	1993	109789	114598	111039	100652	104396	108389	95165	94197	91124	81657	76385
Heavy Duty Vehicles	Diesel TT/AT 40 - 50t	Euro I	1994	1996					79563	92663	99809	121225	138856	122156	112618
Heavy Duty Vehicles	Diesel TT/AT 40 - 50t	Euro II	1997	2001								79383	92909	107228	111665
Heavy Duty Vehicles	Diesel TT/AT 40 - 50t	Euro III	2002	2006											
Heavy Duty Vehicles	Diesel TT/AT 40 - 50t	Euro IV	2007	2009											
Heavy Duty Vehicles	Diesel TT/AT 40 - 50t	Euro V	2010	2013											
Heavy Duty Vehicles	Diesel TT/AT 40 - 50t	Euro VI	2014	9999											
Heavy Duty Vehicles	Diesel TT/AT 50 - 60t	Euro I	1994	1996								138231	135435	116422	110616
Heavy Duty Vehicles	Diesel TT/AT 50 - 60t	Euro II	1997	2001										72370	137521
Heavy Duty Vehicles	Diesel TT/AT 50 - 60t	Euro IV	2007	2009											
Heavy Duty Vehicles	Diesel TT/AT >60t	Euro III	2002	2006											
Heavy Duty Vehicles	Diesel TT/AT >60t	Euro V	2010	2013											
Buses	Gasoline Urban Buses	Conventional	0	9999	29543	28977	24021	21995	17353	26031	23506	19551	21801	20975	18284
Buses	Diesel Urban Buses <15t	Conventional	0	1993	183187	207698	168762	142671	109361	117960	108353	102142	95506	91310	81145
Buses	Diesel Urban Buses <15t	Euro I	1994	1996						96215	142149	144219	157743	147146	129841
Buses	Diesel Urban Buses <15t	Euro II	1997	2001									91225	119788	134162
Buses	Diesel Urban Buses <15t	Euro III	2002	2006											
Buses	Diesel Urban Buses <15t	Euro IV	2007	2009											
Buses	Diesel Urban Buses <15t	Euro V	2010	2013											
Buses	Diesel Urban Buses 15 - 18t	Conventional	0	1993	173862	197367	160377	135588	103937	112696	103596	99477	92134	87205	79413
Buses	Diesel Urban Buses 15 - 18t	Euro I	1994	1996						96215	142765	138879	157646	147436	129279
Buses	Diesel Urban Buses 15 - 18t	Euro II	1997	2001									91225	117036	134971
Buses	Diesel Urban Buses 15 - 18t	Euro III	2002	2006											
Buses	Diesel Urban Buses 15 - 18t	Euro IV	2007	2009											
Buses	Diesel Urban Buses 15 - 18t	Euro V	2010	2013											
Buses	Diesel Urban Buses >18t	Conventional	0	1993	226400	257009	208841	176562	135345	148089	129169	116874	115488	93970	71929
Buses	Diesel Urban Buses >18t	Euro I	1994	1996						96215	172711	115545	162038	150495	138217
Buses	Diesel Urban Buses >18t	Euro II	1997	2001										91104	108084
Buses	Diesel Urban Buses >18t	Euro III	2002	2006											
Buses	Diesel Urban Buses >18t	Euro IV	2007	2009											
Buses	Diesel Urban Buses >18t	Euro V	2010	2013											
Buses	Gasoline Coaches	Conventional	0	9999	16804	17038	14174	13035	12050	13129	16303	18243	17792	17183	16236
Buses	Diesel Coaches <15t	Conventional	0	1993	32231	37854	30875	26219	23552	24864	28995	33414	32601	31028	29777
Buses	Diesel Coaches <15t	Euro I	1994	1996						16383	30705	36060	43701	40996	38739
Buses	Diesel Coaches <15t	Euro II	1997	2001									24161	34986	37030

Buses	Diesel Coaches <15t	Euro III	2002	2006											
Buses	Diesel Coaches <15t	Euro IV	2007	2009											
Buses	Diesel Coaches <15t	Euro V	2010	2013											
Buses	Diesel Coaches 15 - 18t	Conventional	0	1993	46554	54705	44620	37893	34040	32112	38024	44216	43217	41123	39645
Buses	Diesel Coaches 15 - 18t	Euro I	1994	1996						26952	45374	59695	72100	67579	63928
Buses	Diesel Coaches 15 - 18t	Euro II	1997	2001									39748	59444	60740
Buses	Diesel Coaches 15 - 18t	Euro III	2002	2006											
Buses	Diesel Coaches 15 - 18t	Euro IV	2007	2009											
Buses	Diesel Coaches 15 - 18t	Euro V	2010	2013											
Buses	Diesel Coaches >18t	Conventional	0	1993	92397	108585	88569	75217	67569	66502	77292	89552	86760	82331	77766
Buses	Diesel Coaches >18t	Euro I	1994	1996						42878	72260	92808	115142	107985	102154
Buses	Diesel Coaches >18t	Euro II	1997	2001									63234	87431	91335
Buses	Diesel Coaches >18t	Euro III	2002	2006											
Buses	Diesel Coaches >18t	Euro IV	2007	2009											
Buses	Diesel Coaches >18t	Euro V	2010	2013											
Mopeds	2-stroke <50 cm³	Conventional	0	1999	1416	1341	1400	1469	1509	1544	1599	1711	1963	2203	2000
Mopeds	2-stroke <50 cm³	Euro I	2000	2003											
Mopeds	2-stroke <50 cm³	Euro II	2004	2014											
Mopeds	4-stroke <50 cm³	Euro II	2004	2014											
Motorcycles	2-stroke >50 cm³	Conventional	0	1999	6850	6697	6704	7068	7252	7410	7244	6880	6680	6463	6078
Motorcycles	2-stroke >50 cm³	Euro I	2000	2003											
Motorcycles	2-stroke >50 cm³	Euro II	2004	2006											
Motorcycles	2-stroke >50 cm³	Euro III	2007	9999											
Motorcycles	4-stroke <250 cm³	Conventional	0	1999	6850	6697	6704	7068	7252	7410	7244	6880	6680	6463	6078
Motorcycles	4-stroke <250 cm³	Euro I	2000	2003											
Motorcycles	4-stroke <250 cm³	Euro II	2004	2006											
Motorcycles	4-stroke <250 cm³	Euro III	2007	9999											
Motorcycles	4-stroke 250 - 750 cm³	Conventional	0	1999	6850	6697	6704	7068	7252	7410	7244	6880	6680	6463	6078
Motorcycles	4-stroke 250 - 750 cm³	Euro I	2000	2003											
Motorcycles	4-stroke 250 - 750 cm³	Euro II	2004	2006											
Motorcycles	4-stroke 250 - 750 cm³	Euro III	2007	9999											
Motorcycles	4-stroke >750 cm³	Conventional	0	1999	6850	6697	6704	7068	7252	7410	7244	6880	6680	6463	6078
Motorcycles	4-stroke >750 cm³	Euro I	2000	2003											
Motorcycles	4-stroke >750 cm³	Euro II	2004	2006											
Motorcycles	4-stroke >750 cm³	Euro III	2007	9999											

Sector	Subsector	Tech 2	FYear	LYear	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
Passenger Cars	Gasoline 0,8 - 1,4 l	PRE ECE	0	1969	8322	7869	7768	7593	7412	6918	6482	6389	6055	5776	5467	5292	5079
Passenger Cars	Gasoline 0,8 - 1,4 l	ECE 15/00-01	1970	1978	10751	10160	9981	9712	9418	8743	8139	7937	7514	7149	6739	6508	6236
Passenger Cars	Gasoline 0,8 - 1,4 l	ECE 15/02	1979	1980	11910	11293	11141	10904	10631	9932	9290	9135	8680	8283	7831	7578	7270
Passenger Cars	Gasoline 0,8 - 1,4 l	ECE 15/03	1981	1985	13387	12812	12661	12392	12110	11321	10629	10256	9906	9441	8897	8561	8176
Passenger Cars	Gasoline 0,8 - 1,4 l	ECE 15/04	1986	1990	14690	14012	13868	13584	13296	12442	11689	11413	10947	10460	9896	9569	9151
Passenger Cars	Gasoline 0,8 - 1,4 l	Euro I	1991	1996	17076	16211	16019	15672	15300	14294	13402	13193	12561	12020	11395	11056	10627
Passenger Cars	Gasoline 0,8 - 1,4 l	Euro II	1997	2000	17174	18159	17948	17556	17139	15991	14975	14782	13990	13351	12636	12241	11765
Passenger Cars	Gasoline 0,8 - 1,4 l	Euro III	2001	2005		9693	14122	15944	15903	15178	16977	16790	15878	15149	14338	13881	13337
Passenger Cars	Gasoline 0,8 - 1,4 l	Euro IV	2006	2010							9067	13744	14370	14743	13940	15645	15027
Passenger Cars	Gasoline 0,8 - 1,4 l	Euro V	2011	2014												8406	11701
Passenger Cars	Gasoline 1,4 - 2,0 l	PRE ECE	0	1969	10219	9649	9528	9313	9093	8492	7957	7858	7447	7107	6726	6511	6250
Passenger Cars	Gasoline 1,4 - 2,0 l	ECE 15/00-01	1970	1978	13080	12344	12131	11809	11459	10629	9884	9667	9111	8667	8177	7902	7584
Passenger Cars	Gasoline 1,4 - 2,0 l	ECE 15/02	1979	1980	14571	13788	13612	13324	12995	12132	11351	11194	10601	10119	9567	9265	8886
Passenger Cars	Gasoline 1,4 - 2,0 l	ECE 15/03	1981	1985	16262	15470	15291	14950	14592	13613	12754	12461	11886	11331	10708	10340	9916
Passenger Cars	Gasoline 1,4 - 2,0 l	ECE 15/04	1986	1990	17995	17102	16929	16583	16222	15170	14226	13981	13302	12701	12012	11628	11151
Passenger Cars	Gasoline 1,4 - 2,0 l	Euro I	1991	1996	21008	19934	19695	19267	18807	17565	16465	16213	15419	14746	13974	13551	13019
Passenger Cars	Gasoline 1,4 - 2,0 l	Euro II	1997	2000	21342	22231	21965	21480	20967	19565	18328	18051	17112	16331	15454	14966	14373
Passenger Cars	Gasoline 1,4 - 2,0 l	Euro III	2001	2005		11898	17414	20033	20194	19474	20634	20321	19273	18394	17403	16838	16142
Passenger Cars	Gasoline 1,4 - 2,0 l	Euro IV	2006	2010													
Passenger Cars	Gasoline 1,4 - 2,0 l	Euro V	2011	2014							11129	16919	18656	19084	18251	18770	18000
Passenger Cars	Gasoline >2,0 l	PRE ECE	0	1969	11255	10611	10487	10283	10090	9461	8916	8848	8398	8024	7583	7331	7052
Passenger Cars	Gasoline >2,0 l	ECE 15/00-01	1970	1978	14493	13697	13480	13156	12854	11899	10999	10747	10097	9604	9083	8812	8499
Passenger Cars	Gasoline >2,0 l	ECE 15/02	1979	1980	16275	15348	15169	14795	14485	13529	12670	12530	11863	11313	10701	10396	9993
Passenger Cars	Gasoline >2,0 l	ECE 15/03	1981	1985	17994	16987	16805	16432	16038	14958	13996	13837	13074	12465	11786	11400	10935
Passenger Cars	Gasoline >2,0 l	ECE 15/04	1986	1990	20447	19314	19091	18686	18256	17041	15954	15774	14926	14237	13488	13052	12559
Passenger Cars	Gasoline >2,0 l	Euro I	1991	1996	23618	22334	22050	21557	21047	19647	18397	18133	17178	16391	15513	15016	14420
Passenger Cars	Gasoline >2,0 l	Euro II	1997	2000	23100	25204	24880	24318	23732	22144	20744	20336	19307	18424	17425	16853	16145
Passenger Cars	Gasoline >2,0 l	Euro III	2001	2005		13431	19503	21913	22034	21459	23318	22717	21635	20658	19538	18882	18104
Passenger Cars	Gasoline >2,0 l	Euro IV	2006	2010							12528	19167	21128	21833	20987	20999	20164
Passenger Cars	Gasoline >2,0 l	Euro V	2011	2014												11573	18688
Passenger Cars	Diesel 1,4 - 2,0 l	Conventional	0	1990	33586	30342	29663	30687	30638	28016	26119	21907	23444	21533	20732	20558	19115
Passenger Cars	Diesel 1,4 - 2,0 l	Euro I	1991	1996	40694	36121	35088	36087	35846	32663	30053	26974	27319	24898	24043	23919	22366
Passenger Cars	Diesel 1,4 - 2,0 l	Euro II	1997	2000	41596	43922	41235	41711	41252	37510	34057	31516	31270	28407	27452	27266	25465
Passenger Cars	Diesel 1,4 - 2,0 l	Euro III	2001	2005		23955	32964	38627	38821	36272	38243	35241	34761	31564	30561	30354	28230
Passenger Cars	Diesel 1,4 - 2,0 l	Euro IV	2006	2010							20846	28905	33241	33350	33010	33752	31660
Passenger Cars	Diesel 1,4 - 2,0 l	Euro V	2011	2014							20846	25491	26060	28378	27978	30111	31194
Passenger Cars	Diesel >2,0 l	Conventional	0	1990	32416	28512	27974	28962	28929	26435	23948	23759	22352	20170	19613	19575	18370

Passenger Cars	Diesel >2,0 l	Euro I	1991	1996	41527	35930	34584	35270	34807	31561	28456	28427	26535	23936	23289	23340	22116
Passenger Cars	Diesel >2,0 l	Euro II	1997	2000	64167	62355	48724	43301	40719	36289	32454	30870	29733	26845	26007	25874	24141
Passenger Cars	Diesel >2,0 l	Euro III	2001	2005		42598	57973	62445	56784	50073	47421	39034	34785	30679	29446	29083	27332
Passenger Cars	Diesel >2,0 l	Euro IV	2006	2010							32485	44190	49037	41813	37119	34187	30832
Passenger Cars	Diesel >2,0 l	Euro V	2011	2014							32485	38678	44322	43444	43214	41146	38817
Passenger Cars	LPG cars	Conventional	0	1990	25034	22415	21851	20927	20407	18906	17304	15442	14908	13803	12490	12087	10833
Passenger Cars	LPG cars	Euro I	1991	1996	31569	28387	27999	28490	27879	25815	23399	22887	21670	20318	19523	19654	18606
Passenger Cars	LPG cars	Euro II	1997	2000			31538	30754	29812	27773	27787			22727	20577	18904	16990
Passenger Cars	LPG cars	Euro III	2001	2005								14961	25009	25508	23665	20230	17708
Passenger Cars	LPG cars	Euro IV	2006	2010										13953	26497	23106	20696
Passenger Cars	2-Stroke	Conventional	0	9999	13027	12291	12154	11891	11609								
Passenger Cars	Electric cars	Conventional	0	9999	13435	12364	12487	12254	11681	10761	9887	8459	9257	8288	9118	7368	8120
Passenger Cars	Gasoline <0,8 l	PRE ECE	0	1969							6621	6513	6184	5900	5575	5378	5144
Passenger Cars	Gasoline <0,8 l	ECE 15/00-01	1970	1978							7611	7532	7117	6784	6456	6266	6039
Passenger Cars	Gasoline <0,8 l	ECE 15/02	1979	1980								9128	8663	8229	7771	7539	7229
Passenger Cars	Gasoline <0,8 l	ECE 15/03	1981	1985						11385	9948	10021	9507	9069	8583	8109	7883
Passenger Cars	Gasoline <0,8 l	ECE 15/04	1986	1990							11614	11488	10868	10306	9799	9473	9000
Passenger Cars	Gasoline <0,8 l	Euro I	1991	1996						14585	13495	13229	12578	12032	11408	11046	10567
Passenger Cars	Gasoline <0,8 l	Euro II	1997	2000	13907	18697	18487	18086	17658	16460	15353	15127	14287	13621	12890	12497	11976
Passenger Cars	Gasoline <0,8 l	Euro III	2001	2005		9687	13912	15384	15170	14401	17114	16890	15972	15228	14408	13917	13368
Passenger Cars	Gasoline <0,8 l	Euro IV	2006	2010							9067	12702	12193	14852	15507	15582	14979
Passenger Cars	Gasoline <0,8 l	Euro V	2011	2014												8403	12424
Passenger Cars	Diesel <1,4 l	Conventional	0	1990						27755	25870	20195	22357	20190	19276	19727	17962
Passenger Cars	Diesel <1,4 l	Euro I	1991	1996						29641	27891	22719	24739	22721	21814	21622	20127
Passenger Cars	Diesel <1,4 l	Euro II	1997	2000	27578	38048	37220	38424	38221	34809	31433	31584	29436	26541	25851	25907	24498
Passenger Cars	Diesel <1,4 l	Euro III	2001	2005		19592	26227	32202	33707	32256	34314	34528	32138	28977	28231	28300	26835
Passenger Cars	Diesel <1,4 l	Euro IV	2006	2010							18213	24839	28556	30318	30212	30946	29282
Passenger Cars	Diesel <1,4 l	Euro V	2011	2014							18213	22062	22285	26396	25958	27381	26805
Light Duty Vehicles	Gasoline <3,5t	Conventional	0	1994	15170	16229	15923	15590	15077	13995	13045	14073	11585	10837	10352	9636	9103
Light Duty Vehicles	Gasoline <3,5t	Euro I	1995	1998	19114	20185	19629	19084	18392	17019	15867	17181	14296	13550	13216	12643	12341
Light Duty Vehicles	Gasoline <3,5t	Euro II	1999	2001	16055	18632	21916	21295	20510	18964	17677	19124	15905	15061	14682	14031	13689
Light Duty Vehicles	Gasoline <3,5t	Euro III	2002	2006			11722	16961	17824	17877	18004	22096	18384	17401	16968	16226	15847
Light Duty Vehicles	Gasoline <3,5t	Euro IV	2007	2011								12110	18383	19066	18217	17550	17871
Light Duty Vehicles	Gasoline <3,5t	Euro V	2012	2015													10247
Light Duty Vehicles	Diesel <3,5t	Conventional	0	1994	25182	23519	22569	22795	22397	20777	19686	20754	18207	16802	16322	16011	15471
Light Duty Vehicles	Diesel <3,5t	Euro I	1995	1998	33622	31006	29494	29561	28903	26692	25300	26687	23438	21646	21092	20806	20248
Light Duty Vehicles	Diesel <3,5t	Euro II	1999	2001	27919	29302	33883	33943	33169	30594	28989	30560	26835	24753	24119	23778	23156
Light Duty Vehicles	Diesel <3,5t	Euro III	2002	2006			18355	27898	29513	29155	29500	36603	32181	29692	28947	28540	27815

Light Duty Vehicles	Diesel <3,5t	Euro IV	2007	2011								20289	30220	32943	31675	28960	32828
Light Duty Vehicles	Diesel <3,5t	Euro V	2012	2015							18478	24202	27928	32447	29400	33302	27218
Light Duty Vehicles	LPG <3,5t	Conventional	0	1994	16708	15378	14693	13669	12032	10422	10437	10453	8462	8231	7815	7487	7278
Light Duty Vehicles	LPG <3,5t	Euro I	1995	1998	29696	26882											
Light Duty Vehicles	LPG <3,5t	Euro II	1999	2001	16111				26621	24311	23755	24769	22124	20658	19613	19295	18756
Light Duty Vehicles	LPG <3,5t	Euro III	2002	2006							22160	28203	25620	24746	23736	22863	22356
Light Duty Vehicles	LPG <3,5t	Euro IV	2007	2011								15822	23749	27665	26444	25420	24960
Light Duty Vehicles	Electric <3,5t	Conventional	0	9999	11034								11736	9258	11038	8068	8018
Heavy Duty Vehicles	Gasoline >3,5t	Conventional	0	9999	14866	15637	16704	15797	15284	14526	15520	17127	15235	14799	15095	14421	11319
Heavy Duty Vehicles	Diesel RT 3,5 - 7,5t	Conventional	0	1993	16450	17555	15899	15373	15109	11852	10381	9304	7564	6067	5630	5475	4760
Heavy Duty Vehicles	Diesel RT 3,5 - 7,5t	Euro I	1994	1996	30112	31797	28690	27720	27193	21469	19223	17600	14718	12188	11720	11483	10300
Heavy Duty Vehicles	Diesel RT 3,5 - 7,5t	Euro II	1997	2001	36148	41424	39193	37869	37116	29441	26408	24062	20163	16699	16054	15776	14373
Heavy Duty Vehicles	Diesel RT 3,5 - 7,5t	Euro III	2002	2006		26608	33528	40593	44021	36845	36376	35890	30122	24977	24046	23673	21401
Heavy Duty Vehicles	Diesel RT 3,5 - 7,5t	Euro IV	2007	2009							24022	26634	30074	31205	32784	33057	29985
Heavy Duty Vehicles	Diesel RT 3,5 - 7,5t	Euro V	2010	2013								23745	28756	22683	22963	30032	31238
Heavy Duty Vehicles	Diesel RT 3,5 - 7,5t	Euro VI	2014	9999													21027
Heavy Duty Vehicles	Diesel RT 7,5 - 12t	Conventional	0	1993	19695	20999	19041	18508	18160	14262	12574	11252	9064	7309	6670	6230	5295
Heavy Duty Vehicles	Diesel RT 7,5 - 12t	Euro I	1994	1996	36691	38573	34845	33447	32780	25793	23049	20966	17472	14586	13965	13732	12337
Heavy Duty Vehicles	Diesel RT 7,5 - 12t	Euro II	1997	2001	43237	49590	47514	46026	45324	35868	32101	29250	24606	20551	19609	19376	17399
Heavy Duty Vehicles	Diesel RT 7,5 - 12t	Euro III	2002	2006		32748	41535	50625	52691	44235	43099	43431	36868	30995	29704	29326	26389
Heavy Duty Vehicles	Diesel RT 7,5 - 12t	Euro IV	2007	2009		32085	57865	47772	46787	34531	29689	34165	37414	39679	39794	39506	35578
Heavy Duty Vehicles	Diesel RT 7,5 - 12t	Euro V	2010	2013				47115	46144	36317	29099	29348	31124	33722	36569	39808	38722
Heavy Duty Vehicles	Diesel RT 7,5 - 12t	Euro VI	2014	9999													36845
Heavy Duty Vehicles	Diesel RT 12 - 14 t	Conventional	0	1993	17020	17197	15678	15125	14591	13337	11853	10777	8964	7339	7081	6970	6432
Heavy Duty Vehicles	Diesel RT 12 - 14 t	Euro I	1994	1996	30408	30573	27575	26659	25571	23430	21009	18951	15864	13086	12536	12350	11316
Heavy Duty Vehicles	Diesel RT 12 - 14 t	Euro II	1997	2001	37280	39786	36339	35219	33898	31293	28220	25609	21631	17653	17084	16969	15734
Heavy Duty Vehicles	Diesel RT 12 - 14 t	Euro III	2002	2006		25939	33136	40304	41075	39536	40367	40247	33971	28018	27300	27002	24780
Heavy Duty Vehicles	Diesel RT 12 - 14 t	Euro IV	2007	2009					43872	40215	30758	28175	33932	34265	38047	37386	34580
Heavy Duty Vehicles	Diesel RT 12 - 14 t	Euro V	2010	2013									24386	34805	28377	37228	39941
Heavy Duty Vehicles	Diesel RT 14 - 20t	Conventional	0	1993	31119	31354	28485	27785	26718	24659	21953	19799	16454	13399	12849	12500	11184
Heavy Duty Vehicles	Diesel RT 14 - 20t	Euro I	1994	1996	53993	54048	48968	47659	45903	42077	37638	34149	28425	23384	22682	22263	20511
Heavy Duty Vehicles	Diesel RT 14 - 20t	Euro II	1997	2001	63606	70424	66757	64879	62491	57474	51562	47035	39361	32492	31628	31177	28888
Heavy Duty Vehicles	Diesel RT 14 - 20t	Euro III	2002	2006		46965	60842	68989	74458	72743	70620	71117	59772	49349	48258	47464	44004
Heavy Duty Vehicles	Diesel RT 14 - 20t	Euro IV	2007	2009		45403	70832	71755	79578	62868	52425	54598	63084	63136	64166	63437	58695
Heavy Duty Vehicles	Diesel RT 14 - 20t	Euro V	2010	2013						48862	53801	54097	55630	55557	60808	64497	63200
Heavy Duty Vehicles	Diesel RT 14 - 20t	Euro VI	2014	9999													42804
Heavy Duty Vehicles	Diesel RT 20 - 26t	Conventional	0	1993	50550	44699	41095	40144	38234	34823	31048	28241	23423	19199	18217	17701	15824
Heavy Duty Vehicles	Diesel RT 20 - 26t	Euro I	1994	1996	84574	73953	67587	65983	62312	56632	50601	46253	38537	31950	30592	29937	26861

Heavy Duty Vehicles	Diesel RT 20 - 26t	Euro II	1997	2001	99869	96014	91503	89421	84683	77053	68994	63254	52904	43999	42249	41678	37384
Heavy Duty Vehicles	Diesel RT 20 - 26t	Euro III	2002	2006		63686	80427	97199	100972	97719	95376	98317	82562	68864	66537	65879	59414
Heavy Duty Vehicles	Diesel RT 20 - 26t	Euro IV	2007	2009		62376	106351	110845	98270	69114	76473	74125	85219	85664	86853	85925	77330
Heavy Duty Vehicles	Diesel RT 20 - 26t	Euro V	2010	2013						66080	81296	73989	76168	74942	81123	86260	82628
Heavy Duty Vehicles	Diesel RT 20 - 26t	Euro VI	2014	9999													56353
Heavy Duty Vehicles	Diesel RT 26 - 28t	Conventional	0	1993	52379	44939	42151	40044	37721	34306	30520	27764	23315	18204	18548	18059	17251
Heavy Duty Vehicles	Diesel RT 26 - 28t	Euro I	1994	1996	86401	72870	66888	64932	59815	59846	52832	48061	39945	35123	32924		28450
Heavy Duty Vehicles	Diesel RT 26 - 28t	Euro II	1997	2001	107624	94817	86528	83997	79987	72997	64442	58166	51037	40222	44120	42955	36535
Heavy Duty Vehicles	Diesel RT 26 - 28t	Euro III	2002	2006				66051	81831	110192	109123	99270	82506	68531	69828	67985	60359
Heavy Duty Vehicles	Diesel RT 26 - 28t	Euro IV	2007	2009								63653	94454	91708	92144	90976	84108
Heavy Duty Vehicles	Diesel RT 26 - 28t	Euro V	2010	2013								63653	105808	87886	93442	92240	82462
Heavy Duty Vehicles	Diesel RT 28 - 32t	Conventional	0	1993	58442	51503	46870	45155	43125	39576	35629	31681	26713	21646	22642	20693	19388
Heavy Duty Vehicles	Diesel RT 28 - 32t	Euro I	1994	1996	85349	74915	69088	67408	63266	57637	50834	46085	38227	32045	32473	32020	28983
Heavy Duty Vehicles	Diesel RT 28 - 32t	Euro II	1997	2001	96195	97370	96270	93945	88835	81258	72107	65910	55034	46003	47142	45909	40821
Heavy Duty Vehicles	Diesel RT 28 - 32t	Euro III	2002	2006		63397	82736	96181	100949	98168	93655	100171	84589	70681	72224	70649	63020
Heavy Duty Vehicles	Diesel RT 28 - 32t	Euro IV	2007	2009			62598	84545	121119	69919	83240	75075	85403	85633	92136	90551	80466
Heavy Duty Vehicles	Diesel RT 28 - 32t	Euro V	2010	2013							64373	71114	76710	76054	88349	87216	83841
Heavy Duty Vehicles	Diesel RT 28 - 32t	Euro VI	2014	9999													58702
Heavy Duty Vehicles	Diesel RT >32t	Conventional	0	1993	59903	56213	51599	27663	26136	23942	18917	17209	14303				
Heavy Duty Vehicles	Diesel RT >32t	Euro I	1994	1996	78093	69998	70083	68033	64086	58485	51631	46969	39037	32425	35787	34843	30924
Heavy Duty Vehicles	Diesel RT >32t	Euro II	1997	2001	82426	110824	105966										
Heavy Duty Vehicles	Diesel RT >32t	Euro III	2002	2006		62740	115180	99028	102706	87264	82996	79418	73115	76652	78102	87628	75316
Heavy Duty Vehicles	Diesel RT >32t	Euro IV	2007	2009								63653	105808	87886			77379
Heavy Duty Vehicles	Diesel RT >32t	Euro V	2010	2013										87886	68183	76657	85692
Heavy Duty Vehicles	Diesel TT/AT 28 - 34t	Conventional	0	1993	34196	32408	29399	28876	28262	25969	24016	22207	19350	14701	15343	15300	12788
Heavy Duty Vehicles	Diesel TT/AT 28 - 34t	Euro I	1994	1996	58763	55334	50097	49153	48215	44114	40820	38016	33435	25869	27385	27566	23565
Heavy Duty Vehicles	Diesel TT/AT 28 - 34t	Euro II	1997	2001	69272	72203	68380	66970	65637	60193	55632	52022	45994	35734	38112	38572	33017
Heavy Duty Vehicles	Diesel TT/AT 28 - 34t	Euro III	2002	2006		48050	62204	71092	78113	76128	76146	78699	69796	54228	58113	58572	50405
Heavy Duty Vehicles	Diesel TT/AT 28 - 34t	Euro IV	2007	2009		46453	72418	73947	83437	65671	56567	60589	74099	69595	77672	78573	67077
Heavy Duty Vehicles	Diesel TT/AT 28 - 34t	Euro V	2010	2013						51040	57911	59688	64900	61096	73121	79322	72081
Heavy Duty Vehicles	Diesel TT/AT 28 - 34t	Euro VI	2014	9999													48737
Heavy Duty Vehicles	Diesel TT/AT 34 - 40t	Conventional	0	1993	56625	48112	43717	43168	41964	38857	36802	34493	31072	24481	27010	27845	23502
Heavy Duty Vehicles	Diesel TT/AT 34 - 40t	Euro I	1994	1996	90683	78398	71204	70175	67791	62711	58875	54932	48850	38263	42357	44087	37833
Heavy Duty Vehicles	Diesel TT/AT 34 - 40t	Euro II	1997	2001	102103	101183	96985	96677	94115	88121	83090	76632	68056	53129	58008	59761	50589
Heavy Duty Vehicles	Diesel TT/AT 34 - 40t	Euro III	2002	2006		68252	86315	101701	108823	110400	117037	121430	110059	86927	96562	97733	82177
Heavy Duty Vehicles	Diesel TT/AT 34 - 40t	Euro IV	2007	2009		64138	96872	118439	91155	80530	89003	87231	109795	104049	121515	125950	104924
Heavy Duty Vehicles	Diesel TT/AT 34 - 40t	Euro V	2010	2013		63626	116998	116736		74047	89476	89628	108189	100922	122305	130457	122488
Heavy Duty Vehicles	Diesel TT/AT 34 - 40t	Euro VI	2014	9999													87962

Heavy Duty Vehicles	Diesel TT/AT 40 - 50t	Conventional	0	1993	72670	76003	68799	65556	61140	56358	51174	45518	37602	33403	31045	30957	27664
Heavy Duty Vehicles	Diesel TT/AT 40 - 50t	Euro I	1994	1996	105682	110697	100642	96515	90471	84051	76201	69347	57944	54294	52829	54976	51885
Heavy Duty Vehicles	Diesel TT/AT 40 - 50t	Euro II	1997	2001	115340	139916	136798	132096	124360	116363	104944	94121	77669	71987	67695	68312	65525
Heavy Duty Vehicles	Diesel TT/AT 40 - 50t	Euro III	2002	2006		90191	111040	129803	134683	136396	135797	140801	119475	112999	108722	109974	103805
Heavy Duty Vehicles	Diesel TT/AT 40 - 50t	Euro IV	2007	2009		87910	102529	148583	155594	93826	105539	96872	114685	132657	134987	139813	132813
Heavy Duty Vehicles	Diesel TT/AT 40 - 50t	Euro V	2010	2013		87910	123254	159991	151244	138321	90278	93461	110193	123151	126984	133242	137653
Heavy Duty Vehicles	Diesel TT/AT 40 - 50t	Euro VI	2014	9999													95740
Heavy Duty Vehicles	Diesel TT/AT 50 - 60t	Euro I	1994	1996	104015	113754	103723	99758	94304	89492							
Heavy Duty Vehicles	Diesel TT/AT 50 - 60t	Euro II	1997	2001	129314	141422	128952	124022	117242	111259	101197	91076	77175	73318	71764	74183	70542
Heavy Duty Vehicles	Diesel TT/AT 50 - 60t	Euro IV	2007	2009								87504	148296	140884	137898		135551
Heavy Duty Vehicles	Diesel TT/AT >60t	Euro III	2002	2006													126062
Heavy Duty Vehicles	Diesel TT/AT >60t	Euro V	2010	2013									79729	104793		95279	
Buses	Gasoline Urban Buses	Conventional	0	9999	16034	15579	15526	14611	13087	17997	18361	33302	31424	35250	30704	26962	24419
Buses	Diesel Urban Buses <15t	Conventional	0	1993	73676	69484	65605	63013	59405	53446	48573	44675	43353	38028	34078	31692	30675
Buses	Diesel Urban Buses <15t	Euro I	1994	1996	116914	108582	100999	96838	91737	81466	75670	71538	66510	59167	53603	50460	47829
Buses	Diesel Urban Buses <15t	Euro II	1997	2001	132924	131860	127923	123406	117254	103905	97082	91902	85181	77538	71842	68695	65658
Buses	Diesel Urban Buses <15t	Euro III	2002	2006			81916	110773	130640	123957	126569	136452	126717	111962	102346	98407	93387
Buses	Diesel Urban Buses <15t	Euro IV	2007	2009								83706	114450	119020	136850	129204	121754
Buses	Diesel Urban Buses <15t	Euro V	2010	2013											78390	122673	122988
Buses	Diesel Urban Buses 15 - 18t	Conventional	0	1993	72412	67883	64392	62326	59190	52698	50100	47644	44933	40538	37823	36127	34126
Buses	Diesel Urban Buses 15 - 18t	Euro I	1994	1996	115649	106807	98967	94505	89761	79737	74702	70868	65776	58629	53817	50245	47377
Buses	Diesel Urban Buses 15 - 18t	Euro II	1997	2001	129338	130498	130636	125521	119264	105683	98725	93769	87636	78039	71684	68015	64388
Buses	Diesel Urban Buses 15 - 18t	Euro III	2002	2006			81916	130578	137019	128466	120209	135041	125675	111742	102106	96158	91852
Buses	Diesel Urban Buses 15 - 18t	Euro IV	2007	2009								83706	107894	114047	138735	131097	123765
Buses	Diesel Urban Buses 15 - 18t	Euro V	2010	2013											78390	116290	126633
Buses	Diesel Urban Buses >18t	Conventional	0	1993	86260	80649	76134	73463	69148	62605	58948	53947	52718	44128	38307	38724	37862
Buses	Diesel Urban Buses >18t	Euro I	1994	1996	112266	104754	99766	95738	90872	80750	75344	73092	62172	54820	49093	46397	47071
Buses	Diesel Urban Buses >18t	Euro II	1997	2001	103154	119246	142861	137074	129796	115282	107308	101977	94686	84801	77314	73770	68718
Buses	Diesel Urban Buses >18t	Euro III	2002	2006			81916	127231	131471	133178	127581	131055	122668	109347	99994	94649	89982
Buses	Diesel Urban Buses >18t	Euro IV	2007	2009								83706	120106	126273	134902	127556	120401
Buses	Diesel Urban Buses >18t	Euro V	2010	2013											78390	120260	116928
Buses	Gasoline Coaches	Conventional	0	9999	15711	15044	14923	14430	14037	13151	12332	11849	11112	10252	9833	9326	8687
Buses	Diesel Coaches <15t	Conventional	0	1993	27553	26929	26143	26488	26584	24194	22527	21780	20015	17658	17261	16601	15240
Buses	Diesel Coaches <15t	Euro I	1994	1996	35550	34244	32950	33080	33021	29955	27774	26785	24638	21787	21352	20586	19041
Buses	Diesel Coaches <15t	Euro II	1997	2001	37435	38045	40184	40224	40093	36410	33850	32709	30136	26671	26137	25170	23283
Buses	Diesel Coaches <15t	Euro III	2002	2006			23581	37335	40695	39914	39191	42182	38766	34269	33591	32378	29767
Buses	Diesel Coaches <15t	Euro IV	2007	2009								24584	33549	33826	41249	39652	36681
Buses	Diesel Coaches <15t	Euro V	2010	2013											22802	35400	37408

Buses	Diesel Coaches 15 - 18t	Conventional	0	1993	37182	36133	35369	35855	36061	33280	31113	30305	28188	25296	25031	24576	23214
Buses	Diesel Coaches 15 - 18t	Euro I	1994	1996	58754	56477	54178	54405	54233	49181	45474	43832	40202	35668	34826	33482	30934
Buses	Diesel Coaches 15 - 18t	Euro II	1997	2001	60185	63385	66182	66524	66237	59825	55332	53359	49175	43470	42712	41296	38227
Buses	Diesel Coaches 15 - 18t	Euro III	2002	2006			38794	56152	71971	68771	65364	68179	62399	54978	53761	51665	47702
Buses	Diesel Coaches 15 - 18t	Euro IV	2007	2009								40443	50246	54782	68555	65943	60884
Buses	Diesel Coaches 15 - 18t	Euro V	2010	2013											37512	66509	61184
Buses	Diesel Coaches >18t	Conventional	0	1993	72220	69827	67704	68423	68914	62540	58112	56472	52345	45744	45422	44120	40841
Buses	Diesel Coaches >18t	Euro I	1994	1996	93701	90378	87280	87323	86828	78439	73096	70302	64768	57216	56129	53788	49629
Buses	Diesel Coaches >18t	Euro II	1997	2001	95313	97615	107547	108048	107837	97310	89693	86222	79492	70671	69284	66957	62392
Buses	Diesel Coaches >18t	Euro III	2002	2006			61716	86146	105183	102584	104553	110765	101831	90059	88406	84731	78412
Buses	Diesel Coaches >18t	Euro IV	2007	2009								64341	91238	94128	107323	103438	95682
Buses	Diesel Coaches >18t	Euro V	2010	2013											59678	90780	90893
Mopeds	2-stroke <50 cm³	Conventional	0	1999	2010	1565	1591	1581	1545	1518	1516	1511	1487	1472	1430	1395	1343
Mopeds	2-stroke <50 cm³	Euro I	2000	2003	1076	1330	1516	1594	1703	1672	1670	1664	1638	1621	1575	1536	1479
Mopeds	2-stroke <50 cm³	Euro II	2004	2014					914	1315	1498	1600	1664	1716	1701	1688	1661
Mopeds	4-stroke <50 cm³	Euro II	2004	2014					914	1315	1498	1600	1664	1716	1701	1688	1661
Motorcycles	2-stroke >50 cm³	Conventional	0	1999	5740	5282	5050	4710	4284	3891	3541	3280	2996	2759	2725	2667	2613
Motorcycles	2-stroke >50 cm³	Euro I	2000	2003	4322	6296	6494	6371	6664	6007	5412	4965	4513	4151	4097	4007	3923
Motorcycles	2-stroke >50 cm³	Euro II	2004	2006					3554	4496	4601	5465	4967	4568	4508	4408	4315
Motorcycles	2-stroke >50 cm³	Euro III	2007	9999								2858	4149	4381	4530	4548	4513
Motorcycles	4-stroke <250 cm³	Conventional	0	1999	5740	5282	5050	4710	4284	3891	3541	3280	2996	2759	2725	2667	2613
Motorcycles	4-stroke <250 cm³	Euro I	2000	2003	4322	6296	6494	6371	6664	6007	5412	4965	4513	4151	4097	4007	3923
Motorcycles	4-stroke <250 cm³	Euro II	2004	2006					3554	4496	4601	5465	4967	4568	4508	4408	4315
Motorcycles	4-stroke <250 cm³	Euro III	2007	9999								2858	4149	4381	4530	4548	4513
Motorcycles	4-stroke 250 - 750 cm³	Conventional	0	1999	5740	5282	5050	4710	4284	3891	3541	3280	2996	2759	2725	2667	2613
Motorcycles	4-stroke 250 - 750 cm³	Euro I	2000	2003	4322	6296	6494	6371	6664	6007	5412	4965	4513	4151	4097	4007	3923
Motorcycles	4-stroke 250 - 750 cm³	Euro II	2004	2006					3554	4496	4601	5465	4967	4568	4508	4408	4315
Motorcycles	4-stroke 250 - 750 cm³	Euro III	2007	9999								2858	4149	4381	4530	4548	4513
Motorcycles	4-stroke >750 cm³	Conventional	0	1999	5740	5282	5050	4710	4284	3891	3541	3280	2996	2759	2725	2667	2613
Motorcycles	4-stroke >750 cm³	Euro I	2000	2003	4322	6296	6494	6371	6664	6007	5412	4965	4513	4151	4097	4007	3923
Motorcycles	4-stroke >750 cm³	Euro II	2004	2006					3554	4496	4601	5465	4967	4568	4508	4408	4315
Motorcycles	4-stroke >750 cm³	Euro III	2007	9999								2858	4149	4381	4530	4548	4513

Annex 3B-3 EU directive emission limits for road transportation vehicles

Private cars and light duty vehicles I (<1305 kg).

G pr km		EURO 1	EURO 2	EURO 3 ¹⁾	EURO 4	EURO 5	EURO 6
<u>Normal temp.</u>							
CO	Gasoline	2.72	2.2	2.3	1.0	1.0	1.0
	Diesel	2.72	1.0	0.64	0.5	0.5	0.5
HC	Gasoline	-	-	0.20	0.10	0.1	0.1
NMHC	Gasoline	-	-	-	-	0.068	0.068
NO _x	Gasoline	-	-	0.15	0.08	0.06	0.06
	Diesel	-	-	0.5	0.25	0.18	0.08
HC+NO _x	Gasoline	0.97	0.5	-	-	-	-
	Diesel	0.97	0.7/0.9 ²⁾	0.56	0.30	0.23	0.17
Particulates	Diesel	0.14	0.08/0.10 ²⁾	0.05	0.025	0.005	0.005
Particulates (#)		-	-	-	-	-	6x10 ^{11 4)}
<u>Low temp.</u>							
CO	Gasoline	-	-	-	15	15	15
HC	Gasoline	-	-	-	1.8	1.8	1.8
<u>Evaporation</u>							
HC ³⁾	Gasoline	2.0	2.0	2.0	2.0	2.0	2.0

¹⁾ Changed test procedure at normal temperatures (40 s warm-up phase omitted) and for evaporation measurements. ²⁾ Less stringent emission limits for direct injection diesel engines. ³⁾ Unit: g/test. ⁴⁾ Applicable for diesel and gasoline direct injection (GDI). 6x10¹² within first three years of Euro 6 effective dates

Light duty vehicles II (1305-1760 kg)

G pr km		EURO 1	EURO 2	EURO 3 ¹⁾	EURO 4	EURO 5	EURO 6
<u>Normal temp.</u>							
CO	Gasoline	5.17	4.0	4.17	1.81	1.81	1.81
	Diesel	5.17	1.25	0.80	0.63	0.63	0.63
HC	Gasoline	-	-	0.25	0.13	0.13	0.13
NMHC	Gasoline	-	-	-	-	0.9	0.9
NO _x	Gasoline	-	-	0.18	0.10	0.75	0.75
	Diesel	-	-	0.65	0.33	0.235	0.105
HC+NO _x	Gasoline	1.4	0.6	-	-	-	-
	Diesel	1.4	1.0/1.3 ²⁾	0.72	0.39	0.295	0.195
Particulates	Gasoline					0.005	0.005
	Diesel	0.19	0.12/0.14 ²⁾	0.07	0.04	0.005	0.005
Particulates (#)		-	-	-	-	-	6x10 ^{11 4)}
<u>Low temp.</u>							
CO	Gasoline	-	-	-	24	24	24
HC	Gasoline	-	-		2.7	2.7	2.7
<u>Evaporation</u>							
HC ³⁾	Gasoline	2.0	2.0	2.0	2.0	2.0	2.0

¹⁾ Changed test procedure at normal temperatures (40 s warm-up phase omitted) and for evaporation measurements. ²⁾ Less stringent emission limits for direct injection diesel engines. ³⁾ Unit: g/test. ⁴⁾ Applicable for diesel and gasoline direct injection (GDI). 6x10¹² within first three years of Euro 6 effective dates

Light duty vehicles III (>1760 kg).

G pr km		EURO 1	EURO 2	EURO 3 ¹⁾	EURO 4	EURO 5	EURO 6
<u>Normal temp.</u>							
CO	Gasoline	6.9	5.0	5.22	2.27	2.27	2.27
	Diesel	6.9	1.5	0.95	0.74	0.74	0.74
HC	Gasoline	-	-	0.29	0.16	0.16	0.16
NMHC	Gasoline					0.108	0.108
NO _x	Gasoline	-	-	0.21	0.11	0.082	0.082
	Diesel	-	-	0.78	0.39	0.28	0.125
HC+NO _x	Gasoline	1.7	0.7	-	-	-	-
	Diesel	1.7	1.2/1.6 ²⁾	0.86	0.46	0.35	0.215
Particulates	Gasoline					0.005	0.005
	Diesel	0.25	0.17/0.20 ²⁾	0.10	0.06	0.005	0.005
Particulates (#)		-	-	-	-	-	6x10 ^{11 4)}
<u>Low temp.</u>							
CO	Gasoline	-	-	-	30	30	30
HC	Gasoline	-	-	-	3.2	3.2	3.2
<u>Evaporation</u>							
HC ³⁾	Gasoline	2.0	2.0	2.0	2.0	2.0	2.0

¹⁾ Changed test procedure at normal temperatures (40 s warm-up phase omitted) and for evaporation measurements. ²⁾ Less stringent emission limits for direct injection diesel engines. ³⁾ Unit: g/test. ⁴⁾ Applicable for diesel and gasoline direct injection (GDI). 6x10¹² within first three years of Euro 6 effective dates.

Heavy duty diesel vehicles.

(g pr kWh)		EURO						
Test ¹⁾		EURO I	EURO II	EURO III	IV	EURO V	EURO VI	EEV ²⁾
		1993	1996	2001	2006	2009	2014	2000
CO	ECE/ESC	4.5	4.0	2.1	1.5	1.5	1.5	1.5
	ETC	-	-	(5.45)	4.0	4.0	4.0	3.0
HC	ECE/ESC	1.1	1.1	0.66	0.46	0.46	0.13	0.25
	ETC	-	-	(0.78)	0.55	0.55	0.16	0.40
NO _x	ECE/ESC	8.0	7.0	5.0	3.5	2.0	0.4	2.0
	ETC	-	-	(5.0)	3.5	2.0	0.4	2.0
Particulates ³⁾	ECE/ESC	0.36/0.61	0.15/0.25	0.10/0.13	0.02	0.02	0.01	0.02
	ETC	-	-	(0.16/0.21)	0.03	0.03	0.01	0.02
	ELR	-	-	0.8	0.5	0.5		0.15
NH ₃	ECE/ESC						10 (ppm)	
	ETC						10 (ppm)	

¹⁾ Test procedure:

Euro 1 og Euro 2: ECE (stationary)

Euro 3: ESC (stationary) + ELR (load response)

Euro 4, Euro 5 og EEV: ESC (stationary) + ETC (transient) + ELR (load response)

²⁾ EEV: Emission limits for extra environmental friendly vehicles, used as a basis for economical incitements (gas fueled vehicles).

³⁾ For Euro 1, Euro 2 og Euro 3 less stringent emission limits apply for small engines:

Euro 1: <85 kW

Euro 2: <0,7 l

Euro 3: <0,75 l

Annex 3B-4 Basis emission factors (g pr km)

Sector	Subsector	Tech 2	FYear	LYear	FCu	FCr	FCh	COu	COr	COh	PMu	PMr	PMh	NOxu	NOxr	NOxh
Passenger Cars	Gasoline 0,8 - 1,4 l	PRE ECE	0	1969	67,499	55,000	62,743	27,505	19,333	15,520	0,063	0,044	0,041	1,849	2,062	2,023
Passenger Cars	Gasoline 0,8 - 1,4 l	ECE 15/00-01	1970	1978	58,240	44,460	48,600	18,966	14,480	18,620	0,063	0,044	0,041	1,849	2,062	2,023
Passenger Cars	Gasoline 0,8 - 1,4 l	ECE 15/02	1979	1980	53,248	45,170	51,200	15,859	8,200	8,260	0,063	0,044	0,041	1,619	2,102	2,909
Passenger Cars	Gasoline 0,8 - 1,4 l	ECE 15/03	1981	1985	53,248	45,170	51,200	16,752	8,793	7,620	0,042	0,029	0,029	1,680	2,253	3,276
Passenger Cars	Gasoline 0,8 - 1,4 l	ECE 15/04	1986	1990	51,420	43,440	47,700	9,087	4,956	4,292	0,030	0,020	0,020	1,691	2,089	2,662
Passenger Cars	Gasoline 0,8 - 1,4 l	Euro I	1991	1996	47,399	41,954	46,055	1,765	1,372	1,765	0,003	0,002	0,002	0,273	0,281	0,458
Passenger Cars	Gasoline 0,8 - 1,4 l	Euro II	1997	2000	46,486	39,509	44,016	0,659	0,575	0,749	0,003	0,002	0,002	0,154	0,154	0,181
Passenger Cars	Gasoline 0,8 - 1,4 l	Euro III	2001	2005	48,687	42,255	45,323	0,519	0,691	1,148	0,001	0,001	0,001	0,076	0,060	0,052
Passenger Cars	Gasoline 0,8 - 1,4 l	Euro IV	2006	2010	50,038	44,193	48,285	0,195	0,287	0,529	0,001	0,001	0,001	0,054	0,030	0,019
Passenger Cars	Gasoline 0,8 - 1,4 l	Euro V	2011	2014	50,038	44,193	48,285	0,195	0,287	0,529	0,001	0,001	0,001	0,041	0,023	0,014
Passenger Cars	Gasoline 1,4 - 2,0 l	PRE ECE	0	1969	79,277	67,000	76,386	27,505	19,333	15,520	0,063	0,044	0,041	2,164	2,683	3,130
Passenger Cars	Gasoline 1,4 - 2,0 l	ECE 15/00-01	1970	1978	67,779	51,090	60,300	18,966	14,480	18,620	0,063	0,044	0,041	2,164	2,683	3,130
Passenger Cars	Gasoline 1,4 - 2,0 l	ECE 15/02	1979	1980	61,731	50,686	59,680	15,859	8,200	8,260	0,063	0,044	0,041	1,831	2,377	3,283
Passenger Cars	Gasoline 1,4 - 2,0 l	ECE 15/03	1981	1985	61,731	50,686	59,680	16,752	8,793	7,620	0,042	0,029	0,029	1,917	2,580	3,472
Passenger Cars	Gasoline 1,4 - 2,0 l	ECE 15/04	1986	1990	61,652	49,112	52,052	9,087	4,956	4,292	0,030	0,020	0,020	2,122	2,757	3,524
Passenger Cars	Gasoline 1,4 - 2,0 l	Euro I	1991	1996	57,521	48,522	51,518	1,765	1,372	1,765	0,003	0,002	0,002	0,273	0,281	0,458
Passenger Cars	Gasoline 1,4 - 2,0 l	Euro II	1997	2000	56,324	47,687	48,786	0,659	0,575	0,749	0,003	0,002	0,002	0,154	0,154	0,181
Passenger Cars	Gasoline 1,4 - 2,0 l	Euro III	2001	2005	58,259	49,897	53,092	0,519	0,691	1,148	0,001	0,001	0,001	0,076	0,060	0,052
Passenger Cars	Gasoline 1,4 - 2,0 l	Euro IV	2006	2010	60,486	52,793	55,293	0,195	0,287	0,529	0,001	0,001	0,001	0,054	0,030	0,019
Passenger Cars	Gasoline 1,4 - 2,0 l	Euro V	2011	2014	60,486	52,793	55,293	0,195	0,287	0,529	0,001	0,001	0,001	0,041	0,023	0,014
Passenger Cars	Gasoline >2,0 l	PRE ECE	0	1969	96,536	80,000	88,267	27,505	19,333	15,520	0,063	0,044	0,041	2,860	4,090	5,500
Passenger Cars	Gasoline >2,0 l	ECE 15/00-01	1970	1978	73,798	57,090	66,300	18,966	14,480	18,620	0,063	0,044	0,041	2,860	4,090	5,500
Passenger Cars	Gasoline >2,0 l	ECE 15/02	1979	1980	75,270	63,260	70,700	15,859	8,200	8,260	0,063	0,044	0,041	2,066	2,675	3,680
Passenger Cars	Gasoline >2,0 l	ECE 15/03	1981	1985	75,270	63,260	70,700	16,752	8,793	7,620	0,042	0,029	0,029	2,806	3,441	4,604
Passenger Cars	Gasoline >2,0 l	ECE 15/04	1986	1990	71,055	58,080	69,900	9,087	4,956	4,292	0,030	0,020	0,020	2,293	2,750	3,687
Passenger Cars	Gasoline >2,0 l	Euro I	1991	1996	74,616	61,902	65,020	1,765	1,372	1,765	0,003	0,002	0,002	0,273	0,281	0,458
Passenger Cars	Gasoline >2,0 l	Euro II	1997	2000	76,837	65,226	66,732	0,659	0,575	0,749	0,003	0,002	0,002	0,154	0,154	0,181
Passenger Cars	Gasoline >2,0 l	Euro III	2001	2005	70,798	57,424	56,826	0,519	0,691	1,148	0,001	0,001	0,001	0,076	0,060	0,052
Passenger Cars	Gasoline >2,0 l	Euro IV	2006	2010	86,099	67,877	65,859	0,195	0,287	0,529	0,001	0,001	0,001	0,054	0,030	0,019
Passenger Cars	Gasoline >2,0 l	Euro V	2011	2014	86,099	67,877	65,859	0,195	0,287	0,529	0,001	0,001	0,001	0,041	0,023	0,014
Passenger Cars	Diesel 1,4 - 2,0 l	Conventional	0	1990	57,529	41,209	50,089	0,651	0,472	0,384	0,199	0,132	0,170	0,520	0,433	0,528
Passenger Cars	Diesel 1,4 - 2,0 l	Euro I	1991	1996	47,836	42,807	48,388	0,419	0,215	0,208	0,057	0,062	0,107	0,603	0,562	0,663
Passenger Cars	Diesel 1,4 - 2,0 l	Euro II	1997	2000	50,442	44,117	48,779	0,343	0,110	0,035	0,047	0,039	0,050	0,651	0,555	0,665

Passenger Cars	Diesel 1,4 - 2,0 l	Euro III	2001	2005	48,920	43,427	45,585	0,099	0,041	0,012	0,029	0,030	0,045	0,716	0,665	0,750
Passenger Cars	Diesel 1,4 - 2,0 l	Euro IV	2006	2010	48,920	43,427	45,585	0,083	0,034	0,021	0,029	0,024	0,026	0,539	0,424	0,576
Passenger Cars	Diesel 1,4 - 2,0 l	Euro V	2011	2014	48,889	43,371	45,487	0,083	0,034	0,021	0,006	0,005	0,005	0,663	0,521	0,708
Passenger Cars	Diesel >2,0 l	Conventional	0	1990	57,529	41,209	50,089	0,651	0,472	0,384	0,199	0,132	0,170	0,824	0,723	0,861
Passenger Cars	Diesel >2,0 l	Euro I	1991	1996	65,267	58,299	64,360	0,419	0,215	0,208	0,057	0,062	0,107	0,603	0,562	0,663
Passenger Cars	Diesel >2,0 l	Euro II	1997	2000	65,267	58,299	64,360	0,343	0,110	0,035	0,047	0,039	0,050	0,651	0,555	0,665
Passenger Cars	Diesel >2,0 l	Euro III	2001	2005	65,267	58,299	64,360	0,099	0,041	0,012	0,029	0,030	0,045	0,716	0,665	0,750
Passenger Cars	Diesel >2,0 l	Euro IV	2006	2010	65,267	58,299	64,360	0,083	0,034	0,021	0,029	0,024	0,026	0,539	0,424	0,576
Passenger Cars	Diesel >2,0 l	Euro V	2011	2014	65,315	58,336	64,394	0,083	0,034	0,021	0,006	0,005	0,005	0,663	0,521	0,708
Passenger Cars	LPG cars	Conventional	0	1990	59,000	45,000	54,000	2,043	2,373	9,723	0,040	0,030	0,025	2,203	2,584	2,861
Passenger Cars	LPG cars	Euro I	1991	1996	49,145	45,155	54,125	1,310	1,445	3,560	0,040	0,030	0,025	0,340	0,283	0,298
Passenger Cars	LPG cars	Euro II	1997	2000	49,145	45,155	54,125	0,891	0,982	2,421	0,040	0,030	0,025	0,122	0,102	0,107
Passenger Cars	LPG cars	Euro III	2001	2005	49,145	45,155	54,125	0,733	0,809	1,993	0,040	0,030	0,025	0,082	0,068	0,071
Passenger Cars	LPG cars	Euro IV	2006	2010	49,145	45,155	54,125	0,445	0,491	1,210	0,040	0,030	0,025	0,044	0,037	0,039
Passenger Cars	Gasoline <0,8 l	PRE ECE	0	1969	52,252	46,235	56,545	27,505	19,333	15,520	0,063	0,044	0,041	1,849	2,062	2,023
Passenger Cars	Gasoline <0,8 l	ECE 15/00-01	1970	1978	45,084	37,374	43,799	18,966	14,480	18,620	0,063	0,044	0,041	1,849	2,062	2,023
Passenger Cars	Gasoline <0,8 l	ECE 15/02	1979	1980	41,220	37,971	46,142	15,859	8,200	8,260	0,063	0,044	0,041	1,619	2,102	2,909
Passenger Cars	Gasoline <0,8 l	ECE 15/03	1981	1985	41,220	37,971	46,142	16,752	8,793	7,620	0,042	0,029	0,029	1,680	2,253	3,276
Passenger Cars	Gasoline <0,8 l	ECE 15/04	1986	1990	39,805	36,517	42,988	9,087	4,956	4,292	0,030	0,020	0,020	1,691	2,089	2,662
Passenger Cars	Gasoline <0,8 l	Euro I	1991	1996	36,693	35,268	41,505	1,765	1,372	1,765	0,003	0,002	0,002	0,273	0,281	0,458
Passenger Cars	Gasoline <0,8 l	Euro II	1997	2000	35,985	33,213	39,668	0,659	0,575	0,749	0,003	0,002	0,002	0,154	0,154	0,181
Passenger Cars	Gasoline <0,8 l	Euro III	2001	2005	37,689	35,521	40,846	0,519	0,691	1,148	0,001	0,001	0,001	0,076	0,060	0,052
Passenger Cars	Gasoline <0,8 l	Euro IV	2006	2010	38,735	37,150	43,515	0,195	0,287	0,529	0,001	0,001	0,001	0,054	0,030	0,019
Passenger Cars	Gasoline <0,8 l	Euro V	2011	2014	38,735	37,150	43,515	0,195	0,287	0,529	0,001	0,001	0,001	0,041	0,023	0,014
Passenger Cars	Diesel <1,4 l	Conventional	0	1990	36,565	30,792	40,702	0,651	0,472	0,384	0,199	0,132	0,170	0,520	0,433	0,528
Passenger Cars	Diesel <1,4 l	Euro I	1991	1996	30,404	31,986	39,319	0,419	0,215	0,208	0,057	0,062	0,107	0,603	0,562	0,663
Passenger Cars	Diesel <1,4 l	Euro II	1997	2000	32,060	32,965	39,637	0,343	0,110	0,035	0,047	0,039	0,050	0,651	0,555	0,665
Passenger Cars	Diesel <1,4 l	Euro III	2001	2005	31,093	32,450	37,042	0,099	0,041	0,012	0,029	0,030	0,045	0,716	0,665	0,750
Passenger Cars	Diesel <1,4 l	Euro IV	2006	2010	31,093	32,450	37,042	0,083	0,034	0,021	0,029	0,024	0,026	0,539	0,424	0,576
Passenger Cars	Diesel <1,4 l	Euro V	2011	2014	31,073	32,408	36,962	0,083	0,034	0,021	0,006	0,005	0,005	0,663	0,521	0,708
Light Duty Vehicles	Gasoline <3,5t	Conventional	0	1994	82,270	59,883	56,470	14,925	6,075	7,389	0,040	0,040	0,040	2,671	3,118	3,387
Light Duty Vehicles	Gasoline <3,5t	Euro I	1995	1998	96,450	70,388	66,450	4,187	0,862	1,087	0,003	0,002	0,002	0,427	0,400	0,429
Light Duty Vehicles	Gasoline <3,5t	Euro II	1999	2001	96,450	70,388	66,450	2,554	0,526	0,663	0,003	0,002	0,002	0,145	0,136	0,146
Light Duty Vehicles	Gasoline <3,5t	Euro III	2002	2006	96,450	70,388	66,450	2,177	0,448	0,565	0,001	0,001	0,001	0,090	0,084	0,090
Light Duty Vehicles	Gasoline <3,5t	Euro IV	2007	2011	96,450	70,388	66,450	1,172	0,241	0,304	0,001	0,001	0,001	0,043	0,040	0,043
Light Duty Vehicles	Gasoline <3,5t	Euro V	2012	2015	96,450	70,388	66,450	1,172	0,241	0,304	0,001	0,001	0,001	0,032	0,030	0,032
Light Duty Vehicles	Diesel <3,5t	Conventional	0	1994	76,718	65,934	72,142	1,124	1,009	1,060	0,285	0,303	0,322	1,673	0,843	0,834

Light Duty Vehicles	Diesel <3,5t	Euro I	1995	1998	68,860	58,185	63,660	0,393	0,328	0,423	0,070	0,066	0,090	1,138	0,975	1,022
Light Duty Vehicles	Diesel <3,5t	Euro II	1999	2001	68,860	58,185	63,660	0,393	0,328	0,423	0,070	0,066	0,090	1,138	0,975	1,022
Light Duty Vehicles	Diesel <3,5t	Euro III	2002	2006	68,860	58,185	63,660	0,322	0,269	0,347	0,047	0,044	0,061	0,956	0,819	0,859
Light Duty Vehicles	Diesel <3,5t	Euro IV	2007	2011	68,860	58,185	63,660	0,255	0,213	0,275	0,024	0,023	0,032	0,774	0,663	0,695
Light Duty Vehicles	Diesel <3,5t	Euro V	2012	2015	68,860	58,185	63,660	0,255	0,213	0,275	0,001	0,001	0,002	0,558	0,478	0,501
Light Duty Vehicles	LPG <3,5t	Conventional	0	1994	88,500	67,500	81,000	3,064	3,559	14,584	0,060	0,045	0,038	3,305	3,876	4,291
Light Duty Vehicles	LPG <3,5t	Euro II	1999	2001	73,718	67,733	81,188	1,336	1,474	3,631	0,060	0,045	0,038	0,183	0,153	0,161
Light Duty Vehicles	LPG <3,5t	Euro III	2002	2006	73,718	67,733	81,188	1,100	1,214	2,990	0,060	0,045	0,038	0,122	0,102	0,107
Light Duty Vehicles	LPG <3,5t	Euro IV	2007	2011	73,718	67,733	81,188	0,668	0,737	1,815	0,060	0,045	0,038	0,066	0,055	0,058
Heavy Duty Vehicles	Gasoline >3,5t	Conventional	0	9999	225,000	150,000	165,000	70,000	55,000	55,000	0,400	0,400	0,400	4,500	7,500	7,500
Heavy Duty Vehicles	Diesel RT 3,5 - 7,5t	Conventional	0	1993	125,002	110,985	112,984	2,060	1,509	1,351	0,321	0,240	0,216	4,211	4,104	4,476
Heavy Duty Vehicles	Diesel RT 3,5 - 7,5t	Euro I	1994	1996	100,036	91,682	104,222	0,668	0,501	0,546	0,126	0,095	0,090	2,939	2,938	3,316
Heavy Duty Vehicles	Diesel RT 3,5 - 7,5t	Euro II	1997	2001	94,988	88,592	101,003	0,534	0,466	0,461	0,059	0,053	0,061	3,223	3,118	3,414
Heavy Duty Vehicles	Diesel RT 3,5 - 7,5t	Euro III	2002	2006	101,379	92,883	105,924	0,660	0,481	0,452	0,067	0,048	0,041	2,499	2,300	2,498
Heavy Duty Vehicles	Diesel RT 3,5 - 7,5t	Euro IV	2007	2009	98,559	92,910	106,610	0,342	0,270	0,258	0,015	0,013	0,014	1,707	1,645	1,801
Heavy Duty Vehicles	Diesel RT 3,5 - 7,5t	Euro V	2010	2013	99,641	93,536	106,995	0,344	0,270	0,259	0,015	0,013	0,014	1,012	0,972	1,062
Heavy Duty Vehicles	Diesel RT 3,5 - 7,5t	Euro VI	2014	9999	98,248	91,130	103,414	0,355	0,276	0,260	0,001	0,001	0,001	0,233	0,097	0,060
Heavy Duty Vehicles	Diesel RT 7,5 - 12t	Conventional	0	1993	183,253	153,117	150,068	2,358	1,698	1,525	0,330	0,236	0,207	7,928	7,236	7,499
Heavy Duty Vehicles	Diesel RT 7,5 - 12t	Euro I	1994	1996	155,870	135,518	136,666	1,086	0,817	0,766	0,201	0,144	0,131	4,729	4,306	4,464
Heavy Duty Vehicles	Diesel RT 7,5 - 12t	Euro II	1997	2001	148,625	131,263	133,537	0,868	0,727	0,717	0,094	0,080	0,093	5,152	4,593	4,682
Heavy Duty Vehicles	Diesel RT 7,5 - 12t	Euro III	2002	2006	157,573	137,771	138,996	1,084	0,771	0,733	0,104	0,073	0,063	3,997	3,536	3,485
Heavy Duty Vehicles	Diesel RT 7,5 - 12t	Euro IV	2007	2009	151,450	136,152	138,554	0,553	0,418	0,369	0,023	0,019	0,019	2,728	2,512	2,488
Heavy Duty Vehicles	Diesel RT 7,5 - 12t	Euro V	2010	2013	153,617	137,425	139,289	0,560	0,421	0,374	0,023	0,019	0,019	1,647	1,483	1,468
Heavy Duty Vehicles	Diesel RT 7,5 - 12t	Euro VI	2014	9999	151,539	134,140	134,852	0,583	0,431	0,373	0,002	0,002	0,002	0,384	0,154	0,096
Heavy Duty Vehicles	Diesel RT 12 - 14 t	Conventional	0	1993	198,513	163,310	159,212	2,546	1,876	1,693	0,351	0,254	0,233	8,826	7,718	7,748
Heavy Duty Vehicles	Diesel RT 12 - 14 t	Euro I	1994	1996	170,171	144,307	143,334	1,200	0,918	0,866	0,218	0,159	0,147	5,321	4,638	4,638
Heavy Duty Vehicles	Diesel RT 12 - 14 t	Euro II	1997	2001	163,223	140,030	139,590	0,985	0,820	0,804	0,103	0,087	0,103	5,815	4,975	4,889
Heavy Duty Vehicles	Diesel RT 12 - 14 t	Euro III	2002	2006	172,146	146,082	144,611	1,176	0,873	0,835	0,109	0,078	0,071	4,745	3,881	3,702
Heavy Duty Vehicles	Diesel RT 12 - 14 t	Euro IV	2007	2009	163,114	142,925	143,274	0,599	0,448	0,410	0,024	0,020	0,020	3,208	2,754	2,620
Heavy Duty Vehicles	Diesel RT 12 - 14 t	Euro V	2010	2013	165,111	144,096	144,035	0,606	0,452	0,413	0,025	0,020	0,020	1,909	1,634	1,552
Heavy Duty Vehicles	Diesel RT 14 - 20t	Conventional	0	1993	261,662	205,735	193,152	3,512	2,514	2,221	0,483	0,341	0,298	11,287	9,455	9,120
Heavy Duty Vehicles	Diesel RT 14 - 20t	Euro I	1994	1996	212,834	172,142	164,411	1,612	1,206	1,117	0,298	0,209	0,181	6,721	5,601	5,385
Heavy Duty Vehicles	Diesel RT 14 - 20t	Euro II	1997	2001	204,313	167,263	160,324	1,267	1,025	1,002	0,129	0,105	0,122	7,473	6,118	5,804
Heavy Duty Vehicles	Diesel RT 14 - 20t	Euro III	2002	2006	215,351	173,802	164,914	1,601	1,150	1,096	0,153	0,106	0,090	6,139	4,859	4,431
Heavy Duty Vehicles	Diesel RT 14 - 20t	Euro IV	2007	2009	201,093	168,074	161,976	0,829	0,602	0,523	0,031	0,024	0,023	4,079	3,400	3,171
Heavy Duty Vehicles	Diesel RT 14 - 20t	Euro V	2010	2013	205,393	169,743	162,354	0,869	0,625	0,536	0,032	0,025	0,023	2,460	2,028	1,883
Heavy Duty Vehicles	Diesel RT 14 - 20t	Euro VI	2014	9999	200,654	165,826	158,608	0,870	0,625	0,536	0,003	0,002	0,002	0,694	0,262	0,135

Heavy Duty Vehicles	Diesel RT 20 - 26t	Conventional	0	1993	315,898	243,280	222,355	2,558	1,885	1,712	0,482	0,353	0,319	12,251	9,862	9,114
Heavy Duty Vehicles	Diesel RT 20 - 26t	Euro I	1994	1996	269,815	211,940	195,827	2,068	1,563	1,437	0,383	0,264	0,231	8,634	6,952	6,468
Heavy Duty Vehicles	Diesel RT 20 - 26t	Euro II	1997	2001	261,049	207,213	191,812	1,620	1,285	1,399	0,172	0,137	0,157	9,465	7,549	6,947
Heavy Duty Vehicles	Diesel RT 20 - 26t	Euro III	2002	2006	272,733	213,630	195,690	2,025	1,487	1,403	0,189	0,130	0,113	7,649	6,024	5,545
Heavy Duty Vehicles	Diesel RT 20 - 26t	Euro IV	2007	2009	257,598	207,458	192,565	1,003	0,728	0,628	0,041	0,031	0,028	5,146	4,223	3,967
Heavy Duty Vehicles	Diesel RT 20 - 26t	Euro V	2010	2013	260,560	209,253	193,919	1,015	0,735	0,634	0,041	0,031	0,028	3,062	2,508	2,353
Heavy Duty Vehicles	Diesel RT 20 - 26t	Euro VI	2014	9999	255,823	204,120	188,510	1,045	0,753	0,646	0,004	0,003	0,003	0,710	0,287	0,164
Heavy Duty Vehicles	Diesel RT 26 - 28t	Conventional	0	1993	333,975	257,930	233,499	2,703	1,987	1,810	0,512	0,375	0,336	12,868	10,379	9,526
Heavy Duty Vehicles	Diesel RT 26 - 28t	Euro I	1994	1996	286,465	225,388	206,076	2,162	1,647	1,535	0,398	0,281	0,244	9,122	7,308	6,742
Heavy Duty Vehicles	Diesel RT 26 - 28t	Euro II	1997	2001	276,892	220,156	201,909	1,682	1,346	1,457	0,185	0,148	0,167	9,876	7,848	7,164
Heavy Duty Vehicles	Diesel RT 26 - 28t	Euro III	2002	2006	289,328	227,449	206,788	2,121	1,582	1,481	0,201	0,141	0,118	7,733	6,089	5,633
Heavy Duty Vehicles	Diesel RT 26 - 28t	Euro IV	2007	2009	277,178	222,906	203,989	1,044	0,752	0,640	0,044	0,033	0,029	5,258	4,284	4,029
Heavy Duty Vehicles	Diesel RT 26 - 28t	Euro V	2010	2013	280,496	224,942	205,435	1,056	0,760	0,647	0,044	0,033	0,029	3,127	2,544	2,388
Heavy Duty Vehicles	Diesel RT 28 - 32t	Conventional	0	1993	369,813	292,229	265,715	2,928	2,149	2,047	0,567	0,415	0,376	14,515	11,942	11,008
Heavy Duty Vehicles	Diesel RT 28 - 32t	Euro I	1994	1996	324,707	259,936	238,178	2,377	1,862	1,795	0,436	0,314	0,281	10,453	8,509	7,843
Heavy Duty Vehicles	Diesel RT 28 - 32t	Euro II	1997	2001	313,359	251,467	240,101	1,930	1,574	1,563	0,211	0,172	0,195	11,232	9,043	8,280
Heavy Duty Vehicles	Diesel RT 28 - 32t	Euro III	2002	2006	327,617	262,877	239,852	2,325	1,732	1,685	0,214	0,153	0,135	8,883	7,017	6,445
Heavy Duty Vehicles	Diesel RT 28 - 32t	Euro IV	2007	2009	316,735	259,706	237,679	1,145	0,834	0,714	0,049	0,038	0,034	5,978	5,101	4,533
Heavy Duty Vehicles	Diesel RT 28 - 32t	Euro V	2010	2013	320,475	262,095	239,548	1,159	0,844	0,722	0,049	0,038	0,034	3,554	3,030	2,690
Heavy Duty Vehicles	Diesel RT 28 - 32t	Euro VI	2014	9999	314,526	255,828	233,053	1,191	0,863	0,736	0,005	0,004	0,003	0,638	0,290	0,191
Heavy Duty Vehicles	Diesel RT >32t	Euro I	1994	1996	328,394	256,124	232,416	2,482	1,894	1,795	0,453	0,317	0,286	10,614	8,446	7,666
Heavy Duty Vehicles	Diesel RT >32t	Euro III	2002	2006	330,977	257,873	232,502	2,398	1,789	1,725	0,219	0,153	0,135	9,225	7,224	6,550
Heavy Duty Vehicles	Diesel RT >32t	Euro IV	2007	2009	316,444	252,503	229,586	1,151	0,839	0,723	0,049	0,037	0,033	6,270	5,071	4,708
Heavy Duty Vehicles	Diesel RT >32t	Euro V	2010	2013	319,806	254,559	231,118	1,166	0,847	0,729	0,049	0,037	0,033	3,735	3,012	2,790
Heavy Duty Vehicles	Diesel TT/AT 28 - 34t	Conventional	0	1993	332,114	254,391	227,288	2,560	1,899	1,804	0,488	0,361	0,339	13,305	10,460	9,286
Heavy Duty Vehicles	Diesel TT/AT 28 - 34t	Euro I	1994	1996	297,033	229,431	205,352	2,173	1,665	1,602	0,380	0,277	0,261	9,509	7,408	6,570
Heavy Duty Vehicles	Diesel TT/AT 28 - 34t	Euro II	1997	2001	280,137	219,605	203,132	1,746	1,372	1,500	0,191	0,152	0,174	10,046	7,771	6,867
Heavy Duty Vehicles	Diesel TT/AT 28 - 34t	Euro III	2002	2006	294,936	228,574	203,723	2,067	1,559	1,515	0,184	0,132	0,120	8,110	6,154	5,397
Heavy Duty Vehicles	Diesel TT/AT 28 - 34t	Euro IV	2007	2009	283,202	224,159	200,624	0,990	0,709	0,618	0,043	0,031	0,028	5,531	4,329	3,837
Heavy Duty Vehicles	Diesel TT/AT 28 - 34t	Euro V	2010	2013	286,144	226,034	202,156	1,002	0,717	0,625	0,043	0,032	0,028	3,297	2,575	2,277
Heavy Duty Vehicles	Diesel TT/AT 28 - 34t	Euro VI	2014	9999	280,404	220,395	196,620	1,033	0,738	0,629	0,004	0,003	0,003	0,531	0,245	0,167
Heavy Duty Vehicles	Diesel TT/AT 34 - 40t	Conventional	0	1993	385,216	290,623	255,748	3,006	2,216	2,091	0,579	0,419	0,384	15,378	11,908	10,419
Heavy Duty Vehicles	Diesel TT/AT 34 - 40t	Euro I	1994	1996	338,164	257,767	227,915	2,561	1,946	1,861	0,464	0,324	0,293	10,891	8,408	7,387
Heavy Duty Vehicles	Diesel TT/AT 34 - 40t	Euro II	1997	2001	329,707	253,289	223,868	2,056	1,607	1,775	0,227	0,177	0,201	11,695	8,978	7,885
Heavy Duty Vehicles	Diesel TT/AT 34 - 40t	Euro III	2002	2006	341,490	259,512	227,377	2,453	1,826	1,775	0,223	0,155	0,136	9,414	7,197	6,354
Heavy Duty Vehicles	Diesel TT/AT 34 - 40t	Euro IV	2007	2009	327,133	254,126	224,236	1,157	0,830	0,704	0,050	0,036	0,032	6,398	5,061	4,523
Heavy Duty Vehicles	Diesel TT/AT 34 - 40t	Euro V	2010	2013	330,656	256,284	225,882	1,168	0,837	0,715	0,050	0,037	0,032	3,814	3,008	2,681

Heavy Duty Vehicles	Diesel TT/AT 34 - 40t	Euro VI	2014	9999	323,739	249,678	219,585	1,202	0,857	0,727	0,005	0,004	0,003	0,610	0,285	0,191
Heavy Duty Vehicles	Diesel TT/AT 40 - 50t	Conventional	0	1993	427,609	323,566	283,490	3,242	2,400	2,283	0,622	0,462	0,425	17,311	13,363	11,617
Heavy Duty Vehicles	Diesel TT/AT 40 - 50t	Euro I	1994	1996	376,029	287,195	252,542	2,823	2,135	2,079	0,500	0,358	0,333	12,142	9,377	8,189
Heavy Duty Vehicles	Diesel TT/AT 40 - 50t	Euro II	1997	2001	364,063	281,631	253,871	2,313	1,826	1,823	0,257	0,201	0,227	12,955	9,936	8,683
Heavy Duty Vehicles	Diesel TT/AT 40 - 50t	Euro III	2002	2006	380,024	289,287	252,570	2,675	1,999	1,959	0,240	0,170	0,146	10,432	7,969	6,995
Heavy Duty Vehicles	Diesel TT/AT 40 - 50t	Euro IV	2007	2009	367,275	285,007	249,788	1,241	0,894	0,759	0,054	0,040	0,035	7,035	5,657	4,952
Heavy Duty Vehicles	Diesel TT/AT 40 - 50t	Euro V	2010	2013	371,248	287,356	251,512	1,256	0,902	0,765	0,055	0,040	0,035	4,187	3,365	2,944
Heavy Duty Vehicles	Diesel TT/AT 40 - 50t	Euro VI	2014	9999	363,184	279,711	244,208	1,288	0,921	0,779	0,006	0,004	0,004	0,564	0,294	0,214
Heavy Duty Vehicles	Diesel TT/AT 50 - 60t	Euro II	1997	2001	439,443	338,240	299,997	2,783	2,192	2,191	0,317	0,246	0,275	15,566	11,836	10,222
Heavy Duty Vehicles	Diesel TT/AT 50 - 60t	Euro IV	2007	2009	440,973	342,585	300,013	1,445	1,038	0,878	0,064	0,047	0,041	8,477	6,746	5,764
Heavy Duty Vehicles	Diesel TT/AT >60t	Euro III	2002	2006	451,915	343,013	304,759	3,169	2,369	2,341	0,278	0,194	0,184	12,548	9,501	8,223
Buses	Gasoline Urban Buses	Conventional	0	9999	225,000	150,000	165,000	70,000	55,000	55,000	0,400	0,400	0,400	4,500	7,500	7,500
Buses	Diesel Urban Buses <15t	Conventional	0	1993	265,880	211,064	197,424	4,479	3,144	2,830	0,729	0,490	0,413	9,347	7,678	7,133
Buses	Diesel Urban Buses <15t	Euro I	1994	1996	214,880	174,564	162,024	1,568	1,120	0,981	0,261	0,199	0,178	6,945	5,531	4,861
Buses	Diesel Urban Buses <15t	Euro II	1997	2001	207,395	170,373	158,652	1,391	0,958	0,806	0,129	0,107	0,103	7,552	5,971	5,224
Buses	Diesel Urban Buses <15t	Euro III	2002	2006	219,770	179,899	167,027	1,509	1,028	0,926	0,130	0,100	0,093	6,425	4,515	3,631
Buses	Diesel Urban Buses <15t	Euro IV	2007	2009	204,146	174,431	172,127	0,800	0,542	0,422	0,032	0,025	0,022	4,076	3,101	2,593
Buses	Diesel Urban Buses <15t	Euro V	2010	2013	207,620	176,864	174,491	0,813	0,551	0,430	0,032	0,025	0,023	2,432	1,845	1,545
Buses	Diesel Urban Buses 15 - 18t	Conventional	0	1993	338,177	261,819	230,080	4,720	3,242	2,606	0,656	0,439	0,351	15,108	12,139	10,803
Buses	Diesel Urban Buses 15 - 18t	Euro I	1994	1996	288,515	228,326	202,771	2,204	1,612	1,330	0,359	0,258	0,226	9,289	7,392	6,426
Buses	Diesel Urban Buses 15 - 18t	Euro II	1997	2001	279,657	224,821	202,070	1,892	1,310	1,120	0,179	0,146	0,137	9,989	7,828	6,822
Buses	Diesel Urban Buses 15 - 18t	Euro III	2002	2006	293,115	235,088	211,025	2,070	1,382	1,257	0,174	0,132	0,115	8,427	6,044	4,919
Buses	Diesel Urban Buses 15 - 18t	Euro IV	2007	2009	276,404	230,306	217,637	1,045	0,709	0,556	0,044	0,033	0,028	5,452	4,181	3,521
Buses	Diesel Urban Buses 15 - 18t	Euro V	2010	2013	280,396	232,974	220,038	1,057	0,716	0,563	0,044	0,033	0,029	3,250	2,486	2,089
Buses	Diesel Urban Buses >18t	Conventional	0	1993	424,462	330,433	285,157	6,145	4,310	3,420	0,833	0,575	0,455	19,310	15,492	13,433
Buses	Diesel Urban Buses >18t	Euro I	1994	1996	369,176	292,254	253,780	2,882	2,132	1,965	0,451	0,336	0,311	11,840	9,361	8,043
Buses	Diesel Urban Buses >18t	Euro II	1997	2001	358,097	288,482	265,154	2,541	1,716	1,467	0,241	0,194	0,178	12,472	9,751	8,334
Buses	Diesel Urban Buses >18t	Euro III	2002	2006	373,469	299,269	262,705	2,691	1,778	1,703	0,209	0,151	0,142	10,561	7,685	6,305
Buses	Diesel Urban Buses >18t	Euro IV	2007	2009	359,379	300,406	272,408	1,287	0,869	0,664	0,054	0,039	0,032	7,106	5,505	4,635
Buses	Diesel Urban Buses >18t	Euro V	2010	2013	363,797	303,565	275,335	1,297	0,877	0,671	0,054	0,039	0,033	4,234	3,275	2,755
Buses	Gasoline Coaches	Conventional	0	9999	225,000	150,000	165,000	70,000	55,000	55,000	0,400	0,400	0,400	4,500	7,500	7,500
Buses	Diesel Coaches <15t	Conventional	0	1993	306,332	225,195	199,049	2,712	1,738	1,372	0,490	0,328	0,269	11,324	8,822	8,156
Buses	Diesel Coaches <15t	Euro I	1994	1996	280,973	207,851	184,178	2,199	1,466	1,186	0,395	0,260	0,209	8,768	6,699	6,147
Buses	Diesel Coaches <15t	Euro II	1997	2001	279,483	208,488	184,973	1,775	1,203	1,092	0,186	0,137	0,120	10,033	7,549	6,840
Buses	Diesel Coaches <15t	Euro III	2002	2006	303,872	224,218	197,656	2,308	1,464	1,283	0,223	0,145	0,115	8,591	6,046	5,368
Buses	Diesel Coaches <15t	Euro IV	2007	2009	290,989	221,962	197,681	1,241	0,813	0,689	0,048	0,034	0,030	5,666	4,225	3,842
Buses	Diesel Coaches <15t	Euro V	2010	2013	298,215	226,393	200,893	1,288	0,842	0,696	0,049	0,034	0,030	3,434	2,544	2,291

Buses	Diesel Coaches 15 - 18t	Conventional	0	1993	306,332	225,195	199,049	2,712	1,738	1,372	0,490	0,328	0,269	11,324	8,822	8,156
Buses	Diesel Coaches 15 - 18t	Euro I	1994	1996	280,973	207,851	184,178	2,199	1,466	1,186	0,395	0,260	0,209	8,768	6,699	6,147
Buses	Diesel Coaches 15 - 18t	Euro II	1997	2001	279,483	208,488	184,973	1,775	1,203	1,092	0,186	0,137	0,120	10,033	7,549	6,840
Buses	Diesel Coaches 15 - 18t	Euro III	2002	2006	303,872	224,218	197,656	2,308	1,464	1,283	0,223	0,145	0,115	8,591	6,046	5,368
Buses	Diesel Coaches 15 - 18t	Euro IV	2007	2009	290,989	221,962	197,681	1,241	0,813	0,689	0,048	0,034	0,030	5,666	4,225	3,842
Buses	Diesel Coaches 15 - 18t	Euro V	2010	2013	298,215	226,393	200,893	1,288	0,842	0,696	0,049	0,034	0,030	3,434	2,544	2,291
Buses	Diesel Coaches >18t	Conventional	0	1993	371,932	272,817	240,539	3,104	2,042	1,732	0,572	0,388	0,331	14,084	10,772	9,735
Buses	Diesel Coaches >18t	Euro I	1994	1996	329,598	243,565	215,080	2,511	1,722	1,458	0,452	0,302	0,246	10,737	8,049	7,206
Buses	Diesel Coaches >18t	Euro II	1997	2001	323,939	241,571	213,608	2,031	1,395	1,290	0,214	0,161	0,143	11,883	8,817	7,837
Buses	Diesel Coaches >18t	Euro III	2002	2006	335,657	242,331	211,644	2,557	1,669	1,439	0,242	0,156	0,126	9,681	6,781	5,889
Buses	Diesel Coaches >18t	Euro IV	2007	2009	319,737	238,136	211,184	1,328	0,875	0,742	0,052	0,036	0,032	6,428	4,728	4,226
Buses	Diesel Coaches >18t	Euro V	2010	2013	328,400	243,537	215,269	1,363	0,896	0,758	0,053	0,037	0,032	3,881	2,845	2,536
Mopeds	2-stroke <50 cm ³	Conventional	0	1999	25,000	25,000	0,000	14,700	14,700	0,000	0,176	0,176	0,000	0,056	0,056	0,000
Mopeds	2-stroke <50 cm ³	Euro I	2000	2003	20,000	20,000	0,000	4,600	4,600	0,000	0,045	0,045	0,000	0,180	0,180	0,000
Mopeds	2-stroke <50 cm ³	Euro II	2004	2014	20,000	20,000	0,000	2,800	2,800	0,000	0,026	0,026	0,000	0,170	0,170	0,000
Mopeds	4-stroke <50 cm ³	Euro II	2004	2014	20,000	20,000	0,000	4,200	4,200	0,000	0,007	0,007	0,000	0,170	0,170	0,000
Motorcycles	2-stroke >50 cm ³	Conventional	0	1999	27,115	28,317	39,640	15,605	19,285	28,470	0,200	0,200	0,200	0,029	0,030	0,035
Motorcycles	2-stroke >50 cm ³	Euro I	2000	2003	27,115	28,317	39,640	10,315	12,786	18,933	0,080	0,080	0,080	0,029	0,030	0,035
Motorcycles	2-stroke >50 cm ³	Euro II	2004	2006	24,892	25,627	35,438	8,146	10,067	14,890	0,040	0,040	0,040	0,040	0,050	0,060
Motorcycles	2-stroke >50 cm ³	Euro III	2007	9999	24,892	25,627	35,438	4,510	5,593	8,342	0,012	0,012	0,012	0,048	0,058	0,069
Motorcycles	4-stroke <250 cm ³	Conventional	0	1999	24,800	27,499	36,055	15,258	17,209	24,960	0,020	0,020	0,020	0,237	0,428	0,655
Motorcycles	4-stroke <250 cm ³	Euro I	2000	2003	27,015	30,386	40,330	10,391	14,456	24,910	0,020	0,020	0,020	0,304	0,424	0,567
Motorcycles	4-stroke <250 cm ³	Euro II	2004	2006	22,260	25,160	33,756	3,708	5,765	9,135	0,005	0,005	0,005	0,323	0,447	0,598
Motorcycles	4-stroke <250 cm ³	Euro III	2007	9999	19,262	20,359	25,932	2,060	3,201	5,092	0,005	0,005	0,005	0,253	0,382	0,612
Motorcycles	4-stroke 250 - 750 cm ³	Conventional	0	1999	26,648	23,766	26,620	20,461	19,486	22,990	0,020	0,020	0,020	0,196	0,300	0,548
Motorcycles	4-stroke 250 - 750 cm ³	Euro I	2000	2003	37,374	35,472	41,400	10,599	9,003	10,460	0,020	0,020	0,020	0,258	0,400	0,610
Motorcycles	4-stroke 250 - 750 cm ³	Euro II	2004	2006	34,197	33,450	41,276	2,230	2,436	6,092	0,005	0,005	0,005	0,257	0,390	0,577
Motorcycles	4-stroke 250 - 750 cm ³	Euro III	2007	9999	30,983	30,719	38,129	1,228	1,345	3,357	0,005	0,005	0,005	0,076	0,132	0,265
Motorcycles	4-stroke >750 cm ³	Conventional	0	1999	35,731	35,542	43,748	20,461	19,486	22,990	0,020	0,020	0,020	0,019	0,030	0,086
Motorcycles	4-stroke >750 cm ³	Euro I	2000	2003	43,101	41,041	47,500	10,599	9,003	10,460	0,020	0,020	0,020	0,125	0,178	0,392
Motorcycles	4-stroke >750 cm ³	Euro II	2004	2006	42,110	38,004	41,895	2,230	2,436	6,092	0,005	0,005	0,005	0,143	0,244	0,459
Motorcycles	4-stroke >750 cm ³	Euro III	2007	9999	40,343	37,470	43,083	1,228	1,345	3,357	0,005	0,005	0,005	0,104	0,200	0,484

Sector	Subsector	Tech 2	FYear	LYear	CH4u	CH4r	CH4h	N2Ou	N2Or	N2Oh	NH3u	NH3r	NH3h	VOCu	VOCr	VOCh
Passenger Cars	Gasoline 0,8 - 1,4 l	PRE ECE	0	1969	0,092	0,029	0,026	0,010	0,007	0,007	0,002	0,002	0,002	2,354	1,597	1,247
Passenger Cars	Gasoline 0,8 - 1,4 l	ECE 15/00-01	1970	1978	0,092	0,029	0,026	0,010	0,007	0,007	0,002	0,002	0,002	1,862	1,256	1,121
Passenger Cars	Gasoline 0,8 - 1,4 l	ECE 15/02	1979	1980	0,092	0,029	0,026	0,010	0,007	0,007	0,002	0,002	0,002	1,849	1,061	0,950
Passenger Cars	Gasoline 0,8 - 1,4 l	ECE 15/03	1981	1985	0,092	0,029	0,026	0,010	0,007	0,007	0,002	0,002	0,002	1,849	1,061	0,950
Passenger Cars	Gasoline 0,8 - 1,4 l	ECE 15/04	1986	1990	0,092	0,029	0,026	0,010	0,007	0,007	0,002	0,002	0,002	1,480	0,895	0,698
Passenger Cars	Gasoline 0,8 - 1,4 l	Euro I	1991	1996	0,026	0,016	0,014	0,026	0,011	0,005	0,070	0,133	0,074	0,177	0,121	0,111
Passenger Cars	Gasoline 0,8 - 1,4 l	Euro II	1997	2000	0,017	0,013	0,011	0,013	0,005	0,003	0,180	0,150	0,084	0,071	0,047	0,042
Passenger Cars	Gasoline 0,8 - 1,4 l	Euro III	2001	2005	0,003	0,002	0,004	0,001	0,000	0,000	0,002	0,030	0,065	0,015	0,015	0,025
Passenger Cars	Gasoline 0,8 - 1,4 l	Euro IV	2006	2010	0,003	0,003	0,005	0,002	0,000	0,000	0,002	0,029	0,065	0,012	0,014	0,017
Passenger Cars	Gasoline 0,8 - 1,4 l	Euro V	2011	2014	0,003	0,003	0,005	0,002	0,000	0,000	0,002	0,029	0,065	0,012	0,014	0,017
Passenger Cars	Gasoline 1,4 - 2,0 l	PRE ECE	0	1969	0,092	0,029	0,026	0,010	0,007	0,007	0,002	0,002	0,002	2,354	1,597	1,247
Passenger Cars	Gasoline 1,4 - 2,0 l	ECE 15/00-01	1970	1978	0,092	0,029	0,026	0,010	0,007	0,007	0,002	0,002	0,002	1,862	1,256	1,121
Passenger Cars	Gasoline 1,4 - 2,0 l	ECE 15/02	1979	1980	0,092	0,029	0,026	0,010	0,007	0,007	0,002	0,002	0,002	1,849	1,061	0,950
Passenger Cars	Gasoline 1,4 - 2,0 l	ECE 15/03	1981	1985	0,092	0,029	0,026	0,010	0,007	0,007	0,002	0,002	0,002	1,849	1,061	0,950
Passenger Cars	Gasoline 1,4 - 2,0 l	ECE 15/04	1986	1990	0,092	0,029	0,026	0,010	0,007	0,007	0,002	0,002	0,002	1,480	0,895	0,698
Passenger Cars	Gasoline 1,4 - 2,0 l	Euro I	1991	1996	0,026	0,016	0,014	0,026	0,011	0,005	0,070	0,133	0,074	0,177	0,121	0,111
Passenger Cars	Gasoline 1,4 - 2,0 l	Euro II	1997	2000	0,017	0,013	0,011	0,013	0,005	0,003	0,188	0,150	0,084	0,071	0,047	0,042
Passenger Cars	Gasoline 1,4 - 2,0 l	Euro III	2001	2005	0,003	0,002	0,004	0,002	0,000	0,000	0,002	0,030	0,065	0,015	0,015	0,025
Passenger Cars	Gasoline 1,4 - 2,0 l	Euro IV	2006	2010	0,003	0,003	0,005	0,002	0,000	0,000	0,002	0,029	0,065	0,012	0,014	0,017
Passenger Cars	Gasoline 1,4 - 2,0 l	Euro V	2011	2014	0,003	0,003	0,005	0,002	0,000	0,000	0,002	0,029	0,065	0,012	0,014	0,017
Passenger Cars	Gasoline >2,0 l	PRE ECE	0	1969	0,092	0,029	0,026	0,010	0,007	0,007	0,002	0,002	0,002	2,354	1,597	1,247
Passenger Cars	Gasoline >2,0 l	ECE 15/00-01	1970	1978	0,092	0,029	0,026	0,010	0,007	0,007	0,002	0,002	0,002	1,862	1,256	1,121
Passenger Cars	Gasoline >2,0 l	ECE 15/02	1979	1980	0,092	0,029	0,026	0,010	0,007	0,007	0,002	0,002	0,002	1,849	1,061	0,950
Passenger Cars	Gasoline >2,0 l	ECE 15/03	1981	1985	0,092	0,029	0,026	0,010	0,007	0,007	0,002	0,002	0,002	1,849	1,061	0,950
Passenger Cars	Gasoline >2,0 l	ECE 15/04	1986	1990	0,092	0,029	0,026	0,010	0,007	0,007	0,002	0,002	0,002	1,480	0,895	0,698
Passenger Cars	Gasoline >2,0 l	Euro I	1991	1996	0,026	0,016	0,014	0,026	0,011	0,005	0,070	0,133	0,074	0,177	0,121	0,111
Passenger Cars	Gasoline >2,0 l	Euro II	1997	2000	0,017	0,013	0,011	0,013	0,005	0,003	0,190	0,150	0,084	0,071	0,047	0,042
Passenger Cars	Gasoline >2,0 l	Euro III	2001	2005	0,003	0,002	0,004	0,002	0,000	0,000	0,002	0,030	0,065	0,015	0,015	0,025
Passenger Cars	Gasoline >2,0 l	Euro IV	2006	2010	0,003	0,003	0,005	0,002	0,000	0,000	0,002	0,030	0,065	0,012	0,014	0,017
Passenger Cars	Gasoline >2,0 l	Euro V	2011	2014	0,003	0,003	0,005	0,002	0,000	0,000	0,002	0,029	0,065	0,012	0,014	0,017
Passenger Cars	Diesel 1,4 - 2,0 l	Conventional	0	1990	0,028	0,012	0,008	0,000	0,000	0,000	0,001	0,001	0,001	0,145	0,086	0,062
Passenger Cars	Diesel 1,4 - 2,0 l	Euro I	1991	1996	0,011	0,009	0,003	0,002	0,004	0,004	0,001	0,001	0,001	0,053	0,031	0,026
Passenger Cars	Diesel 1,4 - 2,0 l	Euro II	1997	2000	0,007	0,003	0,002	0,004	0,006	0,006	0,001	0,001	0,001	0,034	0,021	0,015
Passenger Cars	Diesel 1,4 - 2,0 l	Euro III	2001	2005	0,003	0,000	0,000	0,009	0,004	0,004	0,001	0,001	0,001	0,018	0,011	0,009
Passenger Cars	Diesel 1,4 - 2,0 l	Euro IV	2006	2010	0,000	0,000	0,000	0,009	0,004	0,004	0,001	0,001	0,001	0,011	0,006	0,006
Passenger Cars	Diesel 1,4 - 2,0 l	Euro V	2011	2014	0,000	0,000	0,000	0,009	0,004	0,004	0,001	0,001	0,001	0,011	0,006	0,006

Passenger Cars	Diesel >2,0 l	Conventional	0	1990	0,028	0,012	0,008	0,000	0,000	0,000	0,001	0,001	0,001	0,145	0,086	0,062
Passenger Cars	Diesel >2,0 l	Euro I	1991	1996	0,011	0,009	0,003	0,002	0,004	0,004	0,001	0,001	0,001	0,080	0,046	0,034
Passenger Cars	Diesel >2,0 l	Euro II	1997	2000	0,007	0,003	0,002	0,004	0,006	0,006	0,001	0,001	0,001	0,098	0,058	0,038
Passenger Cars	Diesel >2,0 l	Euro III	2001	2005	0,003	0,000	0,000	0,009	0,004	0,004	0,001	0,001	0,001	0,038	0,017	0,012
Passenger Cars	Diesel >2,0 l	Euro IV	2006	2010	0,000	0,000	0,000	0,009	0,004	0,004	0,001	0,001	0,001	0,011	0,006	0,006
Passenger Cars	Diesel >2,0 l	Euro V	2011	2014	0,000	0,000	0,000	0,009	0,004	0,004	0,001	0,001	0,001	0,011	0,006	0,006
Passenger Cars	LPG cars	Conventional	0	1990	0,080	0,035	0,025	0,000	0,000	0,000	0,000	0,000	0,000	1,082	0,667	0,490
Passenger Cars	LPG cars	Euro I	1991	1996	0,080	0,035	0,025	0,021	0,013	0,008	0,000	0,000	0,000	0,239	0,071	0,083
Passenger Cars	LPG cars	Euro II	1997	2000	0,019	0,008	0,006	0,013	0,003	0,002	0,000	0,000	0,000	0,050	0,015	0,017
Passenger Cars	LPG cars	Euro III	2001	2005	0,013	0,006	0,004	0,005	0,002	0,001	0,000	0,000	0,000	0,036	0,011	0,012
Passenger Cars	LPG cars	Euro IV	2006	2010	0,004	0,002	0,001	0,005	0,002	0,001	0,000	0,000	0,000	0,007	0,002	0,002
Passenger Cars	Gasoline <0,8 l	PRE ECE	0	1969	0,092	0,029	0,026	0,010	0,007	0,007	0,002	0,002	0,002	2,354	1,597	1,247
Passenger Cars	Gasoline <0,8 l	ECE 15/00-01	1970	1978	0,092	0,029	0,026	0,010	0,007	0,007	0,002	0,002	0,002	1,862	1,256	1,121
Passenger Cars	Gasoline <0,8 l	ECE 15/02	1979	1980	0,092	0,029	0,026	0,010	0,007	0,007	0,002	0,002	0,002	1,849	1,061	0,950
Passenger Cars	Gasoline <0,8 l	ECE 15/03	1981	1985	0,092	0,029	0,026	0,010	0,007	0,007	0,002	0,002	0,002	1,849	1,061	0,950
Passenger Cars	Gasoline <0,8 l	ECE 15/04	1986	1990	0,092	0,029	0,026	0,010	0,007	0,007	0,002	0,002	0,002	1,480	0,895	0,698
Passenger Cars	Gasoline <0,8 l	Euro I	1991	1996	0,026	0,016	0,014	0,023	0,009	0,004	0,070	0,132	0,074	0,177	0,121	0,111
Passenger Cars	Gasoline <0,8 l	Euro II	1997	2000	0,017	0,013	0,011	0,012	0,005	0,003	0,173	0,149	0,084	0,071	0,047	0,042
Passenger Cars	Gasoline <0,8 l	Euro III	2001	2005	0,003	0,002	0,004	0,001	0,000	0,000	0,002	0,030	0,065	0,015	0,015	0,025
Passenger Cars	Gasoline <0,8 l	Euro IV	2006	2010	0,003	0,003	0,005	0,002	0,000	0,000	0,002	0,029	0,065	0,012	0,014	0,017
Passenger Cars	Gasoline <0,8 l	Euro V	2011	2014	0,003	0,003	0,005	0,002	0,000	0,000	0,002	0,029	0,065	0,012	0,014	0,017
Passenger Cars	Diesel <1,4 l	Conventional	0	1990	0,028	0,012	0,008	0,000	0,000	0,000	0,001	0,001	0,001	0,145	0,086	0,062
Passenger Cars	Diesel <1,4 l	Euro I	1991	1996	0,011	0,009	0,003	0,002	0,004	0,004	0,001	0,001	0,001	0,053	0,031	0,026
Passenger Cars	Diesel <1,4 l	Euro II	1997	2000	0,007	0,003	0,002	0,004	0,006	0,006	0,001	0,001	0,001	0,034	0,021	0,015
Passenger Cars	Diesel <1,4 l	Euro III	2001	2005	0,003	0,000	0,000	0,009	0,004	0,004	0,001	0,001	0,001	0,018	0,011	0,009
Passenger Cars	Diesel <1,4 l	Euro IV	2006	2010	0,000	0,000	0,000	0,009	0,004	0,004	0,001	0,001	0,001	0,011	0,006	0,006
Passenger Cars	Diesel <1,4 l	Euro V	2011	2014	0,000	0,000	0,000	0,009	0,004	0,004	0,001	0,001	0,001	0,011	0,006	0,006
Light Duty Vehicles	Gasoline <3,5t	Conventional	0	1994	0,150	0,040	0,025	0,010	0,007	0,007	0,002	0,002	0,002	1,877	0,729	0,446
Light Duty Vehicles	Gasoline <3,5t	Euro I	1995	1998	0,026	0,016	0,014	0,045	0,027	0,013	0,070	0,133	0,074	0,220	0,109	0,078
Light Duty Vehicles	Gasoline <3,5t	Euro II	1999	2001	0,017	0,013	0,011	0,028	0,016	0,010	0,181	0,150	0,084	0,053	0,026	0,019
Light Duty Vehicles	Gasoline <3,5t	Euro III	2002	2006	0,003	0,002	0,004	0,008	0,001	0,001	0,002	0,030	0,065	0,031	0,015	0,011
Light Duty Vehicles	Gasoline <3,5t	Euro IV	2007	2011	0,002	0,002	0,000	0,001	0,000	0,000	0,002	0,029	0,065	0,013	0,007	0,005
Light Duty Vehicles	Gasoline <3,5t	Euro V	2012	2015	0,002	0,002	0,000	0,001	0,000	0,000	0,002	0,029	0,065	0,013	0,007	0,005
Light Duty Vehicles	Diesel <3,5t	Conventional	0	1994	0,028	0,012	0,008	0,000	0,000	0,000	0,001	0,001	0,001	0,131	0,106	0,101
Light Duty Vehicles	Diesel <3,5t	Euro I	1995	1998	0,011	0,009	0,003	0,002	0,004	0,004	0,001	0,001	0,001	0,131	0,106	0,101
Light Duty Vehicles	Diesel <3,5t	Euro II	1999	2001	0,007	0,003	0,002	0,004	0,006	0,006	0,001	0,001	0,001	0,131	0,106	0,101
Light Duty Vehicles	Diesel <3,5t	Euro III	2002	2006	0,003	0,000	0,000	0,009	0,004	0,004	0,001	0,001	0,001	0,081	0,065	0,063

Light Duty Vehicles	Diesel <3,5t	Euro IV	2007	2011	0,000	0,000	0,000	0,009	0,004	0,004	0,001	0,001	0,001	0,030	0,024	0,023
Light Duty Vehicles	Diesel <3,5t	Euro V	2012	2015	0,000	0,000	0,000	0,009	0,004	0,004	0,001	0,001	0,001	0,030	0,024	0,023
Light Duty Vehicles	LPG <3,5t	Conventional	0	1994	0,120	0,053	0,038	0,000	0,000	0,000	0,000	0,000	0,000	1,623	1,000	0,735
Light Duty Vehicles	LPG <3,5t	Euro II	1999	2001	0,029	0,013	0,009	0,020	0,005	0,003	0,000	0,000	0,000	0,075	0,022	0,026
Light Duty Vehicles	LPG <3,5t	Euro III	2002	2006	0,019	0,008	0,006	0,008	0,003	0,002	0,000	0,000	0,000	0,054	0,016	0,019
Light Duty Vehicles	LPG <3,5t	Euro IV	2007	2011	0,006	0,003	0,002	0,008	0,003	0,002	0,000	0,000	0,000	0,011	0,003	0,004
Heavy Duty Vehicles	Gasoline >3,5t	Conventional	0	9999	0,140	0,110	0,070	0,006	0,006	0,006	0,002	0,002	0,002	7,000	5,500	3,500
Heavy Duty Vehicles	Diesel RT 3,5 - 7,5t	Conventional	0	1993	0,085	0,023	0,020	0,030	0,030	0,030	0,003	0,003	0,003	1,298	0,789	0,576
Heavy Duty Vehicles	Diesel RT 3,5 - 7,5t	Euro I	1994	1996	0,085	0,023	0,020	0,030	0,030	0,030	0,003	0,003	0,003	0,253	0,167	0,130
Heavy Duty Vehicles	Diesel RT 3,5 - 7,5t	Euro II	1997	2001	0,054	0,020	0,019	0,030	0,030	0,030	0,003	0,003	0,003	0,171	0,111	0,086
Heavy Duty Vehicles	Diesel RT 3,5 - 7,5t	Euro III	2002	2006	0,048	0,021	0,018	0,030	0,030	0,030	0,003	0,003	0,003	0,162	0,102	0,077
Heavy Duty Vehicles	Diesel RT 3,5 - 7,5t	Euro IV	2007	2009	0,003	0,002	0,001	0,030	0,030	0,030	0,003	0,003	0,003	0,022	0,017	0,017
Heavy Duty Vehicles	Diesel RT 3,5 - 7,5t	Euro V	2010	2013	0,003	0,002	0,001	0,030	0,030	0,030	0,003	0,003	0,003	0,022	0,017	0,017
Heavy Duty Vehicles	Diesel RT 3,5 - 7,5t	Euro VI	2014	9999	0,001	0,000	0,000	0,030	0,030	0,030	0,003	0,003	0,003	0,011	0,008	0,007
Heavy Duty Vehicles	Diesel RT 7,5 - 12t	Conventional	0	1993	0,085	0,023	0,020	0,030	0,030	0,030	0,003	0,003	0,003	0,957	0,589	0,449
Heavy Duty Vehicles	Diesel RT 7,5 - 12t	Euro I	1994	1996	0,085	0,023	0,020	0,030	0,030	0,030	0,003	0,003	0,003	0,389	0,258	0,208
Heavy Duty Vehicles	Diesel RT 7,5 - 12t	Euro II	1997	2001	0,054	0,020	0,019	0,030	0,030	0,030	0,003	0,003	0,003	0,263	0,172	0,137
Heavy Duty Vehicles	Diesel RT 7,5 - 12t	Euro III	2002	2006	0,048	0,021	0,018	0,030	0,030	0,030	0,003	0,003	0,003	0,252	0,157	0,120
Heavy Duty Vehicles	Diesel RT 7,5 - 12t	Euro IV	2007	2009	0,003	0,002	0,001	0,030	0,030	0,030	0,003	0,003	0,003	0,035	0,025	0,022
Heavy Duty Vehicles	Diesel RT 7,5 - 12t	Euro V	2010	2013	0,003	0,002	0,001	0,030	0,030	0,030	0,003	0,003	0,003	0,035	0,026	0,022
Heavy Duty Vehicles	Diesel RT 7,5 - 12t	Euro VI	2014	9999	0,001	0,000	0,000	0,030	0,030	0,030	0,003	0,003	0,003	0,017	0,012	0,010
Heavy Duty Vehicles	Diesel RT 12 - 14 t	Conventional	0	1993	0,085	0,023	0,020	0,030	0,030	0,030	0,003	0,003	0,003	1,012	0,646	0,509
Heavy Duty Vehicles	Diesel RT 12 - 14 t	Euro I	1994	1996	0,085	0,023	0,020	0,030	0,030	0,030	0,003	0,003	0,003	0,429	0,279	0,229
Heavy Duty Vehicles	Diesel RT 12 - 14 t	Euro II	1997	2001	0,054	0,020	0,019	0,030	0,030	0,030	0,003	0,003	0,003	0,281	0,186	0,150
Heavy Duty Vehicles	Diesel RT 12 - 14 t	Euro III	2002	2006	0,048	0,021	0,018	0,030	0,030	0,030	0,003	0,003	0,003	0,260	0,168	0,134
Heavy Duty Vehicles	Diesel RT 12 - 14 t	Euro IV	2007	2009	0,003	0,002	0,001	0,030	0,030	0,030	0,003	0,003	0,003	0,034	0,025	0,024
Heavy Duty Vehicles	Diesel RT 12 - 14 t	Euro V	2010	2013	0,003	0,002	0,001	0,030	0,030	0,030	0,003	0,003	0,003	0,034	0,025	0,024
Heavy Duty Vehicles	Diesel RT 14 - 20t	Conventional	0	1993	0,175	0,080	0,070	0,030	0,030	0,030	0,003	0,003	0,003	1,510	0,971	0,768
Heavy Duty Vehicles	Diesel RT 14 - 20t	Euro I	1994	1996	0,175	0,080	0,070	0,030	0,030	0,030	0,003	0,003	0,003	0,606	0,403	0,325
Heavy Duty Vehicles	Diesel RT 14 - 20t	Euro II	1997	2001	0,112	0,070	0,065	0,030	0,030	0,030	0,003	0,003	0,003	0,409	0,267	0,213
Heavy Duty Vehicles	Diesel RT 14 - 20t	Euro III	2002	2006	0,098	0,074	0,064	0,030	0,030	0,030	0,003	0,003	0,003	0,378	0,243	0,196
Heavy Duty Vehicles	Diesel RT 14 - 20t	Euro IV	2007	2009	0,005	0,006	0,004	0,030	0,030	0,030	0,003	0,003	0,003	0,046	0,032	0,028
Heavy Duty Vehicles	Diesel RT 14 - 20t	Euro V	2010	2013	0,005	0,006	0,004	0,030	0,030	0,030	0,003	0,003	0,003	0,047	0,033	0,029
Heavy Duty Vehicles	Diesel RT 14 - 20t	Euro VI	2014	9999	0,002	0,002	0,001	0,030	0,030	0,030	0,003	0,003	0,003	0,024	0,017	0,015
Heavy Duty Vehicles	Diesel RT 20 - 26t	Conventional	0	1993	0,175	0,080	0,070	0,030	0,030	0,030	0,003	0,003	0,003	0,819	0,517	0,406
Heavy Duty Vehicles	Diesel RT 20 - 26t	Euro I	1994	1996	0,175	0,080	0,070	0,030	0,030	0,030	0,003	0,003	0,003	0,728	0,476	0,380
Heavy Duty Vehicles	Diesel RT 20 - 26t	Euro II	1997	2001	0,112	0,070	0,065	0,030	0,030	0,030	0,003	0,003	0,003	0,489	0,314	0,248

Heavy Duty Vehicles	Diesel RT 20 - 26t	Euro III	2002	2006	0,098	0,074	0,064	0,030	0,030	0,030	0,003	0,003	0,003	0,453	0,287	0,225
Heavy Duty Vehicles	Diesel RT 20 - 26t	Euro IV	2007	2009	0,005	0,006	0,004	0,030	0,030	0,030	0,003	0,003	0,003	0,059	0,040	0,035
Heavy Duty Vehicles	Diesel RT 20 - 26t	Euro V	2010	2013	0,005	0,006	0,004	0,030	0,030	0,030	0,003	0,003	0,003	0,059	0,041	0,035
Heavy Duty Vehicles	Diesel RT 20 - 26t	Euro VI	2014	9999	0,002	0,002	0,001	0,030	0,030	0,030	0,003	0,003	0,003	0,030	0,021	0,017
Heavy Duty Vehicles	Diesel RT 26 - 28t	Conventional	0	1993	0,175	0,080	0,070	0,030	0,030	0,030	0,003	0,003	0,003	0,842	0,541	0,430
Heavy Duty Vehicles	Diesel RT 26 - 28t	Euro I	1994	1996	0,175	0,080	0,070	0,030	0,030	0,030	0,003	0,003	0,003	0,736	0,488	0,394
Heavy Duty Vehicles	Diesel RT 26 - 28t	Euro II	1997	2001	0,112	0,070	0,065	0,030	0,030	0,030	0,003	0,003	0,003	0,499	0,327	0,262
Heavy Duty Vehicles	Diesel RT 26 - 28t	Euro III	2002	2006	0,098	0,074	0,064	0,030	0,030	0,030	0,003	0,003	0,003	0,467	0,304	0,243
Heavy Duty Vehicles	Diesel RT 26 - 28t	Euro IV	2007	2009	0,005	0,006	0,004	0,030	0,030	0,030	0,003	0,003	0,003	0,064	0,045	0,037
Heavy Duty Vehicles	Diesel RT 26 - 28t	Euro V	2010	2013	0,005	0,006	0,004	0,030	0,030	0,030	0,003	0,003	0,003	0,065	0,045	0,037
Heavy Duty Vehicles	Diesel RT 28 - 32t	Conventional	0	1993	0,175	0,080	0,070	0,030	0,030	0,030	0,003	0,003	0,003	0,874	0,560	0,444
Heavy Duty Vehicles	Diesel RT 28 - 32t	Euro I	1994	1996	0,175	0,080	0,070	0,030	0,030	0,030	0,003	0,003	0,003	0,778	0,518	0,419
Heavy Duty Vehicles	Diesel RT 28 - 32t	Euro II	1997	2001	0,112	0,070	0,065	0,030	0,030	0,030	0,003	0,003	0,003	0,523	0,344	0,276
Heavy Duty Vehicles	Diesel RT 28 - 32t	Euro III	2002	2006	0,098	0,074	0,064	0,030	0,030	0,030	0,003	0,003	0,003	0,491	0,317	0,252
Heavy Duty Vehicles	Diesel RT 28 - 32t	Euro IV	2007	2009	0,005	0,006	0,004	0,030	0,030	0,030	0,003	0,003	0,003	0,070	0,051	0,043
Heavy Duty Vehicles	Diesel RT 28 - 32t	Euro V	2010	2013	0,005	0,006	0,004	0,030	0,030	0,030	0,003	0,003	0,003	0,071	0,051	0,043
Heavy Duty Vehicles	Diesel RT 28 - 32t	Euro VI	2014	9999	0,002	0,002	0,001	0,030	0,030	0,030	0,003	0,003	0,003	0,034	0,024	0,020
Heavy Duty Vehicles	Diesel RT >32t	Euro I	1994	1996	0,175	0,080	0,070	0,030	0,030	0,030	0,003	0,003	0,003	0,812	0,527	0,419
Heavy Duty Vehicles	Diesel RT >32t	Euro III	2002	2006	0,098	0,074	0,064	0,030	0,030	0,030	0,003	0,003	0,003	0,496	0,316	0,249
Heavy Duty Vehicles	Diesel RT >32t	Euro IV	2007	2009	0,005	0,006	0,004	0,030	0,030	0,030	0,003	0,003	0,003	0,070	0,048	0,041
Heavy Duty Vehicles	Diesel RT >32t	Euro V	2010	2013	0,005	0,006	0,004	0,030	0,030	0,030	0,003	0,003	0,003	0,070	0,049	0,041
Heavy Duty Vehicles	Diesel TT/AT 28 - 34t	Conventional	0	1993	0,175	0,080	0,070	0,030	0,030	0,030	0,003	0,003	0,003	0,736	0,476	0,380
Heavy Duty Vehicles	Diesel TT/AT 28 - 34t	Euro I	1994	1996	0,175	0,080	0,070	0,030	0,030	0,030	0,003	0,003	0,003	0,678	0,450	0,363
Heavy Duty Vehicles	Diesel TT/AT 28 - 34t	Euro II	1997	2001	0,112	0,070	0,065	0,030	0,030	0,030	0,003	0,003	0,003	0,450	0,296	0,238
Heavy Duty Vehicles	Diesel TT/AT 28 - 34t	Euro III	2002	2006	0,098	0,074	0,064	0,030	0,030	0,030	0,003	0,003	0,003	0,415	0,269	0,215
Heavy Duty Vehicles	Diesel TT/AT 28 - 34t	Euro IV	2007	2009	0,005	0,006	0,004	0,030	0,030	0,030	0,003	0,003	0,003	0,059	0,041	0,036
Heavy Duty Vehicles	Diesel TT/AT 28 - 34t	Euro V	2010	2013	0,005	0,006	0,004	0,030	0,030	0,030	0,003	0,003	0,003	0,060	0,042	0,036
Heavy Duty Vehicles	Diesel TT/AT 28 - 34t	Euro VI	2014	9999	0,002	0,002	0,001	0,030	0,030	0,030	0,003	0,003	0,003	0,029	0,021	0,017
Heavy Duty Vehicles	Diesel TT/AT 34 - 40t	Conventional	0	1993	0,175	0,080	0,070	0,030	0,030	0,030	0,003	0,003	0,003	0,877	0,555	0,438
Heavy Duty Vehicles	Diesel TT/AT 34 - 40t	Euro I	1994	1996	0,175	0,080	0,070	0,030	0,030	0,030	0,003	0,003	0,003	0,805	0,524	0,420
Heavy Duty Vehicles	Diesel TT/AT 34 - 40t	Euro II	1997	2001	0,112	0,070	0,065	0,030	0,030	0,030	0,003	0,003	0,003	0,538	0,343	0,270
Heavy Duty Vehicles	Diesel TT/AT 34 - 40t	Euro III	2002	2006	0,098	0,074	0,064	0,030	0,030	0,030	0,003	0,003	0,003	0,494	0,312	0,244
Heavy Duty Vehicles	Diesel TT/AT 34 - 40t	Euro IV	2007	2009	0,005	0,006	0,004	0,030	0,030	0,030	0,003	0,003	0,003	0,071	0,048	0,041
Heavy Duty Vehicles	Diesel TT/AT 34 - 40t	Euro V	2010	2013	0,005	0,006	0,004	0,030	0,030	0,030	0,003	0,003	0,003	0,072	0,049	0,041
Heavy Duty Vehicles	Diesel TT/AT 34 - 40t	Euro VI	2014	9999	0,002	0,002	0,001	0,030	0,030	0,030	0,003	0,003	0,003	0,035	0,024	0,020
Heavy Duty Vehicles	Diesel TT/AT 40 - 50t	Conventional	0	1993	0,175	0,080	0,070	0,030	0,030	0,030	0,003	0,003	0,003	0,901	0,570	0,450
Heavy Duty Vehicles	Diesel TT/AT 40 - 50t	Euro I	1994	1996	0,175	0,080	0,070	0,030	0,030	0,030	0,003	0,003	0,003	0,844	0,546	0,433

Heavy Duty Vehicles	Diesel TT/AT 40 - 50t	Euro II	1997	2001	0,112	0,070	0,065	0,030	0,030	0,030	0,003	0,003	0,003	0,558	0,358	0,282
Heavy Duty Vehicles	Diesel TT/AT 40 - 50t	Euro III	2002	2006	0,098	0,074	0,064	0,030	0,030	0,030	0,003	0,003	0,003	0,510	0,323	0,253
Heavy Duty Vehicles	Diesel TT/AT 40 - 50t	Euro IV	2007	2009	0,005	0,006	0,004	0,030	0,030	0,030	0,003	0,003	0,003	0,077	0,053	0,045
Heavy Duty Vehicles	Diesel TT/AT 40 - 50t	Euro V	2010	2013	0,005	0,006	0,004	0,030	0,030	0,030	0,003	0,003	0,003	0,078	0,053	0,045
Heavy Duty Vehicles	Diesel TT/AT 40 - 50t	Euro VI	2014	9999	0,002	0,002	0,001	0,030	0,030	0,030	0,003	0,003	0,003	0,038	0,026	0,022
Heavy Duty Vehicles	Diesel TT/AT 50 - 60t	Euro II	1997	2001	0,112	0,070	0,065	0,030	0,030	0,030	0,003	0,003	0,003	0,626	0,406	0,323
Heavy Duty Vehicles	Diesel TT/AT 50 - 60t	Euro IV	2007	2009	0,005	0,006	0,004	0,030	0,030	0,030	0,003	0,003	0,003	0,090	0,063	0,053
Heavy Duty Vehicles	Diesel TT/AT >60t	Euro III	2002	2006	0,098	0,074	0,064	0,030	0,030	0,030	0,003	0,003	0,003	0,568	0,364	0,288
Buses	Gasoline Urban Buses	Conventional	0	9999	0,140	0,110	0,070	0,006	0,006	0,006	0,002	0,002	0,002	7,000	5,500	3,500
Buses	Diesel Urban Buses <15t	Conventional	0	1993	0,175	0,080	0,070	0,030	0,030	0,030	0,003	0,003	0,003	2,628	1,738	1,490
Buses	Diesel Urban Buses <15t	Euro I	1994	1996	0,175	0,080	0,070	0,030	0,030	0,030	0,003	0,003	0,003	0,507	0,364	0,312
Buses	Diesel Urban Buses <15t	Euro II	1997	2001	0,114	0,052	0,046	0,030	0,030	0,030	0,003	0,003	0,003	0,350	0,245	0,209
Buses	Diesel Urban Buses <15t	Euro III	2002	2006	0,103	0,047	0,041	0,030	0,030	0,030	0,003	0,003	0,003	0,318	0,220	0,199
Buses	Diesel Urban Buses <15t	Euro IV	2007	2009	0,005	0,002	0,002	0,030	0,030	0,030	0,003	0,003	0,003	0,043	0,034	0,032
Buses	Diesel Urban Buses <15t	Euro V	2010	2013	0,005	0,002	0,002	0,030	0,030	0,030	0,003	0,003	0,003	0,044	0,034	0,033
Buses	Diesel Urban Buses 15 - 18t	Conventional	0	1993	0,175	0,080	0,070	0,030	0,030	0,030	0,003	0,003	0,003	1,602	0,977	0,762
Buses	Diesel Urban Buses 15 - 18t	Euro I	1994	1996	0,175	0,080	0,070	0,030	0,030	0,030	0,003	0,003	0,003	0,659	0,431	0,351
Buses	Diesel Urban Buses 15 - 18t	Euro II	1997	2001	0,114	0,052	0,046	0,030	0,030	0,030	0,003	0,003	0,003	0,451	0,296	0,248
Buses	Diesel Urban Buses 15 - 18t	Euro III	2002	2006	0,103	0,047	0,041	0,030	0,030	0,030	0,003	0,003	0,003	0,416	0,269	0,232
Buses	Diesel Urban Buses 15 - 18t	Euro IV	2007	2009	0,005	0,002	0,002	0,030	0,030	0,030	0,003	0,003	0,003	0,061	0,045	0,040
Buses	Diesel Urban Buses 15 - 18t	Euro V	2010	2013	0,005	0,002	0,002	0,030	0,030	0,030	0,003	0,003	0,003	0,061	0,046	0,040
Buses	Diesel Urban Buses >18t	Conventional	0	1993	0,175	0,080	0,070	0,030	0,030	0,030	0,003	0,003	0,003	1,666	1,018	0,791
Buses	Diesel Urban Buses >18t	Euro I	1994	1996	0,175	0,080	0,070	0,030	0,030	0,030	0,003	0,003	0,003	0,720	0,477	0,386
Buses	Diesel Urban Buses >18t	Euro II	1997	2001	0,114	0,052	0,046	0,030	0,030	0,030	0,003	0,003	0,003	0,491	0,332	0,263
Buses	Diesel Urban Buses >18t	Euro III	2002	2006	0,103	0,047	0,041	0,030	0,030	0,030	0,003	0,003	0,003	0,446	0,291	0,241
Buses	Diesel Urban Buses >18t	Euro IV	2007	2009	0,005	0,002	0,002	0,030	0,030	0,030	0,003	0,003	0,003	0,074	0,055	0,047
Buses	Diesel Urban Buses >18t	Euro V	2010	2013	0,005	0,002	0,002	0,030	0,030	0,030	0,003	0,003	0,003	0,075	0,056	0,048
Buses	Gasoline Coaches	Conventional	0	9999	0,140	0,110	0,070	0,006	0,006	0,006	0,002	0,002	0,002	7,000	5,500	3,500
Buses	Diesel Coaches <15t	Conventional	0	1993	0,175	0,080	0,070	0,030	0,030	0,030	0,003	0,003	0,003	0,907	0,533	0,393
Buses	Diesel Coaches <15t	Euro I	1994	1996	0,175	0,080	0,070	0,030	0,030	0,030	0,003	0,003	0,003	0,830	0,516	0,397
Buses	Diesel Coaches <15t	Euro II	1997	2001	0,114	0,052	0,046	0,030	0,030	0,030	0,003	0,003	0,003	0,586	0,359	0,272
Buses	Diesel Coaches <15t	Euro III	2002	2006	0,103	0,047	0,041	0,030	0,030	0,030	0,003	0,003	0,003	0,577	0,351	0,271
Buses	Diesel Coaches <15t	Euro IV	2007	2009	0,005	0,002	0,002	0,030	0,030	0,030	0,003	0,003	0,003	0,072	0,048	0,039
Buses	Diesel Coaches <15t	Euro V	2010	2013	0,005	0,002	0,002	0,030	0,030	0,030	0,003	0,003	0,003	0,074	0,049	0,039
Buses	Diesel Coaches 15 - 18t	Conventional	0	1993	0,175	0,080	0,070	0,030	0,030	0,030	0,003	0,003	0,003	0,907	0,533	0,393
Buses	Diesel Coaches 15 - 18t	Euro I	1994	1996	0,175	0,080	0,070	0,030	0,030	0,030	0,003	0,003	0,003	0,830	0,516	0,397
Buses	Diesel Coaches 15 - 18t	Euro II	1997	2001	0,114	0,052	0,046	0,030	0,030	0,030	0,003	0,003	0,003	0,586	0,359	0,272

Buses	Diesel Coaches 15 - 18t	Euro III	2002	2006	0,103	0,047	0,041	0,030	0,030	0,030	0,003	0,003	0,003	0,577	0,351	0,271
Buses	Diesel Coaches 15 - 18t	Euro IV	2007	2009	0,005	0,002	0,002	0,030	0,030	0,030	0,003	0,003	0,003	0,072	0,048	0,039
Buses	Diesel Coaches 15 - 18t	Euro V	2010	2013	0,005	0,002	0,002	0,030	0,030	0,030	0,003	0,003	0,003	0,074	0,049	0,039
Buses	Diesel Coaches >18t	Conventional	0	1993	0,175	0,080	0,070	0,030	0,030	0,030	0,003	0,003	0,003	1,013	0,623	0,482
Buses	Diesel Coaches >18t	Euro I	1994	1996	0,175	0,080	0,070	0,030	0,030	0,030	0,003	0,003	0,003	0,915	0,581	0,457
Buses	Diesel Coaches >18t	Euro II	1997	2001	0,114	0,052	0,046	0,030	0,030	0,030	0,003	0,003	0,003	0,630	0,392	0,305
Buses	Diesel Coaches >18t	Euro III	2002	2006	0,103	0,047	0,041	0,030	0,030	0,030	0,003	0,003	0,003	0,608	0,371	0,286
Buses	Diesel Coaches >18t	Euro IV	2007	2009	0,005	0,002	0,002	0,030	0,030	0,030	0,003	0,003	0,003	0,076	0,050	0,042
Buses	Diesel Coaches >18t	Euro V	2010	2013	0,005	0,002	0,002	0,030	0,030	0,030	0,003	0,003	0,003	0,078	0,051	0,042
Mopeds	2-stroke <50 cm ³	Conventional	0	1999	0,219	0,219	0,000	0,001	0,001	0,001	0,001	0,001	0,001	8,400	8,400	0,000
Mopeds	2-stroke <50 cm ³	Euro I	2000	2003	0,044	0,044	0,000	0,001	0,001	0,001	0,001	0,001	0,001	3,400	3,400	0,000
Mopeds	2-stroke <50 cm ³	Euro II	2004	2014	0,024	0,024	0,000	0,001	0,001	0,001	0,001	0,001	0,001	2,600	2,600	0,000
Mopeds	4-stroke <50 cm ³	Euro II	2004	2014	0,024	0,024	0,000	0,001	0,001	0,001	0,001	0,001	0,001	0,790	0,790	0,000
Motorcycles	2-stroke >50 cm ³	Conventional	0	1999	0,150	0,150	0,150	0,002	0,002	0,002	0,002	0,002	0,002	8,393	7,078	9,800
Motorcycles	2-stroke >50 cm ³	Euro I	2000	2003	0,099	0,107	0,098	0,002	0,002	0,002	0,002	0,002	0,002	8,393	7,078	9,800
Motorcycles	2-stroke >50 cm ³	Euro II	2004	2006	0,030	0,032	0,030	0,002	0,002	0,002	0,002	0,002	0,002	2,593	2,569	4,155
Motorcycles	2-stroke >50 cm ³	Euro III	2007	9999	0,012	0,014	0,012	0,002	0,002	0,002	0,002	0,002	0,002	1,385	1,380	2,244
Motorcycles	4-stroke <250 cm ³	Conventional	0	1999	0,200	0,200	0,200	0,002	0,002	0,002	0,002	0,002	0,002	0,128	0,104	0,138
Motorcycles	4-stroke <250 cm ³	Euro I	2000	2003	0,142	0,144	0,132	0,002	0,002	0,002	0,002	0,002	0,002	1,242	0,866	0,976
Motorcycles	4-stroke <250 cm ³	Euro II	2004	2006	0,136	0,092	0,092	0,002	0,002	0,002	0,002	0,002	0,002	1,042	0,843	0,965
Motorcycles	4-stroke <250 cm ³	Euro III	2007	9999	0,082	0,032	0,028	0,002	0,002	0,002	0,002	0,002	0,002	0,456	0,441	0,511
Motorcycles	4-stroke 250 - 750 cm ³	Conventional	0	1999	0,200	0,200	0,200	0,002	0,002	0,002	0,002	0,002	0,002	0,545	0,487	0,361
Motorcycles	4-stroke 250 - 750 cm ³	Euro I	2000	2003	0,148	0,174	0,156	0,002	0,002	0,002	0,002	0,002	0,002	2,390	1,522	1,079
Motorcycles	4-stroke 250 - 750 cm ³	Euro II	2004	2006	0,156	0,120	0,122	0,002	0,002	0,002	0,002	0,002	0,002	1,326	0,925	0,828
Motorcycles	4-stroke 250 - 750 cm ³	Euro III	2007	9999	0,094	0,042	0,036	0,002	0,002	0,002	0,002	0,002	0,002	0,598	0,499	0,615
Motorcycles	4-stroke >750 cm ³	Conventional	0	1999	0,200	0,200	0,200	0,002	0,002	0,002	0,002	0,002	0,002	0,392	0,337	0,556
Motorcycles	4-stroke >750 cm ³	Euro I	2000	2003	0,092	0,092	0,154	0,002	0,002	0,002	0,002	0,002	0,002	2,495	1,643	1,554
Motorcycles	4-stroke >750 cm ³	Euro II	2004	2006	0,084	0,062	0,102	0,002	0,002	0,002	0,002	0,002	0,002	1,088	0,674	0,656
Motorcycles	4-stroke >750 cm ³	Euro III	2007	9999	0,050	0,022	0,030	0,002	0,002	0,002	0,002	0,002	0,002	0,384	0,309	0,416

Annex 3B-5 Reduction factors

Sector	Subsector	Tech 2	FYear	LYear	FCuR	FCrR	FChR	COuR	COrR	COhR	PMuR	PMrR	PMhR	NOxuR	NOxrR	NOxhR	VOCuR	VOCrR	VOChR
Passenger Cars	Gasoline 0,8 - 1,4 l	PRE ECE	0	1969	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
Passenger Cars	Gasoline 0,8 - 1,4 l	ECE 15/00-01	1970	1978	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
Passenger Cars	Gasoline 0,8 - 1,4 l	ECE 15/02	1979	1980	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
Passenger Cars	Gasoline 0,8 - 1,4 l	ECE 15/03	1981	1985	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
Passenger Cars	Gasoline 0,8 - 1,4 l	ECE 15/04	1986	1990	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
Passenger Cars	Gasoline 0,8 - 1,4 l	Euro I	1991	1996	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
Passenger Cars	Gasoline 0,8 - 1,4 l	Euro II	1997	2000	1,93	5,83	4,43	62,65	58,10	57,55	0,00	0,00	0,00	43,59	45,20	60,45	60,19	61,27	62,09
Passenger Cars	Gasoline 0,8 - 1,4 l	Euro III	2001	2005	-2,72	-0,72	1,59	70,59	49,62	34,95	60,25	54,57	37,37	72,16	78,49	88,69	91,74	87,53	77,02
Passenger Cars	Gasoline 0,8 - 1,4 l	Euro IV	2006	2010	-5,57	-5,34	-4,84	88,95	79,10	70,06	60,25	54,57	37,37	80,12	89,24	95,86	93,34	88,71	84,51
Passenger Cars	Gasoline 0,8 - 1,4 l	Euro V	2011	2014	-5,57	-5,34	-4,84	88,95	79,10	70,06	60,25	54,57	37,37	85,09	91,93	96,90	93,34	88,71	84,51
Passenger Cars	Gasoline 1,4 - 2,0 l	PRE ECE	0	1969	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
Passenger Cars	Gasoline 1,4 - 2,0 l	ECE 15/00-01	1970	1978	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
Passenger Cars	Gasoline 1,4 - 2,0 l	ECE 15/02	1979	1980	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
Passenger Cars	Gasoline 1,4 - 2,0 l	ECE 15/03	1981	1985	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
Passenger Cars	Gasoline 1,4 - 2,0 l	ECE 15/04	1986	1990	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
Passenger Cars	Gasoline 1,4 - 2,0 l	Euro I	1991	1996	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
Passenger Cars	Gasoline 1,4 - 2,0 l	Euro II	1997	2000	2,08	1,72	5,30	62,65	58,10	57,55	0,00	0,00	0,00	43,59	45,20	60,45	60,19	61,27	62,09
Passenger Cars	Gasoline 1,4 - 2,0 l	Euro III	2001	2005	-1,28	-2,83	-3,05	70,59	49,62	34,95	60,25	54,57	37,37	72,16	78,49	88,69	91,74	87,53	77,02
Passenger Cars	Gasoline 1,4 - 2,0 l	Euro IV	2006	2010	-5,15	-8,80	-7,33	88,95	79,10	70,06	60,25	54,57	37,37	80,12	89,24	95,86	93,34	88,71	84,51
Passenger Cars	Gasoline 1,4 - 2,0 l	Euro V	2011	2014	-5,15	-8,80	-7,33	88,95	79,10	70,06	60,25	54,57	37,37	85,09	91,93	96,90	93,34	88,71	84,51
Passenger Cars	Gasoline >2,0 l	PRE ECE	0	1969	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
Passenger Cars	Gasoline >2,0 l	ECE 15/00-01	1970	1978	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
Passenger Cars	Gasoline >2,0 l	ECE 15/02	1979	1980	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
Passenger Cars	Gasoline >2,0 l	ECE 15/03	1981	1985	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
Passenger Cars	Gasoline >2,0 l	ECE 15/04	1986	1990	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
Passenger Cars	Gasoline >2,0 l	Euro I	1991	1996	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
Passenger Cars	Gasoline >2,0 l	Euro II	1997	2000	-2,98	-5,37	-2,63	62,65	58,10	57,55	0,00	0,00	0,00	43,59	45,20	60,45	60,19	61,27	62,09
Passenger Cars	Gasoline >2,0 l	Euro III	2001	2005	5,12	7,23	12,60	70,59	49,62	34,95	60,25	54,57	37,37	72,16	78,49	88,69	91,74	87,53	77,02
Passenger Cars	Gasoline >2,0 l	Euro IV	2006	2010	-15,39	-9,65	-1,29	88,95	79,10	70,06	60,25	54,57	37,37	80,12	89,24	95,86	93,34	88,71	84,51
Passenger Cars	Gasoline >2,0 l	Euro V	2011	2014	-15,39	-9,65	-1,29	88,95	79,10	70,06	60,25	54,57	37,37	85,09	91,93	96,90	93,34	88,71	84,51
Passenger Cars	Diesel 1,4 - 2,0 l	Conventional	0	1990	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
Passenger Cars	Diesel 1,4 - 2,0 l	Euro I	1991	1996	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
Passenger Cars	Diesel 1,4 - 2,0 l	Euro II	1997	2000	-5,45	-3,06	-0,81	18,08	48,77	83,05	17,92	36,92	53,22	-7,94	1,18	-0,20	34,81	33,43	41,61

Passenger Cars	Diesel 1,4 - 2,0 l	Euro III	2001	2005	-2,27	-1,45	5,79	76,38	81,12	94,30	48,53	51,90	58,32	-18,71	-18,46	-12,98	65,94	63,35	66,25
Passenger Cars	Diesel 1,4 - 2,0 l	Euro IV	2006	2010	-2,27	-1,45	5,79	80,09	84,22	89,72	49,02	60,57	75,83	10,60	24,53	13,19	79,38	79,24	77,57
Passenger Cars	Diesel 1,4 - 2,0 l	Euro V	2011	2014	-2,20	-1,32	5,99	80,09	84,22	89,72	89,80	92,11	95,17	-9,96	7,17	-6,78	79,38	79,24	77,57
Passenger Cars	Diesel >2,0 l	Conventional	0	1990	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
Passenger Cars	Diesel >2,0 l	Euro I	1991	1996	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
Passenger Cars	Diesel >2,0 l	Euro II	1997	2000	0,00	0,00	0,00	18,08	48,77	83,05	17,92	36,92	53,22	-7,94	1,18	-0,20	-22,14	-25,38	-11,51
Passenger Cars	Diesel >2,0 l	Euro III	2001	2005	0,00	0,00	0,00	76,38	81,12	94,30	48,53	51,90	58,32	-18,71	-18,46	-12,98	52,23	62,67	63,93
Passenger Cars	Diesel >2,0 l	Euro IV	2006	2010	0,00	0,00	0,00	80,09	84,22	89,72	49,02	60,57	75,83	10,60	24,53	13,19	86,39	86,10	83,20
Passenger Cars	Diesel >2,0 l	Euro V	2011	2014	-0,07	-0,06	-0,05	80,09	84,22	89,72	89,80	92,11	95,17	-9,96	7,17	-6,78	86,39	86,10	83,20
Passenger Cars	LPG cars	Conventional	0	1990	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
Passenger Cars	LPG cars	Euro I	1991	1996	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
Passenger Cars	LPG cars	Euro II	1997	2000	0,00	0,00	0,00	32,00	32,00	32,00	0,00	0,00	0,00	64,00	64,00	64,00	79,00	79,00	79,00
Passenger Cars	LPG cars	Euro III	2001	2005	0,00	0,00	0,00	44,00	44,00	44,00	0,00	0,00	0,00	76,00	76,00	76,00	85,00	85,00	85,00
Passenger Cars	LPG cars	Euro IV	2006	2010	0,00	0,00	0,00	66,00	66,00	66,00	0,00	0,00	0,00	87,00	87,00	87,00	97,00	97,00	97,00
Passenger Cars	Electric cars	Conventional	0	9999	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
Passenger Cars	Gasoline <0,8 l	PRE ECE	0	1969	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
Passenger Cars	Gasoline <0,8 l	ECE 15/00-01	1970	1978	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
Passenger Cars	Gasoline <0,8 l	ECE 15/02	1979	1980	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
Passenger Cars	Gasoline <0,8 l	ECE 15/03	1981	1985	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
Passenger Cars	Gasoline <0,8 l	ECE 15/04	1986	1990	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
Passenger Cars	Gasoline <0,8 l	Euro I	1991	1996	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
Passenger Cars	Gasoline <0,8 l	Euro II	1997	2000	1,93	5,83	4,43	62,65	58,10	57,55	0,00	0,00	0,00	43,59	45,20	60,45	60,19	61,27	62,09
Passenger Cars	Gasoline <0,8 l	Euro III	2001	2005	-2,72	-0,72	1,59	70,59	49,62	34,95	60,25	54,57	37,37	72,16	78,49	88,69	91,74	87,53	77,02
Passenger Cars	Gasoline <0,8 l	Euro IV	2006	2010	-5,57	-5,34	-4,84	88,95	79,10	70,06	60,25	54,57	37,37	80,12	89,24	95,86	93,34	88,71	84,51
Passenger Cars	Gasoline <0,8 l	Euro V	2011	2014	-5,57	-5,34	-4,84	88,95	79,10	70,06	60,25	54,57	37,37	85,09	91,93	96,90	93,34	88,71	84,51
Passenger Cars	Diesel <1,4 l	Conventional	0	1990	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
Passenger Cars	Diesel <1,4 l	Euro I	1991	1996	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
Passenger Cars	Diesel <1,4 l	Euro II	1997	2000	-5,45	-3,06	-0,81	18,08	48,77	83,05	17,92	36,92	53,22	-7,94	1,18	-0,20	34,81	33,43	41,61
Passenger Cars	Diesel <1,4 l	Euro III	2001	2005	-2,27	-1,45	5,79	76,38	81,12	94,30	48,53	51,90	58,32	-18,71	-18,46	-12,98	65,94	63,35	66,25
Passenger Cars	Diesel <1,4 l	Euro IV	2006	2010	-2,27	-1,45	5,79	80,09	84,22	89,72	49,02	60,57	75,83	10,60	24,53	13,19	79,38	79,24	77,57
Passenger Cars	Diesel <1,4 l	Euro V	2011	2014	-2,20	-1,32	5,99	80,09	84,22	89,72	89,80	92,11	95,17	-9,96	7,17	-6,78	79,38	79,24	77,57
Light Duty Vehicles	Gasoline <3,5t	Conventional	0	1994	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
Light Duty Vehicles	Gasoline <3,5t	Euro I	1995	1998	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
Light Duty Vehicles	Gasoline <3,5t	Euro II	1999	2001	0,00	0,00	0,00	39,00	39,00	39,00	0,00	0,00	0,00	66,00	66,00	66,00	76,00	76,00	76,00
Light Duty Vehicles	Gasoline <3,5t	Euro III	2002	2006	0,00	0,00	0,00	48,00	48,00	48,00	60,25	54,57	37,37	79,00	79,00	79,00	86,00	86,00	86,00
Light Duty Vehicles	Gasoline <3,5t	Euro IV	2007	2011	0,00	0,00	0,00	72,00	72,00	72,00	60,25	54,57	37,37	90,00	90,00	90,00	94,00	94,00	94,00
Light Duty Vehicles	Gasoline <3,5t	Euro V	2012	2015	0,00	0,00	0,00	72,00	72,00	72,00	60,25	54,57	37,37	92,50	92,50	92,50	94,00	94,00	94,00

Light Duty Vehicles	Diesel <3,5t	Conventional	0	1994	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
Light Duty Vehicles	Diesel <3,5t	Euro I	1995	1998	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
Light Duty Vehicles	Diesel <3,5t	Euro II	1999	2001	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
Light Duty Vehicles	Diesel <3,5t	Euro III	2002	2006	0,00	0,00	0,00	18,00	18,00	18,00	33,00	33,00	33,00	16,00	16,00	16,00	38,00	38,00	38,00
Light Duty Vehicles	Diesel <3,5t	Euro IV	2007	2011	0,00	0,00	0,00	35,00	35,00	35,00	65,00	65,00	65,00	32,00	32,00	32,00	77,00	77,00	77,00
Light Duty Vehicles	Diesel <3,5t	Euro V	2012	2015	0,00	0,00	0,00	35,00	35,00	35,00	98,25	98,25	98,25	51,00	51,00	51,00	77,00	77,00	77,00
Light Duty Vehicles	LPG <3,5t	Conventional	0	1994	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
Light Duty Vehicles	LPG <3,5t	Euro II	1999	2001	0,00	0,00	0,00	32,00	32,00	32,00	0,00	0,00	0,00	64,00	64,00	64,00	79,00	79,00	79,00
Light Duty Vehicles	LPG <3,5t	Euro III	2002	2006	0,00	0,00	0,00	44,00	44,00	44,00	0,00	0,00	0,00	76,00	76,00	76,00	85,00	85,00	85,00
Light Duty Vehicles	LPG <3,5t	Euro IV	2007	2011	0,00	0,00	0,00	66,00	66,00	66,00	0,00	0,00	0,00	87,00	87,00	87,00	97,00	97,00	97,00
Light Duty Vehicles	Electric <3,5t	Conventional	0	9999	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
Heavy Duty Vehicles	Gasoline >3,5t	Conventional	0	9999	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
Heavy Duty Vehicles	Diesel RT 3,5 - 7,5t	Conventional	0	1993	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
Heavy Duty Vehicles	Diesel RT 3,5 - 7,5t	Euro I	1994	1996	19,97	17,39	7,76	67,55	66,82	59,55	60,69	60,33	58,47	30,21	28,42	25,92	80,53	78,89	77,38
Heavy Duty Vehicles	Diesel RT 3,5 - 7,5t	Euro II	1997	2001	24,01	20,18	10,60	74,08	69,13	65,86	81,61	77,85	71,87	23,47	24,02	23,73	86,86	85,98	85,16
Heavy Duty Vehicles	Diesel RT 3,5 - 7,5t	Euro III	2002	2006	18,90	16,31	6,25	67,98	68,13	66,52	79,28	80,07	80,84	40,66	43,96	44,20	87,56	87,06	86,71
Heavy Duty Vehicles	Diesel RT 3,5 - 7,5t	Euro IV	2007	2009	21,15	16,29	5,64	83,37	82,11	80,91	95,44	94,65	93,58	59,45	59,91	59,77	98,28	97,79	97,11
Heavy Duty Vehicles	Diesel RT 3,5 - 7,5t	Euro V	2010	2013	20,29	15,72	5,30	83,30	82,11	80,82	95,45	94,62	93,60	75,97	76,32	76,28	98,27	97,79	97,13
Heavy Duty Vehicles	Diesel RT 3,5 - 7,5t	Euro VI	2014	9999	21,40	17,89	8,47	82,75	81,73	80,76	99,54	99,46	99,36	94,47	97,63	98,67	99,16	98,98	98,77
Heavy Duty Vehicles	Diesel RT 7,5 - 12t	Conventional	0	1993	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
Heavy Duty Vehicles	Diesel RT 7,5 - 12t	Euro I	1994	1996	14,94	11,49	8,93	53,96	51,87	49,75	39,18	39,03	36,52	40,35	40,49	40,47	59,34	56,17	53,69
Heavy Duty Vehicles	Diesel RT 7,5 - 12t	Euro II	1997	2001	18,90	14,27	11,02	63,18	57,16	53,02	71,53	66,12	55,10	35,02	36,53	37,57	72,55	70,85	69,48
Heavy Duty Vehicles	Diesel RT 7,5 - 12t	Euro III	2002	2006	14,01	10,02	7,38	54,03	54,57	51,94	68,40	69,00	69,32	49,58	51,13	53,53	73,67	73,28	73,21
Heavy Duty Vehicles	Diesel RT 7,5 - 12t	Euro IV	2007	2009	17,35	11,08	7,67	76,57	75,39	75,78	93,10	92,09	90,79	65,60	65,29	66,82	96,38	95,67	95,13
Heavy Duty Vehicles	Diesel RT 7,5 - 12t	Euro V	2010	2013	16,17	10,25	7,18	76,27	75,18	75,47	93,04	92,07	90,83	79,22	79,51	80,43	96,34	95,66	95,14
Heavy Duty Vehicles	Diesel RT 7,5 - 12t	Euro VI	2014	9999	17,31	12,39	10,14	75,26	74,61	75,54	99,29	99,20	99,09	95,16	97,87	98,72	98,22	97,93	97,71
Heavy Duty Vehicles	Diesel RT 12 - 14 t	Conventional	0	1993	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
Heavy Duty Vehicles	Diesel RT 12 - 14 t	Euro I	1994	1996	14,28	11,64	9,97	52,86	51,03	48,87	37,82	37,42	37,06	39,71	39,91	40,14	57,63	56,88	54,97
Heavy Duty Vehicles	Diesel RT 12 - 14 t	Euro II	1997	2001	17,78	14,26	12,32	61,29	56,29	52,49	70,72	65,68	55,94	34,11	35,54	36,90	72,19	71,22	70,63
Heavy Duty Vehicles	Diesel RT 12 - 14 t	Euro III	2002	2006	13,28	10,55	9,17	53,81	53,44	50,67	68,93	69,17	69,39	46,23	49,71	52,23	74,33	74,00	73,71
Heavy Duty Vehicles	Diesel RT 12 - 14 t	Euro IV	2007	2009	17,83	12,48	10,01	76,48	76,13	75,80	93,03	92,25	91,58	63,65	64,32	66,18	96,64	96,09	95,32
Heavy Duty Vehicles	Diesel RT 12 - 14 t	Euro V	2010	2013	16,83	11,77	9,53	76,18	75,91	75,63	92,99	92,24	91,61	78,37	78,82	79,97	96,61	96,08	95,34
Heavy Duty Vehicles	Diesel RT 14 - 20t	Conventional	0	1993	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
Heavy Duty Vehicles	Diesel RT 14 - 20t	Euro I	1994	1996	18,66	16,33	14,88	54,10	52,05	49,69	38,29	38,82	39,43	40,46	40,76	40,96	59,89	58,50	57,65
Heavy Duty Vehicles	Diesel RT 14 - 20t	Euro II	1997	2001	21,92	18,70	17,00	63,94	59,22	54,89	73,27	69,36	59,06	33,79	35,29	36,36	72,92	72,50	72,28
Heavy Duty Vehicles	Diesel RT 14 - 20t	Euro III	2002	2006	17,70	15,52	14,62	54,42	54,24	50,66	68,29	69,01	69,76	45,61	48,61	51,42	74,94	75,03	74,44
Heavy Duty Vehicles	Diesel RT 14 - 20t	Euro IV	2007	2009	23,15	18,31	16,14	76,40	76,04	76,43	93,48	92,87	92,25	63,86	64,05	65,23	96,97	96,70	96,31

Heavy Duty Vehicles	Diesel RT 14 - 20t	Euro V	2010	2013	21,50	17,49	15,95	75,25	75,14	75,86	93,36	92,81	92,21	78,21	78,55	79,35	96,86	96,61	96,25
Heavy Duty Vehicles	Diesel RT 14 - 20t	Euro VI	2014	9999	23,32	19,40	17,88	75,24	75,14	75,87	99,34	99,28	99,22	93,85	97,23	98,52	98,43	98,28	98,07
Heavy Duty Vehicles	Diesel RT 20 - 26t	Conventional	0	1993	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
Heavy Duty Vehicles	Diesel RT 20 - 26t	Euro I	1994	1996	14,59	12,88	11,93	19,16	17,09	16,07	20,49	25,29	27,75	29,52	29,51	29,03	11,17	8,06	6,45
Heavy Duty Vehicles	Diesel RT 20 - 26t	Euro II	1997	2001	17,36	14,83	13,74	36,69	31,82	18,28	64,44	61,13	50,94	22,74	23,45	23,78	40,30	39,31	39,01
Heavy Duty Vehicles	Diesel RT 20 - 26t	Euro III	2002	2006	13,66	12,19	11,99	20,84	21,10	18,05	60,89	63,32	64,58	37,56	38,91	39,16	44,75	44,58	44,64
Heavy Duty Vehicles	Diesel RT 20 - 26t	Euro IV	2007	2009	18,46	14,72	13,40	60,80	61,35	63,31	91,59	91,35	91,18	58,00	57,18	56,47	92,84	92,20	91,51
Heavy Duty Vehicles	Diesel RT 20 - 26t	Euro V	2010	2013	17,52	13,99	12,79	60,33	61,00	62,95	91,54	91,32	91,17	75,01	74,57	74,18	92,77	92,15	91,47
Heavy Duty Vehicles	Diesel RT 20 - 26t	Euro VI	2014	9999	19,02	16,10	15,22	59,13	60,04	62,26	99,14	99,13	99,11	94,21	97,09	98,20	96,40	96,02	95,71
Heavy Duty Vehicles	Diesel RT 26 - 28t	Conventional	0	1993	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
Heavy Duty Vehicles	Diesel RT 26 - 28t	Euro I	1994	1996	14,23	12,62	11,74	20,03	17,07	15,21	22,33	25,06	27,38	29,11	29,59	29,23	12,60	9,85	8,38
Heavy Duty Vehicles	Diesel RT 26 - 28t	Euro II	1997	2001	17,09	14,65	13,53	37,76	32,22	19,50	63,87	60,48	50,30	23,25	24,39	24,79	40,81	39,59	39,01
Heavy Duty Vehicles	Diesel RT 26 - 28t	Euro III	2002	2006	13,37	11,82	11,44	21,52	20,38	18,17	60,68	62,36	64,94	39,91	41,33	40,87	44,52	43,89	43,54
Heavy Duty Vehicles	Diesel RT 26 - 28t	Euro IV	2007	2009	17,01	13,58	12,64	61,39	62,14	64,62	91,49	91,19	91,37	59,14	58,72	57,71	92,39	91,72	91,36
Heavy Duty Vehicles	Diesel RT 26 - 28t	Euro V	2010	2013	16,01	12,79	12,02	60,92	61,74	64,28	91,43	91,15	91,35	75,70	75,49	74,93	92,31	91,66	91,32
Heavy Duty Vehicles	Diesel RT 28 - 32t	Conventional	0	1993	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
Heavy Duty Vehicles	Diesel RT 28 - 32t	Euro I	1994	1996	12,20	11,05	10,36	18,82	13,37	12,31	23,04	24,44	25,14	27,98	28,75	28,76	10,90	7,44	5,56
Heavy Duty Vehicles	Diesel RT 28 - 32t	Euro II	1997	2001	15,27	13,95	9,64	34,08	26,77	23,67	62,75	58,53	48,13	22,62	24,28	24,79	40,10	38,55	37,82
Heavy Duty Vehicles	Diesel RT 28 - 32t	Euro III	2002	2006	11,41	10,04	9,73	20,60	19,39	17,71	62,16	63,07	64,05	38,80	41,24	41,46	43,86	43,33	43,19
Heavy Duty Vehicles	Diesel RT 28 - 32t	Euro IV	2007	2009	14,35	11,13	10,55	60,90	61,17	65,11	91,34	90,97	90,96	58,81	57,29	58,82	91,98	90,92	90,30
Heavy Duty Vehicles	Diesel RT 28 - 32t	Euro V	2010	2013	13,34	10,31	9,85	60,41	60,74	64,74	91,27	90,91	90,93	75,51	74,62	75,57	91,91	90,86	90,25
Heavy Duty Vehicles	Diesel RT 28 - 32t	Euro VI	2014	9999	14,95	12,46	12,29	59,32	59,85	64,03	99,11	99,08	99,09	95,61	97,57	98,26	96,11	95,65	95,39
Heavy Duty Vehicles	Diesel RT >32t	Euro I	1994	1996	12,62	11,53	10,93	16,13	13,50	12,48	20,12	23,79	25,25	28,73	28,95	28,74	8,76	5,59	4,01
Heavy Duty Vehicles	Diesel RT >32t	Euro III	2002	2006	11,93	10,92	10,90	18,97	18,29	15,90	61,41	63,24	64,69	38,06	39,23	39,11	44,27	43,43	42,99
Heavy Duty Vehicles	Diesel RT >32t	Euro IV	2007	2009	15,80	12,78	12,01	61,10	61,71	64,74	91,36	91,17	91,27	57,90	57,34	56,23	92,18	91,31	90,51
Heavy Duty Vehicles	Diesel RT >32t	Euro V	2010	2013	14,90	12,07	11,43	60,60	61,33	64,46	91,29	91,12	91,26	74,92	74,66	74,06	92,09	91,27	90,50
Heavy Duty Vehicles	Diesel TT/AT 28 - 34t	Conventional	0	1993	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
Heavy Duty Vehicles	Diesel TT/AT 28 - 34t	Euro I	1994	1996	10,56	9,81	9,65	15,13	12,34	11,18	22,12	23,23	22,95	28,53	29,18	29,25	7,82	5,31	4,45
Heavy Duty Vehicles	Diesel TT/AT 28 - 34t	Euro II	1997	2001	15,65	13,67	10,63	31,82	27,73	16,83	60,81	57,94	48,69	24,49	25,71	26,04	38,86	37,80	37,36
Heavy Duty Vehicles	Diesel TT/AT 28 - 34t	Euro III	2002	2006	11,19	10,15	10,37	19,25	17,89	15,99	62,26	63,46	64,44	39,05	41,17	41,88	43,54	43,44	43,48
Heavy Duty Vehicles	Diesel TT/AT 28 - 34t	Euro IV	2007	2009	14,73	11,88	11,73	61,31	62,64	65,74	91,27	91,29	91,77	58,43	58,62	58,68	91,95	91,29	90,55
Heavy Duty Vehicles	Diesel TT/AT 28 - 34t	Euro V	2010	2013	13,84	11,15	11,06	60,85	62,25	65,37	91,19	91,23	91,73	75,22	75,38	75,48	91,88	91,23	90,51
Heavy Duty Vehicles	Diesel TT/AT 28 - 34t	Euro VI	2014	9999	15,57	13,36	13,49	59,64	61,13	65,11	99,10	99,11	99,16	96,01	97,66	98,20	96,03	95,67	95,44
Heavy Duty Vehicles	Diesel TT/AT 34 - 40t	Conventional	0	1993	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
Heavy Duty Vehicles	Diesel TT/AT 34 - 40t	Euro I	1994	1996	12,21	11,31	10,88	14,79	12,18	10,98	19,78	22,63	23,62	29,18	29,40	29,10	8,18	5,48	4,19
Heavy Duty Vehicles	Diesel TT/AT 34 - 40t	Euro II	1997	2001	14,41	12,85	12,47	31,60	27,45	15,10	60,77	57,68	47,54	23,95	24,61	24,32	38,67	38,17	38,36
Heavy Duty Vehicles	Diesel TT/AT 34 - 40t	Euro III	2002	2006	11,35	10,70	11,09	18,41	17,61	15,12	61,53	63,00	64,68	38,79	39,57	39,01	43,62	43,84	44,36

Heavy Duty Vehicles	Diesel TT/AT 34 - 40t	Euro IV	2007	2009	15,08	12,56	12,32	61,51	62,53	66,34	91,40	91,31	91,63	58,39	57,50	56,59	91,92	91,32	90,66
Heavy Duty Vehicles	Diesel TT/AT 34 - 40t	Euro V	2010	2013	14,16	11,82	11,68	61,16	62,20	65,81	91,32	91,25	91,60	75,20	74,74	74,26	91,84	91,26	90,62
Heavy Duty Vehicles	Diesel TT/AT 34 - 40t	Euro VI	2014	9999	15,96	14,09	14,14	60,02	61,34	65,21	99,12	99,11	99,15	96,03	97,61	98,17	96,02	95,69	95,45
Heavy Duty Vehicles	Diesel TT/AT 40 - 50t	Conventional	0	1993	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
Heavy Duty Vehicles	Diesel TT/AT 40 - 50t	Euro I	1994	1996	12,06	11,24	10,92	12,93	11,05	8,95	19,65	22,36	21,85	29,86	29,82	29,51	6,26	4,20	3,76
Heavy Duty Vehicles	Diesel TT/AT 40 - 50t	Euro II	1997	2001	14,86	12,96	10,45	28,66	23,92	20,16	58,75	56,55	46,55	25,17	25,64	25,26	38,00	37,21	37,26
Heavy Duty Vehicles	Diesel TT/AT 40 - 50t	Euro III	2002	2006	11,13	10,59	10,91	17,51	16,69	14,19	61,42	63,23	65,66	39,74	40,37	39,79	43,37	43,37	43,77
Heavy Duty Vehicles	Diesel TT/AT 40 - 50t	Euro IV	2007	2009	14,11	11,92	11,89	61,73	62,77	66,78	91,26	91,37	91,76	59,36	57,67	57,38	91,47	90,74	89,97
Heavy Duty Vehicles	Diesel TT/AT 40 - 50t	Euro V	2010	2013	13,18	11,19	11,28	61,26	62,40	66,49	91,17	91,30	91,72	75,82	74,82	74,66	91,38	90,68	89,95
Heavy Duty Vehicles	Diesel TT/AT 40 - 50t	Euro VI	2014	9999	15,07	13,55	13,86	60,29	61,63	65,88	99,10	99,12	99,16	96,74	97,80	98,16	95,81	95,44	95,19
Heavy Duty Vehicles	Diesel TT/AT 50 - 60t	Euro II	1997	2001	15,12	13,22	11,08	27,35	21,99	18,67	57,33	55,25	45,19	26,42	26,84	26,77	38,07	36,01	34,82
Heavy Duty Vehicles	Diesel TT/AT 50 - 60t	Euro IV	2007	2009	14,82	12,10	11,08	62,26	63,07	67,40	91,40	91,54	91,89	59,93	58,30	58,71	91,05	90,11	89,38
Heavy Duty Vehicles	Diesel TT/AT >60t	Euro III	2002	2006	12,71	11,99	9,67	17,25	15,70	13,12	62,64	64,66	63,21	40,69	41,27	41,10	43,87	42,68	41,90
Buses	Gasoline Urban Buses	Conventional	0	9999	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
Buses	Diesel Urban Buses <15t	Conventional	0	1993	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
Buses	Diesel Urban Buses <15t	Euro I	1994	1996	19,18	17,29	17,93	64,98	64,38	65,34	64,18	59,36	56,81	25,70	27,96	31,85	80,69	79,04	79,08
Buses	Diesel Urban Buses <15t	Euro II	1997	2001	22,00	19,28	19,64	68,95	69,53	71,53	82,28	78,21	74,97	19,20	22,23	26,76	86,68	85,91	86,00
Buses	Diesel Urban Buses <15t	Euro III	2002	2006	17,34	14,77	15,40	66,31	67,30	67,27	82,14	79,67	77,43	31,27	41,19	49,10	87,89	87,33	86,64
Buses	Diesel Urban Buses <15t	Euro IV	2007	2009	23,22	17,36	12,81	82,13	82,75	85,07	95,64	94,92	94,57	56,39	59,61	63,64	98,36	98,06	97,83
Buses	Diesel Urban Buses <15t	Euro V	2010	2013	21,91	16,20	11,62	81,85	82,49	84,81	95,57	94,91	94,51	73,98	75,97	78,34	98,34	98,04	97,80
Buses	Diesel Urban Buses 15 - 18t	Conventional	0	1993	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
Buses	Diesel Urban Buses 15 - 18t	Euro I	1994	1996	14,69	12,79	11,87	53,30	50,29	48,97	45,25	41,27	35,52	38,51	39,11	40,52	58,83	55,91	53,90
Buses	Diesel Urban Buses 15 - 18t	Euro II	1997	2001	17,30	14,13	12,17	59,92	59,59	57,00	72,71	66,78	60,84	33,88	35,52	36,86	71,87	69,67	67,40
Buses	Diesel Urban Buses 15 - 18t	Euro III	2002	2006	13,33	10,21	8,28	56,14	57,37	51,77	73,49	70,02	67,33	44,22	50,21	54,47	74,05	72,45	69,52
Buses	Diesel Urban Buses 15 - 18t	Euro IV	2007	2009	18,27	12,04	5,41	77,87	78,14	78,66	93,34	92,51	91,91	63,91	65,56	67,41	96,21	95,37	94,76
Buses	Diesel Urban Buses 15 - 18t	Euro V	2010	2013	17,09	11,02	4,36	77,61	77,91	78,39	93,29	92,44	91,82	78,49	79,52	80,66	96,18	95,33	94,71
Buses	Diesel Urban Buses >18t	Conventional	0	1993	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
Buses	Diesel Urban Buses >18t	Euro I	1994	1996	13,02	11,55	11,00	53,10	50,54	42,55	45,84	41,58	31,51	38,69	39,58	40,12	56,81	53,19	51,17
Buses	Diesel Urban Buses >18t	Euro II	1997	2001	15,64	12,70	7,01	58,65	60,18	57,11	71,05	66,18	60,82	35,41	37,06	37,96	70,55	67,39	66,73
Buses	Diesel Urban Buses >18t	Euro III	2002	2006	12,01	9,43	7,87	56,20	58,74	50,21	74,88	73,79	68,82	45,31	50,39	53,06	73,26	71,41	69,52
Buses	Diesel Urban Buses >18t	Euro IV	2007	2009	15,33	9,09	4,47	79,06	79,83	80,58	93,56	93,24	92,92	63,20	64,47	65,49	95,53	94,57	94,05
Buses	Diesel Urban Buses >18t	Euro V	2010	2013	14,29	8,13	3,44	78,89	79,66	80,38	93,49	93,15	92,82	78,07	78,86	79,49	95,49	94,52	93,98
Buses	Gasoline Coaches	Conventional	0	9999	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
Buses	Diesel Coaches <15t	Conventional	0	1993	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
Buses	Diesel Coaches <15t	Euro I	1994	1996	8,28	7,70	7,47	18,93	15,68	13,60	19,31	20,59	22,44	22,57	24,07	24,64	8,47	3,14	-1,19
Buses	Diesel Coaches <15t	Euro II	1997	2001	8,76	7,42	7,07	34,56	30,80	20,45	62,13	58,30	55,34	11,40	14,43	16,14	35,38	32,63	30,64
Buses	Diesel Coaches <15t	Euro III	2002	2006	0,80	0,43	0,70	14,91	15,75	6,50	54,54	55,87	57,10	24,13	31,47	34,19	36,41	34,05	30,99

Buses	Diesel Coaches <15t	Euro IV	2007	2009	5,01	1,44	0,69	54,25	53,24	49,77	90,25	89,66	88,99	49,96	52,11	52,89	92,04	91,07	90,12
Buses	Diesel Coaches <15t	Euro V	2010	2013	2,65	-0,53	-0,93	52,50	51,58	49,26	90,05	89,49	88,84	69,67	71,16	71,91	91,83	90,88	89,96
Buses	Diesel Coaches 15 - 18t	Conventional	0	1993	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
Buses	Diesel Coaches 15 - 18t	Euro I	1994	1996	8,28	7,70	7,47	18,93	15,68	13,60	19,31	20,59	22,44	22,57	24,07	24,64	8,47	3,14	-1,19
Buses	Diesel Coaches 15 - 18t	Euro II	1997	2001	8,76	7,42	7,07	34,56	30,80	20,45	62,13	58,30	55,34	11,40	14,43	16,14	35,38	32,63	30,64
Buses	Diesel Coaches 15 - 18t	Euro III	2002	2006	0,80	0,43	0,70	14,91	15,75	6,50	54,54	55,87	57,10	24,13	31,47	34,19	36,41	34,05	30,99
Buses	Diesel Coaches 15 - 18t	Euro IV	2007	2009	5,01	1,44	0,69	54,25	53,24	49,77	90,25	89,66	88,99	49,96	52,11	52,89	92,04	91,07	90,12
Buses	Diesel Coaches 15 - 18t	Euro V	2010	2013	2,65	-0,53	-0,93	52,50	51,58	49,26	90,05	89,49	88,84	69,67	71,16	71,91	91,83	90,88	89,96
Buses	Diesel Coaches >18t	Conventional	0	1993	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
Buses	Diesel Coaches >18t	Euro I	1994	1996	11,38	10,72	10,58	19,11	15,65	15,87	21,00	21,97	25,46	23,77	25,28	25,99	9,71	6,78	5,13
Buses	Diesel Coaches >18t	Euro II	1997	2001	12,90	11,45	11,20	34,56	31,66	25,53	62,65	58,54	56,71	15,63	18,15	19,50	37,83	37,12	36,81
Buses	Diesel Coaches >18t	Euro III	2002	2006	9,75	11,17	12,01	17,62	18,24	16,92	57,74	59,71	61,98	31,26	37,05	39,51	39,99	40,51	40,69
Buses	Diesel Coaches >18t	Euro IV	2007	2009	14,03	12,71	12,20	57,22	57,13	57,19	90,88	90,59	90,41	54,36	56,11	56,59	92,52	91,99	91,37
Buses	Diesel Coaches >18t	Euro V	2010	2013	11,70	10,73	10,51	56,10	56,11	56,24	90,69	90,43	90,28	72,44	73,59	73,96	92,33	91,84	91,19
Mopeds	2-stroke <50 cm³	Conventional	0	1999	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
Mopeds	2-stroke <50 cm³	Euro I	2000	2003	20,00	20,00	20,00	68,71	68,71	68,71	74,43	74,43	74,43	-221,43	-221,43	-221,43	59,52	59,52	59,52
Mopeds	2-stroke <50 cm³	Euro II	2004	2014	20,00	20,00	20,00	80,95	80,95	80,95	85,23	85,23	85,23	-203,57	-203,57	-203,57	69,05	69,05	69,05
Mopeds	4-stroke <50 cm³	Euro II	2004	2014	20,00	20,00	20,00	71,43	71,43	71,43	96,02	96,02	96,02	-203,57	-203,57	-203,57	90,60	90,60	90,60
Motorcycles	2-stroke >50 cm³	Conventional	0	1999	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
Motorcycles	2-stroke >50 cm³	Euro I	2000	2003	0,00	0,00	0,00	33,90	33,70	33,50	60,00	60,00	60,00	0,00	0,00	0,00	0,00	0,00	0,00
Motorcycles	2-stroke >50 cm³	Euro II	2004	2006	8,20	9,50	10,60	47,80	47,80	47,70	80,00	80,00	80,00	-38,70	-68,10	-70,70	69,10	63,70	57,60
Motorcycles	2-stroke >50 cm³	Euro III	2007	9999	8,20	9,50	10,60	71,10	71,00	70,70	94,00	94,00	94,00	-64,90	-93,70	-98,30	83,50	80,50	77,10
Motorcycles	4-stroke <250 cm³	Conventional	0	1999	8,20	9,50	10,60	0,00	0,00	0,00	0,00	0,00	0,00	22,10	-0,90	-15,50	89,70	88,00	85,90
Motorcycles	4-stroke <250 cm³	Euro I	2000	2003	0,00	0,00	0,00	31,90	16,00	0,20	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
Motorcycles	4-stroke <250 cm³	Euro II	2004	2006	17,60	17,20	16,30	75,70	66,50	63,40	75,00	75,00	75,00	-6,10	-5,40	-5,50	16,10	2,60	1,10
Motorcycles	4-stroke <250 cm³	Euro III	2007	9999	28,70	33,00	35,70	86,50	81,40	79,60	75,00	75,00	75,00	16,90	9,90	-7,90	63,30	49,10	47,60
Motorcycles	4-stroke 250 - 750 cm³	Conventional	0	1999	28,70	33,00	35,70	0,00	0,00	0,00	0,00	0,00	0,00	24,10	24,90	10,10	77,20	68,00	66,50
Motorcycles	4-stroke 250 - 750 cm³	Euro I	2000	2003	0,00	0,00	0,00	48,20	53,80	54,50	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
Motorcycles	4-stroke 250 - 750 cm³	Euro II	2004	2006	8,50	5,70	0,30	89,10	87,50	73,50	75,00	75,00	75,00	0,20	2,50	5,40	44,50	39,20	23,30
Motorcycles	4-stroke 250 - 750 cm³	Euro III	2007	9999	17,10	13,40	7,90	94,00	93,10	85,40	75,00	75,00	75,00	70,40	67,00	56,50	75,00	67,20	43,00
Motorcycles	4-stroke >750 cm³	Conventional	0	1999	17,10	13,40	7,90	0,00	0,00	0,00	0,00	0,00	0,00	85,00	83,20	78,10	84,30	79,50	64,20
Motorcycles	4-stroke >750 cm³	Euro I	2000	2003	0,00	0,00	0,00	48,20	53,80	54,50	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
Motorcycles	4-stroke >750 cm³	Euro II	2004	2006	2,30	7,40	11,80	89,10	87,50	73,50	75,00	75,00	75,00	-14,20	-37,30	-17,00	56,40	59,00	57,80
Motorcycles	4-stroke >750 cm³	Euro III	2007	9999	6,40	8,70	9,30	94,00	93,10	85,40	75,00	75,00	75,00	16,90	-12,40	-23,50	84,60	81,20	73,20

Annex 3B-6 Deterioration factors in 2012

Sector	Subsector	Tech 2	FYear	LYear	COU	COR	COH	NOxU	NOxR	NOxH	VOCU	VOCR	VOCH
Passenger Cars	Gasoline 0,8 - 1,4 l	PRE ECE	0	1969	1	1	1	1	1	1	1	1	1
Passenger Cars	Gasoline 0,8 - 1,4 l	ECE 15/00-01	1970	1978	1	1	1	1	1	1	1	1	1
Passenger Cars	Gasoline 0,8 - 1,4 l	ECE 15/02	1979	1980	1	1	1	1	1	1	1	1	1
Passenger Cars	Gasoline 0,8 - 1,4 l	ECE 15/03	1981	1985	1	1	1	1	1	1	1	1	1
Passenger Cars	Gasoline 0,8 - 1,4 l	ECE 15/04	1986	1990	1	1	1	1	1	1	1	1	1
Passenger Cars	Gasoline 0,8 - 1,4 l	Euro I	1991	1996	2,45676364	2,5358	2,5358	2,050881818	1,888	1,888	1,862736364	1,5974	1,5974
Passenger Cars	Gasoline 0,8 - 1,4 l	Euro II	1997	2000	2,45676364	2,5358	2,5358	2,050881818	1,888	1,888	1,862736364	1,5974	1,5974
Passenger Cars	Gasoline 0,8 - 1,4 l	Euro III	2001	2005	1,43682734	1,1507381	1,1507381	1	1	1	1,175928207	1	1
Passenger Cars	Gasoline 0,8 - 1,4 l	Euro IV	2006	2010	1,14113537	1,05078521	1,05078521	1	1	1	1,056995935	1	1
Passenger Cars	Gasoline 0,8 - 1,4 l	Euro V	2011	2014	0,93262402	0,98030202	0,98030202	1	1	1	0,973129172	1	1
Passenger Cars	Gasoline 1,4 - 2,0 l	PRE ECE	0	1969	1	1	1	1	1	1	1	1	1
Passenger Cars	Gasoline 1,4 - 2,0 l	ECE 15/00-01	1970	1978	1	1	1	1	1	1	1	1	1
Passenger Cars	Gasoline 1,4 - 2,0 l	ECE 15/02	1979	1980	1	1	1	1	1	1	1	1	1
Passenger Cars	Gasoline 1,4 - 2,0 l	ECE 15/03	1981	1985	1	1	1	1	1	1	1	1	1
Passenger Cars	Gasoline 1,4 - 2,0 l	ECE 15/04	1986	1990	1	1	1	1	1	1	1	1	1
Passenger Cars	Gasoline 1,4 - 2,0 l	Euro I	1991	1996	1,84864636	1,76984	1,76984	2,050881818	1,888	1,888	1,891659091	1,7868	1,7868
Passenger Cars	Gasoline 1,4 - 2,0 l	Euro II	1997	2000	1,84864636	1,76984	1,76984	2,050881818	1,888	1,888	1,891659091	1,7868	1,7868
Passenger Cars	Gasoline 1,4 - 2,0 l	Euro III	2001	2005	1,18858397	1	1	1,281105214	1	1	1	1	1
Passenger Cars	Gasoline 1,4 - 2,0 l	Euro IV	2006	2010	1,10408306	1	1	1,154955172	1	1	1	1	1
Passenger Cars	Gasoline 1,4 - 2,0 l	Euro V	2011	2014	1,00889333	1	1	1,012847962	1	1	1	1	1
Passenger Cars	Gasoline >2,0 l	PRE ECE	0	1969	1	1	1	1	1	1	1	1	1
Passenger Cars	Gasoline >2,0 l	ECE 15/00-01	1970	1978	1	1	1	1	1	1	1	1	1
Passenger Cars	Gasoline >2,0 l	ECE 15/02	1979	1980	1	1	1	1	1	1	1	1	1
Passenger Cars	Gasoline >2,0 l	ECE 15/03	1981	1985	1	1	1	1	1	1	1	1	1
Passenger Cars	Gasoline >2,0 l	ECE 15/04	1986	1990	1	1	1	1	1	1	1	1	1
Passenger Cars	Gasoline >2,0 l	Euro I	1991	1996	1,44665364	1,19748	1,19748	2,050881818	1,888	1,888	1,677460909	1,45388	1,45388
Passenger Cars	Gasoline >2,0 l	Euro II	1997	2000	1,44665364	1,19748	1,19748	2,050881818	1,888	1,888	1,677460909	1,45388	1,45388
Passenger Cars	Gasoline >2,0 l	Euro III	2001	2005	1,19577486	1	1	1,291840383	1	1	1	1	1
Passenger Cars	Gasoline >2,0 l	Euro IV	2006	2010	1,12718711	1	1	1,189446828	1	1	1	1	1
Passenger Cars	Gasoline >2,0 l	Euro V	2011	2014	1,01466181	1	1	1,02145962	1	1	1	1	1
Passenger Cars	Diesel 1,4 - 2,0 l	Conventional	0	1990	1	1	1	1	1	1	1	1	1
Passenger Cars	Diesel 1,4 - 2,0 l	Euro I	1991	1996	1	1	1	1	1	1	1	1	1
Passenger Cars	Diesel 1,4 - 2,0 l	Euro II	1997	2000	1	1	1	1	1	1	1	1	1

Passenger Cars	Diesel 1,4 - 2,0 l	Euro III	2001	2005	1	1	1	1	1	1	1	1	1
Passenger Cars	Diesel 1,4 - 2,0 l	Euro IV	2006	2010	1	1	1	1	1	1	1	1	1
Passenger Cars	Diesel 1,4 - 2,0 l	Euro V	2011	2014	1	1	1	1	1	1	1	1	1
Passenger Cars	Diesel >2,0 l	Conventional	0	1990	1	1	1	1	1	1	1	1	1
Passenger Cars	Diesel >2,0 l	Euro I	1991	1996	1	1	1	1	1	1	1	1	1
Passenger Cars	Diesel >2,0 l	Euro II	1997	2000	1	1	1	1	1	1	1	1	1
Passenger Cars	Diesel >2,0 l	Euro III	2001	2005	1	1	1	1	1	1	1	1	1
Passenger Cars	Diesel >2,0 l	Euro IV	2006	2010	1	1	1	1	1	1	1	1	1
Passenger Cars	Diesel >2,0 l	Euro V	2011	2014	1	1	1	1	1	1	1	1	1
Passenger Cars	LPG cars	Conventional	0	1990	1	1	1	1	1	1	1	1	1
Passenger Cars	LPG cars	Euro I	1991	1996	1	1	1	1	1	1	1	1	1
Passenger Cars	LPG cars	Euro II	1997	2000	1	1	1	1	1	1	1	1	1
Passenger Cars	LPG cars	Euro III	2001	2005	1	1	1	1	1	1	1	1	1
Passenger Cars	LPG cars	Euro IV	2006	2010	1	1	1	1	1	1	1	1	1
Passenger Cars	Electric cars	Conventional	0	9999	1	1	1	1	1	1	1	1	1
Passenger Cars	Gasoline <0,8 l	PRE ECE	0	1969	1	1	1	1	1	1	1	1	1
Passenger Cars	Gasoline <0,8 l	ECE 15/00-01	1970	1978	1	1	1	1	1	1	1	1	1
Passenger Cars	Gasoline <0,8 l	ECE 15/02	1979	1980	1	1	1	1	1	1	1	1	1
Passenger Cars	Gasoline <0,8 l	ECE 15/03	1981	1985	1	1	1	1	1	1	1	1	1
Passenger Cars	Gasoline <0,8 l	ECE 15/04	1986	1990	1	1	1	1	1	1	1	1	1
Passenger Cars	Gasoline <0,8 l	Euro I	1991	1996	1,8900476	1,938392	1,938392	1,549472614	1,456479718	1,456479718	1,52718272	1,365015717	1,365015717
Passenger Cars	Gasoline <0,8 l	Euro II	1997	2000	2,36122605	2,43508833	2,43508833	1,966353722	1,815253853	1,815253853	1,806168374	1,558224411	1,558224411
Passenger Cars	Gasoline <0,8 l	Euro III	2001	2005	1,43155293	1,14895519	1,14895519	1	1	1	1,173806751	1	1
Passenger Cars	Gasoline <0,8 l	Euro IV	2006	2010	1,15421885	1,05520782	1,05520782	1	1	1	1,062258331	1	1
Passenger Cars	Gasoline <0,8 l	Euro V	2011	2014	0,93830805	0,98222339	0,98222339	1	1	1	0,975415386	1	1
Passenger Cars	Diesel <1,4 l	Conventional	0	1990	1	1	1	1	1	1	1	1	1
Passenger Cars	Diesel <1,4 l	Euro I	1991	1996	1	1	1	1	1	1	1	1	1
Passenger Cars	Diesel <1,4 l	Euro II	1997	2000	1	1	1	1	1	1	1	1	1
Passenger Cars	Diesel <1,4 l	Euro III	2001	2005	1	1	1	1	1	1	1	1	1
Passenger Cars	Diesel <1,4 l	Euro IV	2006	2010	1	1	1	1	1	1	1	1	1
Passenger Cars	Diesel <1,4 l	Euro V	2011	2014	1	1	1	1	1	1	1	1	1
Light Duty Vehicles	Gasoline <3,5t	Conventional	0	1994	1	1	1	1	1	1	1	1	1
Light Duty Vehicles	Gasoline <3,5t	Euro I	1995	1998	2,45676364	2,5358	2,5358	2,050881818	1,888	1,888	1,862736364	1,5974	1,5974
Light Duty Vehicles	Gasoline <3,5t	Euro II	1999	2001	2,45676364	2,5358	2,5358	2,050881818	1,888	1,888	1,862736364	1,5974	1,5974
Light Duty Vehicles	Gasoline <3,5t	Euro III	2002	2006	1,16742689	1	1	1,249520161	1	1	1	1	1
Light Duty Vehicles	Gasoline <3,5t	Euro IV	2007	2011	1,08932894	1	1	1,132928986	1	1	1	1	1
Light Duty Vehicles	Gasoline <3,5t	Euro V	2012	2015	0,98987251	1	1	0,984452072	1	1	1	1	1

Annex 3B-7 Final fuel consumption factors (MJ/km) and emission factors (g/km) in 2012

Sector	Subsector	Tech 2	ForecastYear	Milu	Milr	MilH	FCu_MJ	FCr_MJ	FCh_MJ	FCu_g	FCr_g	FCh_g	CO2_u	CO2_r	CO2_h	NOx_u	NOx_r	NOx_h
Passenger Cars	Gasoline 0,8 - 1,4 l	PRE ECE	2012	6948	17596	10159	4,336	2,409	2,748	101	56	64	305	170	194	2,051	2,062	2,023
Passenger Cars	Gasoline 0,8 - 1,4 l	ECE 15/00-01	2012	5025	12724	7347	3,741	1,947	2,129	87	45	50	263	137	150	2,051	2,062	2,023
Passenger Cars	Gasoline 0,8 - 1,4 l	ECE 15/02	2012	680	1722	994	3,420	1,978	2,243	80	46	52	241	139	158	1,796	2,102	2,909
Passenger Cars	Gasoline 0,8 - 1,4 l	ECE 15/03	2012	3957	10022	5786	3,420	1,978	2,243	80	46	52	241	139	158	1,864	2,253	3,276
Passenger Cars	Gasoline 0,8 - 1,4 l	ECE 15/04	2012	17966	45496	26269	3,303	1,903	2,089	77	44	49	233	134	147	1,876	2,089	2,662
Passenger Cars	Gasoline 0,8 - 1,4 l	Euro I	2012	143072	362319	209195	3,045	1,838	2,017	71	43	47	214	129	142	1,148	0,531	0,864
Passenger Cars	Gasoline 0,8 - 1,4 l	Euro II	2012	227787	576853	333062	3,005	1,731	1,928	70	40	45	212	122	136	0,739	0,291	0,342
Passenger Cars	Gasoline 0,8 - 1,4 l	Euro III	2012	335153	848750	490049	3,101	1,851	1,985	72	43	46	218	130	140	0,264	0,060	0,052
Passenger Cars	Gasoline 0,8 - 1,4 l	Euro IV	2012	679359	1720425	993334	2,739	1,563	1,708	64	36	40	193	110	120	0,160	0,030	0,019
Passenger Cars	Gasoline 0,8 - 1,4 l	Euro V	2012	363791	921273	531922	2,625	1,463	1,598	61	34	37	185	103	113	0,146	0,023	0,014
Passenger Cars	Gasoline 1,4 - 2,0 l	PRE ECE	2012	14599	36971	21346	5,093	2,935	3,346	119	69	78	359	207	236	2,401	2,683	3,130
Passenger Cars	Gasoline 1,4 - 2,0 l	ECE 15/00-01	2012	8722	22087	12752	4,354	2,238	2,641	102	52	62	307	158	186	2,401	2,683	3,130
Passenger Cars	Gasoline 1,4 - 2,0 l	ECE 15/02	2012	1131	2864	1654	3,965	2,220	2,614	93	52	61	279	156	184	2,031	2,377	3,283
Passenger Cars	Gasoline 1,4 - 2,0 l	ECE 15/03	2012	7136	18072	10434	3,965	2,220	2,614	93	52	61	279	156	184	2,126	2,580	3,472
Passenger Cars	Gasoline 1,4 - 2,0 l	ECE 15/04	2012	34941	88484	51089	3,960	2,151	2,280	92	50	53	279	151	161	2,354	2,757	3,524
Passenger Cars	Gasoline 1,4 - 2,0 l	Euro I	2012	365004	924345	533695	3,695	2,125	2,257	86	50	53	260	150	159	1,131	0,531	0,864
Passenger Cars	Gasoline 1,4 - 2,0 l	Euro II	2012	742835	1881173	1086146	3,643	2,089	2,137	85	49	50	257	147	150	0,727	0,291	0,342
Passenger Cars	Gasoline 1,4 - 2,0 l	Euro III	2012	754314	1910243	1102930	3,727	2,185	2,325	87	51	54	262	154	164	0,280	0,060	0,052
Passenger Cars	Gasoline 1,4 - 2,0 l	Euro IV	2012	558941	1415476	817263	3,635	2,147	2,249	85	50	52	256	151	158	0,166	0,030	0,019
Passenger Cars	Gasoline 1,4 - 2,0 l	Euro V	2012	82516	208965	120652	3,503	2,031	2,127	82	47	50	247	143	150	0,144	0,023	0,014
Passenger Cars	Gasoline >2,0 l	PRE ECE	2012	6241	15805	9125	6,201	3,504	3,866	145	82	90	437	247	272	3,173	4,090	5,500
Passenger Cars	Gasoline >2,0 l	ECE 15/00-01	2012	4864	12318	7112	4,741	2,501	2,904	111	58	68	334	176	205	3,173	4,090	5,500
Passenger Cars	Gasoline >2,0 l	ECE 15/02	2012	537	1359	785	4,835	2,771	3,097	113	65	72	341	195	218	2,292	2,675	3,680
Passenger Cars	Gasoline >2,0 l	ECE 15/03	2012	1626	4119	2378	4,835	2,771	3,097	113	65	72	341	195	218	3,113	3,441	4,604
Passenger Cars	Gasoline >2,0 l	ECE 15/04	2012	5823	14747	8514	4,564	2,544	3,062	107	59	71	321	179	216	2,543	2,750	3,687
Passenger Cars	Gasoline >2,0 l	Euro I	2012	28584	72387	41795	4,793	2,711	2,848	112	63	66	338	191	201	0,993	0,531	0,864
Passenger Cars	Gasoline >2,0 l	Euro II	2012	59066	149579	86364	4,890	2,857	2,923	114	67	68	344	201	206	0,627	0,291	0,342
Passenger Cars	Gasoline >2,0 l	Euro III	2012	118155	299220	172762	4,626	2,515	2,489	108	59	58	326	177	175	0,237	0,060	0,052
Passenger Cars	Gasoline >2,0 l	Euro IV	2012	84269	213406	123216	5,586	3,202	3,107	130	75	73	393	225	219	0,142	0,030	0,019
Passenger Cars	Gasoline >2,0 l	Euro V	2012	4209	10659	6154	5,517	3,147	3,053	129	73	71	389	222	215	0,119	0,023	0,014
Passenger Cars	Diesel 1,4 - 2,0 l	Euro I	2012	48565	122988	71011	2,705	1,828	2,066	64	43	49	188	127	143	0,743	0,562	0,663
Passenger Cars	Diesel 1,4 - 2,0 l	Euro II	2012	147906	374561	216263	2,852	1,884	2,083	67	44	49	198	131	144	0,803	0,555	0,665
Passenger Cars	Diesel 1,4 - 2,0 l	Euro III	2012	625682	1584491	914849	2,766	1,854	1,946	65	44	46	192	129	135	0,883	0,665	0,750

Passenger Cars	Diesel 1,4 - 2,0 l	Euro IV	2012	714935	1810517	1045350	2,715	1,809	1,899	64	43	45	188	125	132	0,665	0,424	0,576
Passenger Cars	Diesel 1,4 - 2,0 l	Euro V	2012	1415407	3584409	2069555	2,578	1,687	1,769	61	40	42	179	117	123	0,818	0,521	0,708
Passenger Cars	Diesel 1,4 - 2,0 l	Conventional	2012	22421	56780	32784	3,253	1,760	2,139	77	42	51	226	122	148	0,641	0,433	0,528
Passenger Cars	Diesel >2,0 l	Euro I	2012	16926	42865	24749	3,691	2,489	2,748	87	59	65	256	173	191	0,743	0,562	0,663
Passenger Cars	Diesel >2,0 l	Euro II	2012	60694	153704	88745	3,691	2,489	2,748	87	59	65	256	173	191	0,803	0,555	0,665
Passenger Cars	Diesel >2,0 l	Euro III	2012	157901	399873	230877	3,691	2,489	2,748	87	59	65	256	173	191	0,883	0,665	0,750
Passenger Cars	Diesel >2,0 l	Euro IV	2012	146877	371956	214759	4,007	2,772	3,060	95	65	72	278	192	212	0,665	0,424	0,576
Passenger Cars	Diesel >2,0 l	Euro V	2012	202435	512650	295992	3,653	2,455	2,710	86	58	64	253	170	188	0,818	0,521	0,708
Passenger Cars	Diesel >2,0 l	Conventional	2012	5063	12821	7402	3,253	1,760	2,139	77	42	51	226	122	148	1,016	0,723	0,861
Passenger Cars	LPG cars	Euro I	2012	7	19	11	3,316	2,077	2,490	72	45	54	209	131	157	0,314	0,283	0,298
Passenger Cars	LPG cars	Euro II	2012	14	34	20	3,316	2,077	2,490	72	45	54	209	131	157	0,104	0,102	0,107
Passenger Cars	LPG cars	Euro III	2012	11	27	16	3,316	2,077	2,490	72	45	54	209	131	157	0,073	0,068	0,071
Passenger Cars	LPG cars	Euro IV	2012	17	42	24	3,316	2,077	2,490	72	45	54	209	131	157	0,039	0,037	0,039
Passenger Cars	LPG cars	Conventional	2012	11	27	16	3,980	2,070	2,484	87	45	54	251	131	157	2,034	2,584	2,861
Passenger Cars	Gasoline <0,8 l	PRE ECE	2012	220	558	322	3,357	2,025	2,477	78	47	58	236	143	174	2,051	2,062	2,023
Passenger Cars	Gasoline <0,8 l	ECE 15/00-01	2012	198	502	290	2,896	1,637	1,918	68	38	45	204	115	135	2,051	2,062	2,023
Passenger Cars	Gasoline <0,8 l	ECE 15/02	2012	6	15	8	2,648	1,663	2,021	62	39	47	186	117	142	1,796	2,102	2,909
Passenger Cars	Gasoline <0,8 l	ECE 15/03	2012	51	128	74	2,648	1,663	2,021	62	39	47	186	117	142	1,864	2,253	3,276
Passenger Cars	Gasoline <0,8 l	ECE 15/04	2012	63	160	92	2,557	1,599	1,883	60	37	44	180	113	133	1,876	2,089	2,662
Passenger Cars	Gasoline <0,8 l	Euro I	2012	138	348	201	2,357	1,545	1,818	55	36	42	166	109	128	1,011	0,410	0,666
Passenger Cars	Gasoline <0,8 l	Euro II	2012	572	1450	837	2,326	1,455	1,737	54	34	41	164	102	122	0,726	0,280	0,329
Passenger Cars	Gasoline <0,8 l	Euro III	2012	1988	5034	2907	2,401	1,556	1,789	56	36	42	169	110	126	0,264	0,060	0,052
Passenger Cars	Gasoline <0,8 l	Euro IV	2012	6034	15281	8823	2,222	1,412	1,653	52	33	39	156	99	116	0,160	0,030	0,019
Passenger Cars	Gasoline <0,8 l	Euro V	2012	515	1305	753	1,973	1,173	1,374	46	27	32	139	83	97	0,146	0,023	0,014
Passenger Cars	Diesel <1,4 l	Euro I	2012	347	878	507	1,719	1,366	1,679	41	32	40	119	95	116	0,743	0,562	0,663
Passenger Cars	Diesel <1,4 l	Euro II	2012	6688	16936	9779	1,813	1,408	1,692	43	33	40	126	98	117	0,803	0,555	0,665
Passenger Cars	Diesel <1,4 l	Euro III	2012	91206	230972	133358	1,758	1,386	1,582	42	33	37	122	96	110	0,883	0,665	0,750
Passenger Cars	Diesel <1,4 l	Euro IV	2012	140834	356652	205923	1,889	1,522	1,737	45	36	41	131	106	121	0,665	0,424	0,576
Passenger Cars	Diesel <1,4 l	Euro V	2012	273197	691850	399458	1,820	1,450	1,653	43	34	39	126	101	115	0,818	0,521	0,708
Passenger Cars	Diesel <1,4 l	Conventional	2012	40	100	58	2,068	1,315	1,738	49	31	41	143	91	121	0,641	0,433	0,528
Light Duty Vehicles	Gasoline <3,5t	Conventional	2012	5692	14238	7724	5,239	2,623	2,473	122	61	58	369	185	174	2,955	3,118	3,387
Light Duty Vehicles	Gasoline <3,5t	Euro I	2012	18381	45976	24940	6,142	3,083	2,911	143	72	68	433	217	205	1,532	0,755	0,810
Light Duty Vehicles	Gasoline <3,5t	Euro II	2012	14801	37022	20083	6,142	3,083	2,911	143	72	68	433	217	205	0,771	0,257	0,276
Light Duty Vehicles	Gasoline <3,5t	Euro III	2012	35902	89798	48713	6,142	3,083	2,911	143	72	68	433	217	205	0,322	0,084	0,090
Light Duty Vehicles	Gasoline <3,5t	Euro IV	2012	13538	33861	18368	6,142	3,083	2,911	143	72	68	433	217	205	0,167	0,040	0,043
Light Duty Vehicles	Gasoline <3,5t	Euro V	2012	491	1229	667	6,142	3,083	2,911	143	72	68	433	217	205	0,150	0,030	0,032
Light Duty Vehicles	Diesel <3,5t	Conventional	2012	23203	58037	31483	4,309	2,815	3,080	102	66	73	299	195	214	2,052	0,843	0,834

Light Duty Vehicles	Diesel <3,5t	Euro I	2012	109438	273730	148489	3,868	2,484	2,718	91	59	64	268	172	189	1,396	0,975	1,022
Light Duty Vehicles	Diesel <3,5t	Euro II	2012	132946	332530	180386	3,868	2,484	2,718	91	59	64	268	172	189	1,396	0,975	1,022
Light Duty Vehicles	Diesel <3,5t	Euro III	2012	590213	1476262	800824	3,868	2,484	2,718	91	59	64	268	172	189	1,173	0,819	0,859
Light Duty Vehicles	Diesel <3,5t	Euro IV	2012	497535	1244453	675075	3,868	2,484	2,718	91	59	64	268	172	189	0,949	0,663	0,695
Light Duty Vehicles	Diesel <3,5t	Euro V	2012	280459	701493	380537	3,868	2,484	2,718	91	59	64	268	172	189	0,684	0,478	0,501
Light Duty Vehicles	LPG <3,5t	Conventional	2012	6	15	8	5,919	3,105	3,726	129	68	81	373	196	235	3,058	3,876	4,291
Light Duty Vehicles	LPG <3,5t	Euro II	2012	8	19	10	4,930	3,116	3,735	107	68	81	311	197	236	0,171	0,153	0,161
Light Duty Vehicles	LPG <3,5t	Euro III	2012	32	81	44	4,930	3,116	3,735	107	68	81	311	197	236	0,115	0,102	0,107
Light Duty Vehicles	LPG <3,5t	Euro IV	2012	15	39	21	4,930	3,116	3,735	107	68	81	311	197	236	0,059	0,055	0,058
Heavy Duty Vehicles	Gasoline >3,5t	Conventional	2012	582	2117	1092	9,855	6,570	7,227	230	153	169	694	463	509	4,500	7,500	7,500
Heavy Duty Vehicles	Diesel RT 3,5 - 7,5t	Conventional	2012	321	1168	603	5,338	4,739	4,824	126	112	114	370	329	335	4,211	4,104	4,476
Heavy Duty Vehicles	Diesel RT 3,5 - 7,5t	Euro I	2012	222	808	417	4,272	3,915	4,450	101	92	105	296	272	309	2,939	2,938	3,316
Heavy Duty Vehicles	Diesel RT 3,5 - 7,5t	Euro II	2012	1417	5150	2657	4,056	3,783	4,313	96	89	102	281	262	299	3,223	3,118	3,414
Heavy Duty Vehicles	Diesel RT 3,5 - 7,5t	Euro III	2012	3616	13144	6781	4,329	3,966	4,523	102	94	107	300	275	314	2,499	2,300	2,498
Heavy Duty Vehicles	Diesel RT 3,5 - 7,5t	Euro IV	2012	4336	15759	8129	4,208	3,967	4,552	99	94	108	292	275	316	1,707	1,645	1,801
Heavy Duty Vehicles	Diesel RT 3,5 - 7,5t	Euro V	2012	2380	8651	4463	4,076	3,810	4,362	96	90	103	283	264	303	1,957	0,821	0,447
Heavy Duty Vehicles	Diesel RT 3,5 - 7,5t	Euro VI	2012	3	12	6	4,195	3,891	4,416	99	92	104	291	270	306	0,233	0,097	0,060
Heavy Duty Vehicles	Diesel RT 7,5 - 12t	Conventional	2012	838	3045	1571	7,825	6,538	6,408	185	154	151	543	454	444	7,928	7,236	7,499
Heavy Duty Vehicles	Diesel RT 7,5 - 12t	Euro I	2012	679	2469	1274	6,656	5,787	5,836	157	137	138	462	401	405	4,729	4,306	4,464
Heavy Duty Vehicles	Diesel RT 7,5 - 12t	Euro II	2012	2288	8316	4290	6,346	5,605	5,702	150	132	135	440	389	396	5,152	4,593	4,682
Heavy Duty Vehicles	Diesel RT 7,5 - 12t	Euro III	2012	5636	20485	10567	6,728	5,883	5,935	159	139	140	467	408	412	3,997	3,536	3,485
Heavy Duty Vehicles	Diesel RT 7,5 - 12t	Euro IV	2012	5143	18693	9643	6,467	5,814	5,916	153	137	140	449	403	410	2,728	2,512	2,488
Heavy Duty Vehicles	Diesel RT 7,5 - 12t	Euro V	2012	5289	19226	9918	6,317	5,629	5,702	149	133	135	438	390	396	3,114	1,405	0,778
Heavy Duty Vehicles	Diesel RT 7,5 - 12t	Euro VI	2012	6	21	11	6,471	5,728	5,758	153	135	136	449	397	399	0,384	0,154	0,096
Heavy Duty Vehicles	Diesel RT 12 - 14 t	Conventional	2012	110	400	207	8,477	6,973	6,798	200	165	161	588	484	472	8,826	7,718	7,748
Heavy Duty Vehicles	Diesel RT 12 - 14 t	Euro I	2012	83	300	155	7,266	6,162	6,120	172	146	145	504	427	425	5,321	4,638	4,638
Heavy Duty Vehicles	Diesel RT 12 - 14 t	Euro II	2012	263	955	493	6,970	5,979	5,961	165	141	141	483	415	413	5,815	4,975	4,889
Heavy Duty Vehicles	Diesel RT 12 - 14 t	Euro III	2012	270	982	507	7,351	6,238	6,175	174	147	146	510	433	428	4,745	3,881	3,702
Heavy Duty Vehicles	Diesel RT 12 - 14 t	Euro IV	2012	303	1101	568	6,965	6,103	6,118	164	144	144	483	423	424	3,208	2,754	2,620
Heavy Duty Vehicles	Diesel RT 12 - 14 t	Euro V	2012	351	1277	659	6,828	5,922	5,890	161	140	139	474	411	409	3,478	1,582	0,881
Heavy Duty Vehicles	Diesel RT 14 - 20t	Conventional	2012	446	1620	836	11,173	8,785	8,248	264	207	195	775	609	572	11,287	9,455	9,120
Heavy Duty Vehicles	Diesel RT 14 - 20t	Euro I	2012	473	1718	886	9,088	7,350	7,020	215	174	166	630	510	487	6,721	5,601	5,385
Heavy Duty Vehicles	Diesel RT 14 - 20t	Euro II	2012	2391	8690	4483	8,724	7,142	6,846	206	169	162	605	495	475	7,473	6,118	5,804
Heavy Duty Vehicles	Diesel RT 14 - 20t	Euro III	2012	6355	23099	11916	9,195	7,421	7,042	217	175	166	638	515	488	6,139	4,859	4,431
Heavy Duty Vehicles	Diesel RT 14 - 20t	Euro IV	2012	6370	23154	11944	8,587	7,177	6,916	203	170	163	596	498	480	4,079	3,400	3,171
Heavy Duty Vehicles	Diesel RT 14 - 20t	Euro V	2012	8415	30586	15778	8,572	7,045	6,710	202	166	158	595	489	465	4,749	2,641	1,675
Heavy Duty Vehicles	Diesel RT 14 - 20t	Euro VI	2012	10	36	18	8,568	7,081	6,773	202	167	160	594	491	470	0,694	0,262	0,135

Heavy Duty Vehicles	Diesel RT 20 - 26t	Conventional	2012	277	1008	520	13,489	10,388	9,495	319	245	224	936	721	659	12,251	9,862	9,114
Heavy Duty Vehicles	Diesel RT 20 - 26t	Euro I	2012	531	1931	996	11,521	9,050	8,362	272	214	197	799	628	580	8,634	6,952	6,468
Heavy Duty Vehicles	Diesel RT 20 - 26t	Euro II	2012	3222	11710	6041	11,147	8,848	8,190	263	209	193	773	614	568	9,465	7,549	6,947
Heavy Duty Vehicles	Diesel RT 20 - 26t	Euro III	2012	10842	39406	20328	11,646	9,122	8,356	275	215	197	808	633	580	7,649	6,024	5,545
Heavy Duty Vehicles	Diesel RT 20 - 26t	Euro IV	2012	11186	40657	20973	10,999	8,858	8,223	260	209	194	763	614	570	5,146	4,223	3,967
Heavy Duty Vehicles	Diesel RT 20 - 26t	Euro V	2012	14725	53523	27610	10,804	8,631	7,974	255	204	188	749	599	553	5,824	2,914	1,627
Heavy Duty Vehicles	Diesel RT 20 - 26t	Euro VI	2012	9	31	16	10,924	8,716	8,049	258	206	190	758	605	558	0,710	0,287	0,164
Heavy Duty Vehicles	Diesel RT 26 - 28t	Conventional	2012	3	10	5	14,261	11,014	9,970	337	260	235	989	764	692	12,868	10,379	9,526
Heavy Duty Vehicles	Diesel RT 26 - 28t	Euro I	2012	2	8	4	12,232	9,624	8,799	289	227	208	848	668	610	9,122	7,308	6,742
Heavy Duty Vehicles	Diesel RT 26 - 28t	Euro II	2012	3	10	5	11,823	9,401	8,622	279	222	204	820	652	598	9,876	7,848	7,164
Heavy Duty Vehicles	Diesel RT 26 - 28t	Euro III	2012	19	67	35	12,354	9,712	8,830	292	229	209	857	674	612	7,733	6,089	5,633
Heavy Duty Vehicles	Diesel RT 26 - 28t	Euro IV	2012	13	47	24	11,835	9,518	8,710	280	225	206	821	660	604	5,258	4,284	4,029
Heavy Duty Vehicles	Diesel RT 26 - 28t	Euro V	2012	32	115	59	11,469	9,153	8,338	271	216	197	796	635	578	6,554	3,004	1,426
Heavy Duty Vehicles	Diesel RT 28 - 32t	Conventional	2012	14	50	26	15,791	12,478	11,346	373	295	268	1095	866	787	14,515	11,942	11,008
Heavy Duty Vehicles	Diesel RT 28 - 32t	Euro I	2012	59	214	111	13,865	11,099	10,170	327	262	240	962	770	705	10,453	8,509	7,843
Heavy Duty Vehicles	Diesel RT 28 - 32t	Euro II	2012	1135	4125	2128	13,380	10,738	10,252	316	254	242	928	745	711	11,232	9,043	8,280
Heavy Duty Vehicles	Diesel RT 28 - 32t	Euro III	2012	7107	25832	13326	13,989	11,225	10,242	330	265	242	970	779	710	8,883	7,017	6,445
Heavy Duty Vehicles	Diesel RT 28 - 32t	Euro IV	2012	8262	30029	15491	13,525	11,089	10,149	319	262	240	938	769	704	5,978	5,101	4,533
Heavy Duty Vehicles	Diesel RT 28 - 32t	Euro V	2012	9313	33852	17463	13,087	10,653	9,720	309	252	230	908	739	674	6,484	2,684	1,391
Heavy Duty Vehicles	Diesel RT 28 - 32t	Euro VI	2012	9	33	17	13,430	10,924	9,951	317	258	235	932	758	690	0,638	0,290	0,191
Heavy Duty Vehicles	Diesel RT >32t	Euro I	2012	2	9	4	14,022	10,936	9,924	331	258	234	973	759	688	10,614	8,446	7,666
Heavy Duty Vehicles	Diesel RT >32t	Euro III	2012	29	105	54	14,133	11,011	9,928	334	260	234	980	764	689	9,225	7,224	6,550
Heavy Duty Vehicles	Diesel RT >32t	Euro IV	2012	12	43	22	13,512	10,782	9,803	319	255	232	937	748	680	6,270	5,071	4,708
Heavy Duty Vehicles	Diesel RT >32t	Euro V	2012	53	191	99	13,145	10,402	9,416	310	246	222	912	722	653	6,646	2,837	1,521
Heavy Duty Vehicles	Diesel TT/AT 28 - 34t	Conventional	2012	146	1471	1953	14,181	10,863	9,705	335	257	229	984	753	673	13,305	10,460	9,286
Heavy Duty Vehicles	Diesel TT/AT 28 - 34t	Euro I	2012	145	1468	1951	12,683	9,797	8,769	300	231	207	880	680	608	9,509	7,408	6,570
Heavy Duty Vehicles	Diesel TT/AT 28 - 34t	Euro II	2012	728	7355	9770	11,962	9,377	8,674	283	221	205	830	650	602	10,046	7,771	6,867
Heavy Duty Vehicles	Diesel TT/AT 28 - 34t	Euro III	2012	1944	19642	26091	12,594	9,760	8,699	297	231	205	874	677	603	8,110	6,154	5,397
Heavy Duty Vehicles	Diesel TT/AT 28 - 34t	Euro IV	2012	1938	19586	26016	12,093	9,572	8,567	286	226	202	839	664	594	5,531	4,329	3,837
Heavy Duty Vehicles	Diesel TT/AT 28 - 34t	Euro V	2012	2556	25828	34308	11,833	9,300	8,288	279	220	196	821	645	575	5,295	2,493	1,513
Heavy Duty Vehicles	Diesel TT/AT 28 - 34t	Euro VI	2012	3	30	40	11,973	9,411	8,396	283	222	198	831	653	582	0,531	0,245	0,167
Heavy Duty Vehicles	Diesel TT/AT 34 - 40t	Conventional	2012	180	1822	2420	16,449	12,410	10,920	388	293	258	1141	861	758	15,378	11,908	10,419
Heavy Duty Vehicles	Diesel TT/AT 34 - 40t	Euro I	2012	274	2770	3680	14,440	11,007	9,732	341	260	230	1002	763	675	10,891	8,408	7,387
Heavy Duty Vehicles	Diesel TT/AT 34 - 40t	Euro II	2012	1551	15673	20819	14,078	10,815	9,559	333	255	226	977	750	663	11,695	8,978	7,885
Heavy Duty Vehicles	Diesel TT/AT 34 - 40t	Euro III	2012	6038	61004	81032	14,582	11,081	9,709	344	262	229	1011	769	673	9,414	7,197	6,354
Heavy Duty Vehicles	Diesel TT/AT 34 - 40t	Euro IV	2012	6168	62317	82777	13,969	10,851	9,575	330	256	226	969	753	664	6,398	5,061	4,523
Heavy Duty Vehicles	Diesel TT/AT 34 - 40t	Euro V	2012	16507	166787	221545	13,690	10,560	9,283	323	249	219	950	733	644	6,079	2,869	1,752

Heavy Duty Vehicles	Diesel TT/AT 34 - 40t	Euro VI	2012	25	254	337	13,824	10,661	9,376	326	252	221	959	740	650	0,610	0,285	0,191
Heavy Duty Vehicles	Diesel TT/AT 40 - 50t	Conventional	2012	57	574	763	18,259	13,816	12,105	431	326	286	1267	958	840	17,311	13,363	11,617
Heavy Duty Vehicles	Diesel TT/AT 40 - 50t	Euro I	2012	145	1464	1945	16,056	12,263	10,784	379	290	255	1114	851	748	12,142	9,377	8,189
Heavy Duty Vehicles	Diesel TT/AT 40 - 50t	Euro II	2012	1137	11486	15258	15,545	12,026	10,840	367	284	256	1078	834	752	12,955	9,936	8,683
Heavy Duty Vehicles	Diesel TT/AT 40 - 50t	Euro III	2012	10842	109550	145517	16,227	12,353	10,785	383	292	255	1126	857	748	10,432	7,969	6,995
Heavy Duty Vehicles	Diesel TT/AT 40 - 50t	Euro IV	2012	17764	179492	238421	15,683	12,170	10,666	370	287	252	1088	844	740	7,035	5,657	4,952
Heavy Duty Vehicles	Diesel TT/AT 40 - 50t	Euro V	2012	27152	274346	364417	15,260	11,749	10,255	360	277	242	1059	815	711	6,358	2,665	1,525
Heavy Duty Vehicles	Diesel TT/AT 40 - 50t	Euro VI	2012	62	631	838	15,508	11,944	10,428	366	282	246	1076	828	723	0,564	0,294	0,214
Heavy Duty Vehicles	Diesel TT/AT 50 - 60t	Euro II	2012	3	29	39	18,764	14,443	12,810	443	341	303	1302	1002	889	15,566	11,836	10,222
Heavy Duty Vehicles	Diesel TT/AT 50 - 60t	Euro IV	2012	6	56	74	18,830	14,628	12,811	445	345	303	1306	1015	889	8,477	6,746	5,764
Heavy Duty Vehicles	Diesel TT/AT >60t	Euro III	2012	5	52	69	19,297	14,647	13,013	456	346	307	1339	1016	903	12,548	9,501	8,223
Buses	Gasoline Urban Buses	Conventional	2012	123	126	19	9,855	6,570	7,227	230	153	169	694	463	509	4,500	7,500	7,500
Buses	Diesel Urban Buses <15t	Conventional	2012	99	101	16	11,353	9,012	8,430	268	213	199	788	625	585	9,347	7,678	7,133
Buses	Diesel Urban Buses <15t	Euro I	2012	220	224	35	9,175	7,454	6,918	217	176	163	636	517	480	6,945	5,531	4,861
Buses	Diesel Urban Buses <15t	Euro II	2012	1599	1629	251	8,856	7,275	6,774	209	172	160	614	505	470	7,552	5,971	5,224
Buses	Diesel Urban Buses <15t	Euro III	2012	10945	11148	1721	9,384	7,682	7,132	222	181	168	651	533	495	6,425	4,515	3,631
Buses	Diesel Urban Buses <15t	Euro IV	2012	27476	27985	4320	8,717	7,448	7,350	206	176	174	605	517	510	4,076	3,101	2,593
Buses	Diesel Urban Buses <15t	Euro V	2012	22441	22857	3528	8,470	7,314	6,949	200	173	164	588	507	482	4,841	2,676	1,933
Buses	Diesel Urban Buses 15 - 18t	Conventional	2012	2368	2412	372	14,440	11,180	9,824	341	264	232	1002	775	681	15,108	12,139	10,803
Buses	Diesel Urban Buses 15 - 18t	Euro I	2012	3440	3504	541	12,320	9,750	8,658	291	230	204	855	676	601	9,289	7,392	6,426
Buses	Diesel Urban Buses 15 - 18t	Euro II	2012	15359	15643	2415	11,941	9,600	8,628	282	227	204	828	666	599	9,989	7,828	6,822
Buses	Diesel Urban Buses 15 - 18t	Euro III	2012	36728	37408	5775	12,516	10,038	9,011	296	237	213	868	696	625	8,427	6,044	4,919
Buses	Diesel Urban Buses 15 - 18t	Euro IV	2012	35723	36385	5617	11,802	9,834	9,293	279	232	219	819	682	645	5,452	4,181	3,521
Buses	Diesel Urban Buses 15 - 18t	Euro V	2012	32710	33315	5143	11,432	9,470	8,955	270	224	212	793	657	621	6,085	2,796	2,156
Buses	Diesel Urban Buses >18t	Conventional	2012	17	18	3	18,125	14,109	12,176	428	333	288	1257	979	845	19,310	15,492	13,433
Buses	Diesel Urban Buses >18t	Euro I	2012	22	22	3	15,764	12,479	10,836	372	295	256	1093	866	752	11,840	9,361	8,043
Buses	Diesel Urban Buses >18t	Euro II	2012	1516	1544	238	15,291	12,318	11,322	361	291	267	1061	854	785	12,472	9,751	8,334
Buses	Diesel Urban Buses >18t	Euro III	2012	13813	14069	2172	15,947	12,779	11,217	377	302	265	1106	886	778	10,561	7,685	6,305
Buses	Diesel Urban Buses >18t	Euro IV	2012	18428	18769	2897	15,345	12,827	11,632	362	303	275	1064	890	807	7,106	5,505	4,635
Buses	Diesel Urban Buses >18t	Euro V	2012	15048	15326	2366	14,836	12,376	11,243	350	292	266	1029	858	780	5,523	2,620	1,912
Buses	Gasoline Coaches	Conventional	2012	1872	3581	1228	9,855	6,570	7,227	230	153	169	694	463	509	4,500	7,500	7,500
Buses	Diesel Coaches <15t	Conventional	2012	4254	8136	2789	13,080	9,616	8,499	309	227	201	907	667	590	11,324	8,822	8,156
Buses	Diesel Coaches <15t	Euro I	2012	2604	4981	1707	11,998	8,875	7,864	283	210	186	832	616	546	8,768	6,699	6,147
Buses	Diesel Coaches <15t	Euro II	2012	11020	21079	7226	11,934	8,902	7,898	282	210	187	828	618	548	10,033	7,549	6,840
Buses	Diesel Coaches <15t	Euro III	2012	10135	19386	6646	12,975	9,574	8,440	306	226	199	900	664	585	8,591	6,046	5,368
Buses	Diesel Coaches <15t	Euro IV	2012	7627	14589	5001	12,425	9,478	8,441	293	224	199	862	657	586	5,666	4,225	3,842
Buses	Diesel Coaches <15t	Euro V	2012	4004	7660	2626	12,141	9,185	8,144	287	217	192	842	637	565	7,922	3,947	2,471

Buses	Diesel Coaches 15 - 18t	Conventional	2012	2075	3969	1361	13,080	9,616	8,499	309	227	201	907	667	590	11,324	8,822	8,156
Buses	Diesel Coaches 15 - 18t	Euro I	2012	1170	2239	767	11,998	8,875	7,864	283	210	186	832	616	546	8,768	6,699	6,147
Buses	Diesel Coaches 15 - 18t	Euro II	2012	3042	5819	1995	11,934	8,902	7,898	282	210	187	828	618	548	10,033	7,549	6,840
Buses	Diesel Coaches 15 - 18t	Euro III	2012	1591	3043	1043	12,975	9,574	8,440	306	226	199	900	664	585	8,591	6,046	5,368
Buses	Diesel Coaches 15 - 18t	Euro IV	2012	1092	2089	716	12,425	9,478	8,441	293	224	199	862	657	586	5,666	4,225	3,842
Buses	Diesel Coaches 15 - 18t	Euro V	2012	1012	1935	663	12,141	9,185	8,144	287	217	192	842	637	565	7,922	3,947	2,471
Buses	Diesel Coaches >18t	Conventional	2012	298	569	195	15,881	11,649	10,271	375	275	243	1102	808	712	14,084	10,772	9,735
Buses	Diesel Coaches >18t	Euro I	2012	278	532	182	14,074	10,400	9,184	332	246	217	976	721	637	10,737	8,049	7,206
Buses	Diesel Coaches >18t	Euro II	2012	2343	4481	1536	13,832	10,315	9,121	327	244	215	959	716	633	11,883	8,817	7,837
Buses	Diesel Coaches >18t	Euro III	2012	7075	13534	4639	14,333	10,348	9,037	339	244	213	994	718	627	9,681	6,781	5,889
Buses	Diesel Coaches >18t	Euro IV	2012	5175	9899	3393	13,653	10,168	9,018	322	240	213	947	705	626	6,428	4,728	4,226
Buses	Diesel Coaches >18t	Euro V	2012	2853	5457	1871	13,421	9,918	8,750	317	234	207	931	688	607	8,476	4,163	2,596
Mopeds	2-stroke <50 cm³	Conventional	2012	40304	26869	0	1,095	1,095		26	26		77	77		0,056	0,056	
Mopeds	2-stroke <50 cm³	Euro I	2012	22486	14991	0	0,876	0,876		20	20		62	62		0,180	0,180	
Mopeds	2-stroke <50 cm³	Euro II	2012	53127	35418	0	0,876	0,876		20	20		62	62		0,170	0,170	
Mopeds	4-stroke <50 cm³	Euro II	2012	17709	11806	0	0,876	0,876		20	20		62	62		0,170	0,170	
Motorcycles	2-stroke >50 cm³	Conventional	2012	10671	9188	3477	1,188	1,240	1,736	28	29	41	84	87	122	0,029	0,030	0,035
Motorcycles	2-stroke >50 cm³	Euro I	2012	2038	1755	664	1,188	1,240	1,736	28	29	41	84	87	122	0,029	0,030	0,035
Motorcycles	2-stroke >50 cm³	Euro II	2012	2617	2254	853	1,090	1,122	1,552	25	26	36	77	79	109	0,040	0,050	0,060
Motorcycles	2-stroke >50 cm³	Euro III	2012	4228	3640	1378	1,090	1,122	1,552	25	26	36	77	79	109	0,048	0,058	0,069
Motorcycles	4-stroke <250 cm³	Conventional	2012	21578	18579	7032	1,086	1,204	1,579	25	28	37	76	85	111	0,237	0,428	0,655
Motorcycles	4-stroke <250 cm³	Euro I	2012	4121	3548	1343	1,183	1,331	1,766	28	31	41	83	94	124	0,304	0,424	0,567
Motorcycles	4-stroke <250 cm³	Euro II	2012	5293	4557	1725	0,975	1,102	1,479	23	26	35	69	78	104	0,323	0,447	0,598
Motorcycles	4-stroke <250 cm³	Euro III	2012	8549	7361	2786	0,844	0,892	1,136	20	21	27	59	63	80	0,253	0,382	0,612
Motorcycles	4-stroke 250 - 750 cm³	Conventional	2012	59341	51093	19337	1,167	1,041	1,166	27	24	27	82	73	82	0,196	0,300	0,548
Motorcycles	4-stroke 250 - 750 cm³	Euro I	2012	11332	9757	3693	1,637	1,554	1,813	38	36	42	115	109	128	0,258	0,400	0,610
Motorcycles	4-stroke 250 - 750 cm³	Euro II	2012	14555	12532	4743	1,498	1,465	1,808	35	34	42	105	103	127	0,257	0,390	0,577
Motorcycles	4-stroke 250 - 750 cm³	Euro III	2012	23511	20243	7661	1,357	1,345	1,670	32	31	39	96	95	118	0,076	0,132	0,265
Motorcycles	4-stroke >750 cm³	Conventional	2012	26973	23224	8790	1,565	1,557	1,916	37	36	45	110	110	135	0,019	0,030	0,086
Motorcycles	4-stroke >750 cm³	Euro I	2012	5151	4435	1679	1,888	1,798	2,081	44	42	49	133	127	147	0,125	0,178	0,392
Motorcycles	4-stroke >750 cm³	Euro II	2012	6616	5697	2156	1,844	1,665	1,835	43	39	43	130	117	129	0,143	0,244	0,459
Motorcycles	4-stroke >750 cm³	Euro III	2012	10687	9201	3482	1,767	1,641	1,887	41	38	44	124	116	133	0,104	0,200	0,484

Sector	Subsector	Tech 2	ForecastYear	CO_u	CO_r	CO_h	VOC_u	VOC_r	VOC_h	TSP_u	TSP_r	TSP_h	CH4_u	CH4_r	CH4_h	NMVOC_u	NMVOC_r	NMVOC_h	N2O_u	N2O_r	N2O_h
Passenger Cars	Gasoline 0,8 - 1,4 l	PRE ECE	2012	91,825	19,333	15,520	10,514	1,991	1,285	0,063	0,044	0,041	0,219	0,029	0,026	10,295	1,962	1,259	0,010	0,007	0,007
Passenger Cars	Gasoline 0,8 - 1,4 l	ECE 15/00-01	2012	63,319	14,480	18,620	9,073	1,634	1,158	0,063	0,044	0,041	0,219	0,029	0,026	8,854	1,605	1,132	0,010	0,007	0,007
Passenger Cars	Gasoline 0,8 - 1,4 l	ECE 15/02	2012	52,946	8,200	8,260	8,928	1,429	0,985	0,063	0,044	0,041	0,219	0,029	0,026	8,709	1,400	0,959	0,010	0,007	0,007
Passenger Cars	Gasoline 0,8 - 1,4 l	ECE 15/03	2012	55,927	8,793	7,620	8,852	1,423	0,985	0,042	0,029	0,029	0,219	0,029	0,026	8,633	1,394	0,959	0,010	0,007	0,007
Passenger Cars	Gasoline 0,8 - 1,4 l	ECE 15/04	2012	30,337	4,956	4,292	7,844	1,251	0,732	0,030	0,020	0,020	0,219	0,029	0,026	7,625	1,222	0,706	0,010	0,007	0,007
Passenger Cars	Gasoline 0,8 - 1,4 l	Euro I	2012	26,783	3,478	4,477	2,524	0,204	0,178	0,003	0,002	0,002	0,048	0,016	0,014	2,475	0,188	0,164	0,018	0,011	0,005
Passenger Cars	Gasoline 0,8 - 1,4 l	Euro II	2012	17,782	1,457	1,901	1,401	0,084	0,068	0,003	0,002	0,002	0,067	0,013	0,011	1,334	0,071	0,057	0,013	0,005	0,003
Passenger Cars	Gasoline 0,8 - 1,4 l	Euro III	2012	14,663	0,795	1,322	0,778	0,024	0,026	0,001	0,001	0,001	0,033	0,002	0,004	0,745	0,022	0,022	0,004	0,000	0,000
Passenger Cars	Gasoline 0,8 - 1,4 l	Euro IV	2012	4,263	0,301	0,555	0,474	0,021	0,018	0,001	0,001	0,001	0,014	0,003	0,005	0,460	0,019	0,013	0,003	0,000	0,000
Passenger Cars	Gasoline 0,8 - 1,4 l	Euro V	2012	4,222	0,281	0,518	0,492	0,023	0,018	0,001	0,001	0,001	0,014	0,003	0,005	0,478	0,020	0,013	0,003	0,000	0,000
Passenger Cars	Gasoline 1,4 - 2,0 l	PRE ECE	2012	91,825	19,333	15,520	10,331	1,975	1,283	0,063	0,044	0,041	0,219	0,029	0,026	10,112	1,946	1,257	0,010	0,007	0,007
Passenger Cars	Gasoline 1,4 - 2,0 l	ECE 15/00-01	2012	63,319	14,480	18,620	8,932	1,622	1,157	0,063	0,044	0,041	0,219	0,029	0,026	8,713	1,593	1,131	0,010	0,007	0,007
Passenger Cars	Gasoline 1,4 - 2,0 l	ECE 15/02	2012	52,946	8,200	8,260	8,804	1,418	0,984	0,063	0,044	0,041	0,219	0,029	0,026	8,585	1,389	0,958	0,010	0,007	0,007
Passenger Cars	Gasoline 1,4 - 2,0 l	ECE 15/03	2012	55,927	8,793	7,620	8,746	1,413	0,984	0,042	0,029	0,029	0,219	0,029	0,026	8,527	1,384	0,958	0,010	0,007	0,007
Passenger Cars	Gasoline 1,4 - 2,0 l	ECE 15/04	2012	30,337	4,956	4,292	7,747	1,243	0,731	0,030	0,020	0,020	0,219	0,029	0,026	7,528	1,213	0,705	0,010	0,007	0,007
Passenger Cars	Gasoline 1,4 - 2,0 l	Euro I	2012	19,478	2,427	3,124	2,797	0,225	0,199	0,003	0,002	0,002	0,048	0,016	0,014	2,749	0,209	0,185	0,018	0,011	0,005
Passenger Cars	Gasoline 1,4 - 2,0 l	Euro II	2012	12,894	1,017	1,326	1,548	0,092	0,076	0,003	0,002	0,002	0,067	0,013	0,011	1,481	0,079	0,065	0,013	0,005	0,003
Passenger Cars	Gasoline 1,4 - 2,0 l	Euro III	2012	10,670	0,691	1,148	0,854	0,022	0,026	0,001	0,001	0,001	0,033	0,002	0,004	0,821	0,020	0,022	0,004	0,000	0,000
Passenger Cars	Gasoline 1,4 - 2,0 l	Euro IV	2012	3,134	0,287	0,529	0,514	0,020	0,018	0,001	0,001	0,001	0,014	0,003	0,005	0,500	0,018	0,013	0,003	0,000	0,000
Passenger Cars	Gasoline 1,4 - 2,0 l	Euro V	2012	3,115	0,287	0,529	0,521	0,021	0,018	0,001	0,001	0,001	0,014	0,003	0,005	0,507	0,018	0,013	0,003	0,000	0,000
Passenger Cars	Gasoline >2,0 l	PRE ECE	2012	91,825	19,333	15,520	10,241	1,967	1,283	0,063	0,044	0,041	0,219	0,029	0,026	10,022	1,938	1,257	0,010	0,007	0,007
Passenger Cars	Gasoline >2,0 l	ECE 15/00-01	2012	63,319	14,480	18,620	8,862	1,616	1,156	0,063	0,044	0,041	0,219	0,029	0,026	8,643	1,587	1,130	0,010	0,007	0,007
Passenger Cars	Gasoline >2,0 l	ECE 15/02	2012	52,946	8,200	8,260	8,742	1,413	0,984	0,063	0,044	0,041	0,219	0,029	0,026	8,523	1,384	0,958	0,010	0,007	0,007
Passenger Cars	Gasoline >2,0 l	ECE 15/03	2012	55,927	8,793	7,620	8,699	1,409	0,984	0,042	0,029	0,029	0,219	0,029	0,026	8,480	1,380	0,958	0,010	0,007	0,007
Passenger Cars	Gasoline >2,0 l	ECE 15/04	2012	30,337	4,956	4,292	7,697	1,238	0,731	0,030	0,020	0,020	0,219	0,029	0,026	7,478	1,209	0,705	0,010	0,007	0,007
Passenger Cars	Gasoline >2,0 l	Euro I	2012	13,524	1,642	2,114	1,928	0,184	0,162	0,003	0,002	0,002	0,048	0,016	0,014	1,880	0,168	0,148	0,018	0,011	0,005
Passenger Cars	Gasoline >2,0 l	Euro II	2012	8,853	0,688	0,897	1,064	0,076	0,062	0,003	0,002	0,002	0,067	0,013	0,011	0,997	0,063	0,051	0,013	0,005	0,003
Passenger Cars	Gasoline >2,0 l	Euro III	2012	7,422	0,691	1,148	0,584	0,022	0,026	0,001	0,001	0,001	0,033	0,002	0,004	0,551	0,020	0,022	0,004	0,000	0,000
Passenger Cars	Gasoline >2,0 l	Euro IV	2012	2,195	0,287	0,529	0,360	0,020	0,018	0,001	0,001	0,001	0,014	0,003	0,005	0,345	0,017	0,013	0,003	0,000	0,000
Passenger Cars	Gasoline >2,0 l	Euro V	2012	2,173	0,287	0,529	0,364	0,020	0,018	0,001	0,001	0,001	0,014	0,003	0,005	0,349	0,018	0,013	0,003	0,000	0,000
Passenger Cars	Diesel 1,4 - 2,0 l	Euro I	2012	0,746	0,215	0,208	0,140	0,031	0,026	0,145	0,062	0,107	0,019	0,009	0,003	0,120	0,022	0,023	0,002	0,004	0,004
Passenger Cars	Diesel 1,4 - 2,0 l	Euro II	2012	0,611	0,110	0,035	0,091	0,021	0,015	0,119	0,039	0,050	0,006	0,003	0,002	0,085	0,018	0,013	0,003	0,006	0,006
Passenger Cars	Diesel 1,4 - 2,0 l	Euro III	2012	0,176	0,041	0,012	0,048	0,011	0,009	0,075	0,030	0,045	0,003	0,000	0,000	0,045	0,011	0,009	0,016	0,004	0,004
Passenger Cars	Diesel 1,4 - 2,0 l	Euro IV	2012	0,148	0,034	0,021	0,029	0,006	0,006	0,074	0,024	0,026	0,000	0,000	0,000	0,029	0,006	0,006	0,016	0,004	0,004
Passenger Cars	Diesel 1,4 - 2,0 l	Euro V	2012	0,148	0,034	0,021	0,029	0,006	0,006	0,015	0,005	0,005	0,000	0,000	0,000	0,029	0,006	0,006	0,016	0,004	0,004
Passenger Cars	Diesel 1,4 - 2,0 l	Conventional	2012	1,158	0,472	0,384	0,384	0,086	0,062	0,507	0,132	0,170	0,021	0,012	0,008	0,363	0,074	0,054	0,000	0,000	0,000

Passenger Cars	Diesel >2,0 l	Euro I	2012	0,746	0,215	0,208	0,211	0,046	0,034	0,145	0,062	0,107	0,019	0,009	0,003	0,192	0,037	0,031	0,002	0,004	0,004
Passenger Cars	Diesel >2,0 l	Euro II	2012	0,611	0,110	0,035	0,258	0,058	0,038	0,119	0,039	0,050	0,006	0,003	0,002	0,252	0,055	0,036	0,003	0,006	0,006
Passenger Cars	Diesel >2,0 l	Euro III	2012	0,176	0,041	0,012	0,101	0,017	0,012	0,075	0,030	0,045	0,003	0,000	0,000	0,098	0,017	0,012	0,016	0,004	0,004
Passenger Cars	Diesel >2,0 l	Euro IV	2012	0,148	0,034	0,021	0,029	0,006	0,006	0,074	0,024	0,026	0,000	0,000	0,000	0,029	0,006	0,006	0,016	0,004	0,004
Passenger Cars	Diesel >2,0 l	Euro V	2012	0,148	0,034	0,021	0,029	0,006	0,006	0,015	0,005	0,005	0,000	0,000	0,000	0,029	0,006	0,006	0,016	0,004	0,004
Passenger Cars	Diesel >2,0 l	Conventional	2012	1,158	0,472	0,384	0,384	0,086	0,062	0,507	0,132	0,170	0,021	0,012	0,008	0,363	0,074	0,054	0,000	0,000	0,000
Passenger Cars	LPG cars	Euro I	2012	4,312	1,445	3,560	0,456	0,071	0,083	0,040	0,030	0,025	0,080	0,035	0,025	0,376	0,036	0,058	0,041	0,013	0,008
Passenger Cars	LPG cars	Euro II	2012	3,052	0,982	2,421	0,172	0,015	0,017	0,040	0,030	0,025	0,019	0,008	0,006	0,152	0,007	0,011	0,021	0,003	0,002
Passenger Cars	LPG cars	Euro III	2012	2,595	0,809	1,993	0,105	0,011	0,012	0,040	0,030	0,025	0,013	0,006	0,004	0,092	0,005	0,008	0,006	0,002	0,001
Passenger Cars	LPG cars	Euro IV	2012	0,986	0,491	1,210	0,046	0,002	0,002	0,040	0,030	0,025	0,004	0,002	0,001	0,042	0,000	0,001	0,006	0,002	0,001
Passenger Cars	LPG cars	Conventional	2012	6,725	2,373	9,723	2,063	0,667	0,490	0,040	0,030	0,025	0,080	0,035	0,025	1,983	0,632	0,465	0,000	0,000	0,000
Passenger Cars	Gasoline <0,8 l	PRE ECE	2012	91,825	19,333	15,520	10,501	1,990	1,285	0,063	0,044	0,041	0,219	0,029	0,026	10,283	1,961	1,259	0,010	0,007	0,007
Passenger Cars	Gasoline <0,8 l	ECE 15/00-01	2012	63,319	14,480	18,620	9,099	1,636	1,158	0,063	0,044	0,041	0,219	0,029	0,026	8,880	1,607	1,132	0,010	0,007	0,007
Passenger Cars	Gasoline <0,8 l	ECE 15/02	2012	52,946	8,200	8,260	8,931	1,430	0,985	0,063	0,044	0,041	0,219	0,029	0,026	8,713	1,400	0,959	0,010	0,007	0,007
Passenger Cars	Gasoline <0,8 l	ECE 15/03	2012	55,927	8,793	7,620	8,875	1,425	0,985	0,042	0,029	0,029	0,219	0,029	0,026	8,656	1,395	0,959	0,010	0,007	0,007
Passenger Cars	Gasoline <0,8 l	ECE 15/04	2012	30,337	4,956	4,292	7,853	1,252	0,732	0,030	0,020	0,020	0,219	0,029	0,026	7,634	1,223	0,706	0,010	0,007	0,007
Passenger Cars	Gasoline <0,8 l	Euro I	2012	25,783	2,659	3,422	2,465	0,176	0,152	0,003	0,002	0,002	0,048	0,016	0,014	2,416	0,160	0,138	0,016	0,009	0,004
Passenger Cars	Gasoline <0,8 l	Euro II	2012	17,719	1,399	1,825	1,396	0,082	0,066	0,003	0,002	0,002	0,067	0,013	0,011	1,329	0,069	0,055	0,012	0,005	0,003
Passenger Cars	Gasoline <0,8 l	Euro III	2012	14,660	0,794	1,320	0,778	0,024	0,026	0,001	0,001	0,001	0,033	0,002	0,004	0,745	0,022	0,022	0,004	0,000	0,000
Passenger Cars	Gasoline <0,8 l	Euro IV	2012	4,266	0,303	0,558	0,474	0,021	0,018	0,001	0,001	0,001	0,014	0,003	0,005	0,460	0,019	0,013	0,003	0,000	0,000
Passenger Cars	Gasoline <0,8 l	Euro V	2012	4,223	0,282	0,519	0,487	0,023	0,018	0,001	0,001	0,001	0,014	0,003	0,005	0,473	0,020	0,013	0,003	0,000	0,000
Passenger Cars	Diesel <1,4 l	Euro I	2012	0,746	0,215	0,208	0,140	0,031	0,026	0,057	0,062	0,107	0,019	0,009	0,003	0,120	0,022	0,023	0,002	0,004	0,004
Passenger Cars	Diesel <1,4 l	Euro II	2012	0,611	0,110	0,035	0,091	0,021	0,015	0,047	0,039	0,050	0,006	0,003	0,002	0,085	0,018	0,013	0,003	0,006	0,006
Passenger Cars	Diesel <1,4 l	Euro III	2012	0,176	0,041	0,012	0,048	0,011	0,009	0,029	0,030	0,045	0,003	0,000	0,000	0,045	0,011	0,009	0,016	0,004	0,004
Passenger Cars	Diesel <1,4 l	Euro IV	2012	0,148	0,034	0,021	0,029	0,006	0,006	0,029	0,024	0,026	0,000	0,000	0,000	0,029	0,006	0,006	0,016	0,004	0,004
Passenger Cars	Diesel <1,4 l	Euro V	2012	0,148	0,034	0,021	0,029	0,006	0,006	0,006	0,005	0,005	0,000	0,000	0,000	0,029	0,006	0,006	0,016	0,004	0,004
Passenger Cars	Diesel <1,4 l	Conventional	2012	1,158	0,472	0,384	0,384	0,086	0,062	0,199	0,132	0,170	0,021	0,012	0,008	0,363	0,074	0,054	0,000	0,000	0,000
Light Duty Vehicles	Gasoline <3,5t	Conventional	2012	48,873	6,075	7,389	8,118	1,030	0,477	0,040	0,040	0,040	0,208	0,040	0,025	7,910	0,990	0,452	0,010	0,007	0,007
Light Duty Vehicles	Gasoline <3,5t	Euro I	2012	35,603	2,186	2,756	2,364	0,184	0,125	0,003	0,002	0,002	0,048	0,016	0,014	2,316	0,168	0,111	0,047	0,027	0,013
Light Duty Vehicles	Gasoline <3,5t	Euro II	2012	24,502	1,333	1,681	1,229	0,050	0,031	0,003	0,002	0,002	0,066	0,013	0,011	1,163	0,037	0,020	0,074	0,016	0,010
Light Duty Vehicles	Gasoline <3,5t	Euro III	2012	18,238	0,448	0,565	0,706	0,023	0,012	0,001	0,001	0,001	0,032	0,002	0,004	0,674	0,021	0,008	0,011	0,001	0,001
Light Duty Vehicles	Gasoline <3,5t	Euro IV	2012	5,834	0,241	0,304	0,422	0,013	0,005	0,001	0,001	0,001	0,013	0,002	0,000	0,409	0,011	0,005	0,004	0,000	0,000
Light Duty Vehicles	Gasoline <3,5t	Euro V	2012	5,717	0,241	0,304	0,462	0,017	0,006	0,001	0,001	0,001	0,013	0,002	0,000	0,449	0,015	0,006	0,003	0,000	0,000
Light Duty Vehicles	Diesel <3,5t	Conventional	2012	1,976	1,009	1,060	0,339	0,106	0,101	0,715	0,303	0,322	0,021	0,012	0,008	0,318	0,094	0,093	0,000	0,000	0,000
Light Duty Vehicles	Diesel <3,5t	Euro I	2012	0,691	0,328	0,423	0,339	0,106	0,101	0,175	0,066	0,090	0,019	0,009	0,003	0,320	0,097	0,098	0,002	0,004	0,004
Light Duty Vehicles	Diesel <3,5t	Euro II	2012	0,691	0,328	0,423	0,339	0,106	0,101	0,175	0,066	0,090	0,006	0,003	0,002	0,333	0,103	0,099	0,003	0,006	0,006
Light Duty Vehicles	Diesel <3,5t	Euro III	2012	0,566	0,269	0,347	0,210	0,065	0,063	0,117	0,044	0,061	0,003	0,000	0,000	0,207	0,065	0,063	0,016	0,004	0,004

Light Duty Vehicles	Diesel <3,5t	Euro IV	2012	0,449	0,213	0,275	0,078	0,024	0,023	0,061	0,023	0,032	0,000	0,000	0,000	0,078	0,024	0,023	0,016	0,004	0,004
Light Duty Vehicles	Diesel <3,5t	Euro V	2012	0,449	0,213	0,275	0,078	0,024	0,023	0,003	0,001	0,002	0,000	0,000	0,000	0,078	0,024	0,023	0,016	0,004	0,004
Light Duty Vehicles	LPG <3,5t	Conventional	2012	9,896	3,559	14,584	3,055	1,000	0,735	0,060	0,045	0,038	0,120	0,053	0,038	2,935	0,948	0,697	0,000	0,000	0,000
Light Duty Vehicles	LPG <3,5t	Euro II	2012	4,052	1,474	3,631	0,177	0,022	0,026	0,060	0,045	0,038	0,029	0,013	0,009	0,148	0,010	0,017	0,025	0,005	0,003
Light Duty Vehicles	LPG <3,5t	Euro III	2012	1,889	1,214	2,990	0,111	0,016	0,019	0,060	0,045	0,038	0,019	0,008	0,006	0,092	0,008	0,013	0,009	0,003	0,002
Light Duty Vehicles	LPG <3,5t	Euro IV	2012	1,456	0,737	1,815	0,068	0,003	0,004	0,060	0,045	0,038	0,006	0,003	0,002	0,062	0,001	0,002	0,009	0,003	0,002
Heavy Duty Vehicles	Gasoline >3,5t	Conventional	2012	70,000	55,000	55,000	7,000	5,500	3,500	0,400	0,400	0,400	0,140	0,110	0,070	6,860	5,390	3,430	0,006	0,006	0,006
Heavy Duty Vehicles	Diesel RT 3,5 - 7,5t	Conventional	2012	2,060	1,509	1,351	1,298	0,789	0,576	0,321	0,240	0,216	0,085	0,023	0,020	1,213	0,766	0,556	0,030	0,030	0,030
Heavy Duty Vehicles	Diesel RT 3,5 - 7,5t	Euro I	2012	0,668	0,501	0,546	0,253	0,167	0,130	0,126	0,095	0,090	0,085	0,023	0,020	0,168	0,144	0,110	0,030	0,030	0,030
Heavy Duty Vehicles	Diesel RT 3,5 - 7,5t	Euro II	2012	0,534	0,466	0,461	0,171	0,111	0,086	0,053	0,048	0,055	0,054	0,020	0,019	0,116	0,091	0,067	0,030	0,030	0,030
Heavy Duty Vehicles	Diesel RT 3,5 - 7,5t	Euro III	2012	0,660	0,481	0,452	0,162	0,102	0,077	0,062	0,045	0,039	0,048	0,021	0,018	0,114	0,081	0,058	0,030	0,030	0,030
Heavy Duty Vehicles	Diesel RT 3,5 - 7,5t	Euro IV	2012	0,342	0,270	0,258	0,022	0,017	0,017	0,015	0,013	0,014	0,003	0,002	0,001	0,020	0,016	0,015	0,030	0,030	0,030
Heavy Duty Vehicles	Diesel RT 3,5 - 7,5t	Euro V	2012	0,642	0,591	0,589	0,011	0,008	0,007	0,014	0,010	0,009	0,003	0,002	0,001	0,009	0,007	0,006	0,030	0,030	0,030
Heavy Duty Vehicles	Diesel RT 3,5 - 7,5t	Euro VI	2012	0,355	0,276	0,260	0,011	0,008	0,007	0,001	0,001	0,001	0,001	0,000	0,000	0,010	0,008	0,007	0,030	0,030	0,030
Heavy Duty Vehicles	Diesel RT 7,5 - 12t	Conventional	2012	2,358	1,698	1,525	0,957	0,589	0,449	0,330	0,236	0,207	0,085	0,023	0,020	0,872	0,566	0,429	0,030	0,030	0,030
Heavy Duty Vehicles	Diesel RT 7,5 - 12t	Euro I	2012	1,086	0,817	0,766	0,389	0,258	0,208	0,201	0,144	0,131	0,085	0,023	0,020	0,304	0,235	0,188	0,030	0,030	0,030
Heavy Duty Vehicles	Diesel RT 7,5 - 12t	Euro II	2012	0,868	0,727	0,717	0,263	0,172	0,137	0,085	0,072	0,083	0,054	0,020	0,019	0,208	0,152	0,118	0,030	0,030	0,030
Heavy Duty Vehicles	Diesel RT 7,5 - 12t	Euro III	2012	1,084	0,771	0,733	0,252	0,157	0,120	0,096	0,067	0,058	0,048	0,021	0,018	0,204	0,136	0,102	0,030	0,030	0,030
Heavy Duty Vehicles	Diesel RT 7,5 - 12t	Euro IV	2012	0,553	0,418	0,369	0,035	0,025	0,022	0,023	0,019	0,019	0,003	0,002	0,001	0,032	0,024	0,021	0,030	0,030	0,030
Heavy Duty Vehicles	Diesel RT 7,5 - 12t	Euro V	2012	1,003	0,820	0,772	0,019	0,014	0,012	0,024	0,017	0,015	0,003	0,002	0,001	0,017	0,012	0,011	0,030	0,030	0,030
Heavy Duty Vehicles	Diesel RT 7,5 - 12t	Euro VI	2012	0,583	0,431	0,373	0,017	0,012	0,010	0,002	0,002	0,002	0,001	0,000	0,000	0,016	0,012	0,010	0,030	0,030	0,030
Heavy Duty Vehicles	Diesel RT 12 - 14 t	Conventional	2012	2,546	1,876	1,693	1,012	0,646	0,509	0,351	0,254	0,233	0,085	0,023	0,020	0,927	0,623	0,489	0,030	0,030	0,030
Heavy Duty Vehicles	Diesel RT 12 - 14 t	Euro I	2012	1,200	0,918	0,866	0,429	0,279	0,229	0,218	0,159	0,147	0,085	0,023	0,020	0,344	0,256	0,209	0,030	0,030	0,030
Heavy Duty Vehicles	Diesel RT 12 - 14 t	Euro II	2012	0,985	0,820	0,804	0,281	0,186	0,150	0,101	0,086	0,101	0,054	0,020	0,019	0,227	0,166	0,131	0,030	0,030	0,030
Heavy Duty Vehicles	Diesel RT 12 - 14 t	Euro III	2012	1,176	0,873	0,835	0,260	0,168	0,134	0,104	0,075	0,068	0,048	0,021	0,018	0,212	0,147	0,116	0,030	0,030	0,030
Heavy Duty Vehicles	Diesel RT 12 - 14 t	Euro IV	2012	0,599	0,448	0,410	0,034	0,025	0,024	0,024	0,020	0,020	0,003	0,002	0,001	0,031	0,024	0,023	0,030	0,030	0,030
Heavy Duty Vehicles	Diesel RT 12 - 14 t	Euro V	2012	1,137	0,900	0,821	0,019	0,014	0,013	0,024	0,017	0,016	0,003	0,002	0,001	0,017	0,013	0,012	0,030	0,030	0,030
Heavy Duty Vehicles	Diesel RT 14 - 20t	Conventional	2012	3,512	2,514	2,221	1,510	0,971	0,768	0,483	0,341	0,298	0,175	0,080	0,070	1,335	0,891	0,698	0,030	0,030	0,030
Heavy Duty Vehicles	Diesel RT 14 - 20t	Euro I	2012	1,612	1,206	1,117	0,606	0,403	0,325	0,298	0,209	0,181	0,175	0,080	0,070	0,431	0,323	0,255	0,030	0,030	0,030
Heavy Duty Vehicles	Diesel RT 14 - 20t	Euro II	2012	1,267	1,025	1,002	0,409	0,267	0,213	0,117	0,095	0,111	0,112	0,070	0,065	0,297	0,197	0,148	0,030	0,030	0,030
Heavy Duty Vehicles	Diesel RT 14 - 20t	Euro III	2012	1,601	1,150	1,096	0,378	0,243	0,196	0,141	0,097	0,083	0,098	0,074	0,064	0,280	0,168	0,133	0,030	0,030	0,030

Heavy Duty Vehicles	Diesel RT 14 - 20t	Euro IV	2012	0,829	0,602	0,523	0,046	0,032	0,028	0,031	0,024	0,023	0,005	0,006	0,004	0,041	0,026	0,024	0,030	0,030	0,030
Heavy Duty Vehicles	Diesel RT 14 - 20t	Euro V	2012	1,365	1,029	0,901	0,031	0,022	0,019	0,032	0,022	0,019	0,005	0,006	0,004	0,026	0,016	0,015	0,030	0,030	0,030
Heavy Duty Vehicles	Diesel RT 14 - 20t	Euro VI	2012	0,870	0,625	0,536	0,024	0,017	0,015	0,003	0,002	0,002	0,002	0,002	0,001	0,022	0,015	0,014	0,030	0,030	0,030
Heavy Duty Vehicles	Diesel RT 20 - 26t	Conventional	2012	2,558	1,885	1,712	0,819	0,517	0,406	0,482	0,353	0,319	0,175	0,080	0,070	0,644	0,437	0,336	0,030	0,030	0,030
Heavy Duty Vehicles	Diesel RT 20 - 26t	Euro I	2012	2,068	1,563	1,437	0,728	0,476	0,380	0,383	0,264	0,231	0,175	0,080	0,070	0,553	0,396	0,310	0,030	0,030	0,030
Heavy Duty Vehicles	Diesel RT 20 - 26t	Euro II	2012	1,620	1,285	1,399	0,489	0,314	0,248	0,160	0,128	0,146	0,112	0,070	0,065	0,377	0,244	0,183	0,030	0,030	0,030
Heavy Duty Vehicles	Diesel RT 20 - 26t	Euro III	2012	2,025	1,487	1,403	0,453	0,287	0,225	0,171	0,117	0,102	0,098	0,074	0,064	0,355	0,212	0,161	0,030	0,030	0,030
Heavy Duty Vehicles	Diesel RT 20 - 26t	Euro IV	2012	1,003	0,728	0,628	0,059	0,040	0,035	0,041	0,031	0,028	0,005	0,006	0,004	0,053	0,035	0,030	0,030	0,030	0,030
Heavy Duty Vehicles	Diesel RT 20 - 26t	Euro V	2012	1,787	1,376	1,225	0,034	0,024	0,020	0,041	0,028	0,024	0,005	0,006	0,004	0,029	0,018	0,016	0,030	0,030	0,030
Heavy Duty Vehicles	Diesel RT 20 - 26t	Euro VI	2012	1,045	0,753	0,646	0,030	0,021	0,017	0,004	0,003	0,003	0,002	0,002	0,001	0,028	0,019	0,016	0,030	0,030	0,030
Heavy Duty Vehicles	Diesel RT 26 - 28t	Conventional	2012	2,703	1,987	1,810	0,842	0,541	0,430	0,512	0,375	0,336	0,175	0,080	0,070	0,667	0,461	0,360	0,030	0,030	0,030
Heavy Duty Vehicles	Diesel RT 26 - 28t	Euro I	2012	2,162	1,647	1,535	0,736	0,488	0,394	0,398	0,281	0,244	0,175	0,080	0,070	0,561	0,408	0,324	0,030	0,030	0,030
Heavy Duty Vehicles	Diesel RT 26 - 28t	Euro II	2012	1,682	1,346	1,457	0,499	0,327	0,262	0,185	0,148	0,167	0,112	0,070	0,065	0,387	0,257	0,197	0,030	0,030	0,030
Heavy Duty Vehicles	Diesel RT 26 - 28t	Euro III	2012	2,121	1,582	1,481	0,467	0,304	0,243	0,201	0,141	0,118	0,098	0,074	0,064	0,369	0,229	0,179	0,030	0,030	0,030
Heavy Duty Vehicles	Diesel RT 26 - 28t	Euro IV	2012	1,044	0,752	0,640	0,064	0,045	0,037	0,044	0,033	0,029	0,005	0,006	0,004	0,059	0,039	0,033	0,030	0,030	0,030
Heavy Duty Vehicles	Diesel RT 26 - 28t	Euro V	2012	2,008	1,531	1,359	0,031	0,022	0,018	0,046	0,030	0,025	0,005	0,006	0,004	0,025	0,016	0,014	0,030	0,030	0,030
Heavy Duty Vehicles	Diesel RT 28 - 32t	Conventional	2012	2,928	2,149	2,047	0,874	0,560	0,444	0,567	0,415	0,376	0,175	0,080	0,070	0,699	0,480	0,374	0,030	0,030	0,030
Heavy Duty Vehicles	Diesel RT 28 - 32t	Euro I	2012	2,377	1,862	1,795	0,778	0,518	0,419	0,436	0,314	0,281	0,175	0,080	0,070	0,603	0,438	0,349	0,030	0,030	0,030
Heavy Duty Vehicles	Diesel RT 28 - 32t	Euro II	2012	1,930	1,574	1,563	0,523	0,344	0,276	0,182	0,148	0,168	0,112	0,070	0,065	0,411	0,275	0,211	0,030	0,030	0,030
Heavy Duty Vehicles	Diesel RT 28 - 32t	Euro III	2012	2,325	1,732	1,685	0,491	0,317	0,252	0,189	0,135	0,119	0,098	0,074	0,064	0,393	0,243	0,188	0,030	0,030	0,030
Heavy Duty Vehicles	Diesel RT 28 - 32t	Euro IV	2012	1,145	0,834	0,714	0,070	0,051	0,043	0,049	0,038	0,034	0,005	0,006	0,004	0,065	0,045	0,039	0,030	0,030	0,030
Heavy Duty Vehicles	Diesel RT 28 - 32t	Euro V	2012	2,312	1,748	1,527	0,033	0,024	0,020	0,052	0,035	0,029	0,005	0,006	0,004	0,028	0,018	0,016	0,030	0,030	0,030
Heavy Duty Vehicles	Diesel RT 28 - 32t	Euro VI	2012	1,191	0,863	0,736	0,034	0,024	0,020	0,005	0,004	0,003	0,002	0,002	0,001	0,033	0,023	0,019	0,030	0,030	0,030
Heavy Duty Vehicles	Diesel RT >32t	Euro I	2012	2,482	1,894	1,795	0,812	0,527	0,419	0,453	0,317	0,286	0,175	0,080	0,070	0,637	0,447	0,349	0,030	0,030	0,030
Heavy Duty Vehicles	Diesel RT >32t	Euro III	2012	2,398	1,789	1,725	0,496	0,316	0,249	0,219	0,153	0,135	0,098	0,074	0,064	0,398	0,241	0,185	0,030	0,030	0,030
Heavy Duty Vehicles	Diesel RT >32t	Euro IV	2012	1,151	0,839	0,723	0,070	0,048	0,041	0,049	0,037	0,033	0,005	0,006	0,004	0,064	0,043	0,037	0,030	0,030	0,030
Heavy Duty Vehicles	Diesel RT >32t	Euro V	2012	2,346	1,790	1,602	0,034	0,024	0,020	0,050	0,034	0,030	0,005	0,006	0,004	0,029	0,018	0,015	0,030	0,030	0,030
Heavy Duty Vehicles	Diesel TT/AT 28 - 34t	Conventional	2012	2,560	1,899	1,804	0,736	0,476	0,380	0,488	0,361	0,339	0,175	0,080	0,070	0,561	0,396	0,310	0,030	0,030	0,030
Heavy Duty Vehicles	Diesel TT/AT 28 - 34t	Euro I	2012	2,173	1,665	1,602	0,678	0,450	0,363	0,380	0,277	0,261	0,175	0,080	0,070	0,503	0,370	0,293	0,030	0,030	0,030

Heavy Duty Vehicles	Diesel TT/AT 28 - 34t	Euro II	2012	1,746	1,372	1,500	0,450	0,296	0,238	0,174	0,138	0,158	0,112	0,070	0,065	0,338	0,226	0,173	0,030	0,030	0,030
Heavy Duty Vehicles	Diesel TT/AT 28 - 34t	Euro III	2012	2,067	1,559	1,515	0,415	0,269	0,215	0,169	0,121	0,111	0,098	0,074	0,064	0,317	0,195	0,151	0,030	0,030	0,030
Heavy Duty Vehicles	Diesel TT/AT 28 - 34t	Euro IV	2012	0,990	0,709	0,618	0,059	0,041	0,036	0,043	0,031	0,028	0,005	0,006	0,004	0,054	0,036	0,032	0,030	0,030	0,030
Heavy Duty Vehicles	Diesel TT/AT 28 - 34t	Euro V	2012	1,844	1,384	1,147	0,034	0,024	0,020	0,044	0,030	0,026	0,005	0,006	0,004	0,029	0,018	0,016	0,030	0,030	0,030
Heavy Duty Vehicles	Diesel TT/AT 28 - 34t	Euro VI	2012	1,033	0,738	0,629	0,029	0,021	0,017	0,004	0,003	0,003	0,002	0,002	0,001	0,028	0,019	0,016	0,030	0,030	0,030
Heavy Duty Vehicles	Diesel TT/AT 34 - 40t	Conventional	2012	3,006	2,216	2,091	0,877	0,555	0,438	0,579	0,419	0,384	0,175	0,080	0,070	0,702	0,475	0,368	0,030	0,030	0,030
Heavy Duty Vehicles	Diesel TT/AT 34 - 40t	Euro I	2012	2,561	1,946	1,861	0,805	0,524	0,420	0,464	0,324	0,293	0,175	0,080	0,070	0,630	0,444	0,350	0,030	0,030	0,030
Heavy Duty Vehicles	Diesel TT/AT 34 - 40t	Euro II	2012	2,056	1,607	1,775	0,538	0,343	0,270	0,211	0,165	0,188	0,112	0,070	0,065	0,426	0,274	0,205	0,030	0,030	0,030
Heavy Duty Vehicles	Diesel TT/AT 34 - 40t	Euro III	2012	2,453	1,826	1,775	0,494	0,312	0,244	0,200	0,139	0,122	0,098	0,074	0,064	0,396	0,237	0,180	0,030	0,030	0,030
Heavy Duty Vehicles	Diesel TT/AT 34 - 40t	Euro IV	2012	1,157	0,830	0,704	0,071	0,048	0,041	0,050	0,036	0,032	0,005	0,006	0,004	0,066	0,043	0,037	0,030	0,030	0,030
Heavy Duty Vehicles	Diesel TT/AT 34 - 40t	Euro V	2012	2,113	1,626	1,379	0,041	0,028	0,023	0,052	0,035	0,030	0,005	0,006	0,004	0,035	0,022	0,019	0,030	0,030	0,030
Heavy Duty Vehicles	Diesel TT/AT 34 - 40t	Euro VI	2012	1,202	0,857	0,727	0,035	0,024	0,020	0,005	0,004	0,003	0,002	0,002	0,001	0,033	0,022	0,019	0,030	0,030	0,030
Heavy Duty Vehicles	Diesel TT/AT 40 - 50t	Conventional	2012	3,242	2,400	2,283	0,901	0,570	0,450	0,622	0,462	0,425	0,175	0,080	0,070	0,726	0,490	0,380	0,030	0,030	0,030
Heavy Duty Vehicles	Diesel TT/AT 40 - 50t	Euro I	2012	2,823	2,135	2,079	0,844	0,546	0,433	0,500	0,358	0,333	0,175	0,080	0,070	0,669	0,466	0,363	0,030	0,030	0,030
Heavy Duty Vehicles	Diesel TT/AT 40 - 50t	Euro II	2012	2,313	1,826	1,823	0,558	0,358	0,282	0,225	0,176	0,199	0,112	0,070	0,065	0,446	0,288	0,217	0,030	0,030	0,030
Heavy Duty Vehicles	Diesel TT/AT 40 - 50t	Euro III	2012	2,675	1,999	1,959	0,510	0,323	0,253	0,208	0,147	0,127	0,098	0,074	0,064	0,412	0,248	0,189	0,030	0,030	0,030
Heavy Duty Vehicles	Diesel TT/AT 40 - 50t	Euro IV	2012	1,241	0,894	0,759	0,077	0,053	0,045	0,054	0,040	0,035	0,005	0,006	0,004	0,072	0,047	0,041	0,030	0,030	0,030
Heavy Duty Vehicles	Diesel TT/AT 40 - 50t	Euro V	2012	2,525	1,942	1,648	0,037	0,026	0,021	0,058	0,039	0,033	0,005	0,006	0,004	0,032	0,020	0,017	0,030	0,030	0,030
Heavy Duty Vehicles	Diesel TT/AT 40 - 50t	Euro VI	2012	1,288	0,921	0,779	0,038	0,026	0,022	0,006	0,004	0,004	0,002	0,002	0,001	0,036	0,024	0,020	0,030	0,030	0,030
Heavy Duty Vehicles	Diesel TT/AT 50 - 60t	Euro II	2012	2,783	2,192	2,191	0,626	0,406	0,323	0,317	0,246	0,275	0,112	0,070	0,065	0,514	0,337	0,258	0,030	0,030	0,030
Heavy Duty Vehicles	Diesel TT/AT 50 - 60t	Euro IV	2012	1,445	1,038	0,878	0,090	0,063	0,053	0,064	0,047	0,041	0,005	0,006	0,004	0,085	0,057	0,048	0,030	0,030	0,030
Heavy Duty Vehicles	Diesel TT/AT >60t	Euro III	2012	3,169	2,369	2,341	0,568	0,364	0,288	0,278	0,194	0,184	0,098	0,074	0,064	0,470	0,290	0,224	0,030	0,030	0,030
Buses	Gasoline Urban Buses	Conventional	2012	70,000	55,000	55,000	7,000	5,500	3,500	0,400	0,400	0,400	0,140	0,110	0,070	6,860	5,390	3,430	0,006	0,006	0,006
Buses	Diesel Urban Buses <15t	Conventional	2012	4,479	3,144	2,830	2,628	1,738	1,490	0,729	0,490	0,413	0,175	0,080	0,070	2,453	1,658	1,420	0,030	0,030	0,030
Buses	Diesel Urban Buses <15t	Euro I	2012	1,568	1,120	0,981	0,507	0,364	0,312	0,178	0,136	0,121	0,175	0,080	0,070	0,332	0,284	0,242	0,030	0,030	0,030
Buses	Diesel Urban Buses <15t	Euro II	2012	1,391	0,958	0,806	0,350	0,245	0,209	0,076	0,063	0,061	0,114	0,052	0,046	0,236	0,193	0,163	0,030	0,030	0,030
Buses	Diesel Urban Buses <15t	Euro III	2012	1,509	1,028	0,926	0,318	0,220	0,199	0,094	0,072	0,067	0,103	0,047	0,041	0,215	0,173	0,158	0,030	0,030	0,030
Buses	Diesel Urban Buses <15t	Euro IV	2012	0,800	0,542	0,422	0,043	0,034	0,032	0,032	0,025	0,022	0,005	0,002	0,002	0,038	0,031	0,030	0,030	0,030	0,030
Buses	Diesel Urban Buses <15t	Euro V	2012	1,784	1,211	0,942	0,020	0,016	0,016	0,029	0,019	0,019	0,005	0,002	0,002	0,015	0,014	0,014	0,030	0,030	0,030
Buses	Diesel Urban Buses 15 - 18t	Conventional	2012	4,720	3,242	2,606	1,602	0,977	0,762	0,656	0,439	0,351	0,175	0,080	0,070	1,427	0,897	0,692	0,030	0,030	0,030
Buses	Diesel Urban Buses 15 - 18t	Euro I	2012	2,204	1,612	1,330	0,659	0,431	0,351	0,247	0,177	0,155	0,175	0,080	0,070	0,484	0,351	0,281	0,030	0,030	0,030

Buses	Diesel Urban Buses 15 - 18t	Euro II	2012	1,892	1,310	1,120	0,451	0,296	0,248	0,122	0,099	0,093	0,114	0,052	0,046	0,337	0,244	0,203	0,030	0,030	0,030
Buses	Diesel Urban Buses 15 - 18t	Euro III	2012	2,070	1,382	1,257	0,416	0,269	0,232	0,147	0,111	0,097	0,103	0,047	0,041	0,312	0,222	0,191	0,030	0,030	0,030
Buses	Diesel Urban Buses 15 - 18t	Euro IV	2012	1,045	0,709	0,556	0,061	0,045	0,040	0,044	0,033	0,028	0,005	0,002	0,002	0,055	0,043	0,038	0,030	0,030	0,030
Buses	Diesel Urban Buses 15 - 18t	Euro V	2012	2,396	1,601	1,221	0,028	0,021	0,019	0,042	0,027	0,025	0,005	0,002	0,002	0,022	0,018	0,017	0,030	0,030	0,030
Buses	Diesel Urban Buses >18t	Conventional	2012	6,145	4,310	3,420	1,666	1,018	0,791	0,833	0,575	0,455	0,175	0,080	0,070	1,491	0,938	0,721	0,030	0,030	0,030
Buses	Diesel Urban Buses >18t	Euro I	2012	2,882	2,132	1,965	0,720	0,477	0,386	0,451	0,336	0,311	0,175	0,080	0,070	0,545	0,397	0,316	0,030	0,030	0,030
Buses	Diesel Urban Buses >18t	Euro II	2012	2,541	1,716	1,467	0,491	0,332	0,263	0,102	0,082	0,076	0,114	0,052	0,046	0,377	0,280	0,218	0,030	0,030	0,030
Buses	Diesel Urban Buses >18t	Euro III	2012	2,691	1,778	1,703	0,446	0,291	0,241	0,171	0,123	0,116	0,103	0,047	0,041	0,342	0,244	0,200	0,030	0,030	0,030
Buses	Diesel Urban Buses >18t	Euro IV	2012	1,287	0,869	0,664	0,074	0,055	0,047	0,054	0,039	0,032	0,005	0,002	0,002	0,069	0,053	0,045	0,030	0,030	0,030
Buses	Diesel Urban Buses >18t	Euro V	2012	2,964	1,878	1,405	0,034	0,024	0,023	0,054	0,034	0,032	0,005	0,002	0,002	0,028	0,022	0,021	0,030	0,030	0,030
Buses	Gasoline Coaches	Conventional	2012	70,000	55,000	55,000	7,000	5,500	3,500	0,400	0,400	0,400	0,140	0,110	0,070	6,860	5,390	3,430	0,006	0,006	0,006
Buses	Diesel Coaches <15t	Conventional	2012	2,712	1,738	1,372	0,907	0,533	0,393	0,490	0,328	0,269	0,175	0,080	0,070	0,732	0,453	0,323	0,030	0,030	0,030
Buses	Diesel Coaches <15t	Euro I	2012	2,199	1,466	1,186	0,830	0,516	0,397	0,395	0,260	0,209	0,175	0,080	0,070	0,655	0,436	0,327	0,030	0,030	0,030
Buses	Diesel Coaches <15t	Euro II	2012	1,775	1,203	1,092	0,586	0,359	0,272	0,159	0,117	0,103	0,114	0,052	0,046	0,472	0,307	0,227	0,030	0,030	0,030
Buses	Diesel Coaches <15t	Euro III	2012	2,308	1,464	1,283	0,577	0,351	0,271	0,187	0,122	0,097	0,103	0,047	0,041	0,474	0,304	0,230	0,030	0,030	0,030
Buses	Diesel Coaches <15t	Euro IV	2012	1,241	0,813	0,689	0,072	0,048	0,039	0,048	0,034	0,030	0,005	0,002	0,002	0,067	0,045	0,037	0,030	0,030	0,030
Buses	Diesel Coaches <15t	Euro V	2012	2,375	1,556	1,345	0,036	0,024	0,019	0,049	0,027	0,019	0,005	0,002	0,002	0,030	0,021	0,017	0,030	0,030	0,030
Buses	Diesel Coaches 15 - 18t	Conventional	2012	2,712	1,738	1,372	0,907	0,533	0,393	0,490	0,328	0,269	0,175	0,080	0,070	0,732	0,453	0,323	0,030	0,030	0,030
Buses	Diesel Coaches 15 - 18t	Euro I	2012	2,199	1,466	1,186	0,830	0,516	0,397	0,395	0,260	0,209	0,175	0,080	0,070	0,655	0,436	0,327	0,030	0,030	0,030
Buses	Diesel Coaches 15 - 18t	Euro II	2012	1,775	1,203	1,092	0,586	0,359	0,272	0,162	0,120	0,105	0,114	0,052	0,046	0,472	0,307	0,227	0,030	0,030	0,030
Buses	Diesel Coaches 15 - 18t	Euro III	2012	2,308	1,464	1,283	0,577	0,351	0,271	0,195	0,127	0,101	0,103	0,047	0,041	0,474	0,304	0,230	0,030	0,030	0,030
Buses	Diesel Coaches 15 - 18t	Euro IV	2012	1,241	0,813	0,689	0,072	0,048	0,039	0,048	0,034	0,030	0,005	0,002	0,002	0,067	0,045	0,037	0,030	0,030	0,030
Buses	Diesel Coaches 15 - 18t	Euro V	2012	2,375	1,556	1,345	0,036	0,024	0,019	0,049	0,027	0,019	0,005	0,002	0,002	0,030	0,021	0,017	0,030	0,030	0,030
Buses	Diesel Coaches >18t	Conventional	2012	3,104	2,042	1,732	1,013	0,623	0,482	0,572	0,388	0,331	0,175	0,080	0,070	0,838	0,543	0,412	0,030	0,030	0,030
Buses	Diesel Coaches >18t	Euro I	2012	2,511	1,722	1,458	0,915	0,581	0,457	0,452	0,302	0,246	0,175	0,080	0,070	0,740	0,501	0,387	0,030	0,030	0,030
Buses	Diesel Coaches >18t	Euro II	2012	2,031	1,395	1,290	0,630	0,392	0,305	0,169	0,127	0,113	0,114	0,052	0,046	0,516	0,340	0,259	0,030	0,030	0,030
Buses	Diesel Coaches >18t	Euro III	2012	2,557	1,669	1,439	0,608	0,371	0,286	0,190	0,123	0,099	0,103	0,047	0,041	0,505	0,324	0,245	0,030	0,030	0,030
Buses	Diesel Coaches >18t	Euro IV	2012	1,328	0,875	0,742	0,076	0,050	0,042	0,052	0,036	0,032	0,005	0,002	0,002	0,071	0,048	0,040	0,030	0,030	0,030
Buses	Diesel Coaches >18t	Euro V	2012	2,597	1,719	1,483	0,038	0,025	0,021	0,055	0,031	0,023	0,005	0,002	0,002	0,033	0,023	0,019	0,030	0,030	0,030
Mopeds	2-stroke <50 cm³	Conventional	2012	14,700	14,700		8,881	8,569		0,176	0,176		0,219	0,219		8,662	8,350		0,001	0,001	
Mopeds	2-stroke <50 cm³	Euro I	2012	4,600	4,600		3,858	3,561		0,045	0,045		0,044	0,044		3,814	3,517		0,001	0,001	
Mopeds	2-stroke <50 cm³	Euro II	2012	2,800	2,800		3,034	2,753		0,026	0,026		0,024	0,024		3,009	2,728		0,001	0,001	
Mopeds	4-stroke <50 cm³	Euro II	2012	4,200	4,200		1,224	0,943		0,007	0,007		0,024	0,024		1,199	0,918		0,001	0,001	
Motorcycles	2-stroke >50 cm³	Conventional	2012	15,605	19,285	28,470	9,341	7,323	9,836	0,200	0,200	0,200	0,150	0,150	0,150	9,191	7,173	9,686	0,002	0,002	0,002
Motorcycles	2-stroke >50 cm³	Euro I	2012	10,315	12,786	18,933	9,230	7,294	9,832	0,080	0,080	0,080	0,099	0,107	0,098	9,131	7,187	9,734	0,002	0,002	0,002
Motorcycles	2-stroke >50 cm³	Euro II	2012	8,146	10,067	14,890	3,410	2,780	4,186	0,040	0,040	0,040	0,030	0,032	0,030	3,380	2,749	4,156	0,002	0,002	0,002

Motorcycles	2-stroke >50 cm ³	Euro III	2012	4,510	5,593	8,342	2,193	1,589	2,275	0,012	0,012	0,012	0,012	0,014	0,012	2,181	1,575	2,263	0,002	0,002	0,002
Motorcycles	4-stroke <250 cm ³	Conventional	2012	15,258	17,209	24,960	1,075	0,348	0,174	0,020	0,020	0,020	0,200	0,200	0,200	0,875	0,148	-0,026	0,002	0,002	0,002
Motorcycles	4-stroke <250 cm ³	Euro I	2012	10,391	14,456	24,910	2,079	1,082	1,008	0,020	0,020	0,020	0,142	0,144	0,132	1,937	0,938	0,876	0,002	0,002	0,002
Motorcycles	4-stroke <250 cm ³	Euro II	2012	3,708	5,765	9,135	1,859	1,054	0,996	0,005	0,005	0,005	0,136	0,092	0,092	1,723	0,962	0,904	0,002	0,002	0,002
Motorcycles	4-stroke <250 cm ³	Euro III	2012	2,060	3,201	5,092	1,264	0,649	0,542	0,005	0,005	0,005	0,082	0,032	0,028	1,182	0,617	0,514	0,002	0,002	0,002
Motorcycles	4-stroke 250 - 750 cm ³	Conventional	2012	20,461	19,486	22,990	1,492	0,732	0,397	0,020	0,020	0,020	0,200	0,200	0,200	1,292	0,532	0,197	0,002	0,002	0,002
Motorcycles	4-stroke 250 - 750 cm ³	Euro I	2012	10,599	9,003	10,460	3,227	1,738	1,111	0,020	0,020	0,020	0,148	0,174	0,156	3,079	1,564	0,955	0,002	0,002	0,002
Motorcycles	4-stroke 250 - 750 cm ³	Euro II	2012	2,230	2,436	6,092	2,143	1,136	0,859	0,005	0,005	0,005	0,156	0,120	0,122	1,987	1,016	0,737	0,002	0,002	0,002
Motorcycles	4-stroke 250 - 750 cm ³	Euro III	2012	1,228	1,345	3,357	1,405	0,708	0,646	0,005	0,005	0,005	0,094	0,042	0,036	1,311	0,666	0,610	0,002	0,002	0,002
Motorcycles	4-stroke >750 cm ³	Conventional	2012	20,461	19,486	22,990	1,339	0,581	0,592	0,020	0,020	0,020	0,200	0,200	0,200	1,139	0,381	0,392	0,002	0,002	0,002
Motorcycles	4-stroke >750 cm ³	Euro I	2012	10,599	9,003	10,460	3,332	1,859	1,586	0,020	0,020	0,020	0,092	0,092	0,154	3,240	1,767	1,432	0,002	0,002	0,002
Motorcycles	4-stroke >750 cm ³	Euro II	2012	2,230	2,436	6,092	1,904	0,884	0,687	0,005	0,005	0,005	0,084	0,062	0,102	1,820	0,822	0,585	0,002	0,002	0,002
Motorcycles	4-stroke >750 cm ³	Euro III	2012	1,228	1,345	3,357	1,192	0,517	0,447	0,005	0,005	0,005	0,050	0,022	0,030	1,142	0,495	0,417	0,002	0,002	0,002

Annex 3B-8 Fuel consumption (GJ) and emissions (tonnes) per vehicle category and as totals

Sector	Year	FC (PJ)	SO ₂	NO _x	NM VOC	CH ₄	CO	CO ₂	N ₂ O	NH ₃	TSP
Passenger Cars	1985	66	1714	57216	68940	1576	543043	4847	164	48	1689
Passenger Cars	1986	67	1152	58346	68917	1599	515315	4901	166	49	1719
Passenger Cars	1987	67	1149	59333	68717	1621	492705	4914	168	50	1690
Passenger Cars	1988	68	1169	61147	68768	1645	455881	4975	173	51	1641
Passenger Cars	1989	67	848	61153	67298	1631	423852	4932	172	51	1602
Passenger Cars	1990	72	908	65510	70690	1737	431213	5235	183	54	1667
Passenger Cars	1991	76	953	68046	73286	1823	450464	5567	197	146	1704
Passenger Cars	1992	80	672	67915	72817	1838	436546	5816	214	339	1614
Passenger Cars	1993	82	365	66179	70395	1824	423024	5966	225	529	1557
Passenger Cars	1994	85	385	63881	67374	1774	391546	6197	241	807	1498
Passenger Cars	1995	86	387	59513	62856	1679	371759	6256	250	1073	1396
Passenger Cars	1996	86	391	55400	58678	1590	360862	6320	259	1323	1312
Passenger Cars	1997	89	398	52173	53954	1513	324514	6479	270	1663	1183
Passenger Cars	1998	90	407	48139	48596	1435	301585	6608	271	2013	1070
Passenger Cars	1999	91	317	43763	42860	1332	265548	6644	270	2291	984
Passenger Cars	2000	90	207	40012	35465	1237	241082	6603	267	2473	902
Passenger Cars	2001	89	204	36872	32566	1143	230578	6505	258	2479	838
Passenger Cars	2002	90	207	34667	29210	1057	210434	6592	255	2495	787
Passenger Cars	2003	92	211	32642	26639	981	200641	6749	252	2472	796
Passenger Cars	2004	93	213	30307	22720	881	175387	6804	245	2444	764
Passenger Cars	2005	92	42	27004	20423	776	167079	6703	230	2309	730
Passenger Cars	2006	92	42	24371	17301	681	147053	6749	220	2202	699
Passenger Cars	2007	97	44	23359	15080	616	133106	7077	222	2124	702
Passenger Cars	2008	98	45	22085	13224	541	122310	7168	218	1969	723
Passenger Cars	2009	96	44	20585	11501	482	109132	7021	210	1836	665
Passenger Cars	2010	96	44	19951	10345	427	102701	6934	206	1670	666
Passenger Cars	2011	96	43	19933	8517	375	84379	6814	208	1548	621
Passenger Cars	2012	96	42	19535	7537	333	77006	6723	204	1410	589
Light Duty Vehicles	1985	12	2488	5442	1851	81	13182	918	4	4	1383
Light Duty Vehicles	1986	14	1741	6187	2034	92	14486	1059	4	5	1595
Light Duty Vehicles	1987	15	1828	6505	2146	96	15339	1112	5	5	1684
Light Duty Vehicles	1988	15	1883	6663	2158	99	15137	1144	5	6	1676
Light Duty Vehicles	1989	16	1316	6845	2158	102	15084	1189	5	6	1731
Light Duty Vehicles	1990	17	1414	7313	2282	108	15974	1275	5	6	1854
Light Duty Vehicles	1991	18	1462	7623	2425	112	17120	1322	5	6	1960
Light Duty Vehicles	1992	18	936	7672	2505	115	17473	1312	6	6	1903
Light Duty Vehicles	1993	18	372	7946	2636	118	18674	1350	6	7	1992
Light Duty Vehicles	1994	20	400	8446	2778	126	19447	1445	6	7	2110
Light Duty Vehicles	1995	20	402	8344	2693	122	18847	1448	8	11	2045
Light Duty Vehicles	1996	20	411	8293	2567	117	18071	1475	11	19	1957
Light Duty Vehicles	1997	20	417	8186	2373	113	16226	1494	15	28	1737
Light Duty Vehicles	1998	21	424	8193	2265	110	15566	1525	19	38	1591
Light Duty Vehicles	1999	21	240	8161	2126	103	14284	1553	24	48	1432
Light Duty Vehicles	2000	21	50	8182	1939	96	13511	1581	29	58	1287
Light Duty Vehicles	2001	22	51	8292	1988	91	14006	1617	35	72	1184
Light Duty Vehicles	2002	23	53	8356	1873	84	13102	1669	40	78	1062
Light Duty Vehicles	2003	25	57	8788	1849	77	12940	1810	45	79	1036
Light Duty Vehicles	2004	27	62	9256	1759	74	12316	1970	52	80	953
Light Duty Vehicles	2005	28	13	9533	1736	64	12171	2100	58	78	918
Light Duty Vehicles	2006	31	15	10202	1694	57	11671	2319	65	77	900

Light Duty Vehicles	2007	34	16	10722	1632	50	11049	2502	71	73	885
Light Duty Vehicles	2008	31	15	9559	1349	39	9173	2320	67	59	741
Light Duty Vehicles	2009	29	13	8509	1147	32	7920	2110	62	53	629
Light Duty Vehicles	2010	27	13	7989	1065	27	7397	2028	60	48	579
Light Duty Vehicles	2011	26	12	7358	878	22	6060	1847	57	40	489
Light Duty Vehicles	2012	24	11	6687	757	18	5218	1674	53	34	424
Heavy Duty Vehicles	1985	25	5735	24022	1429	180	5850	1819	75	7	884
Heavy Duty Vehicles	1986	28	3881	27058	1579	203	6469	2050	83	8	994
Heavy Duty Vehicles	1987	27	3804	26499	1531	198	6300	2009	81	8	973
Heavy Duty Vehicles	1988	27	3734	25996	1473	195	6108	1972	79	8	953
Heavy Duty Vehicles	1989	28	2593	27058	1520	203	6297	2053	82	8	991
Heavy Duty Vehicles	1990	29	2664	27736	1506	208	6346	2110	83	8	1013
Heavy Duty Vehicles	1991	29	2706	28201	1540	211	6476	2143	85	8	1030
Heavy Duty Vehicles	1992	28	1727	27644	1481	208	6321	2103	82	8	1008
Heavy Duty Vehicles	1993	27	641	26586	1386	201	6027	2032	78	8	969
Heavy Duty Vehicles	1994	29	682	27825	1440	215	6254	2159	83	8	1017
Heavy Duty Vehicles	1995	29	685	27113	1432	220	6182	2169	86	9	999
Heavy Duty Vehicles	1996	30	702	26773	1384	227	6109	2220	88	9	987
Heavy Duty Vehicles	1997	30	713	26566	1316	229	5998	2254	90	9	940
Heavy Duty Vehicles	1998	31	725	26685	1237	231	5913	2292	93	9	877
Heavy Duty Vehicles	1999	32	416	27536	1189	238	5985	2390	98	10	841
Heavy Duty Vehicles	2000	31	73	26453	1064	226	5654	2316	95	10	759
Heavy Duty Vehicles	2001	32	76	26939	1032	231	5760	2403	99	10	726
Heavy Duty Vehicles	2002	32	76	25965	956	225	5706	2392	98	10	673
Heavy Duty Vehicles	2003	34	80	26293	937	233	5955	2513	103	10	658
Heavy Duty Vehicles	2004	35	82	26281	919	239	6121	2595	106	11	633
Heavy Duty Vehicles	2005	36	17	26299	899	244	6316	2690	108	11	613
Heavy Duty Vehicles	2006	38	18	26414	887	248	6541	2801	112	11	598
Heavy Duty Vehicles	2007	39	18	25325	799	223	6432	2879	115	12	539
Heavy Duty Vehicles	2008	36	17	21282	599	164	5497	2672	107	11	405
Heavy Duty Vehicles	2009	32	15	17192	442	117	4626	2345	94	9	302
Heavy Duty Vehicles	2010	33	15	16515	397	103	4705	2427	97	10	260
Heavy Duty Vehicles	2011	34	15	15688	353	90	4866	2426	101	10	240
Heavy Duty Vehicles	2012	32	14	13372	277	70	4580	2209	95	9	199
Buses	1985	7	1683	7284	744	77	3022	541	18	2	312
Buses	1986	8	1087	7830	793	83	3187	581	19	2	335
Buses	1987	8	1065	7674	778	81	3114	569	19	2	329
Buses	1988	8	1069	7710	780	82	3103	572	19	2	330
Buses	1989	8	730	7890	794	83	3132	585	19	2	338
Buses	1990	8	780	8414	845	89	3362	625	21	2	360
Buses	1991	8	781	8430	848	89	3389	626	21	2	361
Buses	1992	8	484	8049	816	85	3314	598	20	2	344
Buses	1993	8	189	8142	828	86	3401	605	20	2	348
Buses	1994	9	200	8526	865	93	3769	645	22	2	362
Buses	1995	9	206	8559	852	97	4034	667	23	2	359
Buses	1996	9	216	8733	843	103	4222	700	24	2	362
Buses	1997	9	215	8523	790	102	4070	695	24	2	343
Buses	1998	9	210	8231	725	97	3865	679	24	2	312
Buses	1999	9	113	7915	655	92	3601	662	24	2	282
Buses	2000	9	20	7567	594	86	3383	640	23	2	254
Buses	2001	9	20	7493	561	84	3242	639	23	2	229
Buses	2002	9	20	7296	523	81	3134	638	23	2	213
Buses	2003	9	21	7454	505	81	3072	672	25	3	200
Buses	2004	9	22	7417	483	80	2977	686	25	3	193
Buses	2005	9	4	6965	437	74	2734	658	24	2	170
Buses	2006	9	4	6762	407	71	2572	654	24	2	159

Buses	2007	9	4	6608	378	67	2426	662	25	2	149
Buses	2008	9	4	6105	328	57	2172	644	24	2	129
Buses	2009	8	4	5536	273	47	1899	619	23	2	108
Buses	2010	8	4	5315	240	41	1780	627	24	2	95
Buses	2011	8	4	5019	206	35	1680	601	24	2	84
Buses	2012	8	4	4580	169	29	1539	558	23	2	72
Mopeds	1985	0	1	12	1832	47	3144	17	0	0	38
Mopeds	1986	0	0	11	1645	42	2821	15	0	0	34
Mopeds	1987	0	0	10	1522	39	2611	14	0	0	31
Mopeds	1988	0	0	9	1435	37	2453	13	0	0	29
Mopeds	1989	0	0	9	1364	35	2326	13	0	0	28
Mopeds	1990	0	0	9	1386	35	2366	13	0	0	28
Mopeds	1991	0	0	9	1419	36	2428	13	0	0	29
Mopeds	1992	0	0	9	1426	36	2441	13	0	0	29
Mopeds	1993	0	0	9	1408	36	2418	13	0	0	29
Mopeds	1994	0	0	9	1390	35	2383	13	0	0	29
Mopeds	1995	0	0	10	1564	40	2683	15	0	0	32
Mopeds	1996	0	1	12	1801	46	3102	17	0	0	37
Mopeds	1997	0	1	15	2220	57	3822	21	0	0	46
Mopeds	1998	0	1	17	2655	68	4587	25	0	0	55
Mopeds	1999	0	1	17	2575	66	4435	24	0	0	53
Mopeds	2000	0	1	19	2507	64	4323	24	0	0	52
Mopeds	2001	0	1	19	1951	48	3310	19	0	0	39
Mopeds	2002	0	1	23	1972	48	3297	20	0	0	39
Mopeds	2003	0	1	25	1901	45	3153	20	0	0	37
Mopeds	2004	0	1	25	1785	42	2940	19	0	0	34
Mopeds	2005	0	0	27	1682	38	2740	19	0	0	32
Mopeds	2006	0	0	29	1606	36	2589	19	0	0	29
Mopeds	2007	0	0	31	1524	33	2437	19	0	0	27
Mopeds	2008	0	0	31	1419	30	2255	18	0	0	25
Mopeds	2009	0	0	32	1323	28	2085	18	0	0	23
Mopeds	2010	0	0	31	1193	24	1872	16	0	0	20
Mopeds	2011	0	0	30	1089	22	1687	15	0	0	18
Mopeds	2012	0	0	31	1001	19	1532	15	0	0	16
Motorcycles	1985	0	1	57	507	53	5437	25	1	1	13
Motorcycles	1986	0	1	57	508	53	5439	25	1	1	13
Motorcycles	1987	0	1	55	494	52	5299	25	1	1	13
Motorcycles	1988	0	1	56	506	53	5365	25	1	1	13
Motorcycles	1989	0	1	55	501	52	5284	25	1	1	13
Motorcycles	1990	0	1	59	536	56	5670	27	1	1	14
Motorcycles	1991	0	1	61	549	57	5839	27	1	1	14
Motorcycles	1992	0	1	65	588	61	6234	29	1	1	15
Motorcycles	1993	0	1	69	614	65	6599	31	1	1	16
Motorcycles	1994	0	1	74	662	69	7041	33	1	1	17
Motorcycles	1995	0	1	76	683	71	7275	34	1	1	17
Motorcycles	1996	0	1	78	692	73	7418	35	1	1	18
Motorcycles	1997	0	1	82	736	76	7784	36	1	1	19
Motorcycles	1998	1	1	85	757	79	8069	38	1	1	19
Motorcycles	1999	1	1	87	780	80	8222	38	1	1	20
Motorcycles	2000	1	1	87	751	79	8078	39	1	1	19
Motorcycles	2001	1	1	90	762	79	7929	39	1	1	19
Motorcycles	2002	1	1	94	788	80	7976	41	1	1	18
Motorcycles	2003	1	1	96	795	79	7880	41	1	1	18
Motorcycles	2004	1	1	97	784	78	7622	41	1	1	17
Motorcycles	2005	1	0	102	792	77	7420	42	1	1	16
Motorcycles	2006	1	0	110	821	79	7360	45	1	1	16

Motorcycles	2007	1	0	117	839	81	7341	48	1	1	15
Motorcycles	2008	1	0	117	811	78	6997	49	1	1	14
Motorcycles	2009	1	0	112	765	73	6538	47	1	1	13
Motorcycles	2010	1	0	113	742	73	6463	47	1	1	13
Motorcycles	2011	1	0	112	726	71	6285	46	1	1	12
Motorcycles	2012	1	0	111	699	70	6148	45	1	1	12
Total	1985	111	11621	94032	75302	2015	573678	8167	261	63	4320
Total	1986	118	7862	99488	75477	2072	547717	8632	274	65	4690
Total	1987	118	7847	100077	75187	2088	525368	8644	274	66	4721
Total	1988	118	7857	101581	75121	2110	488047	8702	276	67	4643
Total	1989	120	5488	103010	73635	2105	455974	8797	279	68	4701
Total	1990	126	5767	109042	77245	2233	464930	9284	292	72	4936
Total	1991	132	5903	112371	80067	2329	485715	9699	308	163	5097
Total	1992	134	3820	111354	79633	2343	472329	9872	322	356	4913
Total	1993	136	1569	108932	77266	2330	460142	9997	330	546	4910
Total	1994	143	1669	108761	74509	2312	430439	10493	353	825	5033
Total	1995	144	1682	103616	70081	2230	410780	10589	368	1096	4849
Total	1996	147	1721	99289	65965	2155	399783	10766	383	1354	4672
Total	1997	149	1744	95544	61389	2090	362414	10979	401	1703	4268
Total	1998	152	1768	91350	56235	2020	339585	11167	409	2063	3924
Total	1999	154	1088	87478	50185	1912	302074	11313	417	2352	3611
Total	2000	153	352	82320	42322	1788	276031	11203	415	2544	3273
Total	2001	153	353	79704	38860	1676	264825	11223	417	2565	3035
Total	2002	155	357	76400	35321	1574	243650	11352	417	2586	2793
Total	2003	161	371	75298	32626	1497	233641	11806	426	2565	2745
Total	2004	165	381	73383	28450	1393	207364	12115	430	2538	2595
Total	2005	166	77	69930	25969	1274	198459	12214	422	2402	2479
Total	2006	171	79	67887	22716	1172	177787	12587	423	2294	2403
Total	2007	179	83	66162	20251	1070	162791	13187	434	2212	2318
Total	2008	175	81	59179	17728	909	148403	12870	417	2043	2038
Total	2009	166	77	51966	15452	778	132200	12160	390	1902	1740
Total	2010	165	76	49914	13982	695	124918	12080	388	1731	1632
Total	2011	165	74	48140	11769	616	104957	11748	391	1602	1464
Total	2012	161	71	44316	10439	538	96022	11224	376	1456	1311

Annex 3B-9 COPERT IV:DEA statistics fuel use ratios and mileage adjustment factors

Sales			1985	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
Fuel ratio	Gasoline	DEA:COPERT IV	1,06	1,01	1,05	1,10	1,13	1,16	1,14	1,12	1,13	1,14	1,12	1,13	1,12	1,13	1,13	1,11	1,10	1,10	1,12	1,11	1,11	1,10	1,08	1,07
	Diesel	DEA:COPERT IV	1,22	1,33	1,36	1,32	1,32	1,40	1,39	1,40	1,42	1,41	1,40	1,35	1,36	1,35	1,42	1,44	1,40	1,40	1,43	1,40	1,34	1,37	1,38	1,37
Consumption																										
Fuel ratio	Gasoline	DEA:COPERT IV	1,11	1,11	1,10	1,11	1,12	1,13	1,13	1,13	1,14	1,15	1,17	1,19	1,16	1,17	1,16	1,14	1,13	1,12	1,12	1,11	1,11	1,13	1,11	1,10
	Diesel	DEA:COPERT IV	1,13	1,22	1,30	1,27	1,26	1,30	1,27	1,28	1,29	1,28	1,28	1,26	1,27	1,25	1,28	1,29	1,26	1,27	1,29	1,26	1,22	1,25	1,24	1,23

Annex 3B-10 Actual vs. representative aircraft types, no. of LTO's from Danish airports, no. of flights between Denmark and Greenland/Faroe Islands, LTO and average cruise fuel consumption and emission factors

Correspondence table between actual aircraft type codes and representative aircraft types

ICAO code	Representative aircraft	Type	ICAO code	Representative aircraft	Type	ICAO code	Representative aircraft	Type
B73	B737 400	L2J	BN2T	Cessna 208 Caravan	L2T	H60	S61	H2T
739	B737 400	L2J	BSTP	S61	H2T	HA4T	RJ 100	L2J
7474	B737 400	L2J	C10T	Cessna 208 Caravan	L1T	HEL	Shorts 360 300	
757	B757	L2J	C130	Lockheed C-130H Hercules	L4T	HF20	RJ 100	L2J
A109	S61	H2T	C141	DC10-30	L4J	HR16	F50	L2T
A124	B747 400	L4J	C160	F50	L2T	HS25	RJ 100	L2J
A139	S61	H2T	C17	A340	L4J	HS74	F50	L2T
A300	A310	L2J	C17C	A340	L4J	HTA0	BAe Jetstream 31	L2T
A304	A310	L2J	C208	Cessna 208 Caravan	L1T	HU30	S61	H1P
A306	A310	L2J	C20A	Shorts 360 300	L2T	HU50	Shorts 360 300	L2T
A30B	A310	L2J	C212	Shorts 360 300	L2T	IL18	Lockheed P-3B Orion	L4T
A310	A310	L2J	C25A	RJ 100	L2J	IL62	B767 300 ER	L4J
A318	A320	L2J	C25B	RJ 100	L2J	IL76	B767 300 ER	L4J
A319	A320	L2J	C27J	Dash8 400	L2T	IL86	A340	L4J
A320	A320	L2J	C30J	Lockheed C-130H Hercules	L4T	IL96	A340	L4J
A321	A320	L2J	C406	Shorts 360 300	L2T	J32	BAe Jetstream 31	L2T
A322	A320	L2J	C425	Reims F406 Caravan II	L2T	J328	RJ 100	L2J
A330	A330	L2J	C441	Reims F406 Caravan II	L2T	JET	RJ 100	
A332	A330	L2J	C5	DC10-30	L4J	JS20	BAe Jetstream 31	L2T
A333	A330	L2J	C500	RJ 100	L2J	JS31	BAe Jetstream 31	L2T
A340	A340	L4J	C501	RJ 100	L2J	JS32	BAe Jetstream 31	L2T
A343	A340	L4J	C510	RJ 100	L2J	JS41	BAe Jetstream 41	L2T
A550	S61	H1P	C525	RJ 100	L2J	JSTA	Shorts 360 300	L2T
A748	Shorts 360 300	L2T	C550	RJ 100	L2J	JSTB	Shorts 360 300	L2T
AB30	A310	L2J	C551	RJ 100	L2J	KA27	S61	H2T
AC14	Shorts 360 300	L2T	C560	RJ 100	L2J	KODI	De Havilland DHC-3 Turbo-Otter	L1T
AC6T	Shorts 360 300	L2T	C56X	RJ 100	L2J	L101	DC10-30	L3J
AC90	Shorts 360 300	L2T	C650	RJ 100	L2J	L188	Dash8 400	L4T
AC95	Shorts 360 300	L2T	C680	RJ 100	L2J	L29A	BAe146	L4J
ALO3	S61	H1T	C750	RJ 100	L2J	L29B	BAe146	L4J
AN12	Dash8 400	L4T	CL3	RJ 100	L2J	L329	BAe146	L4J
AN22	Lockheed C-130H Hercules	L4T	CL30	RJ 100	L2J	L382	Shorts 360 300	L2T

AN24	F50	L2T	CL60	RJ 100	L2J	LJ24	RJ 100	L2J
AN26	Antonov 26	L2T	CL65	RJ 100	L2J	LJ25	RJ 100	L2J
AN28	Shorts 360 300	L2T	CN35	F50	L2T	LJ31	RJ 100	L2J
AN30	Antonov 26	L2T	CRJ	RJ 100	L2J	LJ35	RJ 100	L2J
AN32	Antonov 26	L2T	CRJ1	RJ 100	L2J	LJ36	RJ 100	L2J
AN7	BAC1-11	L2J	CRJ2	RJ 100	L2J	LJ40	RJ 100	L2J
AN72	BAC1-11	L2J	CRJ7	RJ 100	L2J	LJ45	RJ 100	L2J
AN74	BAC1-11	L2J	CRJ9	CRJ9	L2J	LJ55	RJ 100	L2J
ANF	Dash8 400	L4T	CV44	Dash8 400	L2T	LJ60	RJ 100	L2J
APF	ATR 42-320	L2T	CV58	Dash8 400	L2T	LR24	RJ 100	L2J
AS32	S61	H2T	CVLT	Dash8 400	L2T	LR25	RJ 100	L2J
AS35	S61	H1T	D228	Dornier 328-110	L2T	LR31	RJ 100	L2J
AS50	S61	H1T	D328	Shorts 360 300	L2T	LR35	RJ 100	L2J
AS55	S61	H2T	DA10	RJ 100	L2J	LR36	RJ 100	L2J
AS65	S61	H2T	DA20	RJ 100	L2J	LR55	RJ 100	L2J
ASJ	Shorts 360 300	L2T	DA30	RJ 100	L2J	LR60	RJ 100	L2J
ASTR	RJ 100	L2J	DA50	RJ 100	L3J	LYNX	S61	H2T
AT42	ATR 42-320	L2T	DA90	RJ 100	L3J	M20K	Shorts 360 300	L2T
AT43	ATR 42-320	L2T	DC10	DC10-30	L3J	M7T	Cessna 208 Caravan	L1T
AT44	ATR 42-320	L2T	DC8	B767 300 ER	L2J	MD11	DC10-30	L3J
AT45	ATR 42-320	L2T	DC85	B767 300 ER	L4J	MD52	S61	H1T
AT5	ATR 42-320	L2T	DC86	B767 300 ER	L4J	MD80	MD 82	L2J
AT72	ATR 72-200	L2T	DC87	B767 300 ER	L4J	MD81	MD 82	L2J
ATP	S2000	L2T	DC8F	B767 300 ER	L2J	MD82	MD 82	L2J
ATR	ATR 42-320	L2T	DC8S	B767 300 ER	L4J	MD83	MD 82	L2J
ATR4	ATR 42-320	L2T	DC9	DC9	L2J	MD87	MD 82	L2J
ATR7	ATR 72-200	L2T	DC93	RJ 100	L2J	MD88	MD 82	L2J
AVRO	BAe146	L4J	DC94	DC9	L2J	MD90	B737 400	L2J
AW13	S61	H2T	DC95	DC9	L2J	MI14	S61	H2T
B06	S61	H1T	DF2	RJ 100	L2J	MI2	S61	H2T
B105	S61	H2T	DH 7	DHC7	L2T	MI8	S61	H2T
B12	S61	H2T	DH2T	De Havilland DHC-3 Turbo-Otter	L1T	MU2	Shorts 360 300	L2T
B190	Beech 1900C Airliner	L2T	DH4	Dash8 400	L2T	MU20	Shorts 360 300	L2T
B200	Shorts 360 300	L2T	DH6	Shorts 360 300	L2T	MU30	RJ 100	L2J
B206	S61	H1T	DH7	DHC7	L2T	N24A	Shorts 360 300	L2T
B212	S61	H2T	DH8	Dash8 400	L2T	N262	Shorts 360 300	L2T
B222	S61	H2T	DH8A	Dash8 400	L2T	ND26	Shorts 360 300	L2T

B321	A320	L2J	DH8C	Dash8 400	L2T	NH90	S61	L1P
B350	Beech Super King Air 350	L2T	DH8D	Dash8 400	L2T	NOMA	Shorts 360 300	L2T
B378	B737 400	L2J	DHC6	Shorts 360 300	L2T	OTH	F50	L2T
B407	S61	H1T	DHC8	Dash8 400	L2T	P180	Embraer 110P2A	L2T
B412	S61	H2T	E110	Embraer 110P2A	L2T	P46T	Cessna 208 Caravan	L1T
B429	S61	H2T	E120	Shorts 360 300	L2T	P750	Cessna 208 Caravan	L1T
B430	S61	H2T	E121	Embraer 110P2A	L2T	PA42	Reims F406 Caravan II	L2T
B461	BAe146	L4J	E135	RJ 100	L2J	PA60	Cessna 208 Caravan	L2T
B462	BAe146	L4J	E145	RJ 100	L2J	PAT4	Shorts 360 300	L2T
B463	BAe146	L4J	E170	CRJ9	L2J	PAY1	Reims F406 Caravan II	L2T
B46C	BAe146	L4J	E175	CRJ9	L2J	PAY2	Reims F406 Caravan II	L2T
B703	B757	L4J	E19	B737 100	L2J	PAY3	Reims F406 Caravan II	L2T
B707	B757	L2J	E190	B737 100	L2J	PAY4	Shorts 360 300	L2T
B712	B737 100	L2J	E195	B737 100	L2J	PAZT	Shorts 360 300	L2T
B717	DC9	L2J	E70	CRJ9	L2J	PC12	Cessna 208 Caravan	L1T
B720	B757	L4J	E90	B737 100	L2J	PC7	Cessna 208 Caravan	L1T
B721	B727	L3J	EA19	A320	L2J	PC9	Cessna 208 Caravan	L1T
B722	B727	L3J	EA30	A310	L2J	PRM1	RJ 100	L2J
B727	B727	L2J	EA31	A310	L2J	PUMA	S61	H2T
B72S	B727	L3J	EA32	A320	L2J	R22	S61	H1P
B732	B737 400	L2J	EA33	A330	L2J	R44	S61	H1P
B733	B737 400	L2J	EA34	A340	L4J	RH22	Shorts 360 300	L2T
B734	B737 400	L2J	EA50	RJ 100	L2J	RH44	S61	H1P
B735	B737 400	L2J	EC12	S61	H1P	RJ1H	BAe146	L4J
B736	B737 400	L2J	EC20	S61	H1T	RJ70	RJ 100	L4J
B737	B737 400	L2J	EC25	S61	H2T	RJ85	RJ 100	L4J
B738	B737 400	L2J	EC30	S61	H1T	S210	DC9	L2J
B739	B737 400	L2J	EC35	S61	H2T	S269	S61	H1P
B73A	B737 100	L2J	EC45	S61	H2T	S330	S61	H1T
B73B	B737 400	L2J	EC55	S61	H2T	S350	F50	L2T
B73C	B737 400	L2J	EH10	S61	H3T	S355	S61	H1T
B73E	B737 400	L2J	EMB	Shorts 360 300	L2T	S365	S61	H1T
B73G	B737 400	L2J	EN28	S61	H1P	S601	RJ 100	L2J
B73S	B737 400	L2J	ER3	RJ 100	L2J	S61	S61	H2T
B741	B747 100-300	L4J	EXPL	S61	H2T	S65C	S61	H2T
B742	B747 100-300	L4J	F100	F100	L2J	S76	S61	H2T
B743	B747 100-300	L4J	F26T	Cessna 208 Caravan	L1T	S893	Shorts 360 300	L2T

B744	B747 400	L4J	F27	Fokker 27 Friendship	L2T	S92	S61	H2T
B747	B747 400	L4J	F28	F28	L2T	SA22	Shorts 360 300	L2T
B74A	B747 400	L4J	F2TH	RJ 100	L2J	SB05	RJ 100	L2J
B74B	B747 400	L4J	F406	Reims F406 Caravan II	L2T	SB20	S2000	L2T
B74D	B747 400	L4J	F50	F50	L2T	SBR1	RJ 100	L2J
B74F	B747 400	L4J	F70	F28	L2T	SC7	Shorts SC.7 Srs3M-200	L2T
B74S	B747 100-300	L4J	F71	F28	L2J	SF34	Saab 340B	L2T
B752	B757	L2J	F900	RJ 100	L3J	SH33	Shorts 330	L2T
B757	B757	L2J	FA10	RJ 100	L2J	SH36	Shorts 360 300	L2T
B762	B767 300 ER	L2J	FA20	RJ 100	L2J	SH60	S61	H2T
B763	B767 300 ER	L2J	FA50	RJ 100	L3J	SH7	Shorts 360 300	L2T
B764	B767 300 ER	L2J	FA7X	RJ 100	L3J	SK61	S61	H2T
B767	B767 300 ER	L2J	FK10	F100	L2J	SK76	S61	H2T
B772	B777	L2J	FK27	F50	L2T	STAR	Shorts 360 300	L2T
B773	B777	L2J	FK28	F28	L2J	SW2	Swearingen Metro III	L2T
B777	B777	L2J	FK50	F50	L2T	SW3	Swearingen Metro III	L2T
BA11	BAC1-11	L2J	FK70	F28	L2J	SW4	Swearingen Metro III	L2T
BA14	BAe146	L4J	FOUG	RJ 100	L2J	SW4A	S61	H1T
BA31	Shorts 360 300	L2T	G159	Shorts 360 300	L2T	SW4B	S61	H1T
BA32	Shorts 360 300	L2T	G2	Shorts 360 300	L2T	T134	F100	L2J
BA41	Shorts 360 300	L2T	G222	F28	L2T	T154	B727	L3J
BA46	BAe146	L4J	G3	F50	L2T	T204	B757	L2J
BAE1	BAe146	L4J	G4	CRJ9	L2J	TB21	Shorts 360 300	L2T
BATP	F50	L2T	GALX	RJ 100	L2J	TB9	Shorts 360 300	L2T
BE02	Shorts 360 300	L2T	GAZL	S61	H1T	TBM7	Cessna 208 Caravan	L1T
BE10	Beech Super King Air 200B	L2T	GIV	CRJ9	L2J	TBM8	Cessna 208 Caravan	L1T
BE20	Beech Super King Air 200B	L2T	GLEX	RJ 100	L2J	TEX2	Cessna 208 Caravan	L1T
BE30	Beech Super King Air 350	L2T	GLF2	RJ 100	L2J	TOR	RJ 100	L2J
BE40	RJ 100	L2J	GLF3	RJ 100	L2J	TU34	F100	L2J
BE90	Beech Super King Air 200B	L2T	GLF4	RJ 100	L2J	TU54	B757	L2J
BE99	Beech Super King Air 200B	L2T	GLF5	RJ 100	L2J	UH1	S61	H1T
BE9L	Reims F406 Caravan II	L2T	GULF	F50	L2T	VC10	B757	L4J
BE9T	Reims F406 Caravan II	L2T	H25A	RJ 100	L2J	VF14	RJ 100	L2J
BH06	S61	H1T	H25B	RJ 100	L2J	W3	S61	H2T
BH12	S61	H1T	H25C	RJ 100	L2J	WW24	RJ 100	L2J
BH21	S61	H1T	H269	S61	H1P	WW25	F50	L2T
BH41	S61	H1T	H36	S61	H1P	Y12	Shorts 360 300	L2T

BK17	S61	H2T	H46	S61	H2T	YK40	RJ 100	L3J
BN2	Shorts 360 300	L2T	H500	S61	H1T	YK42	DC9	L3J

LTO no. per representative aircraft type for domestic and int. flights (Copenhagen and other airports).

Flight	Airport name	Rep Aircraft	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
Domestic	Copenhagen	A310	37	1	1	3	1	1		1	1			
Domestic	Copenhagen	A320	71	83	110	77	115	237	263	554	536	497	590	1121
Domestic	Copenhagen	A330	4	28	213	228	228	228	232	211	225	223	237	251
Domestic	Copenhagen	A340	7	3	5	1				1			2	1
Domestic	Copenhagen	Antonov 26				91	284	246	253	253	249	63		
Domestic	Copenhagen	ATR 42-320	4494	5333	4951	2933	804	3320	3393	3820	2455	2925	2799	889
Domestic	Copenhagen	ATR 72-200	2358	2783	4495	5218	6664	5775	5449	7005	5697	6763	8108	4238
Domestic	Copenhagen	B727		1										
Domestic	Copenhagen	B737 400	2264	1722	2212	959	514	549	1258	1376	2240	3521	3172	3108
Domestic	Copenhagen	B747 400					1		1			1		
Domestic	Copenhagen	B757	227	264	152	146	100	101	141	154	112	85	4	
Domestic	Copenhagen	B767 300 ER	206	182	24	1			1	24	15	1	4	4
Domestic	Copenhagen	BAC1-11												1
Domestic	Copenhagen	BAe146	491	532	581	665	1034	1286	1078	1171	1032	934	607	171
Domestic	Copenhagen	Beech Super King Air 200B	3	12	5	9	8	7	2	5	7	10	7	4
Domestic	Copenhagen	Cessna 208 Caravan				1		2	1	1				
Domestic	Copenhagen	CRJ9	2	3	2	3	1	1		65	1890	2792	2596	1488
Domestic	Copenhagen	Dash8 400	2016	3849	4188	8107	6686	4152	2462		1	1		210
Domestic	Copenhagen	DC10-30				1		1						
Domestic	Copenhagen	DC9	113	5										
Domestic	Copenhagen	Dornier 328-110						1						
Domestic	Copenhagen	F100					1		39	10				
Domestic	Copenhagen	F28									2			
Domestic	Copenhagen	F50	292	167	20	3	7	1	54	74				108
Domestic	Copenhagen	MD 82	4498	3131	1571	469	1345	1783	2686	2974	2130	1161	941	1646
Domestic	Copenhagen	Reims F406 Caravan II	2	2	8	11	6	3	1	1	1			1
Domestic	Copenhagen	RJ 100	2318	1048	325	327	560	882	1674	1802	1531	1472	1925	1062
Domestic	Copenhagen	S2000	19	10										
Domestic	Copenhagen	S61		1	1	8	3	3	3		4	15	16	12
Domestic	Copenhagen	Shorts 330	7											

Domestic	Copenhagen	Shorts 360 300	948	525	471	378	431	453	19	8	13	7	15	13
Domestic	Copenhagen	Swearingen Metro III	29	27	29	14	13	19	31	10	6	15	29	26
Domestic	Copenhagen	Saab 340B		6	4		16		15	93	372	313		

Flight	Airport name	Rep Aircraft	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
International	Copenhagen	A310	2488	1900	1179	1081	1142	1318	1181	1164	747	614	711	464
International	Copenhagen	A320	3895	7851	11850	17052	16184	18835	21042	24157	22594	24778	27104	28153
International	Copenhagen	A330	363	306	692	804	783	884	854	818	803	841	1043	881
International	Copenhagen	A340	456	1807	1845	2049	2028	1939	1752	1756	1488	1436	1696	1895
International	Copenhagen	Antonov 26	702	516	517	521	552	549	592	542	416	269	254	253
International	Copenhagen	ATR 42-320	2817	1097	1226	666	312	196	1020	821	1160	802	390	326
International	Copenhagen	ATR 72-200	1311	1059	1235	791	571	461	434	651	291	273	712	856
International	Copenhagen	B727	2051	1143	109	2	1	1	1		4	1		1
International	Copenhagen	B737 100						4	219	288	345	448	1557	4955
International	Copenhagen	B737 400	29665	25656	27987	25883	24782	19369	20690	25053	22285	25416	24361	22613
International	Copenhagen	B747 400	718	556	612	726	900	1084	1055	970	922	872	884	823
International	Copenhagen	B757	1701	2062	2285	2189	2011	2082	2625	2228	1867	1840	1644	1369
International	Copenhagen	B767 300 ER	3026	1103	546	91	151	285	414	678	639	731	670	643
International	Copenhagen	B777	40	266	150	157	168	171	242	264	267	394	431	764
International	Copenhagen	BAC1-11	1	1	5		5	4	3	5	1	1	1	1
International	Copenhagen	BAe146	4510	5849	5131	3878	4540	4098	3723	7660	3202	2280	1077	792
International	Copenhagen	Beech Super King Air 200B	13	12	16	16	48	37	60	37	37	30	54	40
International	Copenhagen	Cessna 208 Caravan	6	1		1	4	5	6	10	1	3	8	22
International	Copenhagen	CRJ9	56	48	43	70	443	1054	1398	1451	7235	12981	13811	15662
International	Copenhagen	Dash8 400	8122	10809	13457	14213	13972	14831	11580	630	1620	2071	3157	3671
International	Copenhagen	DC10-30	147	51	154	157	151	69	131	158	148	28	2	4
International	Copenhagen	DC9	5424	277	91	6	15	3	10	2	27	3		
International	Copenhagen	Dornier 328-110		3	6	9		1		1	2	1	1	
International	Copenhagen	Embraer 110P2A												2
International	Copenhagen	F100	625	464	6	307	666	664	750	1250	626	447	389	450
International	Copenhagen	F28	1433	832	716	727	554	648	390	539	430	128	43	83
International	Copenhagen	F50	6511	3335	6075	5107	4292	3268	2901	2634	794	679	767	193
International	Copenhagen	MD 82	32740	32219	23211	28009	28432	26979	24648	22120	15547	11841	13432	12890
International	Copenhagen	Reims F406 Caravan II	6	19	16	12	23	17	24	19	8	8	2	3
International	Copenhagen	RJ 100	5925	6637	7266	8647	8941	9060	9934	13795	12301	9836	9446	7251

International	Copenhagen	S2000	386	1029	346	496	1426	331	33	2	4	203	559	317
International	Copenhagen	S61	3541	3121	3	1	1	2	3	1	4		2	1
International	Copenhagen	Shorts 330	125			1								
International	Copenhagen	Shorts 360 300	545	89	154	137	280	63	73	224	201	157	25	27
International	Copenhagen	Swearingen Metro III	723	963	943	453	459	462	488	468	453	181	12	15
International	Copenhagen	Saab 340B	71	801	1145	1670	509	21	265	695	843	1303	738	1309

Flight	Airport name	Rep Aircraft	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
Domestic	Other airports	A310	31	3			6		1					
Domestic	Other airports	A320	115	126	98	38	156	357	342	573	552	541	635	719
Domestic	Other airports	A330	9	5	2	7	4		2	1	3	1	1	2
Domestic	Other airports	A340	6	2	1		1							1
Domestic	Other airports	Antonov 26		1		83	274	249	254	252	252	63		
Domestic	Other airports	ATR 42-320	3182	4143	5143	3189	1773	3966	3714	3875	2579	3289	3588	1714
Domestic	Other airports	ATR 72-200	2342	2751	4629	5446	7368	5649	5324	6082	5506	6103	7369	4250
Domestic	Other airports	B727						1			1			
Domestic	Other airports	B737 400	2754	1755	2236	798	505	501	1295	1443	2246	3500	3075	3144
Domestic	Other airports	B747 400		1										
Domestic	Other airports	B757	46	41	50	43	16	17	21	9	4	2	2	
Domestic	Other airports	B767 300 ER	3	6	7		1		3	19	19	1	2	1
Domestic	Other airports	BAe Jetstream 31	249	328	349	331	626	699	582	331	147	89	33	51
Domestic	Other airports	BAe Jetstream 41	46	67	43	49	7		1				1	
Domestic	Other airports	BAe146	46	60	62	100	231	261	259	281	173	94	110	80
Domestic	Other airports	Beech 1900C Airliner	135	370	668	928	651	35	5	3	1			
Domestic	Other airports	Beech Super King Air 200B	194	155	245	241	218	231	153	118	80	66	81	78
Domestic	Other airports	Beech Super King Air 350	18	2	6	7	3	1	86	46	11	9	11	28
Domestic	Other airports	Cessna 208 Caravan	11	24	58	86	98	155	101	129	104	75	106	72
Domestic	Other airports	CRJ9								49	1899	2792	2600	1474
Domestic	Other airports	Dash8 400	2038	3828	4192	8105	6705	4157	2462					235
Domestic	Other airports	DC10-30				3								
Domestic	Other airports	DC9	113	6										
Domestic	Other airports	De Havilland DHC-3 Turbo-Otter				1	2							
Domestic	Other airports	Dornier 328-110				2		1	1					1
Domestic	Other airports	Embraer 110P2A	132	118	455	371	457	638	20	47	30	36	43	24
Domestic	Other airports	F100							37	1				
Domestic	Other airports	F50	140	183	9	2	2	1	53	69	4	1		105

Domestic	Other airports	Fokker 27 Friendship	63	1			1	3		8				
Domestic	Other airports	Lockheed C-130H Hercules	17	12	13	46	54	27	46	38	44	69	49	45
Domestic	Other airports	MD 82	4505	3140	1567	454	1358	1782	2692	3033	2155	1265	1148	1803
Domestic	Other airports	Reims F406 Caravan II	264	298	262	159	134	68	109	71	53	21	12	13
Domestic	Other airports	RJ 100	3160	2387	1930	1618	1107	1639	2718	2754	2403	2235	2379	1738
Domestic	Other airports	S2000	93	91	86	41	26	18	2	1				3
Domestic	Other airports	S61	220	3018	4452	4432	4209	4760	5451	4744	4995	4562	4272	4525
Domestic	Other airports	Shorts 330	7											
Domestic	Other airports	Shorts 360 300	389	207	144	63	145	131	317	465	530	266	39	47
Domestic	Other airports	Shorts SC.7 Srs3M-200	173			1		6	4				1	
Domestic	Other airports	Swearingen Metro III	135	155	263	97	124	211	172	89	93	67	155	173
Domestic	Other airports	Saab 340B		510	389		401	892	925	1015	973	888	167	3

Flight	Airport name	Rep Aircraft	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
International	Other airports	A310	151	19	28	20	9	12	13	4	1	2	6	1
International	Other airports	A320	937	1004	834	849	924	1192	1090	1156	1038	1307	1413	1422
International	Other airports	A330	60	11	3	13	3	7	10	5	2	13	17	12
International	Other airports	A340				2	3							
International	Other airports	Antonov 26	2	2		2	12	11	42	17	18	11	4	2
International	Other airports	ATR 42-320	161	242	403	527	1122	715	463	122	109	415	728	616
International	Other airports	ATR 72-200	15	45	82	46	140	264	363	458	431	451	356	218
International	Other airports	B727	82	90	77	26	26	46	1					
International	Other airports	B737 100	6					7	2		2	252	868	931
International	Other airports	B737 400	6906	6492	6680	6839	6734	4575	5592	6866	6179	7698	7544	7321
International	Other airports	B747 100-300		1		2	10		1		2	5	3	1
International	Other airports	B747 400	2	7	10	16	10	5	15	8	8	6	7	10
International	Other airports	B757	107	137	188	150	79	114	88	64	70	266	316	318
International	Other airports	B767 300 ER	48	71	55	69	37	15	19	39	52	51	43	52
International	Other airports	BAC1-11		1	2	2	2	1						
International	Other airports	BAe Jetstream 31	1885	1802	2124	2718	2298	1811	1099	792	876	757	681	992
International	Other airports	BAe Jetstream 41	739	905	983	689	118	2		5	2	3	2	1
International	Other airports	BAe146	284	229	414	229	335	538	506	974	979	225	186	165
International	Other airports	Beech 1900C Airliner	92	1083	579	548	441	32	8	7	6	4	7	3
International	Other airports	Beech Super King Air 200B	89	123	281	288	339	404	349	361	231	203	177	207
International	Other airports	Beech Super King Air 350	162	28	26	22	34	22	30	38	36	52	33	42
International	Other airports	Cessna 208 Caravan	27	33	164	201	208	227	202	391	360	180	162	142

International	Other airports	CRJ9						443	874	261		1	8	6
International	Other airports	Dash8 400	19	147	498	68	97	62	38	31	43	78	174	282
International	Other airports	DC10-30		1	1	1	6	3		1		1		4
International	Other airports	DC9		1	3	6			2		1			
International	Other airports	De Havilland DHC-3 Turbo-Otter			5	2	2		3					
International	Other airports	Dornier 328-110	1	3	7	6	7	9	12	8	26	10	6	5
International	Other airports	Embraer 110P2A	43	24	127	23	18	68	46	94	78	83	110	109
International	Other airports	F100	10		1	2	3	751	838	150	64	62	53	111
International	Other airports	F28						7	254	257	228	261	577	1200
International	Other airports	F50	241	164	59	2	7	38	5	44	300	48	72	10
International	Other airports	Fokker 27 Friendship	551	359	4	1	10	150	5	3	1		1	
International	Other airports	Lockheed C-130H Hercules	4	1	4	4	7	13	8	5	6	3	5	8
International	Other airports	MD 82	141	168	140	227	461	513	979	963	704	411	340	277
International	Other airports	Reims F406 Caravan II	195	410	394	267	268	197	254	131	94	45	34	39
International	Other airports	RJ 100	2740	3047	4544	5980	4083	4827	5706	6999	5866	7296	8466	7173
International	Other airports	S2000	430	472	651	760	811	101	10	14	3	31	119	239
International	Other airports	S61	33	55	108	120	106	163	168	136	104	95	94	92
International	Other airports	Shorts 330	12											
International	Other airports	Shorts 360 300	564	538	127	78	1680	2894	3074	2264	2044	1592	985	1405
International	Other airports	Shorts SC.7 Srs3M-200			5	4	5	7	1	3				
International	Other airports	Swearingen Metro III	290	309	328	290	374	453	481	427	249	306	341	240
International	Other airports	Saab 340B	6	56	112	11	222	713	637	790	407	312	97	66

No. of flights between Danish airports and airports in Greenland and Faroe Islands

Area	Destination	Airport name	Distance NM	Rep Aircraft	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
Greenland	Narsarsuaq	Billund	1694,92	RJ 100											1	
Greenland	Narsarsuaq	Copenhagen	1796,98	B737 400	2	9	10	7	4	5		5	1		26	27
Greenland	Narsarsuaq	Copenhagen	1796,98	B757	68	73	65	63	61	66	77	72	50	39		
Greenland	Narsarsuaq	Copenhagen	1796,98	F50				1								
Greenland	Narsarsuaq	Copenhagen	1796,98	MD 82	4											
Greenland	Narsarsuaq	Copenhagen	1796,98	RJ 100					1	1	1			1	2	
Greenland	Narsarsuaq	Karup	1675,49	B737 400												1
Greenland	Narsarsuaq	Roskilde	1783,48	Lockheed C-130H Hercules										1		
Greenland	Narsarsuaq	Roskilde	1783,48	RJ 100								1				
Greenland	Narsarsuaq	Sønderborg	1739,20	RJ 100										1		1
Greenland	Narsarsuaq	Aalborg	1670,63	A320						1						

Greenland	Narsarsuaq	Aalborg	1670,63	B737 400						1	5	12	12	2	3	2
Greenland	Narsarsuaq	Aalborg	1670,63	B757			1	7	6	8	2					
Greenland	Narsarsuaq	Aalborg	1670,63	MD 82									2	11	14	12
Greenland	Narsarsuaq	Aalborg	1670,63	RJ 100				1								
Greenland	Narsarsuaq	Aarhus	1717,06	RJ 100	1											
Greenland	Søndre Strømfjord	Billund	1766,74	B737 400										1		
Greenland	Søndre Strømfjord	Billund	1766,74	MD 82										1		2
Greenland	Søndre Strømfjord	Billund	1766,74	RJ 100	1	1			1							1
Greenland	Søndre Strømfjord	Copenhagen	1852,59	A310								1				
Greenland	Søndre Strømfjord	Copenhagen	1852,59	A320						47	44					1
Greenland	Søndre Strømfjord	Copenhagen	1852,59	A330		25	209	207	212	212	219	196	222	219	233	247
Greenland	Søndre Strømfjord	Copenhagen	1852,59	A340			1					1			2	
Greenland	Søndre Strømfjord	Copenhagen	1852,59	B737 400	1	12		1		34	3	2	6	3	31	25
Greenland	Søndre Strømfjord	Copenhagen	1852,59	B757	112	136	22	30	30	26	51	77	57	45	4	
Greenland	Søndre Strømfjord	Copenhagen	1852,59	B767 300 ER	191	167	8					8	1		4	3
Greenland	Søndre Strømfjord	Copenhagen	1852,59	DC10-30						1						
Greenland	Søndre Strømfjord	Copenhagen	1852,59	MD 82									1			
Greenland	Søndre Strømfjord	Copenhagen	1852,59	RJ 100		2	1	1	3	1	4	2	2		2	1
Greenland	Søndre Strømfjord	Copenhagen	1852,59	Shorts 360 300								1				
Greenland	Søndre Strømfjord	Roskilde	1842,33	F50								1				
Greenland	Søndre Strømfjord	Roskilde	1842,33	Lockheed C-130H Hercules							1		1			
Greenland	Søndre Strømfjord	Roskilde	1842,33	RJ 100				1					1			
Greenland	Søndre Strømfjord	Sønderborg	1815,87	RJ 100			13	6	6	4	1	13	11	13	19	11
Greenland	Søndre Strømfjord	Aalborg	1724,62	B737 400						1						
Greenland	Søndre Strømfjord	Aalborg	1724,62	RJ 100								4				
Greenland	Thule	Copenhagen	2084,23	A330				12	13	13	12	12				
Greenland	Thule	Copenhagen	2084,23	A340			4	1								
Greenland	Thule	Copenhagen	2084,23	B767 300 ER	12	13	11									
Greenland	Thule	Copenhagen	2084,23	MD 82	1											
Greenland	Thule	Copenhagen	2084,23	RJ 100					1							
Greenland	Thule	Roskilde	2078,83	RJ 100					1							
Greenland	Thule	Sønderborg	2081,53	RJ 100					1							
Faroe Islands	Vagar	Billund	632,83	A320												1
Faroe Islands	Vagar	Billund	632,83	ATR 42-320	2											
Faroe Islands	Vagar	Billund	632,83	ATR 72-200				1								
Faroe Islands	Vagar	Billund	632,83	B737 400	140	132	153	104		1						
Faroe Islands	Vagar	Billund	632,83	BAe146	34	46	56	97	208	214	215	162	119	57	79	20

[illegible]

LTO fuel consumption and emission factors per representative aircraft type for Copenhagen Airport and other airports.

Origin	Representative aircraft	Fuel kg_LTO	Fuel GJ_LTO	SO ₂ kg_LTO	NO _x kg_LTO	VOC kg_LTO	NM VOC kg_LTO	CH ₄ kg_LTO	CO kg_LTO	CO ₂ tons_LTO	N ₂ O kg_LTO	TSP kg_LTO
EKCH	A310	1200,971	52,242	1,201	21,747	2,417	2,175	0,242	11,518	3,761	0,100	0,070
EKCH	A320	609,300	26,505	0,609	9,940	1,595	1,436	0,160	11,029	1,908	0,100	0,070
EKCH	A330	1727,520	75,147	1,728	33,754	0,974	0,877	0,097	9,860	5,411	0,100	0,070
EKCH	A340	1573,488	68,447	1,573	33,462	8,487	7,638	0,849	22,757	4,928	0,100	0,070
EKCH	Antonov 26	143,310	6,234	0,143	0,202	7,559	6,803	0,756	10,907	0,449	0,100	0,070
EKCH	ATR 42-320	120,720	5,251	0,121	1,056	0,000	0,000	0,000	0,926	0,378	0,100	0,070
EKCH	ATR 72-200	144,130	6,270	0,144	1,514	0,000	0,000	0,000	0,775	0,451	0,100	0,070
EKCH	B727	1028,975	44,760	1,029	11,222	3,366	3,029	0,337	12,941	3,223	0,100	0,070
EKCH	B737 100	669,320	29,115	0,669	7,107	0,340	0,306	0,034	2,456	2,096	0,100	0,070
EKCH	B737 400	613,619	26,692	0,614	7,350	0,296	0,267	0,030	5,455	1,922	0,100	0,070
EKCH	B747 100-300	2603,373	113,247	2,603	53,265	16,181	14,563	1,618	34,464	8,154	0,100	0,070

EKCH	B747 400	2638,978	114,796	2,639	52,985	1,170	1,053	0,117	9,011	8,265	0,100	0,070
EKCH	B757	957,844	41,666	0,958	18,518	0,566	0,509	0,057	5,729	3,000	0,100	0,070
EKCH	B767 300 ER	1270,887	55,284	1,271	24,567	0,448	0,403	0,045	3,019	3,980	0,100	0,070
EKCH	B777	2022,840	87,994	2,023	50,760	10,356	9,320	1,036	27,738	6,336	0,100	0,070
EKCH	BAC1-11	474,566	20,644	0,475	4,466	9,648	8,684	0,965	17,460	1,486	0,100	0,070
EKCH	BAe Jetstream 31	47,110	2,049	0,047	0,381	0,048	0,043	0,005	0,551	0,148	0,100	0,070
EKCH	BAe Jetstream 41	64,920	2,824	0,065	0,483	0,096	0,087	0,010	0,884	0,203	0,100	0,070
EKCH	BAe146	422,117	18,362	0,422	3,590	0,528	0,475	0,053	4,714	1,322	0,100	0,070
EKCH	Beech 1900C Airliner	62,630	2,724	0,063	0,262	0,677	0,609	0,068	2,366	0,196	0,100	0,070
EKCH	Beech Super King Air 200B	54,170	2,356	0,054	0,251	0,140	0,126	0,014	0,814	0,170	0,100	0,070
EKCH	Beech Super King Air 350	60,770	2,643	0,061	0,252	0,251	0,226	0,025	2,001	0,190	0,100	0,070
EKCH	Cessna 208 Caravan	29,710	1,292	0,030	0,158	0,028	0,025	0,003	0,306	0,093	0,100	0,070
EKCH	CRJ9	365,221	15,887	0,365	3,877	0,020	0,018	0,002	2,028	1,144	0,100	0,070
EKCH	Dash8 400	124,022	5,395	0,124	0,884	0,605	0,545	0,061	1,432	0,388	0,100	0,070
EKCH	DC10-30	1836,099	79,870	1,836	39,603	10,300	9,270	1,030	27,670	5,751	0,100	0,070
EKCH	DC9	634,784	27,613	0,635	6,463	0,422	0,380	0,042	2,698	1,988	0,100	0,070
EKCH	De Havilland Dash 7	146,920	6,391	0,147	0,781	0,206	0,186	0,021	1,600	0,460	0,100	0,070
EKCH	De Havilland DHC-3 Turbo-Otter	32,400	1,409	0,032	0,177	0,018	0,016	0,002	0,284	0,101	0,100	0,070
EKCH	Dornier 328-110	130,990	5,698	0,131	1,246	0,000	0,000	0,000	0,757	0,410	0,100	0,070
EKCH	Embraer 110P2A	50,490	2,196	0,050	0,284	0,026	0,024	0,003	0,400	0,158	0,100	0,070
EKCH	F100	532,667	23,171	0,533	5,442	0,719	0,647	0,072	6,527	1,668	0,100	0,070
EKCH	F28	468,141	20,364	0,468	4,669	14,505	13,055	1,451	15,260	1,466	0,100	0,070
EKCH	F50	130,370	5,671	0,130	1,293	0,000	0,000	0,000	0,777	0,408	0,100	0,070
EKCH	Fokker 27 Friendship	169,480	7,372	0,169	0,346	1,862	1,676	0,186	8,035	0,531	0,100	0,070
EKCH	Lockheed C-130H Hercules	287,800	12,519	0,288	1,975	0,945	0,851	0,095	2,021	0,901	0,100	0,070
EKCH	Lockheed P-3B Orion	265,340	11,542	0,265	1,792	0,907	0,817	0,091	1,926	0,831	0,100	0,070
EKCH	MD 82	758,573	32,998	0,759	11,365	1,065	0,958	0,106	3,433	2,376	0,100	0,070
EKCH	Reims F406 Caravan II	42,010	1,827	0,042	0,216	0,040	0,036	0,004	0,475	0,132	0,100	0,070
EKCH	RJ 100	161,020	7,004	0,161	1,122	0,236	0,213	0,024	2,542	0,504	0,100	0,070
EKCH	S2000	103,169	4,488	0,103	0,593	0,038	0,034	0,004	0,734	0,323	0,100	0,070
EKCH	S61	48,676	2,117	0,049	0,385	0,028	0,025	0,003	0,378	0,152	0,100	0,070
EKCH	Shorts 330	73,080	3,179	0,073	0,389	0,126	0,113	0,013	0,851	0,229	0,100	0,070
EKCH	Shorts 360 300	86,790	3,775	0,087	0,412	0,738	0,664	0,074	3,440	0,272	0,100	0,070
EKCH	Shorts SC.7 Srs3M-200	25,060	1,090	0,025	0,181	0,714	0,643	0,071	0,529	0,078	0,100	0,070
EKCH	Swearingen Metro III	47,650	2,073	0,048	0,390	0,047	0,043	0,005	0,544	0,149	0,100	0,070
EKCH	Saab 340B	78,190	3,401	0,078	0,510	0,238	0,214	0,024	0,456	0,245	0,100	0,070

Origin	Representative aircraft	Fuel kg_LTO	Fuel GJ_LTO	SO ₂ kg_LTO	NO _x kg_LTO	VOC kg_LTO	NM VOC kg_LTO	CH ₄ kg_LTO	CO kg_LTO	CO ₂ tons_LTO	N ₂ O kg_LTO	TSP kg_LTO
Other airports	A310	1065,140	46,334	1,065	21,167	1,166	1,050	0,117	5,789	3,336	0,100	0,070
Other airports	A320	532,087	23,146	0,532	9,582	1,464	1,317	0,146	8,403	1,666	0,100	0,070
Other airports	A330	1525,920	66,378	1,526	32,805	0,519	0,467	0,052	5,204	4,779	0,100	0,070
Other airports	A340	1394,928	60,679	1,395	32,697	4,380	3,942	0,438	11,634	4,369	0,100	0,070
Other airports	Antonov 26	105,450	4,587	0,105	0,164	3,701	3,331	0,370	5,980	0,330	0,100	0,070
Other airports	ATR 42-320	89,400	3,889	0,089	0,849	0,000	0,000	0,000	0,557	0,280	0,100	0,070
Other airports	ATR 72-200	107,950	4,696	0,108	1,243	0,000	0,000	0,000	0,475	0,338	0,100	0,070
Other airports	B727	875,431	38,081	0,875	10,682	1,832	1,649	0,183	7,569	2,742	0,100	0,070
Other airports	B737 100	569,168	24,759	0,569	6,760	0,244	0,220	0,024	1,511	1,783	0,100	0,070
Other airports	B737 400	528,911	23,008	0,529	6,988	0,148	0,133	0,015	2,905	1,657	0,100	0,070
Other airports	B747 100-300	2279,174	99,144	2,279	52,194	7,752	6,976	0,775	16,956	7,138	0,100	0,070
Other airports	B747 400	2333,706	101,516	2,334	51,524	0,899	0,809	0,090	4,817	7,309	0,100	0,070
Other airports	B757	839,780	36,530	0,840	18,033	0,299	0,269	0,030	3,002	2,630	0,100	0,070
Other airports	B767 300 ER	1132,405	49,260	1,132	23,981	0,275	0,248	0,028	1,796	3,547	0,100	0,070
Other airports	B777	1806,840	78,598	1,807	49,609	5,389	4,850	0,539	14,282	5,659	0,100	0,070
Other airports	BAC1-11	391,766	17,042	0,392	4,280	4,950	4,455	0,495	9,347	1,227	0,100	0,070
Other airports	BAe Jetstream 31	36,250	1,577	0,036	0,330	0,027	0,024	0,003	0,317	0,114	0,100	0,070
Other airports	BAe Jetstream 41	48,600	2,114	0,049	0,401	0,049	0,044	0,005	0,484	0,152	0,100	0,070
Other airports	BAe146	363,161	15,798	0,363	3,348	0,334	0,301	0,033	2,723	1,137	0,100	0,070
Other airports	Beech 1900C Airliner	47,450	2,064	0,047	0,218	0,364	0,328	0,036	1,409	0,149	0,100	0,070
Other airports	Beech Super King Air 200B	42,350	1,842	0,042	0,212	0,066	0,059	0,007	0,472	0,133	0,100	0,070
Other airports	Beech Super King Air 350	47,150	2,051	0,047	0,210	0,125	0,112	0,012	1,214	0,148	0,100	0,070
Other airports	Cessna 208 Caravan	24,250	1,055	0,024	0,138	0,014	0,012	0,001	0,168	0,076	0,100	0,070
Other airports	CRJ9	318,925	13,873	0,319	3,664	0,014	0,013	0,001	1,184	0,999	0,100	0,070
Other airports	Dash8 400	78,842	3,430	0,079	0,712	0,302	0,272	0,030	0,732	0,247	0,100	0,070
Other airports	DC10-30	1618,066	70,386	1,618	38,762	5,286	4,757	0,529	14,088	5,068	0,100	0,070
Other airports	DC9	538,259	23,414	0,538	6,142	0,281	0,253	0,028	1,636	1,686	0,100	0,070
Other airports	De Havilland Dash 7	115,600	5,029	0,116	0,668	0,097	0,087	0,010	0,909	0,362	0,100	0,070
Other airports	De Havilland DHC-3 Turbo-Otter	26,400	1,148	0,026	0,153	0,008	0,008	0,001	0,156	0,083	0,100	0,070
Other airports	Dornier 328-110	93,850	4,082	0,094	0,968	0,000	0,000	0,000	0,456	0,294	0,100	0,070
Other airports	Embraer 110P2A	40,350	1,755	0,040	0,239	0,014	0,013	0,001	0,227	0,126	0,100	0,070
Other airports	F100	447,982	19,487	0,448	5,301	0,440	0,396	0,044	3,666	1,403	0,100	0,070
Other airports	F28	388,971	16,920	0,389	4,460	7,163	6,447	0,716	8,274	1,218	0,100	0,070
Other airports	F50	95,750	4,165	0,096	1,043	0,000	0,000	0,000	0,466	0,300	0,100	0,070
Other airports	Fokker 27 Friendship	132,400	5,759	0,132	0,320	0,977	0,879	0,098	4,649	0,415	0,100	0,070
Other airports	Lockheed C-130H Hercules	214,000	9,309	0,214	1,555	0,501	0,450	0,050	1,170	0,670	0,100	0,070

Other airports	Lockheed P-3B Orion	194,300	8,452	0,194	1,389	0,479	0,431	0,048	1,107	0,609	0,100	0,070
Other airports	MD 82	660,780	28,744	0,661	10,974	0,725	0,652	0,072	2,198	2,070	0,100	0,070
Other airports	Reims F406 Caravan II	32,950	1,433	0,033	0,180	0,021	0,019	0,002	0,270	0,103	0,100	0,070
Other airports	RJ 100	130,900	5,694	0,131	1,007	0,117	0,105	0,012	1,257	0,410	0,100	0,070
Other airports	S2000	64,769	2,817	0,065	0,439	0,018	0,016	0,002	0,388	0,203	0,100	0,070
Other airports	S61	48,676	2,117	0,049	0,385	0,028	0,025	0,003	0,378	0,152	0,100	0,070
Other airports	Shorts 330	58,200	2,532	0,058	0,337	0,059	0,053	0,006	0,482	0,182	0,100	0,070
Other airports	Shorts 360 300	67,650	2,943	0,068	0,354	0,380	0,342	0,038	1,904	0,212	0,100	0,070
Other airports	Shorts SC.7 Srs3M-200	21,700	0,944	0,022	0,173	0,349	0,314	0,035	0,333	0,068	0,100	0,070
Other airports	Swearingen Metro III	37,150	1,616	0,037	0,341	0,027	0,024	0,003	0,319	0,116	0,100	0,070
Other airports	Saab 340B	58,450	2,543	0,058	0,448	0,151	0,136	0,015	0,278	0,183	0,100	0,070

Total distance flown (NM) and average cruise fuel consumption and emission factors per representative aircraft type for cruise flying.

Airport name	Flight	Rep Aircraft	NM total	Fuel kg_NM	Fuel GJ_NM	SO ₂ g_NM	NO _x g_NM	VOC g_NM	NMVOC g_NM	CH ₄ g_NM	CO g_NM	CO ₂ kg_NM	N ₂ O g_NM	TSP g_NM
Copenhagen	Domestic	A320	74250,652	6,738	0,293	6,738	136,499	1,188	1,188	0	8,703	21,103	0,673	1,347
Copenhagen	Domestic	A330	494,598	14,886	0,647	14,886	415,692	16,039	16,039	0	32,448	46,624	1,488	2,977
Copenhagen	Domestic	A340	118,79	14,504	0,63	14,504	338,899	75,634	75,634	0	73,594	45,427	1,45	2,9
Copenhagen	Domestic	ATR 42-320	91366,085	1,746	0,075	1,746	15,257	0	0	0	17,36	5,469	0,174	0,349
Copenhagen	Domestic	ATR 72-200	423916,655	1,716	0,074	1,716	19,469	0	0	0	11,367	5,377	0,171	0,343
Copenhagen	Domestic	B737 400	381174,245	6,204	0,269	6,204	75,157	1,197	1,197	0	19,244	19,433	0,62	1,24
Copenhagen	Domestic	B767 300 ER	128,509	11,275	0,49	11,275	207,091	1,957	1,957	0	28,886	35,314	1,127	2,255
Copenhagen	Domestic	BAC1-11	118,79	5,698	0,247	5,698	78,988	1,409	1,409	0	11,393	17,846	0,569	1,139
Copenhagen	Domestic	BAe146	2453,552	5,399	0,234	5,399	69,36	2,812	2,812	0	11,466	16,91	0,539	1,079
Copenhagen	Domestic	Beech Super King Air 200B	234,339	0,789	0,034	0,789	3,832	2,352	2,352	0	14,024	2,473	0,078	0,157
Copenhagen	Domestic	CRJ9	159381,294	3,856	0,167	3,856	34,358	0,316	0,316	0	7,585	12,078	0,385	0,771
Copenhagen	Domestic	Dash8 400	20136,482	3,393	0,147	3,393	54,08	6,105	6,105	0	15,746	10,628	0,339	0,678
Copenhagen	Domestic	F50	11718,078	2,426	0,105	2,426	33,005	0	0	0	14,817	7,598	0,242	0,485
Copenhagen	Domestic	MD 82	182131,724	8,784	0,382	8,784	149,648	4,801	4,801	0	14,437	27,513	0,878	1,756
Copenhagen	Domestic	Reims F406 Caravan II	125,809	0,584	0,025	0,584	3,148	0,425	0,425	0	5,53	1,831	0,058	0,116
Copenhagen	Domestic	RJ 100	93295,911	2,954	0,128	2,954	29,167	2,361	2,361	0	22,389	9,251	0,295	0,59
Copenhagen	Domestic	S61	739,733	3,521	0,153	3,521	27,816	2,006	2,006	0	27,358	11,027	0,352	0,704
Copenhagen	Domestic	Shorts 360 300	711,658	1,612	0,07	1,612	9,34	7,665	7,665	0	40,802	5,05	0,161	0,322
Copenhagen	Domestic	Swearingen Metro III	3156,029	0,811	0,035	0,811	8,041	0,455	0,455	0	5,751	2,541	0,081	0,162
Copenhagen	International	A310	211613,648	8,994	0,391	8,994	142,038	1,563	1,563	0	7,885	28,171	0,899	1,798
Copenhagen	International	A320	17826903,56	5,477	0,238	5,477	80,851	0,956	0,956	0	5,311	17,154	0,547	1,095
Copenhagen	International	A330	2469946,022	11,97	0,52	11,97	158,439	11,473	11,473	0	16,245	37,49	1,197	2,394
Copenhagen	International	A340	7723021,202	12,795	0,556	12,795	209,498	10,021	10,021	0	13,542	40,076	1,279	2,559

Copenhagen	International	Antonov 26	76482,487	2,71	0,117	2,71	3,578	78,872	78,872	0	174,031	8,49	0,271	0,542
Copenhagen	International	ATR 42-320	96729,373	1,637	0,071	1,637	13,584	0	0	0	15,58	5,129	0,163	0,327
Copenhagen	International	ATR 72-200	301476,455	1,716	0,074	1,716	17,289	0	0	0	9,898	5,376	0,171	0,343
Copenhagen	International	B727	478,401	8,534	0,371	8,534	85,335	4,811	4,811	0	17,935	26,728	0,853	1,706
Copenhagen	International	B737 100	2391300,309	5,637	0,245	5,637	52,975	4,651	4,651	0	11,487	17,657	0,563	1,127
Copenhagen	International	B737 400	14477447,2	5,601	0,243	5,601	54,983	0,553	0,553	0	10	17,544	0,56	1,12
Copenhagen	International	B747 400	3199954,205	19,541	0,85	19,541	276,128	4,858	4,858	0	18,904	61,204	1,954	3,908
Copenhagen	International	B757	1861882,29	7,158	0,311	7,158	101,011	6,93	6,93	0	10,198	22,419	0,715	1,431
Copenhagen	International	B767 300 ER	1997777,181	9,718	0,422	9,718	127,81	4,748	4,748	0	10,992	30,437	0,971	1,943
Copenhagen	International	B777	2324122,056	13,908	0,605	13,908	218,791	13,28	13,28	0	17,285	43,56	1,39	2,781
Copenhagen	International	BAC1-11	1727,321	4,692	0,204	4,692	50,109	0,687	0,687	0	4,188	14,696	0,469	0,938
Copenhagen	International	BAe146	404392,758	5,107	0,222	5,107	40,203	1,924	1,924	0	6,728	15,995	0,51	1,021
Copenhagen	International	Beech Super King Air 200B	11013,476	0,751	0,032	0,751	3,119	3,342	3,342	0	18,848	2,353	0,075	0,15
Copenhagen	International	Cessna 208 Caravan	8748,908	0,55	0,023	0,55	3,189	0,066	0,066	0	1,971	1,725	0,055	0,11
Copenhagen	International	CRJ9	6143050,52	3,5	0,152	3,5	26,672	0,193	0,193	0	4,896	10,962	0,35	0,7
Copenhagen	International	Dash8 400	1079127,34	3,154	0,137	3,154	42,844	5,818	5,818	0	16,527	9,881	0,315	0,63
Copenhagen	International	DC10-30	4507,017	15,78	0,686	15,78	286,541	24,183	24,183	0	25,722	49,426	1,578	3,156
Copenhagen	International	Embraer 110P2A	981,1	0,923	0,04	0,923	5,345	0,199	0,199	0	3,571	2,893	0,092	0,184
Copenhagen	International	F100	222262,376	4,932	0,214	4,932	43,838	1,997	1,997	0	7,46	15,447	0,493	0,986
Copenhagen	International	F28	30325,554	4,731	0,205	4,731	50,654	9,729	9,729	0	9,592	14,817	0,473	0,946
Copenhagen	International	F50	45742,344	2,238	0,097	2,238	28,121	0	0	0	12,117	7,01	0,223	0,447
Copenhagen	International	MD 82	5617566,36	7,245	0,315	7,245	100,906	3,7	3,7	0	10,946	22,691	0,724	1,449
Copenhagen	International	Reims F406 Caravan II	974,081	0,582	0,025	0,582	2,855	0,683	0,683	0	8,114	1,825	0,058	0,116
Copenhagen	International	RJ 100	2919966,28	2,442	0,106	2,442	19,952	1,022	1,022	0	9,259	7,65	0,244	0,488
Copenhagen	International	S2000	132789,712	2,626	0,114	2,626	26,019	0,125	0,125	0	9,812	8,224	0,262	0,525
Copenhagen	International	S61	24,298	3,448	0,15	3,448	27,241	1,965	1,965	0	26,793	10,8	0,344	0,689
Copenhagen	International	Shorts 360 300	24683,032	1,463	0,063	1,463	7,181	4,926	4,926	0	34,542	4,584	0,146	0,292
Copenhagen	International	Swearingen Metro III	13392,54	0,793	0,034	0,793	7,862	0,497	0,497	0	6,958	2,484	0,079	0,158
Copenhagen	International	Saab 340B	268100,259	1,429	0,062	1,429	14,535	3,475	3,475	0	6,666	4,477	0,142	0,285
Other airports	Domestic	A320	80656,683	6,738	0,293	6,738	136,477	1,188	1,188	0	8,701	21,103	0,673	1,347
Other airports	Domestic	A330	237,58	14,897	0,648	14,897	416,925	16,044	16,044	0	32,433	46,657	1,489	2,979
Other airports	Domestic	A340	118,79	14,504	0,63	14,504	338,899	75,634	75,634	0	73,594	45,427	1,45	2,9
Other airports	Domestic	ATR 42-320	116993,431	1,745	0,075	1,745	15,252	0	0	0	17,355	5,468	0,174	0,349
Other airports	Domestic	ATR 72-200	418710,972	1,716	0,074	1,716	19,47	0	0	0	11,368	5,377	0,171	0,343
Other airports	Domestic	B737 400	387206,073	6,205	0,269	6,205	75,171	1,198	1,198	0	19,247	19,434	0,62	1,241
Other airports	Domestic	B767 300 ER	128,509	11,275	0,49	11,275	207,091	1,957	1,957	0	28,886	35,314	1,127	2,255
Other airports	Domestic	BAe Jetstream 31	3328,813	1,034	0,044	1,034	10,272	0,62	0,62	0	7,992	3,239	0,103	0,206

Other airports	Domestic	BAe146	3267,254	5,376	0,233	5,376	67,651	2,768	2,768	0	11,273	16,838	0,537	1,075
Other airports	Domestic	Beech Super King Air 200B	7749,417	0,788	0,034	0,788	3,82	2,368	2,368	0	14,103	2,471	0,078	0,157
Other airports	Domestic	Beech Super King Air 350	2056,68	0,865	0,037	0,865	3,554	4,357	4,357	0	33,972	2,71	0,086	0,173
Other airports	Domestic	Cessna 208 Caravan	6779,67	0,518	0,022	0,518	3,015	0,096	0,096	0	2,098	1,622	0,051	0,103
Other airports	Domestic	CRJ9	158337,026	3,856	0,167	3,856	34,357	0,316	0,316	0	7,585	12,078	0,385	0,771
Other airports	Domestic	Dash8 400	20791,441	3,393	0,147	3,393	54,08	6,105	6,105	0	15,746	10,628	0,339	0,678
Other airports	Domestic	Dornier 328-110	81,533	1,47	0,063	1,47	13,975	0	0	0	11,574	4,605	0,147	0,294
Other airports	Domestic	Embraer 110P2A	1510,784	0,846	0,036	0,846	5	0,183	0,183	0	3,276	2,65	0,084	0,169
Other airports	Domestic	F50	11374,667	2,426	0,105	2,426	33,003	0	0	0	14,816	7,598	0,242	0,485
Other airports	Domestic	Lockheed C-130H Hercules	5366,606	6,638	0,288	6,638	80,274	8,484	8,484	0	20,699	20,791	0,663	1,327
Other airports	Domestic	MD 82	196234,816	8,785	0,382	8,785	149,711	4,802	4,802	0	14,439	27,516	0,878	1,757
Other airports	Domestic	Reims F406 Caravan II	1041,029	0,584	0,025	0,584	3,151	0,423	0,423	0	5,506	1,831	0,058	0,116
Other airports	Domestic	RJ 100	147556,868	2,953	0,128	2,953	29,155	2,359	2,359	0	22,372	9,249	0,295	0,59
Other airports	Domestic	S2000	72,894	2,807	0,122	2,807	29,272	0,299	0,299	0	11,48	8,792	0,28	0,561
Other airports	Domestic	S61	481411,345	3,536	0,153	3,536	27,941	2,016	2,016	0	27,481	11,077	0,353	0,707
Other airports	Domestic	Shorts 360 300	2502,141	1,616	0,07	1,616	9,392	7,728	7,728	0	40,944	5,061	0,161	0,323
Other airports	Domestic	Swearingen Metro III	16364,914	0,811	0,035	0,811	8,044	0,455	0,455	0	5,735	2,542	0,081	0,162
Other airports	Domestic	Saab 340B	191,143	1,477	0,064	1,477	14,831	3,77	3,77	0	7,66	4,627	0,147	0,295
Other Airports	International	A310	1545,356	8,668	0,377	8,668	110,305	1,219	1,219	0	5,427	27,149	0,866	1,733
Other Airports	International	A320	2028122,379	5,084	0,221	5,084	67,477	0,884	0,884	0	4,191	15,923	0,508	1,016
Other Airports	International	A330	30263,493	12,459	0,541	12,459	170,214	11,908	11,908	0	16,681	39,021	1,245	2,491
Other Airports	International	Antonov 26	676,025	2,703	0,117	2,703	3,46	76,95	76,95	0	174,022	8,468	0,27	0,54
Other Airports	International	ATR 42-320	149127,674	1,656	0,072	1,656	13,87	0	0	0	15,884	5,187	0,165	0,331
Other Airports	International	ATR 72-200	69703,438	1,716	0,074	1,716	17,415	0	0	0	9,983	5,376	0,171	0,343
Other Airports	International	B737 100	282334,366	6,06	0,263	6,06	61,091	5,717	5,717	0	14,884	18,981	0,606	1,212
Other Airports	International	B737 400	6575224,668	5,593	0,243	5,593	53,685	0,469	0,469	0	8,842	17,52	0,559	1,118
Other Airports	International	B747 100-300	3110,691	21,107	0,918	21,107	366,338	6,182	6,182	0	19,092	66,11	2,11	4,221
Other Airports	International	B747 400	20271,053	18,845	0,819	18,845	264,175	6,356	6,356	0	22,75	59,024	1,884	3,769
Other Airports	International	B757	154359,99	7,689	0,334	7,689	145,989	7,743	7,743	0	13,061	24,084	0,768	1,537
Other Airports	International	B767 300 ER	110819,629	9,862	0,429	9,862	133,421	4,691	4,691	0	11,724	30,889	0,986	1,972
Other Airports	International	BAe Jetstream 31	245668,034	0,978	0,042	0,978	10,117	0,491	0,491	0	6,771	3,066	0,097	0,195
Other Airports	International	BAe Jetstream 41	743,52	1,355	0,058	1,355	13,563	0,418	0,418	0	7,1	4,246	0,135	0,271
Other Airports	International	BAe146	64998,853	5,129	0,223	5,129	43,716	2,035	2,035	0	7,381	16,066	0,512	1,025
Other Airports	International	Beech 1900C Airliner	1480,56	0,915	0,039	0,915	3,697	8,889	8,889	0	40,681	2,867	0,091	0,183
Other Airports	International	Beech Super King Air 200B	57431,313	0,752	0,032	0,752	3,129	3,329	3,329	0	18,785	2,355	0,075	0,15
Other Airports	International	Beech Super King Air 350	20978,917	0,831	0,036	0,831	3,102	4,059	4,059	0	34,91	2,602	0,083	0,166
Other Airports	International	Cessna 208 Caravan	65538,799	0,553	0,024	0,553	3,204	0,063	0,063	0	1,963	1,734	0,055	0,11

Other Airports	International	CRJ9	4008,636	3,482	0,151	3,482	25,192	0,144	0,144	0	3,901	10,907	0,348	0,696
Other Airports	International	Dash8 400	138736,982	3,084	0,134	3,084	39,518	5,733	5,733	0	16,761	9,661	0,308	0,616
Other Airports	International	DC10-30	8119,869	15,771	0,686	15,771	276,043	16,046	16,046	0	18,887	49,396	1,577	3,154
Other Airports	International	Dornier 328-110	3626,348	1,389	0,06	1,389	11,677	0	0	0	11,161	4,352	0,138	0,277
Other Airports	International	Embraer 110P2A	31467,005	0,905	0,039	0,905	5,266	0,196	0,196	0	3,504	2,837	0,09	0,181
Other Airports	International	F100	36292,094	5,146	0,223	5,146	50,806	2,32	2,32	0	9,293	16,118	0,514	1,029
Other Airports	International	F28	379250,424	4,785	0,208	4,785	51,621	10,639	10,639	0	10,857	14,989	0,478	0,957
Other Airports	International	F50	4612,308	2,136	0,092	2,136	25,468	0	0	0	10,645	6,69	0,213	0,427
Other Airports	International	Lockheed C-130H Hercules	4555,612	6,811	0,296	6,811	72,236	2,759	2,759	0	13,068	21,332	0,681	1,362
Other Airports	International	MD 82	341788,208	6,408	0,278	6,408	76,086	2,934	2,934	0	8,511	20,07	0,64	1,281
Other Airports	International	Reims F406 Caravan II	14881,194	0,582	0,025	0,582	2,825	0,708	0,708	0	8,374	1,824	0,058	0,116
Other Airports	International	RJ 100	3060615,06	2,427	0,105	2,427	19,675	0,982	0,982	0	8,861	7,602	0,242	0,485
Other Airports	International	S2000	43120,819	2,727	0,118	2,727	27,852	0,223	0,223	0	10,753	8,544	0,272	0,545
Other Airports	International	S61	19429,772	3,702	0,161	3,702	29,248	2,11	2,11	0	28,766	11,595	0,37	0,74
Other Airports	International	Shorts 360 300	477262,39	1,503	0,065	1,503	7,746	5,693	5,693	0	36,368	4,708	0,15	0,3
Other Airports	International	Swearingen Metro III	69133,787	0,799	0,034	0,799	7,921	0,483	0,483	0	6,541	2,504	0,079	0,159
Other Airports	International	Saab 340B	18340,692	1,409	0,061	1,409	14,416	3,354	3,354	0	6,257	4,415	0,14	0,281

Annex 3B-11 Basis fuel consumption and emission factors, deterioration factors, transient factors stock and activity data for non road working machinery and equipment, and recreational craft

Basis factors for diesel fuelled non road machinery.

Engine size [P=kW]	Emission Level	NO _x	VOC	CO	N ₂ O [g pr kWh]	NH ₃	TSP	Fuel
P<19	<1981	12.0	5.0	7	0.035	0.002	2.8	300
P<19	1981-1990	11.5	3.8	6	0.035	0.002	2.3	285
P<19	1991-Stage I	11.2	2.5	5	0.035	0.002	1.6	270
P<19	Stage I	11.2	2.5	5	0.035	0.002	1.6	270
P<19	Stage II	11.2	2.5	5	0.035	0.002	1.6	270
P<19	Stage IIIA	11.2	2.5	5	0.035	0.002	1.6	270
P<19	Stage IIIB	11.2	2.5	5	0.035	0.002	1.6	270
P<19	Stage IV	11.2	2.5	5	0.035	0.002	1.6	270
19<=P<37	<1981	18.0	2.5	6.5	0.035	0.002	2	300
19<=P<37	1981-1990	18.0	2.2	5.5	0.035	0.002	1.4	281
19<=P<37	1991-Stage I	9.8	1.8	4.5	0.035	0.002	1.4	262
19<=P<37	Stage I	9.8	1.8	4.5	0.035	0.002	1.4	262
19<=P<37	Stage II	6.5	0.6	2.2	0.035	0.002	0.4	262
19<=P<37	Stage IIIA	6.2	0.6	2.2	0.035	0.002	0.4	262
19<=P<37	Stage IIIB	6.2	0.6	2.2	0.035	0.002	0.4	262
19<=P<37	Stage IV	6.2	0.6	2.2	0.035	0.002	0.4	262
37<=P<56	<1981	7.7	2.4	6	0.035	0.002	1.8	290
37<=P<56	1981-1990	8.6	2.0	5.3	0.035	0.002	1.2	275
37<=P<56	1991-Stage I	11.5	1.5	4.5	0.035	0.002	0.8	260
37<=P<56	Stage I	7.7	0.6	2.2	0.035	0.002	0.4	260
37<=P<56	Stage II	5.5	0.4	2.2	0.035	0.002	0.2	260
37<=P<56	Stage IIIA	3.9	0.4	2.2	0.035	0.002	0.2	260
37<=P<56	Stage IIIB	3.9	0.4	2.2	0.035	0.002	0.0225	260
37<=P<56	Stage IV	3.9	0.4	2.2	0.035	0.002	0.0225	260
56<=P<75	<1981	7.7	2.0	5	0.035	0.002	1.4	290
56<=P<75	1981-1990	8.6	1.6	4.3	0.035	0.002	1	275
56<=P<75	1991-Stage I	11.5	1.2	3.5	0.035	0.002	0.4	260
56<=P<75	Stage I	7.7	0.4	1.5	0.035	0.002	0.2	260
56<=P<75	Stage II	5.5	0.3	1.5	0.035	0.002	0.2	260
56<=P<75	Stage IIIA	4.0	0.3	1.5	0.035	0.002	0.2	260
56<=P<75	Stage IIIB	3.0	0.2	1.5	0.035	0.002	0.0225	260
56<=P<75	Stage IV	0.4	0.2	1.5	0.035	0.002	0.0225	260
75<=P<130	<1981	10.5	2.0	5	0.035	0.002	1.4	280
75<=P<130	1981-1990	11.8	1.6	4.3	0.035	0.002	1	268
75<=P<130	1991-Stage I	13.3	1.2	3.5	0.035	0.002	0.4	255
75<=P<130	Stage I	8.1	0.4	1.5	0.035	0.002	0.2	255
75<=P<130	Stage II	5.2	0.3	1.5	0.035	0.002	0.2	255
75<=P<130	Stage IIIA	3.4	0.3	1.5	0.035	0.002	0.2	255
75<=P<130	Stage IIIB	3.0	0.2	1.5	0.035	0.002	0.0225	255
75<=P<130	Stage IV	0.4	0.2	1.5	0.035	0.002	0.0225	255
130<=P<560	<1981	17.8	1.5	2.5	0.035	0.002	0.9	270
130<=P<560	1981-1990	12.4	1.0	2.5	0.035	0.002	0.8	260
130<=P<560	1991-Stage I	11.2	0.5	2.5	0.035	0.002	0.4	250
130<=P<560	Stage I	7.6	0.3	1.5	0.035	0.002	0.2	250
130<=P<560	Stage II	5.2	0.3	1.5	0.035	0.002	0.1	250
130<=P<560	Stage IIIA	3.4	0.3	1.5	0.035	0.002	0.1	250
130<=P<560	Stage IIIB	3.0	0.2	1.5	0.035	0.002	0.0225	250
130<=P<560	Stage IV	0.4	0.2	1.5	0.035	0.002	0.0225	250

Basis factors for 4-stroke gasoline non road machinery.

Engine	Size code	Size classe [S=ccm]	Emission Level	NO _x	VOC	CO	N ₂ O [g pr kWh]	NH ₃	TSP	Fuel
4-stroke	SH2	20<=S<50	<1981	2.4	33	198	0.002	0.03	0.08	496
4-stroke	SH2	20<=S<50	1981-1990	3.5	27.5	165	0.002	0.03	0.08	474
4-stroke	SH2	20<=S<50	1991-Stage I	4.7	22	132	0.002	0.03	0.08	451
4-stroke	SH2	20<=S<50	Stage I	4.7	22	132	0.002	0.03	0.08	406
4-stroke	SH2	20<=S<50	Stage II	4.7	22	132	0.002	0.03	0.08	406
4-stroke	SH3	S>=50	<1981	2.4	33	198	0.002	0.03	0.08	496
4-stroke	SH3	S>=50	1981-1990	3.5	27.5	165	0.002	0.03	0.08	474
4-stroke	SH3	S>=50	1991-Stage I	4.7	22	132	0.002	0.03	0.08	451
4-stroke	SH3	S>=50	Stage I	4.7	22	132	0.002	0.03	0.08	406
4-stroke	SH3	S>=50	Stage II	4.7	22	132	0.002	0.03	0.08	406
4-stroke	SN1	S<66	<1981	1.2	26.9	822	0.002	0.03	0.08	603
4-stroke	SN1	S<66	1981-1990	1.8	22.5	685	0.002	0.03	0.08	603
4-stroke	SN1	S<66	1991-Stage I	2.4	18	548	0.002	0.03	0.08	603
4-stroke	SN1	S<66	Stage I	4.3	16.1	411	0.002	0.03	0.08	475
4-stroke	SN1	S<66	Stage II	4.3	16.1	411	0.002	0.03	0.08	475
4-stroke	SN2	66<=S<100	<1981	2.3	10.5	822	0.002	0.03	0.08	627
4-stroke	SN2	66<=S<100	1981-1990	3.5	8.7	685	0.002	0.03	0.08	599
4-stroke	SN2	66<=S<100	1991-Stage I	4.7	7	548	0.002	0.03	0.08	570
4-stroke	SN2	66<=S<100	Stage I	4.7	7	467	0.002	0.03	0.08	450
4-stroke	SN2	66<=S<100	Stage II	4.7	7	467	0.002	0.03	0.08	450
4-stroke	SN3	100<=S<225	<1981	2.6	19.1	525	0.002	0.03	0.08	601
4-stroke	SN3	100<=S<225	1981-1990	3.8	15.9	438	0.002	0.03	0.08	573
4-stroke	SN3	100<=S<225	1991-Stage I	5.1	12.7	350	0.002	0.03	0.08	546
4-stroke	SN3	100<=S<225	Stage I	5.1	11.6	350	0.002	0.03	0.08	546
4-stroke	SN3	100<=S<225	Stage II	5.1	9.4	350	0.002	0.03	0.08	546
4-stroke	SN4	S>=225	<1981	1.3	11.1	657	0.002	0.03	0.08	539
4-stroke	SN4	S>=225	1981-1990	2	9.3	548	0.002	0.03	0.08	514
4-stroke	SN4	S>=225	1991-Stage I	2.6	7.4	438	0.002	0.03	0.08	490
4-stroke	SN4	S>=225	Stage I	2.6	7.4	438	0.002	0.03	0.08	490
4-stroke	SN4	S>=225	Stage II	2.6	7.4	438	0.002	0.03	0.08	490

Basis factors for 2-stroke gasoline non road machinery.

Engine	Size code	Size classe [ccm]	Emission Level	NO _x	VOC	CO	N ₂ O [g pr kWh]	NH ₃	TSP	Fuel
2-stroke	SH2	20<=S<50	<1981	1	305	695	0.002	0.01	7	882
2-stroke	SH2	20<=S<50	1981-1990	1	300	579	0.002	0.01	5.3	809
2-stroke	SH2	20<=S<50	1991-Stage I	1.1	203	463	0.002	0.01	3.5	735
2-stroke	SH2	20<=S<50	Stage I	1.5	188	379	0.002	0.01	3.5	720
2-stroke	SH2	20<=S<50	Stage II	1.5	44	379	0.002	0.01	3.5	500
2-stroke	SH3	S>=50	<1981	1.1	189	510	0.002	0.01	3.6	665
2-stroke	SH3	S>=50	1981-1990	1.1	158	425	0.002	0.01	2.7	609
2-stroke	SH3	S>=50	1991-Stage I	1.2	126	340	0.002	0.01	1.8	554
2-stroke	SH3	S>=50	Stage I	2	126	340	0.002	0.01	1.8	529
2-stroke	SH3	S>=50	Stage II	1.2	64	340	0.002	0.01	1.8	500
2-stroke	SN1	S<66	<1981	0.5	155	418	0.002	0.01	2.6	652
2-stroke	SN1	S<66	1981-1990	0.5	155	418	0.002	0.01	2.6	652
2-stroke	SN1	S<66	1991-Stage I	0.5	155	418	0.002	0.01	2.6	652
2-stroke	SN1	S<66	Stage I	0.5	155	418	0.002	0.01	2.6	652
2-stroke	SN1	S<66	Stage II	0.5	155	418	0.002	0.01	2.6	652
2-stroke	SN2	66<=S<100	<1981	0.5	155	418	0.002	0.01	2.6	652
2-stroke	SN2	66<=S<100	1981-1990	0.5	155	418	0.002	0.01	2.6	652
2-stroke	SN2	66<=S<100	1991-Stage I	0.5	155	418	0.002	0.01	2.6	652
2-stroke	SN2	66<=S<100	Stage I	0.5	155	418	0.002	0.01	2.6	652
2-stroke	SN2	66<=S<100	Stage II	0.5	155	418	0.002	0.01	2.6	652
2-stroke	SN3	100<=S<225	<1981	0.5	155	418	0.002	0.01	2.6	652
2-stroke	SN3	100<=S<225	1981-1990	0.5	155	418	0.002	0.01	2.6	652
2-stroke	SN3	100<=S<225	1991-Stage I	0.5	155	418	0.002	0.01	2.6	652
2-stroke	SN3	100<=S<225	Stage I	0.5	155	418	0.002	0.01	2.6	652
2-stroke	SN3	100<=S<225	Stage II	0.5	155	418	0.002	0.01	2.6	652
2-stroke	SN4	S>=225	<1981	0.5	155	418	0.002	0.01	2.6	652
2-stroke	SN4	S>=225	1981-1990	0.5	155	418	0.002	0.01	2.6	652
2-stroke	SN4	S>=225	1991-Stage I	0.5	155	418	0.002	0.01	2.6	652
2-stroke	SN4	S>=225	Stage I	0.5	155	418	0.002	0.01	2.6	652
2-stroke	SN4	S>=225	Stage II	0.5	155	418	0.002	0.01	2.6	652

Fuel consumption and emission factors for LPG fork lifts.

NO _x	VOC	CO	NH ₃	N ₂ O	TSP	FC
[g pr kWh]	[g pr kWh]	[g pr kWh]	[g pr kWh]	[g pr kWh]	[g pr kWh]	[g pr kWh]
19	2.2	1.5	0.003	0.05	0.07	311

Fuel consumption and emission factors for All Terrain Vehicles (ATV's).

ATV type	NO _x	VOC	CO	NH ₃	N ₂ O	TSP	Fuel
	[g pr GJ]	[g pr GJ]	[g pr GJ]	[g pr GJ]	[g pr GJ]	[g pr GJ]	[kg pr hour]
Professional	108	1077	16306	2	2	32	1.125
Private	128	1527	22043	2	2	39	0.75

Fuel consumption and emission factors for recreational craft.

Fuel type	Vessel type	Engine	Engine type	Direktiv	Engine size	CO	VOC	N ₂ O	NH ₃	NO _x	TSP	Fuel
					[kW]	[g pr kWh]						
Gasoline	Other boats (< 20 ft)	Out board	2-stroke	2003/44	8	202.5	45.9	0.01	0.002	2	10	791
Gasoline	Other boats (< 20 ft)	Out board	2-stroke	Konv.	8	427	257.0	0.01	0.002	2	10	791
Gasoline	Other boats (< 20 ft)	Out board	4-stroke	2003/44	8	202.5	24.0	0.03	0.002	7	0.08	426
Gasoline	Other boats (< 20 ft)	Out board	4-stroke	Konv.	8	520	24.0	0.03	0.002	7	0.08	426
Gasoline	Yawls and cabin boats	Out board	2-stroke	2003/44	20	162	36.5	0.01	0.002	3	10	791
Gasoline	Yawls and cabin boats	Out board	2-stroke	Konv.	20	374	172.0	0.01	0.002	3	10	791
Gasoline	Yawls and cabin boats	Out board	4-stroke	2003/44	20	162	14.0	0.03	0.002	10	0.08	426
Gasoline	Yawls and cabin boats	Out board	4-stroke	Konv.	20	390	14.0	0.03	0.002	10	0.08	426
Gasoline	Sailing boats (< 26 ft)	Out board	2-stroke	2003/44	10	189	43.0	0.01	0.002	2	10	791
Gasoline	Sailing boats (< 26 ft)	Out board	2-stroke	Konv.	10	427	257.0	0.01	0.002	2	10	791
Gasoline	Sailing boats (< 26 ft)	Out board	4-stroke	2003/44	10	189	24.0	0.03	0.002	7	0.08	426
Gasoline	Sailing boats (< 26 ft)	Out board	4-stroke	Konv.	10	520	24.0	0.03	0.002	7	0.08	426
Gasoline	Speed boats	In board	4-stroke	2003/44	90	141	10.0	0.03	0.002	12	0.08	426
Gasoline	Speed boats	In board	4-stroke	Konv.	90	346	10.0	0.03	0.002	12	0.08	426
Gasoline	Speed boats	Out board	2-stroke	2003/44	50	145.8	31.8	0.01	0.002	3	10	791
Gasoline	Speed boats	Out board	2-stroke	Konv.	50	374	172.0	0.01	0.002	3	10	791
Gasoline	Speed boats	Out board	4-stroke	2003/44	50	145.8	14.0	0.03	0.002	10	0.08	426
Gasoline	Speed boats	Out board	4-stroke	Konv.	50	390	14.0	0.03	0.002	10	0.08	426
Gasoline	Water scooters	Built in	2-stroke	2003/44	45	147	32.2	0.01	0.002	3	10	791
Gasoline	Water scooters	Built in	2-stroke	Konv.	45	374	172.0	0.01	0.002	3	10	791
Gasoline	Water scooters	Built in	4-stroke	2003/44	45	147	14.0	0.03	0.002	10	0.08	426
Gasoline	Water scooters	Built in	4-stroke	Konv.	45	390	14.0	0.03	0.002	10	0.08	426
Diesel	Motor boats (27-34 ft)	In board		2003/44	150	5	1.7	0.035	0.002	8.6	1	275
Diesel	Motor boats (27-34 ft)	In board		Konv.	150	5.3	2.0	0.035	0.002	8.6	1.2	275
Diesel	Motor boats (> 34 ft)	In board		2003/44	250	5	1.6	0.035	0.002	8.6	1	275
Diesel	Motor boats (> 34 ft)	In board		Konv.	250	5.3	2.0	0.035	0.002	8.6	1.2	275
Diesel	Motor boats (< 27 ft)	In board		2003/44	40	5	1.8	0.035	0.002	9.8	1	281
Diesel	Motor boats (< 27 ft)	In board		Konv.	40	5.5	2.2	0.035	0.002	18	1.4	281
Diesel	Motor sailors	In board		2003/44	30	5	1.9	0.035	0.002	9.8	1	281
Diesel	Motor sailors	In board		Konv.	30	5.5	2.2	0.035	0.002	18	1.4	281
Diesel	Sailing boats (> 26 ft)	In board		2003/44	30	5	1.9	0.035	0.002	9.8	1	281
Diesel	Sailing boats (> 26 ft)	In board		Konv.	30	5.5	2.2	0.035	0.002	18	1.4	281

CH₄ shares of VOC for diesel, gasoline and LPG.

Fuel type	CH ₄ share of VOC
Diesel	0.016
Gasoline 4-stroke	0.1
Gasoline 2-stroke	0.009
LPG	0.05

Deterioration factors for diesel machinery.

Emission Level	NO _x	VOC	CO	TSP
<1981	0.024	0.047	0.185	0.473
1981-1990	0.024	0.047	0.185	0.473
1991-Stage I	0.024	0.047	0.185	0.473
Stage I	0.024	0.036	0.101	0.473
Stage II	0.009	0.034	0.101	0.473
Stage IIIA	0.008	0.027	0.151	0.473
Stage IIIB	0.008	0.027	0.151	0.473
Stage IV	0.008	0.027	0.151	0.473

Deterioration factors for gasoline 2-stroke machinery.

Engine	Size code	Size classe	Emission Level	NO _x	VOC	CO	TSP
2-stroke	SH2	20<=S<50	<1981	0	0.2	0.2	0
2-stroke	SH2	20<=S<50	1981-1990	0	0.2	0.2	0
2-stroke	SH2	20<=S<50	1991-Stage I	0	0.2	0.2	0
2-stroke	SH2	20<=S<50	Stage I	0	0.29	0.24	0
2-stroke	SH2	20<=S<50	Stage II	0	0.29	0.24	0
2-stroke	SH3	S>=50	<1981	-0.031	0.2	0.2	0
2-stroke	SH3	S>=50	1981-1990	-0.031	0.2	0.2	0
2-stroke	SH3	S>=50	1991-Stage I	-0.031	0.2	0.2	0
2-stroke	SH3	S>=50	Stage I	0	0.266	0.231	0
2-stroke	SH3	S>=50	Stage II	0	0.266	0.231	0
2-stroke	SN1	S<66	<1981	-0.6	0.201	0.9	1.1
2-stroke	SN1	S<66	1981-1990	-0.6	0.201	0.9	1.1
2-stroke	SN1	S<66	1991-Stage I	-0.6	0.201	0.9	1.1
2-stroke	SN1	S<66	Stage I	-0.33	0.266	1.109	5.103
2-stroke	SN1	S<66	Stage II	-0.33	0	1.109	5.103
2-stroke	SN2	66<=S<100	<1981	-0.6	0.201	0.9	1.1
2-stroke	SN2	66<=S<100	1981-1990	-0.6	0.201	0.9	1.1
2-stroke	SN2	66<=S<100	1991-Stage I	-0.6	0.201	0.9	1.1
2-stroke	SN2	66<=S<100	Stage I	-0.33	0.266	1.109	5.103
2-stroke	SN2	66<=S<100	Stage II	-0.33	0	1.109	5.103
2-stroke	SN3	100<=S<225	<1981	-0.6	0.201	0.9	1.1
2-stroke	SN3	100<=S<225	1981-1990	-0.6	0.201	0.9	1.1
2-stroke	SN3	100<=S<225	1991-Stage I	-0.6	0.201	0.9	1.1
2-stroke	SN3	100<=S<225	Stage I	-0.33	0.266	1.109	5.103
2-stroke	SN3	100<=S<225	Stage II	-0.33	0	1.109	5.103
2-stroke	SN4	S>=225	<1981	-0.6	0.201	0.9	1.1
2-stroke	SN4	S>=225	1981-1990	-0.6	0.201	0.9	1.1
2-stroke	SN4	S>=225	1991-Stage I	-0.6	0.201	0.9	1.1
2-stroke	SN4	S>=225	Stage I	-0.274	0	0.887	1.935
2-stroke	SN4	S>=225	Stage II	-0.274	0	0.887	1.935

Deterioration factors for gasoline 4-stroke machinery.

Engine	Size code	Size classe	Emission Level	NO _x	VOC	CO	TSP
4-stroke	SN1	S<66	<1981	-0.6	1.1	0.9	1.1
4-stroke	SN1	S<66	1981-1990	-0.6	1.1	0.9	1.1
4-stroke	SN1	S<66	1991-Stage I	-0.6	1.1	0.9	1.1
4-stroke	SN1	S<66	Stage I	-0.3	1.753	1.051	1.753
4-stroke	SN1	S<66	Stage II	-0.3	1.753	1.051	1.753
4-stroke	SN2	66<=S<100	<1981	-0.6	1.1	0.9	1.1
4-stroke	SN2	66<=S<100	1981-1990	-0.6	1.1	0.9	1.1
4-stroke	SN2	66<=S<100	1991-Stage I	-0.6	1.1	0.9	1.1
4-stroke	SN2	66<=S<100	Stage I	-0.3	1.753	1.051	1.753
4-stroke	SN2	66<=S<100	Stage II	-0.3	1.753	1.051	1.753
4-stroke	SN3	100<=S<225	<1981	-0.6	1.1	0.9	1.1
4-stroke	SN3	100<=S<225	1981-1990	-0.6	1.1	0.9	1.1
4-stroke	SN3	100<=S<225	1991-Stage I	-0.6	1.1	0.9	1.1
4-stroke	SN3	100<=S<225	Stage I	-0.3	1.753	1.051	1.753
4-stroke	SN3	100<=S<225	Stage II	-0.3	1.753	1.051	1.753
4-stroke	SN4	S>=225	<1981	-0.6	1.1	0.9	1.1
4-stroke	SN4	S>=225	1981-1990	-0.6	1.1	0.9	1.1
4-stroke	SN4	S>=225	1991-Stage I	-0.6	1.1	0.9	1.1
4-stroke	SN4	S>=225	Stage I	-0.599	1.095	1.307	1.095
4-stroke	SN4	S>=225	Stage II	-0.599	1.095	1.307	1.095
4-stroke	SH2	20<=S<50	<1981	0	0	0	0
4-stroke	SH2	20<=S<50	1981-1990	0	0	0	0
4-stroke	SH2	20<=S<50	1991-Stage I	0	0	0	0
4-stroke	SH2	20<=S<50	Stage I	0	0	0	0
4-stroke	SH2	20<=S<50	Stage II	0	0	0	0
4-stroke	SH3	S>=50	<1981	0	0	0	0
4-stroke	SH3	S>=50	1981-1990	0	0	0	0
4-stroke	SH3	S>=50	1991-Stage I	0	0	0	0
4-stroke	SH3	S>=50	Stage I	0	0	0	0
4-stroke	SH3	S>=50	Stage II	0	0	0	0

Transient factors for diesel machinery.

Emission Level	Load	NO _x	VOC	CO	TSP	Fuel
<1981	High	0.95	1.05	1.53	1.23	1.01
1981-1990	High	0.95	1.05	1.53	1.23	1.01
1991-Stage I	High	0.95	1.05	1.53	1.23	1.01
Stage I	High	0.95	1.05	1.53	1.23	1.01
Stage II	High	0.95	1.05	1.53	1.23	1.01
Stage IIIA	High	0.95	1.05	1.53	1.23	1.01
Stage IIIB	High	1	1	1	1	1
Stage IV	High	1	1	1	1	1
<1981	Low	1.1	2.29	2.57	1.97	1.18
1981-1990	Low	1.1	2.29	2.57	1.97	1.18
1991-Stage I	Low	1.1	2.29	2.57	1.97	1.18
Stage I	Low	1.1	2.29	2.57	1.97	1.18
Stage II	Low	1.1	2.29	2.57	1.97	1.18
Stage IIIA	Low	1.1	2.29	2.57	1.97	1.18
Stage IIIB	Low	1	1	1	1	1
Stage IV	Low	1	1	1	1	1

Annual working hours, load factors and lifetimes for agricultural tractors.

Tractor type	Annual working hours	Load factor	Lifetime (yrs)
Diesel	500 (0-7 years)	0.5	30
	500-100 (7-16 years)		
	100 (>16 years)		
Gasoline (certified)	100	0.4	37
Gasoline (non certified)	50	0.4	37

Annual working hours, load factors and lifetimes for harvesters.

Annual working hours	Load factor	Lifetime (yrs)
250-100 (linear decrease 0-24 years)	0.8	25

Annual working hours, load factors and lifetime for machine pool machinery.

Tractor type	Hours pr yr	Load factor	Lifetime (yrs)
Tractors	750	0.5	7
Harvesters	100	0.8	11
Self-propelled vehicles	500	0.75	6

Operational data for other machinery types in agriculture.

Machinery type	Fuel type	Load factor	Lifetime (yrs)	Hours	Size (kW)
ATV private	Gasoline	-	6	250	-
ATV professional	Gasoline	-	8	400	-
Bedding machines	Gasoline	0.3	10	50	3
Fodder trucks	Gasoline	0.4	10	200	8
Other (gasoline)	Gasoline	0.4	10	50	5
Scrapers	Gasoline	0.3	10	50	3
Self-propelled vehicles	Diesel	0.75	15	150	60
Sweepers	Gasoline	0.3	10	50	3

Annual working hours, load factors and lifetimes for forestry machinery.

Machinery type	Hours	Load factors	Lifetime
Chippers	1200	0.5	6
Tractors (other)	100 (1990)	0.5	15
	400 (2004)		
Tractors (silvicultural)	800	0.5	6
Harvesters	1200	0.5	8
Forwarders	1200	0.5	8
Chain saws (forestry)	800	0.4	3

Annual working hours, load factors and lifetime for fork lifts.

Hours pr yr	Load factor	Lifetime (yrs)
1200 (≥ 50 kW and ≤ 10 years old)	0.27	20
650 (≥ 50 kW and > 10 years old)		
650 (< 50 kW)		

Operational data for construction machinery.

Machinery type	Load factor	Lifetime	Hours	Size
Track type dozers	0.5	10	1100	140
Track type loaders	0.5	10	1100	100 (1990) 150 (2004)
Wheel loaders (0-5 tonnes)	0.5	10	1200	20
Wheel loaders (> 5,1 tonnes)	0.5	10	1200	120
Wheel type excavators	0.6	10	1200	100
Track type excavators (0-5 tonnes)	0.6	10	1100	20
Track type excavators (>5,1 tonnes)	0.6	10	1100	120
Excavators/Loaders	0.45	10	700	50
Dump trucks	0.4	10	900 (1990) 1200 (2004)	60 (1990) 180 (2004)
Mini loaders	0.5	14	700	30
Telescopic loaders	0.5	14	1000	35

Stock and operational data for other machinery types in industry.

Sector	Fuel type	Machinery type	Size (kW)	No Load Factor	Hours
Construction machinery	Diesel	Tampers/Land rollers	30	2800	0.45 600
Construction machinery	Diesel	Generators (diesel)	45	5000	0.5 200
Construction machinery	Diesel	Kompressors (diesel)	45	5000	0.5 500
Construction machinery	Diesel	Pumps (diesel)	75	1000	0.5 5
Construction machinery	Diesel	Asphalt pavers	80	300	0.35 700
Construction machinery	Diesel	Motor graders	100	100	0.4 700
Construction machinery	Diesel	Refuse compressors	160	100	0.25 1300
Construction machinery	Gasoline	Generators (gasoline)	2.5	11000	0.4 80
Construction machinery	Gasoline	Pumps (gasoline)	4	10000	0.4 300
Construction machinery	Gasoline	Kompressors (gasoline)	4	500	0.35 15
Industry	Diesel	Refrigerating units (distribution)	8	3000	0.5 1250
Industry	Diesel	Refrigerating units (long distance)	15	3500	0.5 200
Industry	Diesel	Tractors (transport, industry)	50	3000	0.4 500
Airport GSE and other	Diesel	Airport GSE and other (light duty)	100	500	0.5 400
Airport GSE and other	Diesel	Airport GSE and other (medium duty)	125	350	0.5 300
Airport GSE and other	Diesel	Airport GSE and other (Heavy duty)	175	650	0.5 200
Building and construction	Diesel	Vibratory plates	6	3500	0.6 300
Building and construction	Diesel	Aereal lifts (diesel)	30	150	0.4 400
Building and construction	Diesel	Sweepers (diesel)	30	200	0.4 300
Building and construction	Diesel	High pressure cleaners (diesel)	30	50	0.8 500
Building and construction	Gasoline	Rammers	2.5	3000	0.4 80
Building and construction	Gasoline	Drills	3	100	0.4 10
Building and construction	Gasoline	Vibratory plates (gasoline)	4	2500	0.5 200
Building and construction	Gasoline	Cutters	4	800	0.5 50
Building and construction	Gasoline	Other (gasoline)	5	1000	0.5 40
Building and construction	Gasoline	High pressure cleaners (gasoline)	5	500	0.6 200
Building and construction	Gasoline	Sweepers (gasoline)	10	500	0.4 150
Building and construction	Gasoline	Slicers	10	100	0.7 150
Building and construction	Gasoline	Aereal lifts (gasoline)	20	50	0.4 400

Operational data for the most important types of household and gardening machinery.

Machinery type	Engine	Size (kW)	Hours	Load factor	Lifetime (yrs)
Chain saws (private)	2-stroke	2	5	0.3	10
Chain saws (professional)	2-stroke	3	270	0.4	3
Cultivators (private-large)	4-stroke	3.7	5	0.6	5
Cultivators (private-small)	4-stroke	1	5	0.6	15
Cultivators (professional)	4-stroke	7	360	0.6	8
Hedge cutters (private)	2-stroke	0.9	10	0.5	10
Hedge cutters (professional)	2-stroke	2	300	0.5	4
		2.5	25		
		(2000)			
		3.5			
Lawn movers (private)	4-stroke	(2004)		0.4	8
		2.5	250		
		(2000)			
		3.5			
Lawn movers (professional)	4-stroke	(2004)		0.4	4
Riders (private)	4-stroke	11	50	0.5	12
Riders (professional)	4-stroke	13	330	0.5	5
Shrub clearers (private)	2-stroke	1	15	0.6	10
Shrub clearers (professional)	2-stroke	2	300	0.6	4
Trimmers (private)	2-stroke	0.9	20	0.5	10
Trimmers (professional)	2-stroke	0.9	200	0.5	4

Stock and operational data for other machines in household and gardening.

Machinery type	Engine	No.	Size (kW)	Hours	Load factor	Lifetime (yrs)
Chippers	2-stroke	200	10	100	0.7	10
Garden shredders	2-stroke	500	3	20	0.7	10
Other (gasoline)	2-stroke	200	2	20	0.5	10
Suction machines	2-stroke	300	4	80	0.5	10
Wood cutters	4-stroke	100	4	15	0.5	10

Operational data for recreational craft.

Fuel type	Vessel type	Engine type	Stroke	Hours	Lifetime	Load factor
Gasoline	Other boats (<20 ft)	Out board engine	2-stroke	30	10	0.5
Gasoline	Other boats (<20 ft)	Out board engine	4-stroke	30	10	0.5
Gasoline	Yawls and cabin boats	Out board engine	2-stroke	50	10	0.5
Gasoline	Yawls and cabin boats	Out board engine	4-stroke	50	10	0.5
Gasoline	Sailing boats (<26ft)	Out board engine	2-stroke	5	10	0.5
Gasoline	Sailing boats (<26ft)	Out board engine	4-stroke	5	10	0.5
Gasoline	Speed boats	In board engine	4-stroke	75	10	0.5
Gasoline	Speed boats	Out board engine	2-stroke	50	10	0.5
Gasoline	Speed boats	Out board engine	4-stroke	50	10	0.5
Gasoline	Water scooters	Built in	2-stroke	10	10	0.5
Gasoline	Water scooters	Built in	4-stroke	10	10	0.5
Diesel	Motor boats (27-34 ft)	In board engine		150	15	0.5
Diesel	Motor boats (>34 ft)	In board engine		100	15	0.5
Diesel	Motor boats (<27 ft)	In board engine		75	15	0.5
Diesel	Motor sailers	In board engine		75	15	0.5
Diesel	Sailing boats (<26ft)	In board engine		25	15	0.5

Stock data for diesel tractors 1985-2012.

Size (kW)	Emission Level	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
37	<1981	3882	3792	3542	3543	3403	3234	3106	2922	2861	2610	2605	2273	2193	1918	1796
37	1981-1990	635	731	760	835	855	879	889	883	915	887	945	883	918	869	888
37	1991-Stage I							25	107	153	201	278	354	445	496	554
37	Stage I															
37	Stage II															
37	Stage IIIA															
45	<1981	25988	25387	23709	23718	22781	21650	20796	19563	19154	17475	17441	15219	14684	12840	12025
45	1981-1990	5740	6808	7263	8075	8476	8770	8867	8805	9128	8848	9419	8807	9151	8668	8856
45	1991-Stage I							203	202	209	203	216	202	210	199	203
49	1991-Stage I								154	281	485	602	618	702	749	765
52	1991-Stage I															247
52	Stage I															
52	Stage II															
52	Stage IIIA															
56	1991-Stage I								201	338	428	747	943	1181	1280	1307
60	<1981	54651	53387	49857	49877	47907	45529	43732	41140	40278	36747	36676	32004	30879	27001	25287
60	1981-1990	11751	14613	15795	17797	19395	20542	20770	20624	21380	20725	22063	20628	21434	20304	20744
60	1991-Stage I							863	857	888	861	917	857	891	844	862
63	1991-Stage I								468	855	1325	2014	2384	2837	3011	3076
67	1991-Stage I															671
67	Stage I															
67	Stage II															
67	Stage IIIA															
67	Stage IIIB															
71	1991-Stage I								411	715	1179	1949	2507	3344	3594	3672
78	<1981	14558	14221	13281	13286	12761	12128	11649	10959	10729	9789	9770	8525	8226	7192	6736
78	1981-1990	4592	6152	7196	8559	10026	11323	11448	11368	11785	11424	12162	11371	11815	11192	11434
78	1991-Stage I							1233	1503	1713	1945	2429	2561	2946	2994	3287
78	Stage I															
78	Stage II															
78	Stage IIIA															
78	Stage IIIB															
86	1991-Stage I								108	193	333	589	880	1364	1532	1718
86	Stage I															
86	Stage II															

86	Stage IIIA															
86	Stage IIIB															
93	1991-Stage I															149
93	Stage I															
93	Stage II															
93	Stage IIIA															
93	Stage IIIB															
97	1991-Stage I								71	175	443	962	1556	2327	2638	2695
101	<1981	4659	4551	4250	4252	4084	3881	3728	3507	3433	3132	3126	2728	2632	2302	2156
101	1981-1990	1158	1434	1618	1921	2156	2377	2403	2387	2474	2398	2553	2387	2480	2350	2400
101	1991-Stage I							266	264	274	266	283	264	275	260	696
101	Stage I															
101	Stage II															
101	Stage IIIA															
101	Stage IIIB															
112	1991-Stage I								63	114	166	252	422	690	790	978
112	Stage I															
112	Stage II															
112	Stage IIIA															
112	Stage IIIB															
127	1991-Stage I								12	36	81	193	279	408	457	590
127	Stage I															
127	Stage II															
127	Stage IIIA															
127	Stage IIIB															
131	<1981	798	780	728	728	700	665	639	601	588	537	536	467	451	394	369
131	1981-1990	288	421	500	651	753	887	897	890	923	895	952	890	925	876	895
131	1991-Stage I							97	97	100	97	103	97	100	95	97
157	1981-1990		2	3	6	11	15	15	15	16	15	16	15	16	15	15
157	1991-Stage I							9	23	39	102	232	357	545	648	784
157	Stage I															
157	Stage II															
157	Stage IIIA															
157	Stage IIIB															
186	1991-Stage I															23
186	Stage I															
186	Stage II															

186	Stage IIIA
186	Stage IIIB

Size (kW)	Emission Level	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
37	<1981	1601	1449	1298	1148	993	833	664	504	342	176			
37	1981-1990	871	876	882	892	900	906	903	914	930	959	991	834	667
37	1991-Stage I	568	572	576	582	587	592	590	597	607	626	647	667	688
37	Stage I		33	56	83	84	84	84	85	86	89	92	95	98
37	Stage II					23	53	162	324	330	340	351	362	374
37	Stage IIIA									109	205	333	491	626
45	<1981	10715	9700	8690	7685	6646	5577	4447	3376	2290	1180			
45	1981-1990	8681	8731	8800	8894	8974	9037	9006	9116	9274	9563	9883	8931	7919
45	1991-Stage I	199	200	202	204	206	207	207	209	213	219	227	234	241
49	1991-Stage I	750	754	760	768	775	780	778	787	801	826	853	880	908
52	1991-Stage I	358	360	363	367	370	373	372	376	383	395	408	421	434
52	Stage I		132	242	377	381	383	382	387	393	406	419	432	446
52	Stage II					68	147	241	347	353	364	377	388	401
52	Stage IIIA									86	133	202	290	345
56	1991-Stage I	1281	1289	1299	1313	1325	1334	1329	1346	1369	1412	1459	1504	1552
60	<1981	22533	20397	18273	16162	13976	11729	9351	7099	4815	2482			
60	1981-1990	20333	20451	20612	20834	21019	21167	21096	21353	21723	22401	23150	21220	19172
60	1991-Stage I	845	850	856	866	873	879	876	887	903	931	962	991	1023
63	1991-Stage I	3015	3033	3057	3090	3117	3139	3128	3167	3221	3322	3433	3539	3653
67	1991-Stage I	1343	1351	1361	1376	1388	1398	1393	1410	1435	1479	1529	1576	1627
67	Stage I		533	835	1113	1123	1131	1127	1141	1161	1197	1237	1275	1316
67	Stage II					375	729	1144	1524	1550	1599	1652	1703	1758
67	Stage IIIA									303	472	658	890	919
67	Stage IIIB													162
71	1991-Stage I	3600	3620	3649	3688	3721	3747	3735	3780	3846	3966	4098	4225	4360
78	<1981	6002	5433	4868	4305	3723	3124	2491	1891	1283	661			
78	1981-1990	11208	11273	11361	11484	11586	11668	11628	11770	11974	12348	12761	12450	12123
78	1991-Stage I	3436	3727	3756	3797	3830	3857	3844	3891	3959	4082	4219	4349	4489
78	Stage I			325	329	332	334	333	337	343	354	365	377	389
78	Stage II				227	310	400	463	469	477	492	508	524	541
78	Stage IIIA								63	121	147	183	226	233
78	Stage IIIB													41

86	1991-Stage I	1876	2023	2039	2061	2079	2094	2087	2112	2149	2216	2290	2361	2437
86	Stage I			134	136	137	138	137	139	142	146	151	156	161
86	Stage II				91	343	530	760	769	783	807	834	860	888
86	Stage IIIA								226	434	529	657	811	837
86	Stage IIIB													146
93	1991-Stage I	245	325	327	331	334	336	335	339	345	356	368	379	391
93	Stage I			114	115	116	117	116	118	120	123	128	132	136
93	Stage II				107	186	313	512	518	527	544	562	579	598
93	Stage IIIA								264	470	574	682	836	863
93	Stage IIIB													143
97	1991-Stage I	2642	2657	2678	2707	2731	2750	2741	2774	2822	2911	3008	3101	3200
101	<1981	1921	1739	1558	1378	1191	1000	797	605	410	212			
101	1981-1990	2353	2367	2385	2411	2432	2449	2441	2471	2514	2592	2679	2536	2385
101	1991-Stage I	1116	1567	1579	1596	1611	1622	1616	1636	1664	1716	1774	1828	1887
101	Stage I			232	234	236	238	237	240	244	252	260	268	277
101	Stage II				136	357	635	776	785	799	824	851	878	906
101	Stage IIIA								188	336	410	487	597	617
101	Stage IIIB													102
112	1991-Stage I	1265	1626	1639	1656	1671	1683	1677	1698	1727	1781	1841	1897	1958
112	Stage I			465	470	474	478	476	482	490	505	522	539	556
112	Stage II				337	732	1170	1763	1785	1815	1872	1935	1994	2059
112	Stage IIIA								378	663	823	971	1264	1304
112	Stage IIIB													248
127	1991-Stage I	707	847	854	863	871	877	874	884	900	928	959	988	1020
127	Stage I			152	154	155	156	156	158	161	166	171	176	182
127	Stage II				78	268	453	591	599	609	628	649	669	690
127	Stage IIIA								292	675	880	1048	1254	1295
127	Stage IIIB													166
131	<1981	329	298	267	236	204	171	137	104	70	36			
131	1981-1990	878	883	890	899	907	914	911	922	938	967	999	991	983
131	1991-Stage I	95	96	96	97	98	99	99	100	102	105	108	112	115
157	1981-1990	15	15	15	15	16	16	16	16	16	17	17	18	18
157	1991-Stage I	900	905	912	922	930	937	934	945	961	991	1025	1056	1090
157	Stage I		89	89	90	91	92	91	92	94	97	100	103	107
157	Stage II			149	415	695	1089	1085	1098	1117	1152	1191	1227	1267
157	Stage IIIA							623	1453	2140	2586	3047	3141	3242
157	Stage IIIB												388	826

186	1991-Stage I	53	54	54	55	55	56	55	56	57	59	61	63	65
186	Stage I		47	48	48	49	49	49	49	50	52	54	55	57
186	Stage II			68	207	320	481	480	486	494	509	526	543	560
186	Stage IIIA							272	685	1103	1427	1665	1717	1772
186	Stage IIIB												228	561

Stock data for gasoline tractors 1985-2005.

Size (kW)	Emission Level	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
Certified	<1981	13176	12541	11906	11270	10635	10000	9053	8148	7285	6465	5687	4951	4258	3607	2998
Non certified	<1981	26352	25082	23811	22541	21270	20000	19042	18041	16998	15913	14785	13616	12403	11149	9852
<i>Continued</i>		2000	2001	2002	2003	2004	2005									
Certified	<1981	2432	1908	1427	987	591	236									
Non certified	<1981	8512	7131	5707	4240	2732	1180									

Stock data for harvesters 1985-2012.

FSize	Size Group	Emission Level	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
0	0<S<=50	<1981	26601	24394	22599	22144	19842	18915	17241	15607	14575	12673	10700	9491	6966	5446	3589
0	0<S<=50	1981-1990	519	534	550	582	566	591	594	601	635	636	633	683	641	686	672
50	50<S<=60	<1981	2703	2648	2634	2785	2711	2828	2847	2876	3040	3044	3029	3271	3068	2930	2235
50	50<S<=60	1981-1990	853	1102	1164	1275	1258	1333	1341	1355	1432	1434	1427	1541	1446	1548	1516
50	50<S<=60	1991-Stage I							8	8	8	8	8	9	9	9	9
60	60<S<=70	<1981	1786	1750	1741	1841	1792	1869	1881	1901	2009	2012	2002	2162	2028	2171	2127
60	60<S<=70	1981-1990	1138	1679	1943	2237	2213	2348	2363	2388	2524	2527	2515	2716	2547	2727	2671
60	60<S<=70	1991-Stage I							8	16	18	21	22	24	23	24	24
70	70<S<=80	<1981	929	910	905	958	932	972	979	989	1045	1046	1041	1125	1055	1129	1106
70	70<S<=80	1981-1990	383	699	1026	1165	1318	1493	1502	1518	1604	1606	1598	1726	1619	1733	1698
70	70<S<=80	1991-Stage I							72	77	83	86	87	96	91	98	96
70	70<S<=80	Stage I															1
80	80<S<=90	<1981	323	317	315	333	324	338	340	344	363	364	362	391	367	393	385
80	80<S<=90	1981-1990	383	562	645	967	1107	1466	1475	1491	1575	1577	1570	1695	1590	1702	1667
80	80<S<=90	1991-Stage I							61	158	181	200	200	217	207	222	217
80	80<S<=90	Stage I															1
90	90<S<=100	1981-1990	89	175	235	387	515	670	674	681	720	721	717	775	726	778	762
90	90<S<=100	1991-Stage I							180	257	320	329	351	382	367	393	385
90	90<S<=100	Stage I															1
100	100<S<=120	1981-1990		54	106	219	334	589	592	599	633	634	630	681	639	684	670
100	100<S<=120	1991-Stage I							129	253	316	375	440	567	586	673	660

100	100<S<=120	Stage I													2
120	120<S<=140	1981-1990	4	69	183	184	186	197	197	196	212	199	213	208	
120	120<S<=140	1991-Stage I				70	148	189	215	319	484	626	804	860	
120	120<S<=140	Stage I													21
120	120<S<=140	Stage II													
120	120<S<=140	Stage IIIA													
120	120<S<=140	Stage IIIB													
140	140<S<=160	1991-Stage I					8	36	69	112	271	354	554	632	
140	140<S<=160	Stage II													
140	140<S<=160	Stage IIIA													
140	140<S<=160	Stage IIIB													
160	160<S<=180	1991-Stage I								26	69	200	374	440	
160	160<S<=180	Stage II													
160	160<S<=180	Stage IIIA													
160	160<S<=180	Stage IIIB													
180	180<S<=200	1991-Stage I									20	67	117	193	
180	180<S<=200	Stage II													
180	180<S<=200	Stage IIIA													
180	180<S<=200	Stage IIIB													
200	200<S<=220	1991-Stage I											45	92	
200	200<S<=220	Stage II													
200	200<S<=220	Stage IIIA													
200	200<S<=220	Stage IIIB													
220	220<S<=240	1991-Stage I													3
220	220<S<=240	Stage II													
220	220<S<=240	Stage IIIA													
220	220<S<=240	Stage IIIB													
240	240<S<=260	1991-Stage I													3
240	240<S<=260	Stage II													
240	240<S<=260	Stage IIIA													
240	240<S<=260	Stage IIIB													
260	260<S<=280	1991-Stage I													14
260	260<S<=280	Stage II													
260	260<S<=280	Stage IIIA													
260	260<S<=280	Stage IIIB													
280	280<S<=300	1991-Stage I													
280	280<S<=300	Stage II													

280	280<S<=300	Stage IIIA
280	280<S<=300	Stage IIIB
300	300<S<=320	Stage II
300	300<S<=320	Stage IIIA
300	300<S<=320	Stage IIIB

FSize	Size Group	Emission Level	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
0	0<S<=50	<1981	2873	1854	1275	754	269								
0	0<S<=50	1981-1990	715	758	778	816	882	913	779	628	448	268	78	38	
50	50<S<=60	<1981	1999	1570	1260	897	391								
50	50<S<=60	1981-1990	1612	1711	1755	1841	1990	2060	1856	1645	1335	1034	730	296	170
50	50<S<=60	1991-Stage I	10	10	10	11	12	12	12	12	12	12	13	15	16
60	60<S<=70	<1981	2073	1648	1340	981	482								
60	60<S<=70	1981-1990	2841	3014	3093	3243	3506	3630	3344	3062	2659	2284	1922	1053	539
60	60<S<=70	1991-Stage I	25	27	27	29	31	32	32	32	32	33	35	39	43
70	70<S<=80	<1981	1176	1248	1105	735	216								
70	70<S<=80	1981-1990	1806	1916	1966	2061	2229	2307	2164	2043	1939	1862	1813	1415	806
70	70<S<=80	1991-Stage I	102	109	112	117	126	131	130	129	131	134	141	161	175
70	70<S<=80	Stage I	1	1	1	1	1	1	1	1	1	1	2	2	2
80	80<S<=90	<1981	409	434	445	467	216								
80	80<S<=90	1981-1990	1773	1881	1931	2024	2189	2266	2123	2002	1897	1819	1768	1642	1592
80	80<S<=90	1991-Stage I	231	245	252	264	285	295	294	292	295	303	317	363	394
80	80<S<=90	Stage I	1	1	1	1	1	1	1	1	1	1	2	2	2
90	90<S<=100	1981-1990	810	860	882	925	1000	1035	1031	1023	986	964	957	915	860
90	90<S<=100	1991-Stage I	410	435	446	468	506	524	521	518	523	538	563	643	698
90	90<S<=100	Stage I	1	1	1	1	1	1	1	1	1	1	2	2	2
100	100<S<=120	1981-1990	712	756	775	813	879	910	906	900	909	934	978	1008	979
100	100<S<=120	1991-Stage I	702	744	764	801	866	896	892	886	896	920	963	1100	1195
100	100<S<=120	Stage I	2	2	2	3	3	3	3	3	3	3	3	4	4
120	120<S<=140	1981-1990	222	235	241	253	274	283	282	280	283	291	304	348	378
120	120<S<=140	1991-Stage I	918	977	1003	1051	1137	1177	1172	1163	1176	1208	1264	1444	1569
120	120<S<=140	Stage I	26	31	32	33	36	37	37	37	37	38	40	46	50
120	120<S<=140	Stage II					3	4	4	4	4	4	4	4	5
120	120<S<=140	Stage IIIA							1	1	1	4	5	5	6
120	120<S<=140	Stage IIIB												3	3
140	140<S<=160	1991-Stage I	715	795	816	855	925	957	953	946	957	983	1028	1175	1277
140	140<S<=160	Stage II			20	35	48	56	56	56	56	58	60	69	75

140	140<S<=160	Stage IIIA							5	8	12	16	18	21	23
140	140<S<=160	Stage IIIB												5	8
160	160<S<=180	1991-Stage I	533	602	618	648	700	725	722	716	724	744	779	890	967
160	160<S<=180	Stage II			40	70	91	105	105	104	105	108	113	129	140
160	160<S<=180	Stage IIIA							9	14	20	24	27	31	34
160	160<S<=180	Stage IIIB												4	9
180	180<S<=200	1991-Stage I	249	300	308	323	349	362	360	357	361	371	389	444	482
180	180<S<=200	Stage II			61	91	114	129	128	127	129	132	138	158	172
180	180<S<=200	Stage IIIA							9	14	20	24	27	31	34
180	180<S<=200	Stage IIIB												4	9
200	200<S<=220	1991-Stage I	142	187	192	201	218	225	224	223	225	231	242	277	301
200	200<S<=220	Stage II			40	70	91	105	105	104	105	108	113	129	140
200	200<S<=220	Stage IIIA							9	14	20	24	27	31	34
200	200<S<=220	Stage IIIB												4	9
220	220<S<=240	1991-Stage I	48	151	155	162	175	181	181	179	181	186	195	223	242
220	220<S<=240	Stage II			72	114	164	221	220	219	221	227	238	271	295
220	220<S<=240	Stage IIIA							61	123	196	237	276	315	342
220	220<S<=240	Stage IIIB												45	91
240	240<S<=260	1991-Stage I	71	142	145	152	165	170	170	169	170	175	183	209	227
240	240<S<=260	Stage II			72	125	201	301	299	297	301	309	323	369	401
240	240<S<=260	Stage IIIA							113	232	371	450	525	599	651
240	240<S<=260	Stage IIIB												85	172
260	260<S<=280	1991-Stage I	61	131	134	140	152	157	157	155	157	161	169	193	210
260	260<S<=280	Stage II			72	125	201	301	299	297	301	309	323	369	401
260	260<S<=280	Stage IIIA							113	232	371	450	525	599	651
260	260<S<=280	Stage IIIB												85	172
280	280<S<=300	1991-Stage I		33	34	36	39	40	40	40	40	41	43	49	53
280	280<S<=300	Stage II			72	125	201	301	299	297	301	309	323	369	401
280	280<S<=300	Stage IIIA							113	232	371	450	525	599	651
280	280<S<=300	Stage IIIB												85	172
300	300<S<=320	Stage II				25	60	108	108	107	108	111	116	133	145
300	300<S<=320	Stage IIIA							57	116	185	225	262	300	326
300	300<S<=320	Stage IIIB												43	86

Stock data for fork lifts 1985-2012.

FuelCode	Size (kW)	Emission Level	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
205B	35	<1981	387	361	336	311	285	260	234	209	183	158	133	107	84	58	30
205B	35	1981-1990	120	162	202	239	270	297	297	297	297	297	297	297	297	297	297
205B	35	1991-Stage I							26	49	65	93	131	168	218	247	275
205B	35	Stage II															
205B	35	Stage IIIA															
205B	45	<1981	1612	1506	1400	1294	1188	1082	976	870	764	658	552	446	349	243	126
205B	45	1981-1990	499	674	839	994	1122	1233	1233	1233	1233	1233	1233	1233	1233	1233	1233
205B	45	1991-Stage I							108	203	270	386	544	699	905	1063	1063
205B	45	Stage I															151
205B	45	Stage II															
205B	45	Stage IIIA															
205B	50	<1981	2173	2031	1888	1745	1602	1459	1316	1174	1031	888	745	602	471	328	170
205B	50	1981-1990	673	909	1131	1340	1512	1662	1662	1662	1662	1662	1662	1662	1662	1662	1662
205B	50	1991-Stage I							145	273	363	519	732	940	1217	1469	1469
205B	50	Stage I															240
205B	50	Stage II															
205B	50	Stage IIIA															
205B	75	<1981	497	465	432	399	367	334	301	269	236	203	170	138	108	75	39
205B	75	1981-1990	154	208	259	307	347	382	382	382	382	382	382	382	382	382	382
205B	75	1991-Stage I							33	63	84	120	169	217	281	354	354
205B	75	Stage I															70
205B	75	Stage II															
205B	75	Stage IIIA															
205B	75	Stage IIIB															
205B	120	<1981	111	103	96	89	81	74	67	60	52	45	38	31	24	17	9
205B	120	1981-1990	34	46	57	68	77	85	85	85	85	85	85	85	85	85	85
205B	120	1991-Stage I							7	14	19	27	38	49	63	97	97
205B	120	Stage I															32
205B	120	Stage II															
205B	120	Stage IIIA															
205B	120	Stage IIIB															
3030	33		5420	5427	5390	5323	5265	5215	5156	5068	4947	4863	4835	4792	4732	4765	4712
3030	40		4917	4923	4889	4828	4775	4730	4676	4596	4486	4410	4384	4344	4289	4295	4223
3030	50		2149	2151	2137	2110	2087	2067	2044	2008	1960	1926	1915	1897	1874	1926	1941
3030	78		97	97	96	95	94	93	92	91	89	88	88	87	86	90	92

3030	120	1	2
------	-----	---	---

FuelCode	Size (kW)	Emission Level	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
205B	35	<1981													
205B	35	1981-1990	297	277	249	232	198	177	135	95	58	27			
205B	35	1991-Stage I	304	304	304	304	304	304	304	304	304	304	304	278	255
205B	35	Stage II		23	53	75	89	117	152	152	152	152	152	152	152
205B	35	Stage IIIA								41	76	92	99	126	153
205B	45	<1981													
205B	45	1981-1990	1233	1151	1036	964	820	734	559	394	239	111			
205B	45	1991-Stage I	1063	1063	1063	1063	1063	1063	1063	1063	1063	1063	1063	955	860
205B	45	Stage I	303	422	524	664	664	664	664	664	664	664	664	664	664
205B	45	Stage II					104	232	452	612	612	612	612	612	612
205B	45	Stage IIIA									126	181	225	346	467
205B	50	<1981													
205B	50	1981-1990	1662	1551	1396	1299	1105	989	753	531	322	150			
205B	50	1991-Stage I	1469	1469	1469	1469	1469	1469	1469	1469	1469	1469	1469	1324	1196
205B	50	Stage I	461	682	897	1135	1135	1135	1135	1135	1135	1135	1135	1135	1135
205B	50	Stage II					187	447	818	1134	1134	1134	1134	1134	1134
205B	50	Stage IIIA									181	275	354	562	770
205B	75	<1981													
205B	75	1981-1990	382	357	321	299	255	228	174	123	75	35			
205B	75	1991-Stage I	354	354	354	354	354	354	354	354	354	354	354	321	291
205B	75	Stage I	162	234	311	311	311	311	311	311	311	311	311	311	311
205B	75	Stage II				58	129	208	326	326	326	326	326	326	326
205B	75	Stage IIIA								142	213	252	294	376	376
205B	75	Stage IIIB													82
205B	120	<1981													
205B	120	1981-1990	85	80	72	67	57	51	39	28	17	8			
205B	120	1991-Stage I	97	97	97	97	97	97	97	97	97	97	97	90	83
205B	120	Stage I	71	89	118	118	118	118	118	118	118	118	118	118	118
205B	120	Stage II				16	38	58	112	112	112	112	112	112	112
205B	120	Stage IIIA								58	70	76	140	179	179
205B	120	Stage IIIB													39
3030	33		4718	4677	4655	4595	4494	4345	4220	4154	4043	3941	3746	3644	3572
3030	40		4218	4214	4244	4224	4166	4116	4048	4005	3951	3878	3723	3660	3624

3030	50	1897	1938	2003	2020	2018	2029	2061	2136	2198	2192	2142	2172	2214
3030	78	88	95	98	99	104	104	114	123	147	149	151	161	172
3030	120	2	2	3	3	3	3	3	3	3	3	7	8	9

Stock data for construction machinery 1985-2012.

EquipmentName (Eng)	Emission Level	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
Track type dozers	<1981	125	100	75	50	25										
Track type dozers	1981-1990	125	150	175	200	225	250	221	193	166	139	114	89	66	43	21
Track type dozers	1991-Stage I							25	48	71	93	114	134	153	172	189
Track type dozers	Stage II															
Track type dozers	Stage IIIA															
Track type dozers	Stage IIIB															
Track type loaders	<1981	50	40	30	20	10										
Track type loaders	1981-1990	50	60	70	80	90	100	89	79	68	58	48	38	28	19	9
Track type loaders	1991-Stage I							10	20	29	39	48	57	66	75	83
Track type loaders	Stage II															
Track type loaders	Stage IIIA															
Track type loaders	Stage IIIB															
Wheel loaders (0-5 tons)	1981-1990							186	331	434	496	517	496	434	331	186
Wheel loaders (0-5 tons)	1991-Stage I							21	83	186	331	517	744	1013	1323	1674
Wheel loaders (0-5 tons)	Stage II															
Wheel loaders (0-5 tons)	Stage IIIA															
Wheel loaders (> 5,1 tons)	<1981	1250	1000	750	500	250										
Wheel loaders (> 5,1 tons)	1981-1990	1250	1500	1750	2000	2250	2500	2228	1960	1698	1441	1188	941	698	460	228
Wheel loaders (> 5,1 tons)	1991-Stage I							248	490	728	960	1188	1411	1629	1841	1822
Wheel loaders (> 5,1 tons)	Stage I															228
Wheel loaders (> 5,1 tons)	Stage II															
Wheel loaders (> 5,1 tons)	Stage IIIA															
Wheel loaders (> 5,1 tons)	Stage IIIB															
Wheel type excavators	<1981	500	400	300	200	100										
Wheel type excavators	1981-1990	500	600	700	800	900	1000	862	732	611	498	394	298	211	132	62
Wheel type excavators	1991-Stage I							96	183	262	332	394	447	491	528	493
Wheel type excavators	Stage I															62
Wheel type excavators	Stage II															
Wheel type excavators	Stage IIIA															
Wheel type excavators	Stage IIIB															

Track type excavators (0-5 tons)	1981-1990							459	816	1071	1224	1275	1224	1071	816	459
Track type excavators (0-5 tons)	1991-Stage I							51	204	459	816	1275	1837	2500	3265	4132
Track type excavators (0-5 tons)	Stage II															
Track type excavators (0-5 tons)	Stage IIIA															
Track type excavators (>5,1 tons)	<1981	1000	800	600	400	200										
Track type excavators (>5,1 tons)	1981-1990	1000	1200	1400	1600	1800	2000	1798	1596	1394	1194	993	794	594	396	198
Track type excavators (>5,1 tons)	1991-Stage I							200	399	598	796	993	1190	1387	1583	1581
Track type excavators (>5,1 tons)	Stage I															198
Track type excavators (>5,1 tons)	Stage II															
Track type excavators (>5,1 tons)	Stage IIIA															
Track type excavators (>5,1 tons)	Stage IIIB															
Excavators/Loaders	<1981	2100	1680	1260	840	420										
Excavators/Loaders	1981-1990	2100	2520	2940	3360	3780	4200	3807	3408	3003	2592	2175	1752	1323	888	447
Excavators/Loaders	1991-Stage I							423	852	1287	1728	2175	2628	3087	3552	3575
Excavators/Loaders	Stage I															447
Excavators/Loaders	Stage II															
Excavators/Loaders	Stage IIIA															
Dump trucks	<1981	250	200	150	100	50										
Dump trucks	1981-1990	250	300	350	400	450	500	489	469	441	404	358	304	241	169	89
Dump trucks	1991-Stage I							54	117	189	269	358	455	561	676	711
Dump trucks	Stage I															89
Dump trucks	Stage II															
Dump trucks	Stage IIIA															
Dump trucks	Stage IIIB															
Mini loaders	<1981	1800	1600	1400	1200	1000	800	635	447	235						
Mini loaders	1981-1990	1000	1200	1400	1600	1800	2000	2118	2237	2355	2473	2332	2168	1980	1768	1532
Mini loaders	1991-Stage I							212	447	706	989	1296	1626	1980	2357	2758
Mini loaders	Stage II															
Mini loaders	Stage IIIA															
Telescopic loaders	1981-1990											149	265	348	398	414
Telescopic loaders	1991-Stage I											83	199	348	530	746
Telescopic loaders	Stage II															
Telescopic loaders	Stage IIIA															

EquipmentName (Eng)	Emission Level	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
Track type dozers	<1981													
Track type dozers	1981-1990													
Track type dozers	1991-Stage I	206	201	177	154	132	128	125	116	95	59	27		
Track type dozers	Stage II			20	38	56	86	100	116	126	119	109	105	75
Track type dozers	Stage IIIA							25	58	95	119	137	132	126
Track type dozers	Stage IIIB												26	50
Track type loaders	<1981													
Track type loaders	1981-1990													
Track type loaders	1991-Stage I	91	91	81	71	62	61	71	68	55	38	19		
Track type loaders	Stage II			9	18	26	40	56	68	73	76	75	72	51
Track type loaders	Stage IIIA							14	34	55	76	94	90	85
Track type loaders	Stage IIIB												18	34
Wheel loaders (0-5 tons)	1981-1990													
Wheel loaders (0-5 tons)	1991-Stage I	2067	2046	1984	1881	1736	1444	1269	1045	726	353			
Wheel loaders (0-5 tons)	Stage II		227	496	806	1158	1444	1903	2090	2177	2117	2024	1644	1265
Wheel loaders (0-5 tons)	Stage IIIA								348	726	1058	1349	1644	1897
Wheel loaders (> 5,1 tons)	<1981													
Wheel loaders (> 5,1 tons)	1981-1990													
Wheel loaders (> 5,1 tons)	1991-Stage I	1802	1559	1322	1089	861	677	485	273					
Wheel loaders (> 5,1 tons)	Stage I	450	668	881	871	861	902	969	1092	1174	854	547	266	
Wheel loaders (> 5,1 tons)	Stage II				218	431	677	969	1092	1174	1138	1094	1062	1098
Wheel loaders (> 5,1 tons)	Stage IIIA								273	587	854	1094	1328	1372
Wheel loaders (> 5,1 tons)	Stage IIIB													274
Wheel type excavators	<1981													
Wheel type excavators	1981-1990													
Wheel type excavators	1991-Stage I	459	372	293	223	162	118	74	38					
Wheel type excavators	Stage I	115	160	196	179	162	157	148	152	146	103	62	31	
Wheel type excavators	Stage II				45	81	118	148	152	146	138	124	122	127
Wheel type excavators	Stage IIIA								38	73	103	124	153	159
Wheel type excavators	Stage IIIB													32
Track type excavators (0-5 tons)	1981-1990													
Track type excavators (0-5 tons)	1991-Stage I	5101	5050	4897	4642	4285	3889	3599	3027	2073	995			
Track type excavators (0-5 tons)	Stage II		561	1224	1990	2857	3889	5399	6054	6220	5968	5554	4398	3502
Track type excavators (0-5 tons)	Stage IIIA								1009	2073	2984	3702	4398	5252
Track type excavators (>5,1 tons)	<1981													
Track type excavators (>5,1 tons)	1981-1990													

Track type excavators (>5,1 tons)	1991-Stage I	1579	1380	1181	983	785	683	536	313							
Track type excavators (>5,1 tons)	Stage I	395	591	787	786	785	910	1073	1251	1338	980	623	303			
Track type excavators (>5,1 tons)	Stage II				197	393	683	1073	1251	1338	1307	1245	1213	1252		
Track type excavators (>5,1 tons)	Stage IIIA								313	669	980	1245	1516	1565		
Track type excavators (>5,1 tons)	Stage IIIB														313	
Excavators/Loaders	<1981															
Excavators/Loaders	1981-1990															
Excavators/Loaders	1991-Stage I	3599	3170	2735	2295	1848	1370	938	481							
Excavators/Loaders	Stage I	900	1359	1824	2295	2310	2283	2344	2403	2314	1688	1137	691	319		
Excavators/Loaders	Stage II					462	913	1406	1922	1851	1688	1516	1382	1278		
Excavators/Loaders	Stage IIIA									463	844	1137	1382	1597		
Dump trucks	<1981															
Dump trucks	1981-1990															
Dump trucks	1991-Stage I	745	682	611	530	442	385	301	176							
Dump trucks	Stage I	186	292	407	530	552	642	752	880	943	739	514	319	150		
Dump trucks	Stage II					110	257	451	704	754	739	685	637	600		
Dump trucks	Stage IIIA									189	369	514	637	600		
Dump trucks	Stage IIIB														150	
Mini loaders	<1981															
Mini loaders	1981-1990	1273	990	684	354											
Mini loaders	1991-Stage I	3183	3301	3419	3537	3656	3063	2540	2055	1599	1178	809	504	259		
Mini loaders	Stage II		330	684	1061	1462	1701	1905	1761	1599	1413	1214	1009	777		
Mini loaders	Stage IIIA								294	533	707	809	841	777		
Telescopic loaders	1981-1990	398	348	265	149											
Telescopic loaders	1991-Stage I	994	1160	1326	1491	1657	1740	1837	1846	1687	1343	1009	732	466		
Telescopic loaders	Stage II		116	265	447	663	966	1378	1582	1687	1612	1514	1464	1397		
Telescopic loaders	Stage IIIA								264	562	806	1009	1220	1397		

Stock data for machine pools 1985-2012

EquipmentName (Eng)	Emission Level	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
Tractors (machine pools)	<1981	1236	627													
Tractors (machine pools)	1981-1990	3091	3763	4575	4515	4370	4100	3643	2808	2368	1786	1214	604			
Tractors (machine pools)	1991-Stage I							607	1123	1776	2382	3035	3624	4324	4210	4336
Tractors (machine pools)	Stage I															
Tractors (machine pools)	Stage II															
Tractors (machine pools)	Stage IIIA															

Tractors (machine pools)	Stage IIIB															
Harvesters (machine pools)	<1981	969	776	661	472	287	139									
Harvesters (machine pools)	1981-1990	807	932	1157	1257	1294	1385	1385	1197	927	794	712	512	421	282	162
Harvesters (machine pools)	1991-Stage I							139	266	348	454	593	615	737	751	729
Harvesters (machine pools)	Stage II															
Harvesters (machine pools)	Stage IIIA															
Harvesters (machine pools)	Stage IIIB															
Self-propelled vehicles (machine pools)	1981-1990									72	61	38				
Self-propelled vehicles (machine pools)	1991-Stage I									72	122	190	263	278	277	295
Self-propelled vehicles (machine pools)	Stage II															
Self-propelled vehicles (machine pools)	Stage IIIA															
Self-propelled vehicles (machine pools)	Stage IIIB															

EquipmentName (Eng)	Emission Level	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
Tractors (machine pools)	<1981													
Tractors (machine pools)	1981-1990													
Tractors (machine pools)	1991-Stage I	3956	4069	3323	2566	2066	1421	927	487					
Tractors (machine pools)	Stage I			554	513	517	474	464	487	524				
Tractors (machine pools)	Stage II				513	1033	1421	1855	1946	2094	1985	1535	947	473
Tractors (machine pools)	Stage IIIA								487	1047	1488	2046	2366	2366
Tractors (machine pools)	Stage IIIB													473
Harvesters (machine pools)	<1981													
Harvesters (machine pools)	1981-1990	78												
Harvesters (machine pools)	1991-Stage I	778	779	651	531	472	300	257	211	169	127	85	42	
Harvesters (machine pools)	Stage II			65	118	177	171	172	169	169	169	169	169	169
Harvesters (machine pools)	Stage IIIA							43	85	127	169	211	211	211
Harvesters (machine pools)	Stage IIIB												42	85
Self-propelled vehicles (machine pools)	1981-1990													
Self-propelled vehicles (machine pools)	1991-Stage I	289	314	237	203	153	99	49						
Self-propelled vehicles (machine pools)	Stage II			47	102	153	199	194	189	142	94	47		
Self-propelled vehicles (machine pools)	Stage IIIA							49	94	142	189	236	236	189
Self-propelled vehicles (machine pools)	Stage IIIB												47	94

Stock data for household and gardening 1985-2012.

EquipmentName (Eng)	Emission Level	SNAPCode	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
Lawn movers (private)	<1981	0809	253125	168750	84375												
Lawn movers (private)	1981-1990	0809	421875	506250	590625	675000	675000	675000	590625	506250	421875	337500	253125	168750	84375		
Lawn movers (private)	1991-Stage I	0809							84375	168750	253125	337500	421875	506250	590625	675000	675000
Lawn movers (private)	Stage I	0809															
Lawn movers (private)	Stage II	0809															
Cultivators (private-large)	<1981	0809	73333	66000	58667	51333	44000	36667	29333	22000	14667	7333					
Cultivators (private-large)	1981-1990	0809	36667	44000	51333	58667	66000	73333	73333	73333	73333	73333	73333	66000	58667	51333	44000
Cultivators (private-large)	1991-Stage I	0809							7333	14667	22000	29333	36667	44000	51333	58667	66000
Cultivators (private-large)	Stage II	0809															
Cultivators (private-small)	1981-1990	0809	10000	10000	10000	10000	10000	10000	8000	6000	4000	2000					
Cultivators (private-small)	1991-Stage I	0809							2000	4000	6000	8000	10000	10000	10000	10000	10000
Cultivators (private-small)	Stage II	0809															
Chain saws (private)	<1981	0809	125000	100000	75000	50000	25000										
Chain saws (private)	1981-1990	0809	125000	150000	175000	200000	225000	250000	227250	204000	180250	156000	131250	106000	80250	54000	27250
Chain saws (private)	1991-Stage I	0809							25250	51000	77250	104000	131250	159000	187250	216000	245250
Chain saws (private)	Stage I	0809															
Chain saws (private)	Stage II	0809															
Riders (private)	<1981	0809	40950	35100	29250	23400	17550	11700	5880								
Riders (private)	1981-1990	0809	29250	35100	40950	46800	52650	58500	58796	59388	54248	49167	44056	38828	33392	27660	21544
Riders (private)	1991-Stage I	0809							5880	11878	18083	24583	31469	38828	46748	55320	64631
Riders (private)	Stage I	0809															
Riders (private)	Stage II	0809															
Shrub clearers (private)	<1981	0809	24000	19200	14400	9600	4800										
Shrub clearers (private)	1981-1990	0809	24000	28800	33600	38400	43200	48000	47520	46080	43680	40320	36000	30720	24480	17280	9120
Shrub clearers (private)	1991-Stage I	0809							5280	11520	18720	26880	36000	46080	57120	69120	82080
Shrub clearers (private)	Stage I	0809															
Shrub clearers (private)	Stage II	0809															
Hedge cutters (private)	<1981	0809	6850	5480	4110	2740	1370										
Hedge cutters (private)	1981-1990	0809	6850	8220	9590	10960	12330	13700	15237	16128	16373	15972	14925	13232	10893	7908	4277
Hedge cutters (private)	1991-Stage I	0809							1693	4032	7017	10648	14925	19848	25417	31632	38493
Hedge cutters (private)	Stage I	0809															
Hedge cutters (private)	Stage II	0809															
Trimmers (private)	<1981	0809	25500	20400	15300	10200	5100										
Trimmers (private)	1981-1990	0809	25500	30600	35700	40800	45900	51000	48086	44686	40800	36429	31571	26229	20400	14086	7286
Trimmers (private)	1991-Stage I	0809							5343	11171	17486	24286	31571	39343	47600	56343	65571

Trimmers (private)	Stage I	0809															
Trimmers (private)	Stage II	0809															
Lawn movers (professional)	1981-1990	0811	25000	25000	25000	25000	25000	25000	18750	12500	6250						
Lawn movers (professional)	1991-Stage I	0811							6250	12500	18750	25000	25000	25000	25000	25000	25000
Lawn movers (professional)	Stage I	0811															
Lawn movers (professional)	Stage II	0811															
Cultivators (professional)	<1981	0811	3750	2500	1250												
Cultivators (professional)	1981-1990	0811	6250	7500	8750	10000	10000	10000	8750	7500	6250	5000	3750	2500	1250		
Cultivators (professional)	1991-Stage I	0811							1250	2500	3750	5000	6250	7500	8750	10000	10000
Cultivators (professional)	Stage I	0811															
Cultivators (professional)	Stage II	0811															
Chain saws (professional)	1981-1990	0811	10000	10000	10000	10000	10000	10000	7333	4000							
Chain saws (professional)	1991-Stage I	0811							3667	8000	13000	14000	15000	16000	17000	18000	19000
Chain saws (professional)	Stage I	0811															
Chain saws (professional)	Stage II	0811															
Riders (professional)	1981-1990	0811	4800	4800	4800	4800	4800	4800	3878	2966	2035	1056					
Riders (professional)	1991-Stage I	0811							970	1978	3053	4224	5520	5760	6000	6240	6480
Riders (professional)	Stage I	0811															
Riders (professional)	Stage II	0811															
Shrub clearers (professional)	1981-1990	0811	2000	2000	2000	2000	2000	2000	1650	1200	650						
Shrub clearers (professional)	1991-Stage I	0811							550	1200	1950	2800	3000	3200	3400	3600	3800
Shrub clearers (professional)	Stage I	0811															
Shrub clearers (professional)	Stage II	0811															
Hedge cutters (professional)	1981-1990	0811	1300	1300	1300	1300	1300	1300	1178	920	528						
Hedge cutters (professional)	1991-Stage I	0811							393	920	1583	2380	2650	2920	3190	3460	3730
Hedge cutters (professional)	Stage I	0811															
Hedge cutters (professional)	Stage II	0811															
Trimmers (professional)	1981-1990	0811	9000	9000	9000	9000	9000	9000	7071	4929	2571						
Trimmers (professional)	1991-Stage I	0811							2357	4929	7714	10714	11143	11571	12000	12429	12857
Trimmers (professional)	Stage I	0811															
Trimmers (professional)	Stage II	0811															

EquipmentName (Eng)	Emission Level	SNAPCode	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
Lawn movers (private)	<1981	0809													
Lawn movers (private)	1981-1990	0809													
Lawn movers (private)	1991-Stage I	0809	675000	675000	675000	675000	675000	595000	513750	428125	342500	256875	171250	85625	

Lawn movers (private)	Stage I	0809							85000	171250	256875	256875	256875	256875	256875	256875
Lawn movers (private)	Stage II	0809										85625	171250	256875	342500	428125
Cultivators (private-large)	<1981	0809														
Cultivators (private-large)	1981-1990	0809	36667	29333	22000	14667	7333									
Cultivators (private-large)	1991-Stage I	0809	73333	80667	88000	95333	102667	102667	95333	88000	80667	73333	66000	58667	51333	
Cultivators (private-large)	Stage II	0809						7333	14667	22000	29333	36667	44000	51333	58667	
Cultivators (private-small)	1981-1990	0809														
Cultivators (private-small)	1991-Stage I	0809	10000	10000	10000	10000	10000	8000	6000	4000	2000					
Cultivators (private-small)	Stage II	0809						2000	4000	6000	8000	10000	10000	10000	10000	
Chain saws (private)	<1981	0809														
Chain saws (private)	1981-1990	0809														
Chain saws (private)	1991-Stage I	0809	275000	280750	286500	292250	298000	268200	238400	208600	178800	149000	119200	89400	59600	
Chain saws (private)	Stage I	0809						29800	59600	89400	89400	89400	89400	89400	89400	
Chain saws (private)	Stage II	0809									29800	59600	89400	119200	149000	
Riders (private)	<1981	0809														
Riders (private)	1981-1990	0809	14954	7910												
Riders (private)	1991-Stage I	0809	74771	87015	101775	109920	119360	117741	114313	107663	99047	86666	74285	61904	49523	
Riders (private)	Stage I	0809						10704	22863	23925	24762	24762	24762	24762	24762	
Riders (private)	Stage II	0809								11963	24762	37143	49523	61904	74285	
Shrub clearers (private)	<1981	0809														
Shrub clearers (private)	1981-1990	0809														
Shrub clearers (private)	1991-Stage I	0809	96000	107000	118000	129000	140000	126000	112000	98000	84000	70000	56000	42000	28000	
Shrub clearers (private)	Stage I	0809						14000	28000	42000	42000	42000	42000	42000	42000	
Shrub clearers (private)	Stage II	0809									14000	28000	42000	56000	70000	
Hedge cutters (private)	<1981	0809														
Hedge cutters (private)	1981-1990	0809														
Hedge cutters (private)	1991-Stage I	0809	46000	52900	59800	66700	73600	66240	58880	51520	44160	36800	29440	22080	14720	
Hedge cutters (private)	Stage I	0809						7360	14720	22080	22080	22080	22080	22080	22080	
Hedge cutters (private)	Stage II	0809									7360	14720	22080	29440	36800	
Trimmers (private)	<1981	0809														
Trimmers (private)	1981-1990	0809														
Trimmers (private)	1991-Stage I	0809	75286	77714	80143	82571	85000	76500	68000	59500	51000	42500	34000	25500	17000	
Trimmers (private)	Stage I	0809						8500	17000	25500	25500	25500	25500	25500	25500	
Trimmers (private)	Stage II	0809									8500	17000	25500	34000	42500	
Lawn movers (professional)	1981-1990	0811														
Lawn movers (professional)	1991-Stage I	0811	25000	25000	25000	25000	25000	18750	12500	6250						
Lawn movers (professional)	Stage I	0811						6250	12500	18750	18750	12500	6250			

Lawn movers (professional)	Stage II	0811									6250	12500	18750	25000	25000
Cultivators (professional)	<1981	0811													
Cultivators (professional)	1981-1990	0811													
Cultivators (professional)	1991-Stage I	0811	10000	10000	10000	10000	10000	8750	7500	6250	5000	3750	2500	1250	
Cultivators (professional)	Stage I	0811						1250	2500	3750	3750	3750	3750	3750	3750
Cultivators (professional)	Stage II	0811									1250	2500	3750	5000	6250
Chain saws (professional)	1981-1990	0811													
Chain saws (professional)	1991-Stage I	0811	20000	27500	35000	42500	50000	33333	16667						
Chain saws (professional)	Stage I	0811						16667	33333	50000	50000	33333	16667		
Chain saws (professional)	Stage II	0811										16667	33333	50000	50000
Riders (professional)	1981-1990	0811													
Riders (professional)	1991-Stage I	0811	6720	7802	9726	12492	16100	15728	13398	9444	4800				
Riders (professional)	Stage I	0811						3932	8932	9444	9600	9600	4800		
Riders (professional)	Stage II	0811								4722	9600	14400	19200	24000	24000
Shrub clearers (professional)	1981-1990	0811													
Shrub clearers (professional)	1991-Stage I	0811	4000	5500	7000	8500	10000	7500	5000	2500					
Shrub clearers (professional)	Stage I	0811						2500	5000	7500	7500	5000	2500		
Shrub clearers (professional)	Stage II	0811									2500	5000	7500	10000	10000
Hedge cutters (professional)	1981-1990	0811													
Hedge cutters (professional)	1991-Stage I	0811	4000	4600	5200	5800	6400	4800	3200	1600					
Hedge cutters (professional)	Stage I	0811						1600	3200	4800	4800	3200	1600		
Hedge cutters (professional)	Stage II	0811									1600	3200	4800	6400	6400
Trimmers (professional)	1981-1990	0811													
Trimmers (professional)	1991-Stage I	0811	13286	13714	14143	14571	15000	11250	7500	3750					
Trimmers (professional)	Stage I	0811						3750	7500	11250	11250	7500	3750		
Trimmers (professional)	Stage II	0811									3750	7500	11250	15000	15000

Stock data for small boats and pleasure crafts 1985-2012.

Brændstof	Motortakt	Boat type	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
Diesel		Motor boats (27-34 ft)	1550	1550	1719	1889	2058	2228	2397	2567	2736	2906	3075	3244	3414	3583	3753
Diesel		Motor boats (> 34 ft)	450	450	503	556	608	661	714	767	819	872	925	978	1031	1083	1136
Diesel		Motor boats <(27 ft)	3000	3000	3000	3000	3000	3000	3000	3000	3000	3000	3000	3000	3000	3000	3000
Diesel		Motor sailors	3500	3500	3583	3667	3750	3833	3917	4000	4083	4167	4250	4333	4417	4500	4583
Diesel		Sailing boats (> 26 ft)	7500	7500	7917	8333	8750	9167	9583	10000	10417	10833	11250	11667	12083	12500	12917
Benzin	2-takt	Other boats (< 20 ft)	4000	4000	4056	4111	4167	4222	4278	4333	4389	4444	4500	4556	4565	4527	4439
Benzin	2-takt	Yawls and cabin boats	4000	4000	4056	4111	4167	4222	4278	4333	4389	4444	4500	4556	4565	4527	4439
Benzin	2-takt	Sailing boats (< 26 ft)	19000	19000	18778	18556	18333	18111	17889	17667	17444	17222	17000	16778	16390	15843	15144
Benzin	2-takt	Speed boats	3000	3000	3000	3000	3000	3000	3000	3000	3000	3000	3000	3000	2970	2910	2820
Benzin	2-takt	Water scooters	1000	1000	1000	1000	1000	1000	1000	1000	1000	1000	1000	1000	990	970	940
Benzin	4-takt	Other boats (< 20 ft)														46	140
Benzin	4-takt	Yawls and cabin boats														46	140
Benzin	4-takt	Sailing boats (< 26 ft)														166	490
Benzin	4-takt	Speed boats	3000	3000	3000	3000	3000	3000	3000	3000	3000	3000	3000	3000	3000	3000	3000
Benzin	4-takt	Speed boats														30	90
Benzin	4-takt	Water scooters														10	30

Brændstof	Motortakt	Boat type	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
Diesel		Motor boats (27-34 ft)	3922	4092	4261	4431	4600	4600	4600	4600	4600	4600	4600	4600	4600
Diesel		Motor boats (> 34 ft)	1189	1242	1294	1347	1400	1400	1400	1400	1400	1400	1400	1400	1400
Diesel		Motor boats <(27 ft)	3000	3000	3000	3000	3000	3000	3000	3000	3000	3000	3000	3000	3000
Diesel		Motor sailors	4667	4750	4833	4917	5000	5000	5000	5000	5000	5000	5000	5000	5000
Diesel		Sailing boats (> 26 ft)	13333	13750	14167	14583	15000	15000	15000	15000	15000	15000	15000	15000	15000
Benzin	2-takt	Other boats (< 20 ft)	4300	4108	3862	3560	3200	2750	2250	1800	1400	1050	750	500	300
Benzin	2-takt	Yawls and cabin boats	4300	4108	3862	3560	3200	2750	2250	1800	1400	1050	750	500	300
Benzin	2-takt	Sailing boats (< 26 ft)	14300	13317	12201	10960	9600	8250	6750	5400	4200	3150	2250	1500	900
Benzin	2-takt	Speed boats	2700	2550	2370	2160	1920	1650	1350	1080	840	630	450	300	180
Benzin	2-takt	Water scooters	900	850	790	720	640	550	450	360	280	210	150	100	60
Benzin	4-takt	Other boats (< 20 ft)	478	725	1027	1384	1800	2250	2750	3200	3600	3950	4250	4500	4700
Benzin	4-takt	Yawls and cabin boats	478	725	1027	1384	1800	2250	2750	3200	3600	3950	4250	4500	4700
Benzin	4-takt	Sailing boats (< 26 ft)	1589	2350	3243	4262	5400	6750	8250	9600	10800	11850	12750	13500	14100
Benzin	4-takt	Speed boats	3000	3000	3000	3000	3000	3000	3000	3000	3000	3000	3000	3000	3000
Benzin	4-takt	Speed boats	300	450	630	840	1080	1350	1650	1920	2160	2370	2550	2700	2820
Benzin	4-takt	Water scooters	100	150	210	280	360	450	550	640	720	790	850	900	940

Engine sizes (kW) for recreational craft 1985-2012.

Motor type	Boat type	1985	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004-2012
2-takt	Other boats (< 20 ft)	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8
2-takt	Yawls and cabin boats	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20
2-takt	Sailing boats (< 26 ft)	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10
2-takt	Speed boats	25	31	32	33	35	36	38	39	40	42	43	44	46	47	49	50
2-takt	Water scooters	45	45	45	45	45	45	45	45	45	45	45	45	45	45	45	45
4-takt	Other boats (< 20 ft)									8	8	8	8	8	8	8	8
4-takt	Yawls and cabin boats									20	20	20	20	20	20	20	20
4-takt	Sailing boats (< 26 ft)									10	10	10	10	10	10	10	10
4-takt	Speed boats (in board eng.)	45	55	58	60	63	65	68	70	73	75	78	80	83	85	88	90
4-takt	Speed boats (out board eng.)									40	42	43	44	46	47	49	50
4-takt	Water scooters									45	45	45	45	45	45	45	45
Diesel	Motor boats (27-34 ft)	70	88	92	97	101	106	110	114	119	123	128	132	137	141	146	150
Diesel	Motor boats (> 34 ft)	120	149	156	163	171	178	185	192	199	207	214	221	228	236	243	250
Diesel	Motor boats <(27 ft)	20	24	26	27	28	29	30	31	32	33	34	36	37	38	39	40
Diesel	Motor sailers	20	22	23	23	24	24	25	26	26	27	27	28	28	29	29	30
Diesel	Sailing boats (> 26 ft)	30	30	30	30	30	30	30	30	30	30	30	30	30	30	30	30

Annex 3B-12 Traffic data and different technical and operational data for Danish domestic ferries

Annual traffic data for ferries (no. of round trips) for Danish domestic ferries.

Domestic ferry lines	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999			
Korsør-Nyborg, DSB	9305	9167	9237	8959	8813	8789	8746	3258	0	0			
Korsør-Nyborg, Vognmandsruten	7512	7363	7468	7496	7502	7828	7917	8302	3576	0			
Halsskov-Knudshoved	10601	10582	11701	11767	12420	12970	13539	13612	5732	0			
Kalundborg-Juelsminde	0	1326	1733	1542	1541	1508	856	0	0	0			
Kalundborg-Århus	1907	2400	3162	2921	2913	3540	4962	4888	4483	1454			
Sjællands Odde-Ebeltoft	3908	3978	4008	3988	4325	4569	5712	8153	7851	7720			
Sjællands Odde-Århus	0	0	0	0	0	0	0	0	0	2339			
Hundested-Grenaa	1026	1025	1032	1030	718	602	67	0	0	0			
København-Rønne	558	545	484	412	427	426	437	465	458	506			
Køge-Rønne	0	0	0	0	0	0	0	0	0	0			
Kalundborg-Samsø	873	873	860	881	826	811	813	823	824	850			
Tårs-Spødsbjerg	7656	8835	9488	9535	9402	9562	9000	9129	7052	6442			
Hirtshals-Torshavn	0	0	0	0	0	0	0	0	0	0			
Hanstholm-Torshavn	0	14	15	0	0	0	0	0	0	48			
Esbjerg-Torshavn	9	9	9	15	14	13	0	0	0	0			
Local ferries	176891	179850	181834	178419	202445	209129	182750	197489	200027	202054			
<i>Continued</i>	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
Korsør-Nyborg, DSB	0	0	0	0	0	0	0	0	0	0	0	0	0
Korsør-Nyborg, Vognmandsruten	0	0	0	0	0	0	0	0	0	0	0	0	0
Halsskov-Knudshoved	0	0	0	0	0	0	0	0	0	0	0	0	0
Kalundborg-Juelsminde	0	0	0	0	0	0	0	0	0	0	0	0	0
Kalundborg-Århus	1870	1804	2037	1800	1750	1725	1724	1695	1694	1668	1552	1158	845
Sjællands Odde-Ebeltoft	4775	4226	3597	3191	2906	2889	2690	2670	2577	2454	2409	1960	844
Sjællands Odde-Århus	1799	1817	1825	2359	2863	2795	2853	2810	2814	2810	2735	2796	2304
Hundested-Grenaa	0	0	0	0	0	0	0	0	0	0	0	0	0
København-Rønne	491	430	413	397	293	0	0	0	0	0	0	0	0
Køge-Rønne	0	0	0	0	154	488	436	399	428	407	459	365	370
Kalundborg-Samsø	828	817	833	831	841	867	862	887	921	969	937	919	927
Tårs-Spødsbjerg	6477	6498	6468	6516	6497	6494	6460	6493	6504	6474	6529	6185	6235
Hirtshals-Torshavn	0	0	0	0	0	0	0	0	0	0	13	63	58
Hanstholm-Torshavn	67	94	85	50	59	51	51	48	52	27	20	0	0
Esbjerg-Torshavn	0	0	0	0	0	0	0	0	0	35	30	0	0
Local ferries	201833	200130	208396	208501	206297	205564	203413	205260	210089	209082	205461	202510	204036

Ferry data: Service, name, engine year, main engine MCR (kW), engine type, specific fuel consumption (sfc), aux. engine (kW).

Ferry service	Ferry name	Engine year	Main engine MCR (kW)	Engine type	Sfc (g/kWh)	Fuel type	Aux engine (kW)
Esbjerg-Torshavn	Gamle Norrøna	1973	11768	Medium speed (4-stroke)	239	Diesel	2354
Esbjerg-Torshavn	Nye Norrøna	2003	21600	Medium speed (4-stroke)	190	Fuel	4320
Halsskov-Knudshoved	ARVEPRINS KNUD DRONNING MARGRETHE	1963	8238	Slow speed (2-stroke)	220	Fuel	1666
Halsskov-Knudshoved	II	1973	8826	Medium speed (4-stroke)	230	Diesel	1692
Halsskov-Knudshoved	HEIMDAL	1983	8309	Medium speed (4-stroke)	220	Diesel	740
Halsskov-Knudshoved	KNUDSHOVED	1961	6400	Slow speed (2-stroke)	220	Fuel	1840
Halsskov-Knudshoved	KONG FREDERIK IX	1954	6767	Slow speed (2-stroke)	225	Fuel	1426
Halsskov-Knudshoved	KRAKA	1982	8309	Medium speed (4-stroke)	220	Diesel	740
Halsskov-Knudshoved	LODBROG	1982	8309	Medium speed (4-stroke)	220	Diesel	740
Halsskov-Knudshoved	PRINSESSE ANNE-MARIE	1960	8238	Slow speed (2-stroke)	220	Fuel	1360
Halsskov-Knudshoved	PRINSESSE ELISABETH	1964	8238	Slow speed (2-stroke)	220	Fuel	1360
Halsskov-Knudshoved	ROMSØ	1973	8826	Medium speed (4-stroke)	230	Diesel	1728
Halsskov-Knudshoved	SPROGØ	1962	6400	Slow speed (2-stroke)	220	Fuel	1840
Hanstholm-Torshavn	Gamle Norrøna	1973	11768	Medium speed (4-stroke)	239	Diesel	2354
Hanstholm-Torshavn	Nye Norrøna	2003	21600	Medium speed (4-stroke)	190	Fuel	4320
Hirtshals-Torshavn	Nye Norrøna	2003	21600	Medium speed (4-stroke)	190	Fuel	4320
Hundested-Grenaa	DJURSLAND	1974	9856	Medium speed (4-stroke)	230	Diesel	900
Hundested-Grenaa	KATTEGAT	1995	23200	High speed (4-stroke)	205	Diesel	1223
Hundested-Grenaa	KONG FREDERIK IX	1954	6767	Slow speed (2-stroke)	235	Fuel	1375
Hundested-Grenaa	PRINSESSE ANNE-MARIE	1960	8238	Slow speed (2-stroke)	220	Fuel	1360
Kalundborg-Juelsminde	Mercandia I	1989	2950	High speed (4-stroke)	220	Diesel	0
Kalundborg-Juelsminde	Mercandia II	1989	2950	High speed (4-stroke)	220	Diesel	0
Kalundborg-Juelsminde	Mercandia III	1989	2950	High speed (4-stroke)	220	Diesel	0
Kalundborg-Juelsminde	Mercandia IV	1989	2950	High speed (4-stroke)	220	Diesel	0
Kalundborg-Samsø	HOLGER DANSKE	1976	2354	High speed (4-stroke)	225	Diesel	600
Kalundborg-Samsø	KALUNDBORG	1952	3825	Slow speed (2-stroke)	235	Fuel	570
Kalundborg-Samsø	KYHOLM	1998	2940	High speed (4-stroke)	195	Diesel	864
Kalundborg-Samsø	VESBORG	1995	1770	High speed (4-stroke)	200	Diesel	494
Kalundborg-Århus	ASK	1984	8826	Medium speed (4-stroke)	215	Diesel	2220
Kalundborg-Århus	ASK	1984	8826	Medium speed (4-stroke)	215	Diesel	3000
Kalundborg-Århus	ASK	1984	9840	Medium speed (4-stroke)	215	Diesel	3000
Kalundborg-Århus	CAT-LINK I	1995	17280	High speed (4-stroke)	205	Diesel	1160
Kalundborg-Århus	CAT-LINK II	1995	17280	High speed (4-stroke)	205	Diesel	1160
Kalundborg-Århus	CAT-LINK III	1995	22000	High speed (4-stroke)	205	Diesel	800
Kalundborg-Århus	CAT-LINK IV	1998	28320	High speed (4-stroke)	205	Diesel	920

Kalundborg-Århus	CAT-LINK V	1998	28320	High speed (4-stroke)	205	Diesel	920
Kalundborg-Århus	KATTEGAT SYD	1979	7650	Medium speed (4-stroke)	225	Diesel	1366
Kalundborg-Århus	KNUDSHOVED	1961	6400	Slow speed (2-stroke)	220	Fuel	1840
Kalundborg-Århus	KONG FREDERIK IX	1954	6767	Slow speed (2-stroke)	225	Fuel	1426
Kalundborg-Århus	KRAKA	1982	8309	Medium speed (4-stroke)	220	Diesel	740
Kalundborg-Århus	MAREN MOLS	1996	11700	Slow speed (2-stroke)	180	Diesel	2530
Kalundborg-Århus	METTE MOLS	1996	11700	Slow speed (2-stroke)	180	Diesel	2530
Kalundborg-Århus	NIELS KLIM	1986	12474	Slow speed (2-stroke)	215	Fuel	4440
Kalundborg-Århus	PEDER PAARS	1985	12474	Slow speed (2-stroke)	215	Fuel	4440
Kalundborg-Århus	PRINSESSE ELISABETH	1964	8238	Slow speed (2-stroke)	220	Fuel	1360
Kalundborg-Århus	ROSTOCK LINK	1975	8385	Medium speed (4-stroke)	230	Diesel	2500
Kalundborg-Århus	SØLØVEN/SØBJØRNEN	1992	4000	High speed (4-stroke)	210	Diesel	272
Kalundborg-Århus	URD	1981	8826	Medium speed (4-stroke)	215	Diesel	2220
Kalundborg-Århus	URD	1981	8826	Medium speed (4-stroke)	215	Diesel	3000
Kalundborg-Århus	URD	1981	9840	Medium speed (4-stroke)	215	Diesel	3000
Korsør-Nyborg, DSB	ASA-THOR	1965	6472	Slow speed (2-stroke)	220	Fuel	1305
Korsør-Nyborg, DSB	DRONNING INGRID DRONNING MARGRETHE II	1980	18720	Medium speed (4-stroke)	220	Diesel	2932
Korsør-Nyborg, DSB	II	1973	8826	Medium speed (4-stroke)	230	Diesel	1692
Korsør-Nyborg, DSB	KONG FREDERIK IX	1954	6767	Slow speed (2-stroke)	225	Fuel	1426
Korsør-Nyborg, DSB	KRONPRINS FREDERIK	1981	18720	Medium speed (4-stroke)	220	Diesel	2932
Korsør-Nyborg, DSB	PRINS JOACHIM	1980	18720	Medium speed (4-stroke)	220	Diesel	2932
Korsør-Nyborg, DSB	SPROGØ/KNUDSHOVED	1962	6400	Slow speed (2-stroke)	220	Fuel	1840
Korsør-Nyborg, Vognmandsruten	Superflex Alfa	1989	2950	High speed (4-stroke)	220	Diesel	0
Korsør-Nyborg, Vognmandsruten	Superflex Bravo	1989	2950	High speed (4-stroke)	220	Diesel	0
Korsør-Nyborg, Vognmandsruten	Superflex Charlie	1988	2950	High speed (4-stroke)	220	Diesel	0
København-Rønne	JENS KOFOED	1979	12950	Medium speed (4-stroke)	233	Fuel	2889
København-Rønne	POVL ANKER	1979	12950	Medium speed (4-stroke)	233	Fuel	2889
Køge-Rønne	DUEODDE	2005	8640	Medium speed (4-stroke)	190	Fuel	1545
Køge-Rønne	HAMMERODDE	2005	8640	Medium speed (4-stroke)	190	Fuel	1545
Køge-Rønne	JENS KOFOED	1979	12950	Medium speed (4-stroke)	233	Fuel	2889
Køge-Rønne	POVL ANKER	1979	12950	Medium speed (4-stroke)	233	Fuel	2889
Sjællands Odde-Ebeltoft	MAI MOLS	1996	24800	Gas turbine	263	Diesel	752
Sjællands Odde-Ebeltoft	MAREN MOLS	1975	12062	Medium speed (4-stroke)	230	Fuel	1986
Sjællands Odde-Ebeltoft	MAREN MOLS 2	1996	11700	Slow speed (2-stroke)	180	Diesel	2530
Sjællands Odde-Ebeltoft	MAX MOLS	1998	28320	High speed (4-stroke)	209	Diesel	920
Sjællands Odde-Ebeltoft	METTE MOLS	1975	12062	Medium speed (4-stroke)	230	Fuel	1986
Sjællands Odde-Ebeltoft	METTE MOLS 2	1996	11700	Slow speed (2-stroke)	180	Diesel	2530

Sjællands Odde-Ebeltoft	MIE MOLS	1971	5884	Medium speed (4-stroke)	230	Diesel	
Sjællands Odde-Ebeltoft	MIE MOLS 2	1996	24800	Gas turbine	263	Diesel	752
Sjællands Odde-Århus	KatExpress	2009	36000	High speed (4-stroke)	190	Diesel	1440
Sjællands Odde-Århus	MADS MOLS	1998	28320	High speed (4-stroke)	205	Diesel	920
Sjællands Odde-Århus	MAI MOLS	1996	24800	Gas turbine	263	Diesel	752
Sjællands Odde-Århus	MAX MOLS	1998	28320	High speed (4-stroke)	209	Diesel	920
Sjællands Odde-Århus	MIE MOLS	1996	24800	Gas turbine	263	Diesel	752
Tårs-Spødsbjerg	FRIGG SYDFYEN	1984	1300	Medium speed (4-stroke)	220	Diesel	780
Tårs-Spødsbjerg	Langeland	2011	875	High speed	210	Diesel	
Tårs-Spødsbjerg	Lolland	2011	875	High speed	210	Diesel	
Tårs-Spødsbjerg	ODIN SYDFYEN	1982	1180	Medium speed (4-stroke)	220	Diesel	780
Tårs-Spødsbjerg	SPODSBJERG	1972	1530	Medium speed (4-stroke)	225	Diesel	300
Tårs-Spødsbjerg	SPODSBJERG	2006	1530	Medium speed (4-stroke)	190	Diesel	300
Tårs-Spødsbjerg	THOR SYDFYEN	1978	1176	Medium speed (4-stroke)	225	Diesel	300

Ferry data: Sailing time (single trip).

Ferry service	Ferry name	1985-1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
Esbjerg-Torshavn	Gamle Norrøna	1860	1860	1860	1860	1860	1860	1860	1860	1860	1860
Esbjerg-Torshavn	Nye Norrøna										
Halsskov-Knudshoved	ARVEPRINS KNUD	60	60	60	60	60	60	60	60	60	
Halsskov-Knudshoved	DRONNING MARGRETHE II	60	60	60	60	60	60	60	60	60	
Halsskov-Knudshoved	HEIMDAL	60	60	60	60	60	60	60	60	60	
Halsskov-Knudshoved	KNUDSHOVED	60	60	60	60	60	60	60	60	60	
Halsskov-Knudshoved	KONG FREDERIK IX	60	60	60	60	60	60	60	60	60	
Halsskov-Knudshoved	KRAKA	60	60	60	60	60	60	60	60	60	
Halsskov-Knudshoved	LODBROG	60	60	60	60	60	60	60	60	60	
Halsskov-Knudshoved	PRINSESSE ANNE-MARIE	60	60	60	60	60	60	60	60	60	
Halsskov-Knudshoved	PRINSESSE ELISABETH	60	60	60	60	60	60	60	60	60	
Halsskov-Knudshoved	ROMSØ	60	60	60	60	60	60	60	60	60	
Halsskov-Knudshoved	SPROGØ	60	60	60	60	60	60	60	60	60	
Hanstholm-Torshavn	Gamle Norrøna	1740	1740	1740	1740	1740	1740	1740	1740	1740	1740
Hanstholm-Torshavn	Nye Norrøna										
Hirtshals-Torshavn	Nye Norrøna										
Hundested-Grenaa	DJURLAND	160	160	160	160	160					
Hundested-Grenaa	KATTEGAT						90	90			
Hundested-Grenaa	KONG FREDERIK IX					170					
Hundested-Grenaa	PRINSESSE ANNE-MARIE					165					

Kalundborg-Juelsminde	Mercandia I	160	160	160	160	160	160	160			
Kalundborg-Juelsminde	Mercandia II	160	160	160	160	160	160	160			
Kalundborg-Juelsminde	Mercandia III	160	160	160	160	160	160	160			
Kalundborg-Juelsminde	Mercandia IV	160	160	160	160	160	160	160			
Kalundborg-Samsø	HOLGER DANSKE			120	120	120	120	120	120	120	
Kalundborg-Samsø	KALUNDBORG	120	120	120							
Kalundborg-Samsø	KYHOLM									110	110
Kalundborg-Samsø	VESBORG									120	
Kalundborg-Århus	ASK		195	195	195	195	195	195	195	195	195
Kalundborg-Århus	CAT-LINK I						80	85	90	95	
Kalundborg-Århus	CAT-LINK II						80	85	90	95	
Kalundborg-Århus	CAT-LINK III							85	90	95	
Kalundborg-Århus	CAT-LINK IV									80	80
Kalundborg-Århus	CAT-LINK V									80	80
Kalundborg-Århus	KATTEGAT SYD										195
Kalundborg-Århus	KNUDSHOVED		190								
Kalundborg-Århus	KONG FREDERIK IX		190	190	190	190	190	190			
Kalundborg-Århus	KRAKA									195	
Kalundborg-Århus	MAREN MOLS										
Kalundborg-Århus	METTE MOLS										
Kalundborg-Århus	NIELS KLIM	185	185								
Kalundborg-Århus	PEDER PAARS	185	185								
Kalundborg-Århus	PRINSESSE ELISABETH		185								
Kalundborg-Århus	ROSTOCK LINK										195
Kalundborg-Århus	SØLØVEN/SØBJØRNEN		90	90	90	90	90	90			
Kalundborg-Århus	URD		195	195	195	195	195	195	195	195	195
Korsør-Nyborg, DSB	ASA-THOR	65	65	65	65	65	65	65	65		
Korsør-Nyborg, DSB	DRONNING INGRID	65	65	65	65	65	65	65	65		
Korsør-Nyborg, DSB	DRONNING MARGRETHE II	65	65	65	65	65	65	65	65		
Korsør-Nyborg, DSB	KONG FREDERIK IX	75	75	75	75	75	75	75	75		
Korsør-Nyborg, DSB	KRONPRINS FREDERIK	65	65	65	65	65	65	65	65		
Korsør-Nyborg, DSB	PRINS JOACHIM	65	65	65	65	65	65	65	65		
Korsør-Nyborg, DSB	SPROGØ/KNUDSHOVED	75	75	75	75	75	75	75	75		
Korsør-Nyborg, Vognmandsruten	Superflex Alfa	70	70	70	70	70	70	70	70	70	
Korsør-Nyborg, Vognmandsruten	Superflex Bravo	70	70	70	70	70	70	70	70	70	
Korsør-Nyborg, Vognmandsruten	Superflex Charlie	70	70	70	70	70	70	70	70	70	
København-Rønne	JENS KOFOED	420	420	420	420	420	420	420	420	420	420

København-Rønne	POVL ANKER	420	420	420	420	420	420	420	420	420	420	420
Køge-Rønne	DUEODDE											
Køge-Rønne	HAMMERODDE											
Køge-Rønne	JENS KOFOED											
Køge-Rønne	POVL ANKER											
Sjællands Odde-Ebeltoft	MAI MOLS								45	45	45	45
Sjællands Odde-Ebeltoft	MAREN MOLS	100	100	100	100	100	100	100				
Sjællands Odde-Ebeltoft	MAREN MOLS 2								100	100	100	95
Sjællands Odde-Ebeltoft	MAX MOLS											
Sjællands Odde-Ebeltoft	METTE MOLS	100	100	100	100	100	100	100				
Sjællands Odde-Ebeltoft	METTE MOLS 2								100	100	100	95
Sjællands Odde-Ebeltoft	MIE MOLS	105	105	105	105	105	105	105				
Sjællands Odde-Ebeltoft	MIE MOLS 2								45	45	45	45
Sjællands Odde-Århus	KatExpress											
Sjællands Odde-Århus	MADS MOLS											60
Sjællands Odde-Århus	MAI MOLS											
Sjællands Odde-Århus	MAX MOLS											60
Sjællands Odde-Århus	MIE MOLS											
Tårs-Spødsbjerg	FRIGG SYDFYEN	45	45	45	45	45	45	45	45	45	45	45
Tårs-Spødsbjerg	Langeland											
Tårs-Spødsbjerg	Lolland											
Tårs-Spødsbjerg	ODIN SYDFYEN	45	45	45	45	45	45	45	45	45	45	45
Tårs-Spødsbjerg	SPODSBJERG	45	45	45	45	45	45	45	45	45	45	45
Tårs-Spødsbjerg	THOR SYDFYEN	45	45	45	45	45	17	45	45	45		

Ferry service	Ferry name	2000	2001	2002	2003	2004	2005	2006-2008	2009	2010	2011	2012
Esbjerg-Torshavn	Gamle Norrøna	1860	1860	1860								
Esbjerg-Torshavn	Nye Norrøna				1860	1860	1860		1860	1860	1860	1860
Halsskov-Knudshoved	ARVEPRINS KNUD											
Halsskov-Knudshoved	DRONNING MARGRETHE II											
Halsskov-Knudshoved	HEIMDAL											
Halsskov-Knudshoved	KNUDSHOVED											
Halsskov-Knudshoved	KONG FREDERIK IX											
Halsskov-Knudshoved	KRAKA											
Halsskov-Knudshoved	LODBROG											
Halsskov-Knudshoved	PRINSESSE ANNE-MARIE											

Halsskov-Knudshoved	PRINSESSE ELISABETH													
Halsskov-Knudshoved	ROMSØ													
Halsskov-Knudshoved	SPROGØ													
Hanstholm-Torshavn	Gamle Norrøna	1740	1740	1740										
Hanstholm-Torshavn	Nye Norrøna				1740	1740	1740		1740	1740	1740	1740	1740	
Hirtshals-Torshavn	Nye Norrøna										1740	1740	1740	
Hundested-Grenaa	DJURSLAND													
Hundested-Grenaa	KATTEGAT													
Hundested-Grenaa	KONG FREDERIK IX													
Hundested-Grenaa	PRINSESSE ANNE-MARIE													
Kalundborg-Juelsminde	Mercandia I													
Kalundborg-Juelsminde	Mercandia II													
Kalundborg-Juelsminde	Mercandia III													
Kalundborg-Juelsminde	Mercandia IV													
Kalundborg-Samsø	HOLGER DANSKE													
Kalundborg-Samsø	KALUNDBORG													
Kalundborg-Samsø	KYHOLM	110	110	110	110	110	110		110	110	110	110	110	
Kalundborg-Samsø	VESBORG													
Kalundborg-Århus	ASK													
Kalundborg-Århus	CAT-LINK I													
Kalundborg-Århus	CAT-LINK II													
Kalundborg-Århus	CAT-LINK III													
Kalundborg-Århus	CAT-LINK IV													
Kalundborg-Århus	CAT-LINK V													
Kalundborg-Århus	KATTEGAT SYD													
Kalundborg-Århus	KNUDSHOVED													
Kalundborg-Århus	KONG FREDERIK IX													
Kalundborg-Århus	KRAKA													
Kalundborg-Århus	MAREN MOLS	160	160	155	155	155	155		165	165	165	165	165	
Kalundborg-Århus	METTE MOLS	160	160	155	155	155	155		165	165	165	165	165	
Kalundborg-Århus	NIELS KLIM													
Kalundborg-Århus	PEDER PAARS													
Kalundborg-Århus	PRINSESSE ELISABETH													
Kalundborg-Århus	ROSTOCK LINK													
Kalundborg-Århus	SØLØVEN/SØBJØRNEN													
Kalundborg-Århus	URD													
Korsør-Nyborg, DSB	ASA-THOR													

Korsør-Nyborg, DSB	DRONNING INGRID													
Korsør-Nyborg, DSB	DRONNING MARGRETHE II													
Korsør-Nyborg, DSB	KONG FREDERIK IX													
Korsør-Nyborg, DSB	KRONPRINS FREDERIK													
Korsør-Nyborg, DSB	PRINS JOACHIM													
Korsør-Nyborg, DSB	SPROGØ/KNUDSHOVED													
Korsør-Nyborg, Vognmandsruten	Superflex Alfa													
Korsør-Nyborg, Vognmandsruten	Superflex Bravo													
Korsør-Nyborg, Vognmandsruten	Superflex Charlie													
København-Rønne	JENS KOFOED	420	420	420	420	420								
København-Rønne	POVL ANKER	420	420	420	420	420								
Køge-Rønne	DUEODDE							375	375	375	375			
Køge-Rønne	HAMMERODDE							375	375	375	375	375	375	
Køge-Rønne	JENS KOFOED							375	375					
Køge-Rønne	POVL ANKER							375	375	375				
Sjællands Odde-Ebeltoft	MAI MOLS	45	45	45	45	45	45		50	50	50	50	50	
Sjællands Odde-Ebeltoft	MAREN MOLS													
Sjællands Odde-Ebeltoft	MAREN MOLS 2													
Sjællands Odde-Ebeltoft	MAX MOLS													55
Sjællands Odde-Ebeltoft	METTE MOLS													
Sjællands Odde-Ebeltoft	METTE MOLS 2													
Sjællands Odde-Ebeltoft	MIE MOLS													
Sjællands Odde-Ebeltoft	MIE MOLS 2	45	45	45	45	45	45		50	50	50	50	50	
Sjællands Odde-Århus	KatExpress													72
Sjællands Odde-Århus	MADS MOLS	65	65	65	65	65	65							
Sjællands Odde-Århus	MAI MOLS			65	65	65	65		68	68	68	68	68	
Sjællands Odde-Århus	MAX MOLS	65	65	65	65	65	65		70	70	70	70	70	
Sjællands Odde-Århus	MIE MOLS			65	65	65	65		68	68	68	68	68	
Tårs-Spødsbjerg	FRIGG SYDFYEN	45	45	45	45	45	45		45	45	45	45	45	
Tårs-Spødsbjerg	Langeland													45
Tårs-Spødsbjerg	Lolland													45
Tårs-Spødsbjerg	ODIN SYDFYEN	45	45	45	45	45	45		45	45	45	45	45	
Tårs-Spødsbjerg	SPODSBJERG	45	45	45	45	45	45		45	45	45			
Tårs-Spødsbjerg	THOR SYDFYEN													

Ferry data: Load factor (% MCR).

Ferry service	Ferry name	1985-1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
Esbjerg-Torshavn	Gamle Norrøna	90	90	90	90	90	90	90	90	90	90
Esbjerg-Torshavn	Nye Norrøna										
Halsskov-Knudshoved	ARVEPRINS KNUD DRONNING MARGRETHE	85	85	85	85	85	85	85	85	85	
Halsskov-Knudshoved	II	85	85	85	85	85	85	85	85	85	
Halsskov-Knudshoved	HEIMDAL	85	85	85	85	85	85	85	85	85	
Halsskov-Knudshoved	KNUDSHOVED	85	85	85	85	85	85	85	85	85	
Halsskov-Knudshoved	KONG FREDERIK IX	85	85	85	85	85	85	85	85	85	
Halsskov-Knudshoved	KRAKA	85	85	85	85	85	85	85	85	85	
Halsskov-Knudshoved	LODBROG	85	85	85	85	85	85	85	85	85	
Halsskov-Knudshoved	PRINSESSE ANNE-MARIE	85	85	85	85	85	85	85	85	85	
Halsskov-Knudshoved	PRINSESSE ELISABETH	85	85	85	85	85	85	85	85	85	
Halsskov-Knudshoved	ROMSØ	85	85	85	85	85	85	85	85	85	
Halsskov-Knudshoved	SPROGØ	85	85	85	85	85	85	85	85	85	
Hanstholm-Torshavn	Gamle Norrøna	90	90	90	90	90	90	90	90	90	90
Hanstholm-Torshavn	Nye Norrøna										
Hirtshals-Torshavn	Nye Norrøna										
Hundested-Grenaa	DJURSLAND	80	80	80	80	80					
Hundested-Grenaa	KATTEGAT						85	85			
Hundested-Grenaa	KONG FREDERIK IX					65					
Hundested-Grenaa	PRINSESSE ANNE-MARIE					85					
Kalundborg-Juelsminde	Mercandia I	75	75	75	75	75	75	75			
Kalundborg-Juelsminde	Mercandia II	70	70	70	70	70	70	70			
Kalundborg-Juelsminde	Mercandia III	70	70	70	70	70	70	70			
Kalundborg-Juelsminde	Mercandia IV	70	70	70	70	70	70	70			
Kalundborg-Samsø	HOLGER DANSKE			85	85	85	85	85	85	85	
Kalundborg-Samsø	KALUNDBORG	80	80	80							
Kalundborg-Samsø	KYHOLM									85	85
Kalundborg-Samsø	VESBORG									95	
Kalundborg-Århus	ASK		85	85	85	80	80	80	80	80	80
Kalundborg-Århus	CAT-LINK I						95	90	90	85	
Kalundborg-Århus	CAT-LINK II						95	90	90	85	
Kalundborg-Århus	CAT-LINK III							95	95	90	
Kalundborg-Århus	CAT-LINK IV									95	95
Kalundborg-Århus	CAT-LINK V									95	95
Kalundborg-Århus	KATTEGAT SYD										85

Kalundborg-Århus	KNUDSHOVED	85									
Kalundborg-Århus	KONG FREDERIK IX	85	85	85	85	85	85				
Kalundborg-Århus	KRAKA									85	
Kalundborg-Århus	MAREN MOLS										
Kalundborg-Århus	METTE MOLS										
Kalundborg-Århus	NIELS KLIM	85	85								
Kalundborg-Århus	PEDER PAARS	85	85								
Kalundborg-Århus	PRINSESSE ELISABETH		80								
Kalundborg-Århus	ROSTOCK LINK										80
Kalundborg-Århus	SØLØVEN/SØBJØRNEN		90	90	90	90	90	90			
Kalundborg-Århus	URD		85	85	85	85	85	85	85	80	80
Korsør-Nyborg, DSB	ASA-THOR	85	85	85	85	85	85	85	85		
Korsør-Nyborg, DSB	DRONNING INGRID	60	60	60	60	60	60	60	60		
Korsør-Nyborg, DSB	DRONNING MARGRETHE II	85	85	85	85	85	85	85	85		
Korsør-Nyborg, DSB	KONG FREDERIK IX	70	70	70	70	70	70	70	70		
Korsør-Nyborg, DSB	KRONPRINS FREDERIK	60	60	60	60	60	60	60	60		
Korsør-Nyborg, DSB	PRINS JOACHIM	60	60	60	60	60	60	60	60		
Korsør-Nyborg, DSB	SPROGØ/KNUDSHOVED	70	70	70	70	70	70	70	70		
Korsør-Nyborg, Vognmandsruten	Superflex Alfa	70	70	70	70	70	70	70	70	70	
Korsør-Nyborg, Vognmandsruten	Superflex Bravo	70	70	70	70	70	70	70	70	70	
Korsør-Nyborg, Vognmandsruten	Superflex Charlie	70	70	70	70	70	70	70	70	70	
København-Rønne	JENS KOFOED	30,77	30,77	30,77	30,77	30,77	30,77	30,77	30,77	30,77	30,77
København-Rønne	POVL ANKER	30,77	30,77	30,77	30,77	30,77	30,77	30,77	30,77	30,77	30,77
Køge-Rønne	DUEODDE										
Køge-Rønne	HAMMERODDE										
Køge-Rønne	JENS KOFOED										
Køge-Rønne	POVL ANKER										
Sjællands Odde-Ebeltoft	MAI MOLS							80	80	80	80
Sjællands Odde-Ebeltoft	MAREN MOLS	75	75	75	75	75	75	75			
Sjællands Odde-Ebeltoft	MAREN MOLS 2							80	80	80	85
Sjællands Odde-Ebeltoft	MAX MOLS										
Sjællands Odde-Ebeltoft	METTE MOLS	75	75	75	75	75	75	75			
Sjællands Odde-Ebeltoft	METTE MOLS 2							80	80	80	85
Sjællands Odde-Ebeltoft	MIE MOLS	85	85	85	85	85	85	85			
Sjællands Odde-Ebeltoft	MIE MOLS 2							80	80	80	80
Sjællands Odde-Århus	KatExpress										
Sjællands Odde-Århus	MADS MOLS										90

Sjællands Odde-Århus	MAI MOLS													
Sjællands Odde-Århus	MAX MOLS													90
Sjællands Odde-Århus	MIE MOLS													
Tårs-Spødsbjerg	FRIGG SYDFYEN	80	80	80	80	80	80	80	80	80	80	80	80	
Tårs-Spødsbjerg	Langeland													
Tårs-Spødsbjerg	Lolland													
Tårs-Spødsbjerg	ODIN SYDFYEN	80	80	80	80	80	80	80	80	80	80	80	80	
Tårs-Spødsbjerg	SPODSBJERG	75	80	80	80	80	80	80	80	80	80	80	80	
Tårs-Spødsbjerg	THOR SYDFYEN	80	80	80	80	80	80	80	80	80	80	80	80	

Ferry service	Ferry name	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
Esbjerg-Torshavn	Gamle Norrøna	90	90	90										
Esbjerg-Torshavn	Nye Norrøna				90	90	90	90	90	90	90	90	90	90
Halsskov-Knudshoved	ARVEPRINS KNUD DRONNING MARGRETHE II													
Halsskov-Knudshoved	HEIMDAL													
Halsskov-Knudshoved	KNUDSHOVED													
Halsskov-Knudshoved	KONG FREDERIK IX													
Halsskov-Knudshoved	KRAKA													
Halsskov-Knudshoved	LODBROG													
Halsskov-Knudshoved	PRINSESSE ANNE-MARIE													
Halsskov-Knudshoved	PRINSESSE ELISABETH													
Halsskov-Knudshoved	ROMSØ													
Halsskov-Knudshoved	SPROGØ													
Hanstholm-Torshavn	Gamle Norrøna	90	90	90										
Hanstholm-Torshavn	Nye Norrøna				90	90	90	90	90	90	90	90	90	90
Hirtshals-Torshavn	Nye Norrøna											90	90	90
Hundested-Grenaa	DJURSLAND													
Hundested-Grenaa	KATTEGAT													
Hundested-Grenaa	KONG FREDERIK IX													
Hundested-Grenaa	PRINSESSE ANNE-MARIE													
Kalundborg-Juelsminde	Mercandia I													
Kalundborg-Juelsminde	Mercandia II													
Kalundborg-Juelsminde	Mercandia III													
Kalundborg-Juelsminde	Mercandia IV													

Kalundborg-Samsø	HOLGER DANSKE														
Kalundborg-Samsø	KALUNDBORG														
Kalundborg-Samsø	KYHOLM	85	85	85	85	85	85	85	85	85	85	85	85	85	85
Kalundborg-Samsø	VESBORG														
Kalundborg-Århus	ASK														
Kalundborg-Århus	CAT-LINK I														
Kalundborg-Århus	CAT-LINK II														
Kalundborg-Århus	CAT-LINK III														
Kalundborg-Århus	CAT-LINK IV														
Kalundborg-Århus	CAT-LINK V														
Kalundborg-Århus	KATTEGAT SYD														
Kalundborg-Århus	KNUDSHOVED														
Kalundborg-Århus	KONG FREDERIK IX														
Kalundborg-Århus	KRAKA														
Kalundborg-Århus	MAREN MOLS	85	85	85	85	85	85	82	80	80	80	80	80	80	80
Kalundborg-Århus	METTE MOLS	85	85	85	85	85	85	82	80	80	80	80	80	80	80
Kalundborg-Århus	NIELS KLIM														
Kalundborg-Århus	PEDER PAARS														
Kalundborg-Århus	PRINSESSE ELISABETH														
Kalundborg-Århus	ROSTOCK LINK														
Kalundborg-Århus	SØLØVEN/SØBJØRNEN														
Kalundborg-Århus	URD														
Korsør-Nyborg, DSB	ASA-THOR														
Korsør-Nyborg, DSB	DRONNING INGRID														
Korsør-Nyborg, DSB	DRONNING MARGRETHE II														
Korsør-Nyborg, DSB	KONG FREDERIK IX														
Korsør-Nyborg, DSB	KRONPRINS FREDERIK														
Korsør-Nyborg, DSB	PRINS JOACHIM														
Korsør-Nyborg, DSB	SPROGØ/KNUDSHOVED														
Korsør-Nyborg, Vognmandsruten	Superflex Alfa														
Korsør-Nyborg, Vognmandsruten	Superflex Bravo														
Korsør-Nyborg, Vognmandsruten	Superflex Charlie														
København-Rønne	JENS KOFOED	30,77	30,77	30,77	30,77	30,77									
København-Rønne	POVL ANKER	30,77	30,77	30,77	30,77	30,77									
Køge-Rønne	DUEODDE						69,07	65	65	65	65	65			
Køge-Rønne	HAMMERODDE						69,07	65	65	65	65	65	65	65	65
Køge-Rønne	JENS KOFOED						31,34	31,34							

Køge-Rønne	POVL ANKER					31,34	31,34	45	49	49				
Sjællands Odde-Ebeltoft	MAI MOLS	80	80	80	80	80	80	79	78	75	74	77	78	83
Sjællands Odde-Ebeltoft	MAREN MOLS													
Sjællands Odde-Ebeltoft	MAREN MOLS 2													
Sjællands Odde-Ebeltoft	MAX MOLS													60
Sjællands Odde-Ebeltoft	METTE MOLS													
Sjællands Odde-Ebeltoft	METTE MOLS 2													
Sjællands Odde-Ebeltoft	MIE MOLS													
Sjællands Odde-Ebeltoft	MIE MOLS 2	80	80	80	80	80	80	79	78	75	74	77	78	83
Sjællands Odde-Århus	KatExpress													72
Sjællands Odde-Århus	MADS MOLS	85	85	85	85	85	85							
Sjællands Odde-Århus	MAI MOLS			75	75	75	75	69	69	69	68	71	70	73
Sjællands Odde-Århus	MAX MOLS	85	85	85	85	85	85	67	67	67	64	65	62	64
Sjællands Odde-Århus	MIE MOLS			75	75	75	75	69	69	69	68	71	70	73
Tårs-Spødsbjerg	FRIGG SYDFYEN	80	80	80	80	80	80	80	80	80	80	80	80	80
Tårs-Spødsbjerg	Langeland													80
Tårs-Spødsbjerg	Lolland													80
Tårs-Spødsbjerg	ODIN SYDFYEN	80	80	80	80	80	80	80	80	80	80	80	80	80
Tårs-Spødsbjerg	SPODSBJERG	80	80	80	80	80	80	80	80	80	80	80		
Tårs-Spødsbjerg	THOR SYDFYEN													

Ferry data: Round trip shares (%).

Ferry service	Ferry name	1985-1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
Esbjerg-Torshavn	Gamle Norrøna	100	100	100	100	100	100	100	100	100	100
Esbjerg-Torshavn	Nye Norrøna										
Halsskov-Knudshoved	ARVEPRINS KNUD	21	20	20	20	21	19	19	18	20	
Halsskov-Knudshoved	DRONNING MARGRETHE II	2	0	0	0	0	0	0	0	0	
Halsskov-Knudshoved	HEIMDAL	23	24	22	24	23	21	21	19	22	
Halsskov-Knudshoved	KNUDSHOVED	0	0	0	0	0	0	2	5	0	
Halsskov-Knudshoved	KONG FREDERIK IX	0	0	0	0	0	0	0	0	0	
Halsskov-Knudshoved	KRAKA	24	25	23	23	21	20	20	20	21	
Halsskov-Knudshoved	LODBROG	0	0	0	0	0	0	0	7	14	
Halsskov-Knudshoved	PRINSESSE ANNE-MARIE	0	0	0	0	0	6	2	0	0	
Halsskov-Knudshoved	PRINSESSE ELISABETH	0	0	0	3	0	0	0	0	0	
Halsskov-Knudshoved	ROMSØ	21	22	21	16	20	19	21	21	23	

Halsskov-Knudshoved	SPROGØ	9	9	15	14	15	15	14	11	1	
Hanstholm-Torshavn	Gamle Norrøna	100	100	100	100	100	100	100	100	100	100
Hanstholm-Torshavn	Nye Norrøna										
Hirtshals-Torshavn	Nye Norrøna										
Hundested-Grenaa	DJURSLAND	100	100	100	100	50					
Hundested-Grenaa	KATTEGAT						100	100			
Hundested-Grenaa	KONG FREDERIK IX					5					
Hundested-Grenaa	PRINSESSE ANNE-MARIE					45					
Kalundborg-Juelsminde	Mercandia I	25	25	25	25	25	25	25			
Kalundborg-Juelsminde	Mercandia II	25	25	25	25	25	25	25			
Kalundborg-Juelsminde	Mercandia III	25	25	25	25	25	25	25			
Kalundborg-Juelsminde	Mercandia IV	25	25	25	25	25	25	25			
Kalundborg-Samsø	HOLGER DANSKE			95	100	100	100	100	100	92	
Kalundborg-Samsø	KALUNDBORG	100	100	5							
Kalundborg-Samsø	KYHOLM									6	100
Kalundborg-Samsø	VESBORG									2	
Kalundborg-Århus	ASK		16	32	26	33	27	18	11	12	2
Kalundborg-Århus	CAT-LINK I						17	25	28	11	
Kalundborg-Århus	CAT-LINK II						1	23	28	8	
Kalundborg-Århus	CAT-LINK III							8	24	19	
Kalundborg-Århus	CAT-LINK IV									23	26
Kalundborg-Århus	CAT-LINK V									15	26
Kalundborg-Århus	KATTEGAT SYD										2
Kalundborg-Århus	KNUDSHOVED		4								
Kalundborg-Århus	KONG FREDERIK IX		4	0	7	0	0	2			
Kalundborg-Århus	KRAKA									2	
Kalundborg-Århus	MAREN MOLS										
Kalundborg-Århus	METTE MOLS										
Kalundborg-Århus	NIELS KLIM	50	20								
Kalundborg-Århus	PEDER PAARS	50	16								
Kalundborg-Århus	PRINSESSE ELISABETH		4								
Kalundborg-Århus	ROSTOCK LINK										22
Kalundborg-Århus	SØLØVEN/SØBJØRNEN		21	36	34	34	28	5			
Kalundborg-Århus	URD		16	32	33	33	27	18	11	9	22
Korsør-Nyborg, DSB	ASA-THOR	13	13	13	11	9	9	9	6		
Korsør-Nyborg, DSB	DRONNING INGRID	26	28	26	28	28	29	28	31		
Korsør-Nyborg, DSB	DRONNING MARGRETHE II	3	0	3	1	3	1	2	0		

Korsør-Nyborg, DSB	KONG FREDERIK IX	0	0	0	0	3	4	1	0		
Korsør-Nyborg, DSB	KRONPRINS FREDERIK	27	28	27	29	28	29	29	32		
Korsør-Nyborg, DSB	PRINS JOACHIM	25	27	25	27	27	27	27	28		
Korsør-Nyborg, DSB	SPROGØ/KNUDSHOVED	6	4	5	4	1	1	4	3		
Korsør-Nyborg, Vognmandsruten	Superflex Alfa	33	33	33	33	33	33	33	33	33	
Korsør-Nyborg, Vognmandsruten	Superflex Bravo	33	33	33	33	33	33	33	33	33	
Korsør-Nyborg, Vognmandsruten	Superflex Charlie	34	34	34	34	34	34	34	34	34	
København-Rønne	JENS KOFOED	50	50	50	50	50	50	50	50	50	50
København-Rønne	POVL ANKER	50	50	50	50	50	50	50	50	50	50
Køge-Rønne	DUEODDE										
Køge-Rønne	HAMMERODDE										
Køge-Rønne	JENS KOFOED										
Køge-Rønne	POVL ANKER										
Sjællands Odde-Ebeltoft	MAI MOLS							21	35	35	35
Sjællands Odde-Ebeltoft	MAREN MOLS	40	40	40	40	40	40	15			
Sjællands Odde-Ebeltoft	MAREN MOLS 2							18	15	15	15
Sjællands Odde-Ebeltoft	MAX MOLS										
Sjællands Odde-Ebeltoft	METTE MOLS	40	40	40	40	40	40	17			
Sjællands Odde-Ebeltoft	METTE MOLS 2							15	15	15	15
Sjællands Odde-Ebeltoft	MIE MOLS	20	20	20	20	20	20	5			
Sjællands Odde-Ebeltoft	MIE MOLS 2							9	35	35	35
Sjællands Odde-Århus	KatExpress										
Sjællands Odde-Århus	MADS MOLS										50
Sjællands Odde-Århus	MAI MOLS										
Sjællands Odde-Århus	MAX MOLS										50
Sjællands Odde-Århus	MIE MOLS										
Tårs-Spødsbjerg	FRIGG SYDFYEN	41	40	39	38	36	36	36	32	33	45
Tårs-Spødsbjerg	Langeland										
Tårs-Spødsbjerg	Lolland										
Tårs-Spødsbjerg	ODIN SYDFYEN	41	40	39	38	36	36	36	32	33	45
Tårs-Spødsbjerg	SPODSBJERG	4	2	8	8	9	8	8	19	20	10
Tårs-Spødsbjerg	THOR SYDFYEN	14	18	14	16	19	20	20	17	14	

Ferry service	Ferry name	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
Esbjerg-Torshavn	Gamle Norrøna	100	100	100										
Esbjerg-Torshavn	Nye Norrøna				100	100	100	100	100	100	100	100	100	100
Halsskov-Knudshoved	ARVEPRINS KNUD													
Halsskov-Knudshoved	DRONNING MARGRETHE II													
Halsskov-Knudshoved	HEIMDAL													
Halsskov-Knudshoved	KNUDSHOVED													
Halsskov-Knudshoved	KONG FREDERIK IX													
Halsskov-Knudshoved	KRAKA													
Halsskov-Knudshoved	LODBROG													
Halsskov-Knudshoved	PRINSESSE ANNE-MARIE													
Halsskov-Knudshoved	PRINSESSE ELISABETH													
Halsskov-Knudshoved	ROMSØ													
Halsskov-Knudshoved	SPROGØ													
Hanstholm-Torshavn	Gamle Norrøna	100	100	100										
Hanstholm-Torshavn	Nye Norrøna				100	100	100	100	100	100	100	100	100	100
Hirtshals-Torshavn	Nye Norrøna											100	100	100
Hundested-Grenaa	DJURSLAND													
Hundested-Grenaa	KATTEGAT													
Hundested-Grenaa	KONG FREDERIK IX													
Hundested-Grenaa	PRINSESSE ANNE-MARIE													
Kalundborg-Juelsminde	Mercandia I													
Kalundborg-Juelsminde	Mercandia II													
Kalundborg-Juelsminde	Mercandia III													
Kalundborg-Juelsminde	Mercandia IV													
Kalundborg-Samsø	HOLGER DANSKE													
Kalundborg-Samsø	KALUNDBORG													
Kalundborg-Samsø	KYHOLM	100	100	100	100	100	100	100	100	100	100	100	100	100
Kalundborg-Samsø	VESBORG													
Kalundborg-Århus	ASK													
Kalundborg-Århus	CAT-LINK I													
Kalundborg-Århus	CAT-LINK II													
Kalundborg-Århus	CAT-LINK III													
Kalundborg-Århus	CAT-LINK IV													
Kalundborg-Århus	CAT-LINK V													
Kalundborg-Århus	KATTEGAT SYD													
Kalundborg-Århus	KNUDSHOVED													

Kalundborg-Århus	KONG FREDERIK IX														
Kalundborg-Århus	KRAKA														
Kalundborg-Århus	MAREN MOLS	50	50	50	50	50	50	50	50	50	50	50	50	50	50
Kalundborg-Århus	METTE MOLS	50	50	50	50	50	50	50	50	50	50	50	50	50	50
Kalundborg-Århus	NIELS KLIM														
Kalundborg-Århus	PEDER PAARS														
Kalundborg-Århus	PRINSESSE ELISABETH														
Kalundborg-Århus	ROSTOCK LINK														
Kalundborg-Århus	SØLØVEN/SØBJØRNEN														
Kalundborg-Århus	URD														
Korsør-Nyborg, DSB	ASA-THOR														
Korsør-Nyborg, DSB	DRONNING INGRID														
Korsør-Nyborg, DSB	DRONNING MARGRETHE II														
Korsør-Nyborg, DSB	KONG FREDERIK IX														
Korsør-Nyborg, DSB	KRONPRINS FREDERIK														
Korsør-Nyborg, DSB	PRINS JOACHIM														
Korsør-Nyborg, DSB	SPROGØ/KNUDSHOVED														
Korsør-Nyborg, Vognmandsruten	Superflex Alfa														
Korsør-Nyborg, Vognmandsruten	Superflex Bravo														
Korsør-Nyborg, Vognmandsruten	Superflex Charlie														
København-Rønne	JENS KOFOED	50	50	50	50	50									
København-Rønne	POVL ANKER	50	50	50	50	50									
Køge-Rønne	DUEODDE							25	49	47	46	51	46		
Køge-Rønne	HAMMERODDE							35	49	53	47	49	54	100	100
Køge-Rønne	JENS KOFOED						50	20							
Køge-Rønne	POVL ANKER						50	20	3	1	7				
Sjællands Odde-Ebeltoft	MAI MOLS	50	50	50	50	50	50	50	50	50	50	50	50	50	47
Sjællands Odde-Ebeltoft	MAREN MOLS														
Sjællands Odde-Ebeltoft	MAREN MOLS 2														
Sjællands Odde-Ebeltoft	MAX MOLS														6
Sjællands Odde-Ebeltoft	METTE MOLS														
Sjællands Odde-Ebeltoft	METTE MOLS 2														
Sjællands Odde-Ebeltoft	MIE MOLS														
Sjællands Odde-Ebeltoft	MIE MOLS 2	50	50	50	50	50	50	50	50	50	50	50	50	50	47
Sjællands Odde-Århus	KatExpress														18
Sjællands Odde-Århus	MADS MOLS	95	90	95	60	60	35								
Sjællands Odde-Århus	MAI MOLS			1	10	15	15	20	19	19	20	18	20	6	

Sjællands Odde-Århus	MAX MOLS	5	10	3	20	10	35	60	62	62	60	64	60	70
Sjællands Odde-Århus	MIE MOLS			1	10	15	15	20	19	19	20	18	20	6
Tårs-Spodsbjerg	FRIGG SYDFYEN	45	45	45	45	45	45	45	45	45	45	45	50	21
Tårs-Spodsbjerg	Langeland													40
Tårs-Spodsbjerg	Lolland													29
Tårs-Spodsbjerg	ODIN SYDFYEN	45	45	45	45	45	45	45	45	45	45	45	50	10
Tårs-Spodsbjerg	SPODSBJERG	10	10	10	10	10	10	10	10	10	10	10		
Tårs-Spodsbjerg	THOR SYDFYEN													

Annex 3B-13 Fuel consumption and emission factors, engine specific (NO_x, CO, VOC (NMVOC and CH₄)), and fuel type specific (S-%, SO₂, PM) for ship engines

Specific fuel consumption and NO_x emission factors (g pr kWh) per engine year for diesel ship engines.

Year	High speed 4-stroke sfc (g pr kWh)	Medium speed 4-stroke sfc (g pr kWh)	Slow speed 2-stroke sfc (g pr kWh)	High speed 4-stroke NO _x (g pr kWh)	Medium speed 4-stroke NO _x (g pr kWh)	Slow speed 2-stroke NO _x (g pr kWh)
1949	265.5	255.5	235.5	7.3	8.0	14.5
1950	265.0	255.0	235.0	7.3	8.0	14.5
1951	264.5	254.5	234.5	7.3	8.0	14.5
1952	264.0	254.0	234.0	7.3	8.0	14.5
1953	263.5	253.5	233.5	7.3	8.0	14.5
1954	263.0	253.0	233.0	7.3	8.0	14.5
1955	262.4	252.4	232.4	7.3	8.0	14.5
1956	261.9	251.9	231.9	7.4	8.1	14.6
1957	261.3	251.3	231.3	7.5	8.2	14.7
1958	260.7	250.7	230.7	7.6	8.3	14.8
1959	260.1	250.1	230.1	7.7	8.4	14.9
1960	259.5	249.5	229.5	7.8	8.5	15.0
1961	258.9	248.9	228.9	7.9	8.6	15.1
1962	258.2	248.2	228.2	8.0	8.7	15.1
1963	257.6	247.6	227.6	8.1	8.8	15.2
1964	256.9	246.9	226.9	8.2	8.9	15.3
1965	256.1	246.1	226.1	8.3	9.0	15.4
1966	255.4	245.4	225.4	8.3	9.1	15.5
1967	254.6	244.6	224.6	8.4	9.2	15.6
1968	253.8	243.8	223.8	8.5	9.3	15.7
1969	253.0	243.0	223.0	8.6	9.4	15.8
1970	252.1	242.1	222.1	8.7	9.5	15.9
1971	251.2	241.2	221.2	8.8	9.6	16.0
1972	250.3	240.3	220.3	8.9	9.7	16.1
1973	249.3	239.3	219.3	9.0	9.8	16.2
1974	248.3	238.3	218.3	9.1	9.9	16.3
1975	247.3	237.3	217.3	9.2	10.0	16.4
1976	246.2	236.2	216.2	9.3	10.1	16.4
1977	245.0	235.0	215.0	9.3	10.2	16.5
1978	243.8	233.8	213.8	9.4	10.3	16.6
1979	242.6	232.6	212.6	9.5	10.4	16.7
1980	241.3	231.3	211.3	9.6	10.5	16.8
1981	239.9	229.9	209.9	9.7	10.6	16.9
1982	238.5	228.5	208.5	9.8	10.7	17.0
1983	237.0	227.0	207.0	9.9	10.8	17.4
1984	235.5	225.5	205.5	10.0	10.9	17.8
1985	233.9	223.9	203.9	10.1	11.0	18.2
1986	232.2	222.2	202.2	10.2	11.1	18.6
1987	230.5	220.5	200.5	10.3	11.3	19.0
1988	228.6	218.6	198.6	10.5	11.4	19.3
1989	226.7	216.7	196.7	10.6	11.6	19.5
1990	224.8	214.8	194.8	10.7	11.7	19.8
1991	222.7	212.7	192.7	10.9	11.9	20.0
1992	220.5	210.5	190.5	11.0	12.0	19.8
1993	218.3	208.3	188.3	11.1	12.1	19.6
1994	216.0	206.0	186.0	11.3	12.3	19.4
1995	213.6	203.6	183.6	11.4	12.4	19.3
1996	211.0	201.0	181.0	11.5	12.6	19.1
1997	208.4	198.4	178.4	11.7	12.7	18.9
1998	205.7	195.7	175.7	11.8	12.9	18.7

1999	202.9	192.9	172.9	11.9	13.0	18.5
2000-2010	199.9	189.9	169.9	11.0	12.0	16.0
2011-2012	199.9	189.9	169.9	8.69	9.72	13.6

CO and VOC emission factors (g/kg fuel) for ship engines.

	High speed 4-stroke CO	Medium speed 4-stroke CO	Slow speed 2-stroke CO	High speed 4-stroke VOC	Medium speed 4-stroke VOC	Slow speed 2-stroke VOC
1949	6.03	6.26	6.79	1.88	1.96	2.12
1950	6.04	6.27	6.81	1.89	1.96	2.13
1951	6.05	6.29	6.82	1.89	1.96	2.13
1952	6.06	6.30	6.84	1.89	1.97	2.14
1953	6.07	6.31	6.85	1.90	1.97	2.14
1954	6.08	6.33	6.87	1.90	1.98	2.15
1955	6.10	6.34	6.88	1.91	1.98	2.15
1956	6.11	6.35	6.90	1.91	1.99	2.16
1957	6.12	6.37	6.92	1.91	1.99	2.16
1958	6.14	6.38	6.93	1.92	1.99	2.17
1959	6.15	6.40	6.95	1.92	2.00	2.17
1960	6.17	6.41	6.97	1.93	2.00	2.18
1961	6.18	6.43	6.99	1.93	2.01	2.18
1962	6.20	6.45	7.01	1.94	2.01	2.19
1963	6.21	6.46	7.03	1.94	2.02	2.20
1964	6.23	6.48	7.05	1.95	2.03	2.20
1965	6.25	6.50	7.08	1.95	2.03	2.21
1966	6.26	6.52	7.10	1.96	2.04	2.22
1967	6.28	6.54	7.12	1.96	2.04	2.23
1968	6.30	6.56	7.15	1.97	2.05	2.23
1969	6.32	6.58	7.17	1.98	2.06	2.24
1970	6.35	6.61	7.20	1.98	2.06	2.25
1971	6.37	6.63	7.23	1.99	2.07	2.26
1972	6.39	6.66	7.26	2.00	2.08	2.27
1973	6.42	6.69	7.29	2.01	2.09	2.28
1974	6.44	6.71	7.33	2.01	2.10	2.29
1975	6.47	6.74	7.36	2.02	2.11	2.30
1976	6.50	6.77	7.40	2.03	2.12	2.31
1977	6.53	6.81	7.44	2.04	2.13	2.33
1978	6.56	6.84	7.48	2.05	2.14	2.34
1979	6.60	6.88	7.53	2.06	2.15	2.35
1980	6.63	6.92	7.57	2.07	2.16	2.37
1981	6.67	6.96	7.62	2.08	2.17	2.38
1982	6.71	7.00	7.67	2.10	2.19	2.40
1983	6.75	7.05	7.73	2.11	2.20	2.42
1984	6.79	7.10	7.79	2.12	2.22	2.43
1985	6.84	7.15	7.85	2.14	2.23	2.45
1986	6.89	7.20	7.91	2.15	2.25	2.47
1987	6.94	7.26	7.98	2.17	2.27	2.49
1988	7.00	7.32	8.05	2.19	2.29	2.52
1989	7.06	7.38	8.13	2.21	2.31	2.54
1990	7.12	7.45	8.22	2.22	2.33	2.57
1991	7.18	7.52	8.30	2.25	2.35	2.59
1992	7.25	7.60	8.40	2.27	2.37	2.62
1993	7.33	7.68	8.50	2.29	2.40	2.66
1994	7.41	7.77	8.60	2.31	2.43	2.69
1995	7.49	7.86	8.72	2.34	2.46	2.72
1996	7.58	7.96	8.84	2.37	2.49	2.76
1997	7.68	8.06	8.97	2.40	2.52	2.80

1998	7.78	8.18	9.11	2.43	2.56	2.85
1999	7.89	8.30	9.26	2.46	2.59	2.89
2000	8.00	8.43	9.42	2.50	2.63	2.94

NM VOC and CH₄ emission factors (g/kg fuel) for ship engines.

	High speed 4-stroke NM VOC	Medium speed 4-stroke NM VOC	Slow speed 2-stroke NM VOC	High speed 4-stroke CH ₄	Medium speed 4-stroke CH ₄	Slow speed 2-stroke CH ₄
1949	1.83	1.90	2.06	0.06	0.06	0.06
1950	1.83	1.90	2.06	0.06	0.06	0.06
1951	1.83	1.91	2.07	0.06	0.06	0.06
1952	1.84	1.91	2.07	0.06	0.06	0.06
1953	1.84	1.91	2.08	0.06	0.06	0.06
1954	1.84	1.92	2.08	0.06	0.06	0.06
1955	1.85	1.92	2.09	0.06	0.06	0.06
1956	1.85	1.93	2.09	0.06	0.06	0.06
1957	1.86	1.93	2.10	0.06	0.06	0.06
1958	1.86	1.93	2.10	0.06	0.06	0.07
1959	1.86	1.94	2.11	0.06	0.06	0.07
1960	1.87	1.94	2.11	0.06	0.06	0.07
1961	1.87	1.95	2.12	0.06	0.06	0.07
1962	1.88	1.95	2.13	0.06	0.06	0.07
1963	1.88	1.96	2.13	0.06	0.06	0.07
1964	1.89	1.96	2.14	0.06	0.06	0.07
1965	1.89	1.97	2.14	0.06	0.06	0.07
1966	1.90	1.98	2.15	0.06	0.06	0.07
1967	1.90	1.98	2.16	0.06	0.06	0.07
1968	1.91	1.99	2.17	0.06	0.06	0.07
1969	1.92	2.00	2.17	0.06	0.06	0.07
1970	1.92	2.00	2.18	0.06	0.06	0.07
1971	1.93	2.01	2.19	0.06	0.06	0.07
1972	1.94	2.02	2.20	0.06	0.06	0.07
1973	1.95	2.03	2.21	0.06	0.06	0.07
1974	1.95	2.04	2.22	0.06	0.06	0.07
1975	1.96	2.04	2.23	0.06	0.06	0.07
1976	1.97	2.05	2.24	0.06	0.06	0.07
1977	1.98	2.06	2.26	0.06	0.06	0.07
1978	1.99	2.07	2.27	0.06	0.06	0.07
1979	2.00	2.09	2.28	0.06	0.06	0.07
1980	2.01	2.10	2.30	0.06	0.06	0.07
1981	2.02	2.11	2.31	0.06	0.07	0.07
1982	2.03	2.12	2.33	0.06	0.07	0.07
1983	2.05	2.14	2.34	0.06	0.07	0.07
1984	2.06	2.15	2.36	0.06	0.07	0.07
1985	2.07	2.17	2.38	0.06	0.07	0.07
1986	2.09	2.18	2.40	0.06	0.07	0.07
1987	2.10	2.20	2.42	0.07	0.07	0.07
1988	2.12	2.22	2.44	0.07	0.07	0.08
1989	2.14	2.24	2.47	0.07	0.07	0.08
1990	2.16	2.26	2.49	0.07	0.07	0.08
1991	2.18	2.28	2.52	0.07	0.07	0.08
1992	2.20	2.30	2.55	0.07	0.07	0.08
1993	2.22	2.33	2.58	0.07	0.07	0.08
1994	2.25	2.35	2.61	0.07	0.07	0.08
1995	2.27	2.38	2.64	0.07	0.07	0.08
1996	2.30	2.41	2.68	0.07	0.07	0.08
1997	2.33	2.44	2.72	0.07	0.08	0.08

1998	2.36	2.48	2.76	0.07	0.08	0.09
1999	2.39	2.51	2.81	0.07	0.08	0.09
2000	2.43	2.55	2.85	0.08	0.08	0.09

S-%, SO₂ and PM emission factors (g/kg fuel and g/GJ) per fuel type for diesel ship engines.

Fuel type	SNAPCode	Year	S %	SO ₂ (g/kg)	TSP (g/kg)	PM ₁₀ (g/kg)	PM _{2.5} (g/kg)	SO ₂ (g/GJ)	TSP (g/GJ)	PM ₁₀ (g/GJ)	PM _{2.5} (g/GJ)
Fuel	National sea	1990	2,6	52,8	6,1	6,0	6,0	1291,0	149,2	147,8	147,0
Fuel	National sea	1991	2,4	47,0	4,9	4,9	4,8	1149,1	120,2	119,0	118,4
Fuel	National sea	1992	1,8	36,0	3,3	3,2	3,2	880,2	79,8	79,0	78,6
Fuel	National sea	1993	2,4	47,8	5,1	5,0	5,0	1168,7	123,9	122,6	122,0
Fuel	National sea	1994	2,6	52,4	6,0	6,0	5,9	1281,2	147,0	145,6	144,8
Fuel	National sea	1995	3,0	59,0	7,7	7,6	7,6	1442,5	188,0	186,1	185,2
Fuel	National sea	1996	2,6	51,4	5,8	5,7	5,7	1256,7	141,7	140,2	139,5
Fuel	National sea	1997	2,7	54,8	6,6	6,5	6,5	1339,9	160,8	159,2	158,4
Fuel	National sea	1998	2,0	39,4	3,7	3,7	3,6	963,3	90,6	89,7	89,2
Fuel	National sea	1999	2,0	39,4	3,7	3,7	3,6	963,3	90,6	89,7	89,2
Fuel	National sea	2000	1,8	36,2	3,3	3,3	3,2	885,1	80,4	79,6	79,2
Fuel	National sea	2001	1,7	34,0	3,0	3,0	3,0	831,3	74,1	73,4	73,0
Fuel	National sea	2002	1,5	30,2	2,6	2,6	2,6	738,4	64,3	63,7	63,3
Fuel	National sea	2003	1,6	32,4	2,9	2,8	2,8	792,2	69,8	69,1	68,8
Fuel	National sea	2004	2,0	39,6	3,7	3,7	3,7	968,2	91,3	90,4	89,9
Fuel	National sea	2005	2,0	40,0	3,8	3,8	3,7	978,0	92,6	91,7	91,3
Fuel	National sea	2006	1,9	38,8	3,6	3,6	3,6	948,7	88,6	87,7	87,3
Fuel	National sea	2007	1,2	24,0	2,1	2,1	2,1	586,8	51,0	50,5	50,3
Fuel	National sea	2008	1,2	24,0	2,1	2,1	2,1	586,8	51,0	50,5	50,3
Fuel	National sea	2009	1,2	24,0	2,1	2,1	2,1	586,8	51,0	50,5	50,3
Fuel	National sea	2010	1,0	20,0	1,8	1,8	1,8	489,0	44,0	43,5	43,3
Fuel	National sea	2011	1,0	20,0	1,8	1,8	1,8	489,0	44,0	43,5	43,3
Fuel	National sea	2012	1,0	20,0	1,8	1,8	1,8	489,0	44,0	43,5	43,3
Fuel	International sea	1990	3,0	59,2	7,7	7,7	7,6	1447,4	189,4	187,5	186,6
Fuel	International sea	1991	2,9	57,8	7,4	7,3	7,2	1413,2	179,8	178,0	177,1
Fuel	International sea	1992	2,9	57,6	7,3	7,2	7,2	1408,3	178,5	176,7	175,8
Fuel	International sea	1993	3,2	64,0	9,3	9,2	9,1	1564,8	226,5	224,2	223,1
Fuel	International sea	1994	3,0	60,6	8,2	8,1	8,0	1481,7	199,6	197,6	196,6
Fuel	International sea	1995	3,3	66,0	10,0	9,9	9,8	1613,7	244,0	241,6	240,4
Fuel	International sea	1996	3,4	68,4	10,9	10,8	10,8	1672,4	266,9	264,2	262,9
Fuel	International sea	1997	3,5	69,0	11,2	11,0	11,0	1687,0	272,9	270,2	268,8
Fuel	International sea	1998	3,4	68,4	10,9	10,8	10,8	1672,4	266,9	264,2	262,9
Fuel	International sea	1999	3,5	69,0	11,2	11,0	11,0	1687,0	272,9	270,2	268,8
Fuel	International sea	2000	3,4	67,2	10,4	10,3	10,3	1643,0	255,2	252,6	251,4
Fuel	International sea	2001	3,4	68,4	10,9	10,8	10,8	1672,4	266,9	264,2	262,9
Fuel	International sea	2002	3,4	68,8	11,1	11,0	10,9	1682,2	270,9	268,2	266,8

Fuel	International sea	2003	3,1	62,2	8,7	8,6	8,5	1520,8	211,8	209,7	208,6
Fuel	International sea	2004	3,2	64,0	9,3	9,2	9,1	1564,8	226,5	224,2	223,1
Fuel	International sea	2005	3,5	70,0	11,6	11,5	11,4	1711,5	283,2	280,4	279,0
Fuel	International sea	2006	3,4	67,0	10,4	10,3	10,2	1638,1	253,3	250,8	249,5
Fuel	International sea	2007	1,5	30,0	2,6	2,6	2,6	733,5	63,8	63,2	62,9
Fuel	International sea	2008	1,5	30,0	2,6	2,6	2,6	733,5	63,8	63,2	62,9
Fuel	International sea	2009	1,5	30,0	2,6	2,6	2,6	733,5	63,8	63,2	62,9
Fuel	International sea	2010	1,0	20,0	1,8	1,8	1,8	489,0	44,0	43,5	43,3
Fuel	International sea	2011	1,0	20,0	1,8	1,8	1,8	489,0	44,0	43,5	43,3
Fuel	International sea	2012	1,0	20,0	1,8	1,8	1,8	489,0	44,0	43,5	43,3
Diesel		1990	0,2	4,0	1,0	1,0	1,0	93,7	23,2	23,0	22,9
Diesel		1991	0,2	4,0	1,0	1,0	1,0	93,7	23,2	23,0	22,9
Diesel		1992	0,2	4,0	1,0	1,0	1,0	93,7	23,2	23,0	22,9
Diesel		1993	0,2	4,0	1,0	1,0	1,0	93,7	23,2	23,0	22,9
Diesel		1994	0,2	4,0	1,0	1,0	1,0	93,7	23,2	23,0	22,9
Diesel		1995	0,2	4,0	1,0	1,0	1,0	93,7	23,2	23,0	22,9
Diesel		1996	0,2	4,0	1,0	1,0	1,0	93,7	23,2	23,0	22,9
Diesel		1997	0,2	4,0	1,0	1,0	1,0	93,7	23,2	23,0	22,9
Diesel		1998	0,2	4,0	1,0	1,0	1,0	93,7	23,2	23,0	22,9
Diesel		1999	0,2	4,0	1,0	1,0	1,0	93,7	23,2	23,0	22,9
Diesel		2000	0,2	4,0	1,0	1,0	1,0	93,7	23,2	23,0	22,9
Diesel		2001	0,2	4,0	1,0	1,0	1,0	93,7	23,2	23,0	22,9
Diesel		2002	0,2	4,0	1,0	1,0	1,0	93,7	23,2	23,0	22,9
Diesel		2003	0,2	4,0	1,0	1,0	1,0	93,7	23,2	23,0	22,9
Diesel		2004	0,2	4,0	1,0	1,0	1,0	93,7	23,2	23,0	22,9
Diesel		2005	0,2	4,0	1,0	1,0	1,0	93,7	23,2	23,0	22,9
Diesel		2006	0,2	4,0	1,0	1,0	1,0	93,7	23,2	23,0	22,9
Diesel		2007	0,2	4,0	1,0	1,0	1,0	93,7	23,2	23,0	22,9
Diesel		2008	0,1	2,0	0,9	0,9	0,9	46,8	21,5	21,3	21,2
Diesel		2009	0,1	2,0	0,9	0,9	0,9	46,8	21,5	21,3	21,2
Diesel		2010	0,1	2,0	0,9	0,9	0,9	46,8	21,5	21,3	21,2
Diesel		2011	0,1	2,0	0,9	0,9	0,9	46,8	21,5	21,3	21,2
Diesel		2012	0,1	2,0	0,9	0,9	0,9	46,8	21,5	21,3	21,2

Annex 3B-14 Fuel sales figures from DEA, and further processed fuel consumption data suited for the Danish inventory

Sector	1 985	1 986	1 987	1 988	1 989	1 990	1 991	1 992	1 993	1 994	1 995	1 996	1 997	1 998	1 999
Agriculture, Forestry and Horticulture															
- LPG	96	89	401	358	510	488	459	397	260	227	227	234	204	243	179
- motor gasoline	435	187	321	323	315	284	261	252	231	184	179	210	77	68	62
- gas/diesel oil	10904	10904	10904	10904	11836	11937	12387	12915	12628	12457	12723	13575	14231	13651	14256
Fishery	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
- LPG	0	0	34	29	50	42	34	30	12	18	16	36	5	1	16
- motor gasoline	0	1	2	2	9	9	10	8	7	7	8	7	6	6	60
- petroleum	7	2	9	5	12	26	9	5	4	3	4	3	3	2	0
- gas/diesel oil	9152	10248	8390	9499	10038	10422	10809	10868	8843	8796	8277	8750	8748	9186	9282
- fuel oil	27	5	82	68	251	285	113	231	146	8	19	219	260	27	0
Manyfactoring Industry	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
- LPG	2860	2839	2688	2553	2080	2032	2076	1827	1858	2029	2234	2404	2106	2017	1917
- motor gasoline	262	273	453	326	136	177	161	158	145	138	110	86	82	137	80
- gas/diesel oil	15576	15441	14743	13346	12670	12259	12934	11901	11323	10154	10401	10184	8921	8720	8852
- fuel oil	29465	29451	21518	19056	16741	15989	17133	16694	14600	15438	14000	12632	11009	10943	8704
Construction Industry	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
- LPG	305	343	500	451	575	500	573	708	579	522	501	509	471	575	422
- motorbenzin	19	85	52	48	36	34	26	24	20	23	25	34	27	23	27
- gas/diesel oil	5313	4962	4378	4220	3945	3548	3797	3839	3871	4145	5317	5572	6079	5947	6556
Single family houses	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
- motor gasoline	1006	1046	1073	1114	1128	1131	1146	1158	1168	1194	1233	1258	1299	1317	1357
- gas/diesel oil	74257	69392	68349	59832	46935	41152	45219	38406	45029	39770	40004	41836	36491	34902	32936
Multi-family houses	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
- gas/diesel oil	10584	9968	10112	7266	7350	5311	5420	4507	4938	3909	3284	3460	3105	2948	2739
Road transport, DEA statistics															
- gasoline	66 037	68 670	70 502	73 151	74 152	74 326	75 290	76 084	76 697	78 425	80 998	82 656	85 341	86 520	89 129
- gas/diesel oil	45 609	49 738	49 626	49 686	51 854	54 746	58 427	57 511	56 796	58 755	58 561	59 851	60 528	61 072	63 619
- bioethanol	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
- biodiesel	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Non-road, DEA statistics															
- LPG	2 955	2 929	3 089	2 911	2 590	2 520	2 535	2 224	2 118	2 257	2 461	2 638	2 310	2 260	2 097

- gasoline	1 722	1 590	1 898	1 810	1 616	1 626	1 595	1 592	1 563	1 540	1 547	1 589	1 485	1 545	1 526
- gas/diesel oil	31 793	31 307	30 025	28 469	28 451	27 744	29 118	28 655	27 822	26 755	28 441	29 331	29 231	28 319	29 665
Non-road, NERI model															
- LPG	1232	1233	1225	1209	1196	1185	1172	1151	1124	1105	1099	1088	1075	1086	1077
- gasoline	2998	2950	2903	2856	2813	2770	2702	2641	2587	2550	2521	2499	2479	2463	2456
- gas/diesel oil	26357	26895	26577	27075	26940	26800	26734	26046	26073	25235	25798	25139	25536	24844	24885
Recreational craft, NERI model															
- gasoline	270	270	279	289	299	309	319	329	339	348	358	368	377	385	391
- gas/diesel oil	219	219	247	277	309	343	378	415	454	495	537	581	628	676	726
Non-road, added 0202															
- gas/diesel oil	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Non-road, added 0203 and 0301															
- gas/diesel oil	5436	4412	3448	1395	1510	944	2384	2609	1748	1521	2642	4192	3695	3475	4780
- LPG	1724	1696	1864	1701	1393	1335	1363	1073	994	1152	1362	1549	1235	1175	1020
Non-road, added 0203															
- gas/diesel oil	1864	1537	1252	534	628	406	1014	1176	794	708	1182	1940	1799	1675	2297
- LPG	56	52	242	209	274	259	247	192	122	116	125	137	109	126	87
Non-road, added 0301															
- gas/diesel oil	3572	2875	2196	860	882	538	1370	1433	955	813	1460	2252	1896	1800	2483
- LPG	1668	1644	1622	1492	1119	1076	1116	881	872	1036	1237	1412	1126	1048	933
Non-road, added road transport															
- gasoline	-1276	-1360	-1005	-1046	-1197	-1145	-1107	-1049	-1023	-1010	-975	-909	-994	-918	-931
Fisheries, added national sea transport															
- fuel oil	27	5	82	68	251	285	113	231	146	8	19	219	260	27	0
Fisheries, consumed by recreational craft															
- gasoline	0	1	2	2	9	9	10	8	7	7	8	7	6	6	60

National sea transport, input NERI model

- LPG	3	1	3	-	2	2	2	3	16	1	2	1	2	3	1
- kerosene	5	-	5	3	1	0	2	1	1	1	1	1	0	1	0
- gas/diesel oil	3 074	3 045	3 032	3 230	2 669	2 782	3 313	3 501	4 971	5 035	6 049	6 764	5 899	4 113	3 409
- fuel oil	2 541	3 424	3 922	2 795	4 228	3 845	4 429	3 646	2 797	2 160	1 592	1 379	1 210	1 367	1 435

Fisheries, input NERI model

- LPG	-	-	34	29	50	42	34	30	12	18	16	36	5	1	16
- gasoline	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
- kerosene	7	2	9	5	12	26	9	5	4	3	4	3	3	2	0
- gas/diesel oil	8 932	10 029	8 143	9 222	9 729	10 080	10 431	10 453	8 389	8 301	7 740	8 169	8 120	8 510	8 556

International sea transport, input NERI model

- gas/diesel oil	-	7 867	8 547	9 743	10 514	11 633	12 590	16 881	19 114	24 123	26 743	27 231	25 325	31 243	26 085
- fuel oil	-	12 236	20 883	27 532	27 667	28 543	23 470	20 998	36 988	39 024	39 509	35 739	32 427	26 952	28 526

National sea transport, output NERI model

- gas/diesel oil	5285	5285	5285	5285	5285	5285	6015	6920	6673	6618	7028	8465	8967	7333	6201
- fuel oil	4571	4571	4571	4571	4571	4571	3926	3202	3201	3362	3382	2826	2052	1590	1455
- kerosene	5	0	5	3	1	0	2	1	1	1	1	1	0	1	0
- LPG	3	1	3	0	2	2	2	3	16	1	2	1	2	3	1

Fisheries, output NERI model

- gas/diesel oil	7064	8131	6233	7509	7455	7920	8170	7482	7075	7097	7134	6744	5329	5567	6375
- kerosene	7	2	9	5	12	26	9	5	4	3	4	3	3	2	0
- LPG	0	0	34	29	50	42	34	30	12	18	16	36	5	1	16

International sea transport, output NERI model

- gas/diesel oil	6828	7524	8204	9400	10171	11289	12149	16433	18726	23742	26370	26955	25049	30967	25474
- fuel oil	9394	11507	20155	26804	26938	27815	22742	20269	36259	38296	38780	35010	31698	26223	27797

Færge og fragtskibsestimat taget fra den internationale søtransport

729	729	729	729	729	729	729	729	729	729	729	729	729	729	729	729
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

National sea transport, added 0301

- fuel oil	-2 030	- 419	80	-1 048	386	3	1 233	1 174	325	-473	-1 061	- 718	- 113	506	709
------------	--------	-------	----	--------	-----	---	-------	-------	-----	------	--------	-------	-------	-----	-----

Road transport, NERI excl. traded fuels

- gasoline	64 492	67 041	69 220	71 819	72 664	72 882	73 874	74 714	75 342	77 074	79 674	81 385	83 976	85 223	87 867
- gas/diesel oil	45 609	49 738	49 626	49 686	51 854	54 746	58 427	57 511	56 796	58 755	58 561	59 851	60 528	61 072	63 619
- bioethanol	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
- biodiesel	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

Road transport, input NERI model incl. traded fuels

- gasoline	64 492	62 442	62 716	63 442	62 546	66 279	70 589	74 320	76 459	79 209	80 101	80 958	83 089	84 832	84 506
- gas/diesel oil	45 609	54 939	54 827	54 887	57 055	59 947	61 296	59 950	59 522	63 561	64 013	65 590	66 374	67 206	69 501
- bioethanol	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
- biodiesel	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

Sector	2 000	2 001	2 002	2 003	2 004	2 005	2 006	2 007	2 008	2 009	2 010	2 011	2 012
Agriculture, Forestry and Horticulture													
- LPG	199	210	176	169	152	133	128	121	114	126	138	142	137
- motor gasoline	41	43	31	47	57	58	22	23	22	34	33	28	33
- gas/diesel oil	14387	14018	14235	14019	13422	12871	13437	12899	14142	14855	14984	14609	14528
Fishery	0	0	0	0	0	0	0	0	0	0	0	0	0
- LPG	13	19	21	20	18	20	20	18	12	12	12	11	10
- motor gasoline	67	3	3	0	0	0	1	1	1	1	1	1	1
- petroleum	25	1	1	1	1	1	0	0	0	0	0	0	0
- gas/diesel oil	9347	8908	8888	8428	7337	7340	7362	6854	6258	6075	6037	5739	4659
- fuel oil	0	0	4	84	35	126	86	13	14	17	0	0	0
Manufacturing Industry	0	0	0	0	0	0	0	0	0	0	0	0	0
- LPG	1819	1526	1405	1472	1488	1478	1482	1216	1178	1029	1093	986	1150
- motor gasoline	97	69	42	26	30	21	32	16	15	97	84	118	15
- gas/diesel oil	8635	10099	9155	9964	10515	10022	9132	8170	7449	6141	6242	4902	4234
- fuel oil	8221	7395	7818	6916	6940	6055	8527	6422	5319	4015	5032	3835	3388
Construction Industry	0	0	0	0	0	0	0	0	0	0	0	0	0
- LPG	165	179	236	226	228	224	248	222	172	103	94	98	104
- motor gasoline	33	24	26	27	27	27	27	28	26	20	22	21	16
- gas/diesel oil	5950	6356	6226	6226	6227	6338	6187	6410	6339	5429	5341	5370	4353
Single Family Houses	0	0	0	0	0	0	0	0	0	0	0	0	0
- motor gasoline	1355	1317	1313	1303	1288	1250	1216	1193	1135	1079	1031	958	912
- gas/diesel oil	27929	28996	26967	24932	22863	21712	19572	18012	16585	15625	16536	13698	11162

Multi-family Houses	0	0	0	0	0	0	0	0	0	0	0	0	0
- gas/diesel oil	2346	2511	2031	2095	2427	2151	1625	1411	1610	1658	1630	1305	1388
Road transport, DEA statistics													
- gasoline	88 975	86 474	86 247	85 611	84 629	82 118	79 822	78 325	74 545	70 880	67 720	62 879	59 864
- gas/diesel oil	64 282	66 254	66 814	70 875	75 422	79 476	86 223	93 111	93 437	88 454	92 359	91 185	87 187
- bioethanol	-	-	-	-	-	-	151	252	210	204	1 118	2 124	2 003
- biodiesel	-	-	-	-	-	-	-	-	10	139	16	3 492	6 526
Non-road, DEA statistics													
- LPG	2 018	1 736	1 581	1 641	1 640	1 612	1 610	1 337	1 292	1 155	1 232	1 128	1 287
- gasoline	1 525	1 453	1 412	1 404	1 402	1 356	1 296	1 259	1 199	1 230	1 170	1 125	975
- gas/diesel oil	28 972	30 473	29 616	30 209	30 164	29 232	28 757	27 479	27 929	26 425	26 567	24 881	23 115
Non-road, NERI model													
- LPG	1071	1073	1084	1079	1065	1049	1038	1040	986	817	985	976	974
- gasoline	2458	2622	2833	3090	3391	3604	3807	3923	3975	3942	3982	3974	3948
- gas/diesel oil	24630	24923	25100	25301	25670	26418	27784	29727	30914	27321	30280	30243	30620
Recreational craft, NERI model													
- gasoline	396	400	403	404	404	393	382	371	361	353	346	340	335
- gas/diesel oil	777	831	886	944	1002	1002	1002	1002	1002	1002	1002	1002	1002
Non-road, added 0202													
- gas/diesel oil	0	0	0	0	0	0	0	-2248	-2984	-897	-3712	-5362	-7505
Non-road, added 0203 and 0301													
- gas/diesel oil	4342	5550	4516	4908	4494	2814	974	0	0	0	0	0	0
- LPG	947	662	497	563	575	562	572	298	306	338	247	152	313
Non-road, added 0203													
- gas/diesel oil	2156	2553	2171	2278	2000	1239	455	0	0	0	0	0	0
- LPG	93	80	55	58	53	46	46	27	27	37	28	19	33
Non-road, added 0301													
- gas/diesel oil	2186	2997	2346	2630	2494	1575	519	0	0	0	0	0	0
- LPG	854	582	442	505	522	516	526	271	279	301	219	133	279

Non-road, added road transport													
- gasoline	-932	-1169	-1421	-1686	-1990	-2248	-2511	-2663	-2776	-2712	-2812	-2849	-2973
Fisheries, added national sea transport													
- fuel oil	0	0	4	84	35	126	86	13	14	17	0	0	0
Fisheries, consumed by recreational craft													
- gasoline	67	3	3	0	0	0	1	1	1	1	1	1	1
National sea transport, input NERI model													
- LPG	0	-	-	0	0	0	0	0	-	-	-	-	-
- kerosene	1	1	1	1	1	1	0	-	-	-	-	-	-
- gas/diesel oil	5 348	5 608	5 855	6 009	5 259	6 646	5 986	5 233	6 954	6 489	5 665	5 654	5 595
- fuel oil	1 509	1 513	2 068	1 907	1 704	1 506	1 367	1 110	1 174	1 062	868	732	624
Fisheries, input NERI model													
- LPG	13	19	21	20	18	20	20	18	12	12	12	11	10
- gasoline	-	-	-	-	-	-	-	-	-	-	-	1	1
- kerosene	25	1	1	1	1	1	0	0	0	-	-	-	-
- gas/diesel oil	8 570	8 077	8 001	7 484	6 335	6 338	6 360	5 852	5 256	5 073	5 035	4 737	3 657
International sea transport, input NERI model													
- gas/diesel oil	20 892	19 022	19 505	18 549	14 357	11 630	10 829	9 124	11 218	10 433	11 493	10 432	10 431
- fuel oil	33 165	25 924	17 547	20 462	17 298	20 591	31 565	35 243	27 164	11 091	17 493	18 909	11 278
National sea transport, output NERI model													
- gas/diesel oil	5258	5233	5061	4475	4591	4559	4426	4435	4357	4235	4049	3714	2903
- fuel oil	1444	1400	1387	1862	1853	1859	2026	2004	2148	2287	2456	2375	2353
- kerosene	1	1	1	1	1	1	0	0	0	0	0	0	0
- LPG	0	0	0	0	0	0	0	0	0	0	0	0	10
Fisheries, output NERI model													
- gas/diesel oil	9403	9384	9664	9294	7286	8725	8166	6966	8142	7597	6787	6827	6466
- kerosene	25	1	1	1	1	1	0	0	0	0	0	0	0
- LPG	13	19	21	20	18	20	20	18	12	12	12	11	10

International sea transport, output NERI model													
- gas/diesel oil	20148	18090	18636	18273	14074	11330	10583	8809	10928	10164	11356	10282	10314
- fuel oil	32437	25195	16818	19247	16118	19411	30172	33848	25650	9416	15682	17120	9515
	729	729	729	1215	1180	1180	1393	1396	1514	1675	1812	1789	1764
National sea transport, added 0301													
- fuel oil	794	842	1 409	1 260	1 032	826	734	502	540	450	224	145	34
Road transport, NERI excl. traded fuels													
- gasoline	87 713	84 907	84 426	83 521	82 235	79 477	76 930	75 292	71 409	67 815	64 564	59 691	56 556
- gas/diesel oil	64 282	66 254	66 814	70 875	75 422	79 476	86 223	93 111	93 437	88 454	92 359	91 185	87 187
- bioethanol	-	-	-	-	-	-	151	252	210	204	1 118	2 124	2 003
- biodiesel	-	-	-	-	-	-	-	-	10	139	16	3 492	6 526
Road transport, input NERI model incl. traded fuels													
- gasoline	83 312	81 852	81 963	81 878	80 593	77 835	76 109	75 292	71 409	67 815	62 757	57 884	54 750
- gas/diesel oil	69 196	70 916	72 552	78 766	84 209	88 264	95 010	103 871	103 480	97 421	101 326	101 658	97 661
- bioethanol	-	-	-	-	-	-	151	252	210	204	1 118	2 124	2 003
- biodiesel	-	-	-	-	-	-	-	-	10	139	16	3 492	6 526

Annex 3B-15 Emission factors and total emissions in CollectER format

1990 emission factors for SO₂, NO_x, NMVOC, CH₄, CO, CO₂, N₂O, NH₃, TSP; PM₁₀ and PM_{2.5}.

Year	SNAP ID	Category	Fuel type	SO ₂	NO _x	NMVOC	CH ₄	CO	CO ₂	N ₂ O	NH ₃	TSP	PM ₁₀	PM _{2.5}		
				g pr GJ	g pr GJ	g pr GJ	g pr GJ	g pr GJ	g pr GJ	g pr GJ	g pr GJ	g pr GJ	g pr GJ	g pr GJ	g pr GJ	
1990	A	070101	Passenger cars	Highway	Diesel	93,68	279,53	25,07	3,74	179,70	74,00	0,00	0,47	79,48	79,48	79,48
1990	A	070101	Passenger cars	Highway	Gasoline	2,28	1315,35	381,72	10,89	3817,22	73,00	2,72	0,84	12,84	12,84	12,84
1990	A	070101	Passenger cars	Highway	LPG	0,00	1151,70	187,09	10,06	3914,25	63,10	0,00	0,00	10,06	10,06	10,06
1990	A	070102	Passenger cars	Rural	Diesel	93,68	280,57	42,09	6,82	268,08	74,00	0,00	0,57	75,13	75,13	75,13
1990	A	070102	Passenger cars	Rural	Gasoline	2,28	1148,38	500,38	13,79	4273,20	73,00	3,08	0,95	14,97	14,97	14,97
1990	A	070102	Passenger cars	Rural	LPG	0,00	1248,46	305,18	16,91	1146,38	63,10	0,00	0,00	14,49	14,49	14,49
1990	A	070103	Passenger cars	Urban	Diesel	93,68	220,85	103,04	6,82	344,97	74,00	0,00	0,32	144,24	144,24	144,24
1990	A	070103	Passenger cars	Urban	Gasoline	2,28	558,41	1095,32	56,50	12514,04	73,00	2,71	0,54	12,37	12,37	12,37
1990	A	070103	Passenger cars	Urban	LPG	0,00	528,01	473,05	20,80	1587,66	63,10	0,00	0,00	10,40	10,40	10,40
1990	A	070201	Light duty vehicles	Highway	Diesel	93,68	270,67	30,19	2,60	344,14	74,00	0,00	0,32	104,48	104,48	104,48
1990	A	070201	Light duty vehicles	Highway	Gasoline	2,28	1369,26	170,29	10,11	2987,40	73,00	2,63	0,81	16,17	16,17	16,17
1990	A	070201	Light duty vehicles	Highway	LPG	0,00	1151,70	187,09	10,06	3914,25	63,10	0,00	0,00	10,06	10,06	10,06
1990	A	070202	Light duty vehicles	Rural	Diesel	93,68	299,25	33,22	4,26	358,42	74,00	0,00	0,36	107,73	107,73	107,73
1990	A	070202	Light duty vehicles	Rural	Gasoline	2,28	1188,86	262,59	15,25	2316,18	73,00	2,48	0,76	15,25	15,25	15,25
1990	A	070202	Light duty vehicles	Rural	LPG	0,00	1248,46	305,18	16,91	1146,38	63,10	0,00	0,00	14,49	14,49	14,49
1990	A	070203	Light duty vehicles	Urban	Diesel	93,68	469,24	74,81	4,58	463,37	74,00	0,00	0,23	165,76	165,76	165,76
1990	A	070203	Light duty vehicles	Urban	Gasoline	2,28	547,55	874,08	39,84	9527,46	73,00	1,84	0,37	7,37	7,37	7,37
1990	A	070203	Light duty vehicles	Urban	LPG	0,00	487,91	487,73	19,58	1705,63	63,10	0,00	0,00	9,79	9,79	9,79
1990	A	070301	Heavy duty vehicles	Highway	Diesel	93,68	978,47	41,71	6,44	200,02	74,00	3,08	0,31	35,10	35,10	35,10
1990	A	070301	Heavy duty vehicles	Highway	Gasoline	2,28	1037,78	474,61	9,69	7610,35	73,00	0,83	0,28	55,35	55,35	55,35
1990	A	070302	Heavy duty vehicles	Rural	Diesel	93,68	984,85	57,74	6,78	211,25	74,00	2,92	0,29	35,73	35,73	35,73
1990	A	070302	Heavy duty vehicles	Rural	Gasoline	2,28	1141,55	820,40	16,74	8371,39	73,00	0,91	0,30	60,88	60,88	60,88
1990	A	070303	Heavy duty vehicles	Urban	Diesel	93,68	966,99	86,86	12,42	269,91	74,00	2,30	0,23	42,21	42,21	42,21
1990	A	070303	Heavy duty vehicles	Urban	Gasoline	2,28	456,62	696,09	14,21	7102,99	73,00	0,61	0,20	40,59	40,59	40,59
1990	A	070400	Mopeds	Urban	Gasoline	2,28	18,26	12503,20	200,00	12602,74	73,00	0,91	0,91	171,69	171,69	171,69
1990	A	070501	Motorcycles	Highway	Gasoline	2,28	264,11	1072,19	129,96	16302,60	73,00	1,35	1,35	31,73	31,73	31,73
1990	A	070502	Motorcycles	Rural	Gasoline	2,28	185,41	981,69	159,32	15782,07	73,00	1,66	1,66	38,90	38,90	38,90
1990	A	070503	Motorcycles	Urban	Gasoline	2,28	112,92	1149,21	155,11	15187,59	73,00	1,61	1,61	37,87	37,87	37,87
1990	A	080100	Military		AvGas	22,83	859,00	1242,60	21,90	6972,00	73,00	2,00	1,60	10,00	10,00	10,00
1990	A	080100	Military		Diesel	93,68	719,72	55,39	6,74	268,73	74,00	1,73	0,32	66,49	66,49	66,49
1990	A	080100	Military		Gasoline	2,28	991,97	1128,00	27,57	6765,35	73,00	2,85	0,79	14,34	14,34	14,34

1990	A	080100	Military	Jet fuel	22,99	250,57	24,94	2,65	229,89	72,00	2,30	0,00	1,16	1,16	1,16
1990	A	080200	Railways	Diesel	93,68	1225,13	79,94	3,07	223,21	74,00	2,04	0,20	50,26	50,26	50,26
1990	A	080200	Railways	Kerosene	5,00	50,00	3,00	7,00	20,00	72,00	2,00	0,00	121,95	115,85	110,06
1990	A	080300	Inland waterways	Diesel	93,68	983,64	171,79	2,79	453,65	74,00	2,96	0,17	106,93	106,93	106,93
1990	A	080300	Inland waterways	Gasoline	2,28	291,33	3606,55	50,38	13853,27	73,00	0,78	0,08	182,44	182,44	182,44
1990	A	080402	National sea traffic	Diesel	93,68	1104,18	50,57	1,56	166,83	74,00	4,68	0,00	23,21	22,98	22,87
1990	A	080402	National sea traffic	Kerosene	2,30	50,00	3,00	7,00	20,00	72,00	0,00	0,00	5,00	5,00	5,00
1990	A	080402	National sea traffic	LPG	0,00	1249,00	384,94	20,26	443,00	63,10	0,00	0,00	0,20	0,20	0,20
1990	A	080402	National sea traffic	Residual oil	1290,95	1615,26	53,44	1,65	176,29	78,00	4,89	0,00	149,25	147,75	147,01
1990	A	080403	Fishing	Diesel	93,68	1052,12	49,13	1,52	162,08	74,00	4,68	0,00	23,21	22,98	22,87
1990	A	080403	Fishing	Kerosene	2,30	50,00	3,00	7,00	20,00	72,00	0,00	0,00	5,00	5,00	5,00
1990	A	080403	Fishing	LPG	0,00	1249,00	384,94	20,26	443,00	63,10	0,00	0,00	0,20	0,20	0,20
1990	A	080404	International sea traffic	Diesel	93,68	1208,60	49,46	1,53	163,17	74,00	4,68	0,00	23,21	22,98	22,87
1990	A	080404	International sea traffic	Residual oil	1447,43	1689,57	53,98	1,67	178,09	78,00	4,89	0,00	189,43	187,53	186,59
1990	A	080501	Air traffic, Dom. < 3000 ft.	AvGas	22,83	859,00	1242,60	21,90	6972,00	73,00	2,00	1,60	10,00	10,00	10,00
1990	A	080501	Air traffic, Dom. < 3000 ft.	Jet fuel	22,99	314,51	14,93	1,59	90,41	72,00	5,70	0,00	1,16	1,16	1,16
1990	A	080502	Air traffic, Int. < 3000 ft.	AvGas	22,83	859,00	1242,60	21,90	6972,00	73,00	2,00	1,60	10,00	10,00	10,00
1990	A	080502	Air traffic, Int. < 3000 ft.	Jet fuel	22,99	309,25	16,47	1,75	168,98	72,00	7,10	0,00	1,16	1,16	1,16
1990	A	080503	Air traffic, Dom. > 3000 ft.	Jet fuel	22,99	330,11	12,36	1,31	90,75	72,00	2,30	0,00	1,16	1,16	1,16
1990	A	080504	Air traffic, Int. > 3000 ft.	Jet fuel	22,99	244,20	6,48	0,69	54,10	72,00	2,30	0,00	1,16	1,16	1,16
1990	A	080600	Agriculture	Diesel	93,68	758,87	156,85	2,55	635,53	74,00	2,93	0,17	144,45	144,45	144,45
1990	A	080600	Agriculture	Gasoline	2,28	31,60	949,55	88,42	47524,17	73,00	1,28	0,09	6,56	6,56	6,56
1990	A	080700	Forestry	Diesel	93,68	857,48	156,47	2,54	645,65	74,00	2,97	0,17	149,05	149,05	149,05
1990	A	080700	Forestry	Gasoline	2,28	40,39	7206,91	60,42	18057,40	73,00	0,37	0,07	101,22	101,22	101,22
1990	A	080800	Industry	Diesel	93,68	933,58	178,23	2,90	655,80	74,00	2,94	0,17	154,50	154,50	154,50
1990	A	080800	Industry	Gasoline	2,28	136,27	1610,77	120,61	14797,46	73,00	1,33	0,09	12,40	12,40	12,40
1990	A	080800	Industry	LPG	0,00	1328,11	146,09	7,69	104,85	63,10	3,50	0,21	4,89	4,89	4,89
1990	A	080900	Household and gardening	Gasoline	2,28	63,98	3366,01	95,22	32901,19	73,00	1,15	0,08	20,75	20,75	20,75
1990	A	081100	Commercial and institutional	Gasoline	2,28	68,83	2280,66	97,87	29887,31	73,00	1,09	0,08	24,00	24,00	24,00
1990	P	080501	Air traffic, Dom. < 3000 ft.	AvGas	22,83	859,00	1242,60	21,90	6972,00	73,00	2,00	1,60	10,00	10,00	10,00
1990	P	080501	Air traffic, Dom. < 3000 ft.	Jet fuel	22,99	283,87	20,73	2,20	129,70	72,00	4,58	0,00	1,16	1,16	1,16
1990	P	080502	Air traffic, Int. < 3000 ft.	AvGas	22,83	859,00	1242,60	21,90	6972,00	73,00	2,00	1,60	10,00	10,00	10,00
1990	P	080502	Air traffic, Int. < 3000 ft.	Jet fuel	22,99	324,87	34,25	3,64	157,15	72,00	3,79	0,00	1,16	1,16	1,16
1990	P	080503	Air traffic, Dom. > 3000 ft.	Jet fuel	22,99	314,86	11,78	1,25	84,05	72,00	2,30	0,00	1,16	1,16	1,16
1990	P	080504	Air traffic, Int. > 3000 ft.	Jet fuel	22,99	290,20	10,08	1,07	37,65	72,00	2,30	0,00	1,16	1,16	1,16

1990 emission factors for Arsenic, Cadmium, Chromium, Copper, Mercury, Nickel, Lead, Selenium and Zinc.

Year	SNAP ID	Category	Fuel type	Arsenic mg pr GJ	Cadmium mg pr GJ	Chromium mg pr GJ	Copper mg pr GJ	Mercury mg pr GJ	Nickel mg pr GJ	Lead mg pr GJ	Selenium mg pr GJ	Zinc mg pr GJ		
1990	A	070101	Passenger cars	Highway	Diesel	0,002	0,292	0,926	0,657	0,124	0,296	1,757	0,002	58,606
1990	A	070101	Passenger cars	Highway	Gasoline	0,007	0,273	0,386	1,043	0,199	0,321	1472,004	0,005	54,458
1990	A	070101	Passenger cars	Highway	LPG	0,000	0,300	0,270	1,052	0,000	0,300	0,901	0,000	60,097
1990	A	070102	Passenger cars	Rural	Diesel	0,002	0,355	1,083	0,770	0,124	0,358	2,133	0,002	71,144
1990	A	070102	Passenger cars	Rural	Gasoline	0,007	0,308	0,417	1,165	0,199	0,356	1472,109	0,005	61,472
1990	A	070102	Passenger cars	Rural	LPG	0,000	0,361	0,325	1,262	0,000	0,361	1,082	0,000	72,116
1990	A	070103	Passenger cars	Urban	Diesel	0,002	0,198	0,691	0,488	0,124	0,201	1,192	0,002	39,770
1990	A	070103	Passenger cars	Urban	Gasoline	0,007	0,178	0,300	0,711	0,199	0,226	1471,720	0,005	35,492
1990	A	070103	Passenger cars	Urban	LPG	0,000	0,194	0,175	0,679	0,000	0,194	0,582	0,000	38,818
1990	A	070201	Light duty vehicles	Highway	Diesel	0,002	0,193	0,678	0,478	0,124	0,196	1,161	0,002	38,731
1990	A	070201	Light duty vehicles	Highway	Gasoline	0,007	0,253	0,367	0,972	0,199	0,301	1471,943	0,005	50,405
1990	A	070201	Light duty vehicles	Highway	LPG	0,000	0,198	0,178	0,693	0,000	0,198	0,594	0,000	39,594
1990	A	070202	Light duty vehicles	Rural	Diesel	0,002	0,211	0,723	0,511	0,124	0,214	1,269	0,002	42,337
1990	A	070202	Light duty vehicles	Rural	Gasoline	0,007	0,239	0,355	0,922	0,199	0,287	1471,901	0,005	47,576
1990	A	070202	Light duty vehicles	Rural	LPG	0,000	0,238	0,214	0,831	0,000	0,238	0,713	0,000	47,513
1990	A	070203	Light duty vehicles	Urban	Diesel	0,002	0,135	0,533	0,374	0,124	0,138	0,813	0,002	27,133
1990	A	070203	Light duty vehicles	Urban	Gasoline	0,007	0,118	0,246	0,499	0,199	0,166	1471,538	0,005	23,388
1990	A	070203	Light duty vehicles	Urban	LPG	0,000	0,120	0,108	0,421	0,000	0,120	0,361	0,000	24,068
1990	A	070301	Heavy duty vehicles	Highway	Diesel	0,002	0,150	0,570	0,401	0,124	0,153	0,903	0,002	30,130
1990	A	070301	Heavy duty vehicles	Highway	Gasoline	0,007	0,207	0,326	0,813	0,199	0,255	1471,807	0,005	41,332
1990	A	070302	Heavy duty vehicles	Rural	Diesel	0,002	0,143	0,555	0,390	0,124	0,147	0,866	0,002	28,884
1990	A	070302	Heavy duty vehicles	Rural	Gasoline	0,007	0,220	0,338	0,857	0,199	0,268	1471,845	0,005	43,830
1990	A	070303	Heavy duty vehicles	Urban	Diesel	0,002	0,115	0,483	0,338	0,124	0,118	0,693	0,002	23,142
1990	A	070303	Heavy duty vehicles	Urban	Gasoline	0,007	0,143	0,268	0,586	0,199	0,190	1471,613	0,005	28,351
1990	A	070400	Mopeds	Urban	Gasoline	0,007	0,005	0,144	0,103	0,199	0,053	1471,199	0,005	0,753
1990	A	070501	Motorcycles	Highway	Gasoline	0,007	0,134	0,260	0,556	0,199	0,182	1471,587	0,005	26,666
1990	A	070502	Motorcycles	Rural	Gasoline	0,007	0,163	0,287	0,659	0,199	0,211	1471,675	0,005	32,520
1990	A	070503	Motorcycles	Urban	Gasoline	0,007	0,159	0,283	0,644	0,199	0,207	1471,663	0,005	31,681
1990	A	080100	Military		AvGas	0,007	0,253	0,367	0,972	0,198	0,301	12785,388	0,005	50,452
1990	A	080100	Military		Diesel	0,002	0,172	0,625	0,440	0,124	0,175	1,035	0,002	34,517
1990	A	080100	Military		Gasoline	0,007	0,256	0,371	0,984	0,199	0,304	1471,954	0,005	51,135
1990	A	080100	Military		Jet fuel	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
1990	A	080200	Railways		Diesel	0,002	0,186	0,660	0,465	0,124	0,189	1,118	0,002	37,295
1990	A	080200	Railways		Kerosene	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000

1990	A	080300	Inland waterways	Diesel	0,002	0,186	0,660	0,465	0,124	0,189	1,118	0,002	37,295
1990	A	080300	Inland waterways	Gasoline	0,007	0,253	0,367	0,972	0,198	0,301	0,773	0,005	50,452
1990	A	080402	National sea traffic	Diesel	1,171	0,234	0,937	1,171	1,170	1,639	2,340	4,684	11,710
1990	A	080402	National sea traffic	Kerosene	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
1990	A	080402	National sea traffic	LPG	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
1990	A	080402	National sea traffic	Residual oil	12,225	0,733	4,890	12,225	0,490	733,496	4,890	9,780	22,005
1990	A	080403	Fishing	Diesel	1,171	0,234	0,937	1,171	1,170	1,639	2,340	4,684	11,710
1990	A	080403	Fishing	Kerosene	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
1990	A	080403	Fishing	LPG	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
1990	A	080404	International sea traffic	Diesel	1,171	0,234	0,937	1,171	1,170	1,639	2,340	4,684	11,710
1990	A	080404	International sea traffic	Residual oil	12,225	0,733	4,890	12,225	0,490	733,496	4,890	9,780	22,005
1990	A	080501	Air traffic, Dom. < 3000 ft.	AvGas	0,007	0,253	0,367	0,972	0,198	0,301	13505,692	0,005	50,452
1990	A	080501	Air traffic, Dom. < 3000 ft.	Jet fuel	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
1990	A	080502	Air traffic, Int. < 3000 ft.	AvGas	0,007	0,253	0,367	0,972	0,198	0,301	13505,692	0,005	50,452
1990	A	080502	Air traffic, Int. < 3000 ft.	Jet fuel	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
1990	A	080503	Air traffic, Dom. > 3000 ft.	Jet fuel	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
1990	A	080504	Air traffic, Int. > 3000 ft.	Jet fuel	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
1990	A	080600	Agriculture	Diesel	0,002	0,186	0,660	0,465	0,124	0,189	1,118	0,002	37,295
1990	A	080600	Agriculture	Gasoline	0,007	0,253	0,367	0,972	0,198	0,301	0,773	0,005	50,452
1990	A	080700	Forestry	Diesel	0,002	0,186	0,660	0,465	0,124	0,189	1,118	0,002	37,295
1990	A	080700	Forestry	Gasoline	0,007	0,253	0,367	0,972	0,198	0,301	0,773	0,005	50,452
1990	A	080800	Industry	Diesel	0,002	0,186	0,660	0,465	0,124	0,189	1,118	0,002	37,295
1990	A	080800	Industry	Gasoline	0,007	0,253	0,367	0,972	0,198	0,301	0,773	0,005	50,452
1990	A	080800	Industry	LPG	0,000	0,131	0,118	0,457	0,000	0,131	0,392	0,000	26,126
1990	A	080900	Household and gardening	Gasoline	0,007	0,253	0,367	0,972	0,198	0,301	0,773	0,005	50,452
1990	A	081100	Commercial and institutional	Gasoline	0,007	0,253	0,367	0,972	0,198	0,301	0,773	0,005	50,452
1990	P	080501	Air traffic, Dom. < 3000 ft.	AvGas	0,007	0,253	0,367	0,972	0,198	0,301	13505,692	0,005	50,452
1990	P	080501	Air traffic, Dom. < 3000 ft.	Jet fuel	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
1990	P	080502	Air traffic, Int. < 3000 ft.	AvGas	0,007	0,253	0,367	0,972	0,198	0,301	13505,692	0,005	50,452
1990	P	080502	Air traffic, Int. < 3000 ft.	Jet fuel	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
1990	P	080503	Air traffic, Dom. > 3000 ft.	Jet fuel	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
1990	P	080504	Air traffic, Int. > 3000 ft.	Jet fuel	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000

1990 emission factors for Dioxins/, Flouranthene, Benzo(b), Benzo(k), Benzo(a), Benzo(g,h,i) and indeno(1,2,3-c,d).

Year	SNAP ID	Category	Fuel type	Dioxins/ microg pr GJ	Flouranthene mg pr GJ	Benzo(b) mg pr GJ	Benzo(k) mg pr GJ	Benzo(a) mg pr GJ	Benzo(g,h,i) mg pr GJ	indeno(1,2,3-c,d) mg pr GJ
1990	A 070101	Passenger cars	Highway Diesel	0,001	12,250	0,748	0,678	0,818	1,589	0,771
1990	A 070101	Passenger cars	Highway Gasoline	0,013	8,507	0,553	0,425	0,468	1,106	0,425
1990	A 070101	Passenger cars	Highway LPG	0,000	0,000	0,000	0,000	0,000	0,000	0,000
1990	A 070102	Passenger cars	Rural Diesel	0,001	14,889	0,909	0,824	0,994	1,932	0,937
1990	A 070102	Passenger cars	Rural Gasoline	0,015	9,540	0,620	0,477	0,525	1,240	0,477
1990	A 070102	Passenger cars	Rural LPG	0,000	0,000	0,000	0,000	0,000	0,000	0,000
1990	A 070103	Passenger cars	Urban Diesel	0,001	9,303	0,568	0,515	0,621	1,207	0,586
1990	A 070103	Passenger cars	Urban Gasoline	0,010	6,427	0,418	0,321	0,354	0,835	0,321
1990	A 070103	Passenger cars	Urban LPG	0,000	0,000	0,000	0,000	0,000	0,000	0,000
1990	A 070201	Light duty vehicles	Highway Diesel	0,000	8,505	0,519	0,470	0,568	1,104	0,536
1990	A 070201	Light duty vehicles	Highway Gasoline	0,013	8,086	0,526	0,404	0,445	1,051	0,404
1990	A 070201	Light duty vehicles	Highway LPG	0,000	0,000	0,000	0,000	0,000	0,000	0,000
1990	A 070202	Light duty vehicles	Rural Diesel	0,001	9,306	0,568	0,515	0,622	1,207	0,586
1990	A 070202	Light duty vehicles	Rural Gasoline	0,012	7,625	0,495	0,381	0,419	0,991	0,381
1990	A 070202	Light duty vehicles	Rural LPG	0,000	0,000	0,000	0,000	0,000	0,000	0,000
1990	A 070203	Light duty vehicles	Urban Diesel	0,000	6,954	0,425	0,385	0,464	0,902	0,438
1990	A 070203	Light duty vehicles	Urban Gasoline	0,007	4,558	0,296	0,228	0,251	0,592	0,228
1990	A 070203	Light duty vehicles	Urban LPG	0,000	0,000	0,000	0,000	0,000	0,000	0,000
1990	A 070301	Heavy duty vehicles	Highway Diesel	0,001	2,086	0,526	0,780	0,097	0,078	0,136
1990	A 070301	Heavy duty vehicles	Highway Gasoline	0,000	0,000	0,000	0,000	0,000	0,000	0,000
1990	A 070302	Heavy duty vehicles	Rural Diesel	0,001	2,208	0,557	0,825	0,103	0,082	0,144
1990	A 070302	Heavy duty vehicles	Rural Gasoline	0,000	0,000	0,000	0,000	0,000	0,000	0,000
1990	A 070303	Heavy duty vehicles	Urban Diesel	0,001	1,788	0,451	0,668	0,083	0,067	0,117
1990	A 070303	Heavy duty vehicles	Urban Gasoline	0,000	0,000	0,000	0,000	0,000	0,000	0,000
1990	A 070400	Mopeds	Urban Gasoline	0,000	0,000	0,000	0,000	0,000	0,000	0,000
1990	A 070501	Motorcycles	Highway Gasoline	0,020	12,673	0,824	0,634	0,697	1,647	0,634
1990	A 070502	Motorcycles	Rural Gasoline	0,024	15,176	0,986	0,759	0,834	1,973	0,759
1990	A 070503	Motorcycles	Urban Gasoline	0,024	15,300	0,994	0,765	0,841	1,989	0,765
1990	A 080100	Military	AvGas	0,005	4,329	0,209	0,071	0,114	0,689	0,245
1990	A 080100	Military	Diesel	0,001	4,391	0,571	0,568	0,290	0,550	0,290
1990	A 080100	Military	Gasoline	0,006	5,257	0,277	0,116	0,142	0,825	0,300
1990	A 080100	Military	Jet fuel	0,000	0,000	0,000	0,000	0,000	0,000	0,000
1990	A 080200	Railways	Diesel	0,001	1,366	0,348	0,389	0,057	0,049	0,089
1990	A 080200	Railways	Kerosene	0,000	0,000	0,000	0,000	0,000	0,000	0,000

1990	A	080300	Inland waterways	Diesel	0,001	4,391	0,571	0,568	0,290	0,550	0,290
1990	A	080300	Inland waterways	Gasoline	0,005	4,329	0,209	0,071	0,114	0,689	0,245
1990	A	080402	National sea traffic	Diesel	0,012	7,420	0,640	0,300	0,150	1,430	1,180
1990	A	080402	National sea traffic	Kerosene	0,000	0,000	0,000	0,000	0,000	0,000	0,000
1990	A	080402	National sea traffic	LPG	0,000	0,000	0,000	0,000	0,000	0,000	0,000
1990	A	080402	National sea traffic	Residual oil	0,013	5,190	0,270	0,050	0,020	0,070	0,030
1990	A	080403	Fishing	Diesel	0,012	7,420	0,640	0,300	0,150	1,430	1,180
1990	A	080403	Fishing	Kerosene	0,000	0,000	0,000	0,000	0,000	0,000	0,000
1990	A	080403	Fishing	LPG	0,000	0,000	0,000	0,000	0,000	0,000	0,000
1990	A	080404	International sea traffic	Diesel	0,012	7,420	0,640	0,300	0,150	1,430	1,180
1990	A	080404	International sea traffic	Residual oil	0,013	4,120	0,200	0,090	0,070	0,260	0,200
1990	A	080501	Air traffic, Dom. < 3000 ft.	AvGas	0,005	4,329	0,209	0,071	0,114	0,689	0,245
1990	A	080501	Air traffic, Dom. < 3000 ft.	Jet fuel	0,000	0,000	0,000	0,000	0,000	0,000	0,000
1990	A	080502	Air traffic, Int. < 3000 ft.	AvGas	0,005	4,329	0,209	0,071	0,114	0,689	0,245
1990	A	080502	Air traffic, Int. < 3000 ft.	Jet fuel	0,000	0,000	0,000	0,000	0,000	0,000	0,000
1990	A	080503	Air traffic, Dom. > 3000 ft.	Jet fuel	0,000	0,000	0,000	0,000	0,000	0,000	0,000
1990	A	080504	Air traffic, Int. > 3000 ft.	Jet fuel	0,000	0,000	0,000	0,000	0,000	0,000	0,000
1990	A	080600	Agriculture	Diesel	0,001	4,391	0,571	0,568	0,290	0,550	0,290
1990	A	080600	Agriculture	Gasoline	0,005	4,329	0,209	0,071	0,114	0,689	0,245
1990	A	080700	Forestry	Diesel	0,001	4,391	0,571	0,568	0,290	0,550	0,290
1990	A	080700	Forestry	Gasoline	0,005	4,329	0,209	0,071	0,114	0,689	0,245
1990	A	080800	Industry	Diesel	0,001	4,391	0,571	0,568	0,290	0,550	0,290
1990	A	080800	Industry	Gasoline	0,005	4,329	0,209	0,071	0,114	0,689	0,245
1990	A	080800	Industry	LPG	0,000	0,000	0,000	0,000	0,000	0,000	0,000
1990	A	080900	Household and gardening	Gasoline	0,005	4,329	0,209	0,071	0,114	0,689	0,245
1990	A	081100	Commercial and institutional	Gasoline	0,005	4,329	0,209	0,071	0,114	0,689	0,245
1990	P	080501	Air traffic, Dom. < 3000 ft.	AvGas	0,005	4,329	0,209	0,071	0,114	0,689	0,245
1990	P	080501	Air traffic, Dom. < 3000 ft.	Jet fuel	0,000	0,000	0,000	0,000	0,000	0,000	0,000
1990	P	080502	Air traffic, Int. < 3000 ft.	AvGas	0,005	4,329	0,209	0,071	0,114	0,689	0,245
1990	P	080502	Air traffic, Int. < 3000 ft.	Jet fuel	0,000	0,000	0,000	0,000	0,000	0,000	0,000
1990	P	080503	Air traffic, Dom. > 3000 ft.	Jet fuel	0,000	0,000	0,000	0,000	0,000	0,000	0,000
1990	P	080504	Air traffic, Int. > 3000 ft.	Jet fuel	0,000	0,000	0,000	0,000	0,000	0,000	0,000

1990 emission factors for SO₂, NO_x, NMVOC, CH₄, CO, CO₂, N₂O, NH₃, TSP, PM₁₀ and PM_{2.5}.

Year	SNAP ID	Category	Fuel type	SO ₂ g pr GJ	NO _x g pr GJ	NMVOC g pr GJ	CH ₄ g pr GJ	CO g pr GJ	CO ₂ g pr GJ	N ₂ O g pr GJ	NH ₃ g pr GJ	TSP g pr GJ	PM ₁₀ g pr GJ	PM _{2.5} g pr GJ
1990	A 070101	Passenger cars	Highway Diesel	93,68	279,53	25,07	3,74	179,70	74,00	0,00	0,47	79,48	79,48	79,48
1990	A 070101	Passenger cars	Highway Gasoline	2,28	1315,35	381,72	10,89	3817,22	73,00	2,72	0,84	12,84	12,84	12,84
1990	A 070101	Passenger cars	Highway LPG	0,00	1151,70	187,09	10,06	3914,25	63,10	0,00	0,00	10,06	10,06	10,06
1990	A 070102	Passenger cars	Rural Diesel	93,68	280,57	42,09	6,82	268,08	74,00	0,00	0,57	75,13	75,13	75,13
1990	A 070102	Passenger cars	Rural Gasoline	2,28	1148,38	500,38	13,79	4273,20	73,00	3,08	0,95	14,97	14,97	14,97
1990	A 070102	Passenger cars	Rural LPG	0,00	1248,46	305,18	16,91	1146,38	63,10	0,00	0,00	14,49	14,49	14,49
1990	A 070103	Passenger cars	Urban Diesel	93,68	220,85	103,04	6,82	344,97	74,00	0,00	0,32	144,24	144,24	144,24
1990	A 070103	Passenger cars	Urban Gasoline	2,28	558,41	1095,32	56,50	12514,03	73,00	2,71	0,54	12,37	12,37	12,37
1990	A 070103	Passenger cars	Urban LPG	0,00	528,01	473,05	20,80	1587,66	63,10	0,00	0,00	10,40	10,40	10,40
1990	A 070201	Light duty vehicles	Highway Diesel	93,68	270,67	30,19	2,60	344,14	74,00	0,00	0,32	104,48	104,48	104,48
1990	A 070201	Light duty vehicles	Highway Gasoline	2,28	1369,26	170,29	10,11	2987,40	73,00	2,63	0,81	16,17	16,17	16,17
1990	A 070201	Light duty vehicles	Highway LPG	0,00	1151,70	187,09	10,06	3914,25	63,10	0,00	0,00	10,06	10,06	10,06
1990	A 070202	Light duty vehicles	Rural Diesel	93,68	299,25	33,22	4,26	358,42	74,00	0,00	0,36	107,73	107,73	107,73
1990	A 070202	Light duty vehicles	Rural Gasoline	2,28	1188,86	262,59	15,25	2316,18	73,00	2,48	0,76	15,25	15,25	15,25
1990	A 070202	Light duty vehicles	Rural LPG	0,00	1248,46	305,18	16,91	1146,38	63,10	0,00	0,00	14,49	14,49	14,49
1990	A 070203	Light duty vehicles	Urban Diesel	93,68	469,24	74,81	4,58	463,37	74,00	0,00	0,23	165,76	165,76	165,76
1990	A 070203	Light duty vehicles	Urban Gasoline	2,28	547,55	874,08	39,84	9527,46	73,00	1,84	0,37	7,37	7,37	7,37
1990	A 070203	Light duty vehicles	Urban LPG	0,00	487,91	487,73	19,58	1705,63	63,10	0,00	0,00	9,79	9,79	9,79
1990	A 070301	Heavy duty vehicles	Highway Diesel	93,68	978,47	41,71	6,44	200,02	74,00	3,08	0,31	35,10	35,10	35,10
1990	A 070301	Heavy duty vehicles	Highway Gasoline	2,28	1037,78	474,61	9,69	7610,35	73,00	0,83	0,28	55,35	55,35	55,35
1990	A 070302	Heavy duty vehicles	Rural Diesel	93,68	984,85	57,74	6,78	211,25	74,00	2,92	0,29	35,73	35,73	35,73
1990	A 070302	Heavy duty vehicles	Rural Gasoline	2,28	1141,55	820,40	16,74	8371,39	73,00	0,91	0,30	60,88	60,88	60,88
1990	A 070303	Heavy duty vehicles	Urban Diesel	93,68	966,99	86,86	12,42	269,91	74,00	2,30	0,23	42,21	42,21	42,21
1990	A 070303	Heavy duty vehicles	Urban Gasoline	2,28	456,62	696,09	14,21	7102,99	73,00	0,61	0,20	40,59	40,59	40,59
1990	A 070400	Mopeds	Urban Gasoline	2,28	51,14	7471,23	200,00	13424,66	73,00	0,91	0,91	160,73	160,73	160,73
1990	A 070501	Motorcycles	Highway Gasoline	2,28	264,11	1072,19	129,96	16302,60	73,00	1,35	1,35	31,73	31,73	31,73
1990	A 070502	Motorcycles	Rural Gasoline	2,28	185,41	981,69	159,32	15782,07	73,00	1,66	1,66	38,90	38,90	38,90
1990	A 070503	Motorcycles	Urban Gasoline	2,28	112,92	1149,21	155,11	15187,59	73,00	1,61	1,61	37,87	37,87	37,87
1990	A 080100	Military	AvGas	22,83	859,00	1242,60	21,90	6972,00	73,00	2,00	1,60	10,00	10,00	10,00
1990	A 080100	Military	Diesel	93,68	719,72	55,39	6,74	268,73	74,00	1,73	0,32	66,49	66,49	66,49
1990	A 080100	Military	Gasoline	2,28	992,06	1114,62	27,57	6767,54	73,00	2,85	0,79	14,31	14,31	14,31
1990	A 080100	Military	Jet fuel	22,99	250,57	24,94	2,65	229,89	72,00	2,30	0,00	1,16	1,16	1,16
1990	A 080200	Railways	Diesel	93,68	1225,13	79,94	3,07	223,21	74,00	2,04	0,20	50,26	50,26	50,26
1990	A 080200	Railways	Kerosene	5,00	50,00	3,00	7,00	20,00	72,00	2,00	0,00	121,95	115,85	110,06

1990	A	080300	Inland waterways	Diesel	93,68	983,64	171,79	2,79	453,65	74,00	2,96	0,17	106,93	106,93	106,93
1990	A	080300	Inland waterways	Gasoline	2,28	291,33	3606,55	50,38	13853,27	73,00	0,78	0,08	182,44	182,44	182,44
1990	A	080402	National sea traffic	Diesel	93,68	1104,18	50,57	1,56	166,83	74,00	4,68	0,00	23,21	22,98	22,87
1990	A	080402	National sea traffic	Kerosene	2,30	50,00	3,00	7,00	20,00	72,00	0,00	0,00	5,00	5,00	5,00
1990	A	080402	National sea traffic	LPG	0,00	1249,00	384,94	20,26	443,00	63,10	0,00	0,00	0,20	0,20	0,20
1990	A	080402	National sea traffic	Residual oil	1290,95	1615,26	53,44	1,65	176,29	78,00	4,89	0,00	149,25	147,75	147,01
1990	A	080403	Fishing	Diesel	93,68	1052,12	49,13	1,52	162,08	74,00	4,68	0,00	23,21	22,98	22,87
1990	A	080403	Fishing	Kerosene	2,30	50,00	3,00	7,00	20,00	72,00	0,00	0,00	5,00	5,00	5,00
1990	A	080403	Fishing	LPG	0,00	1249,00	384,94	20,26	443,00	63,10	0,00	0,00	0,20	0,20	0,20
1990	A	080404	International sea traffic	Diesel	93,68	1208,60	49,46	1,53	163,17	74,00	4,68	0,00	23,21	22,98	22,87
1990	A	080404	International sea traffic	Residual oil	1447,43	1689,57	53,98	1,67	178,09	78,00	4,89	0,00	189,43	187,53	186,59
1990	A	080501	Air traffic, Dom. < 3000 ft.	AvGas	22,83	859,00	1242,60	21,90	6972,00	73,00	2,00	1,60	10,00	10,00	10,00
1990	A	080501	Air traffic, Dom. < 3000 ft.	Jet fuel	22,99	314,51	14,86	1,65	90,41	72,00	5,70	0,00	1,16	1,16	1,16
1990	A	080502	Air traffic, Int. < 3000 ft.	AvGas	22,83	859,00	1242,60	21,90	6972,00	73,00	2,00	1,60	10,00	10,00	10,00
1990	A	080502	Air traffic, Int. < 3000 ft.	Jet fuel	22,99	309,25	16,40	1,82	168,98	72,00	7,10	0,00	1,16	1,16	1,16
1990	A	080503	Air traffic, Dom. > 3000 ft.	Jet fuel	22,99	330,11	13,67	0,00	90,75	72,00	2,30	0,00	1,16	1,16	1,16
1990	A	080504	Air traffic, Int. > 3000 ft.	Jet fuel	22,99	244,20	7,17	0,00	54,10	72,00	2,30	0,00	1,16	1,16	1,16
1990	A	080600	Agriculture	Diesel	93,68	758,87	156,85	2,55	635,53	74,00	2,93	0,17	144,45	144,45	144,45
1990	A	080600	Agriculture	Gasoline	2,28	31,60	949,55	88,42	47524,17	73,00	1,28	0,09	6,56	6,56	6,56
1990	A	080700	Forestry	Diesel	93,68	857,48	156,47	2,54	645,65	74,00	2,97	0,17	149,05	149,05	149,05
1990	A	080700	Forestry	Gasoline	2,28	40,39	7206,91	60,42	18057,40	73,00	0,37	0,07	101,22	101,22	101,22
1990	A	080800	Industry	Diesel	93,68	933,58	178,23	2,90	655,80	74,00	2,94	0,17	154,50	154,50	154,50
1990	A	080800	Industry	Gasoline	2,28	136,27	1610,77	120,61	14797,46	73,00	1,33	0,09	12,40	12,40	12,40
1990	A	080800	Industry	LPG	0,00	1328,11	146,09	7,69	104,85	63,10	3,50	0,21	4,89	4,89	4,89
1990	A	080900	Household and gardening	Gasoline	2,28	63,98	3366,01	95,22	32901,19	73,00	1,15	0,08	20,75	20,75	20,75
1990	A	081100	Commercial and institutional	Gasoline	2,28	68,83	2280,66	97,87	29887,31	73,00	1,09	0,08	24,00	24,00	24,00
1990	P	080501	Air traffic, Dom. < 3000 ft.	AvGas	22,83	859,00	1242,60	21,90	6972,00	73,00	2,00	1,60	10,00	10,00	10,00
1990	P	080501	Air traffic, Dom. < 3000 ft.	Jet fuel	22,99	283,87	20,64	2,29	129,70	72,00	4,58	0,00	1,16	1,16	1,16
1990	P	080502	Air traffic, Int. < 3000 ft.	AvGas	22,83	859,00	1242,60	21,90	6972,00	73,00	2,00	1,60	10,00	10,00	10,00
1990	P	080502	Air traffic, Int. < 3000 ft.	Jet fuel	22,99	324,87	34,10	3,79	157,15	72,00	3,79	0,00	1,16	1,16	1,16
1990	P	080503	Air traffic, Dom. > 3000 ft.	Jet fuel	22,99	314,86	13,03	0,00	84,05	72,00	2,30	0,00	1,16	1,16	1,16
1990	P	080504	Air traffic, Int. > 3000 ft.	Jet fuel	22,99	290,20	11,15	0,00	37,65	72,00	2,30	0,00	1,16	1,16	1,16

1990 emission factors for Arsenic, Cadmium, Chromium, Copper, Mercury, Nickel, Lead, Selenium and Zinc.

Year	SNAP ID	Category	Fuel type	Arsenic mg pr GJ	Cadmium mg pr GJ	Chromium mg pr GJ	Copper mg pr GJ	Mercury mg pr GJ	Nickel mg pr GJ	Lead mg pr GJ	Selenium mg pr GJ	Zinc mg pr GJ
1990	A 070101	Passenger cars	Highway Diesel	0,002	0,292	0,926	0,657	0,124	0,296	1,757	0,002	58,606
1990	A 070101	Passenger cars	Highway Gasoline	0,007	0,273	0,386	1,043	0,199	0,321	1472,004	0,005	54,458
1990	A 070101	Passenger cars	Highway LPG	0,000	0,300	0,270	1,052	0,000	0,300	0,901	0,000	60,097
1990	A 070102	Passenger cars	Rural Diesel	0,002	0,355	1,083	0,770	0,124	0,358	2,133	0,002	71,144
1990	A 070102	Passenger cars	Rural Gasoline	0,007	0,308	0,417	1,165	0,199	0,356	1472,109	0,005	61,472
1990	A 070102	Passenger cars	Rural LPG	0,000	0,361	0,325	1,262	0,000	0,361	1,082	0,000	72,116
1990	A 070103	Passenger cars	Urban Diesel	0,002	0,198	0,691	0,488	0,124	0,201	1,192	0,002	39,770
1990	A 070103	Passenger cars	Urban Gasoline	0,007	0,178	0,300	0,711	0,199	0,226	1471,720	0,005	35,492
1990	A 070103	Passenger cars	Urban LPG	0,000	0,194	0,175	0,679	0,000	0,194	0,582	0,000	38,818
1990	A 070201	Light duty vehicles	Highway Diesel	0,002	0,193	0,678	0,478	0,124	0,196	1,161	0,002	38,731
1990	A 070201	Light duty vehicles	Highway Gasoline	0,007	0,253	0,367	0,972	0,199	0,301	1471,943	0,005	50,405
1990	A 070201	Light duty vehicles	Highway LPG	0,000	0,198	0,178	0,693	0,000	0,198	0,594	0,000	39,594
1990	A 070202	Light duty vehicles	Rural Diesel	0,002	0,211	0,723	0,511	0,124	0,214	1,269	0,002	42,337
1990	A 070202	Light duty vehicles	Rural Gasoline	0,007	0,239	0,355	0,922	0,199	0,287	1471,901	0,005	47,576
1990	A 070202	Light duty vehicles	Rural LPG	0,000	0,238	0,214	0,831	0,000	0,238	0,713	0,000	47,513
1990	A 070203	Light duty vehicles	Urban Diesel	0,002	0,135	0,533	0,374	0,124	0,138	0,813	0,002	27,133
1990	A 070203	Light duty vehicles	Urban Gasoline	0,007	0,118	0,246	0,499	0,199	0,166	1471,538	0,005	23,388
1990	A 070203	Light duty vehicles	Urban LPG	0,000	0,120	0,108	0,421	0,000	0,120	0,361	0,000	24,068
1990	A 070301	Heavy duty vehicles	Highway Diesel	0,002	0,150	0,570	0,401	0,124	0,153	0,903	0,002	30,130
1990	A 070301	Heavy duty vehicles	Highway Gasoline	0,007	0,207	0,326	0,813	0,199	0,255	1471,807	0,005	41,332
1990	A 070302	Heavy duty vehicles	Rural Diesel	0,002	0,143	0,555	0,390	0,124	0,147	0,866	0,002	28,884
1990	A 070302	Heavy duty vehicles	Rural Gasoline	0,007	0,220	0,338	0,857	0,199	0,268	1471,845	0,005	43,830
1990	A 070303	Heavy duty vehicles	Urban Diesel	0,002	0,115	0,483	0,338	0,124	0,118	0,693	0,002	23,142
1990	A 070303	Heavy duty vehicles	Urban Gasoline	0,007	0,143	0,268	0,586	0,199	0,190	1471,613	0,005	28,351
1990	A 070400	Mopeds	Urban Gasoline	0,007	0,005	0,144	0,103	0,199	0,053	1471,199	0,005	0,753
1990	A 070501	Motorcycles	Highway Gasoline	0,007	0,134	0,260	0,556	0,199	0,182	1471,587	0,005	26,666
1990	A 070502	Motorcycles	Rural Gasoline	0,007	0,163	0,287	0,659	0,199	0,211	1471,675	0,005	32,520
1990	A 070503	Motorcycles	Urban Gasoline	0,007	0,159	0,283	0,644	0,199	0,207	1471,663	0,005	31,681
1990	A 080100	Military	AvGas	0,007	0,253	0,367	0,972	0,198	0,301	12785,388	0,005	50,452
1990	A 080100	Military	Diesel	0,002	0,172	0,625	0,440	0,124	0,175	1,035	0,002	34,517
1990	A 080100	Military	Gasoline	0,007	0,256	0,371	0,984	0,199	0,304	1471,954	0,005	51,135
1990	A 080100	Military	Jet fuel	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
1990	A 080200	Railways	Diesel	0,002	0,186	0,660	0,465	0,124	0,189	1,118	0,002	37,295
1990	A 080200	Railways	Kerosene	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000

1990	A	080300	Inland waterways	Diesel	0,002	0,186	0,660	0,465	0,124	0,189	1,118	0,002	37,295
1990	A	080300	Inland waterways	Gasoline	0,007	0,253	0,367	0,972	0,198	0,301	0,773	0,005	50,452
1990	A	080402	National sea traffic	Diesel	1,171	0,234	0,937	1,171	1,170	1,639	2,340	4,684	11,710
1990	A	080402	National sea traffic	Kerosene	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
1990	A	080402	National sea traffic	LPG	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
1990	A	080402	National sea traffic	Residual oil	12,225	0,733	4,890	12,225	0,490	733,496	4,890	9,780	22,005
1990	A	080403	Fishing	Diesel	1,171	0,234	0,937	1,171	1,170	1,639	2,340	4,684	11,710
1990	A	080403	Fishing	Kerosene	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
1990	A	080403	Fishing	LPG	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
1990	A	080404	International sea traffic	Diesel	1,171	0,234	0,937	1,171	1,170	1,639	2,340	4,684	11,710
1990	A	080404	International sea traffic	Residual oil	12,225	0,733	4,890	12,225	0,490	733,496	4,890	9,780	22,005
1990	A	080501	Air traffic, Dom. < 3000 ft.	AvGas	0,007	0,253	0,367	0,972	0,198	0,301	13505,692	0,005	50,452
1990	A	080501	Air traffic, Dom. < 3000 ft.	Jet fuel	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
1990	A	080502	Air traffic, Int. < 3000 ft.	AvGas	0,007	0,253	0,367	0,972	0,198	0,301	13505,692	0,005	50,452
1990	A	080502	Air traffic, Int. < 3000 ft.	Jet fuel	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
1990	A	080503	Air traffic, Dom. > 3000 ft.	Jet fuel	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
1990	A	080504	Air traffic, Int. > 3000 ft.	Jet fuel	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
1990	A	080600	Agriculture	Diesel	0,002	0,186	0,660	0,465	0,124	0,189	1,118	0,002	37,295
1990	A	080600	Agriculture	Gasoline	0,007	0,253	0,367	0,972	0,198	0,301	0,773	0,005	50,452
1990	A	080700	Forestry	Diesel	0,002	0,186	0,660	0,465	0,124	0,189	1,118	0,002	37,295
1990	A	080700	Forestry	Gasoline	0,007	0,253	0,367	0,972	0,198	0,301	0,773	0,005	50,452
1990	A	080800	Industry	Diesel	0,002	0,186	0,660	0,465	0,124	0,189	1,118	0,002	37,295
1990	A	080800	Industry	Gasoline	0,007	0,253	0,367	0,972	0,198	0,301	0,773	0,005	50,452
1990	A	080800	Industry	LPG	0,000	0,131	0,118	0,457	0,000	0,131	0,392	0,000	26,126
1990	A	080900	Household and gardening	Gasoline	0,007	0,253	0,367	0,972	0,198	0,301	0,773	0,005	50,452
1990	A	081100	Commercial and institutional	Gasoline	0,007	0,253	0,367	0,972	0,198	0,301	0,773	0,005	50,452
1990	P	080501	Air traffic, Dom. < 3000 ft.	AvGas	0,007	0,253	0,367	0,972	0,198	0,301	13505,692	0,005	50,452
1990	P	080501	Air traffic, Dom. < 3000 ft.	Jet fuel	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
1990	P	080502	Air traffic, Int. < 3000 ft.	AvGas	0,007	0,253	0,367	0,972	0,198	0,301	13505,692	0,005	50,452
1990	P	080502	Air traffic, Int. < 3000 ft.	Jet fuel	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
1990	P	080503	Air traffic, Dom. > 3000 ft.	Jet fuel	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
1990	P	080504	Air traffic, Int. > 3000 ft.	Jet fuel	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000

1990 emission factors for Dioxins/, Flouranthene, Benzo(b), Benzo(k), Benzo(a), Benzo(g,h,i), indeno(1,2,3-c,d), HCB and PCB.

Year	SNAP ID	Category	Fuel type	Dioxins/ microg pr GJ	Flouranthene mg pr GJ	Benzo(b) mg pr GJ	Benzo(k) mg pr GJ	Benzo(a) mg pr GJ	Benzo(g,h,i) mg pr GJ	Indeno (1,2,3-c,d) mg pr GJ	HCB ng pr GJ	PCB ng pr GJ
1990	A 070101	Passenger cars	Highway Diesel	0,001	12,250	0,748	0,678	0,818	1,589	0,771	6,150	21,300
1990	A 070101	Passenger cars	Highway Gasoline	0,013	8,507	0,553	0,425	0,468	1,106	0,425	0,159	1029,731
1990	A 070101	Passenger cars	Highway LPG	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
1990	A 070102	Passenger cars	Rural Diesel	0,001	14,889	0,909	0,824	0,994	1,932	0,937	6,150	21,300
1990	A 070102	Passenger cars	Rural Gasoline	0,015	9,540	0,620	0,477	0,525	1,240	0,477	0,159	1029,731
1990	A 070102	Passenger cars	Rural LPG	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
1990	A 070103	Passenger cars	Urban Diesel	0,001	9,303	0,568	0,515	0,621	1,207	0,586	6,150	21,300
1990	A 070103	Passenger cars	Urban Gasoline	0,010	6,427	0,418	0,321	0,354	0,835	0,321	0,159	1029,731
1990	A 070103	Passenger cars	Urban LPG	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
1990	A 070201	Light duty vehicles	Highway Diesel	0,000	8,505	0,519	0,470	0,568	1,104	0,536	6,150	21,300
1990	A 070201	Light duty vehicles	Highway Gasoline	0,013	8,086	0,526	0,404	0,445	1,051	0,404	0,159	1029,731
1990	A 070201	Light duty vehicles	Highway LPG	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
1990	A 070202	Light duty vehicles	Rural Diesel	0,001	9,306	0,568	0,515	0,622	1,207	0,586	6,150	21,300
1990	A 070202	Light duty vehicles	Rural Gasoline	0,012	7,625	0,495	0,381	0,419	0,991	0,381	0,159	1029,731
1990	A 070202	Light duty vehicles	Rural LPG	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
1990	A 070203	Light duty vehicles	Urban Diesel	0,000	6,954	0,425	0,385	0,464	0,902	0,438	6,150	21,300
1990	A 070203	Light duty vehicles	Urban Gasoline	0,007	4,558	0,296	0,228	0,251	0,592	0,228	0,159	1029,731
1990	A 070203	Light duty vehicles	Urban LPG	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
1990	A 070301	Heavy duty vehicles	Highway Diesel	0,001	2,086	0,526	0,780	0,097	0,078	0,136	6,150	526,000
1990	A 070301	Heavy duty vehicles	Highway Gasoline	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,159	1029,731
1990	A 070302	Heavy duty vehicles	Rural Diesel	0,001	2,208	0,557	0,825	0,103	0,082	0,144	6,150	526,000
1990	A 070302	Heavy duty vehicles	Rural Gasoline	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,159	1029,731
1990	A 070303	Heavy duty vehicles	Urban Diesel	0,001	1,788	0,451	0,668	0,083	0,067	0,117	6,150	526,000
1990	A 070303	Heavy duty vehicles	Urban Gasoline	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,159	1029,731
1990	A 070400	Mopeds	Urban Gasoline	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,159	1029,731
1990	A 070501	Motorcycles	Highway Gasoline	0,020	12,673	0,824	0,634	0,697	1,647	0,634	0,159	1029,731
1990	A 070502	Motorcycles	Rural Gasoline	0,024	15,176	0,986	0,759	0,834	1,973	0,759	0,159	1029,731
1990	A 070503	Motorcycles	Urban Gasoline	0,024	15,300	0,994	0,765	0,841	1,989	0,765	0,159	1029,731
1990	A 080100	Military	AvGas	0,005	4,329	0,209	0,071	0,114	0,689	0,245	0,000	0,000
1990	A 080100	Military	Diesel	0,001	4,391	0,571	0,568	0,290	0,550	0,290	6,150	330,743
1990	A 080100	Military	Gasoline	0,006	5,257	0,277	0,116	0,142	0,825	0,300	0,159	1029,731
1990	A 080100	Military	Jet fuel	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
1990	A 080200	Railways	Diesel	0,001	1,366	0,348	0,389	0,057	0,049	0,089	6,150	526,000
1990	A 080200	Railways	Kerosene	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000

1990	A	080300	Inland waterways	Diesel	0,001	4,391	0,571	0,568	0,290	0,550	0,290	6,150	526,000
1990	A	080300	Inland waterways	Gasoline	0,005	4,329	0,209	0,071	0,114	0,689	0,245	0,159	1029,731
1990	A	080402	National sea traffic	Diesel	0,012	7,420	0,640	0,300	0,150	1,430	1,180	1,950	8,760
1990	A	080402	National sea traffic	Kerosene	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
1990	A	080402	National sea traffic	LPG	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
1990	A	080402	National sea traffic	Residual oil	0,013	5,190	0,270	0,050	0,020	0,070	0,030	3,500	14,000
1990	A	080403	Fishing	Diesel	0,012	7,420	0,640	0,300	0,150	1,430	1,180	1,950	8,760
1990	A	080403	Fishing	Kerosene	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
1990	A	080403	Fishing	LPG	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
1990	A	080404	International sea traffic	Diesel	0,012	7,420	0,640	0,300	0,150	1,430	1,180	1,950	8,760
1990	A	080404	International sea traffic	Residual oil	0,013	4,120	0,200	0,090	0,070	0,260	0,200	3,500	14,000
1990	A	080501	Air traffic, Dom. < 3000 ft.	AvGas	0,005	4,329	0,209	0,071	0,114	0,689	0,245	0,000	0,000
1990	A	080501	Air traffic, Dom. < 3000 ft.	Jet fuel	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
1990	A	080502	Air traffic, Int. < 3000 ft.	AvGas	0,005	4,329	0,209	0,071	0,114	0,689	0,245	0,000	0,000
1990	A	080502	Air traffic, Int. < 3000 ft.	Jet fuel	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
1990	A	080503	Air traffic, Dom. > 3000 ft.	Jet fuel	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
1990	A	080504	Air traffic, Int. > 3000 ft.	Jet fuel	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
1990	A	080600	Agriculture	Diesel	0,001	4,391	0,571	0,568	0,290	0,550	0,290	6,150	526,000
1990	A	080600	Agriculture	Gasoline	0,005	4,329	0,209	0,071	0,114	0,689	0,245	0,159	1029,731
1990	A	080700	Forestry	Diesel	0,001	4,391	0,571	0,568	0,290	0,550	0,290	6,150	526,000
1990	A	080700	Forestry	Gasoline	0,005	4,329	0,209	0,071	0,114	0,689	0,245	0,159	1029,731
1990	A	080800	Industry	Diesel	0,001	4,391	0,571	0,568	0,290	0,550	0,290	6,150	526,000
1990	A	080800	Industry	Gasoline	0,005	4,329	0,209	0,071	0,114	0,689	0,245	0,159	1029,731
1990	A	080800	Industry	LPG	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
1990	A	080900	Household and gardening	Gasoline	0,005	4,329	0,209	0,071	0,114	0,689	0,245	0,159	1029,731
1990	A	081100	Commercial and institutional	Gasoline	0,005	4,329	0,209	0,071	0,114	0,689	0,245	0,159	1029,731
1990	P	080501	Air traffic, Dom. < 3000 ft.	AvGas	0,005	4,329	0,209	0,071	0,114	0,689	0,245	0,000	0,000
1990	P	080501	Air traffic, Dom. < 3000 ft.	Jet fuel	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
1990	P	080502	Air traffic, Int. < 3000 ft.	AvGas	0,005	4,329	0,209	0,071	0,114	0,689	0,245	0,000	0,000
1990	P	080502	Air traffic, Int. < 3000 ft.	Jet fuel	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
1990	P	080503	Air traffic, Dom. > 3000 ft.	Jet fuel	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
1990	P	080504	Air traffic, Int. > 3000 ft.	Jet fuel	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000

2012 emission factors for SO₂, NO_x, NMVOC, CH₄, CO, CO₂, N₂O, NH₃, TSP, PM₁₀ and PM_{2.5}.

Year	SNAP ID	Category	Fuel type	SO ₂ g pr GJ	NO _x g pr GJ	NMVOC g pr GJ	CH ₄ g pr GJ	CO g pr GJ	CO ₂ kg pr GJ	N ₂ O g pr GJ	NH ₃ g pr GJ	TSP g pr GJ	PM ₁₀ g pr GJ	PM _{2.5} g pr GJ		
2012	A	070101	Passenger cars	Highway	Bio ethanol	0,00	133,00	31,74	3,63	695,05	0,00	0,74	32,49	1,02	1,02	1,02
2012	A	070101	Passenger cars	Highway	Biodiesel	0,00	345,09	4,01	0,11	12,92	0,00	2,07	0,51	12,10	12,10	12,10
2012	A	070101	Passenger cars	Highway	Diesel	0,47	345,09	4,01	0,11	12,92	74,00	2,07	0,51	12,10	12,10	12,10
2012	A	070101	Passenger cars	Highway	Gasoline	0,46	133,00	31,74	3,63	695,05	73,00	0,74	32,49	1,02	1,02	1,02
2012	A	070101	Passenger cars	Highway	LPG	0,00	245,42	39,01	4,09	1401,42	63,10	0,78	0,00	10,05	10,05	10,05
2012	A	070102	Passenger cars	Rural	Bio ethanol	0,00	110,36	36,67	3,50	555,44	0,00	1,37	33,65	0,97	0,97	0,97
2012	A	070102	Passenger cars	Rural	Biodiesel	0,00	288,67	5,25	0,21	24,66	0,00	2,22	0,54	10,09	10,09	10,09
2012	A	070102	Passenger cars	Rural	Diesel	0,47	288,67	5,25	0,21	24,66	74,00	2,22	0,54	10,09	10,09	10,09
2012	A	070102	Passenger cars	Rural	Gasoline	0,46	110,36	36,67	3,50	555,44	73,00	1,37	33,65	0,97	0,97	0,97
2012	A	070102	Passenger cars	Rural	LPG	0,00	267,71	59,24	6,87	542,88	63,10	1,57	0,00	14,45	14,45	14,45
2012	A	070103	Passenger cars	Urban	Bio ethanol	0,00	138,64	295,15	11,54	3243,70	0,00	2,06	7,68	0,88	0,88	0,88
2012	A	070103	Passenger cars	Urban	Biodiesel	0,00	290,68	16,07	0,52	71,75	0,00	5,49	0,37	18,31	18,31	18,31
2012	A	070103	Passenger cars	Urban	Diesel	0,47	290,68	16,07	0,52	71,75	74,00	5,49	0,37	18,31	18,31	18,31
2012	A	070103	Passenger cars	Urban	Gasoline	0,46	138,64	295,15	11,54	3243,70	73,00	2,06	7,68	0,88	0,88	0,88
2012	A	070103	Passenger cars	Urban	LPG	0,00	134,04	138,14	9,48	937,64	63,10	3,74	0,00	11,64	11,64	11,64
2012	A	070201	Light duty vehicles	Highway	Bio ethanol	0,00	164,40	19,61	2,76	555,42	0,00	1,88	22,93	1,37	1,37	1,37
2012	A	070201	Light duty vehicles	Highway	Biodiesel	0,00	283,25	18,22	0,18	122,71	0,00	1,51	0,37	18,27	18,27	18,27
2012	A	070201	Light duty vehicles	Highway	Diesel	0,47	283,25	18,22	0,18	122,71	74,00	1,51	0,37	18,27	18,27	18,27
2012	A	070201	Light duty vehicles	Highway	Gasoline	0,46	164,40	19,61	2,76	555,42	73,00	1,88	22,93	1,37	1,37	1,37
2012	A	070201	Light duty vehicles	Highway	LPG	0,00	136,68	20,73	2,25	1046,75	63,10	0,41	0,00	10,04	10,04	10,04
2012	A	070202	Light duty vehicles	Rural	Bio ethanol	0,00	143,84	28,80	3,00	420,31	0,00	3,01	22,65	1,22	1,22	1,22
2012	A	070202	Light duty vehicles	Rural	Biodiesel	0,00	295,92	20,62	0,41	105,18	0,00	1,65	0,40	14,95	14,95	14,95
2012	A	070202	Light duty vehicles	Rural	Diesel	0,47	295,92	20,62	0,41	105,18	74,00	1,65	0,40	14,95	14,95	14,95
2012	A	070202	Light duty vehicles	Rural	Gasoline	0,46	143,84	28,80	3,00	420,31	73,00	3,01	22,65	1,22	1,22	1,22
2012	A	070202	Light duty vehicles	Rural	LPG	0,00	149,37	31,44	3,78	435,27	63,10	0,93	0,00	14,45	14,45	14,45
2012	A	070203	Light duty vehicles	Urban	Bio ethanol	0,00	130,10	200,38	8,08	3760,68	0,00	4,56	4,44	0,74	0,74	0,74
2012	A	070203	Light duty vehicles	Urban	Biodiesel	0,00	275,24	42,60	0,81	141,68	0,00	3,51	0,26	25,22	25,22	25,22
2012	A	070203	Light duty vehicles	Urban	Diesel	0,47	275,24	42,60	0,81	141,68	74,00	3,51	0,26	25,22	25,22	25,22
2012	A	070203	Light duty vehicles	Urban	Gasoline	0,46	130,10	200,38	8,08	3760,68	73,00	4,56	4,44	0,74	0,74	0,74
2012	A	070203	Light duty vehicles	Urban	LPG	0,00	78,75	73,39	5,36	563,91	63,10	1,97	0,00	11,94	11,94	11,94
2012	A	070301	Heavy duty vehicles	Highway	Bio ethanol	0,00	1037,78	474,61	9,69	7610,35	0,00	0,83	0,28	55,35	55,35	55,35
2012	A	070301	Heavy duty vehicles	Highway	Biodiesel	0,00	392,17	7,43	2,04	136,80	0,00	3,13	0,31	6,18	6,18	6,18
2012	A	070301	Heavy duty vehicles	Highway	Diesel	0,47	392,17	7,43	2,04	136,80	74,00	3,13	0,31	6,18	6,18	6,18
2012	A	070301	Heavy duty vehicles	Highway	Gasoline	0,46	1037,78	474,61	9,69	7610,35	73,00	0,83	0,28	55,35	55,35	55,35

2012	A	070302	Heavy duty vehicles	Rural	Bio ethanol	0,00	1141,55	820,40	16,74	8371,39	0,00	0,91	0,30	60,88	60,88	60,88
2012	A	070302	Heavy duty vehicles	Rural	Biodiesel	0,00	454,93	10,02	2,34	134,81	0,00	2,95	0,29	6,64	6,64	6,64
2012	A	070302	Heavy duty vehicles	Rural	Diesel	0,47	454,93	10,02	2,34	134,81	74,00	2,95	0,29	6,64	6,64	6,64
2012	A	070302	Heavy duty vehicles	Rural	Gasoline	0,46	1141,55	820,40	16,74	8371,39	73,00	0,91	0,30	60,88	60,88	60,88
2012	A	070303	Heavy duty vehicles	Urban	Bio ethanol	0,00	456,62	696,09	14,21	7102,99	0,00	0,61	0,20	40,59	40,59	40,59
2012	A	070303	Heavy duty vehicles	Urban	Biodiesel	0,00	570,34	14,75	3,73	149,48	0,00	2,50	0,25	8,10	8,10	8,10
2012	A	070303	Heavy duty vehicles	Urban	Diesel	0,47	570,34	14,75	3,73	149,48	74,00	2,50	0,25	8,10	8,10	8,10
2012	A	070303	Heavy duty vehicles	Urban	Gasoline	0,46	456,62	696,09	14,21	7102,99	73,00	0,61	0,20	40,59	40,59	40,59
2012	A	070400	Mopeds	Urban	Bio ethanol	0,00	145,74	4413,69	91,50	7300,73	0,00	1,06	1,06	76,35	76,35	76,35
2012	A	070400	Mopeds	Urban	Gasoline	0,46	145,74	4413,69	91,50	7300,73	73,00	1,06	1,06	76,35	76,35	76,35
2012	A	070501	Motorcycles	Highway	Bio ethanol	0,00	270,47	651,06	90,62	10392,91	0,00	1,27	1,27	15,72	15,72	15,72
2012	A	070501	Motorcycles	Highway	Gasoline	0,46	270,47	651,06	90,62	10392,91	73,00	1,27	1,27	15,72	15,72	15,72
2012	A	070502	Motorcycles	Rural	Bio ethanol	0,00	192,46	650,59	109,49	9619,73	0,00	1,55	1,55	19,14	19,14	19,14
2012	A	070502	Motorcycles	Rural	Gasoline	0,46	192,46	650,59	109,49	9619,73	73,00	1,55	1,55	19,14	19,14	19,14
2012	A	070503	Motorcycles	Urban	Bio ethanol	0,00	118,83	802,17	115,50	9267,95	0,00	1,51	1,51	18,65	18,65	18,65
2012	A	070503	Motorcycles	Urban	Gasoline	0,46	118,83	802,17	115,50	9267,95	73,00	1,51	1,51	18,65	18,65	18,65
2012	A	080100	Military		AvGas	22,99	859,00	1242,60	21,90	6972,00	73,00	2,00	1,60	10,00	10,00	10,00
2012	A	080100	Military		Diesel	0,44	356,14	12,51	1,14	92,27	74,00	2,81	0,38	11,76	11,76	11,76
2012	A	080100	Military		Gasoline	0,44	127,06	160,98	7,39	1522,54	73,00	1,46	24,95	1,51	1,51	1,51
2012	A	080100	Military		Jet fuel	22,99	250,57	24,94	2,65	229,89	72,00	2,30	0,00	1,16	1,16	1,16
2012	A	080200	Railways		Diesel	0,47	751,17	56,43	2,17	125,99	74,00	2,04	0,20	24,24	24,24	24,24
2012	A	080200	Railways		Gasoline											
2012	A	080200	Railways		Kerosene											
2012	A	080300	Inland waterways		Diesel	46,84	817,03	156,52	2,55	439,34	74,00	2,97	0,17	95,54	95,54	95,54
2012	A	080300	Inland waterways		Gasoline	0,46	567,23	877,36	64,41	11258,30	73,00	1,55	0,10	17,43	17,43	17,43
2012	A	080402	National sea traffic		Diesel	46,84	1081,18	50,38	1,66	111,43	74,00	4,68	0,00	21,55	21,33	21,22
2012	A	080402	National sea traffic		Kerosene											
2012	A	080402	National sea traffic		LPG	0,00	1249,00	384,94	20,26	443,00	63,10	0,00	0,00	0,20	0,20	0,20
2012	A	080402	National sea traffic		Residual oil	489,00	1925,97	63,12	1,95	208,23	78,00	4,89	0,00	43,98	43,54	43,32
2012	A	080403	Fishing		Diesel	46,84	1340,90	57,72	1,79	190,43	74,00	4,68	0,00	21,55	21,33	21,22
2012	A	080403	Fishing		Gasoline											
2012	A	080403	Fishing		Kerosene											
2012	A	080403	Fishing		LPG	0,00	1249,00	384,94	20,26	443,00	63,10	0,00	0,00	0,20	0,20	0,20
2012	A	080404	International sea traffic		Diesel	46,84	1585,62	57,71	1,78	190,37	74,00	4,68	0,00	21,55	21,33	21,22
2012	A	080404	International sea traffic		Residual oil	489,00	2119,92	63,46	1,96	209,36	78,00	4,89	0,00	43,98	43,54	43,32
2012	A	080501	Air traffic, Dom. < 3000 ft.		AvGas	22,83	859,00	1242,60	21,90	6972,00	73,00	2,00	1,60	10,00	10,00	10,00
2012	A	080501	Air traffic, Dom. < 3000 ft.		Jet fuel	22,99	309,39	16,41	1,82	147,19	72,00	9,24	0,00	1,16	1,16	1,16

2012	A	080502	Air traffic, Int. < 3000 ft.	AvGas	22,83	859,00	1242,60	21,90	6972,00	73,00	2,00	1,60	10,00	10,00	10,00
2012	A	080502	Air traffic, Int. < 3000 ft.	Jet fuel	22,99	298,57	38,17	4,24	184,54	72,00	7,24	0,00	1,16	1,16	1,16
2012	A	080503	Air traffic, Dom. > 3000 ft.	Jet fuel	22,99	278,02	6,83	0,00	95,20	72,00	2,30	0,00	1,16	1,16	1,16
2012	A	080504	Air traffic, Int. > 3000 ft.	Jet fuel	22,99	238,74	6,01	0,00	48,19	72,00	2,30	0,00	1,16	1,16	1,16
2012	A	080600	Agriculture	Diesel	0,47	530,77	48,98	0,80	308,69	74,00	3,19	0,18	39,27	39,27	39,27
2012	A	080600	Agriculture	Gasoline	0,46	110,77	1184,09	159,58	21731,05	73,00	1,71	1,51	31,03	31,03	31,03
2012	A	080700	Forestry	Diesel	0,47	347,95	25,87	0,42	226,21	74,00	3,21	0,18	23,40	23,40	23,40
2012	A	080700	Forestry	Gasoline	0,46	54,79	3964,24	30,97	17915,98	73,00	0,46	0,09	82,19	82,19	82,19
2012	A	080800	Industry	Diesel	0,47	489,22	54,19	0,88	306,75	74,00	3,10	0,18	47,52	47,52	47,52
2012	A	080800	Industry	Gasoline	0,46	212,00	1528,15	108,83	14105,02	73,00	1,49	0,10	20,45	20,45	20,45
2012	A	080800	Industry	LPG	0,00	1328,11	146,09	7,69	104,85	63,10	3,50	0,21	4,89	4,89	4,89
2012	A	080900	Household and gardening	Gasoline	0,46	107,75	2283,84	75,45	30678,27	73,00	1,26	0,09	17,21	17,21	17,21
2012	A	081100	Commercial and institutional	Gasoline	0,46	93,28	1548,54	64,42	30913,57	73,00	1,13	0,09	28,53	28,53	28,53
2012	P	080501	Air traffic, Dom. < 3000 ft.	AvGas	22,83	859,00	1242,60	21,90	6972,00	73,00	2,00	1,60	10,00	10,00	10,00
2012	P	080501	Air traffic, Dom. < 3000 ft.	Jet fuel	22,99	302,09	18,59	2,07	191,64	72,00	5,66	0,00	1,16	1,16	1,16
2012	P	080502	Air traffic, Int. < 3000 ft.	AvGas	22,83	859,00	1242,60	21,90	6972,00	73,00	2,00	1,60	10,00	10,00	10,00
2012	P	080502	Air traffic, Int. < 3000 ft.	Jet fuel	22,99	339,36	33,80	3,76	237,36	72,00	3,87	0,00	1,16	1,16	1,16
2012	P	080503	Air traffic, Dom. > 3000 ft.	Jet fuel	22,99	299,41	9,63	0,00	50,21	72,00	2,30	0,00	1,16	1,16	1,16
2012	P	080504	Air traffic, Int. > 3000 ft.	Jet fuel	22,99	309,94	8,90	0,00	31,29	72,00	2,30	0,00	1,16	1,16	1,16

2012 emission factors for Arsenic, Cadmium, Chromium, Copper, Mercury, Nickel, Lead, Selenium and Zinc.

Year	SNAP ID	Category	Fuel type	Arsenic mg pr GJ	Cadmium mg pr GJ	Chromium mg pr GJ	Copper mg pr GJ	Mercury mg pr GJ	Nickel mg pr GJ	Lead mg pr GJ	Selenium mg pr GJ	Zinc mg pr GJ
2012	A 070101	Passenger cars	Highway Bio ethanol	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
2012	A 070101	Passenger cars	Highway Biodiesel	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
2012	A 070101	Passenger cars	Highway Diesel	0,002	0,312	0,977	0,694	0,124	0,316	1,880	0,002	62,685
2012	A 070101	Passenger cars	Highway Gasoline	0,007	0,332	0,438	1,248	0,199	0,380	1,009	0,005	66,216
2012	A 070101	Passenger cars	Highway LPG	0,000	0,294	0,265	1,030	0,000	0,294	0,883	0,000	58,875
2012	A 070102	Passenger cars	Rural Bio ethanol	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
2012	A 070102	Passenger cars	Rural Biodiesel	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
2012	A 070102	Passenger cars	Rural Diesel	0,002	0,335	1,035	0,735	0,124	0,339	2,017	0,002	67,260
2012	A 070102	Passenger cars	Rural Gasoline	0,007	0,351	0,456	1,316	0,199	0,399	1,067	0,005	70,066
2012	A 070102	Passenger cars	Rural LPG	0,000	0,353	0,318	1,235	0,000	0,353	1,059	0,000	70,585
2012	A 070103	Passenger cars	Urban Bio ethanol	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
2012	A 070103	Passenger cars	Urban Biodiesel	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
2012	A 070103	Passenger cars	Urban Diesel	0,002	0,227	0,763	0,540	0,124	0,230	1,366	0,002	45,556
2012	A 070103	Passenger cars	Urban Gasoline	0,007	0,205	0,324	0,804	0,199	0,253	0,628	0,005	40,805
2012	A 070103	Passenger cars	Urban LPG	0,000	0,213	0,192	0,746	0,000	0,213	0,639	0,000	42,625
2012	A 070201	Light duty vehicles	Highway Bio ethanol	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
2012	A 070201	Light duty vehicles	Highway Biodiesel	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
2012	A 070201	Light duty vehicles	Highway Diesel	0,002	0,239	0,793	0,561	0,124	0,242	1,438	0,002	47,956
2012	A 070201	Light duty vehicles	Highway Gasoline	0,007	0,253	0,368	0,973	0,199	0,301	0,774	0,005	50,509
2012	A 070201	Light duty vehicles	Highway LPG	0,000	0,172	0,154	0,601	0,000	0,172	0,515	0,000	34,315
2012	A 070202	Light duty vehicles	Rural Bio ethanol	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
2012	A 070202	Light duty vehicles	Rural Biodiesel	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
2012	A 070202	Light duty vehicles	Rural Diesel	0,002	0,261	0,849	0,602	0,124	0,265	1,572	0,002	52,429
2012	A 070202	Light duty vehicles	Rural Gasoline	0,007	0,239	0,355	0,925	0,199	0,287	0,732	0,005	47,723
2012	A 070202	Light duty vehicles	Rural LPG	0,000	0,206	0,185	0,720	0,000	0,206	0,617	0,000	41,136
2012	A 070203	Light duty vehicles	Urban Bio ethanol	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
2012	A 070203	Light duty vehicles	Urban Biodiesel	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
2012	A 070203	Light duty vehicles	Urban Diesel	0,002	0,168	0,617	0,434	0,124	0,172	1,014	0,002	33,839
2012	A 070203	Light duty vehicles	Urban Gasoline	0,007	0,122	0,250	0,515	0,199	0,170	0,381	0,005	24,327
2012	A 070203	Light duty vehicles	Urban LPG	0,000	0,127	0,115	0,446	0,000	0,127	0,382	0,000	25,489
2012	A 070301	Heavy duty vehicles	Highway Bio ethanol	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
2012	A 070301	Heavy duty vehicles	Highway Biodiesel	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
2012	A 070301	Heavy duty vehicles	Highway Diesel	0,002	0,159	0,593	0,417	0,124	0,162	0,958	0,002	31,954
2012	A 070301	Heavy duty vehicles	Highway Gasoline	0,007	0,260	0,374	0,997	0,199	0,308	0,794	0,005	51,857

2012	A	070302	Heavy duty vehicles	Rural	Bio ethanol	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
2012	A	070302	Heavy duty vehicles	Rural	Biodiesel	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
2012	A	070302	Heavy duty vehicles	Rural	Diesel	0,002	0,152	0,577	0,406	0,124	0,156	0,920	0,002 30,681
2012	A	070302	Heavy duty vehicles	Rural	Gasoline	0,007	0,295	0,405	1,119	0,199	0,343	0,898	0,005 58,799
2012	A	070303	Heavy duty vehicles	Urban	Bio ethanol	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
2012	A	070303	Heavy duty vehicles	Urban	Biodiesel	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
2012	A	070303	Heavy duty vehicles	Urban	Diesel	0,002	0,132	0,527	0,369	0,124	0,136	0,798	0,002 26,630
2012	A	070303	Heavy duty vehicles	Urban	Gasoline	0,007	0,206	0,325	0,806	0,199	0,253	0,630	0,005 40,950
2012	A	070400	Mopeds	Urban	Bio ethanol	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
2012	A	070400	Mopeds	Urban	Gasoline	0,007	0,005	0,144	0,103	0,199	0,053	0,027	0,005 0,753
2012	A	070501	Motorcycles	Highway	Bio ethanol	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
2012	A	070501	Motorcycles	Highway	Gasoline	0,007	0,134	0,260	0,555	0,199	0,182	0,415	0,005 26,609
2012	A	070502	Motorcycles	Rural	Bio ethanol	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
2012	A	070502	Motorcycles	Rural	Gasoline	0,007	0,162	0,286	0,654	0,199	0,210	0,500	0,005 32,240
2012	A	070503	Motorcycles	Urban	Bio ethanol	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
2012	A	070503	Motorcycles	Urban	Gasoline	0,007	0,158	0,282	0,640	0,199	0,206	0,488	0,005 31,445
2012	A	080100	Military		AvGas	0,007	0,253	0,367	0,972	0,198	0,301	12785,390	0,005 50,452
2012	A	080100	Military		Diesel	0,002	0,214	0,718	0,508	0,116	0,217	1,287	0,002 42,931
2012	A	080100	Military		Gasoline	0,007	0,286	0,393	1,086	0,192	0,333	0,872	0,004 57,120
2012	A	080100	Military		Jet fuel	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
2012	A	080200	Railways		Diesel	0,002	0,186	0,660	0,465	0,124	0,189	1,118	0,002 37,295
2012	A	080200	Railways		Gasoline								
2012	A	080200	Railways		Kerosene								
2012	A	080300	Inland waterways		Diesel	0,002	0,186	0,660	0,465	0,124	0,189	1,118	0,002 37,295
2012	A	080300	Inland waterways		Gasoline	0,007	0,253	0,367	0,972	0,198	0,301	0,773	0,005 50,452
2012	A	080402	National sea traffic		Diesel	1,170	0,230	0,940	1,170	1,170	1,640	2,340	4,680 11,710
2012	A	080402	National sea traffic		Kerosene								
2012	A	080402	National sea traffic		LPG	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
2012	A	080402	National sea traffic		Residual oil	12,220	0,730	4,890	12,220	0,490	733,500	4,890	9,780 22,000
2012	A	080403	Fishing		Diesel	1,170	0,230	0,940	1,170	1,170	1,640	2,340	4,680 11,710
2012	A	080403	Fishing		Gasoline								
2012	A	080403	Fishing		Kerosene								
2012	A	080403	Fishing		LPG	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
2012	A	080404	International sea traffic		Diesel	1,170	0,230	0,940	1,170	1,170	1,640	2,340	4,680 11,710
2012	A	080404	International sea traffic		Residual oil	12,220	0,730	4,890	12,220	0,490	733,500	4,890	9,780 22,000
2012	A	080501	Air traffic, Dom. < 3000 ft.		AvGas	0,007	0,253	0,367	0,972	0,198	0,301	13505,692	0,005 50,452
2012	A	080501	Air traffic, Dom. < 3000 ft.		Jet fuel	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000

2012	A	080502	Air traffic, Int. < 3000 ft.	AvGas	0,007	0,253	0,367	0,972	0,198	0,301	13505,692	0,005	50,452
2012	A	080502	Air traffic, Int. < 3000 ft.	Jet fuel	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
2012	A	080503	Air traffic, Dom. > 3000 ft.	Jet fuel	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
2012	A	080504	Air traffic, Int. > 3000 ft.	Jet fuel	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
2012	A	080600	Agriculture	Diesel	0,002	0,186	0,660	0,465	0,124	0,189	1,118	0,002	37,295
2012	A	080600	Agriculture	Gasoline	0,007	0,253	0,367	0,972	0,198	0,301	0,773	0,005	50,452
2012	A	080700	Forestry	Diesel	0,002	0,186	0,660	0,465	0,124	0,189	1,118	0,002	37,295
2012	A	080700	Forestry	Gasoline	0,007	0,253	0,367	0,972	0,198	0,301	0,773	0,005	50,452
2012	A	080800	Industry	Diesel	0,002	0,186	0,660	0,465	0,124	0,189	1,118	0,002	37,295
2012	A	080800	Industry	Gasoline	0,007	0,253	0,367	0,972	0,198	0,301	0,773	0,005	50,452
2012	A	080800	Industry	LPG	0,000	0,131	0,118	0,457	0,000	0,131	0,392	0,000	26,126
2012	A	080900	Household and gardening	Gasoline	0,007	0,253	0,367	0,972	0,198	0,301	0,773	0,005	50,452
2012	A	081100	Commercial and institutional	Gasoline	0,007	0,253	0,367	0,972	0,198	0,301	0,773	0,005	50,452
2012	P	080501	Air traffic, Dom. < 3000 ft.	AvGas	0,007	0,253	0,367	0,972	0,198	0,301	13505,692	0,005	50,452
2012	P	080501	Air traffic, Dom. < 3000 ft.	Jet fuel	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
2012	P	080502	Air traffic, Int. < 3000 ft.	AvGas	0,007	0,253	0,367	0,972	0,198	0,301	13505,692	0,005	50,452
2012	P	080502	Air traffic, Int. < 3000 ft.	Jet fuel	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
2012	P	080503	Air traffic, Dom. > 3000 ft.	Jet fuel	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
2012	P	080504	Air traffic, Int. > 3000 ft.	Jet fuel	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000

2012 emission factors for Dioxins/, Flouranthene, Benzo(b), Benzo(k), Benzo(a), Benzo(g,h,i), indeno(1,2,3-c,d), HCB and PCB.

Year	SNAP ID	Category	Fuel type	Dioxins/ microg pr GJ	Flouranthene mg pr GJ	Benzo(b) mg pr GJ	Benzo(k) mg pr GJ	Benzo(a) mg pr GJ	Benzo(g,h,i) mg pr GJ	indeno(1,2,3-c,d) mg pr GJ	HCB ng pr GJ	PCB ng pr GJ
2012	A 070101	Passenger cars	Highway Bio ethanol	0,000	1,137	0,208	0,252	0,206	0,417	0,299	0,010	0,457
2012	A 070101	Passenger cars	Highway Biodiesel	0,000	12,815	0,782	0,709	0,856	1,663	0,807	6,150	21,300
2012	A 070101	Passenger cars	Highway Diesel	0,000	12,815	0,782	0,709	0,856	1,663	0,807	6,150	21,300
2012	A 070101	Passenger cars	Highway Gasoline	0,000	1,137	0,208	0,252	0,206	0,417	0,299	0,010	0,457
2012	A 070101	Passenger cars	Highway LPG	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
2012	A 070102	Passenger cars	Rural Bio ethanol	0,000	1,251	0,232	0,280	0,229	0,464	0,334	0,010	0,457
2012	A 070102	Passenger cars	Rural Biodiesel	0,001	14,593	0,891	0,807	0,975	1,894	0,919	6,150	21,300
2012	A 070102	Passenger cars	Rural Diesel	0,001	14,593	0,891	0,807	0,975	1,894	0,919	6,150	21,300
2012	A 070102	Passenger cars	Rural Gasoline	0,000	1,251	0,232	0,280	0,229	0,464	0,334	0,010	0,457
2012	A 070102	Passenger cars	Rural LPG	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
2012	A 070103	Passenger cars	Urban Bio ethanol	0,000	0,720	0,129	0,155	0,127	0,257	0,184	0,010	0,457
2012	A 070103	Passenger cars	Urban Biodiesel	0,001	9,684	0,591	0,536	0,647	1,257	0,610	6,150	21,300
2012	A 070103	Passenger cars	Urban Diesel	0,001	9,684	0,591	0,536	0,647	1,257	0,610	6,150	21,300
2012	A 070103	Passenger cars	Urban Gasoline	0,000	0,720	0,129	0,155	0,127	0,257	0,184	0,010	0,457
2012	A 070103	Passenger cars	Urban LPG	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
2012	A 070201	Light duty vehicles	Highway Bio ethanol	0,001	1,029	0,159	0,184	0,154	0,317	0,217	0,010	0,457
2012	A 070201	Light duty vehicles	Highway Biodiesel	0,001	9,234	0,564	0,511	0,617	1,198	0,581	6,150	21,300
2012	A 070201	Light duty vehicles	Highway Diesel	0,001	9,234	0,564	0,511	0,617	1,198	0,581	6,150	21,300
2012	A 070201	Light duty vehicles	Highway Gasoline	0,001	1,029	0,159	0,184	0,154	0,317	0,217	0,010	0,457
2012	A 070201	Light duty vehicles	Highway LPG	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
2012	A 070202	Light duty vehicles	Rural Bio ethanol	0,001	0,971	0,150	0,174	0,146	0,300	0,205	0,010	0,457
2012	A 070202	Light duty vehicles	Rural Biodiesel	0,001	10,103	0,617	0,559	0,675	1,311	0,636	6,150	21,300
2012	A 070202	Light duty vehicles	Rural Diesel	0,001	10,103	0,617	0,559	0,675	1,311	0,636	6,150	21,300
2012	A 070202	Light duty vehicles	Rural Gasoline	0,001	0,971	0,150	0,174	0,146	0,300	0,205	0,010	0,457
2012	A 070202	Light duty vehicles	Rural LPG	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
2012	A 070203	Light duty vehicles	Urban Bio ethanol	0,000	0,561	0,087	0,101	0,084	0,173	0,118	0,010	0,457
2012	A 070203	Light duty vehicles	Urban Biodiesel	0,000	7,261	0,443	0,402	0,485	0,942	0,457	6,150	21,300
2012	A 070203	Light duty vehicles	Urban Diesel	0,000	7,261	0,443	0,402	0,485	0,942	0,457	6,150	21,300
2012	A 070203	Light duty vehicles	Urban Gasoline	0,000	0,561	0,087	0,101	0,084	0,173	0,118	0,010	0,457
2012	A 070203	Light duty vehicles	Urban LPG	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
2012	A 070301	Heavy duty vehicles	Highway Bio ethanol	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,010	0,457
2012	A 070301	Heavy duty vehicles	Highway Biodiesel	0,001	2,030	0,512	0,759	0,095	0,076	0,133	6,150	526,000
2012	A 070301	Heavy duty vehicles	Highway Diesel	0,001	2,030	0,512	0,759	0,095	0,076	0,133	6,150	526,000
2012	A 070301	Heavy duty vehicles	Highway Gasoline	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,010	0,457

2012	A	070302	Heavy duty vehicles	Rural	Bio ethanol	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,010	0,457
2012	A	070302	Heavy duty vehicles	Rural	Biodiesel	0,001	2,066	0,521	0,772	0,096	0,077	0,135	6,150	526,000
2012	A	070302	Heavy duty vehicles	Rural	Diesel	0,001	2,066	0,521	0,772	0,096	0,077	0,135	6,150	526,000
2012	A	070302	Heavy duty vehicles	Rural	Gasoline	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,010	0,457
2012	A	070303	Heavy duty vehicles	Urban	Bio ethanol	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,010	0,457
2012	A	070303	Heavy duty vehicles	Urban	Biodiesel	0,001	1,676	0,423	0,626	0,078	0,063	0,110	6,150	526,000
2012	A	070303	Heavy duty vehicles	Urban	Diesel	0,001	1,676	0,423	0,626	0,078	0,063	0,110	6,150	526,000
2012	A	070303	Heavy duty vehicles	Urban	Gasoline	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,010	0,457
2012	A	070400	Mopeds	Urban	Bio ethanol	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,010	0,457
2012	A	070400	Mopeds	Urban	Gasoline	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,010	0,457
2012	A	070501	Motorcycles	Highway	Bio ethanol	0,020	12,799	0,832	0,640	0,704	1,664	0,640	0,010	0,457
2012	A	070501	Motorcycles	Highway	Gasoline	0,020	12,799	0,832	0,640	0,704	1,664	0,640	0,010	0,457
2012	A	070502	Motorcycles	Rural	Bio ethanol	0,024	15,331	0,996	0,766	0,843	1,993	0,766	0,010	0,457
2012	A	070502	Motorcycles	Rural	Gasoline	0,024	15,331	0,996	0,766	0,843	1,993	0,766	0,010	0,457
2012	A	070503	Motorcycles	Urban	Bio ethanol	0,024	15,500	1,007	0,775	0,852	2,015	0,775	0,010	0,457
2012	A	070503	Motorcycles	Urban	Gasoline	0,024	15,500	1,007	0,775	0,852	2,015	0,775	0,010	0,457
2012	A	080100	Military		AvGas	0,005	4,329	0,209	0,071	0,114	0,689	0,245	0,000	0,000
2012	A	080100	Military		Diesel	0,001	4,350	0,510	0,496	0,256	0,464	0,264	6,150	214,151
2012	A	080100	Military		Gasoline	0,007	2,152	0,180	0,115	0,118	0,358	0,179	0,010	0,457
2012	A	080100	Military		Jet fuel	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
2012	A	080200	Railways		Diesel	0,001	1,411	0,360	0,402	0,059	0,051	0,092	6,150	526,000
2012	A	080200	Railways		Gasoline									
2012	A	080200	Railways		Kerosene									
2012	A	080300	Inland waterways		Diesel	0,001	4,350	0,510	0,496	0,256	0,464	0,264	6,150	526,000
2012	A	080300	Inland waterways		Gasoline	0,005	4,329	0,209	0,071	0,114	0,689	0,245	0,010	0,457
2012	A	080402	National sea traffic		Diesel	0,012	7,420	0,640	0,300	0,150	1,430	1,180	1,950	8,760
2012	A	080402	National sea traffic		Kerosene									
2012	A	080402	National sea traffic		LPG	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
2012	A	080402	National sea traffic		Residual oil	0,013	5,190	0,270	0,050	0,020	0,070	0,030	3,500	14,000
2012	A	080403	Fishing		Diesel	0,012	7,420	0,640	0,300	0,150	1,430	1,180	1,950	8,760
2012	A	080403	Fishing		Gasoline									
2012	A	080403	Fishing		Kerosene									
2012	A	080403	Fishing		LPG	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
2012	A	080404	International sea traffic		Diesel	0,012	7,420	0,640	0,300	0,150	1,430	1,180	1,950	8,760
2012	A	080404	International sea traffic		Residual oil	0,013	4,120	0,200	0,090	0,070	0,260	0,200	3,500	14,000
2012	A	080501	Air traffic, Dom. < 3000 ft.		AvGas	0,005	4,329	0,209	0,071	0,114	0,689	0,245	0,000	0,000
2012	A	080501	Air traffic, Dom. < 3000 ft.		Jet fuel	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000

2012	A	080502	Air traffic, Int. < 3000 ft.	AvGas	0,005	4,329	0,209	0,071	0,114	0,689	0,245	0,000	0,000
2012	A	080502	Air traffic, Int. < 3000 ft.	Jet fuel	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
2012	A	080503	Air traffic, Dom. > 3000 ft.	Jet fuel	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
2012	A	080504	Air traffic, Int. > 3000 ft.	Jet fuel	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
2012	A	080600	Agriculture	Diesel	0,001	4,350	0,510	0,496	0,256	0,464	0,264	6,150	526,000
2012	A	080600	Agriculture	Gasoline	0,005	4,329	0,209	0,071	0,114	0,689	0,245	0,010	0,457
2012	A	080700	Forestry	Diesel	0,001	4,350	0,510	0,496	0,256	0,464	0,264	6,150	526,000
2012	A	080700	Forestry	Gasoline	0,005	4,329	0,209	0,071	0,114	0,689	0,245	0,010	0,457
2012	A	080800	Industry	Diesel	0,001	4,350	0,510	0,496	0,256	0,464	0,264	6,150	526,000
2012	A	080800	Industry	Gasoline	0,005	4,329	0,209	0,071	0,114	0,689	0,245	0,010	0,457
2012	A	080800	Industry	LPG	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
2012	A	080900	Household and gardening	Gasoline	0,005	4,329	0,209	0,071	0,114	0,689	0,245	0,010	0,457
2012	A	081100	Commercial and institutional	Gasoline	0,005	4,329	0,209	0,071	0,114	0,689	0,245	0,010	0,457
2012	P	080501	Air traffic, Dom. < 3000 ft.	AvGas	0,005	4,329	0,209	0,071	0,114	0,689	0,245	0,000	0,000
2012	P	080501	Air traffic, Dom. < 3000 ft.	Jet fuel	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
2012	P	080502	Air traffic, Int. < 3000 ft.	AvGas	0,005	4,329	0,209	0,071	0,114	0,689	0,245	0,000	0,000
2012	P	080502	Air traffic, Int. < 3000 ft.	Jet fuel	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
2012	P	080503	Air traffic, Dom. > 3000 ft.	Jet fuel	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
2012	P	080504	Air traffic, Int. > 3000 ft.	Jet fuel	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000

1990 emissions for SO₂, NO_x, NMVOC, CH₄, CO, CO₂, N₂O, NH₃, TSP, PM₁₀ and PM_{2.5}.

Year	SNAP ID	Category	Fuel type	Fuel PJ	SO ₂ tons	NO _x tons	NMVOC tons	CH ₄ tons	CO tons	CO ₂ ktons	N ₂ O tons	NH ₃ tons	TSP tons	PM ₁₀ tons	PM _{2.5} tons
1990	A 070101	Passenger cars	Highway Diesel	1,84	172,33	514,23	46,12	6,88	330,57	136,13	0,00	0,86	146,22	146,22	146,22
1990	A 070101	Passenger cars	Highway Gasoline	13,64	31,14	17941,01	5206,57	148,53	52065,85	995,70	37,14	11,43	175,16	175,16	175,16
1990	A 070101	Passenger cars	Highway LPG	0,02	0,00	20,08	3,26	0,18	68,23	1,10	0,00	0,00	0,18	0,18	0,18
1990	A 070102	Passenger cars	Rural Diesel	3,74	350,29	1049,16	157,40	25,50	1002,43	276,71	0,00	2,13	280,94	280,94	280,94
1990	A 070102	Passenger cars	Rural Gasoline	29,81	68,05	34230,22	14915,05	411,15	127372,70	2175,93	91,75	28,23	446,30	446,30	446,30
1990	A 070102	Passenger cars	Rural LPG	0,04	0,00	44,81	10,95	0,61	41,15	2,26	0,00	0,00	0,52	0,52	0,52
1990	A 070103	Passenger cars	Urban Diesel	2,57	240,85	567,80	264,93	17,55	886,92	190,26	0,00	0,81	370,85	370,85	370,85
1990	A 070103	Passenger cars	Urban Gasoline	19,93	45,50	11129,18	21829,78	1126,14	249404,91	1454,89	54,00	10,80	246,59	246,59	246,59
1990	A 070103	Passenger cars	Urban LPG	0,03	0,00	13,47	12,07	0,53	40,50	1,61	0,00	0,00	0,27	0,27	0,27
1990	A 070201	Light duty vehicles	Highway Diesel	3,35	314,00	907,29	101,20	8,71	1153,53	248,05	0,00	1,09	350,21	350,21	350,21
1990	A 070201	Light duty vehicles	Highway Gasoline	0,40	0,90	542,61	67,48	4,01	1183,85	28,93	1,04	0,32	6,41	6,41	6,41
1990	A 070201	Light duty vehicles	Highway LPG	0,01	0,00	16,85	2,74	0,15	57,28	0,92	0,00	0,00	0,15	0,15	0,15
1990	A 070202	Light duty vehicles	Rural Diesel	7,98	747,28	2387,17	265,03	34,00	2859,22	590,31	0,00	2,83	859,40	859,40	859,40
1990	A 070202	Light duty vehicles	Rural Gasoline	1,09	2,50	1300,90	287,33	16,69	2534,47	79,88	2,71	0,83	16,69	16,69	16,69
1990	A 070202	Light duty vehicles	Rural LPG	0,03	0,00	39,65	9,69	0,54	36,40	2,00	0,00	0,00	0,46	0,46	0,46
1990	A 070203	Light duty vehicles	Urban Diesel	3,71	347,82	1742,28	277,76	16,99	1720,50	274,76	0,00	0,84	615,45	615,45	615,45
1990	A 070203	Light duty vehicles	Urban Gasoline	0,67	1,53	367,62	586,86	26,75	6396,69	49,01	1,24	0,25	4,95	4,95	4,95
1990	A 070203	Light duty vehicles	Urban LPG	0,02	0,00	9,07	9,07	0,36	31,71	1,17	0,00	0,00	0,18	0,18	0,18
1990	A 070301	Heavy duty vehicles	Highway Diesel	10,37	971,35	10145,87	432,52	66,80	2074,02	767,32	31,95	3,20	363,93	363,93	363,93
1990	A 070301	Heavy duty vehicles	Highway Gasoline	0,03	0,08	35,45	16,21	0,33	259,98	2,49	0,03	0,01	1,89	1,89	1,89
1990	A 070302	Heavy duty vehicles	Rural Diesel	17,72	1659,89	17450,79	1023,06	120,07	3743,13	1311,23	51,77	5,18	633,09	633,09	633,09
1990	A 070302	Heavy duty vehicles	Rural Gasoline	0,09	0,21	103,04	74,05	1,51	755,60	6,59	0,08	0,03	5,50	5,50	5,50
1990	A 070303	Heavy duty vehicles	Urban Diesel	8,67	811,84	8380,30	752,74	107,60	2339,19	641,31	19,90	1,99	365,78	365,78	365,78
1990	A 070303	Heavy duty vehicles	Urban Gasoline	0,08	0,17	34,44	52,51	1,07	535,81	5,51	0,05	0,02	3,06	3,06	3,06
1990	A 070400	Mopeds	Urban Gasoline	0,18	0,40	9,01	1316,72	35,25	2365,94	12,87	0,16	0,16	28,33	28,33	28,33
1990	A 070501	Motorcycles	Highway Gasoline	0,06	0,13	14,85	60,27	7,31	916,37	4,10	0,08	0,08	1,78	1,78	1,78
1990	A 070502	Motorcycles	Rural Gasoline	0,13	0,31	24,98	132,28	21,47	2126,59	9,84	0,22	0,22	5,24	5,24	5,24
1990	A 070503	Motorcycles	Urban Gasoline	0,17	0,39	19,53	198,77	26,83	2626,92	12,63	0,28	0,28	6,55	6,55	6,55
1990	A 080100	Military	AvGas	0,00	0,11	4,22	6,11	0,11	34,26	0,36	0,01	0,01	0,05	0,05	0,05
1990	A 080100	Military	Diesel	0,15	13,69	105,20	8,10	0,99	39,28	10,82	0,25	0,05	9,72	9,72	9,72
1990	A 080100	Military	Gasoline	0,00	0,00	0,98	1,10	0,03	6,67	0,07	0,00	0,00	0,01	0,01	0,01
1990	A 080100	Military	Jet fuel	1,50	34,41	375,06	37,33	3,96	344,09	107,77	3,44		1,74	1,74	1,74
1990	A 080200	Railways	Diesel	4,01	375,64	4912,78	320,54	12,32	895,07	296,74	8,18	0,82	201,55	201,55	201,55
1990	A 080200	Railways	Gasoline	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00

1990	A	080300	Inland waterways	Diesel	0,34	32,10	337,02	58,86	0,96	155,43	25,35	1,01	0,06	36,64	36,64	36,64
1990	A	080300	Inland waterways	Gasoline	0,31	0,71	90,06	1114,91	15,58	4282,54	22,57	0,24	0,02	56,40	56,40	56,40
1990	A	080402	National sea traffic	Diesel	5,29	495,12	5836,01	267,28	8,27	881,74	391,12	24,76		122,69	121,47	120,85
1990	A	080402	National sea traffic	Residual oil	4,57	5901,32	7383,82	244,28	7,56	805,87	356,56	22,35		682,25	675,43	672,02
1990	A	080403	Fishing	Diesel	7,92	741,91	8332,71	389,10	12,03	1283,63	586,07	37,10		183,85	182,01	181,09
1990	A	080403	Fishing	Gasoline	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
1990	A	080403	Fishing	Residual oil	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00		0,00	0,00	0,00
1990	A	080404	International sea traffic	Diesel	11,29	1057,56	13644,52	558,38	17,27	1842,07	835,42	52,88		262,07	259,45	258,14
1990	A	080404	International sea traffic	Residual oil	27,81	40259,78	46994,61	1501,54	46,44	4953,54	2169,54	136,01		5268,82	5216,14	5189,79
1990	A	080501	Air traffic, Dom. < 3000 ft.	AvGas	0,10	2,40	90,15	130,41	2,30	731,69	7,66	0,21	0,17	1,05	1,05	1,05
1990	A	080501	Air traffic, Dom. < 3000 ft.	Jet fuel	0,42	9,71	132,78	6,27	0,70	38,17	30,40	2,40		0,49	0,49	0,49
1990	A	080502	Air traffic, Int. < 3000 ft.	AvGas	0,03	0,70	26,34	38,10	0,67	213,76	2,24	0,06	0,05	0,31	0,31	0,31
1990	A	080502	Air traffic, Int. < 3000 ft.	Jet fuel	0,13	3,04	40,93	2,17	0,24	22,36	9,53	0,94		0,15	0,15	0,15
1990	A	080503	Air traffic, Dom. > 3000 ft.	Jet fuel	1,03	23,59	338,70	14,02	0,00	93,11	73,87	2,36		1,19	1,19	1,19
1990	A	080504	Air traffic, Int. > 3000 ft.	Jet fuel	1,61	37,06	393,62	11,56	0,00	87,20	116,06	3,71		1,87	1,87	1,87
1990	A	080600	Agriculture	Diesel	16,50	1545,32	12518,46	2587,36	42,07	10483,86	1220,72	48,34	2,76	2382,90	2382,90	2382,90
1990	A	080600	Agriculture	Gasoline	0,71	1,62	22,40	673,10	62,68	33688,19	51,75	0,91	0,06	4,65	4,65	4,65
1990	A	080700	Forestry	Diesel	0,15	13,62	124,63	22,74	0,37	93,84	10,76	0,43	0,02	21,66	21,66	21,66
1990	A	080700	Forestry	Gasoline	0,34	0,78	13,79	2460,65	20,63	6165,33	24,92	0,13	0,03	34,56	34,56	34,56
1990	A	080800	Industry	Diesel	10,16	951,61	9483,66	1810,53	29,44	6661,90	751,72	29,87	1,71	1569,49	1569,49	1569,49
1990	A	080800	Industry	Gasoline	0,18	0,40	23,88	282,25	21,13	2592,92	12,79	0,23	0,02	2,17	2,17	2,17
1990	A	080800	Industry	LPG	1,18	0,00	1573,62	173,10	9,11	124,23	74,76	4,14	0,25	5,80	5,80	5,80
1990	A	080900	Household and gardening	Gasoline	0,54	1,22	34,24	1801,26	50,96	17606,46	39,06	0,62	0,04	11,10	11,10	11,10
1990	A	081100	Commercial and institutional	Gasoline	1,01	2,31	69,51	2303,07	98,83	30181,04	73,72	1,10	0,08	24,24	24,24	24,24
1990	P	080501	Air traffic, Dom. < 3000 ft.	AvGas	0,01	0,20	7,42	10,74	0,19	60,25	0,63	0,02	0,01	0,09	0,09	0,09
1990	P	080501	Air traffic, Dom. < 3000 ft.	Jet fuel	0,50	11,54	142,54	10,36	1,15	65,13	36,16	2,30		0,58	0,58	0,58
1990	P	080502	Air traffic, Int. < 3000 ft.	AvGas	0,01	0,13	4,82	6,97	0,12	39,13	0,41	0,01	0,01	0,06	0,06	0,06
1990	P	080502	Air traffic, Int. < 3000 ft.	Jet fuel	2,00	46,00	650,12	68,24	7,58	314,49	144,09	7,58		2,32	2,32	2,32
1990	P	080503	Air traffic, Dom. > 3000 ft.	Jet fuel	1,31	30,00	410,96	17,01	0,00	109,71	93,97	3,00		1,51	1,51	1,51
1990	P	080504	Air traffic, Int. > 3000 ft.	Jet fuel	20,33	467,36	5899,81	226,68	0,00	765,45	1463,78	46,74		23,58	23,58	23,58

1990 emissions for Arsenic, Cadmium, Chromium, Copper, Mercury, Nickel, Lead, Selenium and Zinc.

Year	SNAP ID	Category	Fuel type	Fuel PJ	Arsenic kg	Cadmium kg	Chromium kg	Copper kg	Mercury kg	Nickel kg	Lead kg	Selenium kg	Zinc kg
1990	A 070101	Passenger cars	Highway Diesel	1,84	0,00	0,54	1,70	1,21	0,23	0,54	3,23	0,00	107,81
1990	A 070101	Passenger cars	Highway Gasoline	13,64	0,09	3,72	5,26	14,22	2,71	4,38	20077,75	0,06	742,80
1990	A 070101	Passenger cars	Highway LPG	0,02	0,00	0,01	0,00	0,02	0,00	0,01	0,02	0,00	1,05
1990	A 070102	Passenger cars	Rural Diesel	3,74	0,01	1,33	4,05	2,88	0,46	1,34	7,98	0,01	266,03
1990	A 070102	Passenger cars	Rural Gasoline	29,81	0,20	9,19	12,43	34,73	5,92	10,61	43879,62	0,14	1832,30
1990	A 070102	Passenger cars	Rural LPG	0,04	0,00	0,01	0,01	0,05	0,00	0,01	0,04	0,00	2,59
1990	A 070103	Passenger cars	Urban Diesel	2,57	0,01	0,51	1,78	1,25	0,32	0,52	3,07	0,01	102,25
1990	A 070103	Passenger cars	Urban Gasoline	19,93	0,14	3,55	5,98	14,16	3,96	4,51	29331,40	0,09	707,35
1990	A 070103	Passenger cars	Urban LPG	0,03	0,00	0,00	0,00	0,02	0,00	0,00	0,01	0,00	0,99
1990	A 070201	Light duty vehicles	Highway Diesel	3,35	0,01	0,65	2,27	1,60	0,42	0,66	3,89	0,01	129,82
1990	A 070201	Light duty vehicles	Highway Gasoline	0,40	0,00	0,10	0,15	0,39	0,08	0,12	583,30	0,00	19,97
1990	A 070201	Light duty vehicles	Highway LPG	0,01	0,00	0,00	0,00	0,01	0,00	0,00	0,01	0,00	0,58
1990	A 070202	Light duty vehicles	Rural Diesel	7,98	0,02	1,68	5,77	4,07	0,99	1,71	10,12	0,02	337,73
1990	A 070202	Light duty vehicles	Rural Gasoline	1,09	0,01	0,26	0,39	1,01	0,22	0,31	1610,62	0,00	52,06
1990	A 070202	Light duty vehicles	Rural LPG	0,03	0,00	0,01	0,01	0,03	0,00	0,01	0,02	0,00	1,51
1990	A 070203	Light duty vehicles	Urban Diesel	3,71	0,01	0,50	1,98	1,39	0,46	0,51	3,02	0,01	100,74
1990	A 070203	Light duty vehicles	Urban Gasoline	0,67	0,00	0,08	0,16	0,33	0,13	0,11	987,98	0,00	15,70
1990	A 070203	Light duty vehicles	Urban LPG	0,02	0,00	0,00	0,00	0,01	0,00	0,00	0,01	0,00	0,45
1990	A 070301	Heavy duty vehicles	Highway Diesel	10,37	0,02	1,55	5,91	4,16	1,29	1,59	9,36	0,02	312,42
1990	A 070301	Heavy duty vehicles	Highway Gasoline	0,03	0,00	0,01	0,01	0,03	0,01	0,01	50,28	0,00	1,41
1990	A 070302	Heavy duty vehicles	Rural Diesel	17,72	0,04	2,54	9,83	6,90	2,20	2,60	15,34	0,04	511,81
1990	A 070302	Heavy duty vehicles	Rural Gasoline	0,09	0,00	0,02	0,03	0,08	0,02	0,02	132,85	0,00	3,96
1990	A 070303	Heavy duty vehicles	Urban Diesel	8,67	0,02	0,99	4,19	2,93	1,08	1,03	6,01	0,02	200,56
1990	A 070303	Heavy duty vehicles	Urban Gasoline	0,08	0,00	0,01	0,02	0,04	0,01	0,01	111,01	0,00	2,14
1990	A 070400	Mopeds	Urban Gasoline	0,18	0,00	0,00	0,03	0,02	0,04	0,01	259,28	0,00	0,13
1990	A 070501	Motorcycles	Highway Gasoline	0,06	0,00	0,01	0,01	0,03	0,01	0,01	82,72	0,00	1,50
1990	A 070502	Motorcycles	Rural Gasoline	0,13	0,00	0,02	0,04	0,09	0,03	0,03	198,30	0,00	4,38
1990	A 070503	Motorcycles	Urban Gasoline	0,17	0,00	0,03	0,05	0,11	0,03	0,04	254,55	0,00	5,48
1990	A 080100	Military	AvGas	0,00	0,00	0,00	0,00	0,00	0,00	0,00	62,82	0,00	0,25
1990	A 080100	Military	Diesel	0,15	0,00	0,03	0,09	0,06	0,02	0,03	0,15	0,00	5,05
1990	A 080100	Military	Gasoline	0,00	0,00	0,00	0,00	0,00	0,00	0,00	1,45	0,00	0,05
1990	A 080100	Military	Jet fuel	1,50		0,00	0,00	0,00		0,00		0,00	0,00
1990	A 080200	Railways	Diesel	4,01	0,01	0,74	2,65	1,87	0,50	0,76	4,48	0,01	149,55
1990	A 080200	Railways	Gasoline	0,00		0,00	0,00	0,00		0,00	0,00	0,00	0,00

1990	A	080300	Inland waterways	Diesel	0,34	0,00	0,06	0,23	0,16	0,04	0,06	0,38	0,00	12,78
1990	A	080300	Inland waterways	Gasoline	0,31	0,00	0,08	0,11	0,30	0,06	0,09	0,24	0,00	15,60
1990	A	080402	National sea traffic	Diesel	5,29	6,19	1,24	4,95	6,19	6,18	8,66	12,37	24,76	61,89
1990	A	080402	National sea traffic	Residual oil	4,57	55,88	3,35	22,35	55,88	2,24	3353,02	22,35	44,71	100,59
1990	A	080403	Fishing	Diesel	7,92	9,27	1,85	7,42	9,27	9,27	12,98	18,53	37,10	92,74
1990	A	080403	Fishing	Gasoline	0,00		0,00	0,00	0,00		0,00	0,00	0,00	0,00
1990	A	080403	Fishing	Residual oil	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
1990	A	080404	International sea traffic	Diesel	11,29	13,22	2,64	10,58	13,22	13,21	18,51	26,42	52,88	132,20
1990	A	080404	International sea traffic	Residual oil	27,81	340,03	20,40	136,01	340,03	13,63	20401,92	136,01	272,03	612,06
1990	A	080501	Air traffic, Dom. < 3000 ft.	AvGas	0,10	0,00	0,03	0,04	0,10	0,02	0,03	1417,38	0,00	5,29
1990	A	080501	Air traffic, Dom. < 3000 ft.	Jet fuel	0,42		0,00	0,00	0,00		0,00		0,00	0,00
1990	A	080502	Air traffic, Int. < 3000 ft.	AvGas	0,03	0,00	0,01	0,01	0,03	0,01	0,01	414,08	0,00	1,55
1990	A	080502	Air traffic, Int. < 3000 ft.	Jet fuel	0,13		0,00	0,00	0,00		0,00		0,00	0,00
1990	A	080503	Air traffic, Dom. > 3000 ft.	Jet fuel	1,03		0,00	0,00	0,00		0,00		0,00	0,00
1990	A	080504	Air traffic, Int. > 3000 ft.	Jet fuel	1,61		0,00	0,00	0,00		0,00		0,00	0,00
1990	A	080600	Agriculture	Diesel	16,50	0,04	3,06	10,88	7,67	2,04	3,12	18,44	0,04	615,23
1990	A	080600	Agriculture	Gasoline	0,71	0,00	0,18	0,26	0,69	0,14	0,21	0,55	0,00	35,76
1990	A	080700	Forestry	Diesel	0,15	0,00	0,03	0,10	0,07	0,02	0,03	0,16	0,00	5,42
1990	A	080700	Forestry	Gasoline	0,34	0,00	0,09	0,13	0,33	0,07	0,10	0,26	0,00	17,23
1990	A	080800	Industry	Diesel	10,16	0,02	1,88	6,70	4,73	1,26	1,92	11,36	0,02	378,86
1990	A	080800	Industry	Gasoline	0,18	0,00	0,04	0,06	0,17	0,03	0,05	0,14	0,00	8,84
1990	A	080800	Industry	LPG	1,18	0,00	0,15	0,14	0,54	0,00	0,15	0,46	0,00	30,96
1990	A	080900	Household and gardening	Gasoline	0,54	0,00	0,14	0,20	0,52	0,11	0,16	0,41	0,00	27,00
1990	A	081100	Commercial and institutional	Gasoline	1,01	0,01	0,26	0,37	0,98	0,20	0,30	0,78	0,00	50,95
1990	P	080501	Air traffic, Dom. < 3000 ft.	AvGas	0,01	0,00	0,00	0,00	0,01	0,00	0,00	116,72	0,00	0,44
1990	P	080501	Air traffic, Dom. < 3000 ft.	Jet fuel	0,50		0,00	0,00	0,00		0,00		0,00	0,00
1990	P	080502	Air traffic, Int. < 3000 ft.	AvGas	0,01	0,00	0,00	0,00	0,01	0,00	0,00	75,80	0,00	0,28
1990	P	080502	Air traffic, Int. < 3000 ft.	Jet fuel	2,00		0,00	0,00	0,00		0,00		0,00	0,00
1990	P	080503	Air traffic, Dom. > 3000 ft.	Jet fuel	1,31		0,00	0,00	0,00		0,00		0,00	0,00
1990	P	080504	Air traffic, Int. > 3000 ft.	Jet fuel	20,33		0,00	0,00	0,00		0,00		0,00	0,00

1990 emissions for Dioxins/, Flouranthene, Benzo(b), Benzo(k), Benzo(a), Benzo(g,h,i), indeno(1,2,3-c,d), HCB and PCB.

Year	SNAP ID	Category	Fuel type	Fuel	Dioxins/	Flouranthene	Benzo(b)	Benzo(k)	Benzo(a)	Benzo(g,h,i)	indeno(1,2,3-c,d)	HCB	PCB
				PJ	g	kg	kg	kg	kg	kg	kg	g	g
1990	A 070101	Passenger cars	Highway Diesel	1,84	0,00	22,53	1,38	1,25	1,50	2,92	1,42	0,01	0,04
1990	A 070101	Passenger cars	Highway Gasoline	13,64	0,18	116,03	7,54	5,80	6,38	15,08	5,80	0,00	14,05
1990	A 070101	Passenger cars	Highway LPG	0,02								0,00	0,00
1990	A 070102	Passenger cars	Rural Diesel	3,74	0,00	55,68	3,40	3,08	3,72	7,22	3,51	0,02	0,08
1990	A 070102	Passenger cars	Rural Gasoline	29,81	0,45	284,37	18,48	14,21	15,64	36,97	14,21	0,00	30,69
1990	A 070102	Passenger cars	Rural LPG	0,04								0,00	0,00
1990	A 070103	Passenger cars	Urban Diesel	2,57	0,00	23,92	1,46	1,32	1,60	3,10	1,51	0,02	0,05
1990	A 070103	Passenger cars	Urban Gasoline	19,93	0,20	128,10	8,33	6,40	7,05	16,65	6,40	0,00	20,52
1990	A 070103	Passenger cars	Urban LPG	0,03								0,00	0,00
1990	A 070201	Light duty vehicles	Highway Diesel	3,35	0,00	28,51	1,74	1,58	1,90	3,70	1,80	0,02	0,07
1990	A 070201	Light duty vehicles	Highway Gasoline	0,40	0,01	3,20	0,21	0,16	0,18	0,42	0,16	0,00	0,41
1990	A 070201	Light duty vehicles	Highway LPG	0,01	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
1990	A 070202	Light duty vehicles	Rural Diesel	7,98	0,00	74,23	4,53	4,11	4,96	9,63	4,67	0,05	0,17
1990	A 070202	Light duty vehicles	Rural Gasoline	1,09	0,01	8,34	0,54	0,42	0,46	1,08	0,42	0,00	1,13
1990	A 070202	Light duty vehicles	Rural LPG	0,03	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
1990	A 070203	Light duty vehicles	Urban Diesel	3,71	0,00	25,82	1,58	1,43	1,72	3,35	1,63	0,02	0,08
1990	A 070203	Light duty vehicles	Urban Gasoline	0,67	0,00	3,06	0,20	0,15	0,17	0,40	0,15	0,00	0,69
1990	A 070203	Light duty vehicles	Urban LPG	0,02	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
1990	A 070301	Heavy duty vehicles	Highway Diesel	10,37	0,01	21,63	5,46	8,08	1,01	0,81	1,41	0,06	5,45
1990	A 070301	Heavy duty vehicles	Highway Gasoline	0,03								0,00	0,04
1990	A 070302	Heavy duty vehicles	Rural Diesel	17,72	0,02	39,12	9,87	14,62	1,83	1,46	2,56	0,11	9,32
1990	A 070302	Heavy duty vehicles	Rural Gasoline	0,09								0,00	0,09
1990	A 070303	Heavy duty vehicles	Urban Diesel	8,67	0,01	15,49	3,91	5,79	0,72	0,58	1,01	0,05	4,56
1990	A 070303	Heavy duty vehicles	Urban Gasoline	0,08								0,00	0,08
1990	A 070400	Mopeds	Urban Gasoline	0,18								0,00	0,18
1990	A 070501	Motorcycles	Highway Gasoline	0,06	0,00	0,71	0,05	0,04	0,04	0,09	0,04	0,00	0,06
1990	A 070502	Motorcycles	Rural Gasoline	0,13	0,00	2,04	0,13	0,10	0,11	0,27	0,10	0,00	0,14
1990	A 070503	Motorcycles	Urban Gasoline	0,17	0,00	2,65	0,17	0,13	0,15	0,34	0,13	0,00	0,18
1990	A 080100	Military	AvGas	0,00	0,00	0,02	0,00	0,00	0,00	0,00	0,00	0,00	0,00
1990	A 080100	Military	Diesel	0,15	0,00	0,64	0,08	0,08	0,04	0,08	0,04	0,00	0,05
1990	A 080100	Military	Gasoline	0,00	0,00	0,01	0,00	0,00	0,00	0,00	0,00	0,00	0,00
1990	A 080100	Military	Jet fuel	1,50	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
1990	A 080200	Railways	Diesel	4,01	0,00	5,48	1,40	1,56	0,23	0,20	0,36	0,02	2,11
1990	A 080200	Railways	Gasoline	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00

1990	A	080300	Inland waterways	Diesel	0,34	0,00	1,50	0,20	0,19	0,10	0,19	0,10	0,00	0,18
1990	A	080300	Inland waterways	Gasoline	0,31	0,00	1,34	0,06	0,02	0,04	0,21	0,08	0,00	0,32
1990	A	080402	National sea traffic	Diesel	5,29	0,06	39,22	3,38	1,59	0,79	7,56	6,24	0,01	0,05
1990	A	080402	National sea traffic	Residual oil	4,57	0,06	23,72	1,23	0,23	0,09	0,32	0,14	0,02	0,06
1990	A	080403	Fishing	Diesel	7,92	0,10	58,77	5,07	2,38	1,19	11,33	9,35	0,02	0,07
1990	A	080403	Fishing	Gasoline	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
1990	A	080403	Fishing	Residual oil	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
1990	A	080404	International sea traffic	Diesel	11,29	0,14	83,77	7,23	3,39	1,69	16,14	13,32	0,02	0,10
1990	A	080404	International sea traffic	Residual oil	27,81	0,37	114,60	5,56	2,50	1,95	7,23	5,56	0,10	0,39
1990	A	080501	Air traffic, Dom. < 3000 ft.	AvGas	0,10	0,00	0,45	0,02	0,01	0,01	0,07	0,03	0,00	0,00
1990	A	080501	Air traffic, Dom. < 3000 ft.	Jet fuel	0,42	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
1990	A	080502	Air traffic, Int. < 3000 ft.	AvGas	0,03	0,00	0,13	0,01	0,00	0,00	0,02	0,01	0,00	0,00
1990	A	080502	Air traffic, Int. < 3000 ft.	Jet fuel	0,13	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
1990	A	080503	Air traffic, Dom. > 3000 ft.	Jet fuel	1,03	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
1990	A	080504	Air traffic, Int. > 3000 ft.	Jet fuel	1,61	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
1990	A	080600	Agriculture	Diesel	16,50	0,01	72,44	9,41	9,37	4,78	9,07	4,79	0,10	8,68
1990	A	080600	Agriculture	Gasoline	0,71	0,00	3,07	0,15	0,05	0,08	0,49	0,17	0,00	0,73
1990	A	080700	Forestry	Diesel	0,15	0,00	0,64	0,08	0,08	0,04	0,08	0,04	0,00	0,08
1990	A	080700	Forestry	Gasoline	0,34	0,00	1,48	0,07	0,02	0,04	0,24	0,08	0,00	0,35
1990	A	080800	Industry	Diesel	10,16	0,01	44,61	5,80	5,77	2,94	5,59	2,95	0,06	5,34
1990	A	080800	Industry	Gasoline	0,18	0,00	0,76	0,04	0,01	0,02	0,12	0,04	0,00	0,18
1990	A	080800	Industry	LPG	1,18								0,00	0,00
1990	A	080900	Household and gardening	Gasoline	0,54	0,00	2,32	0,11	0,04	0,06	0,37	0,13	0,00	0,55
1990	A	081100	Commercial and institutional	Gasoline	1,01	0,01	4,37	0,21	0,07	0,12	0,70	0,25	0,00	1,04
1990	P	080501	Air traffic, Dom. < 3000 ft.	AvGas	0,01	0,00	0,04	0,00	0,00	0,00	0,01	0,00		
1990	P	080501	Air traffic, Dom. < 3000 ft.	Jet fuel	0,50	0,00	0,00	0,00	0,00	0,00	0,00	0,00		
1990	P	080502	Air traffic, Int. < 3000 ft.	AvGas	0,01	0,00	0,02	0,00	0,00	0,00	0,00	0,00		
1990	P	080502	Air traffic, Int. < 3000 ft.	Jet fuel	2,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00		
1990	P	080503	Air traffic, Dom. > 3000 ft.	Jet fuel	1,31	0,00	0,00	0,00	0,00	0,00	0,00	0,00		
1990	P	080504	Air traffic, Int. > 3000 ft.	Jet fuel	20,33	0,00	0,00	0,00	0,00	0,00	0,00	0,00		

2012 emissions for SO₂, NO_x, NMVOC, CH₄, CO, CO₂, N₂O, NH₃, TSP, PM₁₀ and PM_{2.5}.

Year	SNAP ID	Category	Fuel type	Fuel PJ	SO ₂ tons	NO _x tons	NMVOC tons	CH ₄ tons	CO tons	CO ₂ ktons	N ₂ O tons	NH ₃ tons	TSP tons	PM ₁₀ tons	PM _{2.5} tons
2012	A 070101	Passenger cars	Highway Bio ethanol	0,51	0,00	67,81	16,18	1,85	354,39	0,00	0,38	16,57	0,52	0,52	0,52
2012	A 070101	Passenger cars	Highway Biodiesel	0,74	0,00	254,21	2,96	0,08	9,52	0,00	1,52	0,37	8,92	8,92	8,92
2012	A 070101	Passenger cars	Highway Diesel	11,03	5,16	3804,87	44,23	1,16	142,42	815,90	22,79	5,59	133,46	133,46	133,46
2012	A 070101	Passenger cars	Highway Gasoline	13,94	6,37	1854,54	442,61	50,59	9691,85	1017,93	10,27	453,08	14,24	14,24	14,24
2012	A 070101	Passenger cars	Highway LPG	0,00	0,00	0,05	0,01	0,00	0,30	0,01	0,00	0,00	0,00	0,00	0,00
2012	A 070102	Passenger cars	Rural Bio ethanol	0,83	0,00	92,05	30,58	2,92	463,27	0,00	1,14	28,06	0,81	0,81	0,81
2012	A 070102	Passenger cars	Rural Biodiesel	1,19	0,00	343,09	6,24	0,25	29,31	0,00	2,64	0,65	11,99	11,99	11,99
2012	A 070102	Passenger cars	Rural Diesel	17,79	8,33	5135,05	93,38	3,72	438,75	1316,38	39,47	9,68	179,51	179,51	179,51
2012	A 070102	Passenger cars	Rural Gasoline	22,81	10,42	2517,30	836,39	79,89	12669,41	1665,10	31,28	767,44	22,21	22,21	22,21
2012	A 070102	Passenger cars	Rural LPG	0,00	0,00	0,08	0,02	0,00	0,17	0,02	0,00	0,00	0,00	0,00	0,00
2012	A 070103	Passenger cars	Urban Bio ethanol	0,57	0,00	79,02	168,23	6,58	1848,83	0,00	1,18	4,38	0,50	0,50	0,50
2012	A 070103	Passenger cars	Urban Biodiesel	0,70	0,00	202,02	11,17	0,36	49,87	0,00	3,82	0,26	12,73	12,73	12,73
2012	A 070103	Passenger cars	Urban Diesel	10,40	4,87	3023,73	167,13	5,36	746,34	769,77	57,15	3,82	190,51	190,51	190,51
2012	A 070103	Passenger cars	Urban Gasoline	15,59	7,12	2161,08	4600,64	179,84	50561,05	1137,88	32,16	119,78	13,77	13,77	13,77
2012	A 070103	Passenger cars	Urban LPG	0,00	0,00	0,03	0,03	0,00	0,19	0,01	0,00	0,00	0,00	0,00	0,00
2012	A 070201	Light duty vehicles	Highway Bio ethanol	0,01	0,00	2,01	0,24	0,03	6,81	0,00	0,02	0,28	0,02	0,02	0,02
2012	A 070201	Light duty vehicles	Highway Biodiesel	0,38	0,00	107,10	6,89	0,07	46,40	0,00	0,57	0,14	6,91	6,91	6,91
2012	A 070201	Light duty vehicles	Highway Diesel	5,66	2,65	1602,96	103,12	0,99	694,44	418,78	8,53	2,08	103,42	103,42	103,42
2012	A 070201	Light duty vehicles	Highway Gasoline	0,34	0,15	55,09	6,57	0,92	186,11	24,46	0,63	7,68	0,46	0,46	0,46
2012	A 070201	Light duty vehicles	Highway LPG	0,00	0,00	0,04	0,01	0,00	0,33	0,02	0,00	0,00	0,00	0,00	0,00
2012	A 070202	Light duty vehicles	Rural Bio ethanol	0,02	0,00	3,44	0,69	0,07	10,06	0,00	0,07	0,54	0,03	0,03	0,03
2012	A 070202	Light duty vehicles	Rural Biodiesel	0,64	0,00	188,52	13,14	0,26	67,01	0,00	1,05	0,26	9,52	9,52	9,52
2012	A 070202	Light duty vehicles	Rural Diesel	9,54	4,47	2821,59	196,64	3,90	1002,88	705,59	15,73	3,83	142,52	142,52	142,52
2012	A 070202	Light duty vehicles	Rural Gasoline	0,65	0,30	94,11	18,85	1,96	275,02	47,77	1,97	14,82	0,80	0,80	0,80
2012	A 070202	Light duty vehicles	Rural LPG	0,00	0,00	0,07	0,02	0,00	0,21	0,03	0,00	0,00	0,01	0,01	0,01
2012	A 070203	Light duty vehicles	Urban Bio ethanol	0,02	0,00	2,48	3,82	0,15	71,68	0,00	0,09	0,08	0,01	0,01	0,01
2012	A 070203	Light duty vehicles	Urban Biodiesel	0,40	0,00	109,10	16,88	0,32	56,16	0,00	1,39	0,10	10,00	10,00	10,00
2012	A 070203	Light duty vehicles	Urban Diesel	5,93	2,78	1632,94	252,72	4,79	840,58	439,03	20,82	1,53	149,62	149,62	149,62
2012	A 070203	Light duty vehicles	Urban Gasoline	0,52	0,24	67,81	104,45	4,21	1960,17	38,05	2,38	2,31	0,38	0,38	0,38
2012	A 070203	Light duty vehicles	Urban LPG	0,00	0,00	0,02	0,02	0,00	0,17	0,02	0,00	0,00	0,00	0,00	0,00
2012	A 070301	Heavy duty vehicles	Highway Bio ethanol	0,00	0,00	0,62	0,28	0,01	4,54	0,00	0,00	0,00	0,03	0,03	0,03
2012	A 070301	Heavy duty vehicles	Highway Biodiesel	0,96	0,00	375,59	7,12	1,96	131,01	0,00	3,00	0,30	5,92	5,92	5,92
2012	A 070301	Heavy duty vehicles	Highway Diesel	14,33	6,71	5621,53	106,53	29,27	1960,92	1060,75	44,88	4,49	88,65	88,65	88,65
2012	A 070301	Heavy duty vehicles	Highway Gasoline	0,02	0,01	16,92	7,74	0,16	124,12	1,19	0,01	0,00	0,90	0,90	0,90

2012	A	070302	Heavy duty vehicles	Rural	Biodiesel	1,14	0,00	518,83	11,43	2,67	153,74	0,00	3,36	0,34	7,57	7,57	7,57
2012	A	070302	Heavy duty vehicles	Rural	Diesel	17,07	8,00	7765,50	171,00	39,94	2301,11	1263,14	50,32	5,03	113,27	113,27	113,27
2012	A	070302	Heavy duty vehicles	Rural	Gasoline	0,04	0,02	42,14	30,28	0,62	309,02	2,69	0,03	0,01	2,25	2,25	2,25
2012	A	070303	Heavy duty vehicles	Urban	Diesel	5,91	2,77	3372,53	87,20	22,08	883,88	437,58	14,79	1,48	47,88	47,88	47,88
2012	A	070303	Heavy duty vehicles	Urban	Gasoline	0,02	0,01	11,19	17,06	0,35	174,10	1,79	0,01	0,00	0,99	0,99	0,99
2012	A	070400	Mopeds	Urban	Bio ethanol	0,01	0,00	1,08	32,67	0,68	54,03	0,00	0,01	0,01	0,57	0,57	0,57
2012	A	070501	Motorcycles	Highway	Gasoline	0,11	0,05	29,05	69,92	9,73	1116,11	7,84	0,14	0,14	1,69	1,69	1,69
2012	A	070502	Motorcycles	Rural	Bio ethanol	0,01	0,00	1,64	5,54	0,93	81,96	0,00	0,01	0,01	0,16	0,16	0,16
2012	A	070502	Motorcycles	Rural	Gasoline	0,23	0,11	44,84	151,59	25,51	2241,45	17,01	0,36	0,36	4,46	4,46	4,46
2012	A	070503	Motorcycles	Urban	Bio ethanol	0,01	0,00	1,21	8,14	1,17	94,09	0,00	0,02	0,02	0,19	0,19	0,19
2012	A	070503	Motorcycles	Urban	Gasoline	0,28	0,13	32,99	222,71	32,07	2573,10	20,27	0,42	0,42	5,18	5,18	5,18
2012	A	080100	Military		AvGas	0,00	0,09	3,21	4,64	0,08	26,02	0,27	0,01	0,01	0,04	0,04	0,04
2012	A	080100	Military		Diesel	0,88	0,39	313,55	11,02	1,00	81,24	65,15	2,47	0,34	10,36	10,36	10,36
2012	A	080100	Military		Gasoline	0,00	0,00	0,19	0,24	0,01	2,30	0,11	0,00	0,04	0,00	0,00	0,00
2012	A	080100	Military		Jet fuel	0,70	16,05	174,89	17,41	1,85	160,46	50,25	1,61	0,00	0,81	0,81	0,81
2012	A	080200	Railways		Diesel	3,37	1,58	2530,89	190,12	7,30	424,51	249,33	6,87	0,67	81,68	81,68	81,68
2012	A	080200	Railways		Gasoline	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
2012	A	080200	Railways		Kerosene	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
2012	A	080300	Inland waterways		Diesel	1,00	46,94	818,78	156,85	2,55	440,28	74,16	2,98	0,17	95,75	95,75	95,75
2012	A	080300	Inland waterways		Gasoline	0,34	0,15	190,06	293,98	21,58	3772,37	24,46	0,52	0,04	5,84	5,84	5,84
2012	A	080402	National sea traffic		Diesel	2,90	135,99	3139,13	146,27	4,81	323,52	214,85	13,60	0,00	62,56	61,93	61,62
2012	A	080402	National sea traffic		Kerosene	0,00											
2012	A	080402	National sea traffic		LPG	0,01	0,00	12,12	3,74	0,20	4,30	0,61	0,00	0,00	0,00	0,00	0,00
2012	A	080402	National sea traffic		Residual oil	2,35	1150,72	4532,21	148,53	4,59	490,00	183,55	11,51		103,50	102,47	101,95
2012	A	080403	Fishing		Diesel	6,47	302,86	8670,24	373,24	11,54	1231,32	478,48	30,29	0,00	139,32	137,93	137,23
2012	A	080403	Fishing		Gasoline	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
2012	A	080403	Fishing		Kerosene	0,00											
2012	A	080403	Fishing		LPG	0,01	0,00	12,12	3,74	0,20	4,30	0,61	0,00	0,00	0,00	0,00	0,00
2012	A	080404	International sea traffic		Diesel	10,31	483,08	16353,79	595,16	18,41	1963,42	763,22	48,31		222,23	220,01	218,90
2012	A	080404	International sea traffic		Residual oil	9,51	4652,66	20170,41	603,83	18,68	1992,00	742,15	46,53		418,48	414,30	412,21
2012	A	080501	Air traffic, Dom. < 3000 ft.		AvGas	0,06	1,42	53,48	77,36	1,36	434,06	4,54	0,12	0,10	0,62	0,62	0,62
2012	A	080501	Air traffic, Dom. < 3000 ft.		Jet fuel	0,23	5,32	71,62	3,80	0,42	34,07	16,67	2,14		0,27	0,27	0,27
2012	A	080502	Air traffic, Int. < 3000 ft.		AvGas	0,00	0,02	0,68	0,98	0,02	5,49	0,06	0,00	0,00	0,01	0,01	0,01
2012	A	080502	Air traffic, Int. < 3000 ft.		Jet fuel	0,33	7,67	99,58	12,73	1,41	61,55	24,01	2,42		0,39	0,39	0,39
2012	A	080503	Air traffic, Dom. > 3000 ft.		Jet fuel	0,49	11,22	135,75	3,34	0,00	46,48	35,15	1,12		0,57	0,57	0,57
2012	A	080504	Air traffic, Int. > 3000 ft.		Jet fuel	3,54	81,38	845,12	21,27	0,00	170,61	254,88	8,14		4,11	4,11	4,11
2012	A	080600	Agriculture		Diesel	17,64	8,26	9364,59	864,10	14,05	5446,35	1305,62	56,21	3,21	692,88	692,88	692,88

2012	A	080600	Agriculture	Gasoline	0,52	0,24	57,46	614,21	82,78	11272,32	37,87	0,89	0,79	16,09	16,09	16,09
2012	A	080700	Forestry	Diesel	0,16	0,07	55,24	4,11	0,07	35,91	11,75	0,51	0,03	3,72	3,72	3,72
2012	A	080700	Forestry	Gasoline	0,07	0,03	3,84	277,81	2,17	1255,55	5,12	0,03	0,01	5,76	5,76	5,76
2012	A	080800	Industry	Diesel	12,82	6,00	6270,79	694,64	11,29	3931,91	948,53	39,70	2,27	609,10	609,10	609,10
2012	A	080800	Industry	Gasoline	0,16	0,07	33,14	238,85	17,01	2204,59	11,41	0,23	0,02	3,20	3,20	3,20
2012	A	080800	Industry	LPG	0,97	0,00	1293,53	142,29	7,49	102,12	61,46	3,40	0,20	4,77	4,77	4,77
2012	A	080900	Household and gardening	Gasoline	0,86	0,39	92,15	1953,14	64,53	26236,07	62,43	1,08	0,08	14,71	14,71	14,71
2012	A	081100	Commercial and institutional	Gasoline	2,35	1,07	219,03	3636,05	151,26	72586,61	171,41	2,65	0,21	66,99	66,99	66,99
2012	P	080501	Air traffic, Dom. < 3000 ft.	AvGas	0,00	0,01	0,43	0,62	0,01	3,48	0,04	0,00	0,00	0,00	0,00	0,00
2012	P	080501	Air traffic, Dom. < 3000 ft.	Jet fuel	0,25	5,83	76,58	4,71	0,52	48,58	18,25	1,44		0,29	0,29	0,29
2012	P	080502	Air traffic, Int. < 3000 ft.	AvGas	0,00	0,00	0,10	0,14	0,00	0,80	0,01	0,00	0,00	0,00	0,00	0,00
2012	P	080502	Air traffic, Int. < 3000 ft.	Jet fuel	2,76	63,36	935,33	93,17	10,35	654,21	198,45	10,67	0,00	3,20	3,20	3,20
2012	P	080503	Air traffic, Dom. > 3000 ft.	Jet fuel	0,80	18,51	241,03	7,75	0,00	40,42	57,96	1,85	0,00	0,93	0,93	0,93
2012	P	080504	Air traffic, Int. > 3000 ft.	Jet fuel	28,22	648,83	8747,67	251,07	0,00	883,26	2032,12	64,88	0,00	32,74	32,74	32,74

2012 emissions for Arsenic, Cadmium, Chromium, Copper, Mercury, Nickel, Lead, Selenium and Zinc.

Year	SNAP ID	Category	Fuel type	Fuel PJ	Arsenic kg	Cadmium kg	Chromium kg	Copper kg	Mercury kg	Nickel kg	Lead kg	Selenium kg	Zinc kg
2012	A 070101	Passenger cars	Highway	Bio ethanol	0,51	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
2012	A 070101	Passenger cars	Highway	Biodiesel	0,74	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
2012	A 070101	Passenger cars	Highway	Diesel	11,03	0,03	3,45	10,78	7,65	1,37	3,48	20,72	0,03 691,14
2012	A 070101	Passenger cars	Highway	Gasoline	13,94	0,10	4,63	6,11	17,41	2,77	5,30	14,07	0,06 923,33
2012	A 070101	Passenger cars	Highway	LPG	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,01
2012	A 070102	Passenger cars	Rural	Bio ethanol	0,83	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
2012	A 070102	Passenger cars	Rural	Biodiesel	1,19	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
2012	A 070102	Passenger cars	Rural	Diesel	17,79	0,04	5,97	18,40	13,08	2,21	6,03	35,88	0,04 1196,48
2012	A 070102	Passenger cars	Rural	Gasoline	22,81	0,16	8,01	10,40	30,01	4,53	9,10	24,34	0,10 1598,17
2012	A 070102	Passenger cars	Rural	LPG	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,02
2012	A 070103	Passenger cars	Urban	Bio ethanol	0,57	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
2012	A 070103	Passenger cars	Urban	Biodiesel	0,70	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
2012	A 070103	Passenger cars	Urban	Diesel	10,40	0,02	2,36	7,94	5,61	1,29	2,40	14,21	0,02 473,89
2012	A 070103	Passenger cars	Urban	Gasoline	15,59	0,11	3,19	5,05	12,53	3,10	3,94	9,79	0,07 636,04
2012	A 070103	Passenger cars	Urban	LPG	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,01
2012	A 070201	Light duty vehicles	Highway	Bio ethanol	0,01	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
2012	A 070201	Light duty vehicles	Highway	Biodiesel	0,38	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
2012	A 070201	Light duty vehicles	Highway	Diesel	5,66	0,01	1,35	4,49	3,18	0,70	1,37	8,14	0,01 271,39
2012	A 070201	Light duty vehicles	Highway	Gasoline	0,34	0,00	0,08	0,12	0,33	0,07	0,10	0,26	0,00 16,92
2012	A 070201	Light duty vehicles	Highway	LPG	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,01
2012	A 070202	Light duty vehicles	Rural	Bio ethanol	0,02	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
2012	A 070202	Light duty vehicles	Rural	Biodiesel	0,64	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
2012	A 070202	Light duty vehicles	Rural	Diesel	9,54	0,02	2,49	8,10	5,74	1,18	2,52	14,99	0,02 499,91
2012	A 070202	Light duty vehicles	Rural	Gasoline	0,65	0,00	0,16	0,23	0,61	0,13	0,19	0,48	0,00 31,23
2012	A 070202	Light duty vehicles	Rural	LPG	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,02
2012	A 070203	Light duty vehicles	Urban	Bio ethanol	0,02	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
2012	A 070203	Light duty vehicles	Urban	Biodiesel	0,40	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
2012	A 070203	Light duty vehicles	Urban	Diesel	5,93	0,01	1,00	3,66	2,58	0,74	1,02	6,02	0,01 200,76
2012	A 070203	Light duty vehicles	Urban	Gasoline	0,52	0,00	0,06	0,13	0,27	0,10	0,09	0,20	0,00 12,68
2012	A 070203	Light duty vehicles	Urban	LPG	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,01
2012	A 070301	Heavy duty vehicles	Highway	Bio ethanol	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
2012	A 070301	Heavy duty vehicles	Highway	Biodiesel	0,96	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
2012	A 070301	Heavy duty vehicles	Highway	Diesel	14,33	0,03	2,28	8,50	5,98	1,78	2,33	13,73	0,03 458,05
2012	A 070301	Heavy duty vehicles	Highway	Gasoline	0,02	0,00	0,00	0,01	0,02	0,00	0,01	0,01	0,00 0,85

2012	A	070302	Heavy duty vehicles	Rural	Biodiesel	1,14	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
2012	A	070302	Heavy duty vehicles	Rural	Diesel	17,07	0,04	2,60	9,85	6,93	2,12	2,66	15,70	0,04 523,71
2012	A	070302	Heavy duty vehicles	Rural	Gasoline	0,04	0,00	0,01	0,01	0,04	0,01	0,01	0,03	0,00 2,17
2012	A	070303	Heavy duty vehicles	Urban	Diesel	5,91	0,01	0,78	3,11	2,18	0,73	0,80	4,72	0,01 157,47
2012	A	070303	Heavy duty vehicles	Urban	Gasoline	0,02	0,00	0,01	0,01	0,02	0,00	0,01	0,02	0,00 1,00
2012	A	070400	Mopeds	Urban	Bio ethanol	0,01	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00 0,00
2012	A	070501	Motorcycles	Highway	Gasoline	0,11	0,00	0,01	0,03	0,06	0,02	0,02	0,04	0,00 2,86
2012	A	070502	Motorcycles	Rural	Bio ethanol	0,01	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00 0,00
2012	A	070502	Motorcycles	Rural	Gasoline	0,23	0,00	0,04	0,07	0,15	0,05	0,05	0,12	0,00 7,51
2012	A	070503	Motorcycles	Urban	Bio ethanol	0,01	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00 0,00
2012	A	070503	Motorcycles	Urban	Gasoline	0,28	0,00	0,04	0,08	0,18	0,06	0,06	0,14	0,00 8,73
2012	A	080100	Military		AvGas	0,00	0,00	0,00	0,00	0,00	0,00	0,00	47,71	0,00 0,19
2012	A	080100	Military		Diesel	0,88	0,00	0,19	0,63	0,45	0,10	0,19	1,13	0,00 37,80
2012	A	080100	Military		Gasoline	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00 0,09
2012	A	080100	Military		Jet fuel	0,70	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00 0,00
2012	A	080200	Railways		Diesel	3,37	0,01	0,63	2,22	1,57	0,42	0,64	3,77	0,01 125,66
2012	A	080200	Railways		Gasoline	0,00		0,00	0,00	0,00		0,00	0,00	0,00 0,00
2012	A	080200	Railways		Kerosene	0,00								
2012	A	080300	Inland waterways		Diesel	1,00	0,00	0,19	0,66	0,47	0,12	0,19	1,12	0,00 37,38
2012	A	080300	Inland waterways		Gasoline	0,34	0,00	0,08	0,12	0,33	0,07	0,10	0,26	0,00 16,91
2012	A	080402	National sea traffic		Diesel	2,90	3,40	0,67	2,73	3,40	3,40	4,76	6,79	13,59 34,00
2012	A	080402	National sea traffic		Kerosene	0,00								
2012	A	080402	National sea traffic		LPG	0,01								
2012	A	080402	National sea traffic		Residual oil	2,35	28,76	1,72	11,51	28,76	1,15	1726,08	11,51	23,01 51,77
2012	A	080403	Fishing		Diesel	6,47	7,57	1,49	6,08	7,57	7,57	10,60	15,13	30,26 75,72
2012	A	080403	Fishing		Gasoline	0,00		0,00	0,00	0,00		0,00	0,00	0,00 0,00
2012	A	080403	Fishing		Kerosene	0,00								
2012	A	080403	Fishing		LPG	0,01								
2012	A	080404	International sea traffic		Diesel	10,31	12,07	2,37	9,69	12,07	12,07	16,91	24,13	48,27 120,77
2012	A	080404	International sea traffic		Residual oil	9,51	116,27	6,95	46,53	116,27	4,66	6979,03	46,53	93,05 209,32
2012	A	080501	Air traffic, Dom. < 3000 ft.		AvGas	0,06	0,00	0,02	0,02	0,06	0,01	0,02	840,83	0,00 3,14
2012	A	080501	Air traffic, Dom. < 3000 ft.		Jet fuel	0,23		0,00	0,00	0,00		0,00	0,00	0,00 0,00
2012	A	080502	Air traffic, Int. < 3000 ft.		AvGas	0,00	0,00	0,00	0,00	0,00	0,00	0,00	10,63	0,00 0,04
2012	A	080502	Air traffic, Int. < 3000 ft.		Jet fuel	0,33		0,00	0,00	0,00		0,00	0,00	0,00 0,00
2012	A	080503	Air traffic, Dom. > 3000 ft.		Jet fuel	0,49		0,00	0,00	0,00		0,00	0,00	0,00 0,00
2012	A	080504	Air traffic, Int. > 3000 ft.		Jet fuel	3,54		0,00	0,00	0,00		0,00	0,00	0,00 0,00
2012	A	080600	Agriculture		Diesel	17,64	0,04	3,27	11,64	8,21	2,19	3,34	19,72	0,04 658,02

2012	A	080600	Agriculture	Gasoline	0,52	0,00	0,13	0,19	0,50	0,10	0,16	0,40	0,00	26,17
2012	A	080700	Forestry	Diesel	0,16	0,00	0,03	0,10	0,07	0,02	0,03	0,18	0,00	5,92
2012	A	080700	Forestry	Gasoline	0,07	0,00	0,02	0,03	0,07	0,01	0,02	0,05	0,00	3,54
2012	A	080800	Industry	Diesel	12,82	0,03	2,38	8,46	5,96	1,59	2,42	14,33	0,03	478,05
2012	A	080800	Industry	Gasoline	0,16	0,00	0,04	0,06	0,15	0,03	0,05	0,12	0,00	7,89
2012	A	080800	Industry	LPG	0,97	0,00	0,13	0,11	0,45	0,00	0,13	0,38	0,00	25,45
2012	A	080900	Household and gardening	Gasoline	0,86	0,01	0,22	0,31	0,83	0,17	0,26	0,66	0,00	43,15
2012	A	081100	Commercial and institutional	Gasoline	2,35	0,02	0,59	0,86	2,28	0,46	0,71	1,81	0,01	118,46
2012	P	080501	Air traffic, Dom. < 3000 ft.	AvGas	0,00	0,00	0,00	0,00	0,00	0,00	0,00	6,73	0,00	0,03
2012	P	080501	Air traffic, Dom. < 3000 ft.	Jet fuel	0,25		0,00	0,00	0,00		0,00		0,00	0,00
2012	P	080502	Air traffic, Int. < 3000 ft.	AvGas	0,00	0,00	0,00	0,00	0,00	0,00	0,00	1,55	0,00	0,01
2012	P	080502	Air traffic, Int. < 3000 ft.	Jet fuel	2,76	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
2012	P	080503	Air traffic, Dom. > 3000 ft.	Jet fuel	0,80	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
2012	P	080504	Air traffic, Int. > 3000 ft.	Jet fuel	28,22	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00

2012 emissions for Dioxins/, Flouranthene, Benzo(b), Benzo(k), Benzo(a), Benzo(g,h,i), indeno(1,2,3-c,d), HCB and PCB.

Year	SNAP ID	Category	Fuel type	Fuel	Dioxins/	Flouranthene	Benzo(b)	Benzo(k)	Benzo(a)	Benzo(g,h,i)	indeno(1,2,3-c,d)	HCB	PCB		
				PJ	g	kg	kg	kg	kg	kg	kg	g	g		
2012	A	070101	Passenger cars	Highway	Bio ethanol	0,51	0,00	0,58	0,11	0,13	0,10	0,21	0,15	0,00	0,00
2012	A	070101	Passenger cars	Highway	Biodiesel	0,74	0,00	9,44	0,58	0,52	0,63	1,23	0,59	0,00	0,02
2012	A	070101	Passenger cars	Highway	Diesel	11,03	0,00	141,29	8,62	7,82	9,44	18,34	8,90	0,07	0,23
2012	A	070101	Passenger cars	Highway	Gasoline	13,94	0,01	15,85	2,91	3,51	2,87	5,81	4,17	0,00	0,01
2012	A	070101	Passenger cars	Highway	LPG	0,00								0,00	0,00
2012	A	070102	Passenger cars	Rural	Bio ethanol	0,83	0,00	1,04	0,19	0,23	0,19	0,39	0,28	0,00	0,00
2012	A	070102	Passenger cars	Rural	Biodiesel	1,19	0,00	17,34	1,06	0,96	1,16	2,25	1,09	0,01	0,03
2012	A	070102	Passenger cars	Rural	Diesel	17,79	0,01	259,60	15,85	14,36	17,34	33,69	16,35	0,11	0,38
2012	A	070102	Passenger cars	Rural	Gasoline	22,81	0,01	28,53	5,28	6,40	5,22	10,57	7,61	0,00	0,01
2012	A	070102	Passenger cars	Rural	LPG	0,00								0,00	0,00
2012	A	070103	Passenger cars	Urban	Bio ethanol	0,57	0,00	0,41	0,07	0,09	0,07	0,15	0,10	0,00	0,00
2012	A	070103	Passenger cars	Urban	Biodiesel	0,70	0,00	6,73	0,41	0,37	0,45	0,87	0,42	0,00	0,01
2012	A	070103	Passenger cars	Urban	Diesel	10,40	0,01	100,73	6,15	5,57	6,73	13,07	6,34	0,06	0,22
2012	A	070103	Passenger cars	Urban	Gasoline	15,59	0,00	11,23	2,00	2,41	1,97	4,01	2,86	0,00	0,01
2012	A	070103	Passenger cars	Urban	LPG	0,00								0,00	0,00
2012	A	070201	Light duty vehicles	Highway	Bio ethanol	0,01	0,00	0,01	0,00	0,00	0,00	0,00	0,00	0,00	0,00
2012	A	070201	Light duty vehicles	Highway	Biodiesel	0,38	0,00	3,49	0,21	0,19	0,23	0,45	0,22	0,00	0,01
2012	A	070201	Light duty vehicles	Highway	Diesel	5,66	0,00	52,26	3,19	2,89	3,49	6,78	3,29	0,03	0,12
2012	A	070201	Light duty vehicles	Highway	Gasoline	0,34	0,00	0,34	0,05	0,06	0,05	0,11	0,07	0,00	0,00
2012	A	070201	Light duty vehicles	Highway	LPG	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
2012	A	070202	Light duty vehicles	Rural	Bio ethanol	0,02	0,00	0,02	0,00	0,00	0,00	0,01	0,00	0,00	0,00
2012	A	070202	Light duty vehicles	Rural	Biodiesel	0,64	0,00	6,44	0,39	0,36	0,43	0,84	0,41	0,00	0,01
2012	A	070202	Light duty vehicles	Rural	Diesel	9,54	0,01	96,33	5,88	5,33	6,43	12,50	6,06	0,06	0,20
2012	A	070202	Light duty vehicles	Rural	Gasoline	0,65	0,00	0,64	0,10	0,11	0,10	0,20	0,13	0,00	0,00
2012	A	070202	Light duty vehicles	Rural	LPG	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
2012	A	070203	Light duty vehicles	Urban	Bio ethanol	0,02	0,00	0,01	0,00	0,00	0,00	0,00	0,00	0,00	0,00
2012	A	070203	Light duty vehicles	Urban	Biodiesel	0,40	0,00	2,88	0,18	0,16	0,19	0,37	0,18	0,00	0,01
2012	A	070203	Light duty vehicles	Urban	Diesel	5,93	0,00	43,08	2,63	2,38	2,88	5,59	2,71	0,04	0,13
2012	A	070203	Light duty vehicles	Urban	Gasoline	0,52	0,00	0,29	0,05	0,05	0,04	0,09	0,06	0,00	0,00
2012	A	070203	Light duty vehicles	Urban	LPG	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
2012	A	070301	Heavy duty vehicles	Highway	Bio ethanol	0,00								0,00	0,00
2012	A	070301	Heavy duty vehicles	Highway	Biodiesel	0,96	0,00	1,94	0,49	0,73	0,09	0,07	0,13	0,01	0,50
2012	A	070301	Heavy duty vehicles	Highway	Diesel	14,33	0,01	29,11	7,34	10,88	1,36	1,09	1,90	0,09	7,54
2012	A	070301	Heavy duty vehicles	Highway	Gasoline	0,02								0,00	0,00

2012	A	070302	Heavy duty vehicles	Rural	Biodiesel	1,14	0,00	2,36	0,59	0,88	0,11	0,09	0,15	0,01	0,60
2012	A	070302	Heavy duty vehicles	Rural	Diesel	17,07	0,02	35,27	8,90	13,18	1,65	1,32	2,31	0,10	8,98
2012	A	070302	Heavy duty vehicles	Rural	Gasoline	0,04								0,00	0,00
2012	A	070303	Heavy duty vehicles	Urban	Diesel	5,91	0,01	9,91	2,50	3,70	0,46	0,37	0,65	0,04	3,11
2012	A	070303	Heavy duty vehicles	Urban	Gasoline	0,02								0,00	0,00
2012	A	070400	Mopeds	Urban	Bio ethanol	0,01								0,00	0,00
2012	A	070501	Motorcycles	Highway	Gasoline	0,11	0,00	1,37	0,09	0,07	0,08	0,18	0,07	0,00	0,00
2012	A	070502	Motorcycles	Rural	Bio ethanol	0,01	0,00	0,13	0,01	0,01	0,01	0,02	0,01	0,00	0,00
2012	A	070502	Motorcycles	Rural	Gasoline	0,23	0,01	3,57	0,23	0,18	0,20	0,46	0,18	0,00	0,00
2012	A	070503	Motorcycles	Urban	Bio ethanol	0,01	0,00	0,16	0,01	0,01	0,01	0,02	0,01	0,00	0,00
2012	A	070503	Motorcycles	Urban	Gasoline	0,28	0,01	4,30	0,28	0,22	0,24	0,56	0,22	0,00	0,00
2012	A	080100	Military		AvGas	0,00	0,00	0,02	0,00	0,00	0,00	0,00	0,00	0,00	0,00
2012	A	080100	Military		Diesel	0,88	0,00	3,83	0,45	0,44	0,23	0,41	0,23	0,01	0,19
2012	A	080100	Military		Gasoline	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
2012	A	080100	Military		Jet fuel	0,70	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
2012	A	080200	Railways		Diesel	3,37	0,00	4,75	1,21	1,35	0,20	0,17	0,31	0,02	1,77
2012	A	080200	Railways		Gasoline	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
2012	A	080200	Railways		Kerosene	0,00								0,00	0,00
2012	A	080300	Inland waterways		Diesel	1,00	0,00	4,36	0,51	0,50	0,26	0,47	0,26	0,01	0,53
2012	A	080300	Inland waterways		Gasoline	0,34	0,00	1,45	0,07	0,02	0,04	0,23	0,08	0,00	0,00
2012	A	080402	National sea traffic		Diesel	2,90	0,03	21,54	1,86	0,87	0,44	4,15	3,43	0,01	0,03
2012	A	080402	National sea traffic		Kerosene	0,00									
2012	A	080402	National sea traffic		LPG	0,01								0,00	0,00
2012	A	080402	National sea traffic		Residual oil	2,35	0,03	12,21	0,64	0,12	0,05	0,16	0,07	0,01	0,03
2012	A	080403	Fishing		Diesel	6,47	0,08	47,98	4,14	1,94	0,97	9,25	7,63	0,01	0,06
2012	A	080403	Fishing		Gasoline	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
2012	A	080403	Fishing		Kerosene	0,00									
2012	A	080403	Fishing		LPG	0,01								0,00	0,00
2012	A	080404	International sea traffic		Diesel	10,31	0,12	76,53	6,60	3,09	1,55	14,75	12,17	0,02	0,09
2012	A	080404	International sea traffic		Residual oil	9,51	0,13	39,20	1,90	0,86	0,67	2,47	1,90	0,03	0,13
2012	A	080501	Air traffic, Dom. < 3000 ft.		AvGas	0,06	0,00	0,27	0,01	0,00	0,01	0,04	0,02	0,00	0,00
2012	A	080501	Air traffic, Dom. < 3000 ft.		Jet fuel	0,23	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
2012	A	080502	Air traffic, Int. < 3000 ft.		AvGas	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
2012	A	080502	Air traffic, Int. < 3000 ft.		Jet fuel	0,33	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
2012	A	080503	Air traffic, Dom. > 3000 ft.		Jet fuel	0,49	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
2012	A	080504	Air traffic, Int. > 3000 ft.		Jet fuel	3,54	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
2012	A	080600	Agriculture		Diesel	17,64	0,01	76,75	9,01	8,75	4,51	8,19	4,66	0,11	9,28

2012	A	080600	Agriculture	Gasoline	0,52	0,00	2,25	0,11	0,04	0,06	0,36	0,13	0,00	0,00
2012	A	080700	Forestry	Diesel	0,16	0,00	0,69	0,08	0,08	0,04	0,07	0,04	0,00	0,08
2012	A	080700	Forestry	Gasoline	0,07	0,00	0,30	0,01	0,00	0,01	0,05	0,02	0,00	0,00
2012	A	080800	Industry	Diesel	12,82	0,01	55,76	6,54	6,36	3,28	5,95	3,39	0,08	6,74
2012	A	080800	Industry	Gasoline	0,16	0,00	0,68	0,03	0,01	0,02	0,11	0,04	0,00	0,00
2012	A	080800	Industry	LPG	0,97								0,00	0,00
2012	A	080900	Household and gardening	Gasoline	0,86	0,00	3,70	0,18	0,06	0,10	0,59	0,21	0,00	0,00
2012	A	081100	Commercial and institutional	Gasoline	2,35	0,01	10,16	0,49	0,17	0,27	1,62	0,57	0,00	0,00
2012	P	080501	Air traffic, Dom. < 3000 ft.	AvGas	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00		
2012	P	080501	Air traffic, Dom. < 3000 ft.	Jet fuel	0,25	0,00	0,00	0,00	0,00	0,00	0,00	0,00		
2012	P	080502	Air traffic, Int. < 3000 ft.	AvGas	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	
2012	P	080502	Air traffic, Int. < 3000 ft.	Jet fuel	2,76	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	
2012	P	080503	Air traffic, Dom. > 3000 ft.	Jet fuel	0,80	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	
2012	P	080504	Air traffic, Int. > 3000 ft.	Jet fuel	28,22	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	

Non-exhaust emission factors, activity data and total non-exhaust emissions of TSP, PM₁₀ and PM_{2.5} in 2012.

Year	Source	Category	Mileage kmkveh	TSP mg pr km	PM ₁₀ mg pr km	PM _{2.5} mg pr km	As µg pr km	Cd µg pr km	Cr µg pr km	Cu µg pr km	Hg µg pr km	Ni µg pr km	Pb µg pr km	Se µg pr km	Zn µg pr km
2012	Brake wear	1	43734320	6,2	6,1	2,4	0,1	0,1	0,7	649,8	0,0	0,7	85,7	0,1	124,7
2012	Brake wear	2	8369840	11,9	11,7	4,7	0,1	0,1	1,3	1254,9	0,0	1,3	165,5	0,2	240,8
2012	Brake wear	3	3155814	29,8	29,2	11,6	0,3	0,1	4,9	224,3	0,0	3,4	12,1	0,6	223,5
2012	Brake wear	4	766075	47,5	46,5	18,5	0,5	0,1	3,0	650,2	0,0	7,5	34,7	0,9	443,3
2012	Brake wear	5	222709	6,2	6,1	2,4	0,1	0,1	0,7	649,2	0,0	0,7	85,6	0,1	124,6
2012	Brake wear	6	475124	4,2	4,1	1,6	0,0	0,0	0,4	439,2	0,0	0,4	57,9	0,1	84,3
2012	Road abrasion	1	43734320	15,0	7,5	4,1	0,0	0,0	0,3	0,1	0,0	0,2	0,7	0,0	1,1
2012	Road abrasion	2	8369840	15,0	7,5	4,1	0,0	0,0	0,3	0,1	0,0	0,2	0,7	0,0	1,1
2012	Road abrasion	3	3155814	76,0	38,0	20,5	0,0	0,0	1,5	0,8	0,0	1,2	3,6	0,0	5,7
2012	Road abrasion	4	766075	76,0	38,0	20,5	0,0	0,0	1,5	0,8	0,0	1,2	3,6	0,0	5,7
2012	Road abrasion	5	222709	6,0	3,0	1,6	0,0	0,0	0,1	0,1	0,0	0,1	0,3	0,0	0,5
2012	Road abrasion	6	475124	6,0	3,0	1,6	0,0	0,0	0,1	0,1	0,0	0,1	0,3	0,0	0,5
2012	Tyre wear	1	43734320	10,8	6,5	4,6	0,0	0,0	0,0	0,2	0,0	0,3	0,9	0,2	118,5
2012	Tyre wear	2	8369840	17,2	10,3	7,2	0,0	0,0	0,1	0,3	0,0	0,4	1,4	0,3	187,7
2012	Tyre wear	3	3155814	65,5	39,3	27,5	0,1	0,2	0,2	1,0	0,0	1,7	5,3	1,3	716,8
2012	Tyre wear	4	766075	61,5	36,9	25,8	0,0	0,2	0,2	1,0	0,0	1,6	4,9	1,2	672,1
2012	Tyre wear	5	222709	14,2	8,5	6,0	0,0	0,0	0,1	0,2	0,0	0,4	1,1	0,3	154,9
2012	Tyre wear	6	475124	17,8	10,7	7,5	0,0	0,0	0,1	0,3	0,0	0,5	1,4	0,4	194,1
2012	Total	1	43734320	32,0	20,1	11,0	0,1	0,1	1,0	650,1	0,0	1,2	87,3	0,3	244,3
2012	Total	2	8369840	44,1	29,5	15,9	0,1	0,2	1,6	1255,3	0,0	1,9	167,5	0,6	429,7
2012	Total	3	3155814	171,3	106,5	59,7	0,3	0,3	6,6	226,1	0,0	6,3	21,0	1,9	946,1
2012	Total	4	766075	184,9	121,4	64,8	0,5	0,3	4,8	651,9	0,0	10,3	43,2	2,2	1121,1
2012	Total	5	222709	26,3	17,6	10,0	0,1	0,1	0,8	649,5	0,0	1,1	87,0	0,4	280,0
2012	Total	6	475124	27,9	17,7	10,7	0,1	0,1	0,6	439,6	0,0	1,0	59,6	0,4	278,9

Year	Source	Category	TSP tonnes	PM ₁₀ tonnes	PM _{2.5} tonnes	As kg	Cd kg	Cr kg	Cu kg	Hg kg	Ni kg	Pb kg	Se kg	Zn kg
2012	Brake wear	1	270	265	105	2,705	2,653	28,453	28417,620		28,480	3746,887	5,409	5453,656
2012	Brake wear	2	100	98	39	1,000	0,981	10,517	10503,469		10,527	1384,891	1,999	2015,732
2012	Brake wear	3	94	92	37	0,939	0,288	15,398	707,779		10,703	38,213	1,878	705,436
2012	Brake wear	4	36	36	14	0,364	0,110	2,329	498,107		5,783	26,571	0,727	339,585
2012	Brake wear	5	1	1	1	0,014	0,013	0,145	144,588		0,145	19,064	0,028	27,748
2012	Brake wear	6	2	2	1	0,020	0,019	0,209	208,685		0,209	27,515	0,040	40,049
2012	Road abrasion	1	656	328	177	0,000	0,062	13,019	6,554	0,037	10,414	30,849	0,000	49,568
2012	Road abrasion	2	126	63	34	0,000	0,012	2,491	1,254	0,007	1,993	5,904	0,000	9,486
2012	Road abrasion	3	240	120	65	0,000	0,023	4,760	2,396	0,014	3,807	11,279	0,000	18,122
2012	Road abrasion	4	58	29	16	0,000	0,006	1,155	0,582	0,003	0,924	2,738	0,000	4,399
2012	Road abrasion	5	1	1	0	0,000	0,000	0,027	0,013	0,000	0,021	0,063	0,000	0,101
2012	Road abrasion	6	3	1	1	0,000	0,000	0,057	0,028	0,000	0,045	0,134	0,000	0,215
2012	Tyre wear	1	474	284	199	0,379	1,232	1,706	7,391		12,082	38,140	9,476	5181,351
2012	Tyre wear	2	144	86	60	0,115	0,374	0,517	2,241		3,664	11,566	2,874	1571,309
2012	Tyre wear	3	207	124	87	0,165	0,538	0,745	3,227		5,275	16,652	4,137	2262,229
2012	Tyre wear	4	47	28	20	0,038	0,122	0,169	0,734		1,201	3,790	0,942	514,867
2012	Tyre wear	5	3	2	1	0,003	0,008	0,011	0,049		0,080	0,254	0,063	34,503
2012	Tyre wear	6	8	5	4	0,007	0,022	0,030	0,132		0,215	0,679	0,169	92,228
2012	Total	1	1400	877	482	3,084	3,947	43,177	28431,564	0,037	50,976	3815,876	14,885	10684,575
2012	Total	2	369	247	133	1,115	1,366	13,525	10506,965	0,007	16,183	1402,361	4,873	3596,528
2012	Total	3	541	336	188	1,104	0,849	20,902	713,402	0,014	19,786	66,144	6,015	2985,788
2012	Total	4	142	93	50	0,401	0,238	3,654	499,423	0,003	7,908	33,099	1,669	858,851
2012	Total	5	6	4	2	0,016	0,022	0,183	144,650	0,000	0,247	19,381	0,091	62,352
2012	Total	6	13	8	5	0,027	0,042	0,296	208,845	0,000	0,469	28,328	0,208	132,493

Annex 3B-16 Fuel consumption and emissions in CRF format

Fuel

IPCC ID	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
Industry-Other (1A2f)	11,7	11,7	11,6	11,6	11,6	11,5	11,5	11,5	11,5	11,5	11,6	11,7	11,7	11,9	11,9
Civil Aviation (1A3a)	3,6	3,3	3,7	3,8	3,6	3,4	2,8	2,7	2,6	2,7	2,8	2,8	2,9	2,7	2,4
Road (1A3b)	111,2	117,5	117,7	118,5	119,7	126,4	132,0	134,4	136,2	142,9	144,2	146,6	149,5	152,1	154,0
Railways (1A3c)	4,9	4,9	4,4	4,6	4,2	4,0	4,1	4,3	4,5	4,1	4,1	4,1	4,0	3,3	3,1
Navigation (1A3d)	10,4	10,3	10,4	10,4	10,5	10,5	10,6	10,9	10,7	10,8	11,3	12,2	12,0	10,0	8,8
Comm./Inst. (1A4a)	1,0	1,0	1,0	1,0	1,0	1,0	1,0	1,0	1,0	1,0	1,1	1,1	1,1	1,1	1,2
Residential (1A4b)	0,5	0,5	0,5	0,5	0,5	0,5	0,5	0,5	0,5	0,5	0,5	0,5	0,6	0,6	0,6
Ag./for./fish. (1A4c)	24,4	26,0	23,8	25,5	25,3	25,7	25,7	24,3	23,8	22,9	23,4	22,2	21,0	20,4	21,1
Military (1A5)	5,5	4,3	5,0	2,7	2,3	1,6	3,9	1,9	3,3	3,5	3,4	2,4	2,3	2,8	2,5
Navigation int. (1A3d)	16,2	19,0	28,4	36,2	37,1	39,1	34,9	36,7	55,0	62,0	65,1	62,0	56,7	57,2	53,3
Civil Aviation int. (1A3a)	19,3	20,9	22,4	24,0	25,1	24,1	22,7	23,5	23,0	25,2	25,9	27,4	27,9	30,0	31,8

IPCC ID	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
Industry-Other (1A2f)	12,0	12,1	12,3	12,4	12,5	13,0	14,0	15,1	15,4	11,4	14,4	14,0	13,9
Civil Aviation (1A3a)	2,1	2,3	2,0	1,9	1,8	1,9	2,0	2,2	2,2	2,1	2,2	2,0	1,8
Road (1A3b)	152,5	152,8	154,5	160,6	164,8	166,1	171,3	179,5	175,1	165,6	165,2	165,2	160,9
Railways (1A3c)	3,1	2,9	2,8	3,0	2,9	3,1	3,1	3,1	3,2	3,1	3,3	3,4	3,4
Navigation (1A3d)	7,9	7,9	7,7	7,7	7,9	7,8	7,8	7,8	7,9	7,9	7,9	7,4	6,6
Comm./Inst. (1A4a)	1,2	1,3	1,5	1,8	2,0	2,2	2,4	2,4	2,4	2,4	2,4	2,3	2,3
Residential (1A4b)	0,6	0,6	0,7	0,7	0,8	0,8	0,8	0,9	0,9	0,9	0,9	0,9	0,9
Ag./for./fish. (1A4c)	23,8	23,9	24,2	23,9	22,2	23,8	23,7	23,3	25,4	25,1	24,5	24,8	24,9
Military (1A5)	1,5	1,3	1,2	1,3	3,3	3,7	1,7	2,4	1,5	2,2	1,5	2,7	1,6
Navigation int. (1A3d)	52,6	43,3	35,5	37,5	30,2	30,7	40,8	42,7	36,6	19,6	27,0	27,4	19,8
Civil Aviation int. (1A3a)	32,6	33,1	28,6	29,7	34,0	35,7	35,9	36,8	36,8	32,2	33,6	34,6	34,9

Emissions

pol_name	IPCC ID	Unit	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
SO ₂	Industry-Other (1A2f)	[tonnes]	2402	1441	1440	1438	956	952	955	957	957	959	968	244	246	249	251
SO ₂	Civil Aviation (1A3a)	[tonnes]	82	77	85	86	83	77	64	62	61	63	63	65	68	62	56
SO ₂	Road (1A3b)	[tonnes]	11621	7862	7847	7857	5488	5767	5903	3820	1569	1669	1682	1721	1744	1768	1088
SO ₂	Railways (1A3c)	[tonnes]	1152	695	618	641	393	376	382	263	105	95	96	95	93	78	40
SO ₂	Navigation (1A3d)	[tonnes]	7480	7480	7484	7228	7231	6429	5111	3506	4410	4974	5588	4400	3649	2283	2051
SO ₂	Comm./Inst. (1A4a)	[tonnes]	2	2	2	2	2	2	2	2	2	2	2	2	3	3	3
SO ₂	Residential (1A4b)	[tonnes]	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
SO ₂	Ag./for./fish. (1A4c)	[tonnes]	4766	3484	3173	3073	2269	2303	2317	2186	2150	2072	2120	978	853	856	931
SO ₂	Military (1A5)	[tonnes]	408	260	193	72	70	48	206	82	76	80	80	56	54	65	47
SO ₂	Navigation int. (1A3d)	[tonnes]	17037	20752	35647	46755	47058	41317	33277	30084	58492	58965	65049	61075	55822	46756	49282
SO ₂	Civil Aviation int. (1A3a)	[tonnes]	444	480	515	551	578	554	521	541	530	580	596	629	642	689	731
NO _x	Industry-Other (1A2f)	[tonnes]	10903	10964	11011	11044	11065	11081	11282	11440	11558	11677	11882	12080	12248	12425	12262
NO _x	Civil Aviation (1A3a)	[tonnes]	1203	1132	1237	1252	1208	1123	920	902	900	940	958	971	998	911	815
NO _x	Road (1A3b)	[tonnes]	94032	99488	100077	101581	103010	109042	112371	111354	108932	108761	103616	99289	95544	91350	87478
NO _x	Railways (1A3c)	[tonnes]	6025	6063	5391	5589	5145	4913	4995	5284	5485	4971	5015	4977	4846	4089	3730
NO _x	Navigation (1A3d)	[tonnes]	13299	13339	13414	13486	13568	13649	13180	12882	12753	12999	13679	14756	13543	11175	8720
NO _x	Comm./Inst. (1A4a)	[tonnes]	66	67	68	70	70	70	75	80	85	89	93	95	98	101	102
NO _x	Residential (1A4b)	[tonnes]	31	32	33	34	34	34	36	38	40	42	43	45	46	48	49
NO _x	Ag./for./fish. (1A4c)	[tonnes]	18159	19915	18153	20143	20342	21066	21722	20824	20763	20524	21442	21138	20176	20119	21495
NO _x	Military (1A5)	[tonnes]	2285	1951	1586	1003	874	485	1755	947	1234	1223	1637	910	1117	1300	1025
NO _x	Navigation int. (1A3d)	[tonnes]	22455	26921	42068	54983	56940	60639	53939	55808	87852	99296	105113	100507	93239	92360	89143
NO _x	Civil Aviation int. (1A3a)	[tonnes]	5663	6129	6569	7035	7313	7016	6586	6846	6702	7317	7517	7904	8058	8662	9204
NM VOC	Industry-Other (1A2f)	[tonnes]	2422	2395	2368	2339	2304	2266	2231	2191	2147	2107	2088	2095	2083	2074	1997
NM VOC	Civil Aviation (1A3a)	[tonnes]	219	216	193	201	196	189	171	166	163	193	208	196	189	171	164
NM VOC	Road (1A3b)	[tonnes]	75302	75477	75187	75121	73635	77245	80067	79633	77266	74509	70081	65965	61389	56235	50185
NM VOC	Railways (1A3c)	[tonnes]	393	396	352	365	336	321	326	345	358	324	327	325	316	267	276
NM VOC	Navigation (1A3d)	[tonnes]	1560	1560	1592	1622	1654	1686	1719	1761	1786	1820	1879	1975	1969	1873	1776
NM VOC	Comm./Inst. (1A4a)	[tonnes]	2347	2333	2318	2303	2303	2303	2314	2302	2265	2285	2367	2458	2547	2636	2741
NM VOC	Residential (1A4b)	[tonnes]	1844	1833	1821	1809	1805	1801	1797	1792	1789	1785	1780	1774	1767	1759	1758
NM VOC	Ag./for./fish. (1A4c)	[tonnes]	6357	6417	6216	6284	6207	6149	5777	5298	4944	4638	4516	4208	3966	3691	3563
NM VOC	Military (1A5)	[tonnes]	564	440	169	467	299	53	158	84	120	117	143	88	99	110	104
NM VOC	Navigation int. (1A3d)	[tonnes]	825	974	1472	1892	1947	2060	1839	1928	2933	3318	3501	3343	3082	3102	2929
NM VOC	Civil Aviation int. (1A3a)	[tonnes]	279	308	334	365	385	354	330	338	330	331	367	385	390	413	424

CH ₄	Industry-Other (1A2f)	[tonnes]	63	63	62	61	61	60	58	57	56	54	53	53	53	53	51
CH ₄	Civil Aviation (1A3a)	[tonnes]	5	5	5	5	5	4	4	4	4	5	5	5	5	4	4
CH ₄	Road (1A3b)	[tonnes]	2015	2072	2088	2110	2105	2233	2329	2343	2330	2312	2230	2155	2090	2020	1912
CH ₄	Railways (1A3c)	[tonnes]	15	15	14	14	13	12	13	13	14	12	13	12	12	10	11
CH ₄	Navigation (1A3d)	[tonnes]	30	30	31	31	32	32	33	34	34	35	36	38	38	35	34
CH ₄	Comm./Inst. (1A4a)	[tonnes]	104	102	100	99	99	99	97	95	92	90	89	89	89	89	90
CH ₄	Residential (1A4b)	[tonnes]	55	54	53	52	51	51	50	49	48	48	47	46	45	45	45
CH ₄	Ag./for./fish. (1A4c)	[tonnes]	155	154	147	146	142	139	132	123	116	110	106	100	94	89	88
CH ₄	Military (1A5)	[tonnes]	28	23	16	17	13	5	17	9	12	12	17	10	12	13	11
CH ₄	Navigation int. (1A3d)	[tonnes]	26	30	46	59	60	64	57	60	91	103	108	103	95	96	91
CH ₄	Civil Aviation int. (1A3a)	[tonnes]	6	7	8	9	10	9	8	8	8	8	11	12	12	13	12
CO	Industry-Other (1A2f)	[tonnes]	9863	9784	9702	9611	9502	9379	9294	9188	9070	8956	8910	8963	8939	8907	8647
CO	Civil Aviation (1A3a)	[tonnes]	1256	1241	1118	1167	1140	1098	989	955	930	1098	1180	1117	1085	973	932
CO	Road (1A3b)	[tonnes]	573678	547717	525368	488047	455974	464930	485715	472329	460142	430439	410780	399783	362414	339585	302074
CO	Railways (1A3c)	[tonnes]	1098	1105	982	1018	937	895	910	963	999	906	914	907	883	745	717
CO	Navigation (1A3d)	[tonnes]	5472	5473	5636	5797	5962	6126	6297	6491	6623	6805	7057	7246	7150	6983	6779
CO	Comm./Inst. (1A4a)	[tonnes]	31348	30972	30583	30181	30181	30181	29610	28987	28319	27809	27575	27800	28012	28211	28817
CO	Residential (1A4b)	[tonnes]	19086	18725	18352	17968	17789	17606	17238	16880	16708	16556	16422	16311	16217	16136	16286
CO	Ag./for./fish. (1A4c)	[tonnes]	61165	59707	57256	55768	53717	51734	48771	45427	42608	39735	37673	34858	32455	29823	27820
CO	Military (1A5)	[tonnes]	4229	3121	1313	3191	1971	424	1019	518	852	874	896	621	605	684	694
CO	Navigation int. (1A3d)	[tonnes]	2722	3214	4855	6243	6424	6796	6065	6361	9677	10946	11548	11030	10168	10233	9662
CO	Civil Aviation int. (1A3a)	[tonnes]	1103	1207	1289	1416	1564	1442	1357	1399	1388	1342	1421	1502	1564	1662	1743
CO ₂	Industry-Other (1A2f)	[ktonnes]	850	849	848	846	843	839	841	841	839	839	846	851	858	865	871
CO ₂	Civil Aviation (1A3a)	[ktonnes]	256	241	268	271	262	243	199	193	190	196	199	205	212	194	174
CO ₂	Road (1A3b)	[ktonnes]	8167	8632	8644	8702	8797	9284	9699	9872	9997	10493	10589	10766	10979	11167	11313
CO ₂	Railways (1A3c)	[ktonnes]	364	366	326	338	311	297	302	319	331	300	303	301	293	247	232
CO ₂	Navigation (1A3d)	[ktonnes]	784	784	787	790	793	796	803	817	803	814	850	917	898	745	655
CO ₂	Comm./Inst. (1A4a)	[ktonnes]	74	74	74	74	74	74	74	75	75	77	78	80	81	83	85
CO ₂	Residential (1A4b)	[ktonnes]	40	40	39	39	39	39	39	39	39	39	40	40	41	41	42
CO ₂	Ag./for./fish. (1A4c)	[ktonnes]	1806	1922	1758	1887	1874	1899	1903	1794	1760	1695	1728	1642	1554	1510	1564
CO ₂	Military (1A5)	[ktonnes]	402	316	361	196	165	119	287	141	237	252	252	176	171	204	182
CO ₂	Navigation int. (1A3d)	[ktonnes]	1238	1454	2179	2786	2854	3005	2673	2797	4214	4744	4976	4725	4326	4337	4053
CO ₂	Civil Aviation int. (1A3a)	[ktonnes]	1391	1503	1613	1725	1809	1736	1632	1693	1659	1818	1867	1971	2010	2159	2290
N ₂ O	Industry-Other (1A2f)	[tonnes]	34	34	34	34	34	34	34	35	35	35	35	36	36	36	37
N ₂ O	Civil Aviation (1A3a)	[tonnes]	10	10	11	11	11	10	9	9	9	9	10	11	11	9	9

N ₂ O	Road (1A3b)	[tonnes]	261	274	274	276	279	292	308	322	330	353	368	383	401	409	417
N ₂ O	Railways (1A3c)	[tonnes]	10	10	9	9	9	8	8	9	9	8	8	8	8	7	6
N ₂ O	Navigation (1A3d)	[tonnes]	48	48	48	48	48	48	49	50	49	49	51	55	54	44	39
N ₂ O	Comm./Inst. (1A4a)	[tonnes]	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
N ₂ O	Residential (1A4b)	[tonnes]	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
N ₂ O	Ag./for./fish. (1A4c)	[tonnes]	81	87	78	85	85	87	88	83	81	79	81	77	71	70	74
N ₂ O	Military (1A5)	[tonnes]	12	9	11	6	5	4	8	4	7	8	7	5	5	6	6
N ₂ O	Navigation int. (1A3d)	[tonnes]	78	92	137	175	179	189	168	176	265	298	313	297	272	273	255
N ₂ O	Civil Aviation int. (1A3a)	[tonnes]	47	50	54	58	61	59	56	58	57	63	64	69	70	75	80
NH ₃	Industry-Other (1A2f)	[tonnes]	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2
NH ₃	Civil Aviation (1A3a)	[tonnes]	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
NH ₃	Road (1A3b)	[tonnes]	63	65	66	67	68	72	163	356	546	825	1096	1354	1703	2063	2352
NH ₃	Railways (1A3c)	[tonnes]	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
NH ₃	Navigation (1A3d)	[tonnes]	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
NH ₃	Comm./Inst. (1A4a)	[tonnes]	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
NH ₃	Residential (1A4b)	[tonnes]	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
NH ₃	Ag./for./fish. (1A4c)	[tonnes]	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3
NH ₃	Military (1A5)	[tonnes]	1	1	0	0	0	0	1	0	0	0	1	0	0	1	1
NH ₃	Navigation int. (1A3d)	[tonnes]		0						0	0						
NH ₃	Civil Aviation int. (1A3a)	[tonnes]	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
TSP	Industry-Other (1A2f)	[tonnes]	1823	1778	1733	1686	1634	1577	1533	1484	1433	1383	1349	1317	1284	1249	1193
TSP	Civil Aviation (1A3a)	[tonnes]	5	5	5	5	5	5	4	4	4	4	4	4	4	4	4
TSP	Road (1A3b)	[tonnes]	4320	4690	4721	4643	4701	4936	5097	4913	4910	5033	4849	4672	4268	3924	3611
TSP	Railways (1A3c)	[tonnes]	247	249	222	229	211	202	205	217	225	204	206	204	199	168	146
TSP	Navigation (1A3d)	[tonnes]	1099	1099	1103	1098	1103	898	710	519	660	762	919	723	670	451	417
TSP	Comm./Inst. (1A4a)	[tonnes]	24	24	24	24	24	24	24	23	22	23	24	25	27	28	29
TSP	Residential (1A4b)	[tonnes]	12	12	12	12	11	11	11	11	11	11	11	11	11	11	11
TSP	Ag./for./fish. (1A4c)	[tonnes]	2783	2820	2673	2723	2665	2628	2534	2362	2300	2119	2087	1892	1783	1633	1576
TSP	Military (1A5)	[tonnes]	101	101	50	18	26	12	114	67	64	55	117	46	73	77	47
TSP	Navigation int. (1A3d)	[tonnes]	2832	3448	5914	7810	7866	5531	4371	3999	8648	8194	10076	9968	9231	7717	8177
TSP	Civil Aviation int. (1A3a)	[tonnes]	23	24	26	28	30	28	27	28	27	29	30	32	32	35	37
PM ₁₀	Industry-Other (1A2f)	[tonnes]	1823	1778	1733	1686	1634	1577	1533	1484	1433	1383	1349	1317	1284	1249	1193
PM ₁₀	Civil Aviation (1A3a)	[tonnes]	5	5	5	5	5	5	4	4	4	4	4	4	4	4	4
PM ₁₀	Road (1A3b)	[tonnes]	4320	4690	4721	4643	4701	4936	5097	4913	4910	5033	4849	4672	4268	3924	3611
PM ₁₀	Railways (1A3c)	[tonnes]	247	249	222	229	211	202	205	217	225	204	206	204	199	168	146

PM ₁₀	Navigation (1A3d)	[tonnes]	1089	1089	1093	1088	1093	890	704	515	655	756	911	717	664	448	414
PM ₁₀	Comm./Inst. (1A4a)	[tonnes]	24	24	24	24	24	24	24	23	22	23	24	25	27	28	29
PM ₁₀	Residential (1A4b)	[tonnes]	12	12	12	12	11	11	11	11	11	11	11	11	11	11	11
PM ₁₀	Ag./for./fish. (1A4c)	[tonnes]	2781	2818	2671	2721	2663	2626	2532	2360	2298	2117	2086	1891	1782	1632	1575
PM ₁₀	Military (1A5)	[tonnes]	101	101	50	18	26	12	114	67	64	55	117	46	73	77	47
PM ₁₀	Navigation int. (1A3d)	[tonnes]	2803	3413	5855	7732	7788	5476	4327	3959	8561	8112	9975	9869	9139	7639	8095
PM ₁₀	Civil Aviation int. (1A3a)	[tonnes]	23	24	26	28	30	28	27	28	27	29	30	32	32	35	37
PM _{2.5}	Industry-Other (1A2f)	[tonnes]	1823	1778	1733	1686	1634	1577	1533	1484	1433	1383	1349	1317	1284	1249	1193
PM _{2.5}	Civil Aviation (1A3a)	[tonnes]	5	5	5	5	5	5	4	4	4	4	4	4	4	4	4
PM _{2.5}	Road (1A3b)	[tonnes]	4320	4690	4721	4643	4701	4936	5097	4913	4910	5033	4849	4672	4268	3924	3611
PM _{2.5}	Railways (1A3c)	[tonnes]	247	249	222	229	211	202	205	217	225	204	206	204	199	168	146
PM _{2.5}	Navigation (1A3d)	[tonnes]	1084	1084	1088	1083	1088	886	701	513	652	753	907	714	662	446	413
PM _{2.5}	Comm./Inst. (1A4a)	[tonnes]	24	24	24	24	24	24	24	23	22	23	24	25	27	28	29
PM _{2.5}	Residential (1A4b)	[tonnes]	12	12	12	12	11	11	11	11	11	11	11	11	11	11	11
PM _{2.5}	Ag./for./fish. (1A4c)	[tonnes]	2780	2817	2670	2720	2662	2625	2531	2359	2297	2116	2085	1890	1781	1631	1574
PM _{2.5}	Military (1A5)	[tonnes]	101	101	50	18	26	12	114	67	64	55	117	46	73	77	47
PM _{2.5}	Navigation int. (1A3d)	[tonnes]	2789	3396	5825	7693	7748	5448	4305	3939	8518	8071	9925	9819	9093	7601	8054
PM _{2.5}	Civil Aviation int. (1A3a)	[tonnes]	23	24	26	28	30	28	27	28	27	29	30	32	32	35	37
Arsenic	Industry-Other (1A2f)	[kg]						0	0	0	0	0	0	0	0	0	0
Arsenic	Civil Aviation (1A3a)	[kg]						0	0	0	0	0	0	0	0	0	0
Arsenic	Road (1A3b)	[kg]						1	1	1	1	1	1	1	1	1	1
Arsenic	Railways (1A3c)	[kg]						0	0	0	0	0	0	0	0	0	0
Arsenic	Navigation (1A3d)	[kg]						62	55	47	47	49	50	44	36	28	25
Arsenic	Comm./Inst. (1A4a)	[kg]						0	0	0	0	0	0	0	0	0	0
Arsenic	Residential (1A4b)	[kg]						0	0	0	0	0	0	0	0	0	0
Arsenic	Ag./for./fish. (1A4c)	[kg]						9	10	9	8	8	8	8	6	7	7
Arsenic	Military (1A5)	[kg]						0	0	0	0	0	0	0	0	0	0
Arsenic	Navigation int. (1A3d)	[kg]						353	292	267	465	496	505	325	417	357	369
Arsenic	Civil Aviation int. (1A3a)	[kg]						0	0	0	0	0	0	0	0	0	0
Cadmium	Industry-Other (1A2f)	[kg]						2	2	2	2	2	2	2	2	2	2
Cadmium	Civil Aviation (1A3a)	[kg]						0	0	0	0	0	0	0	0	0	0
Cadmium	Road (1A3b)	[kg]						27	29	30	30	32	32	32	33	34	34
Cadmium	Railways (1A3c)	[kg]						1	1	1	1	1	1	1	1	1	1
Cadmium	Navigation (1A3d)	[kg]						5	4	4	4	4	4	4	4	3	3
Cadmium	Comm./Inst. (1A4a)	[kg]						0	0	0	0	0	0	0	0	0	0

Cadmium	Residential (1A4b)	[kg]	0	0	0	0	0	0	0	0	0	0
Cadmium	Ag./for./fish. (1A4c)	[kg]	5	5	5	5	5	5	4	4	4	4
Cadmium	Military (1A5)	[kg]	0	0	0	0	0	0	0	0	0	0
Cadmium	Navigation int. (1A3d)	[kg]	23	20	19	31	34	35	20	29	26	26
Cadmium	Civil Aviation int. (1A3a)	[kg]	0	0	0	0	0	0	0	0	0	0
Chromium	Industry-Other (1A2f)	[kg]	7	7	7	7	7	7	7	7	7	7
Chromium	Civil Aviation (1A3a)	[kg]	0	0	0	0	0	0	0	0	0	0
Chromium	Road (1A3b)	[kg]	62	65	65	66	69	70	72	73	74	76
Chromium	Railways (1A3c)	[kg]	3	3	3	3	3	3	3	3	2	2
Chromium	Navigation (1A3d)	[kg]	28	25	23	22	23	24	22	19	15	14
Chromium	Comm./Inst. (1A4a)	[kg]	0	0	0	0	0	0	0	0	0	0
Chromium	Residential (1A4b)	[kg]	0	0	0	0	0	0	0	0	0	0
Chromium	Ag./for./fish. (1A4c)	[kg]	19	19	18	17	17	17	16	15	15	16
Chromium	Military (1A5)	[kg]	0	1	1	1	0	1	0	1	1	1
Chromium	Navigation int. (1A3d)	[kg]	147	123	115	195	210	214	131	178	157	160
Chromium	Civil Aviation int. (1A3a)	[kg]	0	0	0	0	0	0	0	0	0	0
Copper	Industry-Other (1A2f)	[kg]	5	5	5	5	5	5	6	6	6	6
Copper	Civil Aviation (1A3a)	[kg]	0	0	0	0	0	0	0	0	0	0
Copper	Road (1A3b)	[kg]	92	97	100	102	107	108	109	112	114	115
Copper	Railways (1A3c)	[kg]	2	2	2	2	2	2	2	2	2	1
Copper	Navigation (1A3d)	[kg]	63	56	48	47	49	50	45	36	29	26
Copper	Comm./Inst. (1A4a)	[kg]	1	1	1	1	1	1	1	1	1	1
Copper	Residential (1A4b)	[kg]	1	1	1	1	1	1	1	1	1	1
Copper	Ag./for./fish. (1A4c)	[kg]	18	18	17	16	16	16	17	14	14	15
Copper	Military (1A5)	[kg]	0	1	0	0	0	1	0	1	1	0
Copper	Navigation int. (1A3d)	[kg]	353	292	267	465	496	505	325	417	357	369
Copper	Civil Aviation int. (1A3a)	[kg]	0	0	0	0	0	0	0	0	0	0
Mercury	Industry-Other (1A2f)	[kg]	1	1	1	1	1	1	1	1	1	1
Mercury	Civil Aviation (1A3a)	[kg]	0	0	0	0	0	0	0	0	0	0
Mercury	Road (1A3b)	[kg]	21	22	22	23	24	24	24	25	25	25
Mercury	Railways (1A3c)	[kg]	0	1	1	1	1	1	1	0	0	0
Mercury	Navigation (1A3d)	[kg]	9	9	10	9	10	10	11	12	10	8
Mercury	Comm./Inst. (1A4a)	[kg]	0	0	0	0	0	0	0	0	0	0
Mercury	Residential (1A4b)	[kg]	0	0	0	0	0	0	0	0	0	0
Mercury	Ag./for./fish. (1A4c)	[kg]	12	12	11	10	10	10	10	8	8	9

Mercury	Military (1A5)	[kg]	0	0	0	0	0	0	0	0	0	0
Mercury	Navigation int. (1A3d)	[kg]	27	25	29	40	47	50	14	45	49	43
Mercury	Civil Aviation int. (1A3a)	[kg]	0	0	0	0	0	0	0	0	0	0
Nickel	Industry-Other (1A2f)	[kg]	2	2	2	2	2	2	2	2	2	2
Nickel	Civil Aviation (1A3a)	[kg]	0	0	0	0	0	0	0	0	0	0
Nickel	Road (1A3b)	[kg]	31	32	33	34	36	36	36	37	38	38
Nickel	Railways (1A3c)	[kg]	1	1	1	1	1	1	1	1	1	1
Nickel	Navigation (1A3d)	[kg]	3362	2889	2360	2359	2477	2492	2087	1520	1179	1077
Nickel	Comm./Inst. (1A4a)	[kg]	0	0	0	0	0	0	0	0	0	0
Nickel	Residential (1A4b)	[kg]	0	0	0	0	0	0	0	0	0	0
Nickel	Ag./for./fish. (1A4c)	[kg]	16	17	16	15	15	15	14	12	12	13
Nickel	Military (1A5)	[kg]	0	0	0	0	0	0	0	0	0	0
Nickel	Navigation int. (1A3d)	[kg]	20420	16701	14894	26627	28129	28488	19451	23291	19285	20431
Nickel	Civil Aviation int. (1A3a)	[kg]	0	0	0	0	0	0	0	0	0	0
Lead	Industry-Other (1A2f)	[kg]	12	12	12	12	12	12	12	12	12	12
Lead	Civil Aviation (1A3a)	[kg]	1534	1423	1378	1328	1639	1788	1640	1559	1399	1387
Lead	Road (1A3b)	[kg]	97622	75974	68894	29938	129	131	133	136	138	140
Lead	Railways (1A3c)	[kg]	4	5	5	5	5	5	5	4	4	4
Lead	Navigation (1A3d)	[kg]	35	34	33	32	33	34	35	32	26	23
Lead	Comm./Inst. (1A4a)	[kg]	1	1	1	1	1	1	1	1	1	1
Lead	Residential (1A4b)	[kg]	0	0	0	0	0	0	0	0	0	0
Lead	Ag./for./fish. (1A4c)	[kg]	38	38	36	35	34	35	33	30	29	31
Lead	Military (1A5)	[kg]	64	82	63	121	86	104	99	125	118	79
Lead	Navigation int. (1A3d)	[kg]	162	140	138	221	243	251	132	214	201	196
Lead	Civil Aviation int. (1A3a)	[kg]	490	465	452	456	153	175	126	145	145	124
Selenium	Industry-Other (1A2f)	[kg]	0	0	0	0	0	0	0	0	0	0
Selenium	Civil Aviation (1A3a)	[kg]	0	0	0	0	0	0	0	0	0	0
Selenium	Road (1A3b)	[kg]	0	0	0	0	1	1	1	1	1	1
Selenium	Railways (1A3c)	[kg]	0	0	0	0	0	0	0	0	0	0
Selenium	Navigation (1A3d)	[kg]	69	67	64	63	64	66	67	62	50	43
Selenium	Comm./Inst. (1A4a)	[kg]	0	0	0	0	0	0	0	0	0	0
Selenium	Residential (1A4b)	[kg]	0	0	0	0	0	0	0	0	0	0
Selenium	Ag./for./fish. (1A4c)	[kg]	37	38	35	33	33	33	32	25	26	30
Selenium	Military (1A5)	[kg]	0	0	0	0	0	0	0	0	0	0
Selenium	Navigation int. (1A3d)	[kg]	325	279	275	442	486	503	264	427	402	391

Selenium	Civil Aviation int. (1A3a)	[kg]	0	0	0	0	0	0	0	0	0
Zinc	Industry-Other (1A2f)	[kg]	419	420	420	419	419	422	425	428	432
Zinc	Civil Aviation (1A3a)	[kg]	6	5	5	5	6	7	6	6	5
Zinc	Road (1A3b)	[kg]	5466	5741	5915	6014	6314	6399	6463	6620	6730
Zinc	Railways (1A3c)	[kg]	150	152	161	167	151	153	152	148	124
Zinc	Navigation (1A3d)	[kg]	191	187	184	183	188	195	202	193	165
Zinc	Comm./Inst. (1A4a)	[kg]	51	51	52	52	53	54	55	56	57
Zinc	Residential (1A4b)	[kg]	27	27	27	27	27	27	28	28	28
Zinc	Ag./for./fish. (1A4c)	[kg]	766	762	724	717	683	698	664	655	626
Zinc	Military (1A5)	[kg]	5	57	35	31	28	63	26	46	54
Zinc	Navigation int. (1A3d)	[kg]	744	643	638	1017	1121	1162	595	991	940
Zinc	Civil Aviation int. (1A3a)	[kg]	2	2	2	2	1	1	0	1	1
HCB	Industry-Other (1A2f)	[g]	0	0	0	0	0	0	0	0	0
HCB	Civil Aviation (1A3a)	[g]	0	0	0	0	0	0	0	0	0
HCB	Road (1A3b)	[g]	0	0	0	0	0	0	0	0	0
HCB	Railways (1A3c)	[g]	0	0	0	0	0	0	0	0	0
HCB	Navigation (1A3d)	[g]	0	0	0	0	0	0	0	0	0
HCB	Comm./Inst. (1A4a)	[g]	0	0	0	0	0	0	0	0	0
HCB	Residential (1A4b)	[g]	0	0	0	0	0	0	0	0	0
HCB	Ag./for./fish. (1A4c)	[g]	0	0	0	0	0	0	0	0	0
HCB	Military (1A5)	[g]	0	0	0	0	0	0	0	0	0
HCB	Navigation int. (1A3d)	[g]	0	0	0	0	0	0	0	0	0
HCB	Civil Aviation int. (1A3a)	[g]	0	0	0	0	0	0	0	0	0
Dioxins/furans	Industry-Other (1A2f)	[g]	0	0	0	0	0	0	0	0	0
Dioxins/furans	Civil Aviation (1A3a)	[g]	0	0	0	0	0	0	0	0	0
Dioxins/furans	Road (1A3b)	[g]	1	1	1	1	1	1	1	1	1
Dioxins/furans	Railways (1A3c)	[g]	0	0	0	0	0	0	0	0	0
Dioxins/furans	Navigation (1A3d)	[g]	0	0	0	0	0	0	0	0	0
Dioxins/furans	Comm./Inst. (1A4a)	[g]	0	0	0	0	0	0	0	0	0
Dioxins/furans	Residential (1A4b)	[g]	0	0	0	0	0	0	0	0	0
Dioxins/furans	Ag./for./fish. (1A4c)	[g]	0	0	0	0	0	0	0	0	0
Dioxins/furans	Military (1A5)	[g]	0	0	0	0	0	0	0	0	0
Dioxins/furans	Navigation int. (1A3d)	[g]	1	0	0	1	1	1	1	1	1
Dioxins/furans	Civil Aviation int. (1A3a)	[g]	0	0	0	0	0	0	0	0	0
Flouranthene	Industry-Other (1A2f)	[kg]	45	44	45	46	45	46	46	46	46

Flouranthene	Civil Aviation (1A3a)	[kg]	0	0	0	0	1	1	1	0	0	0
Flouranthene	Road (1A3b)	[kg]	855	882	873	857	852	810	772	747	715	687
Flouranthene	Railways (1A3c)	[kg]	5	5	6	6	6	6	6	6	5	4
Flouranthene	Navigation (1A3d)	[kg]	66	68	71	70	70	74	82	82	67	58
Flouranthene	Comm./Inst. (1A4a)	[kg]	4	4	4	4	5	5	5	5	5	5
Flouranthene	Residential (1A4b)	[kg]	2	2	2	2	2	2	2	2	2	2
Flouranthene	Ag./for./fish. (1A4c)	[kg]	136	135	128	127	121	124	117	107	104	110
Flouranthene	Military (1A5)	[kg]	1	7	4	4	3	8	3	6	6	4
Flouranthene	Navigation int. (1A3d)	[kg]	198	184	205	288	334	355	344	316	338	304
Flouranthene	Civil Aviation int. (1A3a)	[kg]	0	0	0	0	0	0	0	0	0	0
Benzo(b) flouranthene	Industry-Other (1A2f)	[kg]	6	6	6	6	6	6	6	6	6	6
Benzo(b) flouranthene	Civil Aviation (1A3a)	[kg]	0	0	0	0	0	0	0	0	0	0
Benzo(b) flouranthene	Road (1A3b)	[kg]	69	71	71	70	71	70	69	68	67	66
Benzo(b) flouranthene	Railways (1A3c)	[kg]	1	1	1	2	1	1	1	1	1	1
Benzo(b) flouranthene	Navigation (1A3d)	[kg]	5	5	6	5	5	6	7	7	6	5
Benzo(b) flouranthene	Comm./Inst. (1A4a)	[kg]	0	0	0	0	0	0	0	0	0	0
Benzo(b) flouranthene	Residential (1A4b)	[kg]	0	0	0	0	0	0	0	0	0	0
Benzo(b) flouranthene	Ag./for./fish. (1A4c)	[kg]	15	15	14	14	13	13	13	12	11	12
Benzo(b) flouranthene	Military (1A5)	[kg]	0	1	1	1	0	1	0	1	1	1
Benzo(b) flouranthene	Navigation int. (1A3d)	[kg]	13	12	15	19	23	25	24	22	25	22
Benzo(b) flouranthene	Civil Aviation int. (1A3a)	[kg]	0	0	0	0	0	0	0	0	0	0
Benzo(k) flouranthene	Industry-Other (1A2f)	[kg]	6	6	6	6	6	6	6	6	6	6
Benzo(k) flouranthene	Civil Aviation (1A3a)	[kg]	0	0	0	0	0	0	0	0	0	0
Benzo(k) flouranthene	Road (1A3b)	[kg]	69	71	71	70	73	72	73	73	72	73
Benzo(k) flouranthene	Railways (1A3c)	[kg]	2	2	2	2	2	2	2	2	1	1
Benzo(k) flouranthene	Navigation (1A3d)	[kg]	2	2	2	2	2	3	3	3	3	2
Benzo(k) flouranthene	Comm./Inst. (1A4a)	[kg]	0	0	0	0	0	0	0	0	0	0
Benzo(k) flouranthene	Residential (1A4b)	[kg]	0	0	0	0	0	0	0	0	0	0
Benzo(k) flouranthene	Ag./for./fish. (1A4c)	[kg]	12	12	11	11	11	11	10	10	9	9
Benzo(k) flouranthene	Military (1A5)	[kg]	0	1	1	1	0	1	0	1	1	1
Benzo(k) flouranthene	Navigation int. (1A3d)	[kg]	6	6	7	9	11	11	11	10	12	10
Benzo(k) flouranthene	Civil Aviation int. (1A3a)	[kg]	0	0	0	0	0	0	0	0	0	0
Benzo(a) pyrene	Industry-Other (1A2f)	[kg]	3	3	3	3	3	3	3	3	3	3
Benzo(a) pyrene	Civil Aviation (1A3a)	[kg]	0	0	0	0	0	0	0	0	0	0
Benzo(a) pyrene	Road (1A3b)	[kg]	49	51	51	51	52	50	49	49	48	48

Benzo(a) pyrene	Railways (1A3c)	[kg]	0	0	0	0	0	0	0	0	0	0
Benzo(a) pyrene	Navigation (1A3d)	[kg]	1	1	1	1	1	1	2	2	1	1
Benzo(a) pyrene	Comm./Inst. (1A4a)	[kg]	0	0	0	0	0	0	0	0	0	0
Benzo(a) pyrene	Residential (1A4b)	[kg]	0	0	0	0	0	0	0	0	0	0
Benzo(a) pyrene	Ag./for./fish. (1A4c)	[kg]	6	6	6	6	5	5	5	5	5	5
Benzo(a) pyrene	Military (1A5)	[kg]	0	0	0	0	0	1	0	0	0	0
Benzo(a) pyrene	Navigation int. (1A3d)	[kg]	4	3	4	5	6	7	6	6	6	6
Benzo(a) pyrene	Civil Aviation int. (1A3a)	[kg]	0	0	0	0	0	0	0	0	0	0
Benzo(g,h,i) perylene	Industry-Other (1A2f)	[kg]	6	6	6	6	5	6	5	5	5	5
Benzo(g,h,i) perylene	Civil Aviation (1A3a)	[kg]	0	0	0	0	0	0	0	0	0	0
Benzo(g,h,i) perylene	Road (1A3b)	[kg]	104	108	108	107	108	104	101	100	97	95
Benzo(g,h,i) perylene	Railways (1A3c)	[kg]	0	0	0	0	0	0	0	0	0	0
Benzo(g,h,i) perylene	Navigation (1A3d)	[kg]	8	9	11	10	10	11	13	14	11	10
Benzo(g,h,i) perylene	Comm./Inst. (1A4a)	[kg]	1	1	1	1	1	1	1	1	1	1
Benzo(g,h,i) perylene	Residential (1A4b)	[kg]	0	0	0	0	0	0	0	0	0	0
Benzo(g,h,i) perylene	Ag./for./fish. (1A4c)	[kg]	21	21	20	19	19	19	18	16	15	16
Benzo(g,h,i) perylene	Military (1A5)	[kg]	0	1	1	1	0	1	0	1	1	0
Benzo(g,h,i) perylene	Navigation int. (1A3d)	[kg]	23	23	29	36	44	48	48	44	51	44
Benzo(g,h,i) perylene	Civil Aviation int. (1A3a)	[kg]	0	0	0	0	0	0	0	0	0	0
indeno(1,2,3-c,d) pyrene	Industry-Other (1A2f)	[kg]	3	3	3	3	3	3	3	3	3	3
indeno(1,2,3-c,d) pyrene	Civil Aviation (1A3a)	[kg]	0	0	0	0	0	0	0	0	0	0
indeno(1,2,3-c,d) pyrene	Road (1A3b)	[kg]	47	49	49	50	51	51	51	51	51	52
indeno(1,2,3-c,d) pyrene	Railways (1A3c)	[kg]	0	0	0	0	0	0	0	0	0	0
indeno(1,2,3-c,d) pyrene	Navigation (1A3d)	[kg]	7	7	8	8	8	9	10	11	9	8
indeno(1,2,3-c,d) pyrene	Comm./Inst. (1A4a)	[kg]	0	0	0	0	0	0	0	0	0	0
indeno(1,2,3-c,d) pyrene	Residential (1A4b)	[kg]	0	0	0	0	0	0	0	0	0	0
indeno(1,2,3-c,d) pyrene	Ag./for./fish. (1A4c)	[kg]	14	15	14	13	13	13	12	11	11	11
indeno(1,2,3-c,d) pyrene	Military (1A5)	[kg]	0	0	0	0	0	1	0	0	0	0
indeno(1,2,3-c,d) pyrene	Navigation int. (1A3d)	[kg]	19	19	23	29	36	39	39	36	42	36
indeno(1,2,3-c,d) pyrene	Civil Aviation int. (1A3a)	[kg]	0	0	0	0	0	0	0	0	0	0
PCB	Industry-Other (1A2f)	[g]	6	6	5	5	5	5	5	6	6	6
PCB	Civil Aviation (1A3a)	[g]	0	0	0	0	0	0	0	0	0	0
PCB	Road (1A3b)	[g]	88	82	73	63	20	21	21	21	22	22
PCB	Railways (1A3c)	[g]	2	2	2	2	2	2	2	2	2	2
PCB	Navigation (1A3d)	[g]	1	1	1	1	0	0	0	0	0	0

PCB	Comm./Inst. (1A4a)	[g]	1	1	1	1	0	0	0	0	0	0
PCB	Residential (1A4b)	[g]	1	0	0	0	0	0	0	0	0	0
PCB	Ag./for./fish. (1A4c)	[g]	10	10	9	9	8	8	8	8	8	8
PCB	Military (1A5)	[g]	0	1	0	0	0	1	0	0	0	0
PCB	Navigation int. (1A3d)	[g]	0	0	0	1	1	1	1	1	1	1
PCB	Civil Aviation int. (1A3a)	[g]	0	0	0	0	0	0	0	0	0	0

pol_name	IPCC ID	Unit	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
SO ₂	Industry-Other (1A2f)	[tonnes]	253	256	258	261	263	28	30	33	33	25	31	6	6
SO ₂	Civil Aviation (1A3a)	[tonnes]	49	52	45	44	41	43	46	51	52	49	50	46	42
SO ₂	Road (1A3b)	[tonnes]	352	353	357	371	381	77	79	83	81	77	76	74	71
SO ₂	Railways (1A3c)	[tonnes]	7	7	7	7	7	1	1	1	1	1	2	2	2
SO ₂	Navigation (1A3d)	[tonnes]	1844	1733	1582	1984	2319	2339	2431	1685	1512	1588	1438	1383	1334
SO ₂	Comm./Inst. (1A4a)	[tonnes]	3	3	4	4	5	1	1	1	1	1	1	1	1
SO ₂	Residential (1A4b)	[tonnes]	1	1	2	2	2	0	0	0	0	0	0	0	0
SO ₂	Ag./for./fish. (1A4c)	[tonnes]	1207	1209	1236	1203	1022	852	800	690	421	396	358	328	311
SO ₂	Military (1A5)	[tonnes]	27	12	19	17	46	57	26	40	19	25	20	37	17
SO ₂	Navigation int. (1A3d)	[tonnes]	55182	43830	30036	30982	26540	34283	50417	25652	19326	7383	8200	8853	5136
SO ₂	Civil Aviation int. (1A3a)	[tonnes]	750	761	657	683	781	822	824	845	845	740	773	796	801
NO _x	Industry-Other (1A2f)	[tonnes]	12096	11869	11617	11214	10744	10706	10843	10827	10142	7247	8665	8046	7597
NO _x	Civil Aviation (1A3a)	[tonnes]	723	752	641	595	551	583	602	693	697	635	623	578	579
NO _x	Road (1A3b)	[tonnes]	82320	79704	76400	75298	73383	69930	67887	66162	59179	51966	49914	48140	44316
NO _x	Railways (1A3c)	[tonnes]	3727	3396	3396	3540	3478	3724	3542	3555	2920	2603	2818	2501	2531
NO _x	Navigation (1A3d)	[tonnes]	8087	8197	8315	8443	8469	8634	8979	9054	9303	9498	9557	9064	8692
NO _x	Comm./Inst. (1A4a)	[tonnes]	104	112	124	138	155	177	199	215	222	220	217	215	219
NO _x	Residential (1A4b)	[tonnes]	50	54	59	64	69	72	76	79	82	84	87	89	92
NO _x	Ag./for./fish. (1A4c)	[tonnes]	25333	25787	26036	25286	22447	24009	22832	20889	22155	20950	19513	19108	18163
NO _x	Military (1A5)	[tonnes]	526	663	463	511	1256	1300	603	760	488	725	456	798	492
NO _x	Navigation int. (1A3d)	[tonnes]	91641	75429	60383	65339	53439	56540	78012	83555	70401	35658	51065	52516	36524
NO _x	Civil Aviation int. (1A3a)	[tonnes]	9446	9600	8724	9084	10472	11025	11158	11402	11292	9843	10110	10466	10628
NMVOC	Industry-Other (1A2f)	[tonnes]	1926	1873	1815	1754	1676	1627	1589	1524	1383	992	1191	1128	1076
NMVOC	Civil Aviation (1A3a)	[tonnes]	158	152	150	142	154	160	151	159	142	122	107	89	98
NMVOC	Road (1A3b)	[tonnes]	42322	38860	35321	32626	28450	25969	22716	20251	17728	15452	13982	11769	10439
NMVOC	Railways (1A3c)	[tonnes]	253	248	243	223	217	235	230	231	205	174	189	175	190

NM VOC	Navigation (1A3d)	[tonnes]	1731	1702	1661	1602	1534	1423	1305	1190	1095	1010	932	840	749
NM VOC	Comm./Inst. (1A4a)	[tonnes]	2845	3504	4188	4897	5631	5775	5922	6022	5844	5159	4423	3636	3636
NM VOC	Residential (1A4b)	[tonnes]	1757	1824	1894	1972	2053	2084	2115	2134	2109	2071	2032	1993	1953
NM VOC	Ag./for./fish. (1A4c)	[tonnes]	3520	3378	3199	2987	2698	2712	2662	2598	2638	2512	2359	2246	2137
NM VOC	Military (1A5)	[tonnes]	55	53	45	45	100	106	51	68	40	55	40	59	33
NM VOC	Navigation int. (1A3d)	[tonnes]	2940	2433	1989	2130	1731	1792	2418	2563	2195	1160	1628	1668	1199
NM VOC	Civil Aviation int. (1A3a)	[tonnes]	437	359	375	368	403	408	414	410	398	341	329	356	379
CH ₄	Industry-Other (1A2f)	[tonnes]	50	49	48	47	46	45	44	43	41	30	38	37	36
CH ₄	Civil Aviation (1A3a)	[tonnes]	4	4	3	3	3	4	3	4	3	3	2	2	2
CH ₄	Road (1A3b)	[tonnes]	1788	1676	1574	1497	1393	1274	1172	1070	909	778	695	616	538
CH ₄	Railways (1A3c)	[tonnes]	10	10	9	9	8	9	9	9	8	7	7	7	7
CH ₄	Navigation (1A3d)	[tonnes]	33	33	34	34	35	35	35	35	35	35	35	34	34
CH ₄	Comm./Inst. (1A4a)	[tonnes]	92	101	113	127	144	157	169	175	174	167	160	151	151
CH ₄	Residential (1A4b)	[tonnes]	45	48	51	55	60	62	64	65	66	66	65	65	65
CH ₄	Ag./for./fish. (1A4c)	[tonnes]	91	90	90	89	85	90	97	104	111	115	116	117	111
CH ₄	Military (1A5)	[tonnes]	5	6	5	5	12	12	6	7	4	5	3	6	3
CH ₄	Navigation int. (1A3d)	[tonnes]	91	75	62	66	54	55	75	79	68	36	50	52	37
CH ₄	Civil Aviation int. (1A3a)	[tonnes]	12	12	12	11	12	12	12	12	13	10	10	11	12
CO	Industry-Other (1A2f)	[tonnes]	8395	8227	8030	7842	7600	7528	7542	7499	7127	5199	6530	6358	6239
CO	Civil Aviation (1A3a)	[tonnes]	895	891	863	835	858	861	842	902	823	717	688	598	607
CO	Road (1A3b)	[tonnes]	276031	264825	243650	233641	207364	198459	177787	162791	148403	132200	124918	104957	96022
CO	Railways (1A3c)	[tonnes]	694	637	627	611	599	648	626	629	526	450	481	398	425
CO	Navigation (1A3d)	[tonnes]	6832	7034	7217	7408	7601	7631	7281	6915	6565	6210	5835	5420	5030
CO	Comm./Inst. (1A4a)	[tonnes]	29423	32889	37681	43798	51239	58128	64197	67870	70290	72227	72338	72458	72587
CO	Residential (1A4b)	[tonnes]	16451	17390	18463	19890	21444	22482	23547	24366	25092	25341	25616	25915	26236
CO	Ag./for./fish. (1A4c)	[tonnes]	26192	24444	22571	20670	18575	17655	17414	18148	19039	19488	19723	19965	19246
CO	Military (1A5)	[tonnes]	399	308	311	302	706	798	380	538	306	414	311	495	270
CO	Navigation int. (1A3d)	[tonnes]	9699	8025	6562	7025	5709	5912	7977	8454	7243	3826	5371	5504	3955
CO	Civil Aviation int. (1A3a)	[tonnes]	1790	1795	1608	1668	1848	1907	1852	1906	1979	1690	1716	1790	1776
CO ₂	Industry-Other (1A2f)	[ktonnes]	877	886	895	905	910	952	1023	1104	1126	834	1051	1023	1021
CO ₂	Civil Aviation (1A3a)	[ktonnes]	154	163	141	138	128	135	143	161	162	153	156	146	133
CO ₂	Road (1A3b)	[ktonnes]	11203	11223	11352	11806	12115	12214	12587	13187	12870	12160	12080	11748	11224
CO ₂	Railways (1A3c)	[ktonnes]	228	211	210	218	216	232	227	228	237	230	242	249	249
CO ₂	Navigation (1A3d)	[ktonnes]	588	587	578	576	588	585	588	586	590	592	591	559	498
CO ₂	Comm./Inst. (1A4a)	[ktonnes]	87	98	112	129	149	162	172	175	176	174	173	171	171

CO ₂	Residential (1A4b)	[ktonnes]	43	46	49	53	57	59	61	62	63	63	63	63	62
CO ₂	Ag./for./fish. (1A4c)	[ktonnes]	1762	1770	1793	1768	1639	1758	1750	1727	1879	1853	1809	1838	1839
CO ₂	Military (1A5)	[ktonnes]	111	97	89	92	239	271	126	175	108	160	107	193	116
CO ₂	Navigation int. (1A3d)	[ktonnes]	4021	3304	2691	2853	2299	2352	3136	3292	2809	1487	2063	2096	1505
CO ₂	Civil Aviation int. (1A3a)	[ktonnes]	2350	2384	2058	2141	2447	2574	2581	2647	2647	2316	2421	2492	2510
N ₂ O	Industry-Other (1A2f)	[tonnes]	37	38	38	38	39	40	43	47	48	35	45	43	43
N ₂ O	Civil Aviation (1A3a)	[tonnes]	8	8	8	8	8	8	8	9	9	8	8	8	7
N ₂ O	Road (1A3b)	[tonnes]	415	417	417	426	430	422	423	434	417	390	388	391	376
N ₂ O	Railways (1A3c)	[tonnes]	6	6	6	6	6	6	6	6	7	6	7	7	7
N ₂ O	Navigation (1A3d)	[tonnes]	34	34	34	33	34	34	34	34	34	34	34	33	29
N ₂ O	Comm./Inst. (1A4a)	[tonnes]	1	1	2	2	2	2	3	3	3	3	3	3	3
N ₂ O	Residential (1A4b)	[tonnes]	1	1	1	1	1	1	1	1	1	1	1	1	1
N ₂ O	Ag./for./fish. (1A4c)	[tonnes]	87	88	90	88	80	87	86	83	92	90	87	88	88
N ₂ O	Military (1A5)	[tonnes]	3	3	3	3	8	9	4	6	4	5	4	7	4
N ₂ O	Navigation int. (1A3d)	[tonnes]	253	208	170	180	145	148	197	207	177	94	130	132	95
N ₂ O	Civil Aviation int. (1A3a)	[tonnes]	82	82	72	75	85	89	89	91	91	79	83	86	86
NH ₃	Industry-Other (1A2f)	[tonnes]	2	2	2	2	2	2	2	3	3	2	3	2	2
NH ₃	Civil Aviation (1A3a)	[tonnes]	0	0	0	0	0	0	0	0	0	0	0	0	0
NH ₃	Road (1A3b)	[tonnes]	2544	2565	2586	2565	2538	2402	2294	2212	2043	1902	1731	1602	1456
NH ₃	Railways (1A3c)	[tonnes]	1	1	1	1	1	1	1	1	1	1	1	1	1
NH ₃	Navigation (1A3d)	[tonnes]	0	0	0	0	0	0	0	0	0	0	0	0	0
NH ₃	Comm./Inst. (1A4a)	[tonnes]	0	0	0	0	0	0	0	0	0	0	0	0	0
NH ₃	Residential (1A4b)	[tonnes]	0	0	0	0	0	0	0	0	0	0	0	0	0
NH ₃	Ag./for./fish. (1A4c)	[tonnes]	3	3	3	3	3	3	3	3	4	4	4	4	4
NH ₃	Military (1A5)	[tonnes]	0	0	0	0	1	1	0	0	1	1	0	0	0
NH ₃	Navigation int. (1A3d)	[tonnes]													
NH ₃	Civil Aviation int. (1A3a)	[tonnes]	0	0	0	0	0	0	0	0	0	0	0	0	0
TSP	Industry-Other (1A2f)	[tonnes]	1135	1121	1098	1075	1037	1011	998	968	883	606	706	660	617
TSP	Civil Aviation (1A3a)	[tonnes]	3	4	3	3	3	3	3	3	3	3	3	3	3
TSP	Road (1A3b)	[tonnes]	3273	3035	2793	2745	2595	2479	2403	2318	2038	1740	1632	1464	1311
TSP	Railways (1A3c)	[tonnes]	141	125	124	119	115	124	120	120	101	84	95	78	82
TSP	Navigation (1A3d)	[tonnes]	383	373	357	387	430	425	421	336	327	325	306	290	268
TSP	Comm./Inst. (1A4a)	[tonnes]	30	38	46	55	63	65	66	66	67	67	67	67	67
TSP	Residential (1A4b)	[tonnes]	11	11	12	13	13	13	14	14	14	14	14	15	15
TSP	Ag./for./fish. (1A4c)	[tonnes]	1553	1498	1429	1351	1244	1213	1144	1076	1051	998	942	910	858

TSP	Military (1A5)	[tonnes]	17	34	16	19	42	36	16	16	13	20	10	15	11
TSP	Navigation int. (1A3d)	[tonnes]	8745	7143	4988	4501	3978	5761	7888	2365	1873	820	934	975	641
TSP	Civil Aviation int. (1A3a)	[tonnes]	38	38	33	35	40	42	42	43	43	37	39	40	40
PM ₁₀	Industry-Other (1A2f)	[tonnes]	1135	1121	1098	1075	1037	1011	998	968	883	606	706	660	617
PM ₁₀	Civil Aviation (1A3a)	[tonnes]	3	4	3	3	3	3	3	3	3	3	3	3	3
PM ₁₀	Road (1A3b)	[tonnes]	3273	3035	2793	2745	2595	2479	2403	2318	2038	1740	1632	1464	1311
PM ₁₀	Railways (1A3c)	[tonnes]	141	125	124	119	115	124	120	120	101	84	95	78	82
PM ₁₀	Navigation (1A3d)	[tonnes]	381	371	355	384	427	422	418	334	325	323	304	289	266
PM ₁₀	Comm./Inst. (1A4a)	[tonnes]	30	38	46	55	63	65	66	66	67	67	67	67	67
PM ₁₀	Residential (1A4b)	[tonnes]	11	11	12	13	13	13	14	14	14	14	14	15	15
PM ₁₀	Ag./for./fish. (1A4c)	[tonnes]	1551	1496	1426	1349	1242	1211	1143	1074	1049	996	941	908	856
PM ₁₀	Military (1A5)	[tonnes]	17	34	16	19	42	36	16	16	13	20	10	15	11
PM ₁₀	Navigation int. (1A3d)	[tonnes]	8658	7072	4938	4456	3938	5703	7809	2341	1854	812	925	965	634
PM ₁₀	Civil Aviation int. (1A3a)	[tonnes]	38	38	33	35	40	42	42	43	43	37	39	40	40
PM _{2.5}	Industry-Other (1A2f)	[tonnes]	1135	1121	1098	1075	1037	1011	998	968	883	606	706	660	617
PM _{2.5}	Civil Aviation (1A3a)	[tonnes]	3	4	3	3	3	3	3	3	3	3	3	3	3
PM _{2.5}	Road (1A3b)	[tonnes]	3273	3035	2793	2745	2595	2479	2403	2318	2038	1740	1632	1464	1311
PM _{2.5}	Railways (1A3c)	[tonnes]	141	125	124	119	115	124	120	120	101	84	95	78	82
PM _{2.5}	Navigation (1A3d)	[tonnes]	379	370	354	383	425	421	417	333	324	322	303	288	265
PM _{2.5}	Comm./Inst. (1A4a)	[tonnes]	30	38	46	55	63	65	66	66	67	67	67	67	67
PM _{2.5}	Residential (1A4b)	[tonnes]	11	11	12	13	13	13	14	14	14	14	14	15	15
PM _{2.5}	Ag./for./fish. (1A4c)	[tonnes]	1549	1495	1425	1348	1242	1210	1142	1073	1048	995	940	907	856
PM _{2.5}	Military (1A5)	[tonnes]	17	34	16	19	42	36	16	16	13	20	10	15	11
PM _{2.5}	Navigation int. (1A3d)	[tonnes]	8614	7036	4913	4434	3918	5675	7770	2330	1845	808	920	960	631
PM _{2.5}	Civil Aviation int. (1A3a)	[tonnes]	38	38	33	35	40	42	42	43	43	37	39	40	40
Arsenic	Industry-Other (1A2f)	[kg]	0	0	0	0	0	0	0	0	0	0	0	0	0
Arsenic	Civil Aviation (1A3a)	[kg]	0	0	0	0	0	0	0	0	0	0	0	0	0
Arsenic	Road (1A3b)	[kg]	1	1	1	1	1	1	1	1	1	1	1	1	1
Arsenic	Railways (1A3c)	[kg]	0	0	0	0	0	0	0	0	0	0	0	0	0
Arsenic	Navigation (1A3d)	[kg]	24	23	23	28	28	28	30	30	31	33	35	33	32
Arsenic	Comm./Inst. (1A4a)	[kg]	0	0	0	0	0	0	0	0	0	0	0	0	0
Arsenic	Residential (1A4b)	[kg]	0	0	0	0	0	0	0	0	0	0	0	0	0
Arsenic	Ag./for./fish. (1A4c)	[kg]	11	11	11	11	9	10	10	8	10	9	8	8	8
Arsenic	Military (1A5)	[kg]	0	0	0	0	0	0	0	0	0	0	0	0	0
Arsenic	Navigation int. (1A3d)	[kg]	420	329	227	257	213	250	381	424	326	127	205	221	128

Arsenic	Civil Aviation int. (1A3a)	[kg]	0	0	0	0	0	0	0	0	0	0	0	0
Cadmium	Industry-Other (1A2f)	[kg]	2	2	2	2	2	2	3	3	3	2	3	3
Cadmium	Civil Aviation (1A3a)	[kg]	0	0	0	0	0	0	0	0	0	0	0	0
Cadmium	Road (1A3b)	[kg]	34	34	35	36	37	37	38	40	39	38	38	39
Cadmium	Railways (1A3c)	[kg]	1	1	1	1	1	1	1	1	1	1	1	1
Cadmium	Navigation (1A3d)	[kg]	3	2	2	3	3	3	3	3	3	3	3	3
Cadmium	Comm./Inst. (1A4a)	[kg]	0	0	0	0	1	1	1	1	1	1	1	1
Cadmium	Residential (1A4b)	[kg]	0	0	0	0	0	0	0	0	0	0	0	0
Cadmium	Ag./for./fish. (1A4c)	[kg]	5	5	5	5	4	5	5	5	5	5	5	5
Cadmium	Military (1A5)	[kg]	0	0	0	0	0	0	0	0	0	0	0	0
Cadmium	Navigation int. (1A3d)	[kg]	28	23	17	18	15	17	24	27	21	9	14	15
Cadmium	Civil Aviation int. (1A3a)	[kg]	0	0	0	0	0	0	0	0	0	0	0	0
Chromium	Industry-Other (1A2f)	[kg]	7	7	7	8	8	8	9	9	10	7	9	9
Chromium	Civil Aviation (1A3a)	[kg]	0	0	0	0	0	0	0	0	0	0	0	0
Chromium	Road (1A3b)	[kg]	75	76	78	82	86	87	91	98	97	94	96	98
Chromium	Railways (1A3c)	[kg]	2	2	2	2	2	2	2	2	2	2	2	2
Chromium	Navigation (1A3d)	[kg]	13	12	12	14	14	14	15	15	15	16	17	16
Chromium	Comm./Inst. (1A4a)	[kg]	0	0	1	1	1	1	1	1	1	1	1	1
Chromium	Residential (1A4b)	[kg]	0	0	0	0	0	0	0	0	0	0	0	0
Chromium	Ag./for./fish. (1A4c)	[kg]	18	18	19	18	17	18	18	17	19	18	18	18
Chromium	Military (1A5)	[kg]	0	1	0	0	1	1	0	0	0	1	0	1
Chromium	Navigation int. (1A3d)	[kg]	178	140	100	111	92	106	157	174	136	56	87	93
Chromium	Civil Aviation int. (1A3a)	[kg]	0	0	0	0	0	0	0	0	0	0	0	0
Copper	Industry-Other (1A2f)	[kg]	6	6	6	6	6	6	7	7	7	5	7	7
Copper	Civil Aviation (1A3a)	[kg]	0	0	0	0	0	0	0	0	0	0	0	0
Copper	Road (1A3b)	[kg]	114	114	116	119	122	120	122	126	123	119	117	117
Copper	Railways (1A3c)	[kg]	1	1	1	1	1	1	1	1	1	1	2	2
Copper	Navigation (1A3d)	[kg]	25	24	24	29	29	29	31	30	32	34	36	34
Copper	Comm./Inst. (1A4a)	[kg]	1	1	1	2	2	2	2	2	2	2	2	2
Copper	Residential (1A4b)	[kg]	1	1	1	1	1	1	1	1	1	1	1	1
Copper	Ag./for./fish. (1A4c)	[kg]	18	18	18	18	16	17	17	16	18	17	16	17
Copper	Military (1A5)	[kg]	0	0	0	0	1	1	0	0	0	1	0	1
Copper	Navigation int. (1A3d)	[kg]	420	329	227	257	213	250	381	424	326	127	205	221
Copper	Civil Aviation int. (1A3a)	[kg]	0	0	0	0	0	0	0	0	0	0	0	0
Mercury	Industry-Other (1A2f)	[kg]	1	1	1	1	1	1	2	2	2	1	2	2

Mercury	Civil Aviation (1A3a)	[kg]	0	0	0	0	0	0	0	0	0	0	0	0
Mercury	Road (1A3b)	[kg]	25	25	25	26	26	26	27	28	27	26	25	24
Mercury	Railways (1A3c)	[kg]	0	0	0	0	0	0	0	0	0	0	0	0
Mercury	Navigation (1A3d)	[kg]	7	7	7	6	6	6	6	6	6	6	6	5
Mercury	Comm./Inst. (1A4a)	[kg]	0	0	0	0	0	0	0	0	0	0	0	0
Mercury	Residential (1A4b)	[kg]	0	0	0	0	0	0	0	0	0	0	0	0
Mercury	Ag./for./fish. (1A4c)	[kg]	13	13	13	13	10	12	12	10	12	11	10	10
Mercury	Military (1A5)	[kg]	0	0	0	0	0	0	0	0	0	0	0	0
Mercury	Navigation int. (1A3d)	[kg]	39	34	30	31	24	23	27	27	25	17	21	20
Mercury	Civil Aviation int. (1A3a)	[kg]	0	0	0	0	0	0	0	0	0	0	0	0
Nickel	Industry-Other (1A2f)	[kg]	2	2	2	2	2	2	3	3	3	2	3	3
Nickel	Civil Aviation (1A3a)	[kg]	0	0	0	0	0	0	0	0	0	0	0	0
Nickel	Road (1A3b)	[kg]	38	38	39	40	41	41	42	44	43	42	41	42
Nickel	Railways (1A3c)	[kg]	1	1	1	1	1	1	1	1	1	1	1	1
Nickel	Navigation (1A3d)	[kg]	1068	1036	1026	1374	1367	1371	1494	1477	1583	1685	1809	1749
Nickel	Comm./Inst. (1A4a)	[kg]	0	0	0	1	1	1	1	1	1	1	1	1
Nickel	Residential (1A4b)	[kg]	0	0	0	0	0	0	0	0	0	0	0	0
Nickel	Ag./for./fish. (1A4c)	[kg]	18	18	19	18	15	17	16	15	17	16	15	15
Nickel	Military (1A5)	[kg]	0	0	0	0	0	0	0	0	0	0	0	0
Nickel	Navigation int. (1A3d)	[kg]	23825	18510	12366	14147	11846	14256	22148	24842	18832	6924	11521	12574
Nickel	Civil Aviation int. (1A3a)	[kg]	0	0	0	0	0	0	0	0	0	0	0	0
Lead	Industry-Other (1A2f)	[kg]	13	13	13	13	13	14	15	16	16	12	15	15
Lead	Civil Aviation (1A3a)	[kg]	1369	1343	1328	1252	1304	1297	1245	1329	1182	991	929	776
Lead	Road (1A3b)	[kg]	140	141	145	153	160	161	168	179	179	173	177	184
Lead	Railways (1A3c)	[kg]	3	3	3	3	3	4	3	3	4	3	4	4
Lead	Navigation (1A3d)	[kg]	21	20	20	21	21	21	22	22	22	22	23	22
Lead	Comm./Inst. (1A4a)	[kg]	1	1	1	1	2	2	2	2	2	2	2	2
Lead	Residential (1A4b)	[kg]	0	0	1	1	1	1	1	1	1	1	1	1
Lead	Ag./for./fish. (1A4c)	[kg]	38	38	39	38	34	37	36	34	38	37	35	36
Lead	Military (1A5)	[kg]	114	89	106	79	84	60	47	81	40	66	80	49
Lead	Navigation int. (1A3d)	[kg]	206	166	126	137	112	121	172	186	151	70	103	108
Lead	Civil Aviation int. (1A3a)	[kg]	118	114	113	106	111	117	22	10	113	52	10	52
Selenium	Industry-Other (1A2f)	[kg]	0	0	0	0	0	0	0	0	0	0	0	0
Selenium	Civil Aviation (1A3a)	[kg]	0	0	0	0	0	0	0	0	0	0	0	0
Selenium	Road (1A3b)	[kg]	1	1	1	1	1	1	1	1	1	1	1	0

Selenium	Railways (1A3c)	[kg]	0	0	0	0	0	0	0	0	0	0	0	0	0
Selenium	Navigation (1A3d)	[kg]	39	38	37	39	40	40	41	40	41	42	43	41	37
Selenium	Comm./Inst. (1A4a)	[kg]	0	0	0	0	0	0	0	0	0	0	0	0	0
Selenium	Residential (1A4b)	[kg]	0	0	0	0	0	0	0	0	0	0	0	0	0
Selenium	Ag./for./fish. (1A4c)	[kg]	44	44	45	44	34	41	38	33	38	36	32	32	30
Selenium	Military (1A5)	[kg]	0	0	0	0	0	0	0	0	0	0	0	0	0
Selenium	Navigation int. (1A3d)	[kg]	412	331	252	274	224	243	345	372	302	140	207	216	141
Selenium	Civil Aviation int. (1A3a)	[kg]	0	0	0	0	0	0	0	0	0	0	0	0	0
Zinc	Industry-Other (1A2f)	[kg]	438	443	447	452	455	476	512	553	564	417	526	512	511
Zinc	Civil Aviation (1A3a)	[kg]	5	5	5	5	5	5	5	5	4	4	3	3	3
Zinc	Road (1A3b)	[kg]	6783	6796	6937	7215	7437	7403	7576	7944	7844	7595	7629	7791	7715
Zinc	Railways (1A3c)	[kg]	115	106	106	110	109	117	114	115	119	116	122	126	126
Zinc	Navigation (1A3d)	[kg]	142	143	143	149	152	152	153	152	154	155	156	150	140
Zinc	Comm./Inst. (1A4a)	[kg]	60	68	77	89	103	112	119	121	122	121	119	118	118
Zinc	Residential (1A4b)	[kg]	29	32	34	36	39	41	42	43	44	44	43	43	43
Zinc	Ag./for./fish. (1A4c)	[kg]	653	658	662	659	645	668	678	698	745	747	746	760	769
Zinc	Military (1A5)	[kg]	14	32	16	22	53	48	24	26	25	44	25	44	38
Zinc	Navigation int. (1A3d)	[kg]	950	766	588	637	519	560	788	848	692	326	478	497	330
Zinc	Civil Aviation int. (1A3a)	[kg]	0	0	0	0	0	0	0	0	0	0	0	0	0
HCB	Industry-Other (1A2f)	[g]	0	0	0	0	0	0	0	0	0	0	0	0	0
HCB	Civil Aviation (1A3a)	[g]	0	0	0	0	0	0	0	0	0	0	0	0	0
HCB	Road (1A3b)	[g]	0	0	0	0	1	1	1	1	1	1	1	1	1
HCB	Railways (1A3c)	[g]	0	0	0	0	0	0	0	0	0	0	0	0	0
HCB	Navigation (1A3d)	[g]	0	0	0	0	0	0	0	0	0	0	0	0	0
HCB	Comm./Inst. (1A4a)	[g]	0	0	0	0	0	0	0	0	0	0	0	0	0
HCB	Residential (1A4b)	[g]	0	0	0	0	0	0	0	0	0	0	0	0	0
HCB	Ag./for./fish. (1A4c)	[g]	0	0	0	0	0	0	0	0	0	0	0	0	0
HCB	Military (1A5)	[g]	0	0	0	0	0	0	0	0	0	0	0	0	0
HCB	Navigation int. (1A3d)	[g]	0	0	0	0	0	0	0	0	0	0	0	0	0
HCB	Civil Aviation int. (1A3a)	[g]	0	0	0	0	0	0	0	0	0	0	0	0	0
Dioxins/furans	Industry-Other (1A2f)	[g]	0	0	0	0	0	0	0	0	0	0	0	0	0
Dioxins/furans	Civil Aviation (1A3a)	[g]	0	0	0	0	0	0	0	0	0	0	0	0	0
Dioxins/furans	Road (1A3b)	[g]	0	0	0	0	0	0	0	0	0	0	0	0	0
Dioxins/furans	Railways (1A3c)	[g]	0	0	0	0	0	0	0	0	0	0	0	0	0
Dioxins/furans	Navigation (1A3d)	[g]	0	0	0	0	0	0	0	0	0	0	0	0	0

Dioxins/furans	Comm./Inst. (1A4a)	[g]	0	0	0	0	0	0	0	0	0	0	0	0	0
Dioxins/furans	Residential (1A4b)	[g]	0	0	0	0	0	0	0	0	0	0	0	0	0
Dioxins/furans	Ag./for./fish. (1A4c)	[g]	0	0	0	0	0	0	0	0	0	0	0	0	0
Dioxins/furans	Military (1A5)	[g]	0	0	0	0	0	0	0	0	0	0	0	0	0
Dioxins/furans	Navigation int. (1A3d)	[g]	1	1	0	0	0	0	1	1	0	0	0	0	0
Dioxins/furans	Civil Aviation int. (1A3a)	[g]	0	0	0	0	0	0	0	0	0	0	0	0	0
Flouranthene	Industry-Other (1A2f)	[kg]	48	48	49	49	50	52	56	61	63	46	58	57	56
Flouranthene	Civil Aviation (1A3a)	[kg]	0	0	0	0	0	0	0	0	0	0	0	0	0
Flouranthene	Road (1A3b)	[kg]	661	640	640	668	689	700	740	820	840	812	839	874	887
Flouranthene	Railways (1A3c)	[kg]	4	4	4	4	4	4	4	4	5	4	5	5	5
Flouranthene	Navigation (1A3d)	[kg]	52	51	50	49	50	50	49	49	49	49	49	46	40
Flouranthene	Comm./Inst. (1A4a)	[kg]	5	6	7	8	9	10	10	10	10	10	10	10	10
Flouranthene	Residential (1A4b)	[kg]	3	3	3	3	3	3	4	4	4	4	4	4	4
Flouranthene	Ag./for./fish. (1A4c)	[kg]	133	133	135	132	119	130	128	123	135	132	127	129	128
Flouranthene	Military (1A5)	[kg]	2	4	2	3	6	6	3	3	3	5	3	5	4
Flouranthene	Navigation int. (1A3d)	[kg]	283	238	208	215	171	164	203	205	187	114	149	147	116
Flouranthene	Civil Aviation int. (1A3a)	[kg]	0	0	0	0	0	0	0	0	0	0	0	0	0
Benzo(b) flouranthene	Industry-Other (1A2f)	[kg]	6	6	6	6	6	6	7	7	7	5	7	7	7
Benzo(b) flouranthene	Civil Aviation (1A3a)	[kg]	0	0	0	0	0	0	0	0	0	0	0	0	0
Benzo(b) flouranthene	Road (1A3b)	[kg]	64	63	63	66	68	69	72	77	77	73	75	77	77
Benzo(b) flouranthene	Railways (1A3c)	[kg]	1	1	1	1	1	1	1	1	1	1	1	1	1
Benzo(b) flouranthene	Navigation (1A3d)	[kg]	4	4	4	4	4	4	4	4	4	4	4	4	3
Benzo(b) flouranthene	Comm./Inst. (1A4a)	[kg]	0	0	0	0	0	0	0	1	1	0	0	0	0
Benzo(b) flouranthene	Residential (1A4b)	[kg]	0	0	0	0	0	0	0	0	0	0	0	0	0
Benzo(b) flouranthene	Ag./for./fish. (1A4c)	[kg]	13	13	13	13	12	13	13	13	14	14	13	13	13
Benzo(b) flouranthene	Military (1A5)	[kg]	0	0	0	0	1	1	0	0	0	1	0	1	0
Benzo(b) flouranthene	Navigation int. (1A3d)	[kg]	19	17	15	16	12	11	13	12	12	8	10	10	9
Benzo(b) flouranthene	Civil Aviation int. (1A3a)	[kg]	0	0	0	0	0	0	0	0	0	0	0	0	0
Benzo(k) flouranthene	Industry-Other (1A2f)	[kg]	6	5	5	6	6	6	6	7	7	5	7	6	6
Benzo(k) flouranthene	Civil Aviation (1A3a)	[kg]	0	0	0	0	0	0	0	0	0	0	0	0	0
Benzo(k) flouranthene	Road (1A3b)	[kg]	71	71	71	75	77	79	82	88	86	81	83	85	84
Benzo(k) flouranthene	Railways (1A3c)	[kg]	1	1	1	1	1	1	1	1	1	1	1	1	1
Benzo(k) flouranthene	Navigation (1A3d)	[kg]	2	2	2	2	2	2	2	2	2	2	2	2	2
Benzo(k) flouranthene	Comm./Inst. (1A4a)	[kg]	0	0	0	0	0	0	0	0	0	0	0	0	0
Benzo(k) flouranthene	Residential (1A4b)	[kg]	0	0	0	0	0	0	0	0	0	0	0	0	0

Benzo(k) flouranthene	Ag./for./fish. (1A4c)	[kg]	10	10	10	10	9	10	10	10	11	11	11	11	11
Benzo(k) flouranthene	Military (1A5)	[kg]	0	0	0	0	1	1	0	0	0	1	0	1	0
Benzo(k) flouranthene	Navigation int. (1A3d)	[kg]	9	8	7	7	6	5	6	6	6	4	5	5	4
Benzo(k) flouranthene	Civil Aviation int. (1A3a)	[kg]	0	0	0	0	0	0	0	0	0	0	0	0	0
Benzo(a) pyrene	Industry-Other (1A2f)	[kg]	3	3	3	3	3	3	3	4	4	3	3	3	3
Benzo(a) pyrene	Civil Aviation (1A3a)	[kg]	0	0	0	0	0	0	0	0	0	0	0	0	0
Benzo(a) pyrene	Road (1A3b)	[kg]	47	46	47	49	51	52	55	61	62	60	62	64	64
Benzo(a) pyrene	Railways (1A3c)	[kg]	0	0	0	0	0	0	0	0	0	0	0	0	0
Benzo(a) pyrene	Navigation (1A3d)	[kg]	1	1	1	1	1	1	1	1	1	1	1	1	1
Benzo(a) pyrene	Comm./Inst. (1A4a)	[kg]	0	0	0	0	0	0	0	0	0	0	0	0	0
Benzo(a) pyrene	Residential (1A4b)	[kg]	0	0	0	0	0	0	0	0	0	0	0	0	0
Benzo(a) pyrene	Ag./for./fish. (1A4c)	[kg]	5	5	5	5	5	5	5	5	6	6	5	6	6
Benzo(a) pyrene	Military (1A5)	[kg]	0	0	0	0	0	0	0	0	0	0	0	0	0
Benzo(a) pyrene	Navigation int. (1A3d)	[kg]	5	4	4	4	3	3	4	4	3	2	3	3	2
Benzo(a) pyrene	Civil Aviation int. (1A3a)	[kg]	0	0	0	0	0	0	0	0	0	0	0	0	0
Benzo(g,h,i) perylene	Industry-Other (1A2f)	[kg]	5	5	5	5	5	6	6	7	7	5	6	6	6
Benzo(g,h,i) perylene	Civil Aviation (1A3a)	[kg]	0	0	0	0	0	0	0	0	0	0	0	0	0
Benzo(g,h,i) perylene	Road (1A3b)	[kg]	93	90	91	95	98	100	105	115	118	114	117	120	122
Benzo(g,h,i) perylene	Railways (1A3c)	[kg]	0	0	0	0	0	0	0	0	0	0	0	0	0
Benzo(g,h,i) perylene	Navigation (1A3d)	[kg]	8	8	8	7	7	7	7	7	7	7	7	6	5
Benzo(g,h,i) perylene	Comm./Inst. (1A4a)	[kg]	1	1	1	1	1	2	2	2	2	2	2	2	2
Benzo(g,h,i) perylene	Residential (1A4b)	[kg]	0	0	0	0	1	1	1	1	1	1	1	1	1
Benzo(g,h,i) perylene	Ag./for./fish. (1A4c)	[kg]	20	20	21	20	17	20	19	18	20	19	18	18	18
Benzo(g,h,i) perylene	Military (1A5)	[kg]	0	0	0	0	1	1	0	0	0	1	0	0	0
Benzo(g,h,i) perylene	Navigation int. (1A3d)	[kg]	37	32	31	31	24	21	23	21	22	17	20	19	17
Benzo(g,h,i) perylene	Civil Aviation int. (1A3a)	[kg]	0	0	0	0	0	0	0	0	0	0	0	0	0
indeno(1,2,3-c,d) pyrene	Industry-Other (1A2f)	[kg]	3	3	3	3	3	3	3	4	4	3	4	3	3
indeno(1,2,3-c,d) pyrene	Civil Aviation (1A3a)	[kg]	0	0	0	0	0	0	0	0	0	0	0	0	0
indeno(1,2,3-c,d) pyrene	Road (1A3b)	[kg]	51	51	52	54	57	58	61	66	67	65	66	67	68
indeno(1,2,3-c,d) pyrene	Railways (1A3c)	[kg]	0	0	0	0	0	0	0	0	0	0	0	0	0
indeno(1,2,3-c,d) pyrene	Navigation (1A3d)	[kg]	7	7	6	6	6	6	6	6	6	5	5	5	4
indeno(1,2,3-c,d) pyrene	Comm./Inst. (1A4a)	[kg]	0	0	0	0	0	1	1	1	1	1	1	1	1
indeno(1,2,3-c,d) pyrene	Residential (1A4b)	[kg]	0	0	0	0	0	0	0	0	0	0	0	0	0
indeno(1,2,3-c,d) pyrene	Ag./for./fish. (1A4c)	[kg]	15	15	15	15	13	14	14	13	14	14	13	13	12
indeno(1,2,3-c,d) pyrene	Military (1A5)	[kg]	0	0	0	0	0	0	0	0	0	0	0	0	0

indeno(1,2,3-c,d) pyrene	Navigation int. (1A3d)	[kg]	30	26	25	25	20	17	19	17	18	14	17	16	14
indeno(1,2,3-c,d) pyrene	Civil Aviation int. (1A3a)	[kg]	0	0	0	0	0	0	0	0	0	0	0	0	0
PCB	Industry-Other (1A2f)	[g]	6	6	6	6	6	6	7	7	7	5	7	7	7
PCB	Civil Aviation (1A3a)	[g]	0	0	0	0	0	0	0	0	0	0	0	0	0
PCB	Road (1A3b)	[g]	22	22	22	23	24	25	26	26	25	22	23	24	22
PCB	Railways (1A3c)	[g]	2	2	1	2	2	2	2	2	2	2	2	2	2
PCB	Navigation (1A3d)	[g]	0	1	1	1	1	1	1	1	1	1	1	1	1
PCB	Comm./Inst. (1A4a)	[g]	0	0	0	0	0	0	0	0	0	0	0	0	0
PCB	Residential (1A4b)	[g]	0	0	0	0	0	0	0	0	0	0	0	0	0
PCB	Ag./for./fish. (1A4c)	[g]	7	7	7	8	8	8	8	8	9	9	9	9	9
PCB	Military (1A5)	[g]	0	0	0	0	0	0	0	0	0	0	0	0	0
PCB	Navigation int. (1A3d)	[g]	1	1	0	0	0	0	1	1	0	0	0	0	0
PCB	Civil Aviation int. (1A3a)	[g]	0	0	0	0	0	0	0	0	0	0	0	0	0

Annex 3B-17 Uncertainty estimates

Uncertainty estimation, CO₂

Gas		Base year emission	Year t emission	Activity data uncertainty	Emission factor uncertainty	Combined uncertainty	Combined uncertainty as % of total national emissions in year t	Type A sensitivity	Type B sensitivity	Uncertainty in trend in national emissions introduced by activity data uncertainty	Uncertainty in trend in national emissions introduced by emission factor uncertainty	Uncertainty introduced into the trend in total national emissions
		Input data	Input data	Input data	Input data							
		Gg	Gg	%	%	%	%	%	%	%	%	%
Road transport	CO ₂	9284	11224	2	5	5,385	3,947	0,05566424	0,8260	0,2783	2,3362	2,3527
Military	CO ₂	119	116	2	5	5,385	0,041	-0,0013494	0,0085	-0,0067	0,0241	0,0250
Railways	CO ₂	297	249	2	5	5,385	0,088	-0,006261	0,0183	-0,0313	0,0519	0,0606
Navigation (small boats)	CO ₂	48	99	41	5	41,304	0,266	0,0032831	0,0073	0,0164	0,4208	0,4211
Navigation (large vessels)	CO ₂	748	399	11	5	12,083	0,315	-0,0326388	0,0294	-0,1632	0,4568	0,4851
Fisheries	CO ₂	591	479	2	5	5,385	0,168	-0,0137183	0,0353	-0,0686	0,0997	0,1210
Agriculture	CO ₂	1272	1343	24	5	24,515	2,151	-0,0066573	0,0989	-0,0333	3,3557	3,3559
Forestry	CO ₂	36	17	30	5	30,414	0,033	-0,001718	0,0012	-0,0086	0,0527	0,0533
Industry (mobile)	CO ₂	839	1021	41	5	41,304	2,755	0,00555693	0,0752	0,0278	4,3583	4,3584
Residential	CO ₂	39	62	35	5	35,355	0,144	0,00135442	0,0046	0,0068	0,2274	0,2275
Commercial/Institutional	CO ₂	74	171	35	5	35,355	0,396	0,00650002	0,0126	0,0325	0,6244	0,6252
Civil aviation	CO ₂	243	133	10	5	11,180	0,097	-0,0103666	0,0098	-0,0518	0,1380	0,1474
		13.589	15314				28,032					36,3011
Total uncertainties				Year (%):			5,294		Trend (%):			6,025

Uncertainty estimation, CH₄

	Gas	Base year emission	Year t emission	Activity data uncertainty	Emission factor uncertainty	Combined uncertainty	Combined uncertainty as % of total national emissions in year t	Type A sensitivity	Type B sensitivity	Uncertainty in trend in national emissions introduced by activity data uncertainty	Uncertainty in trend in national emissions introduced by emission factor uncertainty	Uncertainty introduced into the trend in total national emissions
		Input data	Input data	Input data	Input data							
		Mg	Mg	%	%	%	%	%	%	%	%	%
Road transport	CH ₄	2233	538	2	40	40,050	22,757	-0,0993825	0,2040	-3,9753	0,5771	4,0170
Military	CH ₄	5	3	2	100	100,020	0,311	0,00042476	0,0011	0,0425	0,0032	0,0426
Railways	CH ₄	12	7	2	100	100,020	0,772	0,00109332	0,0028	0,1093	0,0078	0,1096
Navigation (small boats)	CH ₄	17	24	41	100	108,079	2,756	0,00690284	0,0092	0,6903	0,5309	0,8708
Navigation (large vessels)	CH ₄	16	10	11	100	100,603	1,021	0,00148269	0,0036	0,1483	0,0567	0,1587
Fisheries	CH ₄	13	12	2	100	100,020	1,241	0,0026731	0,0045	0,2673	0,0126	0,2676
Agriculture	CH ₄	105	97	24	100	102,840	10,520	0,02245481	0,0367	2,2455	1,2468	2,5684
Forestry	CH ₄	21	2	30	100	104,403	0,247	-0,0020118	0,0008	-0,2012	0,0360	0,2044
Industry (mobile)	CH ₄	60	36	41	100	108,079	4,087	0,00544761	0,0136	0,5448	0,7874	0,9575
Residential	CH ₄	51	65	35	100	105,948	7,223	0,01753449	0,0245	1,7534	1,2117	2,1314
Commercial/Institutional	CH ₄	99	151	35	100	105,948	16,931	0,04390272	0,0574	4,3903	2,8403	5,2289
Civil aviation	CH ₄	4	2	10	100	100,499	0,246	0,00028945	0,0009	0,0289	0,0124	0,0315
		2636	947				708,427					29,1039
Total uncertainties				Year (%):		26,616		Trend (%):		5,395		

Uncertainty estimation, N₂O

	Gas	Base year emission	Year t emission	Activity data uncertainty	Emission factor uncertainty	Combined uncertainty	Combined uncertainty as % of total national emissions in year t	Type A sensitivity	Type B sensitivity	Uncertainty in trend in national emissions introduced by activity data uncertainty	Uncertainty in trend in national emissions introduced by emission factor uncertainty	Uncertainty introduced into the trend in total national emissions
		Input data	Input data	Input data	Input data							
		Mg	Mg	%	%	%	%	%	%	%	%	%
Road transport	N ₂ O	292	376	2	50	50,040	33,754	0,08278395	0,7732	4,1392	2,1870	4,6814
Military	N ₂ O	4	4	2	1000	1000,002	7,344	-0,0003268	0,0084	-0,3268	0,0238	0,3277
Railways	N ₂ O	8	7	2	1000	1000,002	12,343	-0,0051476	0,0141	-5,1476	0,0400	5,1478
Navigation (small boats)	N ₂ O	1	3	41	1000	1000,840	6,278	0,00422799	0,0072	4,2280	0,4169	4,2485
Navigation (large vessels)	N ₂ O	47	25	11	1000	1000,060	45,088	-0,0594191	0,0517	-59,4191	0,8040	59,4246
Fisheries	N ₂ O	37	30	2	1000	1000,002	54,386	-0,0251682	0,0623	-25,1682	0,1763	25,1689
Agriculture	N ₂ O	49	57	24	1000	1000,288	102,572	0,00132531	0,1175	1,3253	3,9895	4,2039
Forestry	N ₂ O	1	1	30	1000	1000,450	0,974	-0,0002064	0,0011	-0,2064	0,0473	0,2117
Industry (mobile)	N ₂ O	34	43	41	1000	1000,840	77,892	0,00840928	0,0892	8,4093	5,1727	9,8728
Residential	N ₂ O	1	1	35	1000	1000,612	1,935	0,00076078	0,0022	0,7608	0,1097	0,7687
Commercial/Institutional	N ₂ O	1	3	35	1000	1000,612	4,770	0,00287152	0,0055	2,8715	0,2705	2,8842
Civil aviation	N ₂ O	10	7	10	1000	1000,050	11,985	-0,0105364	0,0137	-10,5364	0,1943	10,5382
		486	557				23112,406					4458,1569
Total uncertainties		Year (%):		152,028		Trend (%):		66,769				

Annex 3C - Industrial Processes

No annexes for industrial processes in this submission.

Annex 3D - Solvents and Other Product Use

Annex 3D-1:	Tentative list of chemicals
Annex 3D-2:	NMVOC and CO ₂ equivalent emissions, 1990-2011
Annex 3D-3:	Activity data for solvent and product use, 1990-2011
Annex 3D-4:	Emissions from other product use, 1990-2012
Annex 3D-5:	Activity data for other product use, 1990-2012

Annex 3D-1 Tentative list of chemicals

The National Atmospheric Inventory for Great Britain

(<http://www.naei.org.uk/>) covers the following sectors and chemicals:

Sectors:

Total emission
Energy Production
Comm+ Residn Combustn.
Industrial Combustion
Production Processes
Extr & Distrib of Fossil Fuels

Solvent Use
Road Transport
Other Transp & Mach
Waste Treatment & Disp
Nature (Forests)

Chemicals:

- | | |
|---|---|
| 1 (1-methylethyl)cyclohexane | 2 (1-methylpropyl)cyclohexane |
| 3 (2-methyl-1-propyl)acetate | 4 (2-methylbutyl)cyclohexane |
| 5 (2-methylpropyl)cyclohexane | 6 1-(2-butoxy-1-methyl-ethoxy)-2-propanol |
| 7 1-(2-ethoxy-1-methyl-ethoxy)-2-propanol | 8 1-(2-methoxy-1-methyl-ethoxy)2-propanol |
| 9 1-(butoxyethoxy)-2-propanol | 10 1,1,1-trichloroethane |
| 11 1,1,1-trichlorotrifluoroethane | 12 1,1,2,2-tetrachloroethane |
| 13 1,1,2-trimethylcyclohexane | 14 1,1,2-trimethylcyclopentane |
| 15 1,1,3-trimethylcyclohexane | 16 1,1,4,4-tetramethylcyclohexane |
| 17 1,1-dichloroethane | 18 1,1-dichloroethene |
| 19 1,1-dichlorotetrafluoroethane | 20 1,1-dimethylcyclohexane |
| 21 1,1-dimethylcyclopentane | 22 1,2,3,4-tetrahydronaphthalene |
| 23 1,2,3,4-tetramethylbenzene | 24 1,2,3,5-tetramethylbenzene |
| 25 1,2,3,5-tetramethylcyclohexane | 26 1,2,3-trichlorobenzene |
| 27 1,2,3-trimethylbenzene | 28 1,2,3-trimethylcyclohexane |
| 29 1,2,3-trimethylcyclopentane | 30 1,2,4,4-tetramethylcyclopentane |
| 31 1,2,4,5-tetramethylbenzene | 32 1,2,4-trichlorobenzene |
| 33 1,2,4-trimethylcyclopentane | 34 1,2,4-trimethylbenzene |
| 35 1,2,4-trimethylcyclohexane | 36 1,2,4-trimethylcyclopentane |
| 37 1,2-diaminoethane | 38 1,2-dibromoethane |
| 39 1,2-dichlorobenzene | 40 1,2-dichloroethane |
| 41 1,2-dichloroethene | 42 1,2-dichlorotetrafluoroethane |
| 43 1,2-dimethyl-3-isopropylcyclopentane | 44 1,2-dimethylcyclohexane |
| 45 1,2-dimethylcyclopentane | 46 1,2-ethanedioldiacetate |
| 47 1,2-ethylmethylcyclopentane | 48 1,2-propanediol |
| 49 1,3,4,5,6-pentahydroxy-2-hexanone | 50 1,3,5-trichlorobenzene |
| 51 1,3,5-trimethylbenzene | 52 1,3,5-trimethylcyclohexane |
| 53 1,3-butadiene | 54 1,3-dichlorobenzene |
| 55 1,3-diethylbenzene | 56 1,3-dimethyl-4-ethylbenzene |
| 57 1,3-dimethyl-5-propylbenzene | 58 1,3-dimethylcyclohexane |
| 59 1,3-dimethylcyclopentane | 60 1,3-dioxolane |
| 61 1,3-ethylmethylcyclopentane | 62 1,3-hexadiene |
| 63 1,4-butyrolacetone | 64 1,4-dichlorobenzene |
| 65 1,4-diethylbenzene | 66 1,4-dimethyl-2-isopropylbenzene |
| 67 1,4-dimethylcyclohexane | 68 1,4-dimethylpiperazine |
| 69 1,4-dioxane | 70 11-methyl-1-dodecanol |
| 71 1-butanal | 72 1-butanol |
| 73 1-butene | 74 1-butoxy-2-propanol |
| 75 1-butyne | 76 1-chloro-2,3-epoxypropane |
| 77 1-chloro-4-nitrobenzene | 78 1-chloropropane |
| 79 1-decene | 80 1-ethoxy-2-propanol |

81 1-ethoxy-2-propyl acetate	82 1-ethyl-1,4-dimethylcyclohexane
83 1-ethyl-2,2,6-trimethylcyclohexane	84 1-ethyl-2,3-dimethylbenzene
85 1-ethyl-2,3-dimethylcyclohexane	86 1-ethyl-2-propylbenzene
87 1-ethyl-2-propylcyclohexane	88 1-ethyl-3,5-dimethylbenzene
89 1-ethyl-3-methylcyclohexane	90 1-ethyl-4-methylcyclohexane
91 1-ethylpropylbenzene	92 1-heptene
93 1-hexanal	94 1-hexene
95 1-hydrophenol	96 1-methoxy-2-ethanol
97 1-methoxy-2-propanol	98 1-methoxy-2-propyl acetate
99 1-methyl-1-phenylcyclopropane	100 1-methyl-1-propylcyclopentane
101 1-methyl-2-isopropylbenzene	102 1-methyl-2-propylbenzene
103 1-methyl-3-(isopropyl)benzene	104 1-methyl-3-isopropylcyclopentane
105 1-methyl-3-propylbenzene	106 1-methyl-4-isopropylbenzene
107 1-methyl-4-isopropylcyclohexane	108 1-methyl-4-tertbutylbenzene
109 1-methylbutylbenzene	110 1-methylindan
111 1-methylindene	112 1-nonene
113 1-octene	114 1-pentanal
115 1-pentanol	116 1-pentene
117 1-propanal	118 1-propanol
119 2-(2-aminoethylamino)ethanol	120 2-(2-butoxyethoxy)ethanol
121 2-(2-butoxyethoxy)ethyl acetate	122 2-(2-ethoxyethoxy)ethanol
123 2-(2-ethoxyethoxy)ethyl acetate	124 2-(2-hydroxy-ethoxy)ethanol
125 2-(2-hydroxy-propoxy)-1-propanol	126 2-(methoxyethoxy)ethanol
127 2,2,3,3-tetramethylhexane	128 2,2,4,6,6-pentamethylheptane
129 2,2,4-trimethyl-1,3-pentanediol	130 2,2,4-trimethylpentane
131 2,2,5-trimethylhexane	132 2,2-dimethylbutane
133 2,2-dimethylhexane	134 2,2-dimethylpentane
135 2,2-dimethylpropane	136 2,2'-iminodi(ethylamine)
137 2,2'-iminodiethanol	138 2,3,3,4-tetramethylpentane
139 2,3,3-trimethyl-1-butene	140 2,3,4-trimethylhexane
141 2,3,4-trimethylpentane	142 2,3,5-trimethylhexane
143 2,3-dimethylbutane	144 2,3-dimethylfuran
145 2,3-dimethylheptane	146 2,3-dimethylhexane
147 2,3-dimethylnonane	148 2,3-dimethyloctane
149 2,3-dimethylpentane	150 2,3-dimethylundecane
151 2,4,6-trichloro-1,3,5-triazine	152 2,4-difluoroaniline
153 2,4-dimethyl-1-(1-methylethyl)benzene	154 2,4-dimethylfuran
155 2,4-dimethylheptane	156 2,4-dimethylhexane
157 2,4-dimethylpentane	158 2,4-toluene diisocyanate
159 2,5-dimethyldecane	160 2,5-dimethylfuran
161 2,5-dimethylheptane	162 2,5-dimethylhexane
163 2,5-dimethyloctane	164 2,6-dimethyldecane
165 2,6-dimethylheptane	166 2,6-dimethyloctane
167 2,6-dimethylundecane	168 2,6-toluene diisocyanate
169 2,7-dimethyloctane	170 2-[2-(2-ethoxy-ethoxy)-ethoxy]ethanol
171 2-acetoxy-propyl acetate	172 2-aminoethanol
173 2-butanol	174 2-butanone
175 2-butanone oxime	176 2-butene
177 2-butoxyethanol	178 2-butoxyethyl acetate
179 2-chloroethanol	180 2-chloropropane
181 2-chlorotoluene	182 2-ethoxyethanol
183 2-ethoxyethyl acetate	184 2-ethoxypropanol
185 2-ethyl hexanol	186 2-ethyl-1,3-dimethylbenzene
187 2-ethyltoluene	188 2-hexoxyethanol
189 2-hydrophenol	190 2-isopropoxyethanol
191 2-methoxy-2-methylpropane	192 2-methoxyethanol

193 2-methoxyethyl acetate	194 2-methoxypropane
195 2-methyl benzaldehyde	196 2-methyl-1,3-dioxolane
197 2-methyl-1-butene	198 2-methyl-1-butylbenzene
199 2-methyl-1-pentene	200 2-methyl-1-propanol
201 2-methyl-2,4-pentanediol	202 2-methyl-2-butene
203 2-methyl-2-hexene	204 2-methyl-5-ethyloctane
205 2-methylbutanal	206 2-methylbutane
207 2-methyldecalin	208 2-methyldecane
209 2-methylfuran	210 2-methylheptane
211 2-methylhexane	212 2-methylnonane
213 2-methyloctane	214 2-methylpentane
215 2-methylpropanal	216 2-methylpropane
217 2-methylpropenal	218 2-methylpropene
219 2-methylpropyl acetate	220 2-methylpyridine
221 2-methylundecane	222 2-pentanone
223 2-pentene	224 2-phenoxy ethanol
225 2-phenylpropene	226 2-propanol
227 2-propen-1-ol	228 2-propyl acetate
229 3-(2-hydroxy-propoxy)-1-propanol	230 3,3,4-trimethylhexane
231 3,3,5-trimethylheptane	232 3,3-dimethylheptane
233 3,3-dimethyloctane	234 3,3-dimethylpentane
235 3,4-dimethylheptane	236 3,4-dimethylhexane
237 3,5-dimethyloctane	238 3,6-dimethyloctane
239 3,7-dimethylnonane	240 3A,4,7,7A-tetrahydro-4,7-methanoindene
241 3-chloro-4-fluoropicoline	242 3-chloropropene
243 3-chloropyridine	244 3-ethyl-2-methylheptane
245 3-ethyl-2-methylhexane	246 3-ethylheptane
247 3-ethylhexane	248 3-ethyloctane
249 3-ethylpentane	250 3-ethyltoluene
251 3-hydrophenol	252 3-methyl benzaldehyde
253 3-methyl-1-butene	254 3-methylbutanal
255 3-methylbutanol	256 3-methyldecane
257 3-methylfuran	258 3-methylheptane
259 3-methylhexane	260 3-methylnonane
261 3-methyloctane	262 3-methylpentane
263 3-methylundecane	264 3-pentanone
265 4,4-dimethylheptane	266 4,4'-methylenedianiline
267 4,5-dimethylnonane	268 4,6-dimethylindan
269 4,7-dimethylindan	270 4-4'-methylenediphenyl diisocyanate
271 4-bromophenyl acetate	272 4-chlorotoluene
273 4-ethyl morpholine	274 4-ethyl-1,2-dimethylbenzene
275 4-ethyloctane	276 4-ethyltoluene
277 4-methyl benzaldehyde	278 4-methyl-1,3-dioxol-2-one
279 4-methyl-1-pentene	280 4-methyl-2-pentanol
281 4-methyl-2-pentanone	282 4-methyl-4-hydroxy-2-pentanone
283 4-methyldecane	284 4-methylheptane
285 4-methylnonane	286 4-methyloctane
287 4-methylpentene	288 4-propylheptane
289 5-methyl-2-hexanone	290 5-methyldecane
291 5-methylnonane	292 5-methylundecane
293 6-ethyl-2-methyldecane	294 6-ethyl-2-methyloctane
295 6-methylundecane	296 8-methyl-1-nonanol
297 acenaphthene	298 acenaphthylene
299 acetaldehyde	300 acetic acid
301 acetic anhydride	302 acetone
303 acetonitrile	304 acetyl chloride

305 acetylene	306 acrolein
307 acrylamide	308 acrylic acid
309 acrylonitrile	310 aniline
311 anthanthrene	312 anthracene
313 atrazine	314 benzaldehyde
315 benzene	316 benzene-1,2,4-tricarboxylic acid 1,2-
317 benzo (a) anthracene	318 benzo (a) pyrene
319 benzo (b) fluoranthene	320 benzo (c) phenanthrene
321 benzo (e) pyrene	322 benzo (g,h,i) fluoranthene
323 benzo (g,h,i) perylene	324 benzo (k) fluoranthene
325 benzophenone	326 benzopyrenes
327 benzyl alcohol	328 benzyl chloride
329 biphenyl	330 bis(2-hydroxyethyl)ether
331 bis(chloromethyl)ether	332 bis(tributyltin) oxide
333 bromoethane	334 bromoethene
335 bromomethane	336 butane
337 butanethiols	338 butene
339 butoxyl	340 butyl acetate
341 butyl acrylate	342 butyl glycolate
343 butyl lactate	344 butylbenzene
345 butylcyclohexane	346 butyrolactone
347 C10 alkanes	348 C10 alkenes
349 C10 aromatic hydrocarbons	350 C10 cycloalkanes
351 C11 alkanes	352 C11 alkenes
353 C11 aromatic hydrocarbons	354 C11 cycloalkanes
355 C12 alkanes	356 C12 cycloalkanes
357 C13 alkanes	358 C13+ alkanes
359 C13+ aromatic hydrocarbons	360 C14 alkanes
361 C15 alkanes	362 C16 alkanes
363 C2-alkyl-anthracenes	364 C2-alkyl-benzanthracenes
365 C2-alkyl-benzophenanthrenes	366 C2-alkyl-chrysenes
367 C2-alkyl-phenanthrenes	368 C5 alkenes
369 C6 alkenes	370 C7 alkanes
371 C7 alkenes	372 C7 cycloalkanes
373 C8 alkanes	374 C8 alkenes
375 C8 cycloalkanes	376 C9 alkanes
377 C9 alkenes	378 C9 aromatic hydrocarbons
379 C9 cycloalkanes	380 camphor/fenchone
381 carbon disulphide	382 carbon tetrachloride
383 carbonyl sulphide	384 chlorobenzene
385 chlorobutane	386 chlorocyclohexane
387 chlorodifluoromethane	388 chloroethane
389 chloroethene	390 chloroethylene
391 chlorofluoromethane	392 chloromethane
393 chrysene	394 cis-1,3-dimethylcyclopentane
395 cis-2-butene	396 cis-2-hexene
397 cis-2-pentene	398 coronene
399 crotonaldehyde	400 cycloheptane
401 cyclohexanamine	402 cyclohexane
403 cyclohexanol	404 cyclohexanone
405 cyclopenta (c,d) pyrene	406 cyclopenta-anthracenes
407 cyclopentane	408 cyclopenta-phenanthrenes
409 cyclopentene	410 decalin
411 decane	412 diacetoneketogulonic acid
413 diazinon	414 dibenzanthracenes
415 dibenzo (a,h) anthracene	416 dibenzopyrenes

417 dichlorobutenes	418 dichlorodifluoromethane
419 dichlorofluoromethane	420 dichloromethane
421 dichlorvos	422 diethyl disulphide
423 diethyl ether	424 diethyl sulphate
425 diethylamine	426 diethylbenzene
427 difluoromethane	428 dihydroxyacetone
429 diisopropyl ether	430 diisopropylbenzene
431 dimethoxymethane	432 dimethyl disulphide
433 dimethyl esters	434 dimethyl ether
435 dimethyl sulphate	436 dimethyl sulphide
437 dimethylamine	438 dimethylbutene
439 dimethylcyclopentane	440 dimethylformamide
441 dimethylhexene	442 dimethylnonane
443 dimethylpentane	444 dipentene
445 dipropyl ether	446 dodecane
447 ethane	448 ethanethiol
449 ethanol	450 ethofumesate
451 ethyl acetate	452 ethyl acrylate
453 ethyl butanoate	454 ethyl chloroformate
455 ethyl hexanol	456 ethyl lactate
457 ethyl pentanoate	458 ethyl propionate
459 ethylamine	460 ethylbenzene
461 ethylcyclohexane	462 ethylcyclopentane
463 ethyldimethylbenzene	464 ethylene
465 ethylene glycol	466 ethylene oxide
467 ethylisopropylbenzene	468 fenitrothion
469 fluoranthene	470 fluorene
471 formaldehyde	472 formanilide
473 formic acid	474 fumaric acid
475 glycerol	476 glyoxal
477 heptadecane	478 heptane
479 hexachlorocyclohexane	480 hexachloroethane
481 hexadecane	482 hexafluoropropene
483 hexamethylcyclotrisiloxane	484 hexamethyldisilane
485 hexamethyldisiloxane	486 hexamethylenediamine
487 hexane	488 hexylcyclohexane
489 indan	490 indeno (1,2,3-c,d) pyrene
491 iodomethane	492 isobutylbenzene
493 isobutylcyclohexane	494 isopentylbenzene
495 isophorone	496 isoprene
497 isoprene + BVOC (1)	498 isopropylbenzene
499 isopropylcyclohexane	500 limonene
501 malathion	502 maleic anhydride
503 m-cresol	504 menthene
505 methacrylic acid	506 methanethiol
507 methanol	508 methyl acetate
509 methyl acrylate	510 methyl butanoate
511 methyl ethyl ether	512 methyl formate
513 methyl glyoxal	514 methyl methacrylate
515 methyl naphthalenes	516 methyl pentanoate
517 methyl styrene	518 methylamine
519 methyl-anthracenes	520 methyl-benzanthracenes
521 methyl-benzphenanthrenes	522 methylcyclodecane
523 methylcyclohexane	524 methylcyclopentane
525 methylethylbenzene	526 methyl-fluoranthenes
527 methylhexane	528 methylindane

529 methyl-phenanthrenes	530 methylpropene
531 methylpropylbenzene	532 methyltetralin
533 m-xylene	534 N-(hydroxymethyl) acrylamide
535 N,N-diethyl benzenamine	536 N,N-dimethyl benzenamine
537 naphthalene	538 naphthol
539 Nedocromil Sodium	540 nitrobenzene
541 nitromethane	542 nitropentane
543 nitropropane	544 N-methyl pyrrolidone
545 nonane	546 o-cresol
547 octahydroindan	548 octamethylcyclotetrasiloxane
549 octane	550 octylamine
551 o-xylene	552 palmitic acid
553 p-benzoquinone	554 p-cresol
555 pentadecane	556 pentafluoroethane
557 pentane	558 pentanethiols
559 pentylbenzene	560 pentylcyclohexane
561 permethrin	562 perylene
563 phenol	564 phenoxyacetic acid (phenoxy acid)
565 phenylacetic acid	566 phenylacetone nitrile
567 phthalic anhydride	568 pine oil
569 polyethylene glycol	570 polyisobutene
571 polyvinyl chloride	572 potassium phenylacetate
573 propadiene	574 propane
575 propanetriol	576 propanoic acid
577 propionitrile	578 propyl acetate
579 propyl butanoate	580 propyl propionate
581 propylamine	582 propylbenzene
583 propylcyclohexane	584 propylcyclopentane
585 propylene	586 propylene oxide
587 propyne	588 p-xylene
589 pyrene	590 pyridine
591 salicylic acid	592 sec-butylbenzene
593 sec-butylcyclohexane	594 simazine
595 sodium 2-ethylhexanoate	596 sodium acetate
597 sodium phenylacetate	598 styrene
599 sulphanilamide	600 terpenes
601 tert-butylamine	602 tert-butylbenzene
603 tert-butylcyclohexane	604 tert-butylcyclopropane
605 tert-pentylbenzene	606 tetrachloroethene
607 tetradecane	608 tetrafluoroethene
609 tetrahydrofuran	611 tetramethylcyclohexane
612 toluene	613 toluene-2,3-diamine
614 toluene-2,4-diamine	615 toluene-2,4-diisocyanate
616 toluene-2,5-diamine	617 toluene-2,6-diamine
618 toluene-2,6-diisocyanate	619 toluene-3,4-diamine
620 toluene-3,5-diamine	621 trans-2-butene
622 trans-2-hexene	623 trans-2-pentene
624 trans-3-hexene	625 trialkyl phosphate
626 trichloroethene	627 trichlorofluoromethane
628 trichloromethane	629 tridecane
630 triethanolamine	631 triethylamine
632 trifluoroethene	633 trifluoromethane
634 trifluralin	635 trimethylamine
636 trimethylfluorosilane	637 tri-n-butyl phosphate
638 undecane	639 unspeciatiated alcohols
640 unspeciatiated aliphatic hydrocarbons	641 unspeciatiated alkanes

642 unspciated alkenes
644 unspciated aromatic hydrocarbons
646 unspciated cycloalkanes
648 unspciated ketones
650 vinyl acetate

643 unspciated amines
645 unspciated carboxylic acids
647 unspciated hydrocarbons
649 urea
(1) BVOC- biogenic VOCs, such as alpha-pinene and other terpenes

Annex 3D-2 NMVOC and CO₂ equivalent emissions, 1990-2011

Table 3D-2a NMVOC emissions (Gg per year), 1990-1999.

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
3A Paint Application	5,11	5,83	6,37	5,74	6,38	5,90	7,13	6,05	5,95	6,43
3B Degreasing, Dry cleaning and Electronics	7E-05	7E-05	7E-05	7E-05	9E-05	8E-05	7E-05	4E-05	5E-05	3E-05
3C Chemical Products Manufacturing and Processing	8,14	9,32	9,13	7,15	9,25	9,32	9,48	8,04	7,66	7,30
3D Other Use of Solvents and Products	24,7	27,7	30,0	26,6	31,4	30,0	32,8	30,6	27,9	26,9
3D3 Other Product Use	0,083	0,076	0,087	0,077	0,073	0,079	0,084	0,094	0,084	0,087
Total NMVOC	38,0	43,0	45,5	39,5	47,1	45,3	49,5	44,8	41,6	40,8
Total CO ₂ -eqv.	93,5	106	113	97,9	113	109	121	108	101	100

Table 3D-2b NMVOC emissions (Gg per year), 2000-2009.

	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
3A Paint Application	6,40	5,25	5,18	4,99	4,66	4,32	3,73	3,23	3,38	2,85
3B Degreasing, Dry cleaning and	3E-05	1E-05	3E-05	3E-05	2E-05	2E-05	1E-05	2E-05	2E-05	1E-05
3C Chemical Products Manufactur-										
ing and Processing	6,96	6,28	6,58	4,96	6,06	6,25	6,02	6,12	5,91	4,99
3D Other Use of Solvents and										
Products	27,8	24,8	24,4	22,5	21,4	20,8	20,8	18,0	18,4	19,7
3D3 Other Product Use	0,0950	0,0855	0,102	0,115	0,102	0,0950	0,109	0,0838	0,0775	0,0804
Total NMVOC	41,3	36,4	36,2	32,5	32,3	31,5	30,7	27,5	27,8	27,6
Total CO ₂ -eqv.	100	88,1	88,4	80,3	78,2	75,7	71,7	64,3	65,7	65,1

Table 3D-2c NMVOC emissions (Gg per year), 2010-2011.

	2010	2011	2012
3A Paint Application	2,75	2,87	2,86
3B Degreasing, Dry cleaning and Electronics	1E-05	1E-05	3E-06
3C Chemical Products Manufacturing and Processing	5,05	4,81	4,87
3D Other Use of Solvents and Products	19,4	19,3	19,1
3D3 Other Product Use	0,0678	0,0601	0,0817
Total NMVOC	27,3	27,0	27,0
Total CO ₂ -eqv.	63,6	63,4	63,3

Annex 3D-3 Activity data for solvent and product use, 1990-2011

Table 3D-4 Activity data for product use (Gg per year), 1990-2011.

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
3A Paint Application	83,2	96,5	103	90,4	95,7	92,1	110	95,5	97,6	101
3B Degreasing, Dry cleaning and Electronics	1,41	1,31	1,25	1,45	1,88	1,53	1,48	0,892	1,10	0,690
3C Chemical Products Manufacturing and Processing	406	455	569	389	466	504	523	519	528	488
3D Other Use of Solvents and Products	197	224	234	202	239	247	260	249	234	226
3D3 Other Product Use	28,0	28,5	33,1	29,5	30,5	31,4	32,5	35,5	38,6	45,1
Total products	717	805	940	712	833	877	927	900	899	861
<i>Continued</i>	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
3A Paint Application	105	86,4	86,7	79,8	77,7	75,2	64,7	57,3	58,1	48,7
3B Degreasing, Dry cleaning and Electronics	0,586	0,251	0,597	0,578	0,481	0,365	0,292	0,433	0,299	0,263
3C Chemical Products Manufacturing and Processing	567	551	540	513	634	740	749	814	771	683
3D Other Use of Solvents and Products	230	206	218	185	182	204	180	162	169	179
3D3 Other Product Use	46,5	42,0	56,3	61,8	61,4	63,4	63,6	58,5	51,2	52,2
Total products	949	886	902	841	956	1083	1058	1093	1049	963
<i>Continued</i>	2010	2011	2012							
3A Paint Application	45,8	43,8	43,3							
3B Degreasing, Dry cleaning and Electronics	0,247	0,224	0,0547							
3C Chemical Products Manufacturing and Processing	641	640	516							
3D Other Use of Solvents and Products	170	169	169							
3D3 Other Product Use	57,7	49,9	53,6							
Total products	914	903	782							

Annex 3D-4 Emissions from other product use, 1990-2012

Table 3D-5a Emissions of CO₂ and N₂O from other product use, 1990-2009

		1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
CO ₂ emission from											
3D1 & 3D4	Gg	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
Use of fireworks	Gg	0.06	0.07	0.08	0.07	0.08	0.13	0.12	0.09	0.15	0.29
Use of tobacco	Gg	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
Use of candles	Gg	21.66	25.12	27.96	27.06	32.41	26.46	24.87	25.28	39.98	47.31
Use of charcoal for BBQ	Gg	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
Total	Gg	21.72	25.19	28.04	27.13	32.49	26.59	24.98	25.37	40.14	47.60
N ₂ O emission from											
3D1 & 3D4	Mg	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
Use of fireworks	Mg	2.47	3.28	3.54	3.13	3.80	5.80	5.32	4.19	6.82	12.91
Use of tobacco	Mg	0.82	0.76	0.77	0.74	0.73	0.73	0.71	0.71	0.71	0.72
Use of candles	Mg	0.18	0.21	0.23	0.22	0.27	0.22	0.21	0.21	0.33	0.39
Use of charcoal for BBQ	Mg	0.22	0.19	0.29	0.21	0.18	0.24	0.30	0.40	0.31	0.33
Total	Mg	3.61	4.36	4.75	4.23	4.90	6.91	6.47	5.44	8.10	14.28
CO ₂ -equivalents	Gg	22.8	26.5	29.5	28.4	34.0	28.7	27.0	27.1	42.6	52.0
<i>Continued</i>		2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
CO ₂ emission from											
3D1 & 3D4	Gg	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
Use of fireworks	Gg	0.21	0.17	0.20	0.26	0.37	0.16	0.18	0.19	0.19	0.23
Use of tobacco	Gg	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
Use of candles	Gg	49.26	47.54	70.81	71.12	74.19	100.20	85.09	93.44	78.13	75.02
Use of charcoal for BBQ	Gg	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
Total	Gg	49.47	47.71	71.01	71.38	74.57	100.36	85.27	93.64	78.32	75.25
N ₂ O emission from											
3D1 & 3D4	Mg	2.1	7.8	8.6	13.3	19.1	34.2	35.6	40.7	33.0	45.8
Use of fireworks	Mg	9.39	7.41	9.17	11.71	16.72	7.13	8.15	8.66	8.45	10.41
Use of tobacco	Mg	0.73	0.70	0.70	0.72	0.71	0.66	0.66	0.63	0.61	0.60
Use of candles	Mg	0.41	0.39	0.58	0.59	0.61	0.83	0.70	0.77	0.64	0.62
Use of charcoal for BBQ	Mg	0.40	0.33	0.49	0.60	0.49	0.45	0.59	0.36	0.31	0.35
Total	Mg	13.03	16.63	19.54	26.92	37.63	43.27	45.70	51.12	43.02	57.90
CO ₂ equivalents	Gg	53.51	52.87	77.06	79.73	86.24	113.77	99.40	109.47	91.62	104.88

Table 3D-5b Emissions of CO₂ and N₂O from other product use, 2000-2012

<i>Continued</i>		2010	2011	2012
CO ₂ emission from				
3D1 & 3D4	Gg	NO	NO	NO
Use of fireworks	Gg	0.23	0.20	0.15
Use of tobacco	Gg	NO	NO	NO
Use of candles	Gg	102.70	87.80	81.10
Use of charcoal for BBQ	Gg	NO	NO	NO
Total	Gg	102.94	88.00	81.25
N ₂ O emission from				
3D1 & 3D4	Mg	34.4	42.0	30.1
Use of fireworks	Mg	10.49	9.15	6.74
Use of tobacco	Mg	0.59	0.53	0.52
Use of candles	Mg	0.85	0.72	0.67
Use of charcoal for BBQ	Mg	0.24	0.20	0.42
Total	Mg	46.63	52.64	38.5
CO ₂ equivalents	Gg	124.1	103.7	93.19

Annex 3D-5 Activity data for other product use, 1990-2012

Table 3D-6 Activity data for the national use of other products, 1990-2012.

		1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
3D1 & 3D4	Gg	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
Fireworks	Gg	1.3	1.7	1.8	1.6	2.0	3.0	2.8	2.2	3.5	6.7
Tobacco	Gg	12.7	12.0	12.1	11.5	11.4	11.4	11.0	11.2	11.1	11.2
Candles	Gg	7.4	8.6	9.6	9.3	11.1	9.1	8.5	8.7	13.7	16.3
BBQ	Gg	7.2	6.2	9.5	7.1	6.0	7.9	10.2	13.5	10.2	11.0
<i>Continued</i>		2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
3D1 & 3D4	Gg	0.0021	0.0078	0.0086	0.0133	0.0191	0.0342	0.0356	0.0407	0.0330	0.0458
Fireworks	Gg	4.9	3.8	4.7	6.1	8.6	3.7	4.2	4.5	4.4	5.4
Tobacco	Gg	11.4	10.9	10.9	11.3	11.1	10.4	10.3	9.8	9.6	9.4
Candles	Gg	16.9	16.3	24.3	24.4	25.5	34.4	29.2	32.1	26.8	25.8
BBQ	Gg	13.4	10.9	16.4	20.0	16.2	14.9	19.8	12.2	10.4	11.6
<i>Continued</i>		2010	2011	2012							
3D1 & 3D4	Gg	0.0344	0.0420	0.0301							
Fireworks	Gg	5.4	4.7	3.5							
Tobacco	Gg	9.2	8.3	8.2							
Candles	Gg	35.3	30.2	27.9							
BBQ	Gg	7.8	6.7	14.0							

Annex 3E - Agriculture

List of tables

Table 3E-1 Changes in housing type 1990 – 2012

Table 3E-2 Number of animals allocated on subcategories for 1990-2012, 1 000 head.

Table 3E-3 (a-d) NH₃ emission factors for housing units, 2012.

Table 3E-4 NH₃ emission factors for storage units, 2012.

Table 3E-5 Parameters for winter feeding plans.

Table 3E-6 Energy factors used for GE.

Table 3E-7 Feed intake 1990-2012, FU per animal per year.

Table 3E-8 Grazing animals 1990 – 2012, number of days on grass per year.

Table 3E-9a Average gross energy intake (GE) 1990 – 2012, MJ per head per day.

Table 3E-9b Average gross energy intake (GE) 1990 – 2012, MJ per head per day – Subcategories for cattle and swine.

Table 3E-10a VS daily excretion (average) 1990 – 2012, kg dm per head per day – CRF categories.

Table 3E-10b VS daily excretion (average) 1990 – 2012, kg dm per head per day – Subcategories.

Table 3E-11 Basic data from Sommer et al. (2001) used to estimation of lower emission of N₂O and CH₄ from biogas treated slurry.

Table 3E-12a Calculation of lower CH₄ emission as a consequence of biogas treated slurry.

Table 3E-12b Calculation of lower N₂O emission as a consequence of biogas treated slurry.

Table 3E-13 Background data for calculation of N content in nitrogen fixing crops.

Table 3E-14 Estimated share of nitrogen fixing plants in crops.

Table 3E-15 Area of N-fixing crops 1990-2012, ha

Table 3E-16 Background data for estimation of N₂O emission from crop residue 2012.

Table 3E-17 Area of agricultural land, 1990 – 2012, ha

Table 3E-18 Harvested straw, 1990 – 2012, 1 000 tonnes

Figure 3E-1 Model calculation of nitrogen leaching from groundwater nationwide by SKEP/DAISY and N-LES.

3E-Table 18 QA/QC procedure, stage I – III

Table 3E-1 Changes in housing type 1990 – 2012.

Dairy cattle:

Livestock categories	Housing type	1990	1995	2000	2005	2008	2009	2010	2011	2012
Dairy cattle	Tethered with urine and solid manure	35	31	18	6	6	5	5	4	3
	Tethered with slurry	44	42	28	14	9	7	7	6	5
	Loose-holding with beds, slatted floor	13	17	34	43	44	45	45	46	46
	Loose-holding with beds, slatted floor, scrape	1	1	3	11	14	14	14	15	15
	Loose-holding with beds, solid floor	4	3	6	16	20	21	21	21	22
	Loose-holding with beds, drained floor	0	0	0	0	0	0	0	0	0
	Loose-holding with beds, solid floor with tilt	0	0	0	0	1	2	2	3	3
	Deep litter (all)	0	0	0	1	2	2	2	2	3
	Deep litter, long eating space, solid floor	1	1	3	2	1	1	1	1	1
	Deep litter, slatted floor	2	5	7	5	2	2	2	1	1
	Deep litter, slatted floor, scrape	0	0	1	1	2	2	2	1	1

Heifers:

Livestock categories	Housing type	1990	1995	2000	2005	2008	2009	2010	2011	2012
Heifer calves, 0-6 mth.	Deep litter (boxes)	100	100	100	85	96	96	96	96	96
	Deep litter, solid floor	0	0	0	15	4	4	4	4	4
Heifer, 6 mth.-calving	Tethered with urine and solid manure	19	14	9	6	6	6	6	5	4
	Tethered with slurry	19	14	9	4	2	2	2	2	2
	Slatted floor-boxes	40	35	32	32	37	35	35	31	30
	Loose-housing with beds, slatted floor	4	7	14	19	14	16	16	19	21
	Loose-housing with beds, slatted floor, scrape	0	0	0	2	6	6	6	7	8
	Loose-housing with beds, solid floor	0	0	0	2	6	6	6	7	7
	Deep litter (all)	3	1	0	8	22	22	22	21	21
	Deep litter, long eating space, solid floor	1	2	3	2	2	2	2	2	2
	Deep litter, solid floor	9	19	25	19	1	1	1	1	1
	Deep litter, slatted floor	4	7	6	4	2	2	2	2	2
	Deep litter, slatted floor, scrape	1	1	2	2	2	2	2	3	2

Bulls:

Livestock categories	Housing type	1990	1995	2000	2005	2008	2009	2010	2011	2012
Bull calves, 0-6 mth.	Deep litter (boxes)	100	100	100	84	97	97	97	97	97
	Deep litter, solid floor	0	0	0	16	3	3	3	3	3
Bull, 6 mth - 440 kg	Tethered with urine and solid manure	20	14	10	6	4	3	3	2	2
	Tethered with slurry	20	14	10	5	1	1	1	1	1
	Slatted floor-boxes	41	37	33	29	30	27	27	25	23
	Loose-housing with beds, slatted floor	0	0	0	0	0	0	0	3	5
	Loose-housing with beds, slatted floor, scrape	0	0	0	0	0	0	0	3	3
	Loose-housing with beds, solid floor	0	0	0	0	0	0	0	0	1
	Deep litter (all)	3	1	0	19	58	61	61	58	57
	Deep litter, long eating space, solid floor	1	3	3	2	1	1	1	1	1
	Deep litter, solid floor	10	20	33	34	4	4	4	4	4
	Deep litter, slatted floor	4	9	9	4	1	2	2	1	1
	Deep litter, slatted floor, scrape	1	2	2	1	1	1	1	2	2

Table 3E-1 - *Continued* - Changes in housing type 1990 – 2012 – **Cattle.*****Suckling cattle:***

Livestock categories	Housing type	1990	1995	2000	2005	2008	2009	2010	2011	2012
Suckling cattle	Tethered with urine and solid manure	10	10	9	9	16	15	15	13	13
	Tethered with slurry	0	0	0	3	9	9	9	10	9
	Deep litter (all)	73	55	45	51	68	68	69	69	70
	Deep litter, long eating space, solid floor	0	0	0	0	1	1	1	1	1
	Deep litter, solid floor	17	35	46	35	2	3	3	3	3
	Deep litter, slatted floor	0	0	0	1	1	1	1	2	2
	Deep litter, slatted floor, scrape	0	0	0	1	2	2	2	2	2
	Boxes with sloping bedded floor	0	0	0	0	1	1	0	0	0

Table 3E-1 - *Continued* - Changes in housing type 1990 – 2012 – **Swine.**

Livestock categories	Housing type	1990	1995	2000	2005	2008	2009	2010	2011	2012
Sows	Full slatted floor	11	15	17	14	14	15	15	15	14
	Partly slatted floor	56	59	59	65	77	77	77	79	79
	Solid floor	29	16	6	4	1	1	1	0	0
	Deep litter	4	5	7	5	2	1	1	1	1
	Deep litter + slatted floor	0	2	4	6	4	4	4	4	4
	Deep litter + solid floor	0	2	4	4	1	1	1	1	1
	Outdoor sows	0	1	3	2	1	1	1	0	1
Weaners	Fully slatted floor	54	51	38	29	23	22	22	20	19
	Partly slatted floor	20	31	47	57	67	68	68	70	72
	Solid floor	21	11	5	4	1	0	0	1	0
	Deep litter (to-climate housings)	5	5	5	4	2	2	2	1	1
	Deep litter + slatted floor	0	2	5	0	0	0	0	0	0
	Partly slatted and drained floor	0	0	0	6	7	8	8	8	8
Fattening pigs	Fully slatted floor	51	60	58	53	53	54	54	53	52
	Partly slatted floor	23	24	31	0	0	0	0	0	0
	Solid floor	22	11	5	3	3	2	2	1	1
	Deep litter	4	3	1	2	3	2	2	2	2
	Partly slatted floor and partly deep litter	0	2	5	4	0	0	0	1	0
	Partly slatted and drained floor	0	0	0	38	41	42	42	43	45

Table 3E-1 - *Continued* Changes in housing type 1990 – 2012 – **Poultry.**

Livestock categories	1990	1995	2000	2005	2008	2009	2010	2011	2012
Free-range hens	0	5	9	8	6	6	7	8	7
Organic hens	0	3	12	14	16	15	15	16	19
Barn hens	5	15	17	25	19	19	17	17	19
Battery hens, manure shed	24	26	29	32	42	44	45	45	46
Battery hens, manure tank	13	8	5	5	8	7	7	8	5
Battery hens, manure cellar	58	43	28	16	9	9	9	6	4
Hens for production of brood egg	100	100	100	100	100	100	100	100	100
Pullet, consumption, net	17	12	8	6	7	7	7	19	32
Pullet, consumption, floor	57	63	69	70	84	78	78	76	63
Pullet, brood egg, floor	26	25	23	24	9	15	15	5	5
Broilers, (conv. 30 days)	0	0	0	0	0	0	0	0	0
Broilers, (conv. 32 days)	0	0	0	4	2	7	3	11	14
Broilers, (conv. 35 days)	0	0	0	45	49	57	76	86	81
Broilers, (conv. 40 days)	100	100	100	49	49	36	21	3	5
Broilers, (conv. 45 days)	0	0	0	2	0	0	0	0	0
Broilers, barn (56 days)	0	0	0	0	0	0	0	0	0
Organic broilers (81 days)	0	0	0	0	0	0	0	0	0
Turkey, male	50	50	50	50	50	50	50	50	50
Turkey, female	50	50	50	50	50	50	50	50	50
Ducks	100	100	100	100	100	100	100	100	100
Geese	100	100	100	100	100	100	100	100	100
Pheasant	100	100	100	100	100	100	100	100	100

Table 3E-1 - *Continued* Changes in housing type 1990 – 2012 - **Fur farming.**

Livestock categories	Housing type	1990	1995	2000	2005	2008	2009	2010	2011	2012
Mink	Slurry system	18	25	42	73	92	95	97	96	97
	Solid manure and urine	82	75	58	27	8	5	3	4	3
Foxes	Slurry system	0	0	2	0	0	0	0	0	0
	Solid manure and urine	100	100	98	100	100	100	100	100	100

Table 3E-1 - *Continued* Changes in housing type 1990 – 2012 - **Horses, sheep, goats and ostrich.**

Livestock categories	Housing type	1990	1995	2000	2005	2008	2009	2010	2011	2012
Horses, sheep, goats, ostrich	Deep litter	100	100	100	100	100	100	100	100	100

Table 3E-1 - *Continued* Changes in housing type 1990 – 2012 - **Deer and pheasant.**

Livestock categories	Housing type	1990	1995	2000	2005	2008	2009	2010	2011	2012
Deer and pheasant	Pasture	100	100	100	100	100	100	100	100	100

Reference: 1990 – 2004 = The Danish Agricultural Advisory Service, 2005-2012 = Danish AgriFish Agency.

For all years see: <http://envs.au.dk/videnudveksling/luft/emissioner/reportingsectors/agriculture/>.

Table 3E-2 Number of animals allocated on subcategories for 1990-2012, 1 000 head.

	1990	1995	2000	2005	2008	2009	2010	2011	2012
Dairy Cattle	753	702	636	564	558	563	568	565	587
<u>Non-Dairy Cattle</u>									
Bulls 0-6	217	190	150	132	125	117	132	132	133
Bulls 6-	263	213	176	142	139	145	141	138	133
Heifers 0-6	225	215	185	148	151	151	156	158	162
Heifers 6-	695	647	598	483	485	468	473	476	494
Suckling Cattle	87	122	125	101	107	96	101	99	97
<u>Sheep</u>	92	81	112	126	117	116	111	94	90
<u>Goats</u>									
Meat goat	7	7	8	5	10	11	10	8	10
Milk goat	IE	IE	IE	4	4	4	5	4	2
Mohair goat	IE	IE	IE	2	1	1	1	1	0
<u>Horses</u>									
< 300 kg	IE	IE	IE	44	48	44	41	39	39
300-500 kg	80	84	89	60	65	60	56	53	53
500-700 kg	51	54	57	67	72	67	63	59	59
> 700 kg	4	4	5	5	6	5	5	5	5
<u>Swine</u>									
Sows	904	1 015	1 083	1 151	1 059	1 088	1 117	1 063	1 011
Weaners	4 881	5 613	5 330	6 165	5 893	5 882	6 166	6 061	5 847
Fattening pigs	3 712	4 456	5 508	6 218	5 785	5 399	5 890	5 809	5 473
<u>Poultry</u>									
Hens	4 381	4 366	3 720	3 241	3 590	3 345	3 970	3 882	4 037
Pullets	1 315	1 723	1 216	1 928	1 384	1 092	1 278	1 796	1 561
Broilers	9 802	12 585	16 047	11 905	9 737	14 787	12 836	12 528	12 576
Other poultry	750	946	849	456	396	382	431	450	543
<u>Pheasant</u>									
Pheasant hen	63	63	63	63	63	63	63	63	63
Pheasant chicken	1 000	1 000	1 000	1 000	1 000	1 000	1 000	1 000	1 000
<u>Ostrich</u>									
Ostrich hen	NO	0.11	0.30	0.12	0.02	0.01	0.01	0.01	0.01
Ostrich chicken	NO	3.22	8.59	3.54	0.45	0.35	0.35	0.18	0.17
<u>Fur Farming</u>									
Mink	2 233	1 834	2 188	2 547	2 807	2 720	2 698	2 754	2 948
Foxes	31	16	11	5	3	1	2	2	0
<u>Deer</u>	10	10	10	10	10	9	10	8	7

IE = Included else were (main category).

For all years see <http://envs.au.dk/videnudveksling/luft/emissioner/reportingsectors/agriculture/>

Table 3E-3 (a-d) NH₃ emission factors for housing units, 2012.**a) Cattle**

		Urine TAN	Slurry TAN	Solid manure Total N	Deep litter manure Total N
Housing type		pct. loss of TAN ex animal		pct. loss of N ex animal	
Tethered	urine and solid manure	10	-	5	-
	slurry manure	-	6	-	-
Loose-housing with beds	slatted floor	-	16	-	-
	slatted floor and scrape	-	12	-	-
	solid floor	-	20	-	-
	drained floor	-	8	-	-
	solid floor with tilt and scrape	-	8	-	-
	solid floor with tilt	-	12	-	-
Deep litter	All	-	-	-	6
	solid floor	-	-	-	6
	slatted floor	-	16	-	6
	slatted floor and scrape	-	12	-	6
	solid floor and scrape	-	20	-	6
Boxes	sloping bedded floor	-	16	-	-
	slatted floor	-	16	-	-

b) Swine

5) Swine

			Urine TAN	Slurry TAN	Solid manure Total N	Deep litter Total N
	Housing type	Floor or manure type	Pct. loss of TAN ex animal		pct. loss of N ex animal	
<u>Sows</u>	Individual, mating and gestation	Partly slatted floor	-	13	-	-
		Full slatted floor	-	19	-	-
		Solid floor	21	-	16	-
	Group, mating and gestation	Deep litter	-	-	-	15
		Deep litter + slatted floor	-	16	-	15
		Deep litter + solid floor	-	19	-	15
	Farrowing crate	Partly slatted floor	-	16	-	-
		Full slatted floor	-	13	-	-
		Partly slatted floor	-	26	-	-
	Farrowing pen	Solid floor	20	-	15	-
Partly slatted floor		-	22	15	-	
<u>Weaners</u>		Full slatted floor	-	24	-	-
		Drained + partly slatted floor	-	21	-	-
		Deep litter (to-climate housings)	-	10	-	15
		Solid floor	37	-	25	-
		Deep litter	-	-	-	15
<u>Fattening pigs</u>		Partly slatted floor (50-75 % solid)	-	13	-	-
		Partly slatted floor (25-49% solid)	-	17	-	-
		Drained + partly slatted floor	-	21	-	-
		Full slatted floor	-	24	-	-
		Solid floor	27	-	18	-
		Deep litter, divided	-	18	-	15
		Deep litter	-	-	-	15

c) **Poultry**

			Solid manure Total N	Deep litter Total N
	Housing type	Floor or manure type	pct. loss of N ex animal	
Hens and pullets	Free-range, organic and barn	Deep pit	40	25
		Deep litter	-	28
		Manure belt	10	25
	Battery	Deep pit	12	-
		Manure belt	10	-
Broilers	Conventional	Deep litter	-	20
	Organic and barn	Deep litter	-	25
Turkeys, ducks and geese		Deep litter	-	20

d) **Other**

	Urine TAN	Slurry TAN	Solid manure Total N	Deep litter Total N
	Pct. loss of TAN ex animal		pct. loss of N ex animal	
Fur animals	35	47	35	-
Horses, sheep and goats	-	-	-	15

Table 3E-4 NH₃ emission factors for storage units, 2012.

			Urine	Slurry ¹	Solid manure	Deep litter	Pct. of solid manure stored in heap on field
Cattle		Total N	2	2.1	4	1	35
		TAN	2.2	3.5	-	-	-
Pigs	Sows	Total N	2	2.4	19	6.5	50
		TAN	2.2	2.9	-	-	-
	Weaners	Total N	2	2.4	19	9.8	-
		TAN	2.2	2.9	-	-	-
	Fattening pigs	Total N	2	2.4	19	9.8	75
		TAN	2.2	2.9	-	-	-
Poultry	Hens and pullets	Total N	-	2	7.5	4.8	95
	Broilers	Total N	-	-	11.5	6.8	85
	Turkeys, ducks, and geese	Total N	-	-	-	6.8, 8(Turkeys)	-
Fur animals		Total N	0	3.1	11.5	-	-
		TAN	0	3.1	-	-	-
Sheep and goats		Total N	-	-	-	4	-
Horses		Total N	-	-	-	4	-

Table 3E-5 Parameters for winter feeding plans.

		Feeding code*	% dm*	% Crude protein*	% Raw fat*	% Raw ashes*	% Carbo- hydrates	FU/kg dm*	kg dm/day**	MJ/day	GE _{FU}
		PDIR (2002)									
Heifers:	Straw	781	85.0	4.0	1.9	4.5	89.6	0.2	33.4	571.8	
	Maize silage	593	31.0	8.7	2.2	4.2	84.9	0.9	57.5	1 009.0	
	Toasted soya	155	87.5	49.1	3.2	7.4	40.3	1.4	8.1	161.7	
	Total	-	-	-	-	-	-	-	99.0	1 742.4	25.8
Suckling cows:	Straw	781	85.0	4.0	1.9	4.5	89.6	0.2	1.6	119.1	
Period 1 (2 mth)	Toasted soya	155	87.5	49.1	3.2	7.4	40.3	1.4	3.4	49.6	
	Barley	201	85.0	11.2	2.9	2.2	83.7	1.1	1.8	29.2	
Period 2 (4 mth)	Straw	781	85.0	4.0	1.9	4.5	89.6	0.2	3.2	238.2	
	Toasted soya	155	87.5	49.1	3.2	7.4	40.3	1.4	3.0	29.1	
	Barley	202	85.0	11.2	2.9	2.2	83.7	1.1	3.2	52.0	
	Total	-	-	-	-	-	-	-	15.2	517.1	34.0
Horses:	Straw	781	85.0	4.0	1.9	4.5	89.6	0.2	4.0	58.2	
	Hay	665	85.0	12.1	2.6	7.7	77.6	0.6	3.0	44.0	
	Oat	202	86.0	12.1	5.7	2.7	79.5	0.9	2.5	40.1	
	Supplemental		86.4	15.4	4.3	6.6	73.7	1.0	1.0	15.5	
	Total	-	-	-	-	-	-	-	-	157.7	29.8
Sheep and Goats:	Straw	781	85.0	4.0	1.9	4.5	89.6	0.2	1.0	14.6	
	Toasted soya	155	87.5	49.1	3.2	7.4	40.3	1.4	0.1	1.8	
	Barley	202	85.0	11.2	2.9	2.2	83.7	1.1	0.4	6.2	
	Grass pills (dried)	707	92.0	17.0	3.1	11.0	68.9	0.6	1.0	15.7	
	Total	-	-	-	-	-	-	-	-	38.2	30.0
Summer grazing											
Grazing	Clover grass, 2 weeks old	422	18.0	22.0	4.1	9.4	64.5	1.0	1.0	18.8	
	Total	-	-	-	-	-	-	-	1.0	18.8	18.8
Swine:	Full feeding										
	Sows	-	87.1	16.1	5.2	5.5	73.2	1.2	-	64.2	17.5
	Weaners	-	87.4	18.8	5.7	5.5	70.0	1.3	-	2.1	16.5
	Fattening pigs	-	86.9	17.0	4.7	5.1	73.3	1.2	-	9.6	17.3

Table 3E-6 Energy factors used for GE.

	MJ per kg dm
E _{Crude protein}	24.237
E _{Raw fat}	34.116
E _{Carbonhydrates}	17.3

Table 3E-7 Feed intake 1990-2012, FU per animal per year.

	1990	1995	2000	2005	2008	2009	2010	2011	2012
Dairy cattle	5 549	5 896	5 941	6 557	6 687	6 845	6 878	6 804	6 855
<u>Non-dairy cattle:</u>									
Calves, bull	1 190	1 200	1 205	1 228	1 230	1 230	1 230	1 230	1 230
Calves, heifer	1 734	1 743	1 728	1 820	2 043	2 040	2 041	2 040	9 492
Bulls > ½ year	1 995	2 082	1 846	2 275	2 382	2 178	2 227	2 438	2 246
Heifer > ½ year	1 721	1 735	1 737	2 357	2 609	2 606	2 605	2 604	2 604
Suckling cattle	2 515	2 515	2 515	2 378	2 417	2 417	2 417	2 417	2 417
Sheep (mother sheep incl. lambs)	728	728	728	728	728	728	728	728	728
Goats (mother goats incl. kids)	669	669	669	655	664	664	664	665	665
Horses	1 995	1 995	1 995	1 996	1 995	1 995	1 996	1 995	1 995
<u>Swine:</u>									
Sows (incl. pigs < 7.4 kg)	1 300	1 300	1 340	1 450	1 484	1 500	1 520	1 535	1 540
Weaners (7.4 – 32 kg)	128	167	209	206	244	229	226	240	247
Fattening pigs (32 – 107 kg)	1 015	950	826	790	813	816	776	805	800
<u>Other:</u>									
Deer	668	668	668	668	668	668	668	668	668

Table 3E-8 Grazing animals 1990 – 2012, number of days on grass per year.

Livestock category	1990	1995	2000	2005	2008	2009	2010	2011	2012
Dairy cattle	55	55	55	32	18	18	18	18	18
Heifer > ½ year ^a	165	196	196	156	132	132	132	132	132
Suckling cattle	184	224	224	224	224	224	224	224	224
Sheep and goats	265	265	265	265	265	265	265	265	265
Horses	183	183	183	183	183	183	183	183	183

^a feeding days on grass

Table 3E-9a Average gross energy intake (GE) 1990 – 2012, MJ per head per day.

Livestock category	1990	1995	2000	2005	2008	2009	2010	2011	2012
Dairy cattle	278.2	295.6	297.9	328.7	335.3	343.2	344.8	341.1	343.7
Non-dairy cattle (heifer)	107.2	105.2	105.2	115.6	130.5	130.3	130.3	130.2	130.2
Sheep (mother sheep incl. lambs)	43.6	43.6	43.6	43.6	43.6	43.6	43.6	43.6	43.6
Goats (mother goats incl. kids)	40.1	40.1	40.1	39.2	39.8	39.8	39.8	39.9	39.9
Horses	133.0	133.0	133.0	133.0	133.0	133.0	133.0	133.0	133.0
Swine (fattening pigs)	43.3	38.9	38.1	38.9	39.9	40.4	40.3	40.4	40.5
Poultry	NA	NA	NA	NA	NA	NA	NA	NA	NA
<u>Other:</u>									
Deer	34.5	34.5	34.5	34.5	34.5	34.5	34.5	34.5	34.5
Fur farming	NA	NA	NA	NA	NA	NA	NA	NA	NA
Ostrich	NA	NA	NA	NA	NA	NA	NA	NA	NA
Pheasant	NA	NA	NA	NA	NA	NA	NA	NA	NA

Table 3E-9b Average gross energy intake (GE) 1990 – 2012, MJ per head per day – Subcategories for cattle and swine.

Subcategories for cattle and swine	1990	1995	2000	2005	2008	2009	2010	2011	2012
<u>Cattle</u>									
Dairy, large breed	285.8	302.4	304.1	335.9	341.8	350.2	351.7	348.2	350.9
Dairy, Jersey	237.2	250.7	253.2	278.8	290.6	296.3	299.1	296.2	298.6
Calves, bull	59.6	60.2	60.4	61.6	61.7	61.7	61.7	61.7	61.7
Calves, heifer	86.5	87.2	86.5	91.3	102.4	102.3	102.3	102.3	102.2
Bulls > ½ year	113.6	114.3	114.7	115.8	116.0	116.2	116.3	116.2	116.2
Heifer > ½ year	107.2	105.2	105.2	115.6	130.5	130.3	130.3	130.2	130.2
Suckling cattle	181.6	170.2	170.2	160.9	163.6	163.6	163.6	163.6	163.6
<u>Swine</u>									
Sows (incl. pigs < 7.4 kg)	62.3	62.3	64.2	69.5	71.1	71.9	72.8	73.6	73.8
Weaners (7.4 – 32 kg)	11.1	13.2	13.8	13.8	15.2	14.3	14.1	14.2	14.2
Fattening pigs (32 – 107 kg)	43.3	38.9	38.1	38.9	39.9	40.4	40.3	40.4	40.5

Table 3E-10a VS daily excretion (average) 1990 – 2012, kg dm per head per day – CRF categories.

365 housing days	1990	1995	2000	2005	2008	2009	2010	2011	2012
<u>Livestock category</u>									
Dairy cattle	5.54	5.70	6.03	6.59	6.07	6.21	6.23	6.09	6.22
Non-dairy cattle (weighted average)	1.68	1.84	1.95	2.69	2.75	2.76	2.76	2.73	2.70
Sheep (mother sheep incl. lambs)	1.12	1.12	1.12	1.12	1.12	1.12	1.12	1.12	1.05
Goats (mother goats incl. kids)	1.06	1.06	1.06	1.06	1.07	1.07	1.07	1.07	1.01
Horses	3.67	3.67	3.67	3.65	3.65	3.65	3.65	3.65	3.65
Swine (weighted average)	0.23	0.22	0.22	0.22	0.21	0.21	0.21	0.20	0.20
Poultry (weighted average)	0.003	0.003	0.003	0.003	0.003	0.003	0.003	0.003	0.003
Fur farming	0.09	0.09	0.08	0.08	0.08	0.08	0.08	0.09	0.09
Deer	0.72	0.72	0.72	0.72	0.72	0.72	0.72	0.72	0.72

Table 3E-10b VS daily excretion (average) 1990 – 2012, kg dm per head per day – Subcategories.

365 housing days	1990	1995	2000	2005	2008	2009	2010	2011	2012
<u>Cattle:</u>									
Dairy cattle	5.54	5.70	6.03	6.59	6.07	6.21	6.23	6.09	6.22
Calves, bull	1.49	1.50	1.51	1.53	1.53	1.53	1.53	1.53	1.53
Bulls > ½ year	1.77	2.29	2.74	3.40	3.89	4.03	4.04	3.92	3.89
Calves, heifer	1.30	1.30	1.31	1.69	1.80	1.80	1.80	1.80	1.80
Heifer > ½ year	1.69	1.90	1.99	2.72	2.82	2.80	2.80	2.80	2.79
Suckling cattle	6.72	5.67	5.64	5.46	4.19	4.22	4.23	4.24	4.28
<u>Swine:</u>									
Sows (incl. pigs < 7.4 kg)	0.53	0.55	0.62	0.62	0.46	0.46	0.45	0.45	0.45
Weaners (7.4 – 32 kg)	0.11	0.10	0.10	0.11	0.10	0.10	0.10	0.10	0.10
Fattening pigs (32 – 107 kg)	0.34	0.33	0.32	0.34	0.33	0.33	0.33	0.33	0.33
<u>Poultry:</u>									
Hens	0.02	0.02	0.02	0.02	0.02	0.03	0.03	0.03	0.03
Pullets	0.004	0.004	0.004	0.004	0.004	0.004	0.004	0.004	0.003
Broilers	0.002	0.002	0.002	0.002	0.002	0.002	0.002	0.002	0.002
Turkeys, geese and ducks	0.01	0.01	0.02	0.02	0.02	0.02	0.02	0.02	0.02

Table 3E-11 Basic data from Sommer et al. (2001) used to estimation of lower emission of N₂O and CH₄ from biogas treated slurry.

CH ₄ (housing + storage).					
	Reference situation	Biogas scenario I	Difference	Reduction potential	Lower emission compared with the reference situation
	kg CO ₂ eqv/kg VS excreted			Pct.	Pct.
Cattle	1.1458	0.8778	0.2680	23	77
Pigs	1.1481	0.6864	0.4617	40	60
N ₂ O (field)					
	Reference situation	Biogas scenario I	Difference	Reduction potential	Lower emission compared with the reference situation
	kg CO ₂ eqv/kg VS excreted			Pct.	Pct.
Cattle	0.1624	0.1032	0.0592	36	64
Pigs	0.2127	0.1258	0.0869	41	59

Reference: Sommer et al., 2001 (see appendix 3 page 50).

Calculation example of CH₄ emission form biogas treated slurry:

Reference situation (whiteout biogas treatment): $0.7236 + 0.4222 = 1.1458$ kg CO₂ eqv/kg VS ex

Biogas scenario I (biogas treatment): $0.7236 + 0.0770 + 0.0772 = 0.8778$ kg CO₂ eqv/kg VS ex

Reduction potential: $(1.1458 - 0.8778) = 0.2680 \rightarrow 0.2680 / (1.1458 * 100\%) = 23\%$

This leads to an emission from treated cattle slurry, which is 77 % compared with untreated slurry.

Table 3E-12a Calculation of lower CH₄ emission as a consequence of biogas treated slurry.

Year	Biogas treated slurry	Cattle slurry	Pig slurry	VS cattle slurry	VS pig slurry	CH ₄ emission, untreated cattle slurry	CH ₄ emission, treated cattle slurry	Lower CH ₄ emission, cattle slurry	CH ₄ emission, untreated pig slurry	CH ₄ emis- sion, treated pig slurry	Lower CH ₄ emission, pig slurry	Lower CH ₄ emission, total
	1000 Gg			1000 Gg		Gg			Gg			Gg
1990	0.19	0.09	0.10	0.007	0.005	0.113	0.087	0.026	0.154	0.092	0.062	0.088
1991	0.32	0.14	0.18	0.012	0.009	0.191	0.146	0.045	0.259	0.155	0.104	0.149
1992	0.39	0.18	0.21	0.014	0.010	0.233	0.178	0.054	0.316	0.189	0.127	0.181
1993	0.46	0.21	0.25	0.017	0.012	0.274	0.210	0.064	0.372	0.223	0.150	0.214
1994	0.54	0.24	0.30	0.020	0.014	0.322	0.247	0.075	0.437	0.261	0.176	0.251
1995	0.64	0.29	0.35	0.024	0.017	0.382	0.292	0.089	0.518	0.310	0.208	0.298
1996	0.69	0.31	0.38	0.026	0.019	0.411	0.315	0.096	0.558	0.334	0.225	0.321
1997	0.83	0.37	0.46	0.031	0.022	0.495	0.379	0.116	0.672	0.402	0.270	0.386
1998	1.01	0.45	0.56	0.037	0.027	0.602	0.461	0.141	0.817	0.489	0.329	0.470
1999	1.04	0.47	0.57	0.039	0.028	0.620	0.475	0.145	0.842	0.503	0.338	0.483
2000	1.16	0.52	0.64	0.043	0.031	0.692	0.530	0.162	0.939	0.561	0.377	0.539
2001	1.26	0.57	0.69	0.047	0.034	0.751	0.576	0.176	1.020	0.610	0.410	0.586
2002	1.44	0.65	0.79	0.053	0.039	0.859	0.658	0.201	1.165	0.697	0.469	0.669
2003	1.76	0.79	0.97	0.065	0.047	1.049	0.804	0.245	1.424	0.851	0.573	0.818
2004	1.88	0.85	1.03	0.070	0.050	1.121	0.859	0.262	1.521	0.910	0.612	0.874
2005	1.93	0.87	1.06	0.072	0.052	1.151	0.882	0.269	1.562	0.934	0.628	0.897
2006	2.14	0.96	1.18	0.079	0.057	1.276	0.978	0.298	1.732	1.035	0.696	0.995
2007	2.15	0.97	1.18	0.080	0.058	1.282	0.982	0.300	1.740	1.040	0.700	1.000
2008	2.19	0.99	1.20	0.081	0.059	1.306	1.000	0.305	1.772	1.060	0.713	1.018
2009	2.39	1.08	1.31	0.089	0.064	1.425	1.092	0.333	1.934	1.156	0.778	1.111
2010	2.49	1.12	1.37	0.092	0.067	1.485	1.137	0.347	2.015	1.205	0.810	1.158
2011	2.39	1.08	1.31	0.089	0.064	1.425	1.092	0.333	1.934	1.156	0.778	1.111
2012	2.45	1.10	1.35	0.081	0.066	1.461	1.119	0.342	1.983	1.185	0.797	1.139

Table 3E-12b Calculation of lower N₂O emission as a consequence of biogas treated slurry.

Year	Biogas treated slurry	Cattle slurry	Pig slurry	N ₂ O emission, untreated cattle slurry	N ₂ O emission treated cattle slurry	Lower N ₂ O emission, cattle slurry	N ₂ O emission untreated pig slurry	N ₂ O emission treated pig slurry	Lower N ₂ O emission, pig slurry	Lower N ₂ O emission, total
	1000 Gg				Gg			Gg		Gg
1990	0.19	0.09	0.10	0.006	0.004	0.002	0.007	0.004	0.003	0.005
1991	0.32	0.14	0.18	0.010	0.006	0.004	0.012	0.007	0.005	0.008
1992	0.39	0.18	0.21	0.012	0.008	0.004	0.015	0.009	0.006	0.010
1993	0.46	0.21	0.25	0.014	0.009	0.005	0.017	0.010	0.007	0.012
1994	0.54	0.24	0.30	0.016	0.010	0.006	0.020	0.012	0.008	0.014
1995	0.64	0.29	0.35	0.019	0.012	0.007	0.024	0.014	0.010	0.017
1996	0.69	0.31	0.38	0.021	0.013	0.008	0.026	0.015	0.010	0.018
1997	0.83	0.37	0.46	0.025	0.016	0.009	0.031	0.018	0.013	0.022
1998	1.01	0.45	0.56	0.031	0.019	0.011	0.038	0.022	0.015	0.026
1999	1.04	0.47	0.57	0.031	0.020	0.011	0.039	0.023	0.016	0.027
2000	1.16	0.52	0.64	0.035	0.022	0.013	0.043	0.026	0.018	0.030
2001	1.26	0.57	0.69	0.038	0.024	0.014	0.047	0.028	0.019	0.033
2002	1.44	0.65	0.79	0.044	0.028	0.016	0.054	0.032	0.022	0.038
2003	1.76	0.79	0.97	0.053	0.034	0.019	0.065	0.039	0.027	0.046
2004	1.88	0.85	1.03	0.057	0.036	0.021	0.070	0.041	0.029	0.049
2005	1.93	0.87	1.06	0.058	0.037	0.021	0.072	0.042	0.029	0.051
2006	2.14	0.96	1.18	0.065	0.041	0.024	0.080	0.047	0.033	0.056
2007	2.15	0.97	1.18	0.065	0.041	0.024	0.080	0.047	0.033	0.056
2008	2.19	0.99	1.20	0.066	0.042	0.024	0.081	0.048	0.033	0.057
2009	2.39	1.08	1.31	0.072	0.046	0.026	0.089	0.053	0.036	0.063
2010	2.49	1.12	1.37	0.075	0.048	0.027	0.093	0.055	0.038	0.065
2011	2.39	1.08	1.31	0.072	0.046	0.026	0.089	0.053	0.036	0.063
2012	2.45	1.10	1.35	0.074	0.047	0.027	0.091	0.054	0.037	0.064

Table 3E-13 Background data for calculation of N content in nitrogen fixing crops.

Table SE 15 Background data for calculation of N content in nitrogen fixing crops.						
Crop	Dry matter content ¹	N-content in dm ¹	Straw yield in pct. of grain yield ²	Share, root + stubble ³	Share of N in crop which is fixed ³	N-fixed
	pct.	pct.	pct.	pct.	pct.	kg N/tonnes harvested
Based on yield						
Field peas, grain	85	3.97		25	75	
Field peas, straw	87	1.15	60			
Legumes grown to maturity, in total						37.3
Lucerne	21	3.04		60	75	7.7
Crops for silage	23	2.64		25	80	6.1
Legumes, marrow-stem kale and green fodder	23	2.64		25	80	6.1
Grass and clover fields as well as fields sown with an under crop	13	4.00		75	90	8.2
Peas for conservation ⁴	23	2.64		25	80	6.1
Fields with aftermath	13	4.00		75	90	8.2
Based on area						
	kg N/ha/year					
Seed of leguminous grass crops:						
Red clover	200					
White clover	180					
Black medick	180					

¹ Feedstuff table (DAAC, 2000).² Kyllingsbæk (2000).³ Kristensen (2002) and Kyllingsbæk (2000).⁴ Assumed that peas constitute 80% of the total area.

Table 3E-14 Estimated share of nitrogen fixing plants in crops.

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999-2012
	pct.									
Cereals for silage										
Share of peas (whole-crop)	30	30	35	35	40	40	45	45	50	50
Share of peas in whole-crop	40	40	40	40	40	40	40	40	40	40
Legumes, marrow-stem kale and other green fodder										
Share with legumes:	60	60	60	60	60	60	60	60	60	60
of which share with peas	40	40	40	40	40	40	40	40	40	40
Peas for conservation	80	80	80	80	80	80	80	80	80	80
Grass in rotation										
Share of clover grass fields	74	76	78	80	82	84	85	86	87	88
Clover percentage in the clover grass fields	20	20	20	20	20	22	24	26	28	30
Grass not in a rotation										
Clover percentage	5	5	5	5	5	5	5	5	5	5
Fields with aftermath										
Share with clover grass	74	76	78	80	82	84	85	86	87	88
Clover percentage	30	30	30	30	30	30	30	30	30	30

Table 3E-15 Area of N-fixing crops 1990-2012, ha.

	1990	1995	2000	2005	2008	2009	2010	2011	2012
Legumes to maturity	114 354	74 178	35 590	15 819	4 910	6 332	10 349	7 109	6 252
Lucerne	8 494	10 099	5 245	4 575	3 756	5 366	6 405	6 926	4 715
Crops for silage	47 772	87 893	118 763	75 512	52 251	55 851	62 845	56 672	54 333
Legumes/marrow-stem kale	2 584	2 964	585	NO	NO	NO	NO	NO	NO
Grass and clover in rotation	248 815	238 384	246 656	253 007	300 251	305 889	320 914	329 135	326 797
Grass not in rotation	217 235	207 122	166 261	192 968	189 962	192 433	199 859	186 652	200 413
Fields with catch crop	232 000	236 000	309 100	121 800	113 900	115 200	116 600	116 700	84 100
Peas for conservation	8 791	5 529	4 149	2 999	3 592	3 737	2 677	2 935	2 837
Seeds of leguminous grass crops	2 334	3 835	4 603	5 258	4 457	4 542	4 483	3 742	4 536
Total area of N-fixing crops	882 379	866 004	890 952	671 938	673 079	689 350	724 132	709 871	683 983

NO = Not occurring.

Table 3E-16 Background data for estimation of N₂O emission from crop residue 2012.

Crop type	Stubble	Husks	Top	Leafs	Nitrogen content in crop residue	
	kg N per ha				kg N per ha per year	Gg N per year
Winter wheat	6.3	10.7	-	-	17.0	10.01
Spring wheat	6.3	7.4	-	-	13.7	0.42
Winter rye	6.3	10.7	-	-	17.0	0.98
Triticale	6.3	10.7	-	-	17.0	0.67
Winter barley	6.3	5.9	-	-	12.2	1.27
Spring barley	6.3	4.1	-	-	10.4	6.48
Oats	6.3	4.1	-	-	10.4	0.53
Winter rape	4.4	-	-	-	4.4	0.55
Spring rape	4.4	-	-	-	4.4	0.01
Potatoes (top), non-harvest	-	-	48.7	-	48.7	2.03
Beet (top), non-harvest	-	-	56.7 ^a	-	56.7	2.69
Straw, non-harvest	-	-	-	-	6.4 ^a	9.57
Pulse	11.3	-	-	-	11.3	0.07
Lucerne	32.3	-	-	-	10.8	0.05
Maize – for green fodder	6.3	-	-	-	6.3	1.16
Cereal – for green fodder	6.3	-	-	-	6.3	0.34
Peas for conservation	11.3	-	-	-	11.3	0.03
Vegetables	11.3	-	-	-	11.3	0.08
Grass field legumes	11.3	-	-	-	5.7	0.03
Grass- and clover field in rotation	32.3		-	10.0	26.2	8.55
Grass- and clover field out of rotation	38.8		-	20.0	20.0	4.01
Catch crop	6.3	-	-	-	6.3	0.53
Seeds of grass crops	6.3	10.7	-	-	13.9	0.95
Set-a-side	38.8	-	-	15.0	18.9	0.09
Total N from crop residue						51.10

^a express the yield for 2012 - varies from year to year. Based on yield data from Statistics Denmark and N-content from the feeding plan.

Reference: Djurhuus and Hansen (2003).

Table 3E-17 Area of agricultural land, 1990 – 2012, ha.

	1990	1995	2000	2005	2008	2009	2010	2011	2012
Garden centre, fruit & berries	11 687	12 135	11 050	10 682	11 106	9 661	9 410	8 708	8 841
Agriculture crops excl. grass in rotation	2 294 434	2 039 330	2 021 092	2 065 948	2 084 866	2 082 006	2 076 103	2 082 609	2 072 907
Vegetables grown in the open	16 105	12 584	10 628	9 431	11 048	11 463	10 720	11 144	10 219
Permanent grass	217 235	207 122	166 261	192 968	189 962	192 433	199 859	186 652	200 413
Fallow	NO	216 493	191 295	175 200	70 662	5 699	9 874	4 367	5 018
Grass in rotation	248 815	238 384	246 656	253 007	300 251	305 889	320 914	329 135	326 797
Sum	2 788 276	2 726 048	2 646 982	2 707 236	2 667 895	2 607 151	2 626 880	2 622 615	2 624 195

NO = Not occurring.

Table 3E-18 Harvested straw, 1990 – 2012, 1 000 tonnes.


	1990	1995	2000	2005	2008	2009	2010	2011	2012
<u>Straw yield, total</u>	<u>7 032</u>	<u>6 118</u>	<u>6 100</u>	<u>5 524</u>	<u>5 662</u>	<u>6 280</u>	<u>5 469</u>	<u>5 436</u>	<u>5 804</u>
Straw, left on the field	3 354	2 309	2 400	2 271	1 794	2 230	2 161	2 162	2 133
Straw, harvested	3 678	3 809	3 700	3 253	3 868	4 050	3 308	3 274	3 670
- <i>used for fodder</i>	1 063	1 829	1 655	1 038	1 284	1 219	1 074	1 103	1 148
- <i>used for fuel</i>	900	1 089	1 096	1 450	1 795	2 018	1 605	1 507	1 836
- <i>used for bedding</i>	1 715	891	949	765	789	813	628	666	686

Nitrogen leaching and Run-off

Calculations of nitrogen lost by leaching from groundwater are based on two models described in Børgesen and Grant (2003) (in Danish). The model SKEP/DAISY is a dynamic model, N-LES is an empirical model and SKEP is an up scaling model. The SKEP/DAISY calculations were done for 10 scenarios (the years 1984, 1989 and 1995-2002) and the N-LES calculations were done for an 11 year period (1990-2000). Both calculations were up scaled nationwide. The key parameters for the models were land use, nitrogen from synthetic fertilizer and manure, application practice for manure and NH₃ evaporation at application of manure (SKEP/DAISY only). The calculations were normalised to an average climate. A schematic overview of the models is seen below.

Figure 3E-1 Model calculation of nitrogen leaching from groundwater nationwide by SKEP/DAISY and N-LES.

Basic DAISY calculations of N-leaching



Each crop rotation calculates for:

6 climate regions

30 fertilizer plan

4 soil type (here 2 w/w.out water)

38.000 combinations

Data base

Calculation for all combinations for each of 4 climate year


Calculation for 12 combinations for each year in a 11 years period (1989-2001).

N-LES calculations

Model calculations for the crop rotations and fertilizer planes in SKEP plus appurtenant percolations from the DAISY calculations. Model calculations for each of the 11 years in the period 1989-2001, mean of the 11 years is up scaled nationwide by SKEP

Up-scaling by the SKEP model

In the up scaling of DAISY calculations a climate normalisation and yield correction is made



3E-Table 19 QA/QC procedure, stage I – III.

Stage I: Check of input data	Variable	Reference
Livestock production	- number of animal	DSt
	- slaughter data	
Normative figures	- N-excretion	DCA
	- use of straw	
	- amount of manure	
	- feed intake	
	- milk yield	
Housing types	- distribution	DAAS + DAFA
Grazing days		DAAS
Crops	- land use	DSt
	- crop yield	
	- crop production	
Synthetic fertiliser	- N-content	DAFA
	- fertiliser types	
N-leaching	- amount of nitrogen leached	DCE
Atmospheric deposition	- all NH ₃ emission sources	DCE – NH ₃ inventory
Sewage sludge and industrial waste	- Amount of sludge applied to soils	EPA + DAFA
Stage II: Check of IDA data – overall	Emission source	Variable
Recalculation	- CO ₂ eqv. total emission	- compared with latest submission
	- CH ₄ , N ₂ O, NMVOC	
	- emission from field burning	
Time series	- CO ₂ eqv. total emission	- trends
	- CH ₄ , N ₂ O, NMVOC	- jumps and dips
	- emission from field burning	
Stage III: Check of IDA data – specific	Emission source	Variable
CH ₄	- enteric fermentation	- IEF (jumps and dips)
		- Ym (dairy cattle + heifer)
		- GE
CH ₄	- manure management	- IEF (jumps and dips)
		- VS
		- biogas
N ₂ O	- manure management	- trends (jumps and dips)
		- IEF
		- biogas
N ₂ O	- synthetic fertiliser	- trends (jumps and dips)
		- IEF
N ₂ O	- animal waste applied to soil	- trends (jumps and dips)
		- IEF
N ₂ O	- N-fixing crops	- trends (jumps and dips)
		- IEF
N ₂ O	- crop residue	- trends (jumps and dips)
		- IEF
N ₂ O	- pasture, range and paddock	- trends (jumps and dips)
		- IEF
N ₂ O	- atmospheric deposition	- trends (jumps and dips)
		- IEF
N ₂ O	- N-leaching and run-off	- trends (jumps and dips)
		- IEF
N ₂ O	- sewage sludge + industrial waste	- trends (jumps and dips)
		- IEF
NMVOC	- crops	- trends (jumps and dips)

Annex 3F - LULUCF

List of tables

Table 3F.1 Estimation of forest percentage and forest area.

Table 3F.2 Estimation of forest area with a specific characteristic.

Table 3F.3 Estimation of diameter-height equations.

Table 3F.4 Estimation of quadratic mean diameter.

Table 3F.5 Estimation of biomass and carbon of trees.

Table 3F.6 Estimation of total biomass and carbon pools.

Table 3F.7 Estimation of biomass and carbon with a given characteristic.

Table 3F.8 Estimation of biomass and carbon content of dead wood.

Table 3F.9 Estimation of total biomass and carbon pools of dead wood.

Table 3F.10 Estimation of forest floor carbon.

Table 3F.11 Hectares grown in the different areas of Denmark.

Table 3F.12 Crop yield from Statistics Denmark in 2010 distributed regions, Hhg crop ha⁻¹

Table 3F.13 Area input format for Eastern Jutland to C-TOOL in 2010. FK represent the soil type (Color Code (Farve Kode))

Table 3F.14 Average annual temperatures for Denmark, °C

Table 3F.1 Estimation of forest percentage and forest area.

Equation	Description
$X_j = \frac{A_j}{A_{15,j}}$	The forest percentage (X) of the j th sample plot (SSU) is estimated as the forested area (A) divided by the total area of the 15 m radius sample plot ($A_{15,j}$).
$\bar{X}_Z = \frac{1}{n_Z} \sum_Z X_j R_j$	Average forest percentage (\bar{X}) of all inventoried plots (SSU) with forest status Z based on aerial photos. R_j is an indicator variable that is 1 for inventoried plots and 0 otherwise. n_Z is the number of inventoried plots identified as forest or OWL from the air photos.
$\bar{\bar{X}} = \frac{1}{n} \left(\sum_{j=1}^n X_j R_j + N_{21} \bar{X}_1 + N_{22} \bar{X}_2 \right)$	Overall average forest percentage ($\bar{\bar{X}}$). n is the total number of inventoried and non-inventoried sample plots. N_{21} and N_{22} is the number of non-inventoried sample plots with forest and OWL, respectively.
$A_{Forest} = \bar{\bar{X}} \cdot A_{Total}$	Total forest area. A_{Total} is the total land area, $\bar{\bar{X}}$ is the estimated forest percentage and A_{Forest} is the total forest area.

Table 3F.2 Estimation of forest area with a specific characteristic.

Equation	Description
$X_k = \frac{\sum_{j=1}^n R_{jk} A_j}{\sum_{j=1}^n A_j}$	Proportion of the forest area with a given characteristic (X_k). R_{jk} is an indicator variable which is 1 if the forest area on the j th sample plots has the k th characteristic and 0 otherwise. A_j is the sample plot area and n is the total number of inventoried sample plots with forest cover.
$A_k = X_k \cdot A_{Forest}$	Total area with a given characteristic (A_k). X_k is the estimated proportion of the forest area with the k th characteristic and A_{Forest} is the total forest area.

Table 3F.3 Estimation of diameter-height equations.

Equation	Description
$h_{ij} = 13 + (\bar{h}_j - 13) \cdot \exp \left(\alpha_1 \cdot \left(1 - \frac{\bar{d}_j}{d_{ij}} \right) + \alpha_2 \cdot \left(\frac{1}{\bar{d}_j} - \frac{1}{d_{ij}} \right) \right)$	Site specific dh-regression for calculating height of trees not measured for height. h_{ij} and d_{ij} is the height and diameter of the i 'th tree on the j 'th sample plot. \bar{h}_j and \bar{d}_j are the average height and diameter of trees measured for height on the j th sample plot. α_1 and α_2 are species and growth-region specific parameters
$h_{ij} = 13 + \beta_1 \cdot \exp \left(-\frac{\beta_2}{d_{ij}} \right)$	General dh-regression for calculating height of trees not measured for height. h_{ij} and d_{ij} is the height and diameter of the i 'th tree on the j 'th sample plot. β_1 and β_2 are species and growth-region specific parameters

Table 3F.4 Estimation of quadratic mean diameter.

Equation	Description
$g_{ij} = \frac{\pi}{4} d_{ij}^2$	Basal area (g) of the i th tree on the j th plot is calculated from the diameter at breast height (d) (1.3 m above ground) assuming a circular stem form.
$G_j = \sum_{i=1}^m \frac{1}{A_{c,ij}} g_{ij}$	Basal area per hectare (G) the j th sample plot is calculated as the scaled sum of individual tree basal areas. Basal area (g) of the i th tree on the j th sample plot is scaled according to the plot area ($A_{c,ij}$) of the c th concentric circle ($c=3,5; 10; 15$ m).
$N_j = \sum_{i=1}^m \frac{1}{A_{c,ij}}$	Stem number per hectare (N) the j th sample plot is calculated as the scaled number of individual trees. The i th tree on the j th sample plot is scaled according to the plot area ($A_{c,ij}$) of the c th concentric circle ($c=3,5; 10; 15$ m).
$D_{g,j} = \sqrt{\frac{4}{\pi} \frac{G_j}{N_j}}$	The mean squared diameter is calculated from the calculated basal area and stem number for each plot.

Table 3F.5 Estimation of biomass and carbon of trees.

Equation	Description
$v_{ij} = F(d_{ij}, h_{ij}, D_{g,j})$	The volume (v) of the i th tree on the j th sample plots is calculated using the existing volume functions (F) using the tree diameter and height and the quadratic mean diameter.
$B_{ij} = V_{ij} \cdot Density_{ij}$	Biomass (B) of the i th tree on the j th sample plot is estimated as the total volume (V_{tot}) times the species specific density.
$E_{ij} = F(d_{ij}, h_{ij})$	Expansion factor model for beech and Norway spruce
$v_{tot,ij} = B_{ij} \cdot E_{ij}$	The total above and below ground volume (v_{tot}) of the i th tree on the j th sample plot. B_{ij} is the calculated above-ground biomass of the tree and E is the expansion factor.
$C_{ij} = B_{ij} \cdot 0.5$	Carbon of the i th tree on the j th sample plot is calculated as the biomass (B) times 0.5.

Table 3F.6 Estimation of total biomass and carbon pools.

Equation	Description
$V_{cj} = \frac{1}{A_{cj}} \sum_{i=1}^m R_{c,i} v_{ij}$	Volume, biomass or carbon per hectare (V) of the c th concentric circle on the j th sample plot ($c=3,5; 10; 15$ m). R_c is an indicator variable that is 1 if the i th tree is measured on the c th circle and 0 otherwise. $A_{c,ij}$ is the area of the j th sample plot and c th concentric circle; m is the number of trees on the j th sample plot.
$\bar{V}_c = \frac{\sum_{j=1}^n A_{cj} V_{cj}}{\sum_{j=1}^n A_{cj}}$	The average area weighted volume, biomass or carbon per hectare (\bar{V}) of the c th concentric circle. $A_{c,ij}$ is the area of the j th sample plot and c th concentric circle; n is the number of sample plots.
$\bar{\bar{V}} = \bar{V}_{3,5} + \bar{V}_{10} + \bar{V}_{15}$	The overall average volume, biomass or carbon per hectare ($\bar{\bar{V}}$) is estimated as the sum of the average volume, biomass or carbon per hectare (\bar{V}_c) for the three concentric circles ($c=3.5, 10$ and 15)
$V = \bar{\bar{V}} \cdot A_{Skov}$	Total volume, biomass or carbon V is the overall average volume, biomass or carbon per hectare ($\bar{\bar{V}}$) times the forest area A_{Forest} .

Table 3F.7 Estimation of biomass and carbon with a given characteristic.

Equation	Description
$V_{cj,k} = \frac{1}{A_{cj}} \sum_{i=1}^m R_{c,ij} R_{k,ij} v_{ij}$	Volume, biomass or carbon per hectare (V) with the k th characteristic of the c th concentric circle on the j th sample plot ($c=3,5; 10; 15$ m). R_c is an indicator variable that is 1 if the i th tree is measured on the c th circle and 0 otherwise. R_k is an indicator variable that is 1 if the tree has k th characteristic and 0 otherwise. $A_{c,ij}$ is the area of the j th sample plot and c th concentric circle; m is the number of trees on the j th sample plot.
$\bar{V}_{c,k} = \frac{\sum_{j=1}^n A_{cj} V_{cj,k}}{\sum_{j=1}^n A_{cj}}$	The average area weighted volume, biomass or carbon per hectare (\bar{V}) with the k th characteristic of the c th concentric circle. $A_{c,ij}$ is the area of the j th sample plot and c th concentric circle; m is the number of trees on the j th sample plot.
$\bar{\bar{V}}_k = \bar{V}_{3,5,k} + \bar{V}_{10,k} + \bar{V}_{15,k}$	The overall average volume, biomass or carbon per hectare with the k th characteristic ($\bar{\bar{V}}_k$) is estimated as the sum of the average volume, biomass or carbon per hectare ($\bar{V}_{c,k}$) for the three concentric circles ($c=3.5, 10$ and 15)
$V_k = \bar{\bar{V}}_k \cdot A_{Forest}$	Total volume, biomass or carbon with the k th characteristic (V_k) is the overall average volume, biomass or carbon per hectare ($\bar{\bar{V}}_k$) times the forest area A_{Forest} .

Table 3F.8 Estimation of biomass and carbon content of dead wood.

Equation	Description
$v_{s,ij} = F(d_{s,ij}, h_{s,ij}, D_{g,j})$	The volume (v_s) of the i th standing, dead tree on the j th sample plots is calculated using the existing volume functions (F) using the tree diameter and height and the squared mean diameter.
$v_{l,ij} = \frac{\pi}{4} d_{l,ij}^2 \cdot l_{l,ij}$	Volume of lying dead trees (v_l) is calculated as the length (l) and the i th tree on the j th sample plot times the cross sectional area. The cross sectional area is calculated from the mid-diameter (d) of the dead wood.
$B_{s,ij} = v_{s,ij} \cdot D_{ij} \cdot r_{k,ij}$	Biomass of the i th standing (B_s) or lying (B_l) tree on the j th sample plot is calculated as the volume (v_s or v_l) times the species specific density (D) and a the k th reduction factor according to the structural decay of the wood observed in the field.
$B_{l,ij} = v_{l,ij} \cdot D_{ij} \cdot r_{k,ij}$	
$B_{s,tot,ij} = B_{s,ij} \cdot E_{ij}$	The total above and below ground volume ($B_{s,tot}$) of the i th standing, dead tree on the j th sample plot. v_s is the calculated biomass of the tree and E is the expansion factor.
$K_{s,ij} = B_{s,ij} \cdot 0.5$	Carbon in standing or lying dead wood (C_s or C_l) is calculated as the biomass (B_s or B_l) times 0.5.
$K_{l,ij} = B_{l,ij} \cdot 0.5$	

Table 3F.9 Estimation of total biomass and carbon pools of dead wood.

Equation	Description
$V_{D,cj} = \frac{1}{A_{cj}} \sum_{i=1}^m R_c v_{s,ij} + R_c v_{l,ij}$	Deadwood volume, biomass or carbon pools per hectare (V_D) for the c th circle and the j th sample plot. v_s and v_l is the volume of standing and lying deadwood respectively. R_c is an indicator variable that is 1 if the tree is measured in the c th circle and 0 otherwise. A_c is the sample plot area of the c th circle. m is the number of trees within the j th sample plot.
$\bar{V}_{D,c} = \frac{\sum_{j=1}^n A_{cj} V_{D,cj}}{\sum_{j=1}^n A_{cj}}$	The average area weighted deadwood volume, biomass or carbon per hectare (\bar{V}_D) of the c th concentric circle. $A_{c,j}$ is the area of the j th sample plot and c th concentric circle; n is the number of sample plots.
$\bar{\bar{V}}_D = \bar{V}_{D,3.5} + \bar{V}_{D,10} + \bar{V}_{D,15}$	The overall average deadwood volume, biomass or carbon per hectare ($\bar{\bar{V}}_D$) is estimated as the sum of the average volume, biomass or carbon per hectare ($\bar{V}_{D,c}$) for the three concentric circles ($c=3.5, 10$ and 15)
$V_D = \bar{\bar{V}}_D \cdot A_{Forest}$	Total deadwood volume, biomass or carbon V_D is the overall average deadwood volume, biomass or carbon per hectare ($\bar{\bar{V}}_D$) times the forest area A_{Forest} .

Table 3F.10 Estimation of forest floor carbon.

Equation	Description
$C_{floor,s,j} = Depth_j \cdot A_j \cdot B_s \cdot F_{s,j}$	Forest floor carbon ($C_{floor,s,j}$) of the s th species, on the j th plot with an area of A . B_s is the species specific forest floor density and F is the fraction of species s .
$C_{floor,j} = \sum_{s=1}^k C_{floor,s,j}$	Total forest floor carbon on the j th plot.
$C_{floor} = \frac{\sum_{j=1}^n C_{floor,j}}{\sum_{j=1}^n A_j} \cdot A_{Forest}$	Total forest floor carbon is estimated as the area weighted average forest floor carbon content times the total forest area.

Table 3.F.11 Hectares grown in the different areas of Denmark.

	Copenhagen					Southern	Eastern	Western	Northern
	Denmark	area	Bornholm	Sealand	Funen	Jutland	Jutland	Jutland	Jutland
Winter wheat	743911	14984	14941	159521	86704	121584	122810	79914	143452
Spring wheat	13753	583	375	3862	582	2694	1764	1768	2125
Rye	51336	4017	80	4816	2753	9409	9384	7422	13454
Winter barley	142560	2459	2691	16869	13665	30254	32248	21518	22856
Spring barley	425510	6614	4091	97731	30227	86358	41749	96536	62203
Oat	41907	1835	538	2924	1724	9889	4850	7908	12239
Triticale and other cereals for maturity	50192	385	409	4276	1736	10185	8734	11705	12763
Pulses for maturity	10349	80	44	2318	680	1540	1544	2448	1695
Potatoes for seed	5189	43	0	513	146	1260	173	2488	567
Potatoes for starch production	16637	0	0	0	68	3944	412	8793	3420
Potatoes for consumption	16312	493	19	1600	986	4719	1027	5485	1984
Sugarbeet for sugar production	39074	35	0	38571	438	9	13	0	8
Sugarbeet for feeding	4118	67	40	162	133	987	470	1231	1028
Winter rape	163436	6456	2054	38341	22824	25266	28691	14552	25253
Spring rape	1372	73	0	584	42	103	201	35	333
Lin seed	90	1	0	1	4	1	2	82	0
Other industrial seed	823	21	1	537	23	88	84	45	24
Grass and other seeds for seed production	66655	1200	1818	26276	13326	4904	7561	6670	4901
Lucerne	6405	67	87	763	820	3009	493	737	428
Green maize for silage	172168	981	1893	6349	10023	69325	12591	37600	33407
Green cereals for silage	62845	722	210	1747	927	18213	4138	16546	20344
Other green feeding stuff	26	0	0	0	0	0	3	24	0
Grass and cloverfields in rotation	320914	9798	2939	22024	11361	99547	32764	72147	70334
Vegetables in fields	8043	349	24	2106	1502	514	1608	1497	444
Green peas for consumption	2677	51	0	2159	334	31	32	40	32
Flowers and other ornamentals	92	4	0	39	17	7	2	23	0
Apples	1684	60	5	562	746	73	141	34	64
Pears	357	19	1	125	163	18	21	7	5
Strawberries	1137	47	3	369	208	161	164	89	98
Cherries	1743	47	0	1051	619	8	4	1	11
Black currant	1935	91	1	472	672	211	363	16	110
Other fruits and berries	927	29	6	263	377	124	88	18	24

Continued

Nurseries	1521	87	0	154	342	347	172	331	88
Permanent grass outside rotation	199859	10533	1021	28762	12429	47085	23693	32442	43894
Set-a-side with grass	9874	504	164	2155	715	1407	938	1919	2072
Christmastrees	19521	402	67	2697	2618	2802	4607	2764	3566
Other crops	16569	610	49	1893	986	3048	2143	4041	3799
Without crops	24866	707	514	5090	2044	5487	3908	3253	3863
Total agricultural area	2646400	64451	34083	477685	222964	564612	349589	442128	490888
Green houses	490	28	1	90	272	18	59	13	10

Table 3.F.12 Crop yield from Statistics Denmark in 2010 distributed regions, Hhg crop ha⁻¹.

	Denmark	Copenhagen and North Sealand	Bornholm	Sealand	Funen	Southern Jutland	Eastern Jutland	Western Jutland	North- ern Jutland
Winter wheat	66.6	70.3	65.2	73.9	72	66.5	69.3	59	57.2
Spring wheat	46.2	45.6	41.3	46.9	35.4	47.7	45.4	44.6	48.5
Rye	48.9	54.8	44.4	55.4	66.2	48.3	48.3	43.3	45.5
Triticale	48.6	49.2	43.8	58.7	56.4	46	50.4	48.8	46.3
Winter barley	54.3	52.5	57.1	63.1	62.3	55.6	56.7	45.3	46.4
Spring barley	51	47.3	50.7	56.7	51.7	49.3	51.5	47.4	49.7
Oat and mixed cereals	48.1	50.4	48	48.1	49.6	47.6	46.7	46.1	50.3
Winter rape	34.9	33.6	39.9	38	37.7	35.1	33.2	30.9	31.8
Spring rape	22.7	26.1	..	18.2	37.6	32.4	25.9	20.1	23.9
Pulses for maturity	32.3	30.8	30.9	28	34.4	32.1	36.2	31.8	35.8
Straw, gathered	32.8	33.9	33.1	37	36.5	31.9	34.5	28.6	29.8
Potatoes for seed	282	275	..	275	275	275	300	299	238
Potatoes for starch production	413	410	408	450	403	442
Potatoes for consumption	340	406	420	414	289	327	314	348	311
Sugarbeet for sugar production	614	583	..	615	583	583	583	583	583
Sugarbeet for feeding	666	656	703	656	524	654	655	644	727
Lucerne	514	516	706	486	532	533	469	463	496
Green mais for silage	354	468	445	479	317	360	359	332	344
Green cereals for silage	174	199	241	202	169	187	164	165	167
Grass and cloverfields in rotation	438	403	459	404	449	449	453	447	419
Permanent grass outside rotation	158	163	142	170	187	159	152	152	146
Secondary grass crop yields	44	51	24	26	31	51	38	44	46

Table 3.F.13 Area input format for Eastern Jutland to C-TOOL in 2010. FK represent the soil type (Color Code (Farve Kode))

Landsdel/ Region	AFG07_txt	time	FK1	FK2	FK3	FK4	FK5	FK6	FK7	SUM
Eastern Jutland	Set-a-side with grass	2010	120	17	373	208	32	2	186	938
Eastern Jutland	Pulses for maturity	2010	456	75	539	419	45	0	9	1544
Eastern Jutland	Sugarbeet for feeding	2010	67	14	279	78	5	0	27	470
Eastern Jutland	Vegetables in fields	2010	279	40	945	254	1	0	120	1640
Eastern Jutland	Grass and other seeds for seed production	2010	566	188	3274	2812	441	7	273	7561
Eastern Jutland	Grass and cloverfields in rotation	2010	3801	391	15947	6729	1096	32	4770	32767
Eastern Jutland	Oat	2010	804	56	2309	1278	135	8	259	4850
Eastern Jutland	Lin seed	2010	1	0	1	0	0	0	0	2
Eastern Jutland	Strawberries	2010	4	0	99	55	3	0	4	164
Eastern Jutland	Potatoes	2010	245	97	917	254	3	0	97	1612
Eastern Jutland	Green cereals for silage	2010	622	117	2011	775	96	0	518	4138
Eastern Jutland	Lucerne	2010	6	0	327	127	17	0	16	493
Eastern Jutland	Green maize for silage	2010	1900	262	7863	1839	221	11	495	12591
Eastern Jutland	Rye	2010	3253	391	4062	1115	35	0	527	9384
Eastern Jutland	Sugarbeet for sugar production	2010	0	0	0	13	0	0	0	13
Eastern Jutland	Triticale and other cereals for maturity	2010	1261	162	4449	1165	126	18	444	7625
Eastern Jutland	Winter barley	2010	2250	203	17101	10283	1240	35	1114	32226
Eastern Jutland	Winter wheat	2010	4591	1207	55394	46819	8213	311	6275	122810
Eastern Jutland	Winter rape	2010	1490	230	15092	9827	1229	27	796	28691
Eastern Jutland	Spring barley	2010	4567	744	20270	12210	1557	16	2385	41749
Eastern Jutland	Spring wheat	2010	30	0	756	591	43	0	344	1764
Eastern Jutland	Spring rape	2010	39	0	79	109	18	0	39	285
Eastern Jutland	Other crops	2010	880	0	4426	565	24	0	156	6051
Eastern Jutland	Permanent grass outside rotation	2010	4101	701	8855	3284	617	54	6068	23682

Table 3F.14 Average annual temperatures for Denmark, 1977-2012, °C.

Year	Average	Year	Average
1977	7.675464	2000	9.175
1978	7.675464	2001	8.158333
1979	7.675464	2002	9.208333
1980	7.2	2003	8.708333
1981	7.15	2004	8.733333
1982	7.975	2005	8.783333
1983	8.375	2006	9.358333
1984	7.891667	2007	9.416667
1985	6.5	2008	9.366667
1986	6.933333	2009	8.775
1987	6.55	2010	6.908333
1988	8.475	2011	8.916667
1989	9.175	2012	8.275
1990	9.233333		
1991	8.108333		
1992	8.958333		
1993	7.558333		
1994	8.608333		
1995	8.183333		
1996	6.833333		
1997	8.5		
1998	8.2		
1999	8.85		




Figure 3F.1 Average annual temperatures for Denmark, 1977-2012, °C.

Annex 3G - Waste

Annex 3G-1: Emissions from the waste sector, 1990-2012

Annex 3G-2: Solid Waste Disposal on Land, 6A

Annex 3G-3: Wastewater Handling, 6B

Annex 3G-4: Waste Incineration, 6C

Annex 3G-5: Waste Other, 6D

Annex 3G-6: Recalculations to the waste sector

Annex 3G-1 Emissions from the waste sector, 1990-2012

Table 3G-1.1 Emissions for the waste sector, Gg CO₂ equivalents.

		1990	1991	1992	1993	1994	1995	1996
6. A. Solid Waste Disposal on Land	CH ₄	1366	1372	1361	1351	1285	1209	1176
6. B. Wastewater Handling	CH ₄	65	66	66	66	67	68	69
6. B. Wastewater Handling	N ₂ O	105	102	90	108	114	108	91
6. C. Waste incineration	CH ₄	0.01	0.01	0.01	0.01	0.01	0.01	0.01
6. C. Waste incineration	N ₂ O	0.20	0.20	0.20	0.21	0.21	0.21	0.21
6. D. Waste Other	CO ₂	18	18	19	18	18	20	20
6. D. Waste Other	CH ₄	31	34	37	40	43	41	47
6. D. Waste Other	N ₂ O	13	14	16	17	19	23	25
6. Waste	Total	1598	1606	1590	1600	1546	1469	1428
<i>Continued</i>		1997	1998	1999	2000	2001	2002	2003
6. A. Solid Waste Disposal on Land	CH ₄	1105	1044	1057	1047	1041	977	995
6. B. Wastewater Handling	CH ₄	71	71	70	73	73	71	73
6. B. Wastewater Handling	N ₂ O	85	87	84	90	84	97	77
6. C. Waste incineration	CH ₄	0.01	0.01	0.01	0.01	0.01	0.01	0.01
6. C. Waste incineration	N ₂ O	0.21	0.21	0.22	0.22	0.22	0.23	0.22
6. D. Waste Other	CO ₂	19	18	19	18	18	18	19
6. D. Waste Other	CH ₄	55	57	65	70	66	73	76
6. D. Waste Other	N ₂ O	29	59	109	160	154	239	233
6. Waste	Total	1363	1335	1404	1459	1437	1476	1473
<i>Continued</i>		2004	2005	2006	2007	2008	2009	2010
6. A. Solid Waste Disposal on Land	CH ₄	887	862	896	855	828	793	720
6. B. Wastewater Handling	CH ₄	72	72	73	73	72	73	74
6. B. Wastewater Handling	N ₂ O	76	84	73	84	98	73	76
6. C. Waste incineration	CH ₄	0.01	0.01	0.01	0.02	0.02	0.02	0.02
6. C. Waste incineration	N ₂ O	0.23	0.24	0.27	0.28	0.28	0.29	0.29
6. D. Waste Other	CO ₂	18	18	19	19	21	21	18
6. D. Waste Other	CH ₄	69	73	78	86	79	86	88
6. D. Waste Other	N ₂ O	63	62	74	92	90	102	110
6. Waste	Total	1184	1173	1213	1209	1189	1149	1087
<i>Continued</i>		2011	2012					
6. A. Solid Waste Disposal on Land	CH ₄	731	698					
6. B. Wastewater Handling	CH ₄	75	74					
6. B. Wastewater Handling	N ₂ O	79	73					
6. C. Waste incineration	CH ₄	0.01	0.01					
6. C. Waste incineration	N ₂ O	0.27	0.27					
6. D. Waste Other	CO ₂	18	16					
6. D. Waste Other	CH ₄	90	92					
6. D. Waste Other	N ₂ O	118	127					
6. Waste	Total	1113	1080					

Annex 3G-2 Solid Waste Disposal on Land, 6A

The following Table 3G-2.1 shows the total waste production in Denmark, divided after means of handling. (DEPA, 1996a, 1998a, 1999a, 2001a, 2001b, 2002a, 2004a, 2004b, 2005a, 2006a, 2006b, 2008, 2010, 2011a, 2013)

Table 3G-2.1 All nationally produced waste categorised after handling method, collected for the ISAG database 1994-2009 and 2011.

Year	Recycled	Combusted	Landfilled		Special treatment	Temporary storage	Total
	Gg	Gg	Gg	%	Gg	Gg	Gg
1994	6157	2216	2604	23.4	102	0	11105
1995	7046	2306	1957	17.1	145	0	11466
1996	7787	2507	2507	19.4	95	0	12912
1997	8046	2622	2083	16.2	86	0	12857
1998	7542	2740	1859	15.2	84	0	12233
1999	7815	2929	1467	11.9	17	0	12313
2000	8461	3064	1482	11.4	17	0	13031
2001	8101	3221	1300	10.2	20	109	12768
2002	8382	3344	1174	9.0	22	163	13105
2003	8218	3287	966	7.7	20	108	12614
2004	8746	3437	1000	7.5	16	136	13359
2005	9545	3473	957	6.7	18	191	14210
2006	10768	3489	975	6.3	19	181	15459
2007	10480	3584	956	6.3	20	167	15235
2008	10725	3590	1045	6.7	21	167	15575
2009	9536	3386	753	5.4	18	152	13872
2010	-	-	-	-	-	-	-
2011*	5593	2605	549	6	253	102	9102

*In reality the total deposited amount of waste is 2.1 million tonnes, but in the new waste data statistics soil and stones are not included. Furthermore, fly ash, slag from combined heat and power plants and sludge from wastewater treatment plants were not included in the waste data system at the time of publication (DEPA, 2013). Therefore the deposited amount of waste in Table 3G-2.1 differs from the actual amounts as reported in Table 3G-2.2.

Table 3G-2.2 presents the annual net emission of methane generated from the amount of landfilled waste and deducted the recovered methane and the oxidised methane; calculated using the FOD model.

Table 3G-2.2 Annual amounts of deposited waste, gross methane emission, recovered methane collected for biogas production, oxidised methane in the top layer and resulting net emission for the Danish SWDS.

Year	Landfilled waste	Gross methane	Recovered methane	Methane oxidised in the top layers	Net methane emission	
□	Gg	Gg CH ₄	Gg CH ₄	Gg CH ₄	Gg CH ₄	Gg CO ₂ eq
1990	3190	72.8	0.5	7.2	65.0	1366
1991	3050	73.3	0.7	7.3	65.4	1372
1992	2910	73.5	1.5	7.2	64.8	1361
1993	2770	73.3	1.8	7.1	64.3	1351
1994	2630	72.7	4.7	6.8	61.2	1285
1995	1969	71.6	7.6	6.4	57.6	1209
1996	2524	70.6	8.3	6.2	56.0	1176
1997	2103	69.8	11.4	5.8	52.6	1105
1998	1868	68.7	13.5	5.5	49.7	1044
1999	1552	67.6	11.7	5.6	50.3	1057
2000	1489	66.7	11.3	5.5	49.9	1047
2001	1317	65.3	10.2	5.5	49.6	1041
2002	1194	63.2	11.4	5.2	46.5	977
2003	981	60.7	8.1	5.3	47.4	995
2004	1024	58.2	11.3	4.7	42.3	887
2005	983	55.6	9.9	4.6	41.1	862
2006	1002	53.0	5.6	4.7	42.7	896
2007	984	50.8	5.5	4.5	40.7	855
2008	1072	48.7	5.0	4.4	39.4	828
2009	779	46.6	4.7	4.2	37.8	793
2010	1786	44.5	6.4	3.8	34.3	720
2011	2439	42.7	4.0	3.9	34.8	731
2012	1107	41.2	4.3	3.7	33.2	698

Tables 3G-2.3 presents activity data for Solid Waste Disposal on Land allocated according to 18 defined waste types classified according to their content of degradable organic matter, DOC_i , half-life time, $t_{1/2}$. As presented, the basis year of the FOD model is the year 1960. For a detailed description of back-calculation of the time series from the New waste data system (2010-2012) to 1960, the reader is referred to Thomsen et al. (2014).

Table 3G-2.3 Annual amounts of deposited inert and decomposable waste allocated according to 18 identified waste types characterised according to their *DOC*_i and decomposition rate quantified by their half-life times, *t*_½ (cf. Table 8.2.2 in the main report).

Waste type*/Year	1960	1961	1962	1963	1964	1965	1966	1967	1968	1969
Food	71.8	71.8	71.8	71.8	71.8	71.8	71.8	71.8	71.8	71.8
Paper and cardboard	115.7	115.7	115.7	115.7	115.7	115.7	115.7	115.7	115.7	115.7
Wood	94.7	94.7	94.7	94.7	94.7	94.7	94.7	94.7	94.7	94.7
Plastic*	16.0	16.0	16.0	16.0	16.0	16.0	16.0	16.0	16.0	16.0
Textile, fur and leather	2.6	2.6	2.6	2.6	2.6	2.6	2.6	2.6	2.6	2.6
Biodegradable garden waste	85.7	85.7	85.7	85.7	85.7	85.7	85.7	85.7	85.7	85.7
Chemicals, inert*	4.0	4.0	4.0	4.0	4.0	4.0	4.0	4.0	4.0	4.0
Electric & Hazardous*	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3
Glass*	23.0	23.0	23.0	23.0	23.0	23.0	23.0	23.0	23.0	23.0
Metal*	83.3	83.3	83.3	83.3	83.3	83.3	83.3	83.3	83.3	83.3
Scrap vehicles*	53.9	53.9	53.9	53.9	53.9	53.9	53.9	53.9	53.9	53.9
Demolition	145.9	145.9	145.9	145.9	145.9	145.9	145.9	145.9	145.9	145.9
Soil & Stone*	240.4	240.4	240.4	240.4	240.4	240.4	240.4	240.4	240.4	240.4
Particulate matter and dust*	16.6	16.6	16.6	16.6	16.6	16.6	16.6	16.6	16.6	16.6
Sludge, inert*	56.4	56.4	56.4	56.4	56.4	56.4	56.4	56.4	56.4	56.4
Sludge, degradable	131.5	131.5	131.5	131.5	131.5	131.5	131.5	131.5	131.5	131.5
Ash & Slag*	150.1	150.1	150.1	150.1	150.1	150.1	150.1	150.1	150.1	150.1
Other not combustible waste*	353.7	353.7	353.7	353.7	353.7	353.7	353.7	353.7	353.7	353.7
Total, [Gg]	1645.3	1645.3	1645.3	1645.3	1645.3	1645.3	1645.3	1645.3	1645.3	1645.3
Total inert, [Gg]	997.6	997.6	997.6	997.6	997.6	997.6	997.6	997.6	997.6	997.6
Year	1970	1971	1972	1973	1974	1975	1976	1977	1978	1979
Food	71.8	78.3	84.8	91.4	97.9	104.4	110.9	117.5	124.0	130.5
Paper and cardboard	115.7	126.2	136.7	147.2	157.7	168.3	178.8	189.3	199.8	210.3
Wood	94.7	103.3	111.9	120.5	129.1	137.7	146.3	154.9	163.5	172.1
Plastic*	16.0	17.5	18.9	20.4	21.8	23.3	24.7	26.2	27.6	29.1
Textile, fur and leather	2.6	2.8	3.0	3.3	3.5	3.7	4.0	4.2	4.4	4.7
Biodegradable garden waste	85.7	93.5	101.3	109.1	116.9	124.7	132.5	140.3	148.1	155.9
Chemicals, inert*	4.0	4.3	4.7	5.0	5.4	5.8	6.1	6.5	6.8	7.2
Electric & Hazardous*	0.3	0.3	0.3	0.4	0.4	0.4	0.4	0.5	0.5	0.5
Glass*	23.0	25.1	27.1	29.2	31.3	33.4	35.5	37.6	39.7	41.8
Metal*	83.3	90.8	98.4	106.0	113.6	121.1	128.7	136.3	143.8	151.4
Scrap vehicles*	53.9	58.8	63.7	68.6	73.5	78.4	83.3	88.2	93.1	98.0
Demolition	145.9	159.1	172.4	185.6	198.9	212.2	225.4	238.7	251.9	265.2
Soil & Stone*	240.4	262.3	284.1	306.0	327.9	349.7	371.6	393.4	415.3	437.1
Particulate matter and dust*	16.6	18.1	19.6	21.1	22.6	24.1	25.6	27.1	28.6	30.1
Sludge, inert*	56.4	61.5	66.7	71.8	76.9	82.0	87.2	92.3	97.4	102.6
Sludge, degradable	131.5	143.4	155.4	167.3	179.3	191.2	203.2	215.1	227.1	239.0
Ash & Slag*	150.1	163.8	177.4	191.1	204.7	218.4	232.0	245.7	259.3	273.0
Other not combustible waste*	353.7	385.8	418.0	450.1	482.3	514.4	546.6	578.7	610.9	643.1
Total, [Gg]	1645.3	1794.9	1944.5	2094.1	2243.7	2393.2	2542.8	2692.4	2842.0	2991.5
Total inert, [Gg]	997.6	1088.3	1179.0	1269.7	1360.4	1451.1	1541.7	1632.4	1723.1	1813.8
Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989
Food	137.0	143.6	150.1	156.6	163.2	169.7	175.1	181.5	187.9	194.3
Paper and cardboard	220.8	231.4	241.9	252.4	262.9	273.4	283.9	294.4	304.9	315.4
Wood	180.7	189.3	197.9	206.5	215.1	223.7	232.3	240.9	249.5	258.1
Plastic*	30.5	32.0	33.5	34.9	36.4	37.8	39.3	40.7	42.2	43.6
Textile, fur and leather	4.9	5.1	5.4	5.6	5.8	6.1	6.3	6.5	6.7	6.9
Biodegradable garden waste	163.7	171.5	179.3	187.1	194.8	202.6	210.4	218.2	226.0	233.8
Chemicals, inert*	7.6	7.9	8.3	8.6	9.0	9.4	9.7	10.1	10.4	10.8
Electric & Hazardous*	0.5	0.6	0.6	0.6	0.6	0.7	0.7	0.7	0.7	0.7
Glass*	43.9	45.9	48.0	50.1	52.2	54.3	56.4	58.5	60.6	62.7
Metal*	159.0	166.5	174.1	181.7	189.3	196.8	204.4	211.9	219.5	227.0
Scrap vehicles*	102.9	107.8	112.7	117.6	122.5	127.4	132.3	137.2	142.1	147.0

<i>Continued</i>										
Demolition	278.5	291.7	305.0	318.2	331.5	344.8	332.4	320.0	307.6	295.2
Soil & Stone*	459.0	480.9	502.7	524.6	546.4	568.3	547.9	527.4	507.0	486.6
Particulate matter and dust*	31.6	33.1	34.6	36.1	37.6	39.1	37.7	36.3	34.9	33.5
Sludge, inert*	107.7	112.8	117.9	123.1	128.2	133.3	124.2	115.3	106.8	98.6
Sludge, degradable	251.0	262.9	274.9	286.8	298.8	310.7	289.2	268.4	248.4	229.1
Ash & Slag*	286.6	300.3	313.9	327.6	341.2	354.9	383.2	408.5	430.6	449.8
Other not combustible waste*	675.2	707.4	739.5	771.7	803.8	836.0	796.6	757.9	719.9	682.6
Total, [Gg]	3141.1	3290.7	3440.3	3589.8	3739.4	3889.0	3749.2	3609.3	3469.5	3329.6
Total inert, [Gg]	1904.5	1995.2	2085.9	2176.6	2267.3	2358.0	2303.3	2246.3	2187.1	2125.7
Year	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
Food	111.7	101.6	91.9	82.7	74.0	52.0	62.3	48.3	39.7	30.3
Paper and cardboard	180.2	163.9	148.4	133.6	119.6	84.1	100.8	78.2	64.3	49.1
Wood	201.5	196.1	190.4	184.3	178.0	260.9	183.5	182.6	239.2	271.6
Plastic*	27.0	25.0	23.1	21.3	19.6	14.2	17.5	14.1	12.0	9.6
Textile, fur and leather	5.0	4.8	4.5	4.3	4.1	3.1	3.9	3.3	2.9	2.4
Biodegradable garden waste	136.0	124.2	113.0	102.3	92.2	65.2	78.8	61.7	51.3	39.7
Chemicals, inert*	7.7	7.3	7.0	6.7	6.3	4.7	6.1	5.1	4.5	3.7
Electric & Hazardous*	0.5	0.5	0.5	0.5	0.4	0.3	0.4	0.4	0.3	0.3
Glass*	37.3	34.3	31.4	28.6	26.0	18.5	22.6	17.9	15.0	11.8
Metal*	184.3	180.6	176.5	172.0	167.1	127.8	167.5	142.6	129.3	109.7
Scrap vehicles*	104.5	99.9	95.3	90.8	86.2	64.5	82.7	68.9	61.2	50.9
Demolition	282.8	270.4	258.0	245.6	233.2	174.5	223.7	186.4	165.6	137.6
Soil & Stone*	466.1	445.7	425.2	404.8	384.4	308.6	404.2	304.1	368.1	369.7
Particulate matter and dust*	32.1	30.7	29.3	27.9	26.5	0.0	0.0	0.0	0.0	0.8
Sludge, inert*	90.7	83.2	76.0	69.1	62.5	44.5	54.1	42.6	35.7	27.8
Sludge, degradable	210.7	193.0	176.1	159.9	109.8	135.7	155.2	138.3	136.0	143.7
Ash & Slag*	465.8	478.8	488.8	495.7	650.1	145.0	715.4	483.0	215.6	15.5
Other not combustible waste*	645.9	610.0	574.8	540.3	390.6	464.8	244.8	325.3	327.3	278.1
Total, [Gg]	3189.8	3050.0	2910.1	2770.3	2630.4	1968.5	2523.6	2102.8	1868.1	1552.4
Total inert, [Gg]	2062.0	1996.1	1927.9	1857.5	1819.6	1193.0	1715.4	1404.0	1169.1	877.9
Year	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Food	26.5	21.2	17.1	9.3	4.5	4.6	6.1	5.7	3.3	0.8
Paper and cardboard	43.0	34.4	27.9	15.1	7.4	7.5	10.2	9.4	5.6	1.5
Wood	254.8	77.9	17.8	3.6	2.1	2.6	4.5	22.8	4.8	2.2
Plastic*	8.8	7.5	6.5	5.1	5.0	4.6	4.4	4.1	4.2	2.8
Textile, fur and leather	2.3	2.1	1.9	1.1	0.6	0.8	1.3	1.5	1.3	0.6
Biodegradable garden waste	35.2	28.7	23.7	13.2	6.7	7.0	10.1	10.2	6.9	2.4
Chemicals, inert*	3.6	3.2	2.5	1.9	1.8	1.4	1.3	2.9	1.7	1.2
Electric & Hazardous*	0.7	0.6	3.5	103.3	83.7	83.7	90.2	108.3	126.3	6.6
Glass*	10.6	8.8	7.4	5.7	5.5	4.8	4.4	3.9	3.8	2.4
Metal*	107.4	96.8	89.5	74.9	79.7	77.9	80.8	80.9	89.6	66.2
Scrap vehicles*	48.8	72.3	67.2	40.1	26.0	48.7	47.1	10.1	7.1	72.3
Demolition	132.0	116.7	105.8	86.9	90.8	87.1	88.8	87.3	95.0	69.1
Soil & Stone*	271.3	327.4	306.7	171.0	233.9	174.0	157.9	154.8	201.4	203.4
Particulate matter and dust*	0.3	0.5	0.5	0.4	0.3	0.1	0.1	1.9	0.1	0.0
Sludge, inert*	25.0	20.5	17.2	13.0	12.4	10.7	9.8	8.4	7.9	4.9
Sludge, degradable	107.1	81.0	70.6	64.9	48.8	37.7	43.3	48.9	39.0	32.4
Ash & Slag*	8.5	14.5	42.2	63.8	51.0	33.8	38.8	51.7	163.7	45.9
Other not combustible waste*	402.9	402.7	385.7	307.6	363.6	395.9	402.3	371.6	310.4	264.3
Total, [Gg]	1488.8	1316.7	1193.8	980.7	1023.9	983.0	1001.5	984.4	1072.1	778.9
Total inert, [Gg]	887.9	954.8	929.0	786.6	862.8	835.6	837.2	798.6	916.2	670.1
Year	2010	2011	2012							
Food	1.0	0.7	0.9							
Paper and cardboard	1.9	2.6	1.9							
Wood	70.7	96.6	54.7							

<i>Continued</i>			
Plastic*	5.5	7.8	4.3
Textile, fur and leather	2.2	3.1	2.4
Biodegradable garden waste	0.0	22.9	5.2
Chemicals, inert*	1.0	0.6	0.1
Electric & Hazardous*	0.0	0.1	1.4
Glass*	5.4	5.3	2.9
Metal*	134.9	156.0	132.6
Scrap vehicles*	16.0	17.2	1.5
Demolition	94.8	176.7	155.4
Soil & Stone*	1321.0	1774.0	624.1
Particulate matter and dust*	2.5	5.4	25.0
Sludge, inert*	2.2	9.8	11.1
Sludge, degradable	18.5	22.8	13.4
Ash & Slag*	48.4	54.2	23.3
Other not combustible waste*	59.9	82.8	46.6
Total, [Gg]**	1786.0	2438.6	1106.8
Total inert, [Gg]	1596.8	2113.2	872.9

*Waste types characterised as inert, i.e. $DOC_i = 0$

** The reason for the seemingly increased amounts of waste deposited at landfills is due to the fact that only a part of the fraction soil and stones were included in the old ISAG waste statistics, while none is included in the new waste data system as may be observed from table 3G-2.2 (DEPA, 2013). The DEPA report on waste statistics for 2011 (2013) does however include a separate accounting of the soil and stones. In the NIR all waste fraction deposited at landfills are included (Thomsen and Hjelgaard, 2014)

Below the allocation of waste amounts reported according to the European waste codes are presented. For a detailed documentation of the whole time series including back-calculation of the time series, the reader is referred to the methodology report verifying waste amounts and how the allocation of the old ISAG waste categories and types was performed and verified (Thomsen and Hjelgaard, 2014).

Table 3G-2.4 European waste codes allocated according to 18 characterised waste types (based on the reporting years 2010-2012).

Waste types	EWG codes
Food	020199*1/7, 020203, 020299, 020304, 020399, 020601, 020699, 1906xx, 200108
Paper and cardboard	150101, 150106*1/7, 150110*1/7, 191211*1/7, 191212*1/7, 200101
Wood	020107, 020199*1/7, 030105, 030199, 030301, 030399, 150103, 150106*1/7, 150110*1/7, 170201, 170204*1/3, 191207, 191211*1/7, 191212*1/7, 200137, 200138, (200199, 200301, 200307, 200399)*0,5 other waste
Plastic	020104, 020199*1/7, 070299, 120105, 150102, 150106*1/7, 150110*1/7, 160103, 160119, 170203, 170204*1/3, 191204, 191211*1/7, 191212*1/7, 200139
Textile, fur and leather	040108, 150106*1/7, 150110*1/7, 191211*1/7, 191212*1/7
Biodegradable garden waste	1905xx, 200201
Chemicals, inert	020199*1/7, 060204, 060313, 060405, 060899, 080410, 080499, 100109, 200127
Electric & Hazardous*	160605, 170411, 200121, 200135, 200136
Glass*	101112, 101199, 150106*1/7, 150110*1/7, 160120, 170202, 170204*1/3, 191205, 191211*1/7, 191212*1/7, 200102
Metal*	010101, 010305, 010306, 010399, 020110, 020199*1/7, 100299, 100908, 101007, 101012, 101099, 120101, 120102, 150105, (150106+150110)*2/7 (komposit+metal), 170403, 170409, 190102, 191002, 191003, 191005, (191211+191212)*1/7
Scrap vehicles*	160104, 160199
Demolition	
- of which are concrete, bricks, tiles and ceramics	170101, 170102, 170106, 170107
- of which are insulation & asbestos-containing materials	161101, 161106, 170601, 170603, 170604, 170605, 170606
- of which are other demolition	170301, 170302, 170802, 170901, 170902, 170903, 170904
Soil & Stone*	010408, 010499, 010504, 010599, 170503, 170504, 170506, 191211*1/7, 191212*1/7, 200202, 200203, 200303
Particulate matter and dust*	100912, 101299, 101301, 101304, 101306, 101311, 101399, 120116, 120117, 120120, 120121, 120199, 200141
Sludge, inert*	
- of which originates from desanding and screenings at WWTPs*	190801, 190802, 190899
- of which originates from physico-chemical treatments of water(*)	070110, 110109, 190105
- of which originates from anaerobic treatment of waste*	190901, 190902, 190904, 190999
- of which are sludge and filter cakes primarily from gas treatment*	191302
- of which originates from primary filtration & clarification of water*	100215, 101307, 190106
Sludge, degradable	
- of which originates from treatment of urban WW	190805
- of which originates from biological treatment of industrial WW	190812
- of which originates from other treatment of industrial WW	020101, 030305, 030311, 040220, 060503, 190814
- of which originates from physico-chemical treatments of water	190205, 190208, 190210, 190211, 190299
- of which originates from oil/water separators and oil regeneration	130501, 130508, 191106, 191199
- of which is originates from sewage cleaning	200306
- of which is categorised as manure	020106, 020199*1/7
- of which is categorised as organic chemicals	020108, 020109, 020199*1/7, 020702, 070199, 070215, 070399, 070503, 070699, 080112, 120106, 130401, 160306, 160508, 160509, 160799, 180103, 191210, 200128
Ash & Slag*	
- of which is categorised as bottom ash, slag and boiler dust*	100101, 100114, 100115, 100118, 100119, 101313, 190111, 190112
- of is categorised as which fly ash*	100103, 100116, 100117, 190113, 190114, 190402
- of is categorised as which other*	100199, 100399, 190119, 190199, 190203, 190204
Other waste, inert*	
- of which is categorised as not combustible municipal waste	(200199, 200301, 200307, 200399)*0,5

Table 3G-2.5 Fractional distribution of waste types for the whole time series 1990-2012.

Waste types	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
Food	4.36	4.19	4.02	3.85	3.67	3.50	3.33	3.16	2.99	2.81
Paper and cardboard	7.03	6.75	6.48	6.20	5.93	5.65	5.37	5.10	4.82	4.55
Wood	5.75	5.87	5.98	6.09	6.20	6.32	6.43	6.54	6.65	6.77
Plastic*	0.97	0.95	0.92	0.90	0.87	0.85	0.82	0.79	0.77	0.74
Textile, fur and leather	0.16	0.16	0.16	0.16	0.16	0.16	0.16	0.16	0.16	0.16
Biodegradable garden waste	5.21	5.02	4.83	4.64	4.45	4.26	4.07	3.88	3.69	3.50
Chemicals, inert*	0.24	0.24	0.24	0.24	0.24	0.24	0.24	0.24	0.24	0.24
Electric & Hazardous*	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02
Glass*	1.40	1.35	1.31	1.26	1.21	1.17	1.12	1.08	1.03	0.99
Metal*	5.06	5.20	5.35	5.49	5.63	5.78	5.92	6.06	6.21	6.35
Scrap vehicles*	3.28	3.28	3.28	3.28	3.28	3.28	3.28	3.28	3.28	3.28
Demolition*	8.87	8.87	8.87	8.87	8.87	8.87	8.87	8.87	8.87	8.87
- of which is concrete, bricks, tiles and ceramics	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
- of which is insulation & asbestos-containing materials	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.23
- of which is other demolition	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Soil & Stone*	14.6	14.6	14.6	14.6	14.6	14.6	14.6	14.6	14.6	14.6
Particulate matter and dust*	1.01	1.01	1.01	1.01	1.01	1.01	1.01	1.01	1.01	1.01
Sludge, inert*	3.43	3.31	3.19	3.08	2.96	2.84	2.73	2.61	2.49	2.38
- of which is waste from desanding and screenings at WWTPs*	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
- of which is from physico/chemical treatments of water(*)	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
- of which is from anaerobic treatment of waste*	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
- of which is sludges and filter cakes primarily from gas treatment*	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
- of which is from primary filtration & clarification of water*	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Sludge, degradable	7.99	7.71	7.44	7.16	6.88	6.60	6.33	6.05	5.77	4.17
- of which is from treatment of urban WW	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
- of which is from biological treatment of industrial WW	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	1.95
- of which is from other treatment of industrial WW	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
- of which is from physico/chemical treatments of water	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
- of which is from oil/water separators and oil regeneration	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
- of which is waste from sewage cleaning	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
- of which is manure	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
- of which is organic chemicals	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
- of which is other	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.73
- of which has a dry matter content of 10-30%	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	2.33
- of which has a dry matter content above 30%	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	1.54
Ash & Slag	9.12	10.2	11.3	12.4	13.5	14.6	15.7	16.8	17.8	24.7
- of which is bottom ash, slag and boiler dust*	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	24.7
- of which is fly ash*	7.55	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.01
- of which is categorised as other*	1.57	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Other waste, inert*	21.5	21.3	21.0	20.8	20.5	20.3	20.0	19.8	19.5	14.9
- of which is not combustible municipal waste	0.45	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	9.43
- of which is not combustible garden waste	1.65	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.61
- of which is not combustible industrial waste	8.47	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	4.05
Waste types	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Food	1.78	1.61	1.44	0.94	0.44	0.46	0.61	0.57	0.31	0.11
Paper and cardboard	2.89	2.61	2.34	1.54	0.73	0.76	1.01	0.96	0.52	0.19
Wood	17.1	5.92	1.49	0.37	0.21	0.27	0.45	2.32	0.45	0.28
Plastic*	0.59	0.57	0.54	0.52	0.49	0.47	0.44	0.41	0.39	0.36
Textile, fur and leather	0.16	0.16	0.16	0.12	0.06	0.08	0.13	0.16	0.12	0.07
Biodegradable garden waste	2.37	2.18	1.99	1.34	0.65	0.72	1.01	1.04	0.65	0.30
Chemicals, inert*	0.24	0.24	0.21	0.20	0.18	0.14	0.13	0.30	0.16	0.16
Electric & Hazardous*	0.05	0.05	0.30	10.5	8.17	8.51	9.01	11.0	11.8	0.85

<i>Continued</i>										
Glass*	0.71	0.67	0.62	0.58	0.53	0.49	0.44	0.40	0.35	0.31
Metal*	7.21	7.35	7.50	7.64	7.78	7.93	8.07	8.21	8.36	8.50
Scrap vehicles*	3.28	5.49	5.63	4.09	2.54	4.96	4.71	1.03	0.67	9.28
Demolition*	8.87	8.87	8.87	8.87	8.87	8.87	8.87	8.87	8.87	8.87
- of which is concrete, bricks, tiles and ceramics	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
- of which is insulation & asbestos-containing materials	0.59	0.71	1.20	2.00	1.63	2.24	2.07	2.52	3.00	3.96
- of which is other demolition	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Soil & Stone*	18.2	24.9	25.7	17.4	22.9	17.7	15.8	15.7	18.8	26.1
Particulate matter and dust*	0.02	0.04	0.04	0.04	0.03	0.01	0.01	0.19	0.01	0.00
Sludge, inert*	1.68	1.56	1.44	1.33	1.21	1.09	0.98	0.86	0.74	0.62
- of which is waste from desanding and screenings at WWTPs*	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
- of which is from physico/chemical treatments of water(*)	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
- of which is from anaerobic treatment of waste*	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
- of which is sludges and filter cakes primarily from gas treatment*	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
- of which is from primary filtration & clarification of water*	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Sludge, degradable	7.19	6.15	5.91	6.62	4.77	3.83	4.32	4.97	3.64	4.16
- of which is from treatment of urban WW	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
- of which is from biological treatment of industrial WW	3.35	3.27	3.68	3.42	3.26	3.07	3.01	2.94	2.48	2.19
- of which is from other treatment of industrial WW	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
- of which is from physico/chemical treatments of water	0.04	0.16	0.10	0.06	0.06	0.06	0.06	0.04	0.06	0.05
- of which is from oil/water separators and oil regeneration	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
- of which is waste from sewage cleaning	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
- of which is manure	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
- of which is organic chemicals	0.00	0.01	0.00	0.00	0.00	0.03	0.04	0.00	0.00	0.00
- of which is other	0.74	1.01	0.66	0.76	0.61	0.47	0.51	0.39	0.34	0.36
- of which has a dry matter content of 10-30%	3.43	2.09	1.70	1.84	0.88	0.78	1.12	1.55	1.00	1.94
- of which has a dry matter content above 30%	1.31	1.17	1.21	1.85	1.16	0.52	0.56	0.89	0.50	0.24
Ash & Slag	0.57	1.10	3.53	6.50	4.98	3.44	3.88	5.26	15.3	5.90
- of which is bottom ash, slag and boiler dust*	0.57	1.00	3.36	6.21	4.78	3.24	3.64	5.08	15.1	5.60
- of which is fly ash*	0.00	0.10	0.17	0.29	0.19	0.20	0.24	0.18	0.21	0.30
- of which is categorised as other*	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Other waste, inert*	27.1	30.6	32.3	31.4	35.5	40.3	40.2	37.8	29.0	33.9
- of which is not combustible municipal waste	14.1	19.5	20.8	25.3	24.7	30.6	28.8	28.1	22.9	26.5
- of which is not combustible garden waste	0.23	0.23	0.23	0.07	0.04	0.05	0.07	0.03	0.03	0.02
- of which is not combustible industrial waste	6.32	4.93	4.22	0.48	4.13	0.63	1.05	1.65	0.29	0.78
Waste types	2010	2011	2012							
Food	0.06	0.03	0.09							
Paper and cardboard	0.11	0.11	0.17							
Wood	3.96	3.96	4.94							
Plastic*	0.31	0.32	0.38							
Textile, fur and leather	0.12	0.13	0.22							
Biodegradable garden waste	0.00	0.94	0.47							
Chemicals, inert*	0.05	0.02	0.01							
Electric & Hazardous*	0.00	0.00	0.13							
Glass*	0.30	0.22	0.26							
Metal*	7.55	6.40	12.0							
Scrap vehicles*	0.90	0.71	0.14							
Demolition*	5.31	7.25	14.0							
- of which is concrete, bricks, tiles and ceramics	0.28	0.21	0.29							
- of which is insulation & asbestos-containing materials	2.60	3.37	7.17							
- of which is other demolition	2.43	3.67	6.58							
Soil & Stone*	74.0	72.8	56.4							
Particulate matter and dust*	0.14	0.22	2.26							

<i>Continued</i>			
Sludge, inert*	0.12	0.40	1.00
- of which is waste from desanding and screenings at WWTPs*	0.12	0.22	0.75
- of which is from physico/chemical treatments of water(*)	0.00	0.00	0.00
- of which is from anaerobic treatment of waste*	0.00	0.04	0.06
- of which is sludges and filter cakes primarily from gas treatment*	0.00	0.05	0.08

Annex 3G-3 Wastewater Handling, 6B

Table 3G-3.1 presents the methane produced in anaerobic digester tanks, recovered for energy production, emitted from sewer system and WWTPs, primary settling tanks and biological N and P removal processes, fugitive emissions from anaerobic processes and net CH₄ emission from the 6 B. Wastewater handling in Denmark, 1990-2011.

Table 3G-3.1 Produced, recovered and emitted CH₄ from wastewater treatment, Gg, 1990-2010

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
CH ₄ gross methane	13.30	13.27	13.31	13.68	14.85	16.03	19.48	23.05	21.51	19.40
CH ₄ recovered	13.17	13.13	13.17	13.54	14.70	15.87	19.29	22.82	21.30	19.21
CH ₄ emitted from sewer system and WWTP	0.17	0.17	0.17	0.18	0.19	0.21	0.23	0.24	0.26	0.24
CH ₄ emitted from septic tanks	2.81	2.82	2.83	2.84	2.85	2.86	2.87	2.89	2.90	2.91
CH ₄ emission from anaerobic treatment	0.13	0.13	0.13	0.14	0.15	0.16	0.19	0.23	0.22	0.19
Net CH ₄ emission	3.12	3.12	3.13	3.15	3.19	3.23	3.30	3.36	3.37	3.35
<i>Continued</i>	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
CH ₄ gross methane	29.04	28.41	20.05	24.53	20.62	22.94	23.17	24.23	19.69	22.81
CH ₄ recovered	28.75	28.13	19.85	24.29	20.41	22.71	22.94	23.99	19.49	22.58
CH ₄ emitted from sewer system and WWTP	0.27	0.26	0.26	0.27	0.25	0.25	0.26	0.27	0.22	0.25
CH ₄ emitted from septic tanks	2.92	2.93	2.94	2.95	2.96	2.96	2.97	2.98	3.00	3.02
CH ₄ emission from anaerobic treatment	0.29	0.28	0.20	0.25	0.21	0.23	0.23	0.24	0.20	0.23
Net CH ₄ emission	3.48	3.48	3.40	3.47	3.41	3.45	3.46	3.49	3.41	3.50
<i>Continued</i>	2010	2011	2012							
CH ₄ gross methane	23.63	24.63	22.62							
CH ₄ recovered	23.40	24.38	22.39							
CH ₄ emitted from sewer system and WWTP	0.26	0.27	0.24							
CH ₄ emitted from septic tanks	3.03	3.04	3.06							
CH ₄ emission from anaerobic treatment	0.24	0.25	0.23							
Net CH ₄ emission	3.53	3.56	3.52							

Table 3G-3.2 shows the total N₂O emission originating from treatment processes at the Danish WWTPs (direct emissions) and effluents to the Danish surface waters (indirect emissions).

Table 3G-3.2 N₂O emissions from wastewater, Mg, 1990-2010.

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
N ₂ O, indirect	265	252	219	273	268	238	180	158	154	147
N ₂ O, direct	73	77	72	75	99	111	113	116	126	123
N ₂ O, total	339	329	292	348	368	350	292	274	280	271
<i>Continued</i>	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
N ₂ O, indirect	157	134	137	109	119	111	109	116	103	108
N ₂ O, direct	134	137	176	140	125	161	127	154	214	127
N ₂ O, total	292	272	313	249	244	272	236	270	317	235
<i>Continued</i>	2010	2011	2012							
N ₂ O, indirect	109	106	104							
N ₂ O, direct	136	150	131							
N ₂ O, total	246	256	235							

Table 3G-3.3 presents the total degradable organic waste (TOW) calculated by use of the default IPCC method corrected for contribution from industry to the influent TOW (1990-1998) and country-specific data (1999-2011).

Table 3G-3.3 Calculated total degradable organic waste (TOW), 1990-2010.

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
Contribution from industrial inlet BOD	2.5	2.5	2.5	5	13.6	22.2	30.8	39.4	48	41
Population (1000)	5135	5146	5162	5181	5197	5216	5251	5275	5295	5314
TOW [Gg] corrected IPCC method	96.5	96.3	96.6	99.3	107.7	116.3	125.3	134.2	143.0	
TOW [Gg]; country-specific data										136.0
<i>Continued</i>	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Contribution from industrial inlet BOD	42.0	38.0	38.0	37.0	40.5	40.5	40.5	40.5	40.5	40.5
Population (1000)	5330.0	5349.2	5368.4	5383.5	5397.6	5411.4	5427.5	5447.1	5475.8	5511.5
TOW [Gg]; corrected IPCC method										
TOW [Gg]; country-specific data	148.5	145.9	146.4	152.1	139.5	140.9	142.3	148.8	120.9	140.1
<i>Continued</i>	2010	2011	2012							
Contribution from industrial inlet BOD	40.5	40.5	40.5							
Population (1000)	5535	5561	5581							
TOW [Gg] corrected IPCC method										
TOW [Gg]; country-specific data	144.5	150.9	134.6							

*TOW = $(1+I/100) \times (P \times D_{\text{dom}})$, where P is the Population number, $D_{\text{dom}} = 18\ 250$ kg BOD per 1000 persons per year and I is the per cent contribution from industry to the influent wastewater TOW content.

Table 3G-3.4 shows the country-specific emission factor for estimating the methane generated during anaerobic treatment processes.

Table 3G-3.4 Emission factor for estimating the methane generation, kg CH₄/kg BOD.

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
MCf _{AD} *f _{AD}	0.23	0.23	0.23	0.23	0.23	0.23	0.26	0.29	0.25	0.24
EF _{AD} [kg CH ₄ /kg BOD] B ₀ *MCF _{AD} *f _{AD}	0.14	0.14	0.14	0.14	0.14	0.14	0.16	0.17	0.15	0.14
<i>Continued</i>	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
MCf _{AD} *f _{AD}	0.33	0.32	0.23	0.27	0.25	0.27	0.27	0.27	0.27	0.27
EF _{AD} [kg CH ₄ /kg BOD] B ₀ *MCF _{AD} *f _{AD}	0.20	0.19	0.14	0.16	0.15	0.16	0.16	0.16	0.16	0.16
<i>Continued</i>	2010	2011	2012							
MCf _{AD} *f _{AD}	0.27	0.27	0.28							
EF _{AD} [kg CH ₄ /kg BOD] B ₀ *MCF _{AD} *f _{AD}	0.16	0.16	0.17							

Table 3G-3.5 presents the nitrogen content in the influent and effluent wastewater.

Table 3G-3.5 Nitrogen content in the influent and effluent wastewater, Mg.

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
Influent wastewater to WWTPs*	14679	15398	14492	15010	19888	22340	22580	23243	25329	24738
Effluent wastewater from WWTP**	10268	9520	7480	10787	10241	8938	6387	4851	6387	5135
Effluent wastewater, total**	16884	16032	13953	17403	17079	15152	11431	10068	9796	9363
<i>Continued</i>	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Influent wastewater to WWTPs*	26952	27499	35187	28038	24991	32288	25401	30899	42808	25519
Effluent wastewater from WWTP	4653	4221	4528	3614	4027	3831	3634	4358	3575	4025
Effluent wastewater, total**	10005	8553	8740	6927	7589	7038	6935	7381	6557	6878
<i>Continued</i>	2010	2011	2012							
Influent wastewater to WWTPs*	27357	30049	26316							
Effluent wastewater from WWTP	4025	3916	3849							
Effluent wastewater, total**	6960	6770	6597							

*Data on the influent wastewater N load from municipal WWTPs are available from the Danish Water Quality Parameter Database held by the Agency for Spatial and Environmental Planning ** Effluent wastewater, total includes separate industrial discharges, rainwater conditioned effluent, scattered houses, mariculture and fish farming and effluents from WWTPs (DEPA. 1994, 1996a, 1997, 1998, 1999a, 2000, 2001a, 2002, 2003a, 2004b, 2005a, 2005b and ASEP 2007, 2009, 2010, 2011, 2012, 2013).

Table 3G-3.6 presents the per cent uncertainties on the individual parameters used for calculating the uncertainties associated with activity data and emission factors used for estimating the methane and nitrous oxide emissions from category 6.B Wastewater Handling. References are given to the equations presented in Chapter 8.3.2.

Table 3G-3.6 Input parameter uncertainties, %.

Input parameters and equations	Uncertainty, %	Reference
CH₄ (sewer+MB)		Eq. 8.3.2
EF_{sewer+MB}=B_o*MCF_{sewer+MB}	32	
B _o	30	IPCC, 2006
MCF _{sewer+MB}	10	IPCC, 2006
Ac_{sewer+MB}	24	
TOW	24	Table 3G.3
CH_{4, AD}		Eq. 8.3.3
EF_{AD}=B_o*MCF_{AD}*f_{AD}	39	
B _o	30	IPCC, 2000
MCF _{AD}	10	IPCC, 2006
F _{AD}	23	Table 3G.4
Ac_{AD}	24	
TOW	24	Table 3G.3
CH_{4, st}		Eq. 8.3.4
EF_{st}=MCF_{st}*B_o	32	
MCF _{st}	10	IPCC, 2006
B _o	30	IPCC, 2000
Ac_{st}=f_{nc}*P*DOC_{st}	31	
f _{nc}	5	IPCC, 2000
DOC _{st}	30	IPCC, 2006
P	5	IPCC, 2000
N₂O,direct		Eq. 8.3.6
EF _{N2O,direct}	50	Table 3G.5
Ac_{N2O,direct}	22	Table 3G.6
m _{N,influent}	22	Table 3G.6
N₂O,indirect		Eq. 8.3.7
EF _{N2Oindirect}	42	
D_{N,WWTP}	59	Table 3G.6

Annex 3G-4 Waste Incineration, 6C

Table 3G-4.1 presents the greenhouse gas emissions from 6.C Waste Incineration for 1990-2012.

Table 3G-4.1 Overall emission of greenhouse gases from the incineration of human bodies and animal carcasses

		1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
CH ₄ emission from											
Human cremation	Mg	0.48	0.48	0.49	0.51	0.50	0.52	0.51	0.50	0.49	0.50
Animal cremation	Mg	0.03	0.03	0.03	0.03	0.03	0.04	0.04	0.04	0.04	0.07
Total	Mg	0.51	0.51	0.52	0.54	0.54	0.55	0.55	0.54	0.53	0.56
N ₂ O emission from											
Human cremation	Mg	0.60	0.60	0.61	0.63	0.63	0.64	0.64	0.63	0.61	0.62
Animal cremation	Mg	0.03	0.04	0.04	0.04	0.04	0.05	0.05	0.05	0.05	0.08
Total	Mg	0.64	0.63	0.65	0.68	0.67	0.69	0.68	0.68	0.67	0.71
6C. Waste incineration											
CO ₂ equivalents	Gg	0.21	0.21	0.21	0.22	0.22	0.23	0.22	0.22	0.22	0.23
<i>Continued</i>		2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
CH ₄ emission from											
Human cremation	Mg	0.49	0.49	0.50	0.49	0.49	0.48	0.48	0.49	0.49	0.50
Animal cremation	Mg	0.08	0.08	0.08	0.08	0.10	0.14	0.20	0.23	0.24	0.24
Total	Mg	0.57	0.57	0.58	0.58	0.59	0.62	0.69	0.72	0.73	0.74
N ₂ O emission from											
Human cremation	Mg	0.61	0.61	0.63	0.62	0.61	0.60	0.61	0.61	0.61	0.62
Animal cremation	Mg	0.10	0.10	0.10	0.10	0.13	0.17	0.25	0.29	0.30	0.30
Total	Mg	0.71	0.72	0.73	0.72	0.74	0.77	0.86	0.90	0.92	0.93
6C. Waste incineration											
CO ₂ equivalents	Gg	0.23	0.23	0.24	0.24	0.24	0.25	0.28	0.30	0.30	0.30
<i>Continued</i>		2010	2011	2012							
CH ₄ emission from											
Human cremation	Mg	0.49	0.49	0.48							
Animal cremation	Mg	0.26	0.22	0.22							
Total	Mg	0.76	0.71	0.70							
N ₂ O emission from											
Human cremation	Mg	0.62	0.61	0.60							
Animal cremation	Mg	0.33	0.28	0.28							
Total	Mg	0.95	0.88	0.88							
6C. Waste incineration											
CO ₂ equivalents	Gg	0.31	0.29	0.29							

Table 3G-4.2 presents the activity data for human cremation for 1990-2012.

Table 3G-4.2 Activity data for human cremation.

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
Nationally deceased	60926	59581	60821	62809	61099	63127	61043	59898	58453	59179
Cremations	40991	40666	41455	43194	42762	43847	43262	42891	41660	42299
Cremation fraction, %	67.3	68.3	68.2	68.8	70.0	69.5	70.8	71.6	69.1	74.4
<i>Continued</i>										
	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Nationally deceased	57998	58355	58610	57574	55806	54962	55477	55604	54591	54872
Cremations	41651	41707	42539	41997	41555	40758	41233	41766	41788	42408
Cremation fraction, %	71.8	71.5	72.6	72.9	74.5	74.2	74.3	75.1	76.6	77.3
<i>Continued</i>										
	2010	2011	2012							
Nationally deceased	54368	52516	52325							
Cremations	42050	41248	40909							
Cremation fraction, %	77.3	78.6	79.6							

Table 3G-4.3 presents the activity data for animal cremation for 1990-2012.

Table 3G-4.3 Activity data, (direct contact with all Danish pet crematoria).

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
Total, Mg	150	160	170	180	190	200	210	220	235	368
<i>Continued</i>	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Total, Mg	443	452	451	462	571	762	1116	1284	1338	1339
<i>Continued</i>	2010	2011	2012							
Total, Mg	1449	1219	1238							

Annex 3G-5 Waste Other, 6D

Table 3G-5.1a-c presents the national emissions for source category 6D 1990-2012.

Table 3G-5.1a Overall emission of greenhouse gasses from accidental fires and composting, 1990-1999.

		1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
CO ₂ emission from											
Accidental building fires	Gg	63.1	65.1	70.7	62.2	62.6	72.2	73.0	67.5	60.4	64.9
- of which non-biogenic	Gg	11.4	11.8	12.8	11.2	11.3	13.1	13.2	12.2	10.9	11.7
Accidental vehicle fires	Gg	6.1	6.2	6.2	6.4	6.4	6.5	6.7	6.7	6.7	6.8
Total, non-biogenic	Gg	17.5	17.9	19.0	17.7	17.7	19.6	19.9	18.9	17.7	18.5
CH ₄ emission from											
Compost production	Mg	1386.1	1532.5	1678.7	1825.1	1972.9	1860.2	2171.0	2526.6	2628.0	3032.5
Accidental building fires	Mg	64.1	66.2	71.8	63.2	63.6	73.4	74.1	68.5	61.3	65.9
Accidental vehicle fires	Mg	12.8	12.9	12.9	13.4	13.4	13.6	13.9	13.9	14.1	14.2
Total	Mg	1463.0	1611.5	1763.5	1901.7	2049.9	1947.2	2259.0	2609.0	2703.3	3112.6
N ₂ O emission from											
Compost production	Mg	41.5	46.2	51.1	55.9	60.6	72.8	79.4	93.1	190.6	350.0
Accidental building fires	Mg	NAV	NAV	NAV	NAV	NAV	NAV	NAV	NAV	NAV	NAV
Accidental vehicle fires	Mg	NAV	NAV	NAV	NAV	NAV	NAV	NAV	NAV	NAV	NAV
Total	Mg	41.5	46.2	51.1	55.9	60.6	72.8	79.4	93.1	190.6	350.0
6D. Waste other											
CO ₂ equivalents	Gg	61.1	66.1	71.9	74.9	79.6	83.0	91.9	102.5	133.5	192.4

Table 3G-5.1b Overall emission of greenhouse gasses from accidental fires and composting, 2000-2009.

		2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
CO ₂ emission from											
Accidental building fires	Gg	63.8	63.3	61.5	69.5	60.1	62.4	64.2	76.3	72.6	69.6
- of which non-biogenic	Gg	11.5	11.4	11.1	12.6	10.9	11.3	11.6	13.7	13.3	12.6
Accidental vehicle fires	Gg	6.9	6.9	6.8	6.8	6.7	6.9	7.1	5.7	8.2	8.5
Total, non-biogenic	Gg	18.4	18.3	18.0	19.3	17.6	18.1	18.7	19.4	21.5	21.1
CH ₄ emission from											
Compost production	Mg	3240.0	3059.7	3397.1	3534.3	3222.1	3419.9	3627.9	4016.7	3685.7	4011.2
Accidental building fires	Mg	64.9	64.5	62.8	71.0	61.5	63.8	65.6	75.2	74.6	71.3
Accidental vehicle fires	Mg	14.3	14.3	14.2	14.1	14.0	14.3	14.8	11.8	17.0	17.7
Total	Mg	3319.2	3138.5	3474.1	3619.5	3297.6	3497.9	3708.4	4103.7	3777.4	4100.2
N ₂ O emission from											
Compost production	Mg	515.7	498.1	770.9	752.5	201.7	200.2	239.1	295.2	291.8	330.3
Accidental building fires	Mg	NAV	NAV	NAV	NAV	NAV	NAV	NAV	NAV	NAV	NAV
Accidental vehicle fires	Mg	NAV	NAV	NAV	NAV	NAV	NAV	NAV	NAV	NAV	NAV
Total	Mg	515.7	498.1	770.9	752.5	201.7	200.2	239.1	295.2	291.8	330.3
6D. Waste other											
CO ₂ equivalents	Gg	248.0	238.6	329.9	328.6	149.4	153.6	170.7	197.0	191.3	209.6

Table 3G-5.1c Overall emission of greenhouse gasses from accidental fires and composting, 2010-2012.

		2010	2011	2012
CO ₂ emission from				
Accidental building fires	Gg	61.7	67.6	60.5
- of which non-biogenic	Gg	11.1	12.2	10.8
Accidental vehicle fires	Gg	7.3	6.3	5.6
Total, non-biogenic	Gg	18.3	18.4	16.4
CH ₄ emission from				
Compost production	Mg	4094.6	4204.3	4306.3
Accidental building fires	Mg	64.6	68.5	61.7
Accidental vehicle fires	Mg	15.1	13.1	11.6
Total	Mg	4174.3	4285.9	4379.6
N ₂ O emission from				
Compost production	Mg	355.2	381.4	409.1
Accidental building fires	Mg	NAV	NAV	NAV
Accidental vehicle fires	Mg	NAV	NAV	NAV
Total	Mg	355.2	381.4	409.1
6D. Waste other				
CO ₂ equivalents	Gg	216.1	226.7	235.2

Table 3G-5.2 presents the activity data for composting 1990-2012.

Table 3G-5.2 Activity data composting, Gg, 1990-2011.

Year	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
Composting of garden and park waste	288	320	351	383	414	376	452	528	551	634
Composting of organic waste from households and other	16	19	23	26	29	40	38	47	43	49
Composting of sludge	NO	NO	NO	NO	NO	7	6	7	57	134
Home composting of garden and vegetable food waste	20	20	20	20	21	21	21	21	21	21
Total	324	359	394	429	464	444	517	603	672	838
<i>Continued</i>	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Composting of garden and park waste	677	630	685	716	682	737	782	876	795	847
Composting of organic waste from households and other	47	52	63	66	53	45	48	44	46	70
Composting of sludge	218	211	348	336	53	50	67	91	94	107
Home composting of garden and vegetable food waste	21	21	22	22	22	22	22	22	22	23
Total	963	914	1118	1140	810	854	919	1033	957	1047
<i>Continued</i>	2010	2011	2012							
Composting of garden and park waste	877	901	924							
Composting of organic waste from households and other	58	59	59							
Composting of sludge	120	132	145							
Home composting of garden and vegetable food waste	23	23	23							
Total	1078	1115	1151							

NO = Not occurring.

Table 3G-5.3 presents the occurrence of all accidental fires, building fires and vehicle fires, 1990-2012. Building and vehicle fires do not make up for all the national accidental fires. The total number of registered fires also include a portion of fires that does not fit into either building or vehicle fires, these are here called “Other fires” and will include e.g. a chair burning at a marked but mainly consist of “unknown/other” objects at “unknown/other open” locations.

Table 3G-5.3 Occurrence of accidental fires, 1990-2011.

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
All fires	17025	17589	19124	16803	16918	19543	19756	18236	16320	17538
Building fires	10187	10524	11443	10054	10123	11694	11821	10911	9765	10494
Vehicle fires	3354	3465	3767	3310	3333	3850	3892	3592	3215	3455
Other fires	3485	3600	3914	3439	3463	4000	4043	3732	3340	3589
<i>Continued</i>	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
All fires	17174	16894	16362	18443	15927	16551	16965	18263	20643	18930
Building fires	10276	10108	9790	11035	9530	9903	10151	12527	12124	10652
Vehicle fires	3383	3328	3223	3633	3137	3260	3342	3223	4068	3930
Other fires	3515	3458	3349	3775	3260	3387	3472	2513	4451	4348
<i>Continued</i>	2010	2011	2012							
All fires	16728	16157	14084							
Building fires	9325	11447	9932							
Vehicle fires	3459	3255	2889							
Other fires	3944	1455	1263							

Table 3G-5.4 presents the full scale equivalent activity data of accidental building fires.

Table 3G-5.4 Accidental building fires full scale equivalent activity data.

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
Container fires	750	775	842	740	745	861	870	803	719	772
Detached house fires	777	802	873	767	772	892	901	832	745	800
Undetached house fires	231	238	259	228	229	265	268	247	221	237
Apartment building fires	367	379	412	362	365	421	426	393	352	378
Industry building fire	320	331	360	316	318	368	372	343	307	330
Additional building fires	437	451	490	431	434	501	507	468	418	450
<i>Continued</i>	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Container fires	756	744	721	812	701	729	747	958	962	799
Detached house fires	784	771	747	841	727	755	774	757	886	876
Undetached house fires	233	229	222	250	216	224	230	343	278	208
Apartment building fires	370	364	353	398	343	357	366	405	433	413
Industry building fire	323	318	308	347	300	311	319	435	346	344
Additional building fires	440	433	420	473	408	424	435	483	523	466
<i>Continued</i>	2010	2011	2012							
Container fires	594	729	584							
Detached house fires	833	818	742							
Undetached house fires	194	206	181							
Apartment building fires	348	362	327							
Industry building fire	281	334	298							
Additional building fires	429	740	610							

Table 3G-5.5a-c presents emission factors for 1990-2012 for accidental fires in detached houses, undetached houses and apartment buildings respectively.

Table 3G-5.5a Emission factors for accidental detached building fires, 1990-2012.

Detached houses		1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
CO ₂ - total	Mg	30.6	30.5	30.5	30.5	30.5	30.4	30.3	30.4	30.4	30.4
CO ₂ - biogenic	Mg	25.0	24.9	24.8	24.9	24.8	24.8	24.7	24.8	24.7	24.8
CO ₂ - non-biogenic	Mg	5.7	5.7	5.6	5.7	5.6	5.6	5.6	5.6	5.6	5.6
CH ₄	kg	40.6	40.4	40.3	40.4	40.3	40.2	40.2	40.3	40.2	40.3
<i>Continued</i>		2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
CO ₂ - total	Mg	30.7	31.3	31.6	31.8	31.9	31.8	32.0	31.4	31.6	31.7
CO ₂ - biogenic	Mg	25.0	25.5	25.7	25.9	26.0	25.9	26.1	25.6	25.7	25.9
CO ₂ - non-biogenic	Mg	5.7	5.8	5.9	5.9	5.9	5.9	5.9	5.8	5.8	5.9
CH ₄	kg	40.6	41.5	41.8	42.1	42.3	42.1	42.4	41.6	41.8	42.0
<i>Continued</i>		2010	2011	2012							
CO ₂ - total	Mg	32.0	32.3	32.4							
CO ₂ - biogenic	Mg	26.1	26.3	26.4							
CO ₂ - non-biogenic	Mg	5.9	6.0	6.0							
CH ₄	kg	42.3	42.7	43.0							

Table 3G-5.5b Emission factors for accidental undetached building fires, 1990-2012.

Undetached houses		1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
CO ₂ - total	Mg	25.3	25.2	25.2	25.2	25.2	25.2	25.3	25.4	25.5	25.6
CO ₂ - biogenic	Mg	20.6	20.6	20.5	20.5	20.5	20.6	20.6	20.7	20.7	20.8
CO ₂ - non-biogenic	Mg	4.7	4.7	4.7	4.7	4.7	4.7	4.7	4.7	4.7	4.7
CH ₄	kg	33.5	33.4	33.4	33.4	33.4	33.4	33.5	33.6	33.7	33.8
<i>Continued</i>		2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
CO ₂ - total	Mg	25.7	25.7	25.7	25.8	25.8	25.7	25.8	25.9	26.0	26.1
CO ₂ - biogenic	Mg	20.9	20.9	21.0	21.0	21.0	21.0	21.0	21.1	21.2	21.3
CO ₂ - non-biogenic	Mg	4.8	4.8	4.8	4.8	4.8	4.8	4.8	4.8	4.8	4.8
CH ₄	kg	34.0	34.0	34.1	34.1	34.2	34.1	34.2	34.3	34.5	34.6
<i>Continued</i>		2010	2011	2012							
CO ₂ - total	Mg	26.2	26.0	26.2							
CO ₂ - biogenic	Mg	21.4	21.2	21.4							
CO ₂ - non-biogenic	Mg	4.9	4.8	4.9							
CH ₄	kg	34.7	34.4	34.7							

Table 3G-5.5c Emission factors for accidental apartment building fires, 1990-2012.

Apartment buildings		1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
CO ₂ - total	Mg	14.7	14.7	14.7	14.7	14.7	14.7	14.7	14.7	14.7	14.7
CO ₂ - biogenic	Mg	12.0	12.0	12.0	12.0	12.0	12.0	12.0	12.0	12.0	12.0
CO ₂ - non-biogenic	Mg	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7
CH ₄	kg	19.5	19.5	19.5	19.5	19.5	19.5	19.5	19.5	19.5	19.5
<i>Continued</i>		2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
CO ₂ - total	Mg	14.7	14.7	14.8	14.8	14.8	14.8	14.9	15.0	15.0	15.1
CO ₂ - biogenic	Mg	12.0	12.0	12.0	12.0	12.1	12.1	12.1	12.2	12.2	12.3
CO ₂ - non-biogenic	Mg	2.7	2.7	2.7	2.7	2.7	2.7	2.8	2.8	2.8	2.8
CH ₄	kg	19.5	19.5	19.5	19.6	19.6	19.7	19.7	19.8	19.9	20.0
<i>Continued</i>		2010	2011	2012							
CO ₂ - total	Mg	15.1	15.2	15.2							
CO ₂ - biogenic	Mg	12.3	12.4	12.4							
CO ₂ - non-biogenic	Mg	2.8	2.8	2.8							
CH ₄	kg	20.0	20.2	20.2							

Table 3G-5.6 states the average building floor space, 1990-2012.

Table 3G-5.6 Average floor space in building types.

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
Detached houses	156	156	155	155	155	155	155	155	155	155
Undetached houses	129	128	128	128	128	129	129	129	130	130
Apartment buildings	75	75	75	75	75	75	75	75	75	75
Industrial buildings	500	500	500	500	500	500	500	500	500	500
Additional buildings	20	20	20	20	20	20	20	20	20	20
<i>Continued</i>	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Detached houses	156	160	161	162	163	162	163	160	161	162
Undetached houses	131	131	131	131	132	131	132	132	133	133
Apartment buildings	75	75	75	75	75	76	76	76	77	77
Industrial buildings	500	500	500	500	500	500	500	500	500	500
Additional buildings	20	20	20	20	20	20	20	20	20	20
<i>Continued</i>	2010	2011	2012							
Detached houses	163	164	165							
Undetached houses	134	132	134							
Apartment buildings	77	78	78							
Industrial buildings	500	500	500							
Additional buildings	20	20	20							

Table 3G-5.7a-c presents the number of nationally registered vehicles and the number of full scale equivalent accidental vehicle fires, 1990-2012.

Table 3G-5.7a Number of nationally registered vehicles and full scale equivalent vehicle fires

	Passenger Cars		Buses		Light Duty Vehicles		Heavy Duty Vehicles	
	Registered	FSE fires	Registered	FSE fires	Registered	FSE fires	Registered	FSE fires
1990	1645454	479	8109	12	192317	19	45664	58
1991	1649168	480	9989	14	197435	19	45494	58
1992	1659795	483	11259	16	202802	20	45510	58
1993	1678919	488	13513	19	211755	21	46228	59
1994	1672022	486	14261	20	219639	21	47329	60
1995	1733242	504	14371	21	228074	22	48077	61
1996	1792971	522	14594	21	234404	23	48319	61
1997	1840845	535	14690	21	240762	23	48785	62
1998	1877740	546	14894	21	249462	24	49697	63
1999	1905855	554	14953	21	259214	25	50443	64
2000	1916364	557	15051	22	272386	27	50227	64
2001	1932440	562	15005	22	283031	28	49885	63
2002	1946073	566	14971	21	295581	29	49208	62
2003	1948717	567	14989	22	309614	30	48653	62
2004	1967432	572	14997	22	336038	33	48318	61
2005	2012216	585	15131	22	372674	36	49311	63
2006	2093809	609	15243	22	414625	40	50777	64
2007	2155940	518	15052	16	402558	19	51832	46
2008	2187104	666	14854	24	398717	44	50606	71
2009	2201550	729	14794	23	373687	48	46585	67
2010	2246675	646	14577	23	362385	38	44813	60
2011	2281539	584	13915	13	343355	43	43640	54
2012	2326778	514	13177	11	318668	32	42326	53

Table 3G-5.7b Number of nationally registered vehicles and full scale equivalent vehicle fires

	Motorcycles/Mopeds		Caravans		Train		Ship	
	Regis-	FSE fires	Registered	FSE fires	Regis-	FSE fires	Registered	FSE fires
1990	163133	58	86257	24	7156	9	2324	26
1991	162357	57	88278	24	7212	9	2312	26
1992	157912	56	90299	25	7438	9	2307	26
1993	155325	55	93150	26	7496	9	2140	24
1994	153365	54	94551	26	7117	8	2027	22
1995	165272	58	95831	26	6854	8	1911	21
1996	178188	63	97592	27	6631	8	1841	20
1997	191772	68	99931	27	6428	8	1761	19
1998	205129	72	102302	28	5861	7	1696	19
1999	219577	78	104852	29	5525	7	1695	19
2000	233309	82	106935	29	4907	6	1759	19
2001	243020	86	108924	30	4561	5	1797	20
2002	253375	89	110995	30	4169	5	1878	21
2003	256438	91	113338	31	4048	5	1838	20
2004	263472	93	116930	32	3273	4	1783	20
2005	273904	97	121350	33	3195	4	1792	20
2006	287840	102	126011	35	3002	4	1789	20
2007	302900	99	131708	36	2617	2	1755	20
2008	308538	122	136905	45	2588	3	1728	20
2009	307335	128	140366	34	2489	5	1742	22
2010	301562	83	142354	37	2740	2	1773	16
2011	295488	91	142764	34	2943	3	1768	21
2012	295798	82	142654	33	3055	2	1772	14

Table 3G-5.7c Number of nationally registered vehicles and full scale equivalent vehicle fires

	Airplane		Tractor		Combined Harvester		Bicycle	Other Transport	Machine
	Registered	FSE fires	Registered	FSE fires	Regis-	FSE fires	FSE fires	FSE fires	FSE fires
1990	1055	1	131880	82	33594	56			
1991	1059	1	131637	82	32542	54			
1992	1066	1	128205	80	31460	52			
1993	1059	1	129747	81	31502	52			
1994	1063	1	123596	77	29775	49			
1995	1058	1	130028	81	27986	46			
1996	1088	1	120480	75	28609	47			
1997	1094	1	124067	77	25418	42			
1998	1091	1	115509	72	25452	42			
1999	1087	1	115978	72	22961	38			
2000	1070	1	111736	69	23272	39			
2001	1089	1	110300	69	22811	38			
2002	1149	1	108865	68	22349	37			
2003	1083	1	107430	67	21888	36			
2004	1055	1	105994	66	21426	36			
2005	1073	1	104551	65	20965	35			
2006	1039	1	102619	64	20504	34			
2007	1058	1	102619	52	20042	19	2	85	75
2008	1077	1	102619	62	19581	34	4	97	135
2009	1122	1	102619	64	19119	43	3	93	111
2010	1152	1	102619	77	18889	32	4	58	94
2011	1132	0	102619	59	18889	21	3	50	111
2012	1111	0	102619	68	18889	18	2	50	115

Table 3G-5.8 presents the average weight of passenger cars, buses, vans, trucks and motorcycles/mopeds in 1990-2012.

Table 3G-5.8 Average weight of different vehicle categories, kg, 1990-2012.

	Cars	Buses	Vans	Trucks	Motorcycles/ Mopeds
1990	850	10000	2000	15000	86
1991	850	10000	2000	15000	88
1992	850	10000	2000	15000	91
1993	901	10068	2297	14732	93
1994	908	10512	2382	14674	96
1995	923	10807	2492	14801	97
1996	935	10899	2638	14928	98
1997	948	10950	2746	14987	99
1998	964	10960	2848	15111	100
1999	982	11140	2964	15223	102
2000	999	11195	3103	15214	103
2001	1012	11312	3238	14888	105
2002	1024	11387	3333	14486	107
2003	1039	11479	3442	14026	109
2004	1052	11572	3561	13599	112
2005	1068	11560	3793	13258	116
2006	1086	11684	4120	13179	120
2007	1105	11753	4505	13268	124
2008	1122	11700	4710	13246	127
2009	1134	11642	4682	12802	130
2010	1144	11804	4498	11883	133
2011	1154	11907	4296	11291	135
2012	1160	11625	4150	10844	136

The following Table 3G-5.9 shows the annual amount of combusted vehicle in accidental fires.

Table 3G-5.9a Burnt mass of different vehicle and machine categories, Mg

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
Passenger cars	407	408	410	440	442	466	488	508	527	544
Buses	116	143	162	195	215	223	228	231	234	239
Light duty vehicles	37	38	40	47	51	55	60	64	69	75
Heavy duty vehicles	869	865	866	864	881	902	915	927	952	974
Motorcycle, moped	5	5	5	5	5	6	6	7	7	8
Other transport	-	-	-	-	-	-	-	-	-	-
Caravan	30	31	32	35	35	36	38	39	41	42
Train	128	129	133	132	125	121	118	115	106	100
Ship	257	256	255	238	236	228	222	213	205	209
Airplane	12	12	12	11	11	11	12	12	12	12
Bicycle	-	-	-	-	-	-	-	-	-	-
Tractor	164	164	159	185	183	202	198	212	205	214
Combined harvester	530	521	512	520	499	476	494	445	452	413
Machine	-	-	-	-	-	-	-	-	-	-
Total	2555	2572	2585	2673	2683	2727	2778	2774	2811	2832

Table 3G-5.9b Burnt mass of different vehicle and machine categories, Mg

<i>Continued</i>	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Passenger cars	557	569	580	589	602	625	662	572	748	827
Buses	242	244	245	247	249	251	256	182	283	264
Light duty vehicles	82	89	96	104	117	138	166	86	207	223
Heavy duty vehicles	969	942	904	865	833	829	849	608	936	863
Motorcycle, moped	8	9	10	10	10	11	12	12	16	17
Other transport	-	-	-	-	-	-	-	47	54	53
Caravan	44	45	47	48	51	53	56	59	75	57
Train	89	81	72	68	53	51	47	33	39	63
Ship	218	225	236	233	228	229	231	234	230	253
Airplane	12	12	12	11	10	10	10	8	13	13
Bicycle	-	-	-	-	-	-	-	0	0	0
Tractor	216	222	226	230	235	247	263	235	290	301
Combined harvester	425	422	419	416	412	409	405	231	415	533
Machine	-	-	-	-	-	-	-	33	61	50
Total	2863	2863	2849	2825	2805	2858	2963	2367	3397	3537

Table 3G-5.9c Burnt mass of different vehicle and machine categories, Mg

<i>Continued</i>	2010	2011	2012
Passenger cars	739	674	592
Buses	266	160	130
Light duty vehicles	171	185	133
Heavy duty vehicles	715	606	579
Motorcycle, moped	11	12	11
Other transport	33	29	29
Caravan	63	59	57
Train	24	28	23
Ship	189	249	160
Airplane	7	3	5
Bicycle	0	0	0
Tractor	347	254	283
Combined harvester	398	271	236
Machine	43	51	53
Total	3025	2624	2319

Annex 3G-6 Recalculations to the waste sector

Table 3G-6.1 Changes in emissions from Solid Waste Disposal on Land compared with the CRF reported last year.

SWDS	Unit	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
CH ₄ , previous inventory	Gg	70.4	70.4	69.5	68.5	64.7	60.5	58.3	54.2	50.9	51.1
CH ₄ , recalculated	Gg	65.0	65.4	64.8	64.3	61.2	57.6	56.0	52.6	49.7	50.3
Change, CO ₂ equivalents	Gg	-112.0	-106.9	-98.9	-87.5	-72.7	-60.1	-47.6	-34.3	-24.5	-16.4
Change	%	-7.6	-7.2	-6.8	-6.1	-5.4	-4.7	-3.9	-3.0	-2.3	-1.5
<i>Continued</i>	Unit	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
CH ₄ , previous inventory	Gg	50.6	50.3	46.9	47.4	42.2	40.8	42.2	40.0	38.6	36.7
CH ₄ , recalculated	Gg	49.9	49.6	46.5	47.4	42.3	41.1	42.7	40.7	39.4	37.8
Change, CO ₂ equivalents	Gg	-14.9	-14.4	-8.2	-1.9	2.0	5.9	9.6	13.9	17.1	21.2
Change	%	-1.4	-1.4	-0.8	-0.2	0.2	0.7	1.1	1.6	2.1	2.7
<i>Continued</i>	Unit	2010	2011	2012							
CH ₄ , previous inventory	Gg	34.3	33.3								
CH ₄ , recalculated	Gg	34.3	34.8	33.2							
Change, CO ₂ equivalents	Gg	0.2	32.3								
Change	%	0.0	4.6								

Table 3G-6.2 Changes in emissions from Wastewater Handling compared with the CRF reported last year.

Wastewater Handling	Unit	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
CH ₄ , previous inventory	Gg	3.15	3.16	3.17	3.19	3.22	3.27	3.34	3.42	3.43	3.41
CH ₄ , recalculated	Gg	3.12	3.12	3.13	3.15	3.19	3.23	3.30	3.36	3.37	3.35
N ₂ O, previous inventory	Gg	0.34	0.33	0.29	0.35	0.37	0.35	0.29	0.27	0.28	0.27
N ₂ O, recalculated	Gg	0.34	0.33	0.29	0.35	0.37	0.35	0.29	0.27	0.28	0.27
Change, CO ₂ equivalents	Gg	-0.70	-0.70	-0.70	-0.72	-0.78	-0.84	-1.02	-1.21	-1.13	-1.34
Change	%	-0.41	-0.41	-0.45	-0.41	-0.43	-0.48	-0.64	-0.77	-0.71	-0.86
<i>Continued</i>	Unit	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
CH ₄ , previous inventory	Gg	3.52	3.54	3.49	3.56	3.51	3.54	3.53	3.55	3.56	3.59
CH ₄ , recalculated	Gg	3.48	3.48	3.40	3.47	3.41	3.45	3.46	3.49	3.41	3.50
N ₂ O, previous inventory	Gg	0.29	0.27	0.31	0.25	0.24	0.27	0.24	0.27	0.32	0.24
N ₂ O, recalculated	Gg	0.29	0.27	0.31	0.25	0.24	0.27	0.24	0.27	0.32	0.24
Change, CO ₂ equivalents	Gg	-0.90	-1.35	-1.77	-1.94	-2.12	-1.93	-1.56	-1.28	-3.00	-1.87
Change	%	-0.55	-0.85	-1.04	-1.28	-1.42	-1.21	-1.06	-0.81	-1.73	-1.26
<i>Continued</i>	Unit	2010	2011	2012							
CH ₄ , previous inventory	Gg	3.59	3.62								
CH ₄ , recalculated	Gg	3.53	3.56	3.52							
N ₂ O, previous inventory	Gg	0.25	0.26								
N ₂ O, recalculated	Gg	0.25	0.26	0.23							
Change, CO ₂ equivalents	Gg	-1.24	-1.29								
Change	%	-0.82	-0.83								

Table 3G-6.3 Changes in emissions from Waste Incineration compared with the CRF reported last year.

Waste Incineration	Unit	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
CH ₄ , previous inventory	Mg	0.51	0.51	0.52	0.54	0.54	0.55	0.55	0.54	0.53	0.56
CH ₄ , recalculated	Mg	0.51	0.51	0.52	0.54	0.54	0.55	0.55	0.54	0.53	0.56
N ₂ O, previous inventory	Mg	0.64	0.63	0.65	0.68	0.67	0.69	0.68	0.68	0.67	0.71
N ₂ O, recalculated	Mg	0.64	0.63	0.65	0.68	0.67	0.69	0.68	0.68	0.67	0.71
Change, CO ₂ equivalents	Gg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Change	%	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<i>Continued</i>	Unit	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
CH ₄ , previous inventory	Mg	0.57	0.57	0.58	0.58	0.59	0.62	0.69	0.72	0.73	0.74
CH ₄ , recalculated	Mg	0.57	0.57	0.58	0.58	0.59	0.62	0.69	0.72	0.73	0.74
N ₂ O, previous inventory	Mg	0.71	0.72	0.73	0.72	0.74	0.77	0.86	0.90	0.92	0.93
N ₂ O, recalculated	Mg	0.71	0.72	0.73	0.72	0.74	0.77	0.86	0.90	0.92	0.93
Change, CO ₂ equivalents	Gg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Change	%	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<i>Continued</i>	Unit	2010	2011	2012							
CH ₄ , previous inventory	Mg	0.76	0.71								
CH ₄ , recalculated	Mg	0.76	0.71	0.70							
N ₂ O, previous inventory	Mg	0.95	0.88								
N ₂ O, recalculated	Mg	0.95	0.88	0.88							
Change, CO ₂ equivalents	Gg	0.00	0.00								
Change	%	0.00	0.00								

Table 3G-6.4 Changes in emissions from Waste Other compared with the CRF reported last year.

Waste Other	Unit	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
CO ₂ , previous inventory	Gg	18.28	18.65	19.65	18.25	18.29	20.11	20.37	19.30	18.09	18.91
CO ₂ , recalculated	Gg	17.54	17.94	18.99	17.66	17.75	19.60	19.86	18.85	17.65	18.53
CH ₄ , previous inventory	Gg	1.40	1.54	1.68	1.81	1.94	1.80	2.12	2.43	2.52	2.88
CH ₄ , recalculated	Gg	1.46	1.61	1.76	1.90	2.05	1.95	2.26	2.61	2.70	3.11
N ₂ O, previous inventory	Gg	0.04	0.04	0.05	0.05	0.05	0.05	0.06	0.07	0.08	0.11
N ₂ O, recalculated	Gg	0.04	0.05	0.05	0.06	0.06	0.07	0.08	0.09	0.19	0.35
Change, CO ₂ equivalents	Gg	1.61	2.10	2.74	3.26	3.76	9.18	8.35	10.30	36.47	78.67
Change	%	2.70	3.28	3.97	4.54	4.96	12.43	10.00	11.17	37.59	69.17
<i>Continued</i>	Unit	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
CO ₂ , previous inventory	Gg	18.77	18.61	18.20	19.52	17.73	18.20	18.72	19.29	21.43	21.01
CO ₂ , recalculated	Gg	18.40	18.31	17.96	19.35	17.60	18.14	18.71	19.35	21.49	21.07
CH ₄ , previous inventory	Gg	3.06	2.87	3.11	3.25	3.08	3.31	3.50	3.91	3.57	3.80
CH ₄ , recalculated	Gg	3.32	3.14	3.47	3.62	3.30	3.50	3.71	4.10	3.78	4.10
N ₂ O, previous inventory	Gg	0.13	0.13	0.16	0.17	0.10	0.10	0.11	0.13	0.12	0.13
N ₂ O, recalculated	Gg	0.52	0.50	0.77	0.75	0.20	0.20	0.24	0.30	0.29	0.33
Change, CO ₂ equivalents	Gg	123.38	120.45	195.54	189.62	36.15	33.42	43.05	55.38	57.28	67.82
Change	%	99.03	101.91	145.53	136.40	31.93	27.80	33.72	39.09	42.75	47.85
<i>Continued</i>	Unit	2010	2011	2012							
CO ₂ , previous inventory	Gg	18.19	18.21								
CO ₂ , recalculated	Gg	18.35	18.45	16.36							
CH ₄ , previous inventory	Gg	3.91	4.02								
CH ₄ , recalculated	Gg	4.17	4.29	4.38							
N ₂ O, previous inventory	Gg	0.14	0.14								
N ₂ O, recalculated	Gg	0.36	0.38	0.41							
Change, CO ₂ equivalents	Gg	73.10	79.76								
Change	%	51.10	54.28								

Table 3G-6.5 Changes in emissions from the waste sector compared with the CRF reported last year.

Waste	Unit	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
CO ₂ , previous inventory	Gg	18.3	18.6	19.6	18.3	18.3	20.1	20.4	19.3	18.1	18.9
CO ₂ , recalculated	Gg	17.5	17.9	19.0	17.7	17.7	19.6	19.9	18.9	17.7	18.5
CH ₄ , previous inventory	Gg	74.9	75.1	74.4	73.5	69.8	65.5	63.7	60.1	56.8	57.4
CH ₄ , recalculated	Gg	69.6	70.1	69.7	69.4	66.4	62.8	61.6	58.6	55.8	56.8
N ₂ O, previous inventory	Gg	0.4	0.4	0.3	0.4	0.4	0.4	0.4	0.3	0.4	0.4
N ₂ O, recalculated	Gg	0.4	0.4	0.3	0.4	0.4	0.4	0.4	0.4	0.5	0.6
Change, CO ₂ equivalents	Gg	-111.1	-105.5	-96.8	-85.0	-69.8	-51.7	-40.2	-25.2	10.8	60.9
Change	%	-6.5	-6.2	-5.7	-5.0	-4.3	-3.4	-2.7	-1.8	0.8	4.5
<i>Continued</i>	Unit	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
CO ₂ , previous inventory	Gg	18.8	18.6	18.2	19.5	17.7	18.2	18.7	19.3	21.4	21.0
CO ₂ , recalculated	Gg	18.4	18.3	18.0	19.3	17.6	18.1	18.7	19.4	21.5	21.1
CH ₄ , previous inventory	Gg	57.2	56.7	53.5	54.3	48.7	47.6	49.3	47.5	45.7	44.1
CH ₄ , recalculated	Gg	56.7	56.2	53.4	54.4	49.0	48.0	49.9	48.3	46.6	45.4
N ₂ O, previous inventory	Gg	0.4	0.4	0.5	0.4	0.3	0.4	0.4	0.4	0.4	0.4
N ₂ O, recalculated	Gg	0.8	0.8	1.1	1.0	0.4	0.5	0.5	0.6	0.6	0.6
Change, CO ₂ equivalents	Gg	107.5	104.7	185.5	185.8	36.1	37.4	51.1	68.0	71.4	87.1
Change	%	8.0	7.9	14.4	14.4	3.1	3.3	4.4	6.0	6.4	8.2
<i>Continued</i>	Unit	2010	2011	2012							
CO ₂ , previous inventory	Gg	18.19	18.21								
CO ₂ , recalculated	Gg	18.35	18.45	16.36							
CH ₄ , previous inventory	Gg	41.8	40.9								
CH ₄ , recalculated	Gg	42.0	42.7	41.1							
N ₂ O, previous inventory	Gg	0.38	0.40								
N ₂ O, recalculated	Gg	0.60	0.64	0.64							
Change, CO ₂ equivalents	Gg	72.0	110.8								
Change	%	7.1	11.1								

Table 3F.1 Estimation of forest percentage and forest area.

Equation	Description
$X_j = \frac{A_j}{A_{15,j}}$	The forest percentage (X) of the j th sample plot (SSU) is estimated as the forested area (A) divided by the total area of the 15 m radius sample plot ($A_{15,j}$).
$\bar{X}_Z = \frac{1}{n_Z} \sum_Z X_j R_j$	Average forest percentage (\bar{X}) of all inventoried plots (SSU) with forest status Z based on aerial photos. R_j is an indicator variable that is 1 for inventoried plots and 0 otherwise. n_Z is the number of inventoried plots identified as forest or OWL from the air photos.
$\bar{\bar{X}} = \frac{1}{n} \left(\sum_{j=1}^n X_j R_j + N_{21} \bar{X}_1 + N_{22} \bar{X}_2 \right)$	Overall average forest percentage ($\bar{\bar{X}}$). n is the total number of inventoried and non-inventoried sample plots. N_{21} and N_{22} is the number of non-inventoried sample plots with forest and OWL, respectively.
$A_{Forest} = \bar{\bar{X}} \cdot A_{Total}$	Total forest area. A_{Total} is the total land area, $\bar{\bar{X}}$ is the estimated forest percentage and A_{Forest} is the total forest area.

Table 3F.2 Estimation of forest area with a specific characteristic.

Equation	Description
$X_k = \frac{\sum_{j=1}^n R_{jk} A_j}{\sum_{j=1}^n A_j}$	Proportion of the forest area with a given characteristic (X_k). R_{jk} is an indicator variable which is 1 if the forest area on the j th sample plots has the k th characteristic and 0 otherwise. A_j is the sample plot area and n is the total number of inventoried sample plots with forest cover.
$A_k = X_k \cdot A_{Forest}$	Total area with a given characteristic (A_k). X_k is the estimated proportion of the forest area with the k th characteristic and A_{Forest} is the total forest area.

Table 3F.3 Estimation of diameter-height equations.

Equation	Description
$h_{ij} = 13 + (\bar{h}_j - 13) \cdot \exp \left(\alpha_1 \cdot \left(1 - \frac{\bar{d}_j}{d_{ij}} \right) + \alpha_2 \cdot \left(\frac{1}{\bar{d}_j} - \frac{1}{d_{ij}} \right) \right)$	Site specific dh-regression for calculating height of trees not measured for height. h_{ij} and d_{ij} is the height and diameter of the i 'th tree on the j 'th sample plot. \bar{h}_j and \bar{d}_j are the average height and diameter of trees measured for height on the j th sample plot. α_1 and α_2 are species and growth-region specific parameters
$h_{ij} = 13 + \beta_1 \cdot \exp \left(-\frac{\beta_2}{d_{ij}} \right)$	General dh-regression for calculating height of trees not measured for height. h_{ij} and d_{ij} is the height and diameter of the i 'th tree on the j 'th sample plot. β_1 and β_2 are species and growth-region specific parameters

Table 3F.4 Estimation of quadratic mean diameter.

Equation	Description
$g_{ij} = \frac{\pi}{4} d_{ij}^2$	Basal area (g) of the i th tree on the j th plot is calculated from the diameter at breast height (d) (1.3 m above ground) assuming a circular stem form.
$G_j = \sum_{i=1}^m \frac{1}{A_{c,ij}} g_{ij}$	Basal area per hectare (G) the j th sample plot is calculated as the scaled sum of individual tree basal areas. Basal area (g) of the i th tree on the j th sample plot is scaled according to the plot area ($A_{c,ij}$) of the c th concentric circle ($c=3,5; 10; 15$ m).
$N_j = \sum_{i=1}^m \frac{1}{A_{c,ij}}$	Stem number per hectare (N) the j th sample plot is calculated as the scaled number of individual trees. The i th tree on the j th sample plot is scaled according to the plot area ($A_{c,ij}$) of the c th concentric circle ($c=3,5; 10; 15$ m).
$D_{g,j} = \sqrt{\frac{4}{\pi} \frac{G_j}{N_j}}$	The mean squared diameter is calculated from the calculated basal area and stem number for each plot.

Table 3F.5 Estimation of biomass and carbon of trees.

Equation	Description
$v_{ij} = F(d_{ij}, h_{ij}, D_{g,j})$	The volume (v) of the i th tree on the j th sample plots is calculated using the existing volume functions (F) using the tree diameter and height and the quadratic mean diameter.
$B_{ij} = V_{ij} \cdot Density_{ij}$	Biomass (B) of the i th tree on the j th sample plot is estimated as the total volume (V_{tot}) times the species specific density.
$E_{ij} = F(d_{ij}, h_{ij})$	Expansion factor model for beech and Norway spruce
$v_{tot,ij} = B_{ij} \cdot E_{ij}$	The total above and below ground volume (v_{tot}) of the i th tree on the j th sample plot. B_{ij} is the calculated above-ground biomass of the tree and E is the expansion factor.
$C_{ij} = B_{ij} \cdot 0.5$	Carbon of the i th tree on the j th sample plot is calculated as the biomass (B) times 0.5.

Table 3F.6 Estimation of total biomass and carbon pools.

Equation	Description
$V_{cj} = \frac{1}{A_{cj}} \sum_{i=1}^m R_{c,i} v_{ij}$	Volume, biomass or carbon per hectare (V) of the c th concentric circle on the j th sample plot ($c=3,5; 10; 15$ m). R_c is an indicator variable that is 1 if the i th tree is measured on the c th circle and 0 otherwise. $A_{c,ij}$ is the area of the j th sample plot and c th concentric circle; m is the number of trees on the j th sample plot.
$\bar{V}_c = \frac{\sum_{j=1}^n A_{cj} V_{cj}}{\sum_{j=1}^n A_{cj}}$	The average area weighted volume, biomass or carbon per hectare (\bar{V}) of the c th concentric circle. $A_{c,ij}$ is the area of the j th sample plot and c th concentric circle; n is the number of sample plots.
$\bar{\bar{V}} = \bar{V}_{3,5} + \bar{V}_{10} + \bar{V}_{15}$	The overall average volume, biomass or carbon per hectare ($\bar{\bar{V}}$) is estimated as the sum of the average volume, biomass or carbon per hectare (\bar{V}_c) for the three concentric circles ($c=3.5, 10$ and 15)
$V = \bar{\bar{V}} \cdot A_{Skov}$	Total volume, biomass or carbon V is the overall average volume, biomass or carbon per hectare ($\bar{\bar{V}}$) times the forest area A_{Forest} .

Table 3F.7 Estimation of biomass and carbon with a given characteristic.

Equation	Description
$V_{cj,k} = \frac{1}{A_{cj}} \sum_{i=1}^m R_{c,ij} R_{k,ij} v_{ij}$	Volume, biomass or carbon per hectare (V) with the k th characteristic of the c th concentric circle on the j th sample plot ($c=3,5; 10; 15$ m). R_c is an indicator variable that is 1 if the i th tree is measured on the c th circle and 0 otherwise. R_k is an indicator variable that is 1 if the tree has k th characteristic and 0 otherwise. $A_{c,ij}$ is the area of the j th sample plot and c th concentric circle; m is the number of trees on the j th sample plot.
$\bar{V}_{c,k} = \frac{\sum_{j=1}^n A_{cj} V_{cj,k}}{\sum_{j=1}^n A_{cj}}$	The average area weighted volume, biomass or carbon per hectare (\bar{V}) with the k th characteristic of the c th concentric circle. $A_{c,ij}$ is the area of the j th sample plot and c th concentric circle; m is the number of trees on the j th sample plot.
$\bar{\bar{V}}_k = \bar{V}_{3,5,k} + \bar{V}_{10,k} + \bar{V}_{15,k}$	The overall average volume, biomass or carbon per hectare with the k th characteristic ($\bar{\bar{V}}_k$) is estimated as the sum of the average volume, biomass or carbon per hectare ($\bar{V}_{c,k}$) for the three concentric circles ($c=3.5, 10$ and 15)
$V_k = \bar{\bar{V}}_k \cdot A_{Forest}$	Total volume, biomass or carbon with the k th characteristic (V_k) is the overall average volume, biomass or carbon per hectare ($\bar{\bar{V}}_k$) times the forest area A_{Forest} .

Table 3F.8 Estimation of biomass and carbon content of dead wood.

Equation	Description
$v_{s,ij} = F(d_{s,ij}, h_{s,ij}, D_{g,j})$	The volume (v_s) of the i th standing, dead tree on the j th sample plots is calculated using the existing volume functions (F) using the tree diameter and height and the squared mean diameter.
$v_{l,ij} = \frac{\pi}{4} d_{l,ij}^2 \cdot l_{l,ij}$	Volume of lying dead trees (v_l) is calculated as the length (l) and the i th tree on the j th sample plot times the cross sectional area. The cross sectional area is calculated from the mid-diameter (d) of the dead wood.
$B_{s,ij} = v_{s,ij} \cdot D_{ij} \cdot r_{k,ij}$	Biomass of the i th standing (B_s) or lying (B_l) tree on the j th sample plot is calculated as the volume (v_s or v_l) times the species specific density (D) and a the k th reduction factor according to the structural decay of the wood observed in the field.
$B_{l,ij} = v_{l,ij} \cdot D_{ij} \cdot r_{k,ij}$	
$B_{s,tot,ij} = B_{s,ij} \cdot E_{ij}$	The total above and below ground volume ($B_{s,tot}$) of the i th standing, dead tree on the j th sample plot. v_s is the calculated biomass of the tree and E is the expansion factor.
$K_{s,ij} = B_{s,ij} \cdot 0.5$	Carbon in standing or lying dead wood (C_s or C_l) is calculated as the biomass (B_s or B_l) times 0.5.
$K_{l,ij} = B_{l,ij} \cdot 0.5$	

Table 3F.9 Estimation of total biomass and carbon pools of dead wood.

Equation	Description
$V_{D,cj} = \frac{1}{A_{cj}} \sum_{i=1}^m R_c v_{s,ij} + R_c v_{l,ij}$	Deadwood volume, biomass or carbon pools per hectare (V_D) for the c th circle and the j th sample plot. v_s and v_l is the volume of standing and lying deadwood respectively. R_c is an indicator variable that is 1 if the tree is measured in the c th circle and 0 otherwise. A_c is the sample plot area of the c th circle. m is the number of trees within the j th sample plot.
$\bar{V}_{D,c} = \frac{\sum_{j=1}^n A_{cj} V_{D,cj}}{\sum_{j=1}^n A_{cj}}$	The average area weighted deadwood volume, biomass or carbon per hectare (\bar{V}_D) of the c th concentric circle. $A_{c,j}$ is the area of the j th sample plot and c th concentric circle; n is the number of sample plots.
$\bar{\bar{V}}_D = \bar{V}_{D,3.5} + \bar{V}_{D,10} + \bar{V}_{D,15}$	The overall average deadwood volume, biomass or carbon per hectare ($\bar{\bar{V}}_D$) is estimated as the sum of the average volume, biomass or carbon per hectare ($\bar{V}_{D,c}$) for the three concentric circles ($c=3.5, 10$ and 15)
$V_D = \bar{\bar{V}}_D \cdot A_{Forest}$	Total deadwood volume, biomass or carbon V_D is the overall average deadwood volume, biomass or carbon per hectare ($\bar{\bar{V}}_D$) times the forest area A_{Forest} .

Table 3F.10 Estimation of forest floor carbon.

Equation	Description
$C_{floor,s,j} = Depth_j \cdot A_j \cdot B_s \cdot F_{s,j}$	Forest floor carbon ($C_{floor,s,j}$) of the s th species, on the j th plot with an area of A . B_s is the species specific forest floor density and F is the fraction of species s .
$C_{floor,j} = \sum_{s=1}^k C_{floor,s,j}$	Total forest floor carbon on the j th plot.
$C_{floor} = \frac{\sum_{j=1}^n C_{floor,j}}{\sum_{j=1}^n A_j} \cdot A_{Forest}$	Total forest floor carbon is estimated as the area weighted average forest floor carbon content times the total forest area.

Table 3.F.11 Hectares grown in the different areas of Denmark.

	Copenhagen					Southern	Eastern	Western	Northern
	Denmark	area	Bornholm	Sealand	Funen	Jutland	Jutland	Jutland	Jutland
Winter wheat	743911	14984	14941	159521	86704	121584	122810	79914	143452
Spring wheat	13753	583	375	3862	582	2694	1764	1768	2125
Rye	51336	4017	80	4816	2753	9409	9384	7422	13454
Winter barley	142560	2459	2691	16869	13665	30254	32248	21518	22856
Spring barley	425510	6614	4091	97731	30227	86358	41749	96536	62203
Oat	41907	1835	538	2924	1724	9889	4850	7908	12239
Triticale and other cereals for maturity	50192	385	409	4276	1736	10185	8734	11705	12763
Pulses for maturity	10349	80	44	2318	680	1540	1544	2448	1695
Potatoes for seed	5189	43	0	513	146	1260	173	2488	567
Potatoes for starch production	16637	0	0	0	68	3944	412	8793	3420
Potatoes for consumption	16312	493	19	1600	986	4719	1027	5485	1984
Sugarbeet for sugar production	39074	35	0	38571	438	9	13	0	8
Sugarbeet for feeding	4118	67	40	162	133	987	470	1231	1028
Winter rape	163436	6456	2054	38341	22824	25266	28691	14552	25253
Spring rape	1372	73	0	584	42	103	201	35	333
Lin seed	90	1	0	1	4	1	2	82	0
Other industrial seed	823	21	1	537	23	88	84	45	24
Grass and other seeds for seed production	66655	1200	1818	26276	13326	4904	7561	6670	4901
Lucerne	6405	67	87	763	820	3009	493	737	428
Green maize for silage	172168	981	1893	6349	10023	69325	12591	37600	33407
Green cereals for silage	62845	722	210	1747	927	18213	4138	16546	20344
Other green feeding stuff	26	0	0	0	0	0	3	24	0
Grass and cloverfields in rotation	320914	9798	2939	22024	11361	99547	32764	72147	70334
Vegetables in fields	8043	349	24	2106	1502	514	1608	1497	444
Green peas for consumption	2677	51	0	2159	334	31	32	40	32
Flowers and other ornamentals	92	4	0	39	17	7	2	23	0
Apples	1684	60	5	562	746	73	141	34	64
Pears	357	19	1	125	163	18	21	7	5
Strawberries	1137	47	3	369	208	161	164	89	98
Cherries	1743	47	0	1051	619	8	4	1	11
Black currant	1935	91	1	472	672	211	363	16	110
Other fruits and berries	927	29	6	263	377	124	88	18	24

Continued

Nurseries	1521	87	0	154	342	347	172	331	88
Permanent grass outside rotation	199859	10533	1021	28762	12429	47085	23693	32442	43894
Set-a-side with grass	9874	504	164	2155	715	1407	938	1919	2072
Christmastrees	19521	402	67	2697	2618	2802	4607	2764	3566
Other crops	16569	610	49	1893	986	3048	2143	4041	3799
Without crops	24866	707	514	5090	2044	5487	3908	3253	3863
Total agricultural area	2646400	64451	34083	477685	222964	564612	349589	442128	490888
Green houses	490	28	1	90	272	18	59	13	10

Table 3.F.12 Crop yield from Statistics Denmark in 2010 distributed regions, Hhg crop ha⁻¹.

	Denmark	Copenhagen and North Sealand	Bornholm	Sealand	Funen	Southern Jutland	Eastern Jutland	Western Jutland	North- ern Jutland
Winter wheat	66.6	70.3	65.2	73.9	72	66.5	69.3	59	57.2
Spring wheat	46.2	45.6	41.3	46.9	35.4	47.7	45.4	44.6	48.5
Rye	48.9	54.8	44.4	55.4	66.2	48.3	48.3	43.3	45.5
Triticale	48.6	49.2	43.8	58.7	56.4	46	50.4	48.8	46.3
Winter barley	54.3	52.5	57.1	63.1	62.3	55.6	56.7	45.3	46.4
Spring barley	51	47.3	50.7	56.7	51.7	49.3	51.5	47.4	49.7
Oat and mixed cereals	48.1	50.4	48	48.1	49.6	47.6	46.7	46.1	50.3
Winter rape	34.9	33.6	39.9	38	37.7	35.1	33.2	30.9	31.8
Spring rape	22.7	26.1	..	18.2	37.6	32.4	25.9	20.1	23.9
Pulses for maturity	32.3	30.8	30.9	28	34.4	32.1	36.2	31.8	35.8
Straw, gathered	32.8	33.9	33.1	37	36.5	31.9	34.5	28.6	29.8
Potatoes for seed	282	275	..	275	275	275	300	299	238
Potatoes for starch production	413	410	408	450	403	442
Potatoes for consumption	340	406	420	414	289	327	314	348	311
Sugarbeet for sugar production	614	583	..	615	583	583	583	583	583
Sugarbeet for feeding	666	656	703	656	524	654	655	644	727
Lucerne	514	516	706	486	532	533	469	463	496
Green mais for silage	354	468	445	479	317	360	359	332	344
Green cereals for silage	174	199	241	202	169	187	164	165	167
Grass and cloverfields in rotation	438	403	459	404	449	449	453	447	419
Permanent grass outside rotation	158	163	142	170	187	159	152	152	146
Secondary grass crop yields	44	51	24	26	31	51	38	44	46

Table 3.F.13 Area input format for Eastern Jutland to C-TOOL in 2010. FK represent the soil type (Color Code (Farve Kode))

Landsdel/ Region	AFG07_txt	time	FK1	FK2	FK3	FK4	FK5	FK6	FK7	SUM
Eastern Jutland	Set-a-side with grass	2010	120	17	373	208	32	2	186	938
Eastern Jutland	Pulses for maturity	2010	456	75	539	419	45	0	9	1544
Eastern Jutland	Sugarbeet for feeding	2010	67	14	279	78	5	0	27	470
Eastern Jutland	Vegetables in fields	2010	279	40	945	254	1	0	120	1640
Eastern Jutland	Grass and other seeds for seed production	2010	566	188	3274	2812	441	7	273	7561
Eastern Jutland	Grass and cloverfields in rotation	2010	3801	391	15947	6729	1096	32	4770	32767
Eastern Jutland	Oat	2010	804	56	2309	1278	135	8	259	4850
Eastern Jutland	Lin seed	2010	1	0	1	0	0	0	0	2
Eastern Jutland	Strawberries	2010	4	0	99	55	3	0	4	164
Eastern Jutland	Potatoes	2010	245	97	917	254	3	0	97	1612
Eastern Jutland	Green cereals for silage	2010	622	117	2011	775	96	0	518	4138
Eastern Jutland	Lucerne	2010	6	0	327	127	17	0	16	493
Eastern Jutland	Green maize for silage	2010	1900	262	7863	1839	221	11	495	12591
Eastern Jutland	Rye	2010	3253	391	4062	1115	35	0	527	9384
Eastern Jutland	Sugarbeet for sugar production	2010	0	0	0	13	0	0	0	13
Eastern Jutland	Triticale and other cereals for maturity	2010	1261	162	4449	1165	126	18	444	7625
Eastern Jutland	Winter barley	2010	2250	203	17101	10283	1240	35	1114	32226
Eastern Jutland	Winter wheat	2010	4591	1207	55394	46819	8213	311	6275	122810
Eastern Jutland	Winter rape	2010	1490	230	15092	9827	1229	27	796	28691
Eastern Jutland	Spring barley	2010	4567	744	20270	12210	1557	16	2385	41749
Eastern Jutland	Spring wheat	2010	30	0	756	591	43	0	344	1764
Eastern Jutland	Spring rape	2010	39	0	79	109	18	0	39	285
Eastern Jutland	Other crops	2010	880	0	4426	565	24	0	156	6051
Eastern Jutland	Permanent grass outside rotation	2010	4101	701	8855	3284	617	54	6068	23682

Table 3F.14 Average annual temperatures for Denmark, 1977-2012, °C.

Year	Average	Year	Average
1977	7.675464	2000	9.175
1978	7.675464	2001	8.158333
1979	7.675464	2002	9.208333
1980	7.2	2003	8.708333
1981	7.15	2004	8.733333
1982	7.975	2005	8.783333
1983	8.375	2006	9.358333
1984	7.891667	2007	9.416667
1985	6.5	2008	9.366667
1986	6.933333	2009	8.775
1987	6.55	2010	6.908333
1988	8.475	2011	8.916667
1989	9.175	2012	8.275
1990	9.233333		
1991	8.108333		
1992	8.958333		
1993	7.558333		
1994	8.608333		
1995	8.183333		
1996	6.833333		
1997	8.5		
1998	8.2		
1999	8.85		




Figure 3F.1 Average annual temperatures for Denmark, 1977-2012, °C.

Annex 4 - CO₂ reference approach and comparison with sectoral approach, and relevant information on the national energy balance

Please refer to Chapter 3.4 and Annex 3A.

Annex 5 - Assessment of completeness and (potential) sources and sinks of greenhouse gas emissions and removals excluded

GHG inventory

The Danish greenhouse gas emission inventories for 1990-2012 include all sources identified by the Revised 1996 IPCC Guidelines and the 2000 IPCC Good Practice Guidance. Some very minor sources have not been estimated due to lack of methodology, activity data or emission factors, i.e.:

In the solvent and other product use sector currently only N₂O emissions from anaesthesia and some other minor uses are included in CRF category 3D, Denmark will try to obtain activity data for use of N₂O in aerosol cans. N₂O emissions from anaesthesia are only included from 2000 onwards.

Direct and indirect CH₄ emissions from agricultural soils are not estimated. Direct and indirect soil emissions are considered of minor importance for CH₄. No methodology is available in the IPCC Guidelines.

Emissions from harvested wood products are not reported due to lack of data. Several possible sources of CH₄ in the LULUCF sector are also reported as not estimated. For more detail please see Chapter 7.

In the Waste sector CO₂ emissions from managed waste disposal on land are not estimated. According to the 1996 IPCC Guidelines: "Decomposition of organic material derived from biomass sources (e.g., crops, forests), which are regrown on an annual basis is the primary source of CO₂ released from waste. Hence, these CO₂ emissions are not treated as net emissions from waste in the IPCC Methodology."

Emissions of N₂O from accidental fires are reported as not estimated due to lack of emission factors.

KP-LULUCF inventory

The KP-LULUCF inventory is considered complete. Please see Chapter 11 for further documentation.

Annex 6 - Additional information to be considered as part of the annual inventory submission and the supplementary information required under Article 7, paragraph 1, of the Kyoto Protocol or other useful reference information

Tables A6.1 to A6.5 below contain the information publically available in this report. Table A6.6 includes the list of discrepancies identified by the ITL (no discrepancies in the 2014 submission).

Table A6.1 Total quantities of Kyoto Protocol units by account type at beginning of reported year.

Account type	Unit type					
	AAUs	ERUs	RMUs	CERs	tCERs	ICERs
Party holding accounts	152 118 662	6 978 987	3 531 167	1 381 333	NO	NO
Entity holding accounts	3809	66 585	NO	1 477 492	NO	NO
Article 3.3/3.4 net source cancellation accounts	45 099	NO	335 864	NO		
Non-compliance cancellation accounts	NO	NO	NO	NO		
Other cancellation accounts	159 986	13 374	NO	2017	NO	NO
Retirement account	131 237 240	2 284 518	288 245	2 558 075	NO	NO
tCER replacement account for expiry	NO	NO	NO	NO	NO	
ICER replacement account for expiry	NO	NO	NO	NO		
ICER replacement account for reversal of storage	NO	NO	NO	NO		NO
ICER replacement account for non-submission of certification report	NO	NO	NO	NO		NO
Total	283 564 796	9 343 464	4 155 276	5 418 917	NO	NO

Table A6.2a Annual internal transactions.

Transaction type	Additions						Subtractions					
	Unit type						Unit type					
	AAUs	ERUs	RMUs	CERs	tCERs	ICERs	AAUs	ERUs	RMUs	CERs	tCERs	ICERs
Article 6 issuance and conversion												
Party-verified projects		NO					NO		NO			
Independently verified projects		NO					NO		NO			
Article 3.3 and 3.4 issuance or cancellation												
3.3 Afforestation and reforestation			664 882				NO	NO	NO	NO		
3.3 Deforestation			NO				NO	NO	51 586	NO		
3.4 Forest management			NO				NO	NO	NO	NO		
3.4 Cropland management			2 122 565				NO	NO	NO	NO		
3.4 Grazing land management			NO				NO	NO	164 745	NO		
3.4 Revegetation			NO				NO	NO	NO	NO		
Article 12 afforestation and reforestation												
Replacement of expired tCERs							NO	NO	NO	NO	NO	
Replacement of expired ICERs							NO	NO	NO	NO		
Replacement for reversal of storage							NO	NO	NO	NO		NO
Replacement for non-submission of certification report							NO	NO	NO	NO		NO
Other cancellation							1414	NO	NO	3115	NO	NO
Sub-total		NO	2 787 447				1414	NO	216 331	3115	NO	NO

Table A6.2a Annual internal transactions.

Retirement - Unit type						
Transaction type	AAUs	ERUs	RMUs	CERs	tCERs	ICERs
Retirement		NO	NO	NO	NO	NO

Table A6.2b Annual external transactions.

Transfers and acquisitions	Additions - Unit type						Subtractions - Unit type					
	AAUs	ERUs	RMUs	CERs	tCERs	ICERs	AAUs	ERUs	RMUs	CERs	tCERs	ICERs
EU	NO	5 366 847	NO	3 156 052	NO	NO	16 020 508	282 197	NO	2 500 523	NO	NO
CDM	NO	NO	NO	3 656 371	NO	NO	NO	NO	NO	NO	NO	NO
JP	NO	NO	NO	113 193	NO	NO	NO	NO	NO	NO	NO	NO
BG	NO	253 416	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
PL	NO	339 267	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
RU	NO	255 543	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
Sub-total	NO	6 215 073	NO	6 925 616	NO	NO	16 020 508	282 197	NO	2 500 523	NO	NO
Additional information												
Independently verified ERUs							NO					

Table A6.2c Total annual transactions.

	AAUs	ERUs	RMUs	CERs	tCERs	ICERs	AAUs	ERUs	RMUs	CERs	tCERs	ICERs
Total (Sum of tables 2a and 2b)	NO	6 215 073	2 787 447	6 925 616	NO	NO	16 021 922	282 197	216 331	2 503 638	NO	NO

Table A6.3 Expiry, cancellation and replacement.

	Expiry, cancellation and requirement to replace		Replacement					
	Unit type		Unit type					
Transaction or event type	tCERs	ICERs	AAUs	ERUs	RMUs	CERs	tCERs	ICERs
Temporary CERs (tCERS)								
Expired in retirement and replacement accounts	NO							
Replacement of expired tCERs			NO	NO	NO	NO	NO	
Expired in holding accounts	NO							
Cancellation of tCERs expired in holding accounts	NO							
Long-term CERs (ICERs)								
Expired in retirement and replacement accounts		NO						
Replacement of expired ICERs			NO	NO	NO	NO		
Expired in holding accounts		NO						
Cancellation of ICERs expired in holding accounts		NO						
Subject to replacement for reversal of storage		NO						
Replacement for reversal of storage			NO	NO	NO	NO		NO
Subject to replacement for non-submission of certification report		NO						
Replacement for non-submission of certification report			NO	NO	NO	NO		NO
Total			NO	NO	NO	NO	NO	NO

Table A6.4 Total quantities of Kyoto Protocol units by account type at end of reported year.

Account type	Unit type					
	AAUs	ERUs	RMUs	CERs	tCERs	ICERs
Party holding accounts	136 096 740	12 763 022	6 102 283	7 104 476	NO	NO
Entity holding accounts	3809	215 426	NO	176 327	NO	NO
Article 3.3/3.4 net source cancellation accounts	45 099	NO	552 195	NO		
Non-compliance cancellation accounts	NO	NO	NO	NO		
Other cancellation accounts	161 400	13 374	NO	5132	NO	NO
Retirement account	131 237 240	2 284 518	288 245	2 558 075	NO	NO
tCER replacement account for expiry	NO	NO	NO	NO	NO	
ICER replacement account for expiry	NO	NO	NO	NO		
ICER replacement account for reversal of storage	NO	NO	NO	NO		NO
ICER replacement account for non-submission of certification report	NO	NO	NO	NO		NO
Total	267 544 288	15 276 340	6 942 723	9 844 010	NO	NO

Table A6.5 (a). Summary information on additions and subtractions.

	Additions - Unit type						Subtractions – Unit type					
	AAUs	ERUs	RMUs	CERs	tCERs	ICERs	AAUs	ERUs	RMUs	CERs	tCERs	ICERs
Starting values	276 838 955											
Issuance pursuant to Article 3.7 and 3.8												
Non-compliance cancellation							NO	NO	NO	NO		
Carry-over	NO	NO		NO								
Sub-total	276 838 955	NO		NO			NO	NO	NO	NO		
Annual transactions												
Year 0 (2007)	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
Year 1 (2008)	191 772 275	NO	NO	27 439 229	NO	NO	174 391 031	NO	NO	23 971 973	NO	NO
Year 2 (2009)	881 590 260	524 201	NO	32 057 896	NO	NO	874 991 940	185 735	NO	33 349 553	NO	NO
Year 3 (2010)	233 649 660	5 344 875	NO	28 111 141	NO	NO	252 411 415	1389 977	NO	28 008 871	NO	NO
Year 4 (2011)	8 593 901	2 249 840	624 109	3 022 739	NO	NO	6 160 750	NO	335 864	86 669	NO	NO
Year 5 (2012)	1 255 753	4 939 318	3 531 167	3 728 306	NO	NO	2 385 957	2 152 432	NO	3 525 345	NO	NO
Year 6 (2013)	NO	6 215 073	2 787 447	6 925 616	NO	NO	16 021 922	282 197	216 331	2 503 638	NO	NO
Year 7 (2014)	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
Year 8 (2015)	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
Sub-total	1 316 861 849	19 273 307	6 942 723	101 284 927	NO	NO	1 326 363 015	4 010 341	552 195	91 446 049	NO	NO
Total	1 593 700 804	19 273 307	6 942 723	101 284 927	NO	NO	1 326 363 015	4 010 341	552 195	91 446 049	NO	NO

Table A6.5 (b). Summary information on replacement.

	Requirement for Replacement – Unit type		Replacement – Unit type					
	tCERs	ICERs	AAUs	ERUs	RMUs	CERs	tCERs	ICERs
Previous CPs			NO	NO	NO	NO	NO	NO
Year 1 (2008)		NO	NO	NO	NO	NO	NO	NO
Year 2 (2009)		NO	NO	NO	NO	NO	NO	NO
Year 3 (2010)		NO	NO	NO	NO	NO	NO	NO
Year 4 (2011)		NO	NO	NO	NO	NO	NO	NO
Year 5 (2012)	NO	NO	NO	NO	NO	NO	NO	NO
Year 6 (2013)	NO	NO	NO	NO	NO	NO	NO	NO
Year 7 (2014)	NO	NO	NO	NO	NO	NO	NO	NO
Year 8 (2015)	NO	NO	NO	NO	NO	NO	NO	NO
Total	NO	NO	NO	NO	NO	NO	NO	NO

Table A6.5 (c). Summary information on retirement.

Year	Retirement – Unit type						
	AAUs	ERUs	RMUs	CERs	tCERs	ICERs	
Year 1 (2008)		NO	NO	NO	NO	NO	NO
Year 2 (2009)	26 171 207		NO	NO	375 230	NO	NO
Year 3 (2010)	25 322 171		NO	NO	162 743	NO	NO
Year 4 (2011)	24 446 840	1766		NO	822 623	NO	NO
Year 5 (2012)	55 297 022	2 282 752	288 245	1 197 479		NO	NO
Year 6 (2013)		NO	NO	NO	NO	NO	NO
Year 7 (2014)		NO	NO	NO	NO	NO	NO
Year 8 (2015)		NO	NO	NO	NO	NO	NO
Total	131 237 240	2 284 518	288 245	2 558 075		NO	NO

Table A.6.6 List of discrepancies.										
DES Response Code	Average number of occurrences per transaction (x 100.000)		Transaction Number	Proposal Date Time	Transaction Type	Final State	Explanation	Units Involved abbreviated		
	Reported Year	Prior to the Reported Year						Serial Number	Unit Type	Quantity

Annex 7 - Tables 6.1 and 6.2 of the IPCC good practice guidance

IPCC Source category	Gas	Base year emission	Latest year emission	Activity data uncertainty	Emission factor uncertainty	Combined uncertainty	Combined uncertainty as % of total national emissions in year t	Type A sensitivity	Type B sensitivity	Uncertainty in trend in national emissions introduced by emis- sion factor uncertainty	Uncertainty in trend in national emissions introduced by activity data uncertainty	Uncertainty introduced into the trend in total national emissions
		Input data Gg CO ₂ eqv	Input data Gg CO ₂ eqv	Input data %	Input data %	%	%	%	%	%	%	%
Stationary Combustion. Coal	CO ₂	23 834	10 005	1	1	1.030	0.203	-0.085	0.135	-0.042	0.172	0.177
Stationary Combustion. BKB	CO ₂	11	1	3	5	5.831	0.0002	0.000	0.000	0.000	0.000	0.000
Stationary Combustion. Coke	CO ₂	138	74	2	5	5.301	0.008	0.000	0.001	-0.001	0.002	0.003
Stationary Combustion. Fossil waste	CO ₂	573	1397	5	10	11.180	0.307	0.014	0.019	0.135	0.133	0.190
Stationary Combustion. Petroleum coke	CO ₂	415	628	2	5	5.385	0.067	0.005	0.008	0.023	0.024	0.033
Stationary Combustion. Residual oil	CO ₂	2496	571	1	2	2.334	0.026	-0.015	0.008	-0.031	0.013	0.033
Stationary Combustion. Gas oil	CO ₂	4547	742	2	4	4.587	0.067	-0.032	0.010	-0.128	0.032	0.132
Stationary Combustion. Kerosene	CO ₂	366	2	2	5	5.393	0.0002	-0.003	0.000	-0.017	0.000	0.017
Stationary Combustion. LPG	CO ₂	184	91	2	5	5.317	0.010	0.000	0.001	-0.002	0.003	0.004
Stationary Combustion. Refinery gas	CO ₂	816	906	1	2	2.236	0.040	0.005	0.012	0.009	0.017	0.020
Stationary Combustion. Natural gas	CO ₂	4335	8293	1	0.4	1.118	0.182	0.072	0.112	0.029	0.165	0.167
Stationary Combustion. SOLID	CH ₄	13	3	1	100	100.005	0.006	0.000	0.000	-0.008	0.000	0.008
Stationary Combustion. LIQUID	CH ₄	3	1	1	100	100.003	0.002	0.000	0.000	-0.001	0.000	0.001
Stationary Combustion. GAS	CH ₄	3	6	1	100	100.005	0.011	0.000	0.000	0.005	0.000	0.005
Natural gas fuelled engines. GAS	CH ₄	5	120	1	2	2.236	0.005	0.002	0.002	0.003	0.002	0.004
Stationary Combustion. WASTE	CH ₄	1	2	5	100	100.125	0.003	0.000	0.000	0.001	0.000	0.001
Stationary Combustion. BIOMASS	CH ₄	102	135	16	100	101.233	0.269	0.001	0.002	0.088	0.040	0.097
Biogas fuelled engines. BIOMASS	CH ₄	1	31	4	10	10.728	0.007	0.000	0.000	0.004	0.002	0.005
Stationary Combustion. SOLID	N ₂ O	68	27	1	400	400.001	0.216	0.000	0.000	-0.103	0.000	0.103
Stationary Combustion. LIQUID	N ₂ O	44	11	1	1000	1000.000	0.218	0.000	0.000	-0.257	0.000	0.257
Stationary Combustion. GAS	N ₂ O	16	24	1	750	750.001	0.361	0.000	0.000	0.134	0.000	0.134
Stationary Combustion. WASTE	N ₂ O	7	16	5	400	400.031	0.124	0.000	0.000	0.060	0.001	0.060
Stationary Combustion. BIOMASS	N ₂ O	38	91	2	1000	1000.002	1.789	0.001	0.001	0.875	0.004	0.875
Transport. Road transport	CO ₂	9284	11 224	2	5	5.385	1.190	0.066	0.151	0.328	0.428	0.539
Transport. Military	CO ₂	119	116	2	5	5.385	0.012	0.000	0.002	0.002	0.004	0.005
Transport. Railways	CO ₂	297	249	2	5	5.385	0.026	0.001	0.003	0.003	0.010	0.010
Transport. Navigation (small boats)	CO ₂	48	99	41	5	41.304	0.080	0.001	0.001	0.004	0.077	0.077
Transport. Navigation (large vessels)	CO ₂	748	399	11	5	12.083	0.095	-0.002	0.005	-0.008	0.084	0.084
Transport. Fisheries	CO ₂	591	479	2	5	5.385	0.051	0.001	0.006	0.005	0.018	0.019

IPCC Source category	Gas	Base year emission	Latest year emission	Activity data uncertainty	Emission factor uncertainty	Combined uncertainty	Combined uncertainty as % of total national emissions in year t	Type A sensitivity	Type B sensitivity	Uncertainty in trend in national emissions introduced by emis- sion factor uncertainty	Uncertainty in trend in national emissions introduced by activity data uncertainty	Uncertainty introduced into the trend in total national emissions
		Input data Gg CO ₂ eqv	Input data Gg CO ₂ eqv	Input data %	Input data %	%	%	%	%	%	%	%
Transport. Agriculture	CO ₂	1272	1343	24	5	24.515	0.648	0.006	0.018	0.032	0.614	0.615
Transport. Forestry	CO ₂	36	17	30	5	30.414	0.010	0.000	0.000	-0.001	0.010	0.010
Transport. Industry (mobile)	CO ₂	839	1021	41	5	41.304	0.830	0.006	0.014	0.030	0.798	0.798
Transport. Residential	CO ₂	39	62	35	5	35.355	0.043	0.000	0.001	0.002	0.042	0.042
Transport. Commercial/institutional	CO ₂	74	171	35	5	35.355	0.119	0.002	0.002	0.008	0.114	0.115
Transport. Civil aviation	CO ₂	243	133	10	5	11.180	0.029	0.000	0.002	-0.002	0.025	0.025
Transport. Road transport	CH ₄	47	11	2	40	40.050	0.009	0.000	0.000	-0.011	0.000	0.011
Transport. Military	CH ₄	0.1	0.1	2	100	100.020	0.0001	0.000	0.000	0.000	0.000	0.000
Transport. Railways	CH ₄	0.3	0.2	2	100	100.020	0.0003	0.000	0.000	0.000	0.000	0.000
Transport. Navigation (small boats)	CH ₄	0.3	1	41	100	108.079	0.001	0.000	0.000	0.000	0.000	0.001
Transport. Navigation (large vessels)	CH ₄	0.3	0.2	11	100	100.603	0.0004	0.000	0.000	0.000	0.000	0.000
Transport. Fisheries	CH ₄	0.3	0.2	2	100	100.020	0.000	0.000	0.000	0.000	0.000	0.000
Transport. Agriculture	CH ₄	2	2	24	100	102.840	0.004	0.000	0.000	0.001	0.001	0.001
Transport. Forestry	CH ₄	0.4	0.05	30	100	104.403	0.0001	0.000	0.000	0.000	0.000	0.000
Transport. Industry (mobile)	CH ₄	1	1	41	100	108.079	0.002	0.000	0.000	0.000	0.001	0.001
Transport. Residential	CH ₄	1	1	35	100	105.948	0.003	0.000	0.000	0.001	0.001	0.001
Transport. Commercial/institutional	CH ₄	2	3	35	100	105.948	0.007	0.000	0.000	0.002	0.002	0.003
Transport. Civil aviation	CH ₄	0.1	0.05	10	100	100.499	0.0001	0.000	0.000	0.000	0.000	0.000
Transport. Road transport	N ₂ O	91	116	2	50	50.040	0.115	0.001	0.002	0.037	0.004	0.037
Transport. Military	N ₂ O	1	1	2	1000	1000.002	0.025	0.000	0.000	0.006	0.000	0.006
Transport. Railways	N ₂ O	3	2	2	1000	1000.002	0.042	0.000	0.000	0.005	0.000	0.005
Transport. Navigation (small boats)	N ₂ O	0.4	1	41	1000	1000.840	0.021	0.000	0.000	0.011	0.001	0.011
Transport. Navigation (large vessels)	N ₂ O	15	8	11	1000	1000.060	0.153	0.000	0.000	-0.030	0.002	0.030
Transport. Fisheries	N ₂ O	11	9	2	1000	1000.002	0.185	0.000	0.000	0.020	0.000	0.020
Transport. Agriculture	N ₂ O	15	18	24	1000	1000.288	0.349	0.000	0.000	0.098	0.008	0.098
Transport. Forestry	N ₂ O	0.2	0.2	30	1000	1000.450	0.003	0.000	0.000	0.001	0.000	0.001
Transport. Industry (mobile)	N ₂ O	11	13	41	1000	1000.840	0.265	0.000	0.000	0.083	0.010	0.084
Transport. Residential	N ₂ O	0.2	0.3	35	1000	1000.612	0.007	0.000	0.000	0.003	0.000	0.003
Transport. Commercial/institutional	N ₂ O	0.3	1	35	1000	1000.612	0.016	0.000	0.000	0.008	0.001	0.008

IPCC Source category	Gas	Base year emission	Latest year emission	Activity data uncertainty	Emission factor uncertainty	Combined uncertainty	Combined uncertainty as % of total national emissions in year t	Type A sensitivity	Type B sensitivity	Uncertainty in trend in national emissions introduced by emis- sion factor uncertainty	Uncertainty in trend in national emissions introduced by activity data uncertainty	Uncertainty introduced into the trend in total national emissions
		Input data Gg CO ₂ eqv	Input data Gg CO ₂ eqv	Input data %	Input data %	%	%	%	%	%	%	%
Transport. Civil aviation	N ₂ O	3	2	10	1000	1000.050	0.041	0.000	0.000	-0.002	0.000	0.002
1.B.2 Flaring in refinery	CO ₂	23	22	11	2	11.180	0.005	0.000	0.000	0.000	0.005	0.005
1.B.2 Flaring off-shore	CO ₂	302	195	8	2	7.762	0.030	0.000	0.003	0.000	0.028	0.028
1.B.2 Land based activities	CO ₂	0.003	0.01	2	40	40.050	4.2E-06	0.000	0.000	0.000	0.000	0.000
1.B.2 Off-shore activities	CO ₂	2	4	2	30	30.067	0.002	0.000	0.000	0.001	0.000	0.001
1.B.2 Transmission of natural gas	CO ₂	0.003	0.0005	15	2	15.133	1.4E-07	0.000	0.000	0.000	0.000	0.000
1.B.2 Distribution of natural gas	CO ₂	0.005	0.005	25	10	26.926	2.5E-06	0.000	0.000	0.000	0.000	0.000
1.B.2 Venting in gas storage	CO ₂	0.001	0.001	15	2	15.133	3.7E-07	0.000	0.000	0.000	0.000	0.000
1.B.2. Flaring in refinery	CH ₄	0.2	0.2	11	15	18.601	0.00006	0.000	0.000	0.000	0.000	0.000
1.B.2. Flaring off-shore	CH ₄	1	0.3	8	125	125.225	0.001	0.000	0.000	0.000	0.000	0.000
1.B.2 Refinery processes	CH ₄	1	47	1	125	125.004	0.115	0.001	0.001	0.078	0.001	0.078
1.B.2 Land based activities	CH ₄	17	18	2	40	40.050	0.014	0.000	0.000	0.003	0.001	0.003
1.B.2 Off-shore activities	CH ₄	15	37	2	30	30.067	0.022	0.000	0.000	0.011	0.001	0.011
1.B.2 Transmission of natural gas	CH ₄	4	0.3	15	2	15.133	0.000	0.000	0.000	0.000	0.000	0.000
1.B.2 Distribution of natural gas	CH ₄	5	3	25	10	26.926	0.002	0.000	0.000	0.000	0.001	0.001
1.B.2 Venting in gas storage	CH ₄	1	1	15	2	15.133	0.0004	0.000	0.000	0.000	0.000	0.000
1.B.2 Flaring in refinery	N ₂ O	0.1	0.1	11	1000	1000.060	0.001	0.000	0.000	0.000	0.000	0.000
1.B.2 Flaring off-shore	N ₂ O	1	0.5	8	1000	1000.028	0.009	0.000	0.000	0.000	0.000	0.000
2A1 Cement production	CO ₂	882	871	1	2	2.236	0.038	0.004	0.012	0.007	0.017	0.018
2A2 Lime production	CO ₂	116	40	5	5	7.071	0.006	-0.001	0.001	-0.003	0.004	0.005
2A3 Limestone and dolomite use	CO ₂	14	26	5	5	7.071	0.004	0.000	0.000	0.001	0.002	0.003
2A5 Asphalt roofing	CO ₂	0.02	0.02	5	25	25.495	0.00001	0.000	0.000	0.000	0.000	0.000
2A6 Road paving with asphalt	CO ₂	2	2	5	25	25.495	0.001	0.000	0.000	0.000	0.000	0.000
2A7a Glass and Glass wool	CO ₂	17	10	5	2	5.385	0.001	0.000	0.000	0.000	0.001	0.001
2A7b Yellow bricks	CO ₂	23	17	5	2	5.385	0.002	0.000	0.000	0.000	0.002	0.002
2A7c Expanded clay	CO ₂	15	6	5	2	5.385	0.001	0.000	0.000	0.000	0.001	0.001
2B5 Catalysts/Fertilizers. Pesticides and Sulphuric acid	CO ₂	1	1	5	5	7.071	0.0002	0.000	0.000	0.000	0.000	0.000
2C1 Iron and steel production	CO ₂	28	0	5	5	7.071	0.0E+00	0.000	0.000	-0.001	0.000	0.001

IPCC Source category	Gas	Base year emission	Latest year emission	Activity data uncertainty	Emission factor uncertainty	Combined uncertainty	Combined uncertainty as % of total national emissions in year t	Type A sensitivity	Type B sensitivity	Uncertainty in trend in national emissions introduced by emis- sion factor uncertainty	Uncertainty in trend in national emissions introduced by activity data uncertainty	Uncertainty introduced into the trend in total national emissions
		Input data Gg CO ₂ eqv	Input data Gg CO ₂ eqv	Input data %	Input data %	%	%	%	%	%	%	%
2D2 Food and Drink	CO ₂	4	2	5	5	7.071	0.0003	0.000	0.000	0.000	0.000	0.000
2G Lubricants	CO ₂	50	32	2	5	5.385	0.003	0.000	0.000	0.000	0.001	0.001
2B2 Nitric acid production	N ₂ O	1043	0	2	25	25.080	0.000	-0.010	0.000	-0.240	0.000	0.240
2F Consumption of HFC	HFC	218	657	10	50	50.990	0.660	0.007	0.009	0.342	0.125	0.365
2F Consumption of PFC	PFC	1	9	10	50	50.990	0.009	0.000	0.000	0.006	0.002	0.006
2F Consumption of SF ₆	SF ₆	107	118	10	50	50.990	0.118	0.001	0.002	0.030	0.022	0.037
3A Paint application	CO ₂	13	7	10	15	18.028	0.003	0.000	0.000	0.000	0.001	0.001
3B Degreasing and dry cleaning	CO ₂	0.00004	0.000001	10	15	18.028	5.2E-10	0.000	0.000	0.000	0.000	0.000
3C Chemical products. manufacturing and processing	CO ₂	19	12	10	15	18.028	0.004	0.000	0.000	0.000	0.002	0.002
3D5 Other	CO ₂	61	44	10	20	22.361	0.019	0.000	0.001	0.001	0.008	0.008
3D5 Consumption of fireworks	CO ₂	0.1	0.2	8	100	100.319	0.0003	0.000	0.000	0.000	0.000	0.000
3D5 Use of candles	CO ₂	22	81	10	20	22.361	0.036	0.001	0.001	0.018	0.015	0.024
3D1 Other - Use of N ₂ O for Anaesthesia	N ₂ O	0	9	5	5	7.071	0.001	0.000	0.000	0.001	0.001	0.001
3D5 Use of tobacco	N ₂ O	0.3	0.2	20	30	36.056	0.0001	0.000	0.000	0.000	0.000	0.000
3D5 Use of charcoal for BBQ	N ₂ O	0.1	0.1	10	100	100.499	0.0003	0.000	0.000	0.000	0.000	0.000
3D5 Consumption of fireworks	N ₂ O	1	2	8	100	100.319	0.004	0.000	0.000	0.002	0.000	0.002
3D5 Use of candles	N ₂ O	0.1	0.2	10	20	22.361	0.0001	0.000	0.000	0.000	0.000	0.000
4A Enteric Fermentation	CH ₄	3247	2904	2	20	20.100	1.149	0.009	0.039	0.184	0.111	0.214
4B Manure Management	CH ₄	985	1297	5	20	20.616	0.526	0.008	0.017	0.168	0.124	0.208
4F Field burning of agricultural residues	CH ₄	2	2	25	50	55.902	0.003	0.000	0.000	0.001	0.001	0.001
4.B Manure Management	N ₂ O	600	391	22	100	102.470	0.788	0.000	0.005	-0.027	0.166	0.169
4.D1.1 Synthetic Fertilizer	N ₂ O	2354	1103	25	100	103.121	2.240	-0.007	0.015	-0.683	0.529	0.864
4.D1.2 Animal waste applied to soils	N ₂ O	1112	1161	30	100	104.403	2.387	0.005	0.016	0.539	0.664	0.855
4.D1.3 N-fixing crops	N ₂ O	269	256	20	100	101.980	0.514	0.001	0.003	0.097	0.098	0.137
4.D1.4 Crop Residue	N ₂ O	361	311	20	100	101.980	0.625	0.001	0.004	0.086	0.119	0.147
4.D1.5 Cultivation of histosols	N ₂ O	290	198	20	100	101.980	0.398	0.000	0.003	0.000	0.076	0.076
4.D.2 Grassing animals	N ₂ O	334	211	25	100	103.199	0.428	0.000	0.003	-0.024	0.102	0.105
4.D3 Atmospheric deposition	N ₂ O	496	295	19	100	101.736	0.590	-0.001	0.004	-0.061	0.105	0.121

IPCC Source category	Gas	Base year emission	Latest year emission	Activity data uncertainty	Emission factor uncertainty	Combined uncertainty	Combined uncertainty as % of total national emissions in year t	Type A sensitivity	Type B sensitivity	Uncertainty in trend in national emissions introduced by emis- sion factor uncertainty	Uncertainty in trend in national emissions introduced by activity data uncertainty	Uncertainty introduced into the trend in total national emissions
		Input data Gg CO ₂ eqv	Input data Gg CO ₂ eqv	Input data %	Input data %	%	%	%	%	%	%	%
4.D3 Leaching	N ₂ O	2447	1430	20	100	101.980	2.870	-0.003	0.019	-0.330	0.545	0.637
4.D1.6 Sewage sludge and Industrial waste used as fertiliser	N ₂ O	28	39	20	100	101.980	0.078	0.000	0.001	0.026	0.015	0.030
4.F Field Burning of Agricultural Residues	N ₂ O	1	1	25	50	55.902	0.001	0.000	0.000	0.000	0.000	0.001
5.A.1 Forest remaining forest	CO ₂	50	-4491	15	2	15.133	-1.338	-0.061	-0.061	-0.122	-1.284	1.289
5.A.2 Land converted to forest	CO ₂	77	38	15	9	17.360	0.013	0.000	0.001	-0.002	0.011	0.011
5(II) Forest Land.	N ₂ O	16	12	30	10	31.623	0.008	0.000	0.000	0.000	0.007	0.007
5.B Cropland. Living biomass	CO ₂	-74	193	10	50	50.990	0.194	0.003	0.003	0.164	0.037	0.168
5.B Cropland. Dead organic matter	CO ₂	3	12	10	50	50.990	0.012	0.000	0.000	0.007	0.002	0.007
5.B Cropland. Mineral soils	CO ₂	1415	577	10	75	75.664	0.860	-0.005	0.008	-0.396	0.110	0.411
5.B Cropland. Organic soils	CO ₂	2887	1981	10	90	90.554	3.531	0.000	0.027	0.006	0.377	0.377
5(III) Disturbance. Land converted to cropland	N ₂ O	0.3	1	50	75	90.139	0.0018	0.000	0.000	0.001	0.001	0.001
5.C Grassland. Living biomass	CO ₂	75	463	10	50	50.990	0.465	0.006	0.006	0.277	0.088	0.291
5.C Grassland. Dead organic matter	CO ₂	2	7	10	50	50.990	0.007	0.000	0.000	0.004	0.001	0.004
5.C Grassland. Mineral soils	CO ₂	0.2	5	10	75	75.664	0.007	0.000	0.000	0.004	0.001	0.005
5.C Grassland. Organic soils	CO ₂	107	79	10	90	90.554	0.140	0.000	0.001	0.007	0.015	0.017
5.D Wetlands. Living biomass	CO ₂	3	-0.2	10	50	50.990	0.000	0.000	0.000	-0.001	0.000	0.001
5.D Wetlands. Dead organic matter	CO ₂	1	0	10	100	100.499	0.000	0.000	0.000	0.000	0.000	0.000
5.D Wetlands. Soils	CO ₂	85	2	10	100	100.499	0.005	-0.001	0.000	-0.075	0.000	0.075
5(II) Wetlands	N ₂ O	0.1	0.1	10	100	100.499	0.000	0.000	0.000	0.000	0.000	0.000
5.E Settlements. Living biomass	CO ₂	11	56	10	50	50.990	0.056	0.001	0.001	0.032	0.011	0.034
5.E Settlements. Dead organic matter	CO ₂	1	0.03	10	50	50.990	0.000	0.000	0.000	0.000	0.000	0.000
5.E Settlements. Soils	CO ₂	1	35	10	50	50.990	0.035	0.000	0.000	0.023	0.007	0.024
5(IV) Cropland Limestone	CO ₂	623	192	5	50	50.249	0.190	-0.003	0.003	-0.158	0.018	0.159
5(V) Biomass Burning	CH ₄	1	0.02	50	30	58.310	0.000	0.000	0.000	0.000	0.000	0.000
5(V) Biomass Burning	N ₂ O	0.4	0.03	50	30	58.310	0.00004	0.000	0.000	0.000	0.000	0.000
6 A. Solid Waste Disposal on Land	CH ₄	1366	698	10	118	118.323	1.626	-0.003	0.009	-0.376	0.133	0.399
6 B. Wastewater Handling	CH ₄	65	74	24	32	39.678	0.058	0.000	0.001	0.012	0.034	0.036
6 B. Wastewater Handling - Direct	N ₂ O	23	41	22	50	54.145	0.043	0.000	0.001	0.017	0.017	0.024

IPCC Source category	Gas	Base year emission	Latest year emission	Activity data uncertainty	Emission factor uncertainty	Combined uncertainty	Combined uncertainty as % of total national emissions in year t	Type A sensitivity	Type B sensitivity	Uncertainty in trend in national emissions introduced by emis- sion factor uncertainty	Uncertainty in trend in national emissions introduced by activity data uncertainty	Uncertainty introduced into the trend in total national emissions
		Input data Gg CO ₂ eqv	Input data Gg CO ₂ eqv	Input data %	Input data %	%	%	%	%	%	%	%
6 B. Wastewater Handling - Indirect	N ₂ O	82	32	59	42	72.558	0.046	0.000	0.000	-0.014	0.036	0.039
6.D Accidental fires. buildings	CO ₂	11	11	10	300	300.167	0.064	0.000	0.000	0.012	0.002	0.012
6.D Accidental fires. vehicles	CO ₂	6	6	10	500	500.100	0.055	0.000	0.000	0.009	0.001	0.009
6.C Incineration of corpses	CH ₄	0.01	0.01	1	150	150.003	0.00003	0.000	0.000	0.000	0.000	0.000
6.C Incineration of carcasses	CH ₄	0.001	0.005	40	150	155.242	0.00001	0.000	0.000	0.000	0.000	0.000
6.D Compost production	CH ₄	29	90	40	100	107.703	0.192	0.001	0.001	0.095	0.069	0.117
6.D Accidental fires. buildings	CH ₄	1	1	10	500	500.100	0.013	0.000	0.000	0.003	0.000	0.003
6.D Accidental fires. vehicles	CH ₄	0.3	0.2	10	700	700.071	0.003	0.000	0.000	0.001	0.000	0.001
6.C Incineration of corpses	N ₂ O	0.2	0.2	1	150	150.003	0.001	0.000	0.000	0.000	0.000	0.000
6.C Incineration of carcasses	N ₂ O	0.0	0.1	40	150	155.242	0.0003	0.000	0.000	0.000	0.000	0.000
6.D Compost production	N ₂ O	13	127	40	100	107.703	0.269	0.002	0.002	0.159	0.097	0.186
Total		74 225	50 799				47.587					6.949
Total uncertainties		Overall uncertainty in the year (%):				6.898	Trend uncertainty (%):				2.636	

Annex 8 - Methodology applied for the greenhouse gas inventory for the Faroe Islands

Introduction

This report covers the Faroese part of the National Inventory Report for the Kingdom of Denmark.

The report is made by Umhvørvisstovan, the Faroese Environment Agency (FEA).

Background information on greenhouse gas inventories and climate change

Each year the Faroe Islands is obligated to report its emission of greenhouse gases (GHG), according to the requirements of the United Nations Framework Convention on Climate Change (UNFCCC). The Kingdom of Denmark (which includes Denmark, Greenland and the Faroe Islands as geographical areas) has signed the UNFCCC. The Faroese emission figures are part of the emission total for the Kingdom of Denmark.

When Denmark ratified the Kyoto Protocol, it was with territorial reservation for the Faroe Islands. Since the reservation has not been lifted, the requirements for reporting are only those related to the Convention.

The first emission inventories for the Faroe Islands were made using an average method based upon the total use of fossil fuels in the Faroe Islands and consequently the inventories have only included total estimates of CO₂ emissions. Later, the inventories were done according to IPCC guidelines. The FEA has since 2008 yearly reported GHG emissions to Danish Centre for Environment and Energy (DCE), Dep. of Environmental Science (ENVS).

The GHGs reported are:

- Carbon dioxide CO₂
- Methane CH₄
- Nitrous Oxide N₂O
- Hydrofluorocarbons HFCs
- Perfluorocarbons PFCs
- Sulphur hexafluoride SF₆

A description of the institutional arrangement for inventory preparation

FEA, an agency under the Ministry of Fisheries, is responsible for the annual preparation and submission to the UNFCCC of the Faroe Islands' contribution to the Kingdom of Denmark's National Inventory Report and the GHG inventories in the Common Reporting Format in accordance with the UNFCCC Guidelines. The inventory is done with guidance from and in co-operation with DCE.

The work concerning the annual greenhouse gas emission inventory is carried out in co-operation with other Faroese ministries, research institutes, organisations and companies:

- *Statistics Faroe Islands (Ministry of Finance)* Annual statistics on liquid fuel sale, fuel usage for electricity and heat production, and statistics on livestock (sheep and cows).
- *Municipal Waste Plants* Data on amount of incinerated waste.
- *Electricity producing company* Data on import of F-gases (SF₆).
- *Airline Company* Data for fuel bunkers for domestic flights and international flights to and from the Faroe Islands.
- *Refrigeration companies* Data on import of F-gases (HFCs).
- *Oil companies – license holders* Data on use of fuel oil in connection with exploration (deep water) drilling in Faroese territorial waters.

In January 2010, DCE and FEA made a formal agreement about data delivery.

Brief description of the process of inventory preparation. Data collection and processing, data storage and archiving

The activity data for fuel sale and for fuel usage by combustion plants, as well as for the number of livestock (sheep and cows) are collected and stored at Statistics Faroe Islands. Each year, FEA receives new data for fuel sale and fuel usage for the previous year. Numbers of livestock is accessible on the homepage of Statistics Faroe Islands, www.hagstova.fo

Other activity data are delivered by plants owned by municipalities or private companies.

After receiving the data, the material is placed on servers at FEA. The servers are subject to routine backup services. Material that has been backed up is archived safely. All collected data is also archived in the electronic journal of the agency.

The emission factors are yearly received from DCE Denmark, sent by email to the FEA as Excel files. In addition to copying the factors to spread sheet files, the e-mails are archived in the electronic journal.

Since the 2008 submission, all subsequent submissions have been reported in the Common Reporting Format of UNFCCC (CRF). The new format has meant improvements, higher data security and limited the potential for errors in the reporting. The emission inventory is both reported in the form of an xml file and as CRF Excel tables.

Brief general description of methodologies and data sources used

The GHG inventory for the Faroe Islands includes the following sectors:

- Energy sector (CRF sector 1)
- Industrial processes (CRF sector 2)
- Agriculture (CRF sector 4)
- Waste (CRF sector 6)

Since the emissions in the Waste sector all are allocated to the Energy sector, table 1 also includes methods applied and emission factors for calculating GHG emissions related to the Waste sector.

The applied methodologies follow the IPCC Guidelines and IPCC Good Practice Guidance, and the Tier 1 method is always applied.

The methods and the emission factors used in the inventory are shown in Table 1 (emission factors for CO₂, CH₄ and N₂O in the Energy and Agriculture sector) and in Table 2 (emission factors for HFCs and SF₆ in the sector for Industrial Processes). A brief general description of methodologies is included below for the different sectors.

Table 1 Methods applied and emission factors used for calculating CO₂, CH₄ and N₂O emissions in the Energy and Agriculture sectors.

GHG CATEGORIES	CO ₂		CH ₄		N ₂ O	
	Method applied	Emission factor	Method applied	Emission factor	Method applied	Emission factor
1. Energy	T1	CS	T1	CS	T1	CS
A. Fuel Combustion	T1	CS	T1	CS	T1	CS
1. Energy Industries	T1	CS	T1	CS	T1	CS
2. Manufacturing Industries and Construction	T1	CS	T1	CS	T1	CS
3. Transport	T1	CS	T1	CS	T1	CS
4. Other Sectors	T1	CS	T1	CS	T1	CS
4. Agriculture			T1	D	T1	D
A. Enteric Fermentation			T1	D		
B. Manure Management			T1	D	T1	D

Table 2 Methods and Emission factors used for calculating HFCs and SF₆ emissions in the Industrial Processes sector.

GHG CATEGORIES	HFCs		SF ₆	
	Method applied	Emission factor	Method applied	Emission factor
2. Industrial Processes	T1	D	T1	D
F. Consumption of Halocarbons and SF ₆	T1	D	T1	D

Energy sector

All emissions in the Energy sector are from Fuel combustion (1.A.1), and in these categories:

- 1A1a Public Electricity and Heat Production (incl. Waste)
- 1A1c Manufacture of Solid fuels and Other Energy Industries
- 1A2 Manufacturing Industry and Construction
- 1A3a Civil Aviation
- 1A3b Road Transportation
- 1A3d Navigation
- 1A4a Commercial/Institutional
- 1A4b Residential
- 1A4c Agriculture, Forestry and Fishing

Statistics Faroe Islands provides the information on fuel sales by fuel type (in m³) and divided into eight main groups (original titles: Fishing vessels, Other ships, Transportation, Industry, Trading and Service, Residential and Communities, Institutions and Public power), each group again divided into subgroups.

The fuel data delivered by Statistics Faroe Islands originate from several sources. The main data sources are the two main oil companies in the Faroe Islands. Fuel data not included in sales information from the oil companies are delivered by the industry to FEA.

Since the delivered data on fuel sale are not fully arranged according to IPCC guidelines, the FEA rearranges the data to comply with the guidelines.

Emission factors

Emissions from fuel combustion can be divided into two main sources: stationary and mobile combustion. Stationary combustion means fuel combustion related to e.g. industry on land, house heating and oil exploration. Mobile combustion includes the combustion in engines used for propulsion in the various modes of transport such as road transport, marine activities and aviation. The emission factors used for stationary, transport, waste and aviation are country specific and provided by DCE. All emissions factors used in the inventory are found in Annex 2 and 3.

Emissions are calculated by multiplying fuel consumption data with an emission factor (e.g. in tonnes emission per GJ fuel).

Public Electricity and Heat Production (1A1a)

The activity data used for calculations of emissions of GHG from for Public Electricity and Heat Production are data for usage of residual oil and diesel oil at electricity producing plant on the Faroe Islands. The emission factors are calculated and delivered by DCE, see Table 10 in Annex 2.

Manufacture of Solid fuels and Other Energy Industries (1A1c)

This category only covers the emissions of GHG from activity related to exploration drilling in Faroese territory. The activity data (usage of diesel on the rigs) are delivered by the operators. The emission factors are calculated and delivered by DCE, see Table 10 in Annex 2.

Manufacturing Industry and Construction (1A2)

The activity data for oil usage are delivered by Statistics Faroe Islands. The emission factors are calculated and delivered by DCE, see Table 10 in Annex 2.

Civil aviation (1A3a)

The Faroese airline company, Atlantic Airways, delivers data for jet fuel bunkered in the Faroe Islands. As the Faroe Islands has accepted the United Nations Framework Convention on Climate Change as a part of the Kingdom of Denmark, aviation between Denmark and the Faroe Islands is to be reported as domestic aviation. The data is thus divided by destination: flights to destinations inside the Kingdom of Denmark, i.e., Denmark and Greenland (Domestic Aviation), and outside the Danish Kingdom, e.g., Iceland, Norway and Great Britain (International Aviation). Fuel refuelled outside the Faroe Islands is not included in the Faroese inventory.

The emission factors for aviation are made by DCE, see Table 12 in Annex 3.

Road transport (1A3b)

The activity data for road transport is data for sale of gasoline and diesel to all types of vehicle at all filling stations in the Faroe Islands. The data is delivered by the Statistics Faroe Islands. The emission factors for road traffic are calculated by DCE. The Danish results are modified for Faroese traffic conditions such as other gross vehicle weights for heavy-duty vehicles and no highway driving conditions. The emissions factors are also modified because biofuel is not used in the Faroe Islands, unlike in Denmark. The emission factors are shown in Table 12 in Annex 3.

Navigation (1A3d)

The activity data for oil usage used in navigation are delivered by Statistics Faroe Islands. The emission factors are calculated and delivered by DCE, see Table 13 in Annex 3.

Commercial and Institutional (1A4a) and Residential (1A4b)

The activity data for oil usage used to calculate the GHG emissions from the Commercial and Institutional and Residential categories are delivered by Statistics Faroe Islands. The emission factors are calculated and delivered by DCE, and found in Table 10 in Annex 2

Agriculture, Forestry and Fishing (1A4c)

Very little fuel consumption is related to Agriculture and Forestry on the Faroe Islands. The fuel used in most probably bought on filling stations and therefore allocated to road transport. Therefore all emissions in this category are related to Fishing. The activity data (sale of oil to fishing vessels) is delivered by Statistics Faroe Islands. The emission factors are calculated and delivered by DCE, and found in Annex 3.

Until this year's delivery of data, in some few cases, it has not been possible to rearrange the data from Statistics Faroe Islands to fully comply with the IPCC guidelines. This was the case for foreign fishing vessels. According to the guidelines all emissions resulting from fuel used in coastal and deep sea fishing should be allocated to the country delivering the fuel. When oil is sold to foreign vessels, the oil companies do not always, or have not always, registered whether the ship is a fishing vessel or another type of vessel. Even though most foreign vessels today bunkering in the Faroe Islands are fishing vessels, the emission from foreign vessels have until now and for all years been allocated to International Bunkers. This means that for all previous data deliveries, the emission from fishing vessels in reality were higher than in the inventory and emission from International bunkering were lower. This has been change in the last delivery. Through direct communication with the oil companies, the Environmental Agency has received more detailed information about sale of oil to foreign fishing vessels, enough to make a fairly good estimation of the amount of oil sold to foreign fishing vessels in the years 2001-2011. . This has resulted in higher emissions from fishing vessels and lower emissions in International Bunkers for the year 2001-2011. The same new estimations for the years 1990-2000 remains to be done.

The inventory includes all oil bunkered on Faroese territory, excluding oil bunkered at open sea, or on other more near-coast sites, by international companies, i.e., from foreign supplier to foreign customer.

Industrial processes

Emissions from Industrial processes are allocated to these categories:

- 2F1 Refrigeration and Air Conditioning
- 2G1 Electrical Equipment

The inventory follows the principles in the IPCC Guidelines and the IPCC Good Practice Guidance, with a Tier 1 methodology. The emissions factors are IPCC default.

The activity data origin from FEA surveys on the consumption (import) of HFCs and SF₆ which have been conducted annually since 2003. An estimate of the consumption has been done for the years 1990-2002.

There has been no consumption of PFCs in the Faroe Islands.

Solvent and other product use

Since no data are available, emissions from solvent and other product use are not calculated.

Agriculture

GHG emissions from agriculture are calculated for following categories:

- 4A Enteric fermentation
- 4B Manure management
- 4D2 Agricultural Soil - Grassing animals

The inventory follows the principles in the IPCC Guidelines and the IPCC Good Practice Guidance. Tier 1 method is always used. All emission factors used for agriculture are IPCC standard values. The emissions are calculated with support from DCE. Activity data is accessible on the homepage of Statistics Faroe Islands.

Waste

The GHG emission from waste incineration is calculated using country specific methodology. All emissions in the Waste sector have been allocated to the Energy sector. Emission factors relative to emissions of CO₂, N₂O and CH₄ from waste incineration in 1990-2012 are listed in Table 11 in Annex 2. Heating values for waste incineration are listed in Table 3 below.

Table 3 Heating values (GJ pr t) for waste.

Year	Heating values
	GJ pr t
1990-91	8,2
1992	9,0
1993-94	9,4
1995	10,0
1996-2012	10,5

Brief description of key categories

No key category analysis (KCA) has been carried out for the Faroe Islands inventory.

Information on QA/QC plan including verification and treatment of confidential issues where relevant

A number of measures are in place to ensure the quality of the greenhouse gas inventory for the Faroe Islands.

The general QC activities include:

- Check that data from Statistics Faroe Islands and other data deliverers are correctly transferred to emissions spread sheets.
- Check that data are correctly moved between data processing steps, e.g., it is ensured that the data are imported correctly from the emission spread sheets /databases to the CRF Reporter.
- The time series are analysed. Any large fluctuations are investigated and explained /corrected.
- The completeness of the inventory is checked utilising the completeness checker incorporated in the CRF Reporter.

These types of QC checks are recommended as Tier 1 QC checks in the IPCC Good Practice Guidance (IPCC, 2000).

No confidential issues are relevant.

General uncertainty evaluation, including data on the overall uncertainty for the inventory totals

No uncertainty evaluation has been made for the Faroese inventory.

General assessment of the completeness

In general, the inventory is complete.

References

Lastein, L. & Winther, M. 2003: Emissions of greenhouse gases and long-range transboundary air pollutants in the Faroe Islands 1990-2001. National Environmental Research Institute, Denmark. 62 p. NERI Technical Report no. 477. Available at:
<http://technicalreports.dmu.dk>

Winther, M. 2001: 1998 Fuel Use and Emissions for Danish IFR Flights. Environmental Project no. 628, 2001. 112 p. Danish EPA. Prepared by the National Environmental Research Institute (NERI), Denmark. Electronic report at homepage of Danish EPA. Available at :
<http://www.mst.dk/udgiv/Publications/2001/87-7944-661-2/html/>

Trends in Greenhouse Gas Emissions

The trends present in this Chapter cover the emissions from the Faroe Islands.

The emission trend tables 1990, 1995, 2000, 2005, 2010 and 2012 for CO₂, CH₄, N₂O and F-gases (CRF: Table10) and emission trend summary table 1990, 1995, 2000, 2005, 2010 and 2012 are presented in Annex 1.

Description and interpretation of emission trends for aggregated greenhouse gas emissions

The greenhouse gas emissions are estimated according to the IPCC guidelines and are aggregated into four main sectors: Energy, Agriculture,

ture, Waste and Industrial Processes. All emissions from the Waste sector are allocated to the Energy sector. The greenhouse gases include CO₂, CH₄, N₂O, HFCs and SF₆. Figure 1 shows the estimated total greenhouse gas emissions in CO₂ equivalents from 1990 to 2012. The total greenhouse gas emission in CO₂ equivalents has increased by 3.9 % from 1990 to 2012. Comments on the overall trends etc. are given in the sections below.




Figure 1 Greenhouse gas emissions in CO₂ equivalents distributed on main sectors for 2012 and time series for 1990 to 2012.

Figure 2 shows the composition of greenhouse gas emissions (CO₂, N₂O, CH₄ and F-gases) in 2012, calculated in GWP values. CO₂ is the most important greenhouse gas contributing in 2012 with 93.4 %, followed by N₂O with 2.8 %, CH₄ 2.2 % and F-gases (HFCs and SF₆) with 1.6 %.




Figure 2 Emissions of GHG in CO₂ equivalents in 2012 distributed on type of gas.

Figure 3 shows the total emissions of greenhouse gases and the emission of CO₂, N₂O, CH₄ and F-gases (in CO₂ equivalents) in the time period 1990-2012. From 1990 to 1993 a decrease is observed, due to an economic crisis in the Faroe Islands. From 2001 to 2007, the emissions were rather stable. In 2008-2011 the emissions from Faroese fishing ship were significantly lower than previous years. The decrease is concealed by emissions related to new bunkering activity starting in 2009 that has led to a substantial increase in the number of foreign fishing vessels bunkering in the Faroe Island. In 2012, the emissions were 22.8 % above the base year.




Figure 3 GHG emission in CO₂ equivalents, time series 1990-2012.

Description and interpretation of emission trends by gas

Carbon dioxide

The emission of CO₂ on the Faroe Islands is from fuel consumption only. The trend in the total emission of CO₂ (Figure 4) is nearly identical with the trend of the total emission of GHG in the Faroe Islands (Figure 3) showing the trends in CO₂ emissions in the period from 1990 to 2012. After the economic decline in the 1990s the emissions rose and were rather constant until 2007. From 2008 to 2012 the effort in the Faroese fishing fleet was significantly lower than previous years, also meaning a significant reduction in oil consumption. The reduction in the emissions for fisheries in 2009 and 2011 is not visible because a new oil bunkering activity (mostly used by foreign fishing vessels) started up in 2009, increasing the emissions.




Figure 4 Total CO₂ emissions, time series for 1990-2012.

Figure 5 shows how the emissions are distributed between categories. In 2012 42 % of the CO₂ emissions came from fishing vessels. Public electricity and heat production accounted for 16 %, households for 14 % and road transport for 11 % of the total CO₂ emission.




Figure 5 Emissions of CO₂ in the Energy sector, divided in fuel consumption categories, 2012.

Nitrous oxide

Figure 6 shows the emissions of nitrous oxide in the Faroe Islands 1990-2012. Most of the N₂O is from the agriculture sector, especially from animals grazing on agricultural soils.




Figure 6 N₂O emissions in tonnes distributed on sector and time series for 1990-2012.

Methane

Figure 7 shows the emissions of methane in the Faroe Islands 1990-2012. Most of the methane emission is from the agriculture sector, especially from enteric fermentation (87 %). Most of the emission of CH₄ in the energy sector is due to aviation activity.




Figure 7 CH₄ emissions in tonnes distributed on sectors and time series for 1990-2012.

HFCs, PFCs and SF₆

Figure 8 shows the emissions of F-gases, HFCs and SF₆ respectively in the years 1990-2012. Most of the emission is HFCs, which are used for refrigeration purposes, as substitutes for HCFCs. After the emissions increased in the period 1996-2005, the emissions were rather stable at

around 12,000 tonnes of CO₂ equivalents pr. year until 2012, where the emissions of HFC were 14,200 CO₂ equivalents. This is due to higher use of HFC-125 and HFC-143a, both components in the HFC-blend HFC-507a, which in recent years has been used as a substitute when phasing out HCFC-22 (ozone depleting freezing agent) on fishing vessels.




Figure 8 F-gas emissions in CO₂ equivalents, contribution from type of F-gas and time series for 1990-2012.

In 2012 the actual emission of SF₆ was 194 tonnes CO₂ equivalents.

PFC has never been in use in the Faroe Islands.

Description and interpretation of emission trends by source

In 2012, nearly 95 % of all GHG emissions were from the Energy sector, including waste-incineration. Almost 4 % were from Agriculture and nearly 2 % from Industrial processes, see Figure 9.




Figure 9 Emissions of GHG in CO₂ equivalents distributed by main sectors, 2012.

The fluctuations in the GHG emissions in the Energy sector are decisive for the fluctuations in the total GHG emissions, see Figure 10. The emissions from the Agriculture sector and from Industrial processes are relative small and constant.




Figure 10 GHG emissions in CO₂ equivalents, main sectors, time series 1990-2012.

Description and interpretation of emission trends for indirect greenhouse gases and SO₂

Emission trends for indirect greenhouse gases and SO₂ have not been made for the Faroe Islands.

Energy (CRF sector 1)

Overview of the sector

Fuel consumption on the Faroe Islands can be seen in Figure 11. Most of the fuel is used by fishing vessels.




Figure 11 Fuel consumption (tonnes) in the Energy sector, including waste incineration, 1990-2012.

Figure 12 shows the GHG emissions in the Energy sector on the Faroe Islands 1990-2012. The trend in Figure 12 is just the same as in Figure 11.




Figure 12 GHG emissions in CO₂ equivalents, categories in the Energy sector, 1990-2012.

Figure 13 shows how the emission of GHG in 2012 was distributed between groups of fuel users. Fishing vessels, Electricity production, Residential and Road transport had 42, 16, 14 and 11 %, respectively, of the emissions in the Energy sector in 2012.

Waste incineration has been included under sector 1A1a (Electricity and Heat production).




Figure 13 GHG emissions in CO₂ equivalents; Energy sector divided in categories, 2012.

Fugitive emissions (CRF sector 1B)

Fugitive emissions of GHG gases are estimated to be very limited on the Faroe Islands. These emissions have not been estimated.

Industrial Processes (CRF Sector 2)

There is no chemical industry, no metal production, no production of F-gases and no mineral production (other than road paving with asphalt) on the Faroe Islands. The only industrial processes leading to GHG emissions on the Faroe Islands is the use of F-gases. Since no data is available on paving roads with asphalt, the emissions of GHG from road paving are not included in the inventory.

Overview of the sector

Figure 14 shows the GHG emissions from industrial processes on the Faroe Islands. The increase in emissions, starting in 1996 is due to use of HFCs in refrigeration. See figure 8.




Figure 14 GHG emissions in CO₂ equivalents, Industrial processes, 1990-2012.

Mineral products (2A)

There is no mineral production in the Faroe Islands, other than paving roads with asphalt. No data is available for paving roads with asphalt.

Chemical industry (2B)

No chemical industry with GHG emission is located in the Faroe Islands.

Metal production (2C)

No metal production industry is located in the Faroe Islands.

Production of Halocarbons and SF₆ (2E)

There is no production of halocarbons and SF₆ in the Faroe Islands.

Metal Production (2C) and Consumption of Halocarbons and SF₆ (2F)

There is no metal production on the Faroe Islands.

Of the total GHG emissions 2 % are emissions related to consumption of halocarbons and SF₆. The major part of the emission (99%) is HFC gasses, which are used for refrigeration purposes and the rest (1 % of the emission) is SF₆ used in electrical equipment. See Figure 8

Time series of the emission (tonnes) of HFCs 1990-2012, are seen in Table 4.

Table 4 Emissions of HFCs from Refrigeration and Air Conditioning, 1990-2012 (tonnes).

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
Domestic refrigeration											
HFC-134a	NO	NO	NO	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
Commercial refrigeration											
HFC-134a	NO	NO	NO	0,00	0,00	0,00	0,00	0,01	0,02	0,03	0,05
HFC-32	NO	NO	NO	0,00	0,00	0,00	0,00	0,00	0,01	0,05	0,09
HFC-125	NO	NO	NO	0,00	0,00	0,00	0,00	0,01	0,03	0,09	0,15
HFC-143a	NO	NO	NO	0,00	0,00	0,00	0,00	0,01	0,02	0,04	0,06
Industrial refrigeration											
HFC-134a	NO	NO	NO	0,00	0,00	0,00	0,00	0,03	0,06	0,11	0,16
HFC-125	NO	NO	NO	0,00	0,00	0,00	0,00	0,07	0,12	0,23	0,33
HFC-143a	NO	NO	NO	0,00	0,00	0,00	0,00	0,08	0,15	0,28	0,39
Mobile Air Conditioning											
HFC-134a	NO	NO	NO	NO	NO	NO	NO	NO	0,01	0,70	0,70

<i>Continued</i>	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
Domestic refrigeration												
HFC-134a	0,00	0,00	0,01	0,01	0,01	0,01	0,01	0,01	0,01	0,01	0,01	0,01
Commercial refrigeration												
HFC-134a	0,07	0,10	0,12	0,13	0,14	0,13	0,13	0,14	0,14	0,17	0,20	0,19
HFC-32	0,16	0,22	0,28	0,33	0,32	0,31	0,30	0,28	0,26	0,27	0,24	0,22
HFC-125	0,25	0,35	0,43	0,50	0,51	0,50	0,50	0,57	0,59	0,74	0,81	1,29
HFC-143a	0,09	0,12	0,15	0,17	0,19	0,19	0,22	0,32	0,35	0,51	0,62	1,14
Industrial refrigeration												
HFC-134a	0,28	0,36	0,43	0,48	0,45	0,39	0,36	0,34	0,34	0,35	0,36	0,28
HFC-125	0,59	0,75	0,88	0,99	0,97	1,03	1,06	1,01	0,86	0,77	0,68	0,58
HFC-143a	0,70	0,89	1,05	1,17	1,15	1,22	1,25	1,19	1,02	0,91	0,80	0,68
Mobile Air Conditioning												
HFC-134a	0,70	0,70	0,70	0,68	0,59	0,64	0,76	0,83	0,89	0,94	0,97	1,00

The HFC emissions are reported with the following assumptions:

- Domestic refrigeration is use in freezers and refrigerators.
- Commercial refrigeration is use in landbased units.
- Industrial refrigeration is use on ships.
- Mobile air conditioning is use in cars, buses and trucks.

Figure 15 shows the emissions of SF₆ and four specific HFCs.




Figure 15 Emission of F-gases (HFCs and SF₆) in CO₂ equivalents, time series for 1990-2012.

Other (2G)

No emissions are in the category “Other”.

Uncertainty

Estimations of the uncertainties for Industrial processes have not been done.

Solvents and other product use (CRF Sector 3)

Overview of the sector

Since no data are available for this sector, no emissions are calculated. The expected emissions are low.

Agriculture (CRF Sector 4)

Overview

The emission of greenhouse gases from agricultural activities includes:

- CH₄ emission from manure management and enteric fermentation.
- N₂O emission from manure management and agricultural soil.

Nearly 4 % of the total GHG emissions on the Faroe Islands are due to agriculture. The sources are cattle and sheep. Figure 16 shows the number of cattle in the Faroe Islands from 1990 to 2012. The number of sheep is around 78,940, which is the carrying capacity for sheep on the islands. There are no data on the exact number of sheep.




Figure 16 Number of cattle (dairy and non-dairy), time series for 1990-2012.

Figure 17 shows the total emissions from the Agriculture sector.




Figure 17 GHG emissions in CO₂ equivalents, in the Agriculture sector, 1990-2012.

CH₄ emission from Enteric Fermentation (CRF Sector 4A)

Figure 18 shows emissions of CH₄ from enteric fermentation on the Faroe Islands, 1990-2012.




Figure 18 CH₄ emissions in CO₂ equivalents from enteric fermentation, 1990-2012.

CH₄ and N₂O emission from Manure Management (CRF Sector 4B)

Figure 19 shows emissions of N₂O and CH₄ from manure management on the Faroe Islands.




Figure 19 N₂O and CH₄ emission in CO₂ equivalents from Manure management, time series 1990-2012.

N₂O emission from Agricultural Soils (CRF Sector 4D)

The emission from sheep and cows grazing on agricultural soil is 44 tonnes N₂O per year. This corresponds to 13,700 tonnes of CO₂ equivalents. Figure 20 shows the N₂O emissions from agricultural soil. Since the number of sheep is more or less constant over time, the emissions are also constant.




Figure 20 N₂O emissions (tonnes) from Agricultural Soils, grazing animals, time series 1990-2012.

NMVOC emission

The emission of NMVOC is not calculated.

Uncertainties

The uncertainties have not been calculated.

Recalculation

No recalculations were made in the Agriculture sector in the 2014 submission.

Planned improvements

Include emissions from animal categories other than cattle and sheep.

Land Use, Land Use Change and Forestry (CRF Sector 5)

No emissions are calculated for land use, land-use change and forestry.

Waste Sector (CRF Sector 6)

Overview of the Waste sector

Waste incineration is the only source in the Waste sector with significant emission. The emissions have been allocated to the energy sector in accordance with the IPCC Guidelines.

Solid Waste Disposal on Land (CRF Source Category 6A)

A number of land-based solid waste disposals facilities are located on the Faroe Islands. The GHG emissions from these depots have not been calculated.

Wastewater Handling (CRF Source Category 6B)

In the Faroe Islands, most households have a septic tank (mechanical treatment). Industrial wastewater, e.g. from the fishing industry, is treated mechanically (oil/fat separation). Only a few wastewater handling plants are treating the wastewater chemically and/or biologically.

GHG emissions from wastewater handling are not calculated.

Waste Incineration (CRF Source Category 6C)

There are two waste incineration plants on the Faroe Islands, one in Hoyvík and one in Leirvík. Both plants are considered energy recovery operations and therefore the emissions have been allocated to the energy sector (1A1a) in accordance with the IPCC Guidelines.

Figure 21 shows the amounts of waste incinerated on the Faroe Islands 1990-2012.




Figure 21 Incineration of municipal waste on the Faroe Islands, 1990-2012.

Waste Other (CRF Source Category 6D)

There are no activities and emissions in Waste Other.

Other (CRF sector 7)

In CRF sector 7, there are no activities and emissions or removals for the inventory of the Faroe Islands.

Recalculations and improvements

Nearly all recalculations in the 2014 submission for the Faroe Islands are due to changes in emissions factors, and in all these cases the changes are the same as in the inventory for Denmark.

Explanations and justifications for recalculations

The following recalculations and improvements to the emission inventories have been made since the reporting in 2012.

Energy

Energy Industries

Due to updates in the emissions factors for CO₂ a minor recalculation has been made for the emissions in Public Electricity and Heat Production for the year 2012. The updates are the same as in the inventory for Denmark.

Civil aviation

The emissions factors for CH₄ for the years 1990-2001 and for N₂O for the years 2010 and 2011 have been updated due to small changes in the DCE Denmark flight model.

Industrial Processes – F-gases

Due to importers late delivery of data for imports of HFC in 2010 and in 2011 some minor additions of activity data for HFC were made in the 2014 submission. The recalculations implied only minor changes in the emission.

Implications for emission levels

The recalculations in the 2014 submission have substantial implications for emission levels, especially for the fishing category and for the International bunkers. Earlier, almost all emissions from foreign ships have been classified as International bunkers. Based on information from the oil companies, the amount of oil sold to foreign fishing vessels has now been estimated, and the activity data corrected corre-

spondently. Usually around one third of the total emission used to be from fishing vessels. After the recalculations, the fishing vessels represent around 40% of the total emissions. Since activity data only have been reallocated, the total emissions levels have not changed due to these recalculations.

Implications for emission trends, including time series consistency

Unfortunately the recalculation mentioned above could for certain reasons only be done for the time-series 2001-2011. Thus the time series for fishing vessels, 2001-2012, is inconsistent with the time series 1990-2000.

Annexes

Annex 1 Trend tables 1990, 1995, 2000, 2005, 2010 and 2012 for CO₂, CH₄, N₂O, F-gases (CO₂ equivalents) and Trend tables 1990, 1995, 2000, 2005, 2010 and 2012 for Summary (all gases)

The tables are copied from the CRF 2012 spreadsheet file, Tables 10.1-10.5.

Table 5 EMISSION TRENDS CO₂ - Inventory 2012 - Submission 2014 v1.1 - FAROE ISLANDS.

GREENHOUSE GAS SOURCE AND SINK CATEGORIES	1990	1995	2000	2005	2010	2012
	(Gg)	(Gg)	(Gg)	(Gg)	(Gg)	(Gg)
1. Energy	667,67	539,31	664,33	773,89	839,23	814,23
A. Fuel Combustion (Sectoral Approach)	667,67	539,31	664,33	773,89	839,23	814,23
1. Energy Industries	96,90	79,23	119,26	112,26	163,99	150,08
2. Manufacturing Industries and Construction	61,86	31,89	59,30	65,26	43,28	48,54
3. Transport	105,12	97,89	99,17	107,30	125,71	129,82
4. Other Sectors	403,79	330,30	386,60	489,08	506,25	485,79
5. Other	NA	NA	NA	NA	NA	NA
B. Fugitive Emissions from Fuels	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO
1. Solid Fuels	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO
2. Oil and Natural Gas	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO
2. Industrial Processes	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO
A. Mineral Products	NE,NO	NE,NO	NE,NO	NE,NO	NE,NO	NE,NO
B. Chemical Industry	NO	NO	NO	NO	NO	NO
C. Metal Production	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO
D. Other Production	NA	NA	NA	NA	NA	NA
E. Production of Halocarbons and SF ₆						
F. Consumption of Halocarbons and SF ₆						
G. Other	NA	NA	NA	NA	NA	NA
3. Solvent and Other Product Use	NA,NE	NA,NE	NA,NE	NA,NE	NA,NE	NA,NE
4. Agriculture						
5. Land Use, Land-Use Change and Forestry(2)	NA,NE	NA,NE	NA,NE	NA,NE	NA,NE	NA,NE
A. Forest Land	NE	NE	NE	NE	NE	NE
B. Cropland	NE	NE	NE	NE	NE	NE
C. Grassland	NE	NE	NE	NE	NE	NE
D. Wetlands	NE	NE	NE	NE	NE	NE
E. Settlements	NE	NE	NE	NE	NE	NE
F. Other Land	NE	NE	NE	NE	NE	NE
G. Other	NA	NA	NA	NA	NA	NA
6. Waste	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO
A. Solid Waste Disposal on Land	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO
B. Waste-water Handling						
C. Waste Incineration	NA	NA	NA	NA	NA	NA
D. Other	NA	NA	NA	NA	NA	NA
7. Other	NA	NA	NA	NA	NA	NA
Total CO ₂ emissions including net CO ₂ from LULUCF	667,67	539,31	664,33	773,89	839,23	814,23
Total CO ₂ emissions excluding net CO ₂ from LULUCF	667,67	539,31	664,33	773,89	839,23	814,23
Memo Items:						
International Bunkers	NA,NE,NO	131,72	136,46	85,37	43,25	60,87
Aviation	NE,NO	0,13	0,88	1,21	0,77	1,20
Marine	NA,NE,NO	131,59	135,59	84,17	42,48	59,67
Multilateral Operations	NO	NO	NO	NO	NO	NO

Table 6 EMISSION TRENDS CH₄ - Inventory2012 - Submission 2014 v1.1 - FAROE ISLANDS.

GREENHOUSE GAS SOURCE AND SINK CATEGORIES	Base year	1995	2000	2005	2010	2012
	(Gg)	(Gg)	(Gg)	(Gg)	(Gg)	(Gg)
1. Energy	0,05	0,05	0,06	0,13	0,10	0,08
A. Fuel Combustion (Sectoral Approach)	0,05	0,05	0,06	0,13	0,10	0,08
1. Energy Industries	0,00	0,00	0,00	0,00	0,00	0,00
2. Manufacturing Industries and Const	0,00	0,00	0,00	0,00	0,00	0,00
3. Transport	0,04	0,05	0,05	0,12	0,09	0,07
4. Other Sectors	0,01	0,01	0,01	0,01	0,01	0,01
5. Other	NA	NA	NA	NA	NA	NA
B. Fugitive Emissions from Fuels	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO
1. Solid Fuels	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO
2. Oil and Natural Gas	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO
2. Industrial Processes	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO
A. Mineral Products	NO	NO	NO	NO	NO	NO
B. Chemical Industry	NO	NO	NO	NO	NO	NO
C. Metal Production	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO
D. Other Production						
E. Production of Halocarbons and SF ₆						
F. Consumption of Halocarbons and SF ₆						
G. Other	NA	NA	NA	NA	NA	NA
3. Solvent and Other Product Use						
4. Agriculture	0,82	0,84	0,84	0,82	0,81	0,82
A. Enteric Fermentation	0,78	0,81	0,80	0,79	0,77	0,78
B. Manure Management	0,04	0,04	0,04	0,04	0,03	0,04
C. Rice Cultivation	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO
D. Agricultural Soils	NA,NE	NA,NE	NA,NE	NA,NE	NA,NE	NA,NE
E. Prescribed Burning of Savannas	NA	NA	NA	NA	NA	NA
F. Field Burning of Agricultural Residues	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO
G. Other	NA	NA	NA	NA	NA	NA
5. Land Use, Land-Use Change and Forestry	NA,NE	NA,NE	NA,NE	NA,NE	NA,NE	NA,NE
A. Forest Land	NE	NE	NE	NE	NE	NE
B. Cropland	NE	NE	NE	NE	NE	NE
C. Grassland	NE	NE	NE	NE	NE	NE
D. Wetlands	NE	NE	NE	NE	NE	NE
E. Settlements	NA,NE	NA,NE	NA,NE	NA,NE	NA,NE	NA,NE
F. Other Land	NA,NE	NA,NE	NA,NE	NA,NE	NA,NE	NA,NE
G. Other	NA	NA	NA	NA	NA	NA
6. Waste	IE,NA,NE,NO	IE,NA,NE,NO	IE,NA,NE,NO	IE,NA,NE,NO	IE,NA,NE,NO	IE,NA,NE,NO
A. Solid Waste Disposal on Land	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO
B. Waste-water Handling	NA,NE	NA,NE	NA,NE	NA,NE	NA,NE	NA,NE
C. Waste Incineration	IE,NA	IE,NA	IE,NA	IE,NA	IE,NA	IE,NA
D. Other	NA	NA	NA	NA	NA	NA
7. Other	NA	NA	NA	NA	NA	NA
Total CH₄ emissions including CH₄ from LULUCF	0,87	0,90	0,89	0,96	0,91	0,90
Total CH₄ emissions excluding CH₄ from LULUCF	0,87	0,90	0,89	0,96	0,91	0,90
Memo Items:						
International Bunkers	NA,NE,NO	0,00	0,01	0,01	0,01	0,01
Aviation	NE,NO	0,00	0,01	0,01	0,01	0,01
Marine	NA,NE,NO	0,00	0,00	0,00	0,00	0,00
Multilateral Operations	NO	NO	NO	NO	NO	NO
CO₂ Emissions from Biomass						

Table 7 EMISSION TRENDS N₂O - Inventory 2012 - Submission 2014 v1.1 - FAROE ISLANDS

GREENHOUSE GAS SOURCE AND SINK CATEGORIES	1990	1995	2000	2005	2010	2012
	(Gg)	(Gg)	(Gg)	(Gg)	(Gg)	(Gg)
1. Energy	0,02	0,02	0,02	0,03	0,03	0,03
A. Fuel Combustion (Sectoral Approach)	0,02	0,02	0,02	0,03	0,03	0,03
1. Energy Industries	0,00	0,00	0,00	0,00	0,00	0,00
2. Manufacturing Industries and Constr.	0,00	0,00	0,00	0,00	0,00	0,00
3. Transport	0,00	0,00	0,00	0,00	0,00	0,00
4. Other Sectors	0,02	0,01	0,01	0,02	0,02	0,02
5. Other	NA	NA	NA	NA	NA	NA
B. Fugitive Emissions from Fuels	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO
1. Solid Fuels	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO
2. Oil and Natural Gas	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO
2. Industrial Processes	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO
A. Mineral Products	NO	NO	NO	NO	NO	NO
B. Chemical Industry	NO	NO	NO	NO	NO	NO
C. Metal Production	NA	NA	NA	NA	NA	NA
D. Other Production						
E. Production of Halocarbons and SF ₆						
F. Consumption of Halocarbons and SF ₆						
G. Other	NA	NA	NA	NA	NA	NA
3. Solvent and Other Product Use	NA,NE	NA,NE	NA,NE	NA,NE	NA,NE	NA,NE
4. Agriculture	0,05	0,05	0,05	0,05	0,05	0,05
A. Enteric Fermentation						
B. Manure Management	0,00	0,00	0,00	0,00	0,00	0,00
C. Rice Cultivation						
D. Agricultural Soils	0,04	0,04	0,04	0,04	0,04	0,04
E. Prescribed Burning of Savannas	NA	NA	NA	NA	NA	NA
F. Field Burning of Agricultural Residues	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO	NA,NO
G. Other	NA	NA	NA	NA	NA	NA
5. Land Use, Land-Use Change and Forestry	NA,NE	NA,NE	NA,NE	NA,NE	NA,NE	NA,NE
A. Forest Land	NE	NE	NE	NE	NE	NE
B. Cropland	NE	NE	NE	NE	NE	NE
C. Grassland	NE	NE	NE	NE	NE	NE
D. Wetlands	NE	NE	NE	NE	NE	NE
E. Settlements	NA,NE	NA,NE	NA,NE	NA,NE	NA,NE	NA,NE
F. Other Land	NA,NE	NA,NE	NA,NE	NA,NE	NA,NE	NA,NE
G. Other	NA	NA	NA	NA	NA	NA
6. Waste	IE,NA,NE	IE,NA,NE	IE,NA,NE	IE,NA,NE	IE,NA,NE	IE,NA,NE
A. Solid Waste Disposal on Land						
B. Waste-water Handling	NA,NE	NA,NE	NA,NE	NA,NE	NA,NE	NA,NE
C. Waste Incineration	IE,NA	IE,NA	IE,NA	IE,NA	IE,NA	IE,NA
D. Other	NA	NA	NA	NA	NA	NA
7. Other (as specified in Summary 1.A)	NA	NA	NA	NA	NA	NA
Total N₂O emissions including N₂O from LULUCF	0,07	0,07	0,07	0,07	0,08	0,07
Total N₂O emissions excluding N₂O from LULUCF	0,07	0,07	0,07	0,07	0,08	0,07
Memo Items:						
International Bunkers	NA,NE,NO	0,01	0,01	0,01	0,00	0,00
Aviation	NE,NO	0,00	0,00	0,00	0,00	0,00
Marine	NA,NE,NO	0,01	0,01	0,01	0,00	0,00
Multilateral Operations	NO	NO	NO	NO	NO	NO
CO₂ Emissions from Biomass						

Table 8 EMISSION TRENDS HFCs, PFCs and SF₆ - Inventory 2012 - Submission 2014 v1.1 - FAROE ISLANDS.

GREENHOUSE GAS SOURCE AND SINK CATEGORIES	Base year (1990)	1995	2000	2005	2009	2010	2012
	(Gg)	(Gg)	(Gg)	(Gg)	(Gg)	(Gg)	(Gg)
Emissions of HFCs - (Gg CO₂ equivalent)	NA,NE,NO	0,02	4,35	11,20	11,61	12,12	14,22
HFC-23	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO
HFC-32	NA,NE,NO	NA,NE,NO	0,00	0,00	0,00	0,00	0,00
HFC-41	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO
HFC-43-10mee	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO
HFC-125	NA,NE,NO	NA,NE,NO	0,00	0,00	0,00	0,00	0,00
HFC-134	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO
HFC-134a	NA,NE,NO	0,00	0,00	0,00	0,00	0,00	0,00
HFC-152a	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO
HFC-143	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO
HFC-143a	NA,NE,NO	NA,NE,NO	0,00	0,00	0,00	0,00	0,00
HFC-227ea	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO
HFC-236fa	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO
HFC-245ca	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO
Unspecified mix of listed HFCs- (Gg CO ₂ equivalent)	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO
Emissions of PFCs - (Gg CO₂ equivalent)	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO
CF ₄	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO
C ₂ F ₆	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO
C ₃ F ₈	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO
C ₄ F ₁₀	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO
c-C ₄ F ₈	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO
C ₅ F ₁₂	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO
C ₆ F ₁₄	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO
Unspecified mix of listed PFCs - (Gg CO ₂ equivalent)	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO
Emissions of SF₆- (Gg CO₂ equivalent)	NA,NE,NO	0,15	0,08	0,15	0,21	0,17	0,19
SF ₆	NA,NE,NO	0,00	0,00	0,00	0,00	0,00	0,00

Table 9 EMISSION TRENDS SUMMARY - Inventory 2012 - Submission 2014 v1.1 - FAROE ISLANDS.

GREENHOUSE GAS EMISSIONS	Base year (1990)	1995	2000	2005	2010	2012
	CO ₂ equivalent (Gg)	CO ₂ equivalent (Gg)	CO ₂ equivalent (Gg)	CO ₂ equivalent (Gg)	CO ₂ equivalent (Gg)	CO ₂ equivalent (Gg)
CO ₂ emissions including net CO ₂ from LULUCF	667,67	539,31	664,33	773,89	839,23	814,23
CO ₂ emissions excluding net CO ₂ from LULUCF	667,67	539,31	664,33	773,89	839,23	814,23
CH ₄ emissions including CH ₄ from LULUCF	18,35	18,85	18,79	20,13	19,10	18,83
CH ₄ emissions excluding CH ₄ from LULUCF	18,35	18,85	18,79	20,13	19,10	18,83
N ₂ O emissions including N ₂ O from LULUCF	21,98	20,65	22,00	22,77	23,26	22,84
N ₂ O emissions excluding N ₂ O from LULUCF	21,98	20,65	22,00	22,77	23,26	22,84
HFCs	NA,NE,NO	0,02	4,35	11,20	12,12	14,22
PFCs	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO	NA,NE,NO
SF ₆	NA,NE,NO	0,15	0,08	0,15	0,17	0,19
Total (including LULUCF)	708,00	578,99	709,55	828,14	893,88	870,32
Total (excluding LULUCF)	708,00	578,99	709,55	828,14	893,88	870,32

GREENHOUSE GAS SOURCE AND SINK CATEGORIES	Base year (1990)	1995	2000	2005	2010	2012
	CO ₂ equivalent (Gg)	CO ₂ equivalent (Gg)	CO ₂ equivalent (Gg)	CO ₂ equivalent (Gg)	CO ₂ equivalent (Gg)	CO ₂ equivalent (Gg)
1. Energy	675,97	546,19	672,67	784,64	849,84	823,97
2. Industrial Processes	NA,NE,NO	0,18	4,43	11,36	12,29	14,41
3. Solvent and Other Product Use	NA,NE	NA,NE	NA,NE	NA,NE	NA,NE	NA,NE
4. Agriculture	32,04	32,62	32,45	32,14	31,76	31,93
5. Land Use, Land-Use Change and Forestry	NA,NE	NA,NE	NA,NE	NA,NE	NA,NE	NA,NE
6. Waste	IE,NA,NE,NO	IE,NA,NE,NO	IE,NA,NE,NO	IE,NA,NE,NO	IE,NA,NE,NO	IE,NA,NE,NO
7. Other	NA	NA	NA	NA	NA	NA
Total (including LULUCF)	708,00	578,99	709,55	828,14	893,88	870,32

Annex 2a Emissions factors – stationary combustion

The emissions factors used for calculating the Faroese emission in following stationary combustion categories:

- 1A1a Public Electricity and Heat Production
- 1A2 Manufacturing Industry and Construction
- 1A4a Commercial/Institutional
- 1A4b Residential

are found in Table 10.

Table 10 Emission factors for stationary combustion.

Category	Fuel	Pollutant	1990-2005	2006-2012
Public electricity and heat production	Gas/diesel oil	CH ₄ (g/GJ)	0,9	0,9
		CO ₂ (kg/GJ)	74	74
		N ₂ O (g/GJ)	0,4	0,4
	Heavy fuel oil	CH ₄ (g/GJ)	0,9	0,9
		CO ₂ (kg/GJ)	78,4	78,1-79,25
		N ₂ O (g/GJ)	0,3	0,3
Manufacturing industries and construction	Gas/diesel oil	CH ₄ (g/GJ)	0,2	0,2
		CO ₂ (kg/GJ)	74	74
		N ₂ O (g/GJ)	0,4	0,4
	Heavy fuel oil	CH ₄ (g/GJ)	1,3	1,3
		CO ₂ (kg/GJ)	77,4	77,4
		N ₂ O (g/GJ)	5	5
	Kerosene	CH ₄ (g/GJ)	0,2	0,2
		CO ₂ (kg/GJ)	71,9	71,9
		N ₂ O (g/GJ)	0,4	0,4
Commercial/Institutional	Gas/diesel oil	CH ₄ (g/GJ)	0,7	0,7
		CO ₂ (kg/GJ)	74	74
		N ₂ O (g/GJ)	0,4	0,4
	Kerosene	CH ₄ (g/GJ)	0,7	0,7
		CO ₂ (kg/GJ)	71,9	71,9
		N ₂ O (g/GJ)	0,4	0,4
Residential	Gas/diesel oil	CH ₄ (g/GJ)	0,7	0,7
		CO ₂ (kg/GJ)	74	74
		N ₂ O (g/GJ)	0,6	0,6
	Kerosene	CH ₄ (g/GJ)	0,7	0,7
		CO ₂ (kg/GJ)	71,9	71,9
		N ₂ O (g/GJ)	0,6	0,6

The emissions factors for calculating the Faroese emissions from the Waste sector are found in Table 11.

Table 11 Emission factors for waste incineration.

Year	Fossil waste %	CO ₂ EMF - fossil Kg pr GJ	CO ₂ EMF - biogen Kg pr GJ	CH ₄ EMF - tot g pr GJ	N ₂ O EMF - tot g pr GJ
1990	32,2	37	86,7	6	4
1991	32,2	37	86,7	6	4
1992	35,4	37	84,2	6	4
1993	36,9	37	83,0	6	4
1994	36,9	37	83,0	6	4
1995	39,3	37	81,1	6	4
1996-2012	41,2	37	79,6	6	4

Annex 2b Emissions factors – mobile combustion

The emissions factors used for calculating the Faroese emission in following mobile combustion categories:

- 1A3a Civil aviation
- 1A3b Road transport
- 1A3d Navigation
- 1A4c Agriculture, Forestry and Fishing

are found in Table 12, Table 13 and Table 14.

Table 11 Emission factors for aviation, 1990-2012.

	CH ₄ - g pr GJ	CO ₂ - Kg pr GJ	N ₂ O - g pr GJ
1990	465,9	72,0	2,680
1991	465,9	72,0	2,680
1992	465,9	72,0	2,680
1993	465,9	72,0	2,680
1994	465,9	72,0	2,680
1995	465,9	72,0	2,680
1996	465,9	72,0	2,680
1997	465,9	72,0	2,680
1998	465,9	72,0	2,680
1999	465,9	72,0	2,680
2000	465,9	72,0	2,680
2001	465,9	72,0	2,608
2002	473,7	72,0	2,611
2003	475,0	72,0	2,611
2004	523,3	72,0	2,624
2005	718,0	72,0	2,675
2006	716,6	72,0	2,669
2007	719,3	72,0	2,669
2008	716,6	72,0	2,669
2009	716,6	72,0	2,669
2010	720,2	72,0	2,669
2011	720,2	72,0	2,669
2012	465,9	72,0	2,680

Table 13 Emission factors for road transport, 1990-2012.

	Diesel			Gasoline		
	CH ₄	CO ₂	N ₂ O	CH ₄	CO ₂	N ₂ O
1990	6,741	74	1,729	27,566	73	2,848
1991	6,688	74	1,725	27,153	73	2,872
1992	6,671	74	1,720	26,115	73	2,946
1993	6,608	74	1,685	25,296	73	3,003
1994	6,626	74	1,697	23,852	73	3,095
1995	6,695	74	1,770	22,470	73	3,174
1996	6,706	74	1,838	21,181	73	3,244
1997	6,624	74	1,918	19,857	73	3,290
1998	6,443	74	2,007	18,701	73	3,232
1999	6,206	74	2,105	17,515	73	3,202
2000	5,868	74	2,163	16,587	73	3,189
2001	5,629	74	2,263	15,600	73	3,128
2002	5,284	74	2,317	14,526	73	3,036
2003	4,937	74	2,378	13,536	73	2,917
2004	4,640	74	2,435	12,437	73	2,787
2005	4,315	74	2,480	11,480	73	2,604
2006	3,940	74	2,528	10,459	73	2,393
2007	3,251	74	2,568	9,685	73	2,216
2008	2,523	74	2,621	9,047	73	2,033
2009	2,017	74	2,659	8,547	73	1,921
2010	1,698	74	2,716	8,183	73	1,766
2011	1,423	74	2,782	7,766	73	1,635
2012	1,139	74	2,811	7,386	73	1,459

Table 14 Emission factors for Navigation (diesel and residual) and Fisheries (diesel), 1990-2012.

	Navigation - diesel			Navigation and Fisheries - Residual			Fisheries - diesel		
	CH ₄	CO ₂	N ₂ O	CH ₄	CO ₂	N ₂ O	CH ₄	CO ₂	N ₂ O
1990	1,564	74	4,684	1,653	78	4,890	1,519	74	4,684
1991	1,571	74	4,684	1,645	78	4,890	1,530	74	4,684
1992	1,579	74	4,684	1,642	78	4,890	1,541	74	4,684
1993	1,582	74	4,684	1,646	78	4,890	1,553	74	4,684
1994	1,586	74	4,684	1,649	78	4,890	1,565	74	4,684
1995	1,600	74	4,684	1,651	78	4,890	1,578	74	4,684
1996	1,588	74	4,684	1,668	78	4,890	1,592	74	4,684
1997	1,502	74	4,684	1,694	78	4,890	1,606	74	4,684
1998	1,493	74	4,684	1,712	78	4,890	1,622	74	4,684
1999	1,461	74	4,684	1,724	78	4,890	1,639	74	4,684
2000	1,466	74	4,684	1,737	78	4,890	1,656	74	4,684
2001	1,486	74	4,684	1,753	78	4,890	1,673	74	4,684
2002	1,519	74	4,684	1,767	78	4,890	1,689	74	4,684
2003	1,513	74	4,684	1,820	78	4,890	1,704	74	4,684
2004	1,506	74	4,684	1,828	78	4,890	1,718	74	4,684
2005	1,510	74	4,684	1,869	78	4,890	1,731	74	4,684
2006	1,487	74	4,684	1,897	78	4,890	1,743	74	4,684
2007	1,498	74	4,684	1,906	78	4,890	1,753	74	4,684
2008	1,511	74	4,684	1,912	78	4,890	1,762	74	4,684
2009	1,515	74	4,684	1,925	78	4,890	1,770	74	4,684
2010	1,509	74	4,684	1,934	78	4,890	1,775	74	4,684
2011	1,502	74	4,684	1,943	78	4,890	1,780	74	4,684
2012	1,658	74	4,684	1,952	78	4,890	1,785	74	4,684

DENMARK'S NATIONAL INVENTORY REPORT 2014

Emission Inventories 1990-2012 - Submitted under
the United Nations Framework Convention on
Climate Change and the Kyoto Protocol

This report is Denmark's National Inventory Report 2014.
The report contains information on Denmark's emission in-
ventories for all years' from 1990 to 2012 for CO₂, CH₄, N₂O,
HFCs, PFCs and SF₆, NO_x, CO, NMVOC, SO₂.

ISBN: 978-87-7156-071-8
ISSN: 2245-0203