



THE PARTICLE PROJECT

2021

Scientific Report from DCE – Danish Centre for Environment and Energy

No. 500

2022



AARHUS
UNIVERSITY

DCE – DANISH CENTRE FOR ENVIRONMENT AND ENERGY

THE PARTICLE PROJECT 2021

Scientific Report from DCE – Danish Centre for Environment and Energy

No. 500

2022

Thomas Ellermann
Andreas Massling
Rossana Bossi
Claus Nordstrøm

Aarhus University, Department for Environmental Science



AARHUS
UNIVERSITY

DCE – DANISH CENTRE FOR ENVIRONMENT AND ENERGY

Data sheet

Series title and no.:	Scientific Report from DCE – Danish Centre for Environment and Energy No. 500
Title:	The Particle Project 2021
Authors:	Thomas Ellermann, Andreas Massling, Rossana Bossi and Claus Nordstrøm
Institution:	Aarhus University, Department for Environmental Science
Publisher:	Aarhus University, DCE – Danish Centre for Environment and Energy ©
URL:	http://dce.au.dk/en
Year of publication:	June 2022
Editing completed:	June 2022
Referee:	Matthias Ketzel
Quality assurance, DCE:	Vibeke Vestergaard Nielsen
Linguistic QA:	Vibeke Vestergaard Nielsen
Financial support:	Ministry of the Environment
Please cite as:	Ellermann, T., Massling, A., Bossi, R., & Nordstrøm, C. 2022. The Particle Project 2021. Aarhus University, DCE – Danish Centre for Environment and Energy, 38 pp. Scientific Report from DCE – Danish Centre for Environment and Energy No. 500. http://dce2.au.dk/pub/SR500.pdf
	Reproduction permitted provided the source is explicitly acknowledged
Abstract:	The Particle Project 2021 continues the measurements of the long-term trends of particle number concentrations and size distributions for submicron particles as well as the concentrations of elemental carbon in the ambient fine particle fraction (PM _{2.5}) at the Copenhagen urban background measurement station HCØ. The results from the measurements at urban background are compared to results from urban street, suburban and rural locations. The results show decreasing concentrations for both particle number concentrations and elemental carbon, which are mainly due to decreasing emissions on national as well as international level. The report also presents results from an analysis of the temporal variations of the particulate air pollution.
Keywords:	Particulate air pollution, PM _{2.5} , particle number and size distribution, elemental carbon, status 2021, temporal variation, long term trends, urban background,
Layout:	Majbritt Pedersen-Ulrich
Front page photo:	Thomas Ellermann
ISBN:	978-87-7156-693-2
ISSN (electronic):	2244-9981
Number of pages:	38
Internet version:	The report is available in electronic format (pdf) at http://dce2.au.dk/pub/SR500.pdf

Contents

Sammenfatning	5
Partikelantal	5
Elementært kulstof, EC	6
Tidsmæssige variationer	6
Summary	8
Particle number	8
Elemental carbon, EC	9
Temporal variations	9
1. Introduction	11
2. Temporal variation of PM2.5 and PM10 based on TEOM measurements	12
2.1 Diurnal variation	12
3. Elemental carbon (EC) mass concentration	16
3.1 Long-term trends	16
3.2 Monthly variations in EC	18
4. Measurements of particle number concentrations and particle size distribution	21
4.1 Particle number size distribution	22
4.2 Long-term trends for particle number fractions	24
4.3 Long term trend for total particle number	26
4.4 Monthly variations in particle number concentrations	27
4.5 Diurnal variations in particle number concentrations	31
References	37

Sammenfatning

I *The Particle Project 2021* rapporteres resultater af målinger af partikelstørrelsesfordelingen fra 11 nm op til 478/550 nm, partikelantal samt elementært kulstof i Københavns bybaggrund. Disse målinger er et supplement til luftovervågningsprogrammet under NOVANA og bidrager med forskning omkring kilderne til partikelforureningen i Danmark. Resultaterne for bybaggrund sammenholdes med tilsvarende resultater fra en landligt baseret station nord for Roskilde (RISØ), en station i den Københavnske forstad Hvidovre (HVID), samt fra en station i en trafikeret gade i København (HCAB). I rapporten præsenteres resultater for niveauer og udviklingstendenser for partikelantal og elementært kulstof sammen med en analyse af sæson- og døgnvariation af disse partikelkomponenter. Analysen af den tidsmæssige variation suppleres med resultaterne fra analyse af døgnvariation i PM_{2,5} og PM₁₀ baseret på målinger af PM med høj tidsopløsning via TEOM-metoden (Tapered Element Oscillating Microbalance).

Partikelantal

Langtransporterede partikler og deres gasformige forstadier bidrager til partikelantal for partikler med en diameter mindre end 1 µm. Langtransporterede partikler udgør den største andel på de landlige stationer og en mindre andel på stationerne i bybaggrund og forstad, hvor lokale kilder spiller en relativt større rolle. Mindst er andelen af langtransporterede partikler målt på gadestationerne, hvor den største andel udgør bidraget fra lokal trafik. Det relativt største bidrag fra langtransporterede partikler ser man i landlig baggrund på målestationen RISØ. Det næststørste relative bidrag fra langtransporterede partikler finder man på HCØ (bybaggrund), dernæst kommer HVID (forstad), mens dette bidrag udgør den mindste relative andel på HCAB (trafikeret gade).

Set over lang tid er der målt et fald i partikelantal ved alle målestationerne, men der er forskel i udviklingstendensen for de forskellige partikelstørrelsesfraktioner og målestationer. Der er også forskel på dataseriernes længde, hvor målingerne på målestationerne HCØ og HCAB begyndte i 2002, RISØ i 2005 og HVID først i 2015. Dataserien for forstadsmålestationen HVID er for kort til at kunne vurdere udviklingstendenserne. Grundet tekniske problemer med udstyret er der endvidere huller i tidsserierne fra 2016-2019, som varierer fra målestation til målestation. Disse huller gælder for den mindste partikel-fraktion (11-41 nm), hvilket også medfører huller i tidsserien for måleområdet 11 – 478/550nm.

Partikler mellem 11 og 41 nm: For de seneste 10 til 15 år ses en tendens til, at partikelantallet ligger på et relativt stabilt niveau ved bybaggrundsmålestationen HCØ og landbaggrundsmålestationen RISØ. Ved gademålestationen HCAB ses et betydeligt fald i partikelantallet, som dog også flader ud inden for de seneste år. Ved gademålestationen kommer den største andel af partiklerne i denne fraktion fra udstødning og det store fald hænger sammen med den øgede brug af partikelfiltre på køretøjerne.

Partikler mellem 41 og 110 nm og partikler mellem 110 og 478/550 nm: Ved bybaggrundsmålestationen HCØ, landbaggrundsmålestationen RISØ og gademå-

lestationen HCAB måles et generelt fald i partikelantal for partikler med diameter mellem 41 og 110 nm og partikler mellem 110 og 478/550 nm i løbet af de seneste 15-20 år. Ved bybaggrundsmålestationen HCØ og landbaggrundsmålestationen (RISØ) har der dog været et nogenlunde stabilt niveau gennem de seneste fem år. I 2018 blev der målt en noget højere værdi sammenlignet med de omkringliggende år, hvilket formentligt skyldes, at 2018 var et usædvanligt tørt år, hvor sommernedbøren var 40% lavere end for de tilsvarende perioder i 2017 og 2019. Ved gademålestationen HCAB måles fortsat et fald i partikelantal gennem de seneste fem år, hvilket skyldes, at det løbende fald i udledningerne som følge af forbedring af vognparken slår tydeligere igennem på gademålestationen end ved baggrundsmålestationerne.

Partikler mellem 11 – 478/550nm: Når partikelantallet for det samlede måleområde vurderes, ses der et mindre fald i partikelantallet for bybaggrundsmålestationen (HCØ) og landbaggrundsmålestationen (RISØ), mens der for gademålestationen (HCAB) ses et betydeligt fald i partikelantallet – navnlig i den første del af måleserien (2002-2012), hvorefter faldet flader ud.

Elementært kulstof, EC

Årsmiddelkoncentrationer for EC blev i 2021 målt til 0,24 µg/m³ ved bybaggrundsmålestationen HCØ, hvilket er omkring 10% lavere end målt i 2020. Ved den landlige målestation RISØ var faldet mindre (3%), mens faldet var markant højere ved gademålestationen HCAB (18%). Da forstadsmålestationen HVID skulle flyttes i sommeren 2021, var der kun data for omkring 6,5 måneder og dermed ikke data til beregning årsmiddelværdier.

Siden 2015 er koncentrationerne af EC faldet med omkring 35% ved baggrundsmålestationen HCØ. Den vigtigste årsag til faldet i koncentrationerne er de generelle reduktioner i udledningerne af EC på både nationalt og internationalt plan. Ved gademålestationen HCAB er der målt et stort fald på omkring 76% i perioden fra 2010 til 2021, hvilket hovedsageligt skyldes fald i udledningerne fra vejtrafik.

EC er faldet med omkring 45% ved den landlige målestation RISØ i perioden fra 2010 til 2021. Ved forstadsmålestationen HVID er der målt et fald på omkring 29% siden 2016. Faldet i koncentrationerne af EC har siden 2016 forløbet meget ensartet ved bybaggrundsmålestationen, landmålestationen og målestationen i forstad, hvilket indikerer, at det er de generelle reduktioner af udledningerne på nationalt og internationalt niveau, som er årsag til hovedparten af faldet i koncentrationerne.

Tidsmæssige variationer

Denne årsrapport præsenterer resultater fra analyse af de tidsmæssige variationer i luftforureningen med partikler. Døgnvariationen i partikelmassen er undersøgt ud fra målinger af PM_{2,5} og PM₁₀ med TEOM-metoden, som måler partikelmassen på timemiddelniveau ved den landlige målestation RISØ (PM₁₀) og ved gademålestation HCAB (PM_{2,5} og PM₁₀). Målingerne af EC udføres som døgnmiddelmålinger, og disse er anvendt til fastlæggelse af månedsvariationen af EC ved alle fire målestationer. Partikelantal måles på timemiddelniveau, og disse data er anvendt til både at bestemme måneds- og døgnvariationen.

Analysen af månedsvariationen af EC viser, at EC har de højeste koncentrationer om vinteren og de laveste koncentrationer om sommeren. Dette hænger sammen med, at EC hovedsageligt stammer fra udledninger fra vejtrafik og boligopvarmning baseret på brændefyring. Månedsvariationen er mindst ved gademålestationen HCAB og højest ved forstadsmålestationen HVID, hvorefter følger bybaggrundsmålestationen HCØ og den landlige målestation RISØ.

I modsætning til dette, så er partikelantal højere om sommeren end om vinteren ved bybaggrundsmålestationen HCØ og den landlige målestation RISØ. Ved gademålestationen HCAB ses omtrent ens partikelantal sommer og vinter. Årsagen til disse månedsvariationer er, at partikler med diameter mellem 11 og 478 nm stammer fra udledninger fra vejtrafik og boligopvarmning med brændefyring samt fotokemiske reaktioner i atmosfæren. De fotokemiske reaktioner forløber kun, når der er sollys, og derfor ses et relativt stort bidrag fra de fotokemiske reaktioner om sommeren. Ved gademålestationen HCAB er partikelantallet stærkt domineret af udledninger fra vejtrafikken, og bidraget fra de fotokemiske reaktioner er derfor formentligt for lille til at kunne blive observeret.

Døgnvariationen af PM_{10} ved gademålestationen HCAB viste det forventede mønster præget af udledningerne fra vejtrafikken, men der blev også observeret en uventet bred top fra sent på formiddagen til midt på eftermiddagen i den årlige gennemsnitlige døgnvariation. Analyserne viste, at denne brede top stammede fra vintersaltning af vejen i den første del af februar, hvor der var en periode med frostvejr og ringe nedbør. $PM_{2,5}$ ved gademålestationen HCAB viste en mindre tydelig døgnvariation i overensstemmelse med, at størstedelen af $PM_{2,5}$ stammer fra langtransporterede luftforurening og kun en mindre del stammer fra vejtrafik. Ved den landlige målestation RISØ blev der ikke observeret en klar døgnvariation, hvilket hænger sammen med, at langt hovedparten af $PM_{2,5}$ stammer fra langtransporteret luftforurening.

Døgnvariationen i partikelantal viste mange interessante mønstre som for en stor del kan tilskrives de forskellige kilder til partikelantal. Eksempler på dette ses på døgnvariationen ved gademålestationen HCAB, hvor der tydeligt kan identificeres toppe relateret til myldretid, og toppe der kan tilskrives udledninger i forbindelse med de sociale aktiviteter lørdag aften. Ved bybaggrundsmålestationen HCØ, ses en tydelig top i partikel antal fra sidst på formiddag til ud på eftermiddagen om sommeren (april-september). En lignende top kan ikke ses om vinteren (januar-marts og oktober-december). Denne top kan bedst forklares ved de fotokemiske reaktioner i atmosfæren, som fører til dannelse af partikler, når solen skinner om sommeren. Ved forstadsmålestationen HVID blev der observeret relativt højt partikelantal på vinteraftner, hvilket formentligt skal tilskrives boligopvarmning med brændefyring. Da forstadsmålestationen HVID blev flyttet i sommeren 2021, var det ikke muligt at analysere forskellen mellem sommer og vinter ved denne målestation. Analysen af døgnvariationen af partikelantal vil blive gentaget, når der foreligger data fra hele 2022 for alle fire målestationer for på den måde at forbedre datagrundlaget for analysen.

Summary

The *Particle Project 2021* reports time series from measurements of particle number concentrations of submicrometer particles and Elemental Carbon (EC) in fine particles with diameter smaller than $2.5\ \mu\text{m}$ ($\text{PM}_{2.5}$) in urban background in Copenhagen. In order to increase knowledge on the sources of these particulate air pollutants the measurements are carried out as a supplement to the Danish air quality monitoring program under NOVANA. Trends in urban background (HCØ) are compared to rural location (RISØ), suburban location (HVID) and urban street (HCAB).

Particle number

Regional and long-range transported aerosols and precursors hereof contribute to the particle number concentration in the submicrometer size range. The highest relative contribution from long-range transported particles is found at rural background locations while the other locations are more influenced by local sources. For these reasons, long-range transported particles make up a relative smaller fraction at the suburban station HVID and the urban background station HCØ and an even smaller fraction at the street station HCAB, where the highest particle number concentrations are measured due to the significant contributions from local traffic.

A trend of decreasing particle number concentrations is observed at all stations when considering periods of the order of a decade. A general decreasing trend is observed for the rural background station RISØ, the urban background station HCØ and the street station HCAB when considering the measurements of particles in the range 41 - 110 nm and 110 - 478/550 nm of the last 15 - 20 years. However, the measurements show for these fractions very stable numbers for the rural station RISØ and the urban background station HCØ in the last five years with an increased value in 2018. This might be due to an unusual dry year in 2018 where summer precipitation was about 40% lower than summer precipitation in 2017 and 2019, thus leading to reduced wet deposition. In contrast, the decreasing trend is still observed at the street station HCAB for these two size fractions in the last five years where also local contributions from traffic are expected. At the suburban station HVID only measurements covering six years are available.

Data for particle numbers in the size range 11 - 41 nm show a gap due to instrumental problems. The general tendency shows stable particle numbers for this small size fraction in the last ten to fifteen years at the rural background station RISØ and the urban background station HCØ. A decreasing trend can be observed at the street station HCAB which levels more and more out in the last years.

The total number of particles for the size range 11 - 478/550nm shows a small decrease at the rural background station RISØ and the urban background station HCØ while the decrease is much more pronounced at the street station HCAB, especially between 2002 and the following ten years. It is still not possible to make a firm conclusion for the suburban station HVID due to the limited length of the data series.

Elemental carbon, EC

The annual mean concentration of EC in 2021 at the urban background station HCØ in Copenhagen was $0.24 \mu\text{g}/\text{m}^3$, which is about 10% lower than measured in 2020. At the rural station RISØ the change was smaller (3%), while the decrease was more pronounced at the urban street station HCAB (18%). The highest decrease is expected at the street station where the impact from road traffic is largest. The suburban station HVID had to be relocated in 2021 and therefore there were only data for about 6.5 months.

Since 2015, EC has decreased with about 35% at the background station HCØ. The main reason for the decrease in the concentrations at HCØ is the general reductions in the emissions both at national and international level.

EC has decreased by 45% at the urban background station RISØ over the period 2010 to 2021. At the suburban station HVID there has been a reduction from 2016 to 2020 of about 29%. Since 2016, the decreasing trends have been quite similar at urban background, suburban and rural stations, indicating that the general reductions in emissions on national and international level can explain the major part of the decreasing trends.

At HCAB, there has been a pronounced reduction of about 76% ($1.8 \mu\text{g}/\text{m}^3$) in the annual average concentrations over the period 2010 to 2021, predominantly due to a reduction in local road traffic emissions, caused by introduction of cars with newer and cleaner engine technology and exhaust after treatment as particle filters.

Temporal variations

This annual report presents results from an analysis of the temporal variations of the particulate air pollution. The diurnal variation of particulate mass is analysed based on measurements of $\text{PM}_{2.5}$ and PM_{10} using the TEOM method (Tapered Element Oscillating Microbalance) that measures particulate mass with hourly time resolution at the rural station RISØ (PM_{10}) and the street station HCAB ($\text{PM}_{2.5}$ and PM_{10}). The measurements of EC are carried out with a time resolution of 24 h and these data are used to determine the monthly variations at all four stations. Particle number is measured with hourly variation and these data are used to study both the monthly and diurnal variation at all four stations.

The analysis of the monthly variations of EC show that EC has highest concentrations during winter and lowest concentrations during summer. This is because EC mainly originates from emissions from road traffic and household warming using wood burning. The smallest variation is at the street station HCAB and the highest variation is at the suburban station HVID followed by the urban background HCØ and regional background station RISØ.

In contrast to EC variation, the particle number is higher during summer than during winter at the urban background station HCØ and regional background station RISØ. At the street station HCAB, the particle number is roughly equal summer and winter. The reason for these monthly variations are the particle number originates from road traffic and wood burning and additionally from photochemical reactions in the atmosphere. These reactions require sunlight and therefore present an additional source to the particle number during summer. At the street station HCAB the particle number is

strongly dominated by the road traffic emissions and the contribution from the chemical formation of particles is most likely too small to be seen in the monthly variations.

The diurnal variations of PM_{10} at the street station HCAB shows the expected patterns consistent with road traffic being the main source of PM_{10} although there also was an unexpected broad peak from late morning to middle afternoon. This peak was due to winter salting of the street during the first part of February. $PM_{2.5}$ at HCAB showed a less pronounced diurnal variation in agreement with the main part of $PM_{2.5}$ originating from long-range transport and only minor contributions comes from road traffic emissions. At the rural background station RISØ, there were no clear diurnal variation consistent with the main source being long-range transport.

The diurnal variation of particle number showed many interesting patterns that to a large part could be assigned to the sources. Examples on these are the strong rush hour peaks at the street station HCAB and the impact from social activities during Saturday evenings. At the urban background station HCØ there also exists a large peak during midday and afternoon during the summer half year and that peak could not be seen during the winter half year. This peak can best be explained by the atmospheric photochemical reactions that leads to particle formation during sunlight periods in the summer half year. This peak was also seen at the rural background station RISØ but not at the street station HCAB. At the suburban station HVID there were also relatively high concentrations during winter evenings that most likely are due to household warming using wood burning. Due to relocation of HVID in 2021, it was not possible to analyze the difference between summer and winter season at this station. The analysis of the diurnal variations in particle number will be refined during the next annual report where data from 2022 will be available for the entire year for all four stations.

1. Introduction

DCE – Danish Centre for Environment and Energy has since 2001 carried out particle research in Denmark through a series of projects funded by the Danish Environmental Protection Agency (as an example see Nøjgaard et al., 2020 or Ellermann et al., 2021). These projects have enabled DCE to obtain fundamental new knowledge on the particulate air pollution in Denmark including knowledge on the contributions to particulate air pollution from wood burning and traffic. Besides this, the projects have been the basis for establishment of unique time series on the air pollution with elemental carbon and submicrometer particles focusing on particle number and size distribution. In popular terms these air pollution components are often termed soot and ultrafine particles. These air pollution components are associated with health effects and the knowledge established through the particle projects has been used in connection to studies of the health impact of airborne particle pollution in Denmark.

This report presents the results from the particle project in 2021 where the main aim has been to continue the time series based on measurements of elemental carbon, particle size distribution and particle number concentration at the urban background station at H.C. Ørsted Institute in Copenhagen (HCØ). These time series are important because they are fundamental to research concerned with the impact of air pollution on health. The measurements in urban background are especially important since the level of air pollution in urban background has been regarded as the best available measured proxy for the exposure of citizens to air pollution. Moreover, these time series can be used to evaluate the impact of emission regulations on the levels of these air pollution components in urban areas. The long-term measurements of particle number at all four Danish locations have also been extremely valuable as validation dataset during the introduction of particle number as new pollutant in the DCE modelling system (DEHM-UBM-AirGIS; Frohn et al. 2021 and Ketzel et al. 2021)

The main purpose of the project is the measurements in urban background (HCØ). In order to set these results into context, they will throughout the report be compared to the results from the street station at H.C. Andersens Boulevard (HCAB) in Copenhagen, the suburban station in Hvidovre (HVID) and the rural background station at Risø (RISØ) north of Roskilde.

The report presents also results from analysis of temporal variation in $PM_{2.5}$ and PM_{10} based on the results from the measurements of $PM_{2.5}$ and PM_{10} at hourly resolution using the TEOM method (Tapered Element Oscillating Microbalance). These measurements are part of the Danish air quality monitoring program, and they are useful in connection to the evaluation of the reasons behind the observed temporal variations in particle number concentrations and EC.

The analysis of $PM_{2.5}$ and PM_{10} is presented first in the report since this analysis serve as a background for the understanding of the temporal variations in particle number and EC. Chapter 3 presents results on EC and chapter 4 on particle number.

2. Temporal variation of PM_{2.5} and PM₁₀ based on TEOM measurements

In the Danish air quality monitoring program, the main measurements of PM_{2.5} and PM₁₀ are carried out on a diurnal basis using low volume sampling and gravimetric determination of the particle mass. These measurements follow the reference method specified in the EU air quality directive (EU, 2008).

PM_{2.5} and PM₁₀ are also measured using the TEOM method (Tapered Element Oscillating Microbalance) at the street station HCAB (PM_{2.5} and PM₁₀) and the rural background station RISØ (PM₁₀). No measurements are carried out with the TEOM method at the urban background station HCØ.

The reason for use of this additional particulate matter measurement method is that the TEOM method enables measurements of hourly averages of PM_{2.5} and PM₁₀ and hence these measurements can be used for the near real time information of the public about the actual status of the air quality of PM_{2.5} and PM₁₀. Moreover, the results from the TEOM method can be used for analysis of the temporal variations in PM_{2.5} and PM₁₀.

The TEOM method has higher uncertainty than the gravimetric method and one of the main reasons for this is that part of the particle mass can evaporate during the measurement and this has to be kept in mind when the results from the TEOM method is used. Consequently, the long-term trends of PM_{2.5} and PM₁₀ are only determined using the reference method and not the TEOM method. The long-term trends are reported in the annual reporting from the monitoring program (see Ellermann et al., 2022a for the most recently updated reporting).

In the following, diurnal variations in PM_{2.5} and PM₁₀ will be presented based on the data from the TEOM method.

2.1 Diurnal variation

The average diurnal variation in the concentrations of particulate matter is shown in Figure 2.1 - 2.3. The average diurnal variations are calculated by averaging all the hourly concentrations measured from 00:00 - 1:00 for the entire period of interest and then do the same for all 24 diurnal hours. If the period is long enough (typically 2-3 years), it is possible to average out the variations in meteorology and the final variation will illustrate eventual diurnal variations in emissions or chemical formations of an air pollutant. If the period is too short some "random" variation due to the meteorological variations will be left in the calculated average diurnal variation.

The diurnal variations have been divided into working days, and Saturdays and Sundays in order to capture the difference in the diurnal variations of emissions on the different days of the week. There are only 52 Saturdays or Sundays during a year compared to 260 working days, hence there will be higher remains from the meteorological variations left in the curves for Saturdays and Sundays compared to working days.

Figure 2.1 shows the average diurnal variation for PM_{10} at the rural background station RISØ. The majority of PM_{10} at RISØ originates from long-range transport and it is therefore expected that there will be no clear diurnal variations and no variation between different days of the week. The observed variations seen in Figure 2.1 do not show a regular pattern and there seems to be small difference between season and between days of week. The variations seen in Figure 2.1 are therefore most likely due to the natural variations in the meteorological conditions that was not completely averaged out due to use of a too short averaging period. In the next reporting when we include 2022 in the averaging we will better be able to average out the meteorological variability and this will most likely underpin this conclusion.

Figure 2.2 shows the average diurnal variations for $PM_{2.5}$ at the street station at HCAB. About 75% of $PM_{2.5}$ at HCAB originates from long-range transport while road traffic is responsible for the main part of the remaining 25% of $PM_{2.5}$ (Ellermann et al., 2022b). Hence, it is expected that the average diurnal variation is the sum of a large quite constant part from the long-range transported $PM_{2.5}$ and a smaller contribution reflecting the impact of the diurnal variation of road traffic on $PM_{2.5}$.

The measured diurnal variation for working days shows the expected pattern with a morning peak 5:00 – 9:00 corresponding to the morning rush hour. The afternoon rush hour is less clearly seen because of higher meteorological dispersion of the emissions during the afternoon and most likely also because of the afternoon rush hour period is spread over a longer time period than the morning rush hour.

There is relatively high concentration of $PM_{2.5}$ concentration during the evening. This might be because of increased road traffic due to leisure activities (i.e. Friday) and for the winter period it might also be due to a contribution from use of wood burning as household warming, but this has to be investigated further.

Saturdays and Sundays show different patterns than working days with one of the main differences being that the working day rush hours are lacking in the average diurnal variations. In addition, there is an increase in $PM_{2.5}$ from late morning until the late evening and early night reflecting a more even road traffic during weekends with a later start in the morning compared to working days. Moreover, there is a strong peak at 22:00-23:00 during Saturday evenings that occur at the same time as the fireworks in Tivoli. This illustrates that Tivoli's firework activities and other social activities influence $PM_{2.5}$ at HCAB.

Figure 2.3 shows the average diurnal pattern of PM_{10} at H.C. Andersens Boulevard. Overall, as expected the pattern for PM_{10} looks like the pattern for $PM_{2.5}$ but with a stronger influence from road traffic since road traffic makes up a larger part of PM_{10} than $PM_{2.5}$. The measured average diurnal variation for 2021 do not look quite as it is expected due to a large broad peak during midday and early afternoon. It is most clearly seen on Saturdays and Sundays, but it is also seen on working days. A more detailed analysis of the data shows that this peak can be assigned to an episode with frosty conditions and no precipitation during the first part of February. Due to salting of the roads, there is a rapid and large increase in PM_{10} late morning and midday (Figure 2.4). The onset of this increase in PM_{10} is most likely correlated with the drying of the road surface during the day and a strong resuspension of salt and road

dust (Denby et al., 2015). Figure 2.3 also shows the curves for the winter average diurnal variation if the February winter episode is omitted from the calculations (dashed black curve). The dashed curves show that this single episode with winter salting is the main explanation for the unexpected high PM_{10} during midday and early afternoon at all days of the week.




Figure 2.1. Average annual diurnal variations of PM_{10} in 2021 at the rural background station RISØ divided into working days, Saturdays and Sundays.




Figure 2.2. Average annual diurnal variations of $PM_{2.5}$ in 2021 at the street station HCAB divided into working days, and Saturdays and Sundays.




Figure 2.3. Average annual diurnal variations of PM_{10} in 2021 at the street station HCAB divided into working days, and Saturdays and Sundays. The dashed curve shows the diurnal variations when the winter episode from 4-15 February has been omitted.




Figure 2.4. $PM_{2.5}$ and PM_{10} measured using TEOM at the street station HCAB from 25 January to 20 February 2021. The sharp high peaks of PM_{10} from 4 February to 15 February are due to winter salting of the street and a strong resuspension of salt and road dust during the dry up of the street.

3. Elemental carbon (EC) mass concentration

Soot, the blackish or brownish substance formed during incomplete combustion (Andrea and Gelencsér, 2006) is typically measured by exploiting its light absorbing properties as Black Carbon (BC) or its chemical inertness as elemental carbon (EC). In the Danish Air Quality Monitoring Programme, EC is measured using the European standard thermal optical protocol EUSAAR2 (Cavalli et al., 2010) based on a thermal optical technique where the carbon atoms are combusted in the presence of oxygen at temperatures higher than 500°C (Birch and Cary, 1996; Cavalli et al., 2010).

Low Volume Samplers (LVS) equipped with PM_{2.5} inlets are located at the street station HCAB, urban background station HCØ, rural background station RISØ and suburban station HVID. Atmospheric aerosols are collected on quartz fiber filters using 24-hour time resolution. The filters are subsequently weighted to measure PM_{2.5} mass concentration in the case of HCØ station. At the three other stations LVS are dedicated to OC/EC sampling and thus the filters for EC determination are not weighted. Punches of the filters (1.48 cm²) are analyzed for EC using a Thermal/Optical Carbon Analyzer (Sunset Laboratory, Oregon USA) according to the EUSAAR 2 protocol (Cavalli et al., 2010).

EC has been monitored routinely at RISØ and HCAB from 2009/2010 and onwards. Monitoring of EC was extended to urban background in Copenhagen by September 2014, and to a suburban location in Hvidovre from October 2015. In 2021 the suburban station HVID had to be moved to another location in Hvidovre (roughly 600 m north) and in connection with this, the station was renovated and the new location had to be prepared. This caused a data gap at HVID of about 5.5 months.

3.1 Long-term trends

Table 3.1 presents the annual averages measured at the four stations in 2021. In 2021, annual mean values were: rural EC 0.22 µg/m³ < urban background EC 0.24 µg/m³ < suburban EC 0.36 µg/m³ < urban street EC 0.59 µg/m³ (Table 3.1). However, it should be noted that the average for HVID do not reflect a full year of measurements and that the value given most likely is somewhat higher than the annual average.

This pattern is consistent with the fact that the highest concentrations are measured at the street locations due to the strong emissions of EC from road traffic and that the lowest concentrations are measured in rural areas with only few local sources. The emissions in the city increase the concentrations at urban background station HCØ with about 13% compared to the rural station RISØ. The street station has the highest concentrations and here the concentrations are 173% higher or nearly three times higher than at RISØ.

Table 3.1. 2021 annual statistics for EC at Danish measurement sites. Annual mean value for EC and annual mean value for EC at the urban background station HCØ, suburban station HVID and street station (HCAB) relative to the rural background station RISØ (=100%). Data from the suburban station HVID covers only roughly 6.5 months due to relocation of the measurements station and hence the annual average has to be regarded as indicative and most likely it is giving a too high value since data mainly covers the cold months.

	Data coverage %	EC $\mu\text{g}/\text{m}^3$	EC relative to rural site (RISØ = 100%)
Street			
HCAB	94%	0.59	273%
Suburban			
HVID	48%	0.36	164%
Urban background			
HCØ	92%	0.24	113%
Rural background			
RISØ	98%	0.22	100%

Figure 3.1 shows the long-term trends for the annual averages of EC at the four stations. Urban background measurements of EC were not available until 2015 when initiated in *The Particle Project 2014-2016* (Nøjgaard et al., 2017). Since 2015, EC has decreased with about 35% ($0.14 \mu\text{g}/\text{m}^3$) at the urban background station HCØ. The main reason for the decrease in the concentrations at HCØ is the general reductions in the emissions at national as well as at international level.

At HCAB, there has been a pronounced reduction of about 76% ($1.8 \mu\text{g}/\text{m}^3$) in the annual average concentrations over the period 2010 to 2021, predominantly due to a reduction in road traffic emissions. This is a result of improved combustion technology in general, and the fact that the share of particle filters within the fleet of diesel vehicles has increased with more stringent emission standards for newer vehicles. The general reduction in emissions in Europe and urban areas has also contributed to the reductions at HCAB although these reductions play a minor role at HCAB.

EC has decreased by about 45% ($0.23 \mu\text{g}/\text{m}^3$) at rural background station RISØ over the period 2010 to 2021. At the suburban station HVID there has been a reduction from 2016 to 2020 of about 29% ($0.11 \mu\text{g}/\text{m}^3$). Since 2016, the reductions have been quite similar at the urban background, suburban and rural background stations indicating that it is the general reductions on national and international level that explain the main part of the reductions.




Figure 3.1. Long-term trends in annual average concentrations of EC at urban back-ground station HCØ, suburban station HVID, rural background station RISØ and street station HCAB. There is no annual average for 2021 from the suburban station HVID since it was relocated and this caused a gap in the data.

3.2 Monthly variations in EC

Figure 3.2 shows the measured time series of diurnal averages from all four stations since 2017. A clear seasonal variation can be seen in urban back-ground and at the rural and suburban locations with higher concentrations during winter compared to summer. This variation is mainly caused by biomass combustion, including residential wood combustion, as concluded in *The Particle Project 2017-2018* (Nøjgaard et al., 2018). In contrast, the seasonal variation at the curb site station HCAB is less pronounced. Here the emissions of EC from the road traffic are the most dominant source and hence the seasonal variation is much smaller compared to the other three stations. More striking is the clear decreasing trend in EC, which matches the trend in particle number (see next chapter Figure 4.2).

Figure 3.3 shows monthly averages for the four stations in 2021 and Figure 3.4 the monthly variation averaged over the period from 2017 to 2021. The seasonal variation with higher winter concentrations and lower summer concentrations is evident for the three stations in urban background and at the rural and suburban locations both for 2021 and the five years average. Based on five years of measurements at HCØ, mean winter concentrations average $0.38 \mu\text{g}/\text{m}^3$ during December-January-February, while summer concentrations show mean values of $0.23 \mu\text{g}/\text{m}^3$ during June-July-August. Autumn concentrations are slightly higher than spring concentrations, i.e. 0.32 vs. $0.28 \mu\text{g}/\text{m}^3$. Peak concentrations of 1 to $2 \mu\text{g}/\text{m}^3$ are occasionally observed at HCØ (Figure 3.2).

At the street station HCAB, only a small seasonal variation is observed due to the low seasonal variation in the emissions from the road traffic. The difference between winter and summer is less than 10% for the five-year averages (winter $0.93 \mu\text{g}/\text{m}^3$ vs. summer $0.87 \mu\text{g}/\text{m}^3$). The relatively low concentrations

in July and December are most likely due to the summer and Christmas vacations.







Figure 3.2. Time series for the diurnal average concentrations of EC at urban background station HCØ, suburban station HVID, rural background station RISØ and street station HCAB. The lack of data from the suburban station HVID are due to relocation of the measurement station.




Figure 3.3. Monthly averaged concentrations of EC in 2021 at the urban background station HCØ, suburban station HVID, rural background station RISØ and urban street station HCAB. The lack of data for HVID is due to the relocation of the station.




Figure 3.4. Monthly averaged concentrations of EC based on the period 01.01.2017 – 31.12.2021 at the urban background station HCØ, suburban station HVID, rural background station RISØ and urban street station HCAB. The error bars show the standard deviation of the monthly averages for the five-year period. The suburban measurement station HVID covers only 01.01.2017 – 31.12.2020 due to data gap in connection with the relocation of the measurement station in the middle of 2021.

4. Measurements of particle number concentrations and particle size distribution

Custom built DMPS instruments (Differential Mobility Particle Sizer) have been used from 2001/2002 and onwards during several Particle Projects to measure particle number size distributions in the submicrometer size regime. In our case particles with diameters in the range from 11 nm and up to 478-700 nm (see further details below). Submicrometer particles originate mainly from combustion processes where emissions from road traffic and household warming using wood burning are the most important sources. Submicrometer particles can also originate from biogenic sources and they can be formed via the chemical and physical processes in the atmosphere based on biogenic and anthropogenic precursors.

Particle number concentrations - similar to other air pollution concentrations - generally decrease with distance to major sources of aerosol particles. This is a result of polluted air being diluted with cleaner air that in turn leads to a general decrease in particle numbers. These mixing processes go along with particle dynamics processes as condensation, evaporation, coagulation and cloud processing, leading to changes in the general particle population (also called “aging” of aerosols). A result of this particle aging process is an increase in the mean particle diameter with distance to the major sources (Nøjgaard et al., 2015). It is expected that local sources in urban areas mainly contribute to particle number concentrations, and especially to the sizes around and smaller than 100 nm in diameter.

Particle number size distributions of diameters 6 - 700 nm were measured at the rural station RISØ (previously Lille Valby), urban background HCØ and street station HCAB. From 2017 and onwards, the instruments at HCAB and RISØ were replaced with commercial instruments delivered by TSI (Model 3938) (Table 4.1). These are SMPS instruments (Scanning Mobility Particle Sizer) and measures in the size range 11 - 478 nm. At HCØ, one of the original DMPS instruments was still in use in 2017 and 2018 but was exchanged in the beginning of 2019 with a new SMPS system (Model 3938). At the suburban station (HVID) an additional new SMPS system of the same type was used since the end of 2015. The new instruments were connected to new inlet systems, which unfortunately introduced losses, which has turned out to affect the general uncertainty of particle number concentrations (Nøjgaard et al., 2018). In addition, some problems occurred with some of the DMAs (Differential Mobility Analyzer), which are a part of the new SMPS systems. During 2019, these inlets and DMAs have been exchanged, implying that data from smaller size regimes can be used from February 2020 and onwards. In 2021, the data coverage was very good for all stations except the suburban station HVID where the data coverage was only 46% (Table 4.2). The suburban station HVID had to be relocated during the middle of 2021 and data for June-October are lacking.

As discussed in *The particle Project 2017-2018* (Nøjgaard et al., 2018), the slightly different measurement ranges between the new and the old instruments have implications for data comparison. The size range 11 - 550 nm will be discussed for the old DMPS instruments, whereas the size range 11 - 478 nm will be discussed for the new SMPS instruments. Only in this way a comparison of historical and new data is possible, and moreover, particles in the range 478 -

550 nm have very little impact on the particle number. However, the problems with particles in the range 11 - 41 nm, which were observed based on issues with the inlets and DMAs as discussed in the former Particle Project, were not solved until the beginning of 2020. Hence, data reported in the years 2017 to 2019 (RISØ, HCAB), 2019 (HVID) and 2015 to 2019 (HVID) is based on size ranges from 41 to 478 nm and particle numbers within this size range.

Table 4.1. Operation of instruments (old and new) at the four measurement stations and valid size ranges during the years 2002 to 2021.

	Rural background (RISØ)	Suburban (HVID)	Urban background (HCØ)	Street (HCAB)
Old instrument 11 nm – 550 nm	2005-2016	-	2002-2018	2002-2016
New instrument 41 nm – 478 nm	2017-2019	2015-2019	2019	2017-2019
New instrument 11 nm – 478 nm	2020-2021	2020-2021	2020-2021	2020-2021

Table 4.2. Data coverage in 2021 for SMPS measurements at rural background station RISØ, suburban station HVID, urban background station HCØ and street station HCAB.

	Rural background (RISØ)	Suburban (HVID)	Urban background (HCØ)	Street (HCAB)
Data coverage, %	92	46	92	94

4.1 Particle number size distribution

Figure 4.1 shows the particle number size distribution in the size range from 11 - 478 nm for the four stations in 2021 and for comparison also the results from 2020 are shown. The figure shows the annual average particle number size distributions that are calculated from the measurements with a time resolution of 30 minutes.

The data for 2020 covers only 11 months from February to December. January had to be left out since there were still problems with the instruments that were solved from February and ongoing. Data for 2021 covers the full year except for the suburban station HVID where data are lacking for June-October. The particle number size distribution at HVID is therefore uncertain as it only represents parts of the year 2021.

It has to be stated that in general particle numbers measured below 20 nm using typical SMPS instruments that cover large size ranges of the submicrometer size range are highly uncertain as calibration protocols in this range are challenging and require extremely sophisticated laboratory set-ups. Nevertheless, for larger size ranges SMPS instruments as used in our study give quite good results regarding general uncertainties, which can be estimated smaller than about 10 - 15 %.

In Figure 4.1 the derivative $dN/d(\log D_p)$ is plotted against the logarithm of the particle diameter D_p , which has the large advantage that the area under the curve (numeric integral) corresponds to the particle number. In that way, it is clearly visualized that e.g. the number of particles at HCAB is much higher compared to the other three stations (Figure 4.1). This difference is predominantly pronounced for the smallest fraction of the particles below about

100 nm which are also called ultrafine particles and where emissions from traffic at the street station strongly impact the particle number size distribution and thus also the particle number.

At the street station HCAB, the particle number size distribution looks similar in 2020 and 2021 for particles with diameters above about 25 nm (Figure 4.1). For particles in the range from 11 to about 25 nm the values are higher in 2021 compared to 2020. Particles in this size range predominantly originate from road traffic and the difference between 2020 and 2021 may therefore be related to the Covid-19 restrictions that were much more radical in 2020 compared to 2021, but it might also be due to the natural variations in the meteorological conditions from year to year.

At the urban background station HCØ and the rural background station RISØ the particle number size distributions look similar although the particle numbers are a little higher at HCØ compared to RISØ. At both stations the numbers of particles in the size range from about 15 nm to 60-70 nm were slightly lower in 2021 compared to 2020 while the levels were roughly the same for the particles with diameters above 70-80 nm. This difference between 2020 and 2021 may be related to the natural variations in the meteorological conditions. To an example the number of hours with sunshine were lower in 2021 compared to 2020 and this might have led to fewer photochemically formed particles in the atmosphere. However, it might also be due to small variations in the emissions from year to year, but it is not likely that it is related to Covid-19 restrictions.

At the suburban station HVID the particle number size distribution for 2021 differs compared to 2020 in much the same way as for HCØ and RISØ. However, since the data coverage is very low the particle number size distribution is very uncertain and it is therefore not possible to make a proper evaluation of the difference between 2020 and 2021.




Figure 4.1. Annual average of particle number size distributions at the rural background station RISØ, suburban station HVID, urban background station HCØ and street station HCAB during 2021 compared to 2020. Note the logarithmic x-axis. HVID is uncertain due to low data coverage in 2021.

4.2 Long-term trends for particle number fractions

For a more detailed analysis of the long-term trends, particle number concentrations were determined in specific size regimes. In this report, we have divided particles in three size fractions:

- Small size fraction: Diameters between 11 – 41 nm (both DMPS and SMPS)
- Medium size fraction: Diameters between 41 – 110 nm (both DMPS and SMPS),
- Large size fraction: Diameters between 110 – 550 nm for the old instruments (DMPS) and diameters between 110 – 478 nm for the new instruments (SMPS). Table 4.1 gives the details on the type of instruments used at the different stations.

Figure 4.2 shows the long-term trends based on annual averages for the three size fractions. Because of previously stated problems with inlets and DMAs, data for the small size fraction ($D_p = 11 - 41$ nm) are missing for the years 2017 – 2019 (RISØ and HCAB), 2019 (HCØ), and 2015 -2019 (HVID) for the new instruments. In addition, data are missing for 2021 for the suburban station HVID due to the low data coverage in connection with the relocation of the station.




Figure 4.2. Annual mean particle number concentrations in specific size regimes Dp = 11 - 41 nm (upper Figure), Dp = 41 – 110 nm (middle Figure), and Dp = 110 - 478/550nm (lower Figure) combined for old and new instruments at the rural background station RISØ, suburban station HVID, urban background station HCØ and street station HCAB during 2002 to 2021. Note that values for the size range 11-41 nm were not available at RISØ and HCAB from 2017-2019, HCØ in 2019 and HVID is only available for 2020.

For HCAB, the largest particle number is typically found in the smallest size regime from 11 to 41 nm (Wåhlin, 2009). The small particles mostly originate from traffic emissions and thus the highest concentrations are found close to their sources. Especially in the first ten years of measurements, the particle number concentrations in this size regime have been significantly decreasing to about 50% which is a result of technological development of vehicle engines and changes in fuel composition (Figure 4.2).

No clear trend can be observed for the small size fraction at the rural background station RISØ and the urban background station HCØ. This may be explained by the fact that one of the main sources of the small size particles (11 to 41 nm) at these background locations are natural emissions of precursors and formation of particles through atmospheric processes and only to a minor extent these particles come from anthropogenic sources.

The tendency of decreasing particle numbers is evident at HCAB also for the medium size regime (41 – 110 nm) over all the years. A general decrease in particle number is in accordance with the European trend of avoiding particulate emissions especially from road traffic, for which new environmental regulations and cleaner technologies have continuously been introduced over the years. We observe a slight decreasing trend also at the urban background and the rural measurement site, HCØ and RISØ, respectively, but the particle numbers have been stabilizing in the last five years. Similarly, this finding is observed at the suburban station HVID, where concentrations seem to be more stable but only measurements are available for the last five years. At HVID, trends need to be investigated in more detail when longer time series are available. It is possible, that at the suburban station concentrations in the middle size regime are influenced by wood combustion emissions in the area. This could explain higher concentrations compared to the urban background and the rural stations.

For the large size regime (110 – 478/550 nm), a strong decreasing trend is observed at HCAB with respect to the entire measurement period from 2002 to 2021. Similarly, such trend is observed at HCØ, HVID, and RISØ. Nevertheless, some variations are observed for some years which may be linked to this particular meteorological year. In general, decreasing trends are expected as the background aerosol is affected by general emission decreases in Europe and worldwide because of emission regulations. This implies that especially at the street station similar trends as for the smaller size regimes are observed. On the other hand, also natural processes result in relatively stable values over shorter time periods (few years). Such processes include e.g. the emission of VOCs from vegetation that in turn contribute to the formation of new particles through chemical conversion in the atmosphere. Such sources are rather more stable over shorter periods, but may also change because of changing climate over larger time periods as e.g. decades. These freshly formed particles will after some atmospheric transformation as condensation, coagulation and cloud processing enter into the large size regime where their lifetimes in the atmosphere is longer.

4.3 Long term trend for total particle number

In Figure 4.3 the total particle number is presented for the rural background station RISØ, the suburban station HVID, the urban background station HCØ and the street station HCAB in the corresponding size ranges for the old instruments ($D_p = 11\text{--}550\text{ nm}$) and for the new instruments ($D_p = 11\text{--}478\text{ nm}$).




Figure 4.3. Annual mean particle number concentrations for the total size range $D_p = 11 - 550\text{nm}$ (old instruments) and $D_p = 11 - 478\text{ nm}$ (new instruments) combined at the rural station RISØ, suburban station HVID, urban background station HCØ and street station HCAB during 2002 to 2021. Note that values were not available at RISØ and HCAB from 2017-2019, HCØ in 2019 and at HVID only data from 2020 are available.

The general patterns in terms of annual trends reflect the picture that is shown in detail in Figure 4.2 (upper, middle, lower panel). An overall decline in sub-micrometer particle number is observed at all three stations RISØ, HCØ, and HCAB. The particle number has since 2002 declined with about 68% and 64% at HCAB and HCØ, respectively. At RISØ the decrease is about 37% since the beginning of these measurements in 2005. There is a tendency to a smaller decrease with the distance to anthropogenic sources of submicron particles. The rural background station RISØ is predominantly affected by long-range transported aerosols, which show a general decline in Europe as emission regulations have been implemented and effective for various sectors, especially the transport sector. At the urban background this decrease in detected background aerosol concentrations is also reflected while at the same time this station receives more influence from the city of Copenhagen, where emission reductions have a larger impact on the trend of total particle number. Finally, the largest decrease is observed at the urban street station HCAB, which is largely influenced by direct traffic emissions from passing light and heavy-duty vehicles. For this reason, any emission reductions with respect to the changing car fleet, implementation of technological developments as e.g. exhaust after treatment or changes in fuel composition do affect the total particle number directly at this station and especially in the smallest size fraction where emitted particle numbers are typically highest at street locations.

4.4 Monthly variations in particle number concentrations

Figure 4.4 shows the monthly average concentrations for the particle number concentrations in 2021 at the urban background station HCØ, rural background station RISØ, street station HCAB and suburban station HVID. The suburban station had to be relocated in 2021 and hence there is only data from seven months. The graphs show the monthly averages for the full particle range measured with the instruments (11-478 nm) and for the small

size fraction (diameters between 11 – 41 nm), medium size fraction (diameters between 41 – 110 nm) and large size fraction (diameters between 110 – 478 nm).

At the urban background station HCØ the highest particle number concentration is observed during the summer half year with a maximum peak of about 4000 particles per cm³ in June for particles in the size range from 11-478 nm. The average particle number concentration is 22% lower in the winter half year than the summer half year. This seasonal variation is also seen for the small and medium size fractions while the large size fraction shows approximately equal particle number concentrations for the winter and summer half years.

The small size fraction originates mainly from emissions from road traffic and wood combustion in Copenhagen. Based on these sources it would have been expected that the particle number concentrations would have been higher during winter than summer or on an equal level. In addition to this meteorology also on average leads to higher concentrations during winter due to lower turbulent mixing of the air during the colder winter half year compared to the warmer summer half year.

The higher particle number concentration during summer is most likely due to a significant contribution from particles formed in the atmosphere by photochemical processes initiated by sunlight. Since these processes requires sunlight they occur mainly during the summer half year where there is much more sunlight available than during the winter half year (Figure 4.5).

The month to month variations observed at the urban background station HCØ is mainly due to the natural variations in meteorology where precipitation (Figure 4.6) is one of the main factors because wet deposition “washes” particles out of the air. Solar light has also an important impact on the particle number concentrations due to the photochemical production of particles during summer. Figure 4.5 shows the seasonal variation in the hours of sunlight. However, there is no simple relationship between particle number concentrations, precipitation and solar light since wind speed and direction, humidity and temperature also have impact on the processes that determines the amount of air borne particles transported and formed at the site.

The monthly variations observed at the rural background station RISØ follows in general the same pattern as at the urban background station HCØ although the month to month variations during the first half year are stronger compared to HCØ. Moreover, the levels are only 15% lower at Risø for the particle size range 11-478 nm. The largest difference is seen for the small size fraction (17%) and the least difference for the large size fraction (7%). The similarities between HCØ and RISØ show that it is the same sources that are responsible for the observed levels of particle number concentrations and that both stations experience similar meteorological conditions when compared on monthly basis.

The particle number concentration is about three times higher at the street station HCAB than at the urban background station HCØ and the monthly variations show a very different pattern with equal levels during winter and summer half year. Moreover, it is especially the small size fraction that is considerably higher at HCAB (about 450% higher) while the large size fraction is only about 50% higher. These observations are in line with the large impact

that emissions from road traffic have on the particle levels. Road traffic leads to high emissions of particles in the small size fraction (Wählin et al., 2009) and the seasonal variation in traffic intensity is small (Figure 4.7, Vejdirektoratet, 2022). Hence, there is no clear seasonal variation in the local traffic emissions and consequently there is no clear seasonal variation in the particle number concentrations due to variations in the most important emissions. Moreover, it is the meteorological variations that are the main reason for the month to month variations in the particle number concentrations.

The Covid-19 restrictions were not as strict in 2021 as in 2020. However, there were restrictions and these may also contribute to the monthly variations in 2021. Figure 4.7 shows the road traffic index from 2019 to 2021 (Vejdirektoratet, 2022) and from this data it can be seen that the monthly road traffic index was about 17% lower in January 2021 compared to 2019 and that the road traffic index returns to more normal levels in June 2021 in parallel with the gradual lifting of the restrictions. The street station HCAB is the station that will have the largest impact from the reduced traffic during the first five months in 2021. Since particles do not originate from traffic alone it is estimated that the particle number concentrations would have been less than 10% higher during the first five months if there had not been any Covid-19 restrictions.

Since there is only data for 6.5 months from the suburban station HVID it is not possible to evaluate the monthly variation for 2021. However, the data for the first five months follow the pattern observed at RISØ. The monthly variations will be analyzed further when there is a dataset next year.




Figure 4.4. Monthly variation in the particle number concentration in 2021. The graphs show the full particle range measured with the instruments (11-478 nm) and the small size fraction (diameters between 11 – 41 nm), medium size fraction (diameters between 41 – 110 nm) and large size fraction (diameters between 110 – 478 nm).




Figure 4.5. Monthly hours with sunshine in Copenhagen Municipality during 2021 (DMI, 2022).




Figure 4.6. Monthly precipitation in Copenhagen Municipality in 2021 (DMI, 2022).




Figure 4.7. Monthly road traffic index in Denmark for 2019, 2020 and 2021 (Vejdirektoratet, 2022). The average annual road traffic index for 2019 is set to 100%. 2020 and 2021 is clearly impacted by the Covid-19 restrictions while 2019 was the last normal year before the Covid-19 pandemic began.

4.5 Diurnal variations in particle number concentrations

Figure 4.8 to 4.11 shows the average diurnal variations of particle number concentrations for the four stations with measurements of particle number concentrations. As for PM_{2.5} and PM₁₀ the data have been divided into working days, and Saturdays and Sundays and in addition to this they have been split into winter (January-March and October-December) and summer half year (April-September). The aim of this is to analyze the differences between the warm and cold seasons in order to see the impact from the photochemical formation of particles. For the suburban station HVID only the data from winter half year are shown due to relocation of the stations and lack of data for a large part of the summer half year. The most important findings from the analysis of the diurnal patterns are as follows:

All stations show lowest particle number concentrations during the late night.

On the working day mornings, there is a clear peak that can be attributed to rush hour traffic. The morning rush hour peak (5:00-9:00) is seen both in winter and summer and it is shifted to one hour earlier during summer because of summertime. This is because the instruments always are operated on Danish wintertime. This peak is most pronounced at the street station HCAB and for the small size fraction. The morning rush hour peaks are not seen on Sundays and Saturdays.

The noon and afternoon patterns differ significantly between stations, season and day of the week. At the urban background station HCØ, the particle number concentrations decrease to a relatively stable pattern during winter until the particle number concentration begins to increase slowly during the late afternoon and reach an early evening maximum around 19:00. During summer, the particle number concentrations are higher than during winter. The particle number concentrations begin to increase from noon and the particle number concentrations reaches a local maximum during the early afternoon. The same pattern is seen at the rural background station RISØ and here the early afternoon peak is more clearly pronounced. Moreover, roughly the same level of particle number concentrations are observed on Saturdays and Sundays as on working days at RISØ. This early afternoon summer peak can best be explained by photochemically formed particles.

At the street station HCAB the influence from road traffic is strong. On working days the particle number concentrations reach a stable plateau around noon and begin slowly to decrease during the end of the afternoon. This is seen both in summer and in winter. There is no distinct afternoon rush hour peak because the afternoon rush hour is spread out over longer time periods and in addition there is a higher turbulent dilution of the particles during the late afternoon compared to the morning. Photochemically formed particles will also contribute to the particle population at HCAB but the diurnal variation is dominated by the emissions from road traffic.

On Saturdays and Sundays at the street station HCAB, there is an increase in particle number concentrations over the day reflecting that traffic activities especially in the city begins later in weekends. On Saturdays, the highest concentrations are reached around 18:00-19:00 during winter and around 21:00-22:00 during summer reflecting the road traffic in connection to the higher leisure activities in summer compared to winter. Tivoli may also add to this observation because a late evening pronounced peak in particle number is

seen at 22:00. This peak can best be explained by the fireworks in Tivoli, performed every Saturday from June to October. Sundays do not have as high particle number concentrations during the evening as observed on Saturdays.

At the urban background station HCØ and the regional background station RISØ, there are also high particle number concentrations during the working and Saturday evenings. At HCØ the highest particle concentrations are observed during the Saturday summer evenings. These high summer evening particle number concentrations are most likely due to road traffic in relation to the leisure activities during evenings in combination with the remains of the photochemically formed particles.

During winter, quite high concentrations are observed during the evening at most of the stations when looking on the medium and large size fractions. Especially at the suburban station HVID, the highest particle number concentrations are seen during the evening and on Saturdays and Sundays. These high particle number concentrations at winter evenings are most likely due to use of wood burning for household warming at this suburban site.

The findings presented above are only based on data from 2021 and for the suburban station HVID there is only data for about 6.5 months. It is therefore important to continue the analysis of the temporal variations in particle number concentrations in order to further strengthen and consolidate the conclusions about the sources of the particle number concentrations. This will be done by use of the data from 2022 where there will be a full set of data for all four stations and where Covid-19 restrictions can be completely neglected. In addition, we will also look on the long-term trends for some of the features presented above.




Figure 4.8. The diurnal variation of the particle number concentrations in 2021 at the urban background station HCØ. The diurnal variation is divided in winter and summer half years and in working days (left column), Saturdays (middle column) and Sundays (right column). The upper row shows results for the full range (11-478 nm), the second row the small size fraction (11-41 nm), the third row the medium size fraction (41-110 nm) and the fourth row the large size fraction (110-478 nm),




Figure 4.9. The diurnal variation of the particle number concentrations in 2021 at the rural background station RISØ. The diurnal variation is divided in winter and summer half years and in working days (left column), Saturdays (middle column) and Sundays (right column). The upper row shows results for the full range (11-478 nm), the second row the small size fraction (11-41 nm), the third row the medium size fraction (41-110 nm) and the fourth row the large size fraction (110-478 nm),




Figure 4.10. The diurnal variation of the particle number concentrations in 2021 at the street station HCAB. The diurnal variation is divided in winter and summer half years and in working days (left column), Saturdays (middle column) and Sundays (right column). The upper row shows results for the full range (11-478 nm), the second row the small size fraction (11-41 nm), the third row the medium size fraction (41-110 nm) and the forth row the large size fraction (110-478 nm),




Figure 4.11. The diurnal variation of the particle number concentrations in 2021 at the suburban station HVID for the winter half year. The diurnal variation is divided in working days (left column), Saturdays (middle column) and Sundays (right column). The upper row show results for the full range (11-478 nm), the second row the small size fraction (11-41 nm), the third row the medium size fraction (41-110 nm) and the forth row the large size fraction (110-478 nm),

References

Andrea, M.O., & Gelencser, A., 2006. Black carbon or brown carbon? The nature of light-absorbing carbonaceous aerosols. *Atmospheric Chemistry and Physics* 6, 3131 – 3148.

Birch, M.E., & Cary, R.A., 1996. Elemental carbon-based method for monitoring occupational exposures to particulate diesel exhaust. *Aerosol Science and Technology* 25, 221 – 241.

Cavalli, F., Viana, M., Genberg, J., & Putaud, J.-P., 2010. Toward a standardised thermal-optical protocol for measuring atmospheric organic and elemental carbon: the EUSAAR protocol. *Atmospheric Measurement Techniques* 3, 79-89.

DCE, 2021: [Air emissions \(au.dk\)](#)

Denby, B.R., Ketzel, M., Ellermann, T., Stojiljkovic, A., Kupiainen, K., Niemi, J.V., Norman, M., Johansson, C., Gustafsson, M., Blomqvist, G., Janhäll, S. & Sundvor, I., 2016. Road salt emissions: A comparison of measurements and modelling using the NORTRIP road dust emission model. *Atmos Environ*, 141, 508-522.

DMI, 2020. https://www.dmi.dk/fileadmin/user_upload/Afrapportering/Seasonsammendrag/Sammendrag_2019_sommer.pdf

DMI, 2021. [Sammendrag 2020.pdf \(dmi.dk\)](#)

Ellermann, T., Massling, A., Bossi, R., & Nøjgaard, J.K., T., 2021. The Particle Project 2020. Aarhus University, DCE – Danish Centre for Environment and Energy, xx pp. Scientific Report from DCE – Danish Centre for Environment and Energy No. 450. <http://dce2.au.dk/pub/SR450.pdf>

Ellermann, T., Nordstrøm, C., & Sørensen, M.B., 2022a. Status for måling af luftkvalitet i 2021. Aarhus University, DCE – Danish Centre for Environment and Energy, xx pp. Scientific Report from DCE – Danish Centre for Environment and Energy No. xxx., In preparation.

Ellermann, T., Nordstrøm, C., Brandt, J., Christensen, J., Ketzel, M., Massling, A., Bossi, R., Frohn, L.M., Geels, C., Jensen, S.S., Nielsen, O.-K., Winther, M., Bech Poulsen, M., Monies, C. & Sørensen, M.B., 2022b. Luftkvalitet 2020. Status for den nationale luftkvalitetsovervågning. Aarhus Universitet, DCE – Nationalt Center for Miljø og Energi, 144 s. - Videnskabelig rapport nr. 467. <http://dce2.au.dk/pub/SR470.pdf>

Frohn, L.M., Ketzel, M., Christensen, J.H., Brandt, J., Im, U., Massling, A., Andersen, C., Plejdrup, M.S., Nielsen, O.-K., Manders, A. and Raaschou-Nielsen, O., 2021: "Modelling ultrafine particle number concentrations at address resolution in Denmark from 1979 to 2018 – Part 1: regional and urban scale modelling and evaluation". *Atmospheric Environment*, vol. 264, 118631. <https://doi.org/10.1016/j.atmosenv.2021.118631>

Ketzel, M., Frohn, L.M., Christensen, J.H., Brandt, J., Massling, A., Andersen, C., Im, U., Jensen, S.S., Khan, J., Nielsen, O.-K., Plejdrup, M.S., Manders, A., van der Gon, H.D., Kumar, P., Raaschou-Nielsen, O. (2021): Modelling ultrafine particle number concentrations at address resolution in Denmark from 1979 to 2018 - Part 2: Local and street scale modelling and evaluation. *Atmospheric Environment*, vol. 264, 118633. <https://doi.org/10.1016/j.atmosenv.2021.118633>

Nøjgaard, J. K., Massling, A., Christensen, J. H., Nordstrøm, C., & Ellermann, T. 2015. The Particle Project 2011-2013, Aarhus University, DCE – Danish Centre for Environment and Energy, Scientific Report from DCE, No. 156, 55p.

Nøjgaard, J. K., Massling, A., & Ellermann, T., 2017. The Particle Project 2014-2016. Aarhus University, DCE – Danish Centre for Environment and Energy, 40 pp. Scientific Report from DCE – Danish Centre for Environment and Energy No. 233. <http://dce2.au.dk/pub/SR233.pdf>

Nøjgaard, J. K., Massling, A., & Ellermann, T., 2018. The Particle Project 2017-2018. Aarhus University, DCE – Danish Centre for Environment and Energy, 30 pp. Scientific Report from DCE – Danish Centre for Environment and Energy No. 285. [The Particle Project 2017-2018 \(au.dk\)](http://dce2.au.dk/pub/SR285.pdf)

Nøjgaard, J. K., Massling, A., & Ellermann, T., 2020. The Particle Project 2019. Aarhus University, DCE – Danish Centre for Environment and Energy, 22 pp. Scientific Report from DCE – Danish Centre for Environment and Energy No. 385. <http://dce2.au.dk/pub/SR385.pdf>

Vejdirektoratet, 2022. Nøgletal om vejtrafik, [Trafikkens udvikling i tal | Vejdirektoratet](#), Vejdirektoratet, København. Data downloaded April 2022.

Wählin, P. 2009. Measured reduction of curbside ultrafine particle number concentrations in Copenhagen, *Atmos Environ*, 43, 3645 – 3647.

THE PARTICLE PROJECT 2021

The Particle Project 2021 continues the measurements of the long-term trends of particle number concentrations and size distributions for submicron particles as well as the concentrations of elemental carbon in the ambient fine particle fraction ($PM_{2.5}$) at the Copenhagen urban background measurement station HCØ. The results from the measurements at urban background are compared to results from urban street, suburban and rural locations. The results show decreasing concentrations for both particle number concentrations and elemental carbon, which are mainly due to decreasing emissions on national as well as international level. The report also presents results from an analysis of the temporal variations of the particulate air pollution.

ISBN: 978-87-7156-693-2
ISSN: 2244-9981